1

Anodal transcranial direct current stimulation reduces collinear lateral inhibition in normal peripheral vision

Rajkumar Nallour Raveendran[*1,2], Katelyn Tsang[2], Dilraj Tiwana[2], Benjamin Thompson[2]

[1]Envision Research Institute, Wichita, Kansas, USA;

[2]School of Optometry & Vision Science, University of Waterloo, Waterloo, Ontario, Canada

* Corresponding author

Email: sachinrajopto@gmail.com

Abstract

Collinear flanking stimuli can reduce the detectability of a Gabor target presented in peripheral vision. This phenomenon is called collinear lateral inhibition and it may contribute to crowding in peripheral vision. Perceptual learning can reduce collinear lateral inhibition in peripheral vision, however intensive training is required. Our aim was to assess whether modulation of collinear lateral inhibition can be achieved within a short time-frame using a single 20-minute session of primary visual cortex anodal transcranial direct current stimulation (a-tDCS). Thirteen observers with normal vision performed a 2AFC contrast detection task with collinear flankers positioned at a distance of 2λ from the target (lateral inhibition) or 6λ (control condition). The stimuli were presented 6° to the left of a central cross and fixation was monitored with an infra-red eye tracker. Participants each completed two randomly sequenced, single-masked stimulation sessions; real anodal tDCS and sham tDCS. For the 2λ separation condition, a-tDCS induced a significant reduction in detection threshold (reduced lateral inhibition). Sham stimulation had no effect. No effects of a-tDCS were observed for the 6λ separation condition. This result lays the foundation for future work investigating whether a-tDCS may be useful as a visual rehabilitation tool for individuals with central vision loss who are reliant on peripheral vision.

Introduction

Peripheral vision is susceptible to a phenomenon called crowding, whereby it is difficult to segregate a target object from other objects that are in close proximity.[1–4] Crowding is a particular concern for patients with macular degeneration who lose central vision and are forced to rely on peripheral vision. These patients often develop a preferred retinal locus (PRL), a specific region of the peripheral retina that is used for fixation.[5–7] Crowding impairs spatial vision at the PRL leading to problems with everyday visual activates such as reading.

Crowding in peripheral vision involves cortical mechanisms that can be modulated. For example, perceptual learning can reduce letter crowding in central vision for observers with amblyopia[8,9] and in peripheral vision for observers with macular degeneration[10–13]. However, perceptual learning typically requires a large number of training trials [14] which may be a barrier for patients. In addition, the learning does not always transfer to non-trained stimuli.[15–17] Interventions that can directly modulate mechanisms within visual cortex that may contribute to crowding could complement perceptual learning techniques and enable improved vision in patients with central vision loss.

Transcranial direct current stimulation (tDCS) is a non-invasive brain stimulation technique[18–20] that has the potential to modulate neural mechanisms that contribute to crowding. tDCS involves passing a weak 1-2mA electrical current through two head-mounted electrodes (the anode and cathode) and can induce regional changes in cortical excitability and neurotransmitter concentration that outlast the duration of stimulation. For example, anodal tDCS (a-tDCS) of the motor cortex increases cortical excitability[21] and causes a regional reduction in the concentration of the inhibitory neurotransmitter GABA [22–24]. When applied to the primary visual cortex, a-tDCS modulates contrast sensitivity[25,26], visually evoked potential amplitude[25] and the visual cortex BOLD response[26]. Of particular importance for crowding, a-tDCS can immediately improve Vernier acuity and reduce surround

suppression within the near-periphery[27–29], possibly by modulating inhibition within the visual

cortex[27].

Lateral masking involves the presentation of a central target Gabor patch between two flanker

patches.[12] When the patches have collinear orientation, contrast detection thresholds for the target can

be increased (collinear inhibition or lateral masking) or reduced (collinear facilitation) depending on

target/flanker separation. Collinear inhibition is distinct from crowding, which involves impaired object

recognition rather than elevated detection thresholds. However, collinear inhibition represents a well-

established psychophysical technique for assessing low-level inhibitory mechanisms that may contribute

to crowding. Collinear inhibition and facilitation arise within the primary visual cortex[30,31]and are

present in central [8] and peripheral vision[12,32]. Maniglia et al. [12] observed that collinear lateral

inhibition could be reduced by ≈40% in normal peripheral vision after perceptual learning (20

sessions/week over 8 weeks), indicating that collinear inhibition mechanisms within the periphery are

plastic.

In this study, we took a first step towards evaluating whether visual cortex a-tDCS has the potential to

reduce crowding in peripheral vision by assessing the acute effects of a single stimulation session on

lateral inhibition in participants with normal vision. We observed that 20 min of visual cortex a-tDCS

reduced lateral inhibition in the visual periphery of healthy observers by ≈30%. This finding paves the

way for future studies designed to evaluate the possibility of inducing long-lasting changes in lateral

inhibition using a-tDCS and, in the longer term, the potential use of a-tDCS to reduce crowding in

individuals with central vision loss.

Materials and methods

Apparatus and visual stimuli

13 participants with best corrected visual acuity of ≤20/20 agreed to participate in the study. All

participants provided written, informed consent. The study was approved by the University of Waterloo

research ethics committee. All the procedures involved in this research adhered to the tenets of the

Declaration of Helsinki. Participants were instructed to fixate a 0.2°central cross and respond to visual

stimuli by pressing a key on a keyboard. Visual stimuli were created using PsychoPy[33,34] (available for

free download: http://www.psychopy.org). The stimuli were presented on a 27" LCD (ASUS -

https://www.asus.com/ca-en/Monitors/ROG-SWIFT-PG278QR/) placed at the distance of 50cm from a

chinrest. The LCD background luminance was 32 cd/m^2.

The visual stimuli consisted of a central target, Gabor patch (visible extent size: 1°, σ: 5, and spatial

frequency: 7 cpd) presented 6° to the left of the fixation cross. We selected a spatial frequency of 7 cpd

because this spatial frequency produced detection thresholds that were measurable with our 8-bit-

luminance-contrast stimulus display system. We presented our stimuli at an eccentricity of 6° because

about 70% of macular degeneration patients have a scotoma size of <5° [5] and this experiment was the

first step towards investigating the effects of a-tDCS in individuals with macular degeneration. Two

flanker Gabor patches (7 cpd, 0.8 contrast) were presented above and below the target. The flanker Gabor

patches were positioned at a distance of 2λ (lateral inhibition) or 6λ (control). Throughout the procedure,

fixation was monitored in real-time using an infrared video-based eye tracker (EyeLink II, SR Research,

Osgoode, Canada, 500 Hz sampling rate).

Psychophysical task

The 2AFC lateral masking task involved the measurement of contrast detection thresholds for the central

target Gabor patch. The initial contrast of the central Gabor patch was set at 0.5 and its contrast was

altered using a 2 down and 1 up staircase procedure with a fixed step size of 0.05. The staircase was

terminated either after 50 trials or 5 reversals. The contrast detection threshold was determined as the

mean of the last 4 reversals. Two staircases were completed and averaged for each threshold measurement. The same procedure was followed for both flanker distances of 2λ and 6λ. These particular flanker distances were selected based on a previous study reporting collinear inhibition for 2λ but not for 6λ[12] and our own pilot data supporting this observation. In fact, a 6λ flanker distance may induce facilitation.[32,35] The 2λ separation was our experimental condition and the 6λ condition was a control condition to test for any general effects of a-tDCS on contrast sensitivity for the target stimulus. A training session was provided for all participants prior to data collection. Any trials in which eye position deviated by more than 1° from fixation were immediately repeated.

Brain-stimulation (tDCS)

tDCS was delivered by a DC Stimulator MC from NeuroConn gmbh (https://www.rogue-resolutions.com/catalogue/neuro-modulation/dc-stimulator-tes/) using a pair of rubber electrodes (5cm x 5cm) placed inside saline soaked sponges. The electrodes were secured in place by the head strap of the eye tracker over electrode positions Oz (anodal electrode) and Cz (cathodal electrode) identified using the standard 10-20 EEG method (Fig 1). Participants each completed two randomly sequenced stimulation sessions conducted at least 48 hours apart; real 2mA anodal tDCS of the primary visual cortex for 20 minutes and sham tDCS where the current was ramped up and then immediately ramped down with the electrodes kept in place for 20 minutes. Participants were masked to the type of stimulation. During each test session, participants completed a block of four threshold measurements for each flanker distance pre-, during-, 5mins post- and 30mins post-stimulation. The sequence of 2λ and 6λ separation measurements was randomized within each test block.

Data Analysis

Prior to analysis, the standard deviation across reversals was calculated for each individual staircase. Any staircase with a standard deviation of 2 or greater was considered to be unreliable and excluded from analysis. An ANOVA with within-subject factors of stimulation type (anodal vs. sham) and time (pre-, during-, 5 minutes post- and 30 minutes post-stimulation) was applied to the log contrast thresholds. Post-

hoc Tukey HSD was used to compare the log contrast thresholds between different stimulation sessions.

Paired t-tests were used to compare session 1 and session 2 baselines to test for task learning. 2λ and 6λ

baselines were also compared to ensure that the 2λ separation induced collinear lateral inhibition. A p

value of <0.05 was considered statistically significant.

Fig 1. Experimental design. A) Collinear configuration of Gabor patches that has the same orientation

and phase. B) Sample picture of a participant wearing the eye tracker and the electrodes of tDCS secured

using the head strap of the eye tracker. C) Timeline of the experiment. The same timeline was used for

anodal and sham stimulations.

Results

One staircase had a SD greater than 2 and was removed from further analysis (see supporting material).

Baseline contrast detection thresholds for the central Gabor patch differed significantly between the two

flanker separation conditions, with higher thresholds for the 2λ separation than the 6λ separation (mean ±

SEM; 2λ 0.36±0.03, 6λ 0.14±0.02, t_{12} = 8.2, p < 0.001). This is consistent with previous observations of

collinear inhibition within peripheral vision at a target/flanker separation of 2λ[31,32] but not at a 6λ

separation. No significant difference between the baseline measures for session 1 and session 2 were

observed for either the 2λ (session 1, 0.39±0.15; session 2, 0.34±0.16) (t_{12} = 1.1, p=0.30) or 6λ (session 1,

0.13±0.15; session 2, 0.15±0.11) (t_{12} = 0.83, p=0.42) conditions. This indicates the absence of task

learning from one session to the next.

Fig 2 shows raw individual participant data for the a-tDCS and sham stimulation sessions for the 2λ (top

row) and 6λ (bottom row) flanker separation conditions (a table of data with mean contrast threshold and

standard deviation of staircase reversals for every individual participant is provided in the supporting

material: S1 and S2 Tables). Fig 3 shows baseline-normalized group means. For the 2λ separation, a-

tDCS significantly improved contrast detection thresholds for the central target (reduced collinear

inhibition) whereas sham stimulation had no effect. There was a significant interaction between

stimulation type [anodal vs. sham] and time [pre, during, post 5 min, post 30 min] ($F_{3, 36}$ = 3.01, p=0.042,

partial η^2 = 0.21), with post-hoc Tukey HSD comparisons revealing significantly reduced contrast

thresholds relative to baseline for the a-tDCS session during stimulation (p=0.001) and 30 min post

stimulation (p=0.01). Thresholds at the 5 min post stimulation timepoint did not differ significantly from

baseline (p=0.23). Thresholds within the sham stimulation session did not differ from baseline for any

timepoint. For the 6λ separation, there was no significant interaction between stimulation type and time

($F_{3, 36}$ = 0.46, p=0.71, partial η^2 = 0.04) indicating no difference between a-tDCS and sham on contrast

detection thresholds.

Fig 2. Contrast threshold. Log contrast detection thresholds (db) for each participant for the 2λ (top

row) and 6λ (bottom row) flanker separations during the active (left column) and sham (right column)

stimulation sessions.

Fig 3. Reduction of collinear inhibition. Reduction of collinear inhibition using anodal-tDCS. Mean

change in contrast detection threshold from baseline for the 2λ (red) and 6λ (blue) flanker separations for

the anodal (solid line) and sham (dashed line) stimulation conditions. Error bars represent ±1 SEM and

asterisk symbols represent statistical significance (p<0.05). DS, during stimulation; PS, post stimulation.

Discussion

The purpose of the study was to test the hypothesis that anodal tDCS would reduce collinear lateral

inhibition in peripheral vision of observers with normal vision. The hypothesis was based on previous

reports of improved peripheral Vernier acuity, Snellen acuity and contrast sensitivity[29] along with

reduced center-surround suppression[27] following occipital lobe a-tDCS in participants with normal

vision. Enhanced contrast sensitivity and modulation of visual cortex activity following a-tDCS have also

been observed in patients with amblyopia.[25,26] We observed a significant reduction of collinear

inhibition during and 30 min after a single 20 min application of a-tDCS to the occipital lobe. The data

collected 5 min post active stimulation exhibited that same trend as data collected during active

stimulation and 30 min post active stimulation, but they did not differ significantly from baseline or the

sham condition. These results could be likely due to noise inherent in both tDCS effects and

psychophysical tasks involving peripheral vision. The effects of a single a-tDCS session are transient, but,

overall, our results suggest that a-tDCS is able to modulate low-level lateral interactions in early visual

cortex that may contribute to crowding in peripheral vision. The reduction of collinear inhibition that we

observed for the measurements made offline (after stimulation) is consistent with previous studies

reporting stronger offline than online (during stimulation) primary visual cortex a-tDCS effects [36].

However, in addition to the offline effect, we also observed a significant reduction of collinear inhibition

for the online measurements. This is in agreement with a previous report of primary visual cortex a-tDCS

effects on surround suppression. In general, offline effects are likely to be more important for the

potential use of a-tDCS to improve vision in people with central vision loss as they indicate a lasting

influence of a-tDCS on visual cortex function.

A number of explanations have been proposed for the effects of visual cortex a-tDCS. These include

changes in response gain[29], stochastic resonance leading to increased signal strength[37], and reduced

GABA-mediated inhibition[22,24]. Our results are most clearly aligned with a reduction in cortical

inhibition as we observed an effect for the lateral inhibition condition but not the control condition that

would have also benefitted from response gain and stochastic resonance changes. Our results also support

previous work indicating that lateral inhibition takes place in V1[31,38], the primary target of our

stimulation.

Previous studies have observed that collinear lateral inhibition can be reduced using perceptual learning in

observers with normal vision[12] and observers with macular degeneration[10]. Maniglia et al.[12]

reported an approximately 40% reduction of peripheral collinear lateral inhibition after training in healthy

observers (an approximate absolute change in contrast threshold of 0.06). However, in order to achieve

this reduction, each participant underwent 160 sessions over the course of 8 weeks ($\approx$ 7600 trials/week).

In this study, we observed that a single session of anodal tDCS reduced collinear lateral inhibition by

approximately 30% (an absolute change in contrast threshold of 0.13 from baseline to 30 min post active

stimulation). This suggests that a-tDCS may enhance the effects of perceptual learning paradigms

designed to reduce collinear lateral inhibition. Indeed, a very recent study of healthy adults by Contemori

et al. observed that the combination of a different tDCS protocol, transcranial random noise stimulation,

and perceptual learning reduced peripheral crowding for trigram stimuli to a greater extent than

perceptual learning alone.[39] Furthermore, tDCS increased the transfer of learning to other tasks. Taken

together, the present results and the results of Contemori et al.[39] provide a strong foundation for the

future application of non-invasive brain stimulation to the rehabilitation of patients with central vision

loss, for whom the limitations of peripheral vision represent a major cause of visual disability [40].

One limitation of our study is that there is no consensus on whether lateral masking and crowding are

associated, although they share similar features such as increasing strength with eccentricity[41] and a

recent study showed that crowding and lateral masking are related and share similar neural

mechanisms.[42] In particular, the mechanism of crowding is likely to involve higher visual processing

centers.[41] Nonetheless, it is plausible that enhancing the early stages of visual processing by reducing

collinear inhibition will improve higher-level visual processing of crowded stimuli.[43,44] In addition, in

this study we used a 6λ target flanker separation distance as a control condition. This separation distance

was chosen because it does not induce lateral inhibition, however, flankers at this separation may induce

collinear lateral facilitation.[32,35] An alternative control condition could have involved the presentation

of orthogonally oriented flankers at a 2λ target-flanker separation.[32,35,45] The fact that we found no

effect of a-tDCS for the 6λ separation may suggest that lateral facilitation is not affected by V1 a-tDCS,

perhaps because lateral facilitation involves visual areas downstream from V1.[46,47] A study focused

directly on lateral facilitation is required to address this interesting possibility. Finally, this study

demonstrates only an acute effect of a-tDCS on lateral inhibition. The next stage in the development of

this research area will be to study the possibility of long-lasting effects, perhaps by administering multiple

a-tDCS sessions[48,49] and/or combining a-tDCS with perceptual learning[50].

Acknowledgement

The authors would like to thank Dr. Andy Silva for his assistance with PsychoPy.

References

1.   Bouma H. Interaction Effects in Parafoveal Letter Recognition. Nature. Nature Publishing Group;

1970;226: 177–178. doi:10.1038/226177a0

2.   Levi DM. Visual crowding. Curr Biol. Elsevier; 2011;21: R678–R679.

doi:10.1016/j.cub.2011.07.025

3.   Levi DM. Crowding--an essential bottleneck for object recognition: a mini-review. Vision Res.

2008;48: 635–54. doi:10.1016/j.visres.2007.12.009

4.   Pelli DG. Crowding: a cortical constraint on object recognition. Curr Opin Neurobiol. Elsevier

Current Trends; 2008;18: 445–451. doi:10.1016/J.CONB.2008.09.008

5.   Fletcher DC, Schuchard RA. Preferred retinal loci relationship to macular scotomas in a low-

vision population. Ophthalmology. Elsevier; 1997;104: 632–8. doi:10.1016/S0161-

6420(97)30260-7

6.   Timberlake GGT, Sharma MKM, Grose SAS, Gobert DD V, Guach J, Maino JJH, et al. Retinal

Location of the Preferred Retinal Locus Relative to the Fovea in Scanning Laser. Optom Vis Sci.

2005;82: 177–185. Available: http://www.ncbi.nlm.nih.gov/pubmed/15767869

7.   Cheung S-H, Legge GE. Functional and cortical adaptations to central vision loss. Vis Neurosci.

2005;22: 187–201. doi:10.1007/s11103-011-9767-z.Plastid

8.   Polat U, Ma-Naim T, Belkin M, Sagi D. Improving vision in adult amblyopia by perceptual

learning. Proc Natl Acad Sci U S A. National Academy of Sciences; 2004;101: 6692–7.

doi:10.1073/pnas.0401200101

9.   Levi DM, Li RW. Perceptual learning as a potential treatment for amblyopia: a mini-review.

Vision Res. NIH Public Access; 2009;49: 2535–49. doi:10.1016/j.visres.2009.02.010

10.   Maniglia M, Soler V, Cottereau B, Trotter Y. Spontaneous and training-induced cortical plasticity

in MD patients: Hints from lateral masking. Sci Rep. Nature Publishing Group; 2018;8: 90.

doi:10.1038/s41598-017-18261-6

11. Maniglia M, Pavan A, Sato G, Contemori G, Montemurro S, Battaglini L, et al. Perceptual

learning leads to long lasting visual improvement in patients with central vision loss. Restor

Neurol Neurosci. 2016;34: 697–720. doi:10.3233/RNN-150575

12. Maniglia M, Pavan A, Cuturi LF, Campana G, Sato G, Casco C. Reducing Crowding by

Weakening Inhibitory Lateral Interactions in the Periphery with Perceptual Learning. Goldreich D,

editor. PLoS One. Public Library of Science; 2011;6: 1–9. doi:10.1371/journal.pone.0025568

13. Chung STL, Li RW, Levi DM. Crowding between first- and second-order letter stimuli in normal

foveal and peripheral vision. J Vis. NIH Public Access; 2007;7: 10.1-13. doi:10.1167/7.2.10

14. Chung STL. Learning to identify crowded letters: Does it improve reading speed? Vision Res.

Pergamon; 2007;47: 3150–3159. doi:10.1016/J.VISRES.2007.08.017

15. Lu Z-L, Dosher BA. Mechanisms of perceptual learning. Learn Percept. NIH Public Access;

2009;1: 19–36. doi:10.1556/LP.1.2009.1.3

16. Maniglia M, Seitz AR. Towards a whole brain model of Perceptual Learning [Internet]. Current

Opinion in Behavioral Sciences. Elsevier; 2018. pp. 47–55. doi:10.1016/j.cobeha.2017.10.004

17. Fahle M. Perceptual learning: specificity versus generalization. Curr Opin Neurobiol. 2005;15:

154–160. doi:10.1016/j.conb.2005.03.010

18. Nitsche MA, Cohen LG, Wassermann EM, Priori A, Lang N, Antal A, et al. Transcranial direct

current stimulation: State of the art 2008. Brain Stimul. 2008;1: 206–223.

doi:10.1016/j.brs.2008.06.004

19. Reinhart RMG, Cosman JD, Fukuda K, Woodman GF. Using transcranial direct-current

stimulation (tDCS) to understand cognitive processing. Attention, Perception, Psychophys.

Springer US; 2017;79: 3–23. doi:10.3758/s13414-016-1224-2

20. Woods AJ, Antal A, Bikson M, Boggio PS, Brunoni AR, Celnik P, et al. A technical guide to

tDCS, and related non-invasive brain stimulation tools. Clin Neurophysiol. NIH Public Access;

2016;127: 1031–1048. doi:10.1016/j.clinph.2015.11.012

21.   Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak

281   transcranial direct current stimulation. J Physiol. Wiley-Blackwell; 2000;527 Pt 3: 633–9.

282   doi:10.1111/J.1469-7793.2000.T01-1-00633.X

283   22.   Stagg CJ, Best JG, Stephenson MC, O'Shea J, Wylezinska M, Kincses ZT, et al. Polarity-sensitive

284   modulation of cortical neurotransmitters by transcranial stimulation. J Neurosci. 2009;29: 5202–6.

285   doi:10.1523/JNEUROSCI.4432-08.2009

286   23.   Kim S, Stephenson MC, Morris PG, Jackson SR. tDCS-induced alterations in GABA

287   concentration within primary motor cortex predict motor learning and motor memory: a 7 T

288   magnetic resonance spectroscopy study. Neuroimage. Elsevier; 2014;99: 237–43.

289   doi:10.1016/j.neuroimage.2014.05.070

290   24.   Antonenko D, Schubert F, Bohm F, Ittermann B, Aydin S, Hayek D, et al. tDCS-Induced

291   Modulation of GABA Levels and Resting-State Functional Connectivity in Older Adults. J

292   Neurosci. 2017;37: 4065–4073. doi:10.1523/JNEUROSCI.0079-17.2017

293   25.   Ding Z, Li J, Spiegel DP, Chen Z, Chan L, Luo G, et al. The effect of transcranial direct current

294   stimulation on contrast sensitivity and visual evoked potential amplitude in adults with amblyopia.

295   Sci Rep. Nature Publishing Group; 2016;6: 19280. doi:10.1038/srep19280

296   26.   Spiegel DP, Byblow WD, Hess RF, Thompson B. Anodal transcranial direct current stimulation

297   transiently improves contrast sensitivity and normalizes visual cortex activation in individuals with

298   amblyopia. Neurorehabil Neural Repair. 2013;27: 760–9. doi:10.1177/1545968313491006

299   27.   Spiegel DP, Hansen BC, Byblow WD, Thompson B. Anodal Transcranial Direct Current

300   Stimulation Reduces Psychophysically Measured Surround Suppression in the Human Visual

301   Cortex. de Beeck HO p., editor. PLoS One. Public Library of Science; 2012;7: e36220.

302   doi:10.1371/journal.pone.0036220

303   28.   Bonder T, Gopher D, Yeshurun Y. The Joint Effects of Spatial Cueing and Transcranial Direct

304   Current Stimulation on Visual Acuity. Front Psychol. Frontiers Media SA; 2018;9: 159.

305   doi:10.3389/fpsyg.2018.00159

29.   Reinhart RMG, Xiao W, McClenahan LJ, Woodman GF. Electrical Stimulation of Visual Cortex

Can Immediately Improve Spatial Vision. Curr Biol. 2016;26: 1867–1872.

doi:10.1016/j.cub.2016.05.019

30.   Polat U, Mizobe K, Pettet MW, Kasamatsu T, Norcia AM. Collinear stimuli regulate visual

responses depending on cell's contrast threshold. Nature. Nature Publishing Group; 1998;391:

580–584. doi:10.1038/35372

31.   Polat U, Norcia AM. Neurophysiological Evidence for Contrast Dependent Long-range

Facilitation and Suppression in the Human Visual Cortex. Vision Res. Pergamon; 1996;36: 2099–

2109. doi:10.1016/0042-6989(95)00281-2

32.   Lev M, Polat U. Collinear facilitation and suppression at the periphery. Vision Res. Pergamon;

2011;51: 2488–2498. doi:10.1016/j.visres.2011.10.008

33.   Peirce JW. Generating stimuli for neuroscience using PsychoPy. Front Neuroinform. Frontiers;

2008;2: 10. doi:10.3389/neuro.11.010.2008

34.   Peirce JW. PsychoPy—Psychophysics software in Python. J Neurosci Methods. Elsevier;

2007;162: 8–13. doi:10.1016/J.JNEUMETH.2006.11.017

35.   Maniglia M, Pavan A, Aedo-Jury F, Trotter Y. The spatial range of peripheral collinear

facilitation. Sci Rep. Nature Publishing Group; 2015;5: 1–14. doi:10.1038/srep15530

36.   Pirulli C, Fertonani A, Miniussi C. The role of timing in the induction of neuromodulation in

perceptual learning by transcranial electric stimulation. Brain Stimul. Elsevier Ltd; 2013;6: 683–

689. doi:10.1016/j.brs.2012.12.005

37.   Schwarzkopf DS, Silvanto J, Rees G. Stochastic resonance effects reveal the neural mechanisms

of transcranial magnetic stimulation. J Neurosci. Europe PMC Funders; 2011;31: 3143–7.

doi:10.1523/JNEUROSCI.4863-10.2011

38.   Chen J, He Y, Zhu Z, Zhou T, Peng Y, Zhang X, et al. Attention-Dependent Early Cortical

Suppression Contributes to Crowding. J Neurosci. Society for Neuroscience; 2014;34: 10465–

10474. doi:10.1523/jneurosci.1140-14.2014

39.   Contemori G, Trotter Y, Cottereau BR, Maniglia M. tRNS boosts perceptual learning in peripheral

vision. Neuropsychologia. Pergamon; 2019;125: 129–136.

doi:10.1016/J.NEUROPSYCHOLOGIA.2019.02.001

40.   Wallace JM, Chung STL, Tjan BS. Object crowding in age-related macular degeneration. J Vis.

The Association for Research in Vision and Ophthalmology; 2017;17: 33. doi:10.1167/17.1.33

41.   Pelli DG, Palomares M, Majaj NJ. Crowding is unlike ordinary masking: Distinguishing feature

integration from detection. J Vis. The Association for Research in Vision and Ophthalmology;

2004;4: 12. doi:10.1167/4.12.12

42.   Lev M, Polat U. Space and time in masking and crowding. J Vis. The Association for Research in

Vision and Ophthalmology; 2015;15: 10. doi:10.1167/15.13.10

43.   Levi DM. Visual Processing in Amblyopia: Human Studies. Strabismus. Taylor & Francis;

2006;14: 11–19. doi:10.1080/09273970500536243

44.   Kiorpes L. Visual Processing in Amblyopia: Animal Studies. Strabismus. 2006;14: 3–10.

doi:10.1080/09273970500536193

45.   Shani R, Sagi D. Eccentricity effects on lateral interactions. Vision Res. Pergamon; 2005;45:

2009–2024. doi:10.1016/J.VISRES.2005.01.024

46.   Maniglia M, Trotter Y, Aedo-Jury F. TMS reveals inhibitory extrastriate cortico-cortical feedback

modulation of V1 activity in humans. Brain Struct Funct. Springer; 2019;224: 3399–3408.

doi:10.1007/s00429-019-01964-z

47.   Freeman E, Sagi D, Driver J. Lateral interactions between targets and flankers in low-level vision

depend on attention to the flankers. Nat Neurosci. Nature Publishing Group; 2001;4: 1032–1036.

doi:10.1038/nn728

48.   Rushmore RJ, Desimone C, Valero-Cabré A. Multiple sessions of transcranial direct current

stimulation to the intact hemisphere improves visual function after unilateral ablation of visual

cortex. Eur J Neurosci. NIH Public Access; 2013;38: 3799–3807. doi:10.1111/ejn.12373

49.   Behrens JR, Kraft A, Irlbacher K, Gerhardt H, Olma MC, Brandt SA. Long-Lasting Enhancement

of Visual Perception with Repetitive Noninvasive Transcranial Direct Current Stimulation. Front

Cell Neurosci. Frontiers Media S.A.; 2017;11: 238. doi:10.3389/fncel.2017.00238

50. Spiegel DP, Li J, Hess RF, Byblow WD, Deng D, Yu M, et al. Transcranial direct current

stimulation enhances recovery of stereopsis in adults with amblyopia. Neurotherapeutics. 2013;10:

831–9. doi:10.1007/s13311-013-0200-y

Figures:

Supporting Information

S1 Table. Mean contrast threshold values (db) and SD of reversals for each participant in 2λ condition

S2 Table Mean contrast threshold values (db) and SD of reversals for each participant in 6λ condition

		2λ							
		BL_A	DS_A	PS5_A	PS30_A	BL_S	DS_S	PS5_S	PS30_S
S1	Avg. threshold	0.100	0.088	0.144	0.100	0.132	0.108	0.092	0.084
	SD of reversals	0.047	0.041	0.089	0.032	0.054	0.048	0.018	0.043
S2	Avg. threshold	0.249	0.177	0.229	0.223	0.235	0.208	0.213	0.213
	SD of reversals	0.032	0.034	0.028	0.056	0.045	0.064	0.051	0.047
S3	Avg. threshold	0.765	0.400	0.453	0.553	0.343	0.443	0.447	0.500
	SD of reversals	0.084	0.089	0.024	0.043	0.053	0.088	0.059	0.042
S4*	Avg. threshold	0.535	0.210	0.360	0.280	0.280	0.377	0.277	0.227
	SD of reversals	0.019	0.012	0.016	0.016	0.018	0.046	0.056	0.024
S5	Avg. threshold	0.435	0.095	0.220	0.165	0.270	0.305	0.185	0.120
	SD of reversals	0.025	0.019	0.028	0.034	0.026	0.030	0.100	0.037
S6	Avg. threshold	0.385	0.265	0.290	0.225	0.393	0.267	0.187	0.260
	SD of reversals	0.025	0.044	0.012	0.019	0.031	0.042	0.031	0.020
S7	Avg. threshold	0.277	0.223	0.169	0.214	0.385	0.375	0.355	0.450
	SD of reversals	0.069	0.063	0.063	0.043	0.066	0.038	0.019	0.012
S8	Avg. threshold	0.420	0.389	0.349	0.349	0.240	0.205	0.235	0.345
	SD of reversals	0.026	0.020	0.061	0.047	0.016	0.019	0.050	0.038
S9	Avg. threshold	0.525	0.415	0.490	0.410	0.405	0.443	0.513	0.408
	SD of reversals	0.034	0.034	0.026	0.012	0.062	0.038	0.123	0.030
S10	Avg. threshold	0.212	0.240	0.200	0.168	0.343	0.275	0.295	0.350
	SD of reversals	0.018	0.024	0.079	0.041	0.025	0.064	0.082	0.105
S11	Avg. threshold	0.285	0.200	0.285	0.205	0.385	0.310	0.305	0.300
	SD of reversals	0.038	0.037	0.019	0.034	0.082	0.062	0.019	0.059
S12	Avg. threshold	0.344	0.248	0.336	0.328	0.550	0.375	0.405	0.420
	SD of reversals	0.059	0.059	0.055	0.050	0.012	0.019	0.053	0.028
S13	Avg. threshold	0.504	0.420	0.344	0.388	0.437	0.417	0.277	0.477
	SD of reversals	0.074	0.077	0.022	0.052	0.070	0.052	0.021	0.031
Overall Mean ± SD		**0.387±0.173**	**0.259±0.115**	**0.297±0.105**	**0.277±0.124**	**0.338±0.107**	**0.316±0.102**	**0.291±0.116**	**0.319±0.133**

372 Abbreviations: BL – baseline; DS – during stimulation; PS5 and PS30 – post-stimulation after 5min and 30min respectively. A – active; S – Sham. *denotes the participant with a
373 staircase removed due to unreliability.

		6λ							
		BL_A	DS_A	PS5_A	PS30_A	BL_S	DS_S	PS5_S	PS30_S
S1	Avg. threshold	0.028	0.024	0.028	0.064	0.018	0.028	0.033	0.020
	SD of reversals	0.011	0.017	0.018	0.062	0.023	0.021	0.028	0.019
S2	Avg. threshold	0.037	0.080	0.075	0.069	0.033	0.043	0.040	0.043
	SD of reversals	0.035	0.060	0.063	0.034	0.032	0.033	0.067	0.029
S3	Avg. threshold	0.409	0.423	0.378	0.383	0.140	0.368	0.355	0.278
	SD of reversals	0.053	0.029	0.033	0.117	0.075	0.021	0.056	0.039
S4	Avg. threshold	0.289	0.311	0.186	0.274	0.150	0.310	0.190	0.048
	SD of reversals	0.066	0.045	0.085	0.025	0.035	0.082	0.078	0.028
S5	Avg. threshold	0.058	0.086	0.093	0.063	0.370	0.128	0.228	0.118
	SD of reversals	0.036	0.082	0.117	0.069	0.060	0.059	0.063	0.132
S6	Avg. threshold	0.145	0.230	0.355	0.145	0.186	0.155	0.260	0.328
	SD of reversals	0.019	0.012	0.066	0.019	0.063	0.093	0.034	0.043
S7	Avg. threshold	0.026	0.131	0.069	0.033	0.045	0.063	0.050	0.086
	SD of reversals	0.015	0.169	0.047	0.024	0.066	0.071	0.051	0.061
S8	Avg. threshold	0.188	0.055	0.253	0.163	0.070	0.077	0.085	0.097
	SD of reversals	0.164	0.033	0.038	0.105	0.039	0.098	0.050	0.091
S9	Avg. threshold	0.125	0.130	0.115	0.100	0.280	0.405	0.178	0.185
	SD of reversals	0.026	0.024	0.030	0.024	0.057	0.100	0.048	0.058
S10	Avg. threshold	0.116	0.172	0.084	0.108	0.140	0.278	0.255	0.283
	SD of reversals	0.038	0.127	0.043	0.175	0.034	0.140	0.023	0.063
S11	Avg. threshold	0.015	0.187	0.170	0.265	0.265	0.070	0.370	0.318
	SD of reversals	0.019	0.031	0.012	0.034	0.085	0.100	0.176	0.063
S12	Avg. threshold	0.093	0.098	0.108	0.073	0.120	0.048	0.115	0.053
	SD of reversals	0.054	0.055	0.111	0.084	0.030	0.026	0.195	0.032
S13	Avg. threshold	0.180	0.090	0.127	0.117	0.158	0.148	0.228	0.195
	SD of reversals	0.075	0.049	0.037	0.043	0.125	0.076	0.122	0.048
Overall Mean ± SD		**0.131±0.115**	**0.155±0.112**	**0.157±0.109**	**0.152±0.104**	**0.143±0.104**	**0.163±0.132**	**0.183±0.114**	**0.158±0.113**

374   Abbreviations: BL – baseline; DS – during stimulation; PS5 and PS30 – post-stimulation after 5min and 30min respectively. A – active; S – Sham.