Genome-Wide Polygenic Risk Scores and Prediction of Gestational Diabetes in South Asian Women

Amel Lamri[1,2], Shihong Mao[2], Dipika Desai[2], Milan Gupta[1,3], Guillaume Paré[2,4], Sonia S. Anand[1,2,5]*

1. Department of Medicine, McMaster University Hamilton, Ontario, Canada

2. Population Health Research Institute (PHRI), Hamilton, Ontario, Canada

3. Canadian Collaborative Research Network (CCRN), Brampton, ON, Canada

4. Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario, Canada

5. Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada

Corresponding Author: Dr. Sonia S Anand,

McMaster University, MDCL 3202,

1280 Main St W, Hamilton,

Ontario L8S 4K1, Canada.

Tel.: +1-905-525-9140, ext. 21523;

Fax: +1-905-528-2814.

e-mail: anands@mcmaster.ca

Keywords: Genome-wide polygenic risk score; gestational diabetes, type 2 diabetes, South Asian, women, heritability.

Word Count: 4373

ABBREVIATIONS

1KG	1000 Genomes
AUC	Area Under the Curve
BMI	Body Mass Index
CI	Confidence Interval
DIAGRAM	DIAbetes Genetics Replication and Meta-analysis
GDM	Gestational Diabetes
PRS	Polygenic Risk Score
GraBLD	Gradient Boosted and LD adjusted
GRM	Genomic Relationship Matrix
GREML	Genomic relatedness matrix residual maximum likelihood
GRS	Genetic Risk Score
GWAMA	Genome-Wide Association Meta-Analysis
GWAS	Genome-Wide Association Study
LD	Linkage Disequilibrium
LDMS	LD and MAF stratified
MAF	Minor Allele frequency
MAGIC	Meta-Analyses of Glucose and Insulin-related traits Consortium
MS	MAF stratified
OR	Odds Ratio
PCA	Principal Component analysis
P+T	Pruning and Thresholding
ROC	Receiver Operating Characteristic
SC	Single component
SE	Standard Error
SNP	Single Nucleotide polymorphism
START	South Asian Birth Cohort
T2D	Type 2 Diabetes

ABSTRACT

Gestational diabetes Mellitus (GDM) affects 1 in 7 births and is associated with numerous adverse health outcomes for both mother and child. GDM is suspected to share a large common genetic background with type 2 diabetes (T2D). The first aim of this study, was to characterize different GDM polygenic risk scores (PRSs) using data from the South Asian Birth Cohort (START). The second aim of this study was to estimate the heritability of GDM.

PRSs were derived for 832 South Asian women from START using the pruning and thresholding (P+T), LDpred, and GraBLD methods. Weights were derived from multi-ethnic (Mahajan *et al.*, 2014) and white Caucasian (Scott *et al.*, 2017) studies of the DIAGRAM consortium. Association with GDM was tested using logistic regression. Heritability of GDM was estimated using the GRMEL approach. Results were replicated in samples from the UK Biobank (UKB) study.

The top P+T, LDpred and GraBLD PRSs were all based on Mahajan *et al*. The best PRS was highly associated with GDM in START (AUC=0.62, OR=1.60 [95% CI=1.44–1.69]), and in South Asian (AUC=0.65) and white British (AUC=0.58) women from UKB. Heritability of GDM approximated 0.55±0.83 in START and 0.18±0.22 in white British women from UKB.

Our results highlight the importance of combining genome-wide genotypes and summary statistics from large multi-ethnic studies to optimize PRSs in South Asians.

INTRODUCTION

Gestational diabetes mellitus (GDM) is defined as dysglycemia due to elevated blood glucose levels first identified during pregnancy, and is specifically defined based on glucose response to an oral glucose challenge test in pregnancy. GDM has been associated with numerous adverse health outcomes affecting mother and child, both during and after pregnancy [1,2]. Because of its increasing prevalence (~ 1 in 7 births), GDM has become a major health concern worldwide [3]. Nevertheless, the prevalence of GDM largely varies from one region of the globe to the other, and South Asian women have been shown to be at higher risk of GDM than white British women.

Although GDM is thought to have a strong genetic component, to our knowledge, no studies have estimated the heritability of GDM. Nevertheless, since GDM and T2D have similar risk factors and share common pathophysiological pathways, the heritability of GDM is thought to be similar to that of T2D.

Numerous genome-wide association studies (GWASs) and genome-wide association meta-analysis (GWAMAs) of glucose related traits and T2D have been conducted in non-gravid populations, and summary statistics from large consortia (e.g., MAGIC and DIAGRAM) are publicly available [4-13]. By contrast, few studies of genetic determinants of GDM have been conducted or published. For instance, only two studies sought to identify genes associated with dysglycemia, GDM, and diabetes during pregnancy by GWAS [14,15]. Top signals from these studies were located within/near *CDKAL1*, *MTNR1B, GCKR*, *PCSK1, PPP1R3B* and *G6PC2*, which were previously known for their association with glucose metabolism and T2D [14,15]. In addition, other T2D associated loci (e.g., *TCF7L2, PPARG, CDKN2A/B, KCNQ1, GCK*, etc.) were also significantly associated with GDM when tested separately [16-39], or combined in genetic risk scores (GRS) [33,34,40-42].

GRS are used to capture genetic information at one or more loci. Most of published studies

interested in complex traits/diseases and using GRS typically combine data for a small

number of single nucleotide polymorphisms (SNPs), and the predictive power of these GRS is

sub-optimal [43]. However, with the increased availability of genome-wide genotypes and

publicly available data from large consortia, GRS with a larger number of variants are being

used, and the predictive value of these genome-wide polygenic risk scores (PRSs) has

substantially improved [44,45].

PRSs can be derived using different approaches, however, these require both summary

statistics from an external GWAS and genetic data for a reference panel for between-variants

linkage disequilibrium LD (LD) calculations. Pruning and thresholding (P+T) is a commonly

used heuristic approach to derive PRSs in which variants are filtered based on an empirically

determined P-value threshold. Linked variants are further clustered in different groups and

SNPs with the highest significance (lowest P values) in each group are prioritized and

included in the PRS, while variants of less significance within the group are pruned out [46].

Other programs have been shown to improve the predictive value of the score by allowing the

inclusion of a larger number of independent as well as linked variants into the score using

different approaches. For instance, LDpred, another commonly used method, estimates the

mean weight of each variant, assuming a prior knowledge of the genetic architecture of the

trait (fraction causal), and using a Bayesian approach [47]. More recently, we developed the

gradient boosted and LD adjusted (GraBLD), a new PRS building approach which applies

principles of machine-learning to estimate SNP weights (gradient boosted regression trees),

and regional LD adjustment [48].

The first objective of this study was to determine the optimal gene scores and investigate the

association of genetic variants combined in these PRSs with GDM in South Asian women

participating to the South Asian Birth Cohort (START). We considered several parameters: 1)

two different sources of summary statistics, namely Mahajan *et al.*, 2014 (trans-ethnic

GWAMA) [5] *vs.* Scott *et al.*, 2017 [6] (GWAMA in white Caucasians); 2) two templates for LD

calculation (1000 Genomes phase 3 [49] *vs.* START genotypes); 3) different minimal values of

the number of samples in each SNP's analysis in the consortia; 4) weighted *vs.* unweighted

PRSs; 5) three methods to derive the PRSs; Pruning and Thresholding (P+T),. LDpred and

GraBLD and; 6) different P-value thresholds (for P+T and LDpred PRSs only). The second

objective was to estimate the heritability of GDM from: 1) genome-wide data; 2) common

variants; and 3) SNPs in the best P+T PRS. PRS results were further validated in South Asian

women from UK Biobank, and both PRS and heritability results were replicated in a subset of

white British women from UK Biobank[50].

78   RESULTS:

79   Population characteristics:

Table 1 shows the characteristics of START and UK Biobank women included in the

analyses. Because of major differences in recruitment strategies, inclusion criteria and study

protocols, selected participants from the UK Biobank were of older age, higher weight, and

body mass index (BMI) compared to START participants. Furthermore, the proportion of

participants with GDM was significantly lower in South Asian women from the UK Biobank

and even more so in European women of the same study.

86   Minimum sample size per SNP in consortium data:

Summary statistics were derived from a trans-ethnic (Mahajan *et al.*, 2014 [5]) and white

Caucasian (Scott *et al.*, 2017 [6]) GWAMAs of the DIAGRAM consortium. In each study, the

number of participants tested for association with T2D was different for each SNP, and

ranged between 25 – 110,219 and 4,731 – 158,186 in Mahajan *et al.* and Scott *et al*

respectively (Supplementary Figure 3, Supplementary Table 1).

We derived several PRSs for which the list of variants was restricted to SNPs tested in at least

0, 85, 90, 95 and 98% of the maximum sample size in the consortium GWAMA of interest.

The number of SNPs used in the PRSs and the percentage of SNP loss for each one of these

thresholds are shown in Supplementary Table 1. The percentage of variants lost after this

filtering was the most substantial in PRSs based on Mahajan *et al.*, with only 346,290

polymorphisms remaining when keeping variants tested in ≥ 95% samples (Supplementary

Table 1).

For both Mahajan *et al.* and Scott *et al.* based PRSs, the optimal minimum percent of

participants to keep varied depending on the method used (P+T, LDpred or GraBLD), the

source of LD estimates (LD_{START} or LD_{1KG}) and the consortium P-value threshold (Figure 1,

Figure 2, Supplementary Figure 3 and Table 2).

With an AUC of 0.62, the best GraBLD PRS included 1,305,596 SNPs and was derived using

weights from Mahajan *et al.* ($W_{Mahajan}$); LD from START (LD_{START}); SNPs tested in ≥ 90% of

the samples in the consortium data ($N_{90\%}$). This top GraBLD PRS will hereafter be referred to

as $GraBLD\_PRS_1\_W_{Mahajan}\_LD_{START}\_N_{90\%}$. The best P+T was comprised of 9,274 SNPs,

showed an AUC of 0.62 and was derived using the following parameters: Weights from

Mahajan *et al.* ($W_{Mahajan}$); LD from 1KG (LD_{1KG}); SNPs tested in ≥ 85% of the maximum

sample size ($N_{85\%}$) and a maximum P-value of 0.016 in its reference consortium GWAMA

($P_{0.016}$). This top P+T PRS will be referred to as $PT\_PRS_1\_W_{Mahajan}\_LD_{1KG}\_N_{85\%min}\_P_{0.016.}$

Finally, with an AUC of 0.62 as well, the best LDpred PRS included 1,290,525 SNPs and was

derived using weights from Mahajan *et al.* ($W_{Mahajan}$); LD from 1KG (LD_{1KG}); SNPs tested in

≥ 85% of the samples in the consortium data ($N_{85\%}$). This top LDpred PRS will hereafter be

referred to as $LDpred\_PRS_1\_W_{Mahajan}\_LD_{1KG}\_N_{85\%}$. Detailed characteristics and rankings of

the best PRSs are shown in Supplementary Table 2

Pruning and Threshold PRSs

AUCs of unweighted P+T PRSs were similar to AUCs for their weighted counterpart at very

low P-value thresholds (clump $P \leq 0.004$). Interestingly, the inclusion of variants with

association P-values that are less significant than the usual GWAS significance threshold ($5 \times$

10^{-8}) always resulted in a considerable increase in the predictive value of the scores. Optimal

AUCs were reached at clump P-values ranging from 0.016 - 0.20 depending on the source of

LD or the consortium used (Figure 1, Supplementary figure 2 and Table 2). Passed these

maximal points, the difference between weighted and unweighted PRSs gradually increased,

with the weighted PRSs performing better than their unweighted counterparts (Figure 1,

Supplementary figure 2).

LDpred PRSs:

Similarly to P+T PRSs, the increase in the fraction of causal SNPs (from P values of 5×10^{-8}

to P = 0.0005 corresponding to 0.5 and 7.5 % of consortium SNPs in Mahajan *et al.* and Scott

et al. respectively) highly improved the predictive value of the PRSs (Figure 2). The increase

in the fraction causal passed this point was not associated with a significant change in the

AUC of LDpred PRSs (Figure 2).

GraBLD vs. P+T vs. LDpred PRSs: As previously mentioned, whether the performance of a

PRS derived using a given method was better than that of its different counterparts (other two

methods) largely depended on the consortium data, the source of LD, the minimum % of

participants, and the maximum clump P-value cutoffs used. Nevertheless, when comparing

the best PRSs derided from each method, no significant difference was observed between

GraBLD, LDpred and P+T (AUCs=0.62, Table 2, $P_{\text{pairwise differences}} = 0.95$). When comparing

P+T to LDpred only, AUCs were higher and more stable in LDpred PRSs than in P+T PRS

for high P-value thresholds (> 0.1) (Figure 3).

Mahajan et al. vs. Scott et al. PRSs: The predictive value of most Mahajan-based P+T PRSs

were higher than that of their Scott-based counterparts (Figure 4). For GraBLD and LDpred

PRSs, all Mahajan based PRSs had higher AUCs than Scott based PRSs (Figure 4, data not

shown). Finally, the best Mahajan-based PRSs always outperformed the top Scott-based PRSs

for all three methods (Table 2, Supplementary Table 2)

LD from 1000 Genomes vs. START: AUCs of the best LD_{START} PRSs were not significantly

different from AUCs of their LD_{1KG} counterpart for P+T, LDpred and GraBLD PRSs (Figure

5, Table 2).

Association with GDM:

The association results of the top PRSs ($GraBLD\_PRS_{1\_}W_{Mahajan\_}LD_{START\_}N_{90\%,}$

$PT\_PRS_{1\_}W_{Mahajan\_}LD_{1KG\_}N_{85\%\_}P_{0.016}$ and $LDpred\_PRS_{1\_}W_{Mahajan\_}LD_{1KG\_}N_{85\%}$) with GDM

(univariate models) are shown in Table 2 (continuous PRSs) and Table 3 (categorical PRSs).

The odds of developing GDM was 2 to 2.5 fold higher in participants with the highest PRSs

(top 25%) compared to the rest (75%) of the study population, depending on the type of PRS

used. When analyzing participants with high and low PRSs values only, our results show that

participants with the highest PRS values (top 25%) had between 3 and 3.4 fold increase in

their risk of GDM compared to the participants with the lowest PRS values (bottom 25%).

These results were similar in both UK Biobank South Asian and white British replication

datasets (Table 3).

Heritability

In order to better characterize the genetic architecture of GDM, heritability was estimated from 1) genome-wide genotype data ($h^2_{WG\_SNPs}$); 2) common variants (MAF $\geq$ 0.01); and 3) SNPs included in our top P+T PRS, in South Asian women from START, as well as for a subgroup of white British women from UK Biobank. The results are shown in Table 4. Due to a lack of power, standard errors for our heritability estimates were large, and most lower and upper bound values of the 95% confidence intervals were close to, or crossed the respective 0 and 1 boundaries. The proportion of the variance in GDM explained by genome-wide data ($h^2_{WG\_SNPs}$) in South Asian women from the START study was 0.57 $\pm$ 0.88 and $h^2_{WG\_SNPs}$ estimated in women from white British UK Biobank samples was 0.15 $\pm$ 0.22 (Table 4). Heritability attributed to common variants (MAF $\geq$ 1%) reached 0.5 $\pm$ 0.78 and 0.11 $\pm$ 0.14 in our START and UK Biobank samples, which explained 88% and 73% of the $h^2_{WG\_SNPs}$ respectively. Heritability estimated from the SNPs used in our top P+T PRS explained 31.6% and 20 % of $h^2_{WG\_SNPs}$ in START and UK Biobank respectively.

DISCUSSION

In this study, we built several thousands of GDM PRS using genome-wide genotypes, large consortium data and 3 different methods. Our best PRS was built using the LDpred method, with weights extracted from Mahajan *et al.* and LD calculated using 1KG genotypes. This PRS was significantly associated with GDM in South Asian women from the START study, an observation that was successfully replicated in South Asian and white British women from UK Biobank. Participants with the highest PRS values had an increased risk of GDM when compared to the other groups.

We observed a considerable difference in the proportion of participants with GDM between South Asian women from the START study (36.2 %), South Asian women from UK Biobank

(2.2%), and white British women from UK Biobank (0.43%). This disparity is likely due to

major differences in the study design, recruitment strategies and definition of GDM between

the two studies involved. For instance, the definition of GDM status in START was based on

glucose levels measurements performed during pregnancy in response to an oral glucose

challenge, and South Asian-specific diagnosis cutoffs were used. On the other hand, GDM

status was retrospectively self-reported by UK Biobank participants, which most likely

resulted in an increased number of misclassifications and a reduced number of reported GDM

cases. In an effort to refine the phenotype in UK Biobank, our control group was restricted to

women without GDM who also went through at least one live birth. Nevertheless, the lack of

sensitivity and specificity of the GDM phenotype likely remains an issue in UK Biobank.

Summary statistics from two large T2D GWAMAs were used to build our PRSs. One of the

major advantages in using data from Mahajan *et al.* was that ~20% of its participants in their

publically available data originated from the South Asian sub-continent. The study also had a

large maximum number of cases and controls, but many of the SNPs included in the meta-

analysis were tested in a much smaller sample (Supplementary figure 3, Supplementary Table

1). On the other hand, no South Asian participants were included in the GWAMA performed

by Scott *et al.* but the average number of samples tested for each SNP was larger than in

Mahajan *et al*. Our results show that Mahajan-based PRSs consistently outperformed their

Scott-based counterparts in spite of a lower genome coverage and smaller average number of

participants per SNP. This highlights the importance of using consortium data of the same

ethnic group than the study at hand whenever possible. However, since Mahajan *et al.*'s

summary statistics were derived from a blend of participants of different ethnicities, our top

PRS could likely be improved if built based on summary statistics derived from an equally

powered GWAMA performed in South Asians only.

Several reports suggest that T2D and GDM share a common genetic background. In the

absence of publicly available data of large GDM GWASs, summary statistics from a T2D

consortium were used to derive our scores. Our results show that a T2D PRSs can be highly

predictive of GDM in South Asian and European-origin women, hence confirming the

hypothesis of a common genetic background between these two diseases. However, the effect

size of the genetic variants could be different between the two conditions, and some loci could

be specific to each disease. Although these differences should not affect our models

comparisons, we expect that the predictive value of GDM PRSs will be further improved if

built using weights from a large GDM GWAS or GWAMA.

A significant conclusion derived from this study is that, whatever the consortium or the

method used, restricting the list of SNPs to GWAS significant variants (P value $\leq 5 \times 10^{-8}$)

drastically reduces the predictive value of the PRSs. Unfortunately, many studies still rely on

this threshold to select their loci of interest to derive their risk scores. We recommend the use

of higher P-value thresholds (> 0.01 in our case) whenever possible in order to increase the

predictive value of the PRSs.

Based on our results, weighted PRSs perform similarly, or better than their unweighted

counterparts in general. For P+T PRS, the predictive value of the unweighted PRSs are

especially lower than for weighted PRS at high P-value cutoffs. Hence, we recommend the

use weighted PRSs whenever possible. If unweighted PRSs are used, P-value cutoffs should

be set between 0.01 and 0.1.

When comparing the best PRSs, our results suggest that the GraBLD, P+T and LDpred

methods perform equally well in terms of disease prediction as measured by the AUC.

Nevertheless, the identification of the optimal P+T, and LDpred PRSs required the test of

several thousand predictors (n = 2,560 and 1280 respectively), when a similar result was

achieved by testing 40 GraBLD models only. While this may lead one to slightly favor the use

of LDpred or GraBLD for the building of PRSs, the P+T remains the method of choice in our

opinion, given the fact that it required less SNPs and was easier to implement using the

PLINK software.

T2D's SNP-based heritability has recently been estimated at 0.54 [95%CI: 0.47 - 0.61] [51]. Our

results from START suggest that the heritability of GDM is similarly high ($h^2_{WG\_SNPs}$ in

START=0.58, SE=0.43) although the standard error is large. Heritability estimates were

considerably smaller in European participants in UK Biobank. This could be explained by the

difference between the disease prevalence in START and UK Biobank as previously

discussed, and potentially, by differences in the environments and lifestyles of the participants

included in the two studies. Another potentially interesting observation derived from our

heritability results is that a large proportion of the genetic variance of GDM (between 73.3%

and 87.7% depending on the ethnicity) can be explained by common SNPs. Furthermore, our

results suggest that a relatively large proportion of the genetic variance (between 20% and

31.6%) is captured by an even smaller fraction of SNPs (~2.5%, N=9,274) that are included in

our top P+T PRS. However, given the small number of cases in our studies, our GREML tests

are likely underpowered, resulting in very large standard errors. Hence, our heritability results

should be interpreted with caution, and larger studies are needed in order estimate the

heritability of GDM with higher accuracy. An interesting future analysis would be to further

partition these heritability estimates into maternal and fetal genetic effects. Indeed, since

maternal glucose cut-offs used to define GDM also rely on the fetal phenotypes (size and

adiposity related measures), we strongly suspect that the fetus's genome will also contribute

to the mother's risk of developing the disease. Unfortunately, this analysis could not be

performed in our study because of our small sample size.

In conclusion, our results show that the predictive value of polygenic risk scores in South

Asian women can be greatly improved by combining genome-wide genotyping data and by

extracting summary statistics from large multi-ethnic genome-wide meta-analysis.

METHODS

Study design and participants

The South Asian Birth Cohort (START) study: START is a prospective cohort designed to

evaluate the environmental and genetic determinants of cardiometabolic traits of South Asian

pregnant women and their offspring living in Ontario, Canada. The rationale and study design

are described elsewhere [52]. In brief, 1,012 South Asian (people who originate from the Indian

subcontinent) pregnant women, between the ages of 18 and 40 years old, were recruited

during their second trimester of pregnancy from the Peel Region (Ontario, Canada) through

physician referrals between July 11, 2011 and Nov. 10, 2015. All START participants signed

an informed consent including genetic consent, and the study was approved by local ethics

committees (Hamilton Integrated Research Ethics Bard, William Osler Health System, and

Trillium Health Partners). A detailed description of the maternal measurements has been

published previously [53].

UK Biobank: The UK Biobank is a large population-based study which includes over

500,000 participants living in the United Kingdom [50]. Men and Women aged 40 - 69 years

were recruited between 2006 and 2010 and extensive phenotypic and genotypic data about the

participants was collected, including ethnicity and a question regarding past history of GDM.

Details of this study are available online (https://www.ukbiobank.ac.uk) [50]. Data from UK

Biobank were used in order to validate the results from the START study.

Derived variables

START Study: GDM status was determined using the South Asian specific cutoffs as defined in the Born in Bradford study (fasting glucose level of 5.2 mmol/L or higher, or a 2-hour post load level of 7.2 mmol/L or higher) [54]. Self-reported GDM status was used if these measures were unavailable. Participants with a history of T2D prior to pregnancy were excluded. Using these criteria, 832 START participants with known GDM status (301 cases and 531 controls) and available genotypes were included in the analysis. The South Asian ethnicity/ancestry of participants was validated using genetic data.

UK Biobank: Participants in the UK Biobank completed questionnaires at several time points (questionnaire of initial assessment visit, 2006-2010; questionnaire of first repeat assessment visit, 2012-2013; questionnaire of imaging visit, 2014 onwards). For the purpose of our study, GDM cases were defined as women who reported having had diabetes during pregnancy only, collected at any time point using questionnaires. The control group was comprised of women who: 1) had at least one child (self-reported, live births only) 2) had never been diagnosed with diabetes or GDM in all assessments. The ethnicity/ancestry of participants was validated using genetic data.

DNA extraction, genotyping, imputation, filtering and SNP extraction:

START: DNA was extracted and genotyped from a total of 867 samples (START mothers) using the Illumina Human CoreExome-24 and Infinium CoreExome-24 arrays (Illumina, San-Digeo, CA, USA). Data was cleaned using standard quality control (QC) procedures [55] and 837 women samples passed the QC. Genotypes were subsequently phased using SHAPEIT v2.12 [56], and imputed with the IMPUTE v2.3.2 software [57], using the 1000 Genomes (phase 3) data as a reference panel [49]. Variants with an info score $\geq$ 0.7 were kept for analysis.

Addition data manipulation and SNP selection criteria for the building of the PRSs are

detailed in Supplementary Information and Supplementary Figure 1.

UK Biobank: A total of ~500,000 participants from the UK biobank were genotyped using

the UK BiLEVE or UK Biobank Affymetrix Axiom arrays. Detailed QC, phasing and

imputation procedures have previously been described [58]. As a result, 3,169 and 220,703

unrelated South Asian and white British women respectively passed QC. Among these, 2,386

and 184,869 participants had available GDM status respectively, and were used to replicate

our results from the START study. Genotypes for > 98% of SNPs included in our top START

GDM PRSs were available (info score ≥ 0.6) and were extracted for the replication.

Consortium data

Summary statistics (P-value, effect size) from the following two DIAGRAM sources were

used in order to build the PRSs:

1) Mahajan et al., Nature Genetics, 2014

This trans-ethnic GWAMA included up to 12,171 T2D cases and 56,862 controls of

European ancestry; 6,952 cases and 11,865 controls of East Asian ancestry; 5,561 cases

and 14,458 controls South Asian ancestry and 1,804 cases and 779 controls of Mexican

and Mexican American ancestry [5]. This study was selected for its multi-ethnic

composition and its inclusion of South Asian participants.

2) Scott et al., Diabetes, 2017

This study combines data from 18 GWASs, for a total of 26,676 T2D cases and 132,532

controls of European ancestry [6]. This study was selected for its large sample size and its

relatively homogenous samples (100% white Caucasians).

Summary statistics of these studies were downloaded from DIAGRAM's main website (http://www.diagram-consortium.org).

Templates for LD calculation

LD calculations used to build the PRSs were derived from the following two genotyping datasets: 1) START study (LD source hereafter referred to as LD_{START}); and 2) the 1000 Genomes consortium (LD source hereafter referred to as LD_{1KG}) phase 3. Genotypes of 1000 Genomes participants were downloaded from the project's data portal (http://www.internationalgenome.org), and a subset of participants was created in order to match the proportion of the ethnicities represented in each consortium study.

Pruning and thresholding PRSs

Both weighted and unweighted PRSs were built using GNU Parallel [59] and PLINK v1.9 (https://www.cog-genomics.org/plink2). 64 different clump P-value cutoffs ranging from 5×10^{-8} to 1 were tested in order to identify the optimal index variant's significance threshold. All other parameters were set to default. A diagram of the different P+T PRSs built is show in Supplementary Figure 2.

LDpred

LDpred PRSs were derived using the LDpred software v0.9.9 (https://github.com/bvilhjal/ldpred) [47]. The fractions of causal variants assumed a prior were similar to the P value thresholds used for the P+T PRSs. Since the number of SNPs was different between the PRSs, The LD radius was adjusted accordingly in each model using the recommended formula (N SNP/3000). All other parameters were kept on their default setting. A diagram of the different P+T PRSs built is shown in Supplementary Figure 2.

GraBLD

GraBLD PRSs were built using several functions available in the GraBLD R package (https://github.com/GMELab/GraBLD) [48]. Data of all the women participating in the START study were used for the calibration. All parameters were set to default.

Association analysis

The association of each PRS with GDM was assessed using a univariate logistic regression model, and areas under the receiver-operating characteristic (ROC) curves (AUCs, c-statistics) were compared in order to determine the PRS with the highest predictive value of GDM. Continuous PRSs were also divided into quartiles in order to compare the participants with highest PRS values to the other groups. Statistical significance of the difference between the predictive values of two PRSs was tested using the DeLong's test for two correlated ROC curves. Analyses were performed using GNU Parallel [59] and R v3.3 [60].

Heritability

The proportion of the variance of GDM explained by: 1) genome-wide genotypes (minor allele count ≥ 10); 2) common variants (MAF ≥ 0.01); 3) SNPs included in our top P+T PRS; was estimated in unrelated participants (identity by state (IBS) < 0.05) by using the GCTA software [61-64]. Single component GREML models were tested. Because of the large number of UK Biobank white British participants, heritability of GDM was estimated in subgroup of participants selected for inclusion in a case-cohort study (N = 623 cases of GDM and 9,037 controls). Hence, reported values for this study were adjusted for a disease prevalence of 0.4%, as estimated in all of the white British women with GDM data in UK Biobank (Table 1). Heritability was not estimated in South Asian women from UK biobank given the small number of GDM cases in this group. All models were adjusted for the first 3 PCA axes.

ACKNOWLEDGEMENTS

The authors thank the UK Biobank for making their data available. The South Asian Birth Cohort (START) study data were collected as part of a bilateral program funded by the Indian Council of Medical Research / Canadian Institutes of Health Research (Project grant number: 298104, Foundation Scheme grant number: FDN-143255, Study grant numbers: INC 109205). This study was also supported by Heart and Stroke Foundation of Canada (grant NA7283. Sonia Anand is supported by a Tier 1 Canada Research Chair in Ethnic Diversity and Cardiovascular Disease, and a Heart and Stroke Foundation/Michael G. DeGroote Chair in Population Health Research at McMaster University. Guillaume Paré is supported by a Canada Research Chair.

AUTHOR CONTRIBUTIONS

A.L performed all the analysis, wrote the main manuscript and prepared all figures and tables. S.M. provided guidance for the GRABLD analysis. D.D is the study coordinator for the START birth cohort, and she provided comments on the manuscript. M.G is a START co-principal investigator and he provided comments on the manuscript. G.P provided guidance for all analyses and reviewed the manuscript. S.A is a START co-principal investigator. She oversaw the development, analysis, and writing of this manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1   Melchior H, Kurch-Bek D & M., M. The Prevalence of Gestational Diabetes: A Population-
Based Analysis of a Nationwide Screening Program. *Dtsch Aerzteblatt Int.* (2017).

2   Farrar, D. *et al.* Hyperglycaemia and risk of adverse perinatal outcomes: systematic review
and meta-analysis. *BMJ* **354**, i4694, doi:10.1136/bmj.i4694 (2016).

3   International Diabetes Federation. IDF Diabetes Atlas, 8th edn. Brussels, Belgium:
International Diabetes Federation. . (2017).

4   Morris, A. P. *et al.* Large-scale association analysis provides insights into the genetic
architecture and pathophysiology of type 2 diabetes. *Nat Genet* **44**, 981-990,
doi:10.1038/ng.2383 (2012).

5   Mahajan, A. *et al.* Genome-wide trans-ancestry meta-analysis provides insight into the genetic
architecture of type 2 diabetes susceptibility. *Nat Genet* **46**, 234-244, doi:10.1038/ng.2897
(2014).

6   Scott, R. A. *et al.* An Expanded Genome-Wide Association Study of Type 2 Diabetes in
Europeans. *Diabetes* **66**, 2888-2902, doi:10.2337/db16-1253 (2017).

7   Wheeler, E. *et al.* Impact of common genetic determinants of Hemoglobin A1c on type 2
diabetes risk and diagnosis in ancestrally diverse populations: A transethnic genome-wide
meta-analysis. *PLoS Med* **14**, e1002383, doi:10.1371/journal.pmed.1002383 (2017).

8   Prokopenko, I. *et al.* A central role for GRB10 in regulation of islet function in man. *PLoS
Genet* **10**, e1004235, doi:10.1371/journal.pgen.1004235 (2014).

9   Manning, A. K. *et al.* A genome-wide approach accounting for body mass index identifies
genetic variants influencing fasting glycemic traits and insulin resistance. *Nat Genet* **44**, 659-
669, doi:10.1038/ng.2274 (2012).

10   Strawbridge, R. J. *et al.* Genome-wide association identifies nine common variants associated
with fasting proinsulin levels and provides new insights into the pathophysiology of type 2
diabetes. *Diabetes* **60**, 2624-2634, doi:10.2337/db11-0415 (2011).

11   Dupuis, J. *et al.* New genetic loci implicated in fasting glucose homeostasis and their impact
on type 2 diabetes risk. *Nat Genet* **42**, 105-116, doi:10.1038/ng.520 (2010).

12   Saxena, R. *et al.* Genetic variation in GIPR influences the glucose and insulin responses to an
oral glucose challenge. *Nat Genet* **42**, 142-148, doi:10.1038/ng.521 (2010).

13   Soranzo, N. *et al.* Common variants at 10 genomic loci influence hemoglobin A(1)(C) levels
via glycemic and nonglycemic pathways. *Diabetes* **59**, 3229-3239, doi:10.2337/db10-0502
(2010).

14   Kwak, S. H. *et al.* A genome-wide association study of gestational diabetes mellitus in Korean
women. *Diabetes* **61**, 531-541, doi:10.2337/db11-1034 (2012).

15   Hayes, M. G. *et al.* Identification of HKDC1 and BACE2 as genes influencing glycemic traits
during pregnancy through genome-wide association studies. *Diabetes* **62**, 3282-3291,
doi:10.2337/db12-1692 (2013).

16   Tarnowski, M. *et al.* GCK, GCKR, FADS1, DGKB/TMEM195 and CDKAL1 Gene
Polymorphisms in Women with Gestational Diabetes. *Can J Diabetes* **41**, 372-379,
doi:10.1016/j.jcjd.2016.11.009 (2017).

17   Anghebem-Oliveira, M. I. *et al.* Type 2 diabetes-associated genetic variants of FTO, LEPR,
PPARg, and TCF7L2 in gestational diabetes in a Brazilian population. *Arch Endocrinol Metab*
61, 238-248, doi:10.1590/2359-3997000000258 (2017).

429	18	de Melo, S. F. *et al.* Polymorphisms in FTO and TCF7L2 genes of Euro-Brazilian women
430		with gestational diabetes. *Clin Biochem* **48**, 1064-1067,
431		doi:10.1016/j.clinbiochem.2015.06.013 (2015).

19   Kasuga, Y. *et al.* Association of common polymorphisms with gestational diabetes mellitus in
Japanese women: A case-control study. *Endocr J* **64**, 463-475, doi:10.1507/endocrj.EJ16-0431
(2017).

20   Kanthimathi, S. *et al.* Association of recently identified type 2 diabetes gene variants with
Gestational Diabetes in Asian Indian population. *Mol Genet Genomics* **292**, 585-591,
doi:10.1007/s00438-017-1292-6 (2017).

21   Tarnowski, M., Malinowski, D., Safranow, K., Dziedziejko, V. & Pawlik, A.
CDC123/CAMK1D gene rs12779790 polymorphism and rs10811661 polymorphism upstream
of the CDKN2A/2B gene in women with gestational diabetes. *J Perinatol* **37**, 345-348,
doi:10.1038/jp.2016.249 (2017).

22   Wang, X. *et al.* Association study of the miRNA-binding site polymorphisms of CDKN2A/B
genes with gestational diabetes mellitus susceptibility. *Acta Diabetol* **52**, 951-958,
doi:10.1007/s00592-015-0768-2 (2015).

23   Wang, Y. *et al.* Association of six single nucleotide polymorphisms with gestational diabetes
mellitus in a Chinese population. *PLoS One* **6**, e26953, doi:10.1371/journal.pone.0026953
(2011).

24   Lauenborg, J. *et al.* Common type 2 diabetes risk gene variants associate with gestational
diabetes. *J Clin Endocrinol Metab* **94**, 145-150, doi:10.1210/jc.2008-1336 (2009).

25   Fatima, S. S., Chaudhry, B., Khan, T. A. & Farooq, S. KCNQ1 rs2237895 polymorphism is
associated with Gestational Diabetes in Pakistani Women. *Pak J Med Sci* **32**, 1380-1385,
doi:10.12669/pjms.326.11052 (2016).

26   Kanthimathi, S. *et al.* Hexokinase Domain Containing 1 (HKDC1) Gene Variants and their
Association with Gestational Diabetes Mellitus in a South Indian Population. *Ann Hum Genet*
80, 241-245, doi:10.1111/ahg.12155 (2016).

27   Al-Hakeem, M. M. Implication of SH2B1 gene polymorphism studies in gestational diabetes
mellitus in Saudi pregnant women. *Saudi J Biol Sci* **21**, 610-615,
doi:10.1016/j.sjbs.2014.07.007 (2014).

28   Kwak, S. H. *et al.* Polymorphisms in KCNQ1 are associated with gestational diabetes in a
Korean population. *Horm Res Paediatr* **74**, 333-338, doi:10.1159/000313918 (2010).

29   Shin, H. D. *et al.* Association of KCNQ1 polymorphisms with the gestational diabetes
mellitus in Korean women. *J Clin Endocrinol Metab* **95**, 445-449, doi:10.1210/jc.2009-1393
(2010).

30   Cho, Y. M. *et al.* Type 2 diabetes-associated genetic variants discovered in the recent genome-
wide association studies are related to gestational diabetes mellitus in the Korean population.
Diabetologia **52**, 253-261, doi:10.1007/s00125-008-1196-4 (2009).

31   Reyes-Lopez, R., Perez-Luque, E. & Malacara, J. M. Metabolic, hormonal characteristics and
genetic variants of TCF7L2 associated with development of gestational diabetes mellitus in
Mexican women. *Diabetes Metab Res Rev* **30**, 701-706, doi:10.1002/dmrr.2538 (2014).

32   Lin, P. C., Chou, P. L. & Wung, S. F. Geographic diversity in genotype frequencies and meta-
analysis of the association between rs1801282 polymorphisms and gestational diabetes
mellitus. *Diabetes Res Clin Pract* **143**, 15-23, doi:10.1016/j.diabres.2018.05.050 (2018).

33   Ding, M. *et al.* Genetic variants of gestational diabetes mellitus: a study of 112 SNPs among
8722 women in two independent populations. *Diabetologia* **61**, 1758-1768,
doi:10.1007/s00125-018-4637-8 (2018).

34 Ekelund, M. *et al.* Genetic prediction of postpartum diabetes in women with gestational
diabetes mellitus. *Diabetes Res Clin Pract* **97**, 394-398, doi:10.1016/j.diabres.2012.04.020
(2012).

35 Frigeri, H. R. *et al.* The polymorphism rs2268574 in Glucokinase gene is associated with
gestational Diabetes mellitus. *Clin Biochem* **47**, 499-500,
doi:10.1016/j.clinbiochem.2014.01.024 (2014).

36 Pagan, A. *et al.* A gene variant in the transcription factor 7-like 2 (TCF7L2) is associated with
an increased risk of gestational diabetes mellitus. *Eur J Obstet Gynecol Reprod Biol* **180**, 77-
82, doi:10.1016/j.ejogrb.2014.06.024 (2014).

37 Shaat, N. *et al.* A variant in the transcription factor 7-like 2 (TCF7L2) gene is associated with
an increased risk of gestational diabetes mellitus. *Diabetologia* **50**, 972-979,
doi:10.1007/s00125-007-0623-2 (2007).

38 Watanabe, R. M. *et al.* Transcription factor 7-like 2 (TCF7L2) is associated with gestational
diabetes mellitus and interacts with adiposity to alter insulin secretion in Mexican Americans.
Diabetes **56**, 1481-1485, doi:10.2337/db06-1682 (2007).

39 Papadopoulou, A. *et al.* Gestational diabetes mellitus is associated with TCF7L2 gene
polymorphisms independent of HLA-DQB1*0602 genotypes and islet cell autoantibodies.
Diabet Med **28**, 1018-1027, doi:10.1111/j.1464-5491.2011.03359.x (2011).

40 Kawai, V. K. *et al.* A genetic risk score that includes common type 2 diabetes risk variants is
associated with gestational diabetes. *Clin Endocrinol (Oxf)* **87**, 149-155,
doi:10.1111/cen.13356 (2017).

41 Kwak, S. H. *et al.* Prediction of type 2 diabetes in women with a history of gestational
diabetes using a genetic risk score. *Diabetologia* **56**, 2556-2563, doi:10.1007/s00125-013-
3059-x (2013).

42 Cormier, H. *et al.* An explained variance-based genetic risk score associated with gestational
diabetes antecedent and with progression to pre-diabetes and type 2 diabetes: a cohort study.
BJOG **122**, 411-419, doi:10.1111/1471-0528.12937 (2015).

43 Ripatti, S. *et al.* A multilocus genetic risk score for coronary heart disease: case-control and
prospective cohort analyses. *Lancet* **376**, 1393-1400, doi:10.1016/S0140-6736(10)61267-6
(2010).

44 Abraham, G. *et al.* Genomic prediction of coronary heart disease. *Eur Heart J* **37**, 3267-3278,
doi:10.1093/eurheartj/ehw450 (2016).

45 Khera, A. V. *et al.* Genome-wide polygenic scores for common diseases identify individuals
with risk equivalent to monogenic mutations. *Nat Genet* **50**, 1219-1224, doi:10.1038/s41588-
018-0183-z (2018).

46 International Schizophrenia, C. *et al.* Common polygenic variation contributes to risk of
schizophrenia and bipolar disorder. *Nature* **460**, 748-752, doi:10.1038/nature08185 (2009).

47 Vilhjalmsson, B. J. *et al.* Modeling Linkage Disequilibrium Increases Accuracy of Polygenic
Risk Scores. *Am J Hum Genet* **97**, 576-592, doi:10.1016/j.ajhg.2015.09.001 (2015).

48 Pare, G., Mao, S. & Deng, W. Q. A machine-learning heuristic to improve gene score
prediction of polygenic traits. *Sci Rep* **7**, 12665, doi:10.1038/s41598-017-13056-1 (2017).

49 Consortium, T. G. P. *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74,
doi:10.1038/nature15393 (2015).

50 Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide
range of complex diseases of middle and old age. *PLoS Med* **12**, e1001779,
doi:10.1371/journal.pmed.1001779 (2015).

51 Speed, D. *et al.* Reevaluation of SNP heritability in complex human traits. *Nat Genet* **49**, 986-
992, doi:10.1038/ng.3865 (2017).

52 Anand, S. S. *et al.* Rationale and design of South Asian Birth Cohort (START): a Canada-
India collaborative study. *BMC Public Health* **13**, 79, doi:10.1186/1471-2458-13-79 (2013).

53 Anand, S. S. *et al.* Causes and consequences of gestational diabetes in South Asians living in
Canada: results from a prospective cohort study. *CMAJ Open* **5**, E604-E611,
doi:10.9778/cmajo.20170027 (2017).

54 Farrar, D. *et al.* Association between hyperglycaemia and adverse perinatal outcomes in south
Asian and white British women: analysis of data from the Born in Bradford cohort. *Lancet*
Diabetes Endocrinol **3**, 795-804, doi:10.1016/S2213-8587(15)00255-7 (2015).

55 Anderson, C. A. *et al.* Data quality control in genetic case-control association studies. *Nat*
Protoc **5**, 1564-1573, doi:10.1038/nprot.2010.116 (2010).

56 Delaneau, O., Marchini, J., Genomes Project, C. & Genomes Project, C. Integrating sequence
and array data to create an improved 1000 Genomes Project haplotype reference panel. *Nat*
Commun **5**, 3934, doi:10.1038/ncomms4934 (2014).

57 Howie, B. N., Donnelly, P. & Marchini, J. A flexible and accurate genotype imputation
method for the next generation of genome-wide association studies. *PLoS Genet* **5**, e1000529,
doi:10.1371/journal.pgen.1000529 (2009).

58 Bycroft, C. *et al.* Genome-wide genetic data on ~500,000 UK Biobank participants. *BioRxiv*
(2017).

59 Tange, O. GNU Parallel - The Command-Line Power Tool. *;login: The USENIX Magazine*
36, 42-47 (2011).

60 R: A language and environment for statistical computing v. 3.3 (R Foundation for Statistical
Computing, Vienna, Austria, 2016).

61 Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide
complex trait analysis. *Am J Hum Genet* **88**, 76-82, doi:10.1016/j.ajhg.2010.11.011 (2011).

62 Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human height.
Nat Genet **42**, 565-569, doi:10.1038/ng.608 (2010).

63 Yang, J. *et al.* Genetic variance estimation with imputed variants finds negligible missing
heritability for human height and body mass index. *Nat Genet* **47**, 1114-1120,
doi:10.1038/ng.3390 (2015).

64 Evans, L. M. *et al.* Comparison of methods that use whole genome data to estimate the
heritability and genetic architecture of complex traits. *Nat Genet* **50**, 737-745,
doi:10.1038/s41588-018-0108-x (2018).

	START[*]	UK Biobank – SAW	UK Biobank – WBW
Number of Participants with GDM data	832	2,386	184,869
GDM, n (%)	301 (36.2%)	52 (2.2 %)	627 (0.43 %)
Age, years	30.2 (4.0)	53.0 (8.1)[‡]	57.6 (7.8)[‡]
Height, cm	162.3 (6.2)[¥]	156.8 (5.9)[‡]	162.5 (6.1)[‡]
Weight, kg	62.6 (12.0) [¥]	67.7 (12.5)[‡]	71.0 (13.2)[‡]
BMI, kg/m^2	23.8 (4.4)	27.5 (4.9) [‡]	26.9 (4.9)[‡]
Family history of diabetes, n (%)	334 (40.2)	1,556 (49.1)	25656 (17.57)

Table 1: Characteristics of women participants from the START and UK Biobank studies with available GDM and Genotype data. [*]South Asian Women with GDM status. Data are mean (SD) unless otherwise indicated. [¥] pre-pregnancy values, [‡] Values from baseline data. Abbreviations: BMI, Body mass index; GDM, gestational diabetes; SAW, South Asian women, WBW, white British women; START, South Asian Birth Cohort.

PRS Type	Consortium	LD source	START - SAW				UK Biobank - SAW				UK Biobank - WBW			
			Beta	SE	P-value	AUC	Beta	SE	P-value	AUC	Beta	SE	P-value	AUC
P+T	**Mahajan et al, 2014**	**1KG**	**0.445**	**0.08**	$\mathbf{8.7 \times 10^{-9}}$	**0.62**	**0.423**	**0.14**	**0.003**	**0.61**	**0.303**	**0.04**	$\mathbf{4.63 \times 10^{-14}}$	**0.58**
		START	0.448	0.08	6.55×10^{-9}	0.62	0.512	0.14	0.0003	0.64	0.291	0.04	3.84×10^{-13}	0.58
	Scott et al, 2017	1KG	0.370	0.07	7.86×10^{-7}	0.60	0.280	0.14	0.05	0.57	0.296	0.04	1.45×10^{-13}	0.58
		START	0.351	0.07	2.98×10^{-6}	0.60	0.300	0.14	0.03	0.59	0.270	0.04	1.88×10^{-11}	0.57
GraBLD	**Mahajan et al, 2014**	1KG	0.465	0.08	1.8×10^{-9}	0.62	0.520	0.14	0.0003	0.64	0.343	0.04	1.14×10^{-17}	0.59
		START	**0.470**	**0.08**	$\mathbf{1.32 \times 10^{-9}}$	**0.62**	**0.510**	**0.14**	**0.0005**	**0.63**	**0.351**	**0.04**	$\mathbf{1.86 \times 10^{-18}}$	**0.59**
	Scott et al, 2017	1KG	0.317	0.07	1.61×10^{-5}	0.59	0.388	0.14	0.006	0.61	0.344	0.04	8.08×10^{-18}	0.59
		START	0.342	0.07	3.93×10^{-6}	0.59	0.387	0.14	0.006	0.61	0.348	0.04	4.15×10^{-18}	0.59
LDpred	**Mahajan et al, 2014**	**1KG**	**0.461**	**0.07**	$\mathbf{2.18 \times 10^{-9}}$	**0.62**	**0.527**	**0.14**	**0.0002**	**0.65**	**0.305**	**0.04**	$\mathbf{2.63 \times 10^{14}}$	**0.58**
		START	0.470	0.08	1.32×10^{-9}	0.62	0.44	0.14	0.002	0.61	0.278	0.04	3.32×10^{-2}	0.57
	Scott et al, 2017	1KG	0.347	0.07	4.05×10^{-6}	0.59	0.382	0.14	0.006	0.61	0.377	0.04	1.80×10^{-21}	0.60
		START	0.281	0.07	0.00015	0.57	0.225	0.13	0.10	0.55	0.323	0.04	1.90×10^{-16}	0.59

Table 2: Characteristics and GDM association results of the top weighted P+T and GraBLD PRSs in South Asian women from the START and UK Biobank studies. Results are from univariate association tests with GDM. The top 3 PRSs are shown in bold. Abbreviations: 1KG, 1000Genomes; AUC area under the curve; GraBLD, Gradient Boosted and LD adjusted; LD, Linkage Disequilibrium; NA, Non applicable; P+T, pruning and thresholding; PRS, Polygenic Risk Score; SAW, South Asian Women; SE, Standard Error; START, South Asian Birth Cohort; WBW, White British Women.

			START – SAW			UK Biobank – SAW			UK Biobank – WBW		
High PRS definition	Reference group	PRS type	OR	95% CI	P value	OR	95% CI	P value	OR	95% CI	P value
Top 25%	Remaining 75%	GraBLD	2.51	1.82 - 3.47	1.75×10^{-8}	2.66	1.51 - 4.63	0.0006	1.75	1.48 – 2.05	2.07×10^{-11}
		P+T	2.08	1.51 – 2.87	7.44×10^{-6}	1.80	0.99 – 3.17	0.05	1.65	1.40 – 1.94	2.41×10^{-09}
		LDpred	2.00	1.45 – 2.76	2.11×10^{-5}	2.61	1-16 – 3.60	0.01	1.72	1.46 – 2.02	7.66×10^{-11}
Top 25%	Lowest 25%	GraBLD	3.40	2.25 - 5.17	7.30×10^{-9}	5.30	2.17 - 15.88	0.0008	2.11	1.69 – 2.66	1.09×10^{-10}
		P+T	3.09	2.10 – 4.74	1.47×10^{-7}	4.21	1.67 – 12.82	0.005	2.22	1.76 - 2.82	3.08×10^{-11}
		LDpred	3.06	2.02 – 4.69	1.77×10^{-7}	3.59	1.53 – 9.84	0.006	2.08	1.66 – 2.62	3.09×10^{-10}

Table 3: Association results of Top PRSs (categories) with GDM in South Asian women from the START and UK Biobank studies.

GraBLD PRS used: $GraBLD\_PRS_1\_W_{Mahajan}\_LD_{START}\_N_{90\%}$; P+T PRS used: $PT\_PRS_1\_W_{Mahajan}\_LD_{1KG}\_N_{85\%min}\_P_{0.016}$; LDpred PRS used: $LDpred\_PRS_1\_W_{Mahajan}\_LD_{1KG}\_N_{85\%}$. Abbreviations: CI, confidence interval; PRS, Polygenic Risk Score; GraBLD, Gradient Boosted and LD adjusted; OR, Odds ratio; P+T, Pruning and thresholding; SAW, South Asian Women; START, South Asian Birth Cohort; WBW, White British Women.

	Allele count / MAF cutoff	START - SAW		UK Biobank - WBW *	
		h^2	SE	h^2	SE
All SNPs in the study	MAC 10	0.57	0.43	0.15	0.11
	MAF 0.01	0.50	0.40	0.11	0.07
SNPs in top P+T PRS	NA	0.18	0.14	0.03	0.03

Table 4: Heritability estimates for GDM in women from the START and UK Biobank studies. Models are adjusted for the first 3 eigenvectors from the principal component analysis. * heritability estimates on the liability scale (disease prevalence = 0.4%) from the case-cohort study. Abbreviations: MAC, Minor allele count; MAF, Minor allele frequency; NA, Not applicable; P+T, Pruning and thresholding; SAW, South Asian women; SE, Standard Error; SNP, Single Nucleotide Polymorphism; START, South Asian Birth Cohort; WBW, White British Women.

Figure 1: AUCs of the different weighted and unweighted LD$_{START}$ P+T PRSs based on Mahajan *et al.* and Scott *et al. Results from association tests with GDM. Abbreviations: 0 85 90 95 and 98%, PRSs including a subset of SNPs tested in at least 0 85 90 95 and 98% of the total samples of the consortium study respectively; AUC area under the curve; PRS, Polygenic Risk Score; LD, Linkage disequilibrium; P+T, Pruning and thresholding; SNP, Single Nucleotide Polymorphism; START, South Asian Birth Cohort; ROC, receiver operating characteristic; uPRS, unweighted PRSs; wPRS, weighted PRSs.

Figure 2: AUCs of the different LDpred PRSs based on Mahajan *et al.* and Scott *et al.* Results from association tests with GDM. Abbreviations: 0 85 90 95 and 98%, PRSs including a subset of SNPs tested in at least 0 85 90 95 and 98% of the total samples of the consortium study respectively; 1KG, 1000 Genomes; AUC area under the curve; PRS, Polygenic Risk Score; LD, Linkage disequilibrium; SNP, Single Nucleotide Polymorphism; ROC, receiver operating characteristic; wPRS, weighted PRSs.

Figure 3: AUCs of the best Mahajan *et al.* P+T and LDpred PRSs in START.
Abbreviations: 85 %, PRSs including a subset of SNPs tested in at least 85 % of the total
samples of the consortium study respectively; 1KG, 1000 Genomes; AUC area under the curve;
PRS, Polygenic Risk Score; LD, Linkage disequilibrium; P+T, Pruning and thresholding; START,
South Asian Birth Cohort; ROC, receiver operating characteristic; wPRS, weighted PRSs

Figure 4: AUCs of the different weighted P+T and LDpred PRSs based on Mahajan *et al*. and Scott *et al*. Results from association tests with GDM. Abbreviations: 0 85 90 95 and 98%, PRSs including a subset of SNPs tested in at least 0 85 90 95 and 98% of the total samples of the consortium study respectively; 1KG, 1000 Genomes; AUC area under the curve; PRS, Genome-wide Polygenic Risk Score; LD, Linkage disequilibrium; P+T, Pruning and thresholding; SNP, Single Nucleotide Polymorphism; START, South Asian Birth Cohort; ROC, receiver operating characteristic; uPRS, unweighted PRSs; wPRS, weighted PRSs.

Figure 5: AUCs of Mahajan *et al.* N$_{85\%}$ based LD$_{START}$ P+T and LDpred PRS and their LD$_{1KG}$ counterparts. Abbreviations: 85 and 95%, PRSs including a subset of SNPs tested in at least 85 and 95% of the total samples of the consortium study respectively; 1KG, 1000 Genomes; AUC area under the curve; PRS, Polygenic Risk Score; LD, Linkage disequilibrium; P+T, Pruning and thresholding; START, South Asian Birth Cohort; ROC, receiver operating characteristic; wPRS, weighted PRSs.