1

Forward modelling reveals dynamics of neural orientation tuning to unconscious visual stimuli during binocular rivalry

Matthew F. Tang[1,2], Ehsan Arabzadeh[2,3] & Jason B. Mattingley[1,2,4,5]

1. Queensland Brain Institute, The University of Queensland, St Lucia, QLD, Australia

2. Australian Research Council Centre of Excellence for Integrative Brain Function, Victoria, Australia

3. Eccles Institute of Neuroscience, John Curtin School of Medical Research, The Australian National University, Canberra, ACT, Australia

4. School of Psychology, The University of Queensland, St Lucia, QLD, Australia

5. Canadian Institute for Advanced Research (CIFAR)

Corresponding author

Matthew F. Tang

Queensland Brain Institute

The University of Queensland

St Lucia, QLD, Australia

Email: m.tang1@uq.edu.au

23   **Abstract**

When different visual stimuli are presented to the two eyes, they typically compete for

access to conscious perception, a phenomenon known as *binocular rivalry*. Previous

studies of binocular rivalry have shown that neural responses to consciously

suppressed stimuli are markedly diminished in magnitude, though they may still be

encoded to some extent. Here we employed multivariate forward modelling of human

electroencephalography (EEG) data to quantify orientation-selective responses to

visual gratings during binocular rivalry. We found robust orientation tuning to both

conscious and unconscious gratings. This tuning was enhanced for the suppressed

stimulus well before it was available for conscious report. The same pattern was

evident in the overall magnitude of neural responses, and it emerged even earlier than

the changes in neural tuning. Taken together, our findings suggest that rivalry

suppression affects broadband, non-orientation selective aspects of neural activity

before refining fine-grained feature-selective information.

There is considerable evidence that the human brain can encode sensory events

that elude conscious detection, either following acquired lesions of the sensory or

association areas[1,2], or due to masking[3] or attentional manipulations of relevant

stimuli[4,5]. A particularly striking example occurs during binocular rivalry, in which the

presentation of different images to the two eyes leads to alternating periods during

which the image to one eye is clearly perceived while the other is suppressed from

awareness. One reason for the continued interest in binocular rivalry is that it provides

a window onto the neural bases of consciousness[6-8] by allowing investigators to

compare patterns of brain activity elicited by pairs of stimuli that are held constant

while conscious perceptual reports alternate stochastically over time. Although

previous research has shown that neural responses to suppressed stimuli are reduced

in overall amplitude, no study to date has determined how the basic features that

define such stimuli, such as their colour or orientation, are affected under conditions of

binocular suppression. Here we addressed this question directly by recording neural

activity with electroencephalography (EEG) while observers viewed rivalrous displays in

which differently coloured and oriented gratings were projected to each eye. We used

frequency tagging to track neural activity to each grating over time, and applied

multivariate forward encoding modelling to recover orientation tuning profiles for the

two stimuli as they alternated between conscious and unconscious perception.

A persistent question in the literature on binocular rivalry concerns the extent to

which the brain represents stimuli that are currently suppressed from awareness.

Invasive recordings in macaques[9,10] have shown that neurons in early cortical areas

continue to respond to stimuli that are under binocular suppression. For some neurons,

firing rates appear to track the animal's perceptual reports, becoming lower when the

stimulus is suppressed and higher when it is perceived. Likewise, human neuroimaging

studies[6,11-15] have shown that the overall magnitude of neural responses in early cortical

areas is modulated according to observers' subjective perceptual report. For example,

studies using EEG to track the time course of neural activity during binocular rivalry

have revealed that response profiles to perceived and suppressed stimuli are anti-

correlated, and that perceived stimuli elicit a larger neural response overall[13,14,16,17].

Studies using functional magnetic resonance imaging (fMRI) have localised these

effects to relatively early cortical areas[6,11,18]. In higher areas, activity induced by

suppressed inputs may still be present, if for example the stimuli carry biologically

salient information (e.g., fearful faces[19,20]), while it can be completely absent in other

brain regions[6,19] This work suggests that any neural activity elicited by the suppressed

stimulus is simply too weak to cross the threshold for conscious report.

All previous studies in humans have focused exclusively on the overall *level* of

neural activity during binocular rivalry, rather than on the time-varying patterns of

activity associated with changes in the features of suppressed visual inputs. However,

other aspects of the stimulus' representation, in addition to the overall magnitude of

response, are known to be critical for conscious detection of targets[21]. Here we asked

whether orientation selectivity, which can be defined along several dimensions,

including the amplitude or gain of stimulus evoked activity as well as the fidelity or

sharpness of neural tuning, is modulated for a fixed stimulus as it alternates between

periods of conscious perception and suppression. Behavioural work in humans has

suggested that stimulus selectivity is affected during rivalry[7,22], but no study to date has

examined how rivalry affects neural representations of specific stimulus features.

To address this question we used multivariate forward modelling[23-26] of human

electroencephalography (EEG) data to quantify orientation-selective responses to

visual gratings that were undergoing changes in awareness during binocular rivalry. To

anticipate the results, we observed robust orientation tuning to both conscious and

unconscious gratings. Strikingly, orientation tuning began to increase for the currently

suppressed stimulus more than 300 ms before observers were aware of it, while at the

same time orientation tuning began to decline for the currently perceived grating that

was about to be suppressed from awareness. The same pattern was evident for the

overall magnitude of neural responses, and this effect emerged even earlier than the

changes in neural tuning (~800 ms before observers' button press indicating a switch

in percept). Taken together, our findings suggest that rivalry suppression first affects

the broadband, and non-orientation selective aspects of the neural information, before

refining the fine-grained feature selective information.

Results

To quantify the time-course and extent of neural information processing of

conscious and unconscious stimuli, we presented spatially overlapping red and green

gratings dichoptically to the left and right eyes. (Figure 1A; Movie 1). Dichoptic

presentation causes interocular suppression, such that observers perceive a single

coherent image that oscillates stochastically between the two stimuli over time, with

occasional mixed percepts in between (Figure 1B)[27,28]. To measure time-resolved

neural responses to the stimuli, we recorded neural activity using EEG, while the two

orthogonally oriented gratings counter-phase flickered at two different frequencies (20

and 24 Hz). The two gratings generated distinct steady-state visual evoked potentials

(SSVEPs) that were tracked continuously over each 30 s trial. As described in detail

below, the orientations of the two gratings varied from 0° to 160° (in 20° steps) across

the trials to enable modelling of orientation-selective responses as a function of

perceptual awareness. Participants (N = 22) reported whether they perceived the red or

green grating, or a mixed percept, while we measured SSVEP responses. The

magnitude of SSVEPs generated by rivalrous gratings are typically anti-correlated,

exhibiting a larger response to the grating that is consciously perceived than to the

grating that is suppressed from awareness[16].

Figure 1. The SSVEP paradigm used to measure neural activity elicited by conscious
and unconscious grating stimuli during binocular rivalry. **(A)** A red and green grating
were presented on a computer screen while neural activity was recorded using EEG. A
mirror stereoscope and a dividing board ensured that each eye received visual
information exclusively from one of the two gratings, yielding binocular rivalry. **(B)**
Under binocular rivalry observers typically perceive a slowly alternating percept in
which one of the two gratings dominates in perception and the other is suppressed.
Here observers occasionally reported a combination of either colour or orientation
information (i.e., a mixed percept). **(C)** To separately track the brain's response to the
gratings, the two stimuli counter-phase flickered at 20 and 24 Hz, producing unique
frequency tags in the EEG. An illustrative segment from a typical trial of the rivalry task
showing neural responses to the gratings (i.e. the power at the tagged frequencies) as
a function of the participant's percept. **(D)** Mean frequency response for all participants
across the entire 30 s trial (averaged over electrodes Oz, O1, O2, POz). Grey shading
indicates ∓1 standard error of the mean. **(E)** Distribution of SSVEP responses for the
20 and 24 Hz gratings across the scalp for all participants.

Binocular rivalry affects the overall strength of neural responses

The median duration of perceptual dominance, during which one grating was

perceived while the other was suppressed, lasted for 1390 ms (interquartile range =

1024 to 2349 ms; Figure 2A). To determine the effect of binocular rivalry on neural

responses to the two gratings, SSVEP responses were epoched into bins that

extended from 2s before the participant indicated a change in percept to 2 s after this

response. The data were then combined across the flicker frequencies of the two

presented gratings to produce 'aware' and 'unaware' epochs (see Methods for further

details). The label of the response (i.e., aware versus unaware) was defined relative to

the percept reported at the time of the button press (0 ms). As shown in Figure 2B, the

SSVEP elicited by the currently suppressed grating began to increase around 1500 ms

before the stimulus was consciously perceived. At the same time, the SSVEP elicited

by the currently perceived stimulus began to decline sharply after it had been reported,

and the response to the suppressed stimulus began to rise. Thus, around 1000 ms

after the observers' button press, the neural response to the unaware stimulus was

now larger than the response to the stimulus that was currently perceived. This timing

fits with the behavioural observation that periods of stimulus dominance lasted on

average around 1390 ms. As shown in Figure 2C, SSVEPs elicited by the grating

stimuli were most pronounced over occipital-parietal electrode sites, as expected for

such square-wave grating stimuli[6,16].

Figure 2. Behavioural and neural results associated with changes in awareness during binocular rivalry. **(A)** Distribution of durations of perceptual dominance across the 22 participants. **(B)** Time series of relative EEG power elicited by the grating stimuli, plotted separately for aware and unaware periods of the trial, averaged across three occipital electrodes (Oz, POz, Pz). The 'aware' response corresponds to the grating reported as the dominant percept at the time of the button press (0 ms), and the 'unaware' response corresponds to the suppressed stimulus at the same time point. The black horizontal lines indicate periods of significant difference between the conditions (two-tailed cluster-permutation, alpha $p < .05$, cluster alpha $p < .05$, N permutations = 20,000). **(C)** Headmaps of EEG data showing changes in relative power for periods in which gratings were consciously perceived (Aware, top row) and suppressed (Unaware, middle row), and the difference between these states (bottom row). Circles indicate clusters of electrodes with significantly lower activity, and crosses indicate clusters with significantly greater activity in the Aware vs. Unaware conditions (alpha $p < .05$, cluster $p < .025$, N permutations = 1500). **(D)** Accuracy of linear discriminant analysis used to predict observers' grating perception at time 0 ms (corresponding to the participants' button press). Coloured horizontal bars indicate times during which classification accuracy was significantly better than chance for Aware and Unaware epochs. Coloured shading in C and D represents ∓ 1 standard error of the mean.

179
Stimulus-evoked SSVEPs predict perceptual reports prior to awareness

We next sought to determine whether SSVEPs could be used to predict which

percept the participant would report on a single-trial basis. To do this, we used a

multivariate linear discriminant analysis to determine whether there were patterns of

activity distributed across the scalp that predicted which grating was about to be

consciously perceived. Separately, for each SSVEP frequency and at each time point

across the epoch around when the percept changed, the classifier was trained on the

percept reported at time zero, which coincided with the participants' indicated

perceptual report. Again, the results were combined across frequencies to produce

unaware and unaware responses. Neural responses to consciously perceived gratings

predicted observers' perceptual reports around 800 ms before the report occurred

(Figure 2D), whereas neural responses to the unaware gratings did not reliably predict

observers' reports at this same time point. By contrast, at around 1000 ms after

observers made their response, SSVEP power was able to predict observers'

perceptual reports for both the aware and unaware stimuli.

Binocular rivalry affects visual orientation selectivity

Next, we sought to answer the critical question of whether orientation selectivity

for the rivalrous gratings was influenced by observers' awareness of the frequency-

tagged stimulus. To do this, we applied forward encoding modelling, which has

previously been used to recover orientation selectivity from both EEG and MEG

activity[23-26,29]. Broadly, this technique identifies multivariate patterns of EEG activity that

are selective for defined stimulus features, in this case orientation (see Figure 3 for an

example). We convolved grating orientation in each epoch against a bank of nine

canonical, orientation-selective tuning functions, each maximally sensitive to a different

orientation, to produce regression coefficients for that trial (Figure 3B). Then at each

time point in the epoch, multivariate regression was used to estimate the spatial map

(or *weight matrix*) sensitive to orientation in a training set of data. This weight matrix

(Figure 3C) was inverted and multiplied against a test set of data (Figure 3D) to

determine orientation selectivity (Figure 3E) in the test data. The procedure was

iterated over all time points in the epoch, and a cross-validation procedure was used

such that all epochs served in both training and test data.

Figure 3. Overview of the forward encoding modelling procedure used to measure orientation selectivity in the binocular rivalry paradigm. **(A)** Patterns of SSVEP activity (here separated into real and imaginary components) are measured in response to a range of orientations. **(B)** Regression coefficients for the orientations are estimated for each trial from orientation selective functions. **(C)** Least-squares regression is used to estimate weights for all EEG electrodes for each of the channels in the orientation model. The channel weights are shown here for three example occipital-parietal electrodes. **(D)** The channel weights are inverted and multiplied against a set of independent training data. **(E)** The model will reconstruct the presented orientation if there is an orientation selective response. In this case, the stimulus reconstruction shows a tuned response to a presented grating of 20° orientation. **(F)** Average orientation reconstructions over time for the two SSVEP frequencies (collapsed across aware and unaware conditions) for all participants for each of the nine presented orientations, indicated by the black dotted line. These reconstructions were realigned so we could combine the representations of the different orientations.

Figure 3F shows the results of the forward encoding modelling for each of the

presented orientations (20° to 180° in 20° steps) for the two SSVEP frequencies of 20

and 24 Hz. As shown, the forward encoding model successfully recovered the

presented orientation across all time points in the epoch. Because we were not

interested in any specific orientation, but rather in orientation selectivity in general, and

to increase the signal to noise ratio, we re-centred all the representations to the

presented orientation (0° for consistency) in that trial. To quantify orientation selectivity

between different conditions, these outputs were fit with a Gaussian function, with the

amplitude providing an index of orientation selectivity (see Methods for details).

The forward encoding model revealed a strong orientation-selective neural

response that was tuned to the presented orientation (Figure 4A). Strikingly, we found

orientation-selective response to both the currently perceived grating and to the

grating that was suppressed due to binocular rivalry. Critically, this result suggests that

the visual system encodes feature-selective information about the presented stimulus

even when that stimulus is not consciously perceived. Moreover, just prior to the

observers' button press indicating a switch in percept, the grating that was currently

suppressed but was about to be consciously reported was associated with

significantly greater orientation selectivity than the grating that was perceived (but was

about to be suppressed).

Figure 4. Results of forward encoding modelling for orientation selectivity under binocular rivalry. **(A)** Orientation decoding (given by height of tuning curves) over time for SSVEP responses elicited by aware and unaware gratings. The 'aware' response corresponds to the grating reported as the dominant percept at the time of the button press (0 ms), and the 'unaware' response corresponds to the suppressed stimulus at the same time point. The horizontal black bars indicate significant differences between the aware and unaware conditions (two-tailed cluster-permutation, alpha $p < .05$, cluster alpha $p < .05$, N permutations = 20,000). Coloured shadings show ∓ 1 standard error of the mean. **(B)** Orientation tuning curves collapsed across significant time points shown in A. Error bars show ∓ 1 standard error of the mean. The lines are fitted Gaussians (Equation 4) to data averaged across observers.

To further determine how orientation selectivity was affected by awareness, we

averaged the time-resolved tuning curves over the significant time points (black lines in

Figure 4A) to recover the tuning curve, and fitted Gaussian functions to quantify the

change in neural representations (Figure 4B). This again revealed robust orientation-

selective responses to both aware and unaware stimuli, shown by the tuned response

at the presented orientation (0°). The analysis also confirmed that the amplitude of the

orientation-tuning functions was reliably larger for grating stimuli when they were aware

than when they were unaware ($p = .01$), suggesting that conscious perception is

associated with a boost in the gain of orientation selectivity (or a suppression in the

gain for suppressed stimuli). Crucially, however, at the same time points relative to

observers' perceptual reports, there was no change in the width of the tuning functions

($p = .37$), suggesting that awareness does not alter the fidelity or tuning of the

underlying neural representation (aware $M = 33.70$, $SE = 2.97$; unaware $M = 31.58$, SE

$= 1.25$).

274   **Discussion**

We investigated how binocular rivalry affects the neural representation of visual

orientation information in human observers. Previous work in non-human primates has

shown that the selectivity of neurons in visual areas tracks changes in awareness, such

that suppressed stimuli evoke a smaller selective response than dominant (perceived)

stimuli. It has remained unclear whether the same changes in feature-selectivity occur

in humans because conventional neuroimaging approaches focus on overall levels of

activity, rather than tracking the time-varying changes in selectivity for elementary

stimulus features. Here we used forward encoding modelling to determine how

orientation selectivity was influenced under conditions in which orthogonally oriented

pairs of gratings were presented dichoptically to induce binocular rivalry.

We found significant orientation selectivity in neural activity associated with both

aware and unaware gratings, and this response increased just before the reported

change in awareness. Forward encoding modelling allowed us to quantify orientation

tuning curves. These analyses revealed that while the gain (amplitude) of orientation

tuning was affected by rivalry, the fidelity (or sharpness) was not. We also found that

the overall neural response to the gratings, as indexed by the power of the frequency-

tagged SSVEP, began to increase ~1000 ms before orientation-selectivity changed. On

a single trial basis, from the power of the SSVEP response alone, it was possible to

predict an observer's choice ~800 ms before the individual reported the change. As

with most previous investigations of binocular rivalry[6,14,16,30], we used observers' button-

press responses to demarcate changes in awareness, and these perceptual switches

typically precede overt responses by 300 to 400 ms[30]. Such response time delays

cannot account for the current findings, however, as SSVEP power predicted

observers' percepts ~800 ms before their responses. This observation suggests that

during rivalry, alternations in awareness of rivalrous stimuli are initially signalled by

broadband changes which are not selective for orientation in neural activity, followed

by a change in fine-grained stimulus selectivity immediately prior to a change in

conscious report.

Our results are broadly consistent with findings from invasive recordings in

macaque visual cortex during binocular rivalry[10,31]. Around 300 ms before the currently

suppressed stimulus broke into awareness, we found that orientation selectivity for the

grating began to increase. This anticipatory neural response in humans is consistent

with findings from neuronal recordings in macaque area V4, which have shown that the

firing rate of neurons during binocular rivalry increases around 300-500 ms before the

animal reports a change in the perceived stimulus. Our work goes beyond these

findings by showing that, in humans, orientation-selective information is reliably

available to the system even when a rivalrous stimulus is not consciously perceived.

More broadly, our findings are consistent with the influential global workspace

theory of consciousness[32,33], which proposes that while all stimulus information is

represented by early sensory regions, only those that get fed through to higher cortical

regions activate a global brain network and thus reach awareness. On this account,

stimuli reach awareness because they induce high synchrony within a global network

of more local networks. Once an item has activated the global network, it can inhibit

alternative and competing stimuli. Our results potentially add to this account, as we

found that during rivalry broadband neural responses first change with awareness, and

are followed several hundred milliseconds later by more fine-grained (feature-specific)

changes in stimulus representations. One way this might occur is via feedback

connections from higher brain areas, which are predicted to have access to the global

workspace, to lower-level sensory regions which represent stimulus-selective

information such as visual orientation. This idea could be tested by disrupting higher

cortical areas – for example using transcranial magnetic stimulation – and measuring

changes in orientation tuning to rivalrous stimuli. Future studies could also apply the

forward encoding approach used here to assess feature-specific neural responses in

different experimental paradigms in which conscious reports are altered, such as visual

masking[3], or in cases of neural pathologies such as blindsight [2] or unilateral spatial

neglect[1,34].

Methods

Participants

A group of 26 healthy adult observers (age range = 18 to 37 yr) participated in

exchange for partial course credit or financial reimbursement (AUD$20/hr). All

participants provided written informed consent, and all reported normal or corrected-

to-normal vision and normal colour perception. The study was approved by The

University of Queensland Human Research Ethics Committee and was conducted in

accordance with the Declaration of Helsinki.

Experimental setup

340   The experiment was conducted inside a dimly illuminated room with participants

341 seated in a comfortable chair. The stimuli were displayed on a 22-inch LED monitor

342 (resolution 1920 x 1080 pixels, refresh rate 120 Hz) using PsychToolbox presentation

343 software[35,36] for MATLAB (v7.3). Viewing distance was maintained at 54 cm using a

344 chinrest, meaning each pixel subtended 1.8' x 1.8'.

Task

Prior to the start of the experiment, participants aligned the mirror stereoscope

(Fig 1A) so images presented to the left and right eyes appeared overlapping. Identical

circular checkerboards were placed around the stimulus in both eyes to encourage

fusion of the images. Each trial began with presentation of these checkerboards and a

central Gaussian fixation dot. After 1500 ms, green and red square-wave gratings were

presented to the left and right eyes, so that they appeared within the centre of the

surrounding checkerboards. The colours were presented at 100% luminance intensity.

To generate SSVEPs, the gratings counter-phase flickered at 20 and 24 Hz for 30 s.

Counter-phase flickering square-wave gratings were used because these produce

large SSVEP responses and produce orientation-selective neural activity that can be

tracked with EEG [13]. Participants indicated whenever the percept (red, green or

mixed) changed during the trial using separate keys on a standard keyboard. For half

the participants, the red gratings flickered at 20 Hz and the green gratings flickered at

24 Hz; for the remaining participants these contingencies were reversed.

Across trials the orientations of the gratings were varied between 0° and 160° (in

20° steps) so that forward encoding modelling could be employed to determine

orientation selectivity contained with the EEG data. The orientations of the pairs of

gratings presented in each trial were always perpendicular to ensure strong rivalry

between the stimuli[31,37,38]. There were 90 x 30 s trials in total across the experiment,

meaning that each orientation was repeated 10 times.

EEG acquisition and pre-processing

Continuous EEG data were recorded using a BioSemi Active Two system

(BioSemi, Amsterdam, Netherlands). The signal was digitised at 1024 Hz sampling rate

with a 24-bit A/D conversion. The 64 active scalp Ag/AgCl electrodes were arranged

according to the international standard 10–20 system for electrode placement [26]

using a nylon head cap. As per BioSemi system design, the common mode sense and

driven right leg electrodes served as the ground, and all scalp electrodes were

referenced to the common mode sense during recording.

Offline EEG pre-processing was performed using EEGLAB in accordance with

best practice procedures[39,40]. The data were initially down sampled to 256 Hz and

subjected to a 0.5 Hz high-pass filter to remove slow baseline drifts. Electrical line

noise was removed using the *clean_line.m,* and *clean_rawdata.m* in EEGLAB [29] was

used remove bad channels (identified using Artifact Subspace Reconstruction), which

were then interpolated from the neighbouring electrodes. Data were then re-referenced

to the common average before being epoched into trials (-1 to 31 s). Systematic

artefacts from eye blinks, movements and muscle activity were identified and

regressed out of the signal using semi-automated procedures in the SASICA toolbox

[30]. After this stage, any trial with a peak voltage exceeding ±100 *uV* was excluded

from the analysis.

Data analysis

A wavelet transform (150 ms time-domain standard deviation) was applied to

the recorded EEG data in each trial to extract the neural response to the two gratings.

The data were then epoched into 4 s time periods symmetrically around each button

press (-2 s to + 2 s), which observers used to indicate a change in the perceived

grating. Epochs were only included if the preceding epoch was not a mixed percept, so

we were certain the preceding epoch was the previously aware stimulus. We did not

include periods with mixed percepts because these are by definition ambiguous with

respect to stimulus awareness (i.e., both stimuli are perceived to some extent). Epochs

were also only included if the button press occurred between 2 and 28 s into the trial

(thus ensuring a robust frequency tagged signal). To ensure the encoding reflected

periods of stable perception, only dominance periods lasting longer than 500 ms were

included. Two participants were excluded because most responses were between one

colour and mixed. Two further participants were excluded because they reported

significantly fewer switch (<100, SD < 2 SD below average). This produced a mean of

1276 epochs for the remaining (N = 22) participants (range = 445 to 1963 across the

participants). For each participant the periods of dominance were relatively evenly split

between the two colors; there were relatively fewer periods in which the percept was

mixed, and these tended to be shorter-lived in duration (Figure 2A). To maintain

consistent phase information in the SSVEPs, we aligned the nearest counter-phase

flicker within the epoch to the participants' button press, separately for the 20 and 24

Hz frequencies.

Forward encoding modelling

We used forward encoding modelling to extract the orientation selective

response from the patterns of EEG activity. This technique transforms sensor-level

activity distributed across the scalp into tuned 'feature-channels' that are selective for

a specific feature dimension, in this case orientation[23,24,26,29]. To do this, we used the

orientations of the epoched data segments (4 s each) to construct a regression matrix

with 9 regression coefficients for each of the orientations. This regression matrix was

convolved with a tuned set of nine basis functions (half cosine functions raised to the

eighth power) cantered from 0° and 160° (in 20° steps). This tuned regression matrix

was used to measure orientation at each time point in the 4 s epoched segments. This

was done by solving the linear equation (1):

$$B_1 = WC_1 \tag{1}$$

Where B_1 (64 sensors x N training trials) is the electrode data for the training set, C_1 (9

channels x N training trials) is the tuned channel response across the training trials, and

W is the weight matrix for the sensors to be estimated (64 sensors x 9 channels).

Following an approach recently introduced for the analysis of MEG data, we separately

estimated the weights associated with each channel individually [31]. W was estimated

using least square regression to solve equation (2):

$$W = (C_1 C_1^T)^{-1} C_1^T B_1 \qquad (2)$$

Following this previous work[41,42] we also sought to remove the correlation between the

sensors which hinders finding the true solution to the linear equation. To do this, we

first estimated the noise correlation between electrodes and removed this component

through regularisation[43]. The channel response in the test set C_2 (9 channels x N test

trials) was estimated using the weights in (2), and was applied to activity in B_2 (64

sensors x N test trials).

$$C^2 = (W W^T) W^T B^2 \qquad (3)$$

To avoid overfitting, we used a five-fold cross validation procedure in which 80% of

epochs were used to train the model, and the remaining 20% of epochs were used for

testing. This procedure was repeated until all epochs had served as both training and

test trials. We repeated this procedure for both of the SSVEP frequencies and at each

time-point in the epoch to produce a time-resolved orientation-selectivity function. We

then shifted all trials to a common orientation, so that $0°$ corresponded to the

orientation presented on each trial. These reconstructions were smoothed with a

Gaussian temporal kernel to increase signal-to-noise[29].

To quantify orientation selectivity, we fit the results of the forward encoding

model with a Gaussian function (4) using least square regression to quantify the

amount of orientation selective activity.

$$444 \qquad G(x) = A \, exp(-\frac{(x-\phi)^2}{2\sigma^2}) + C \qquad\qquad (4)$$

445 Where *A* is the amplitude representing the amount of orientation selective activity, ϕ is

446 the orientation the function is centred on (in degrees), σ is the width (degrees) and *C* is

447 a constant used to account for non-orientation selective baseline shifts.

Acknowledgements

This work was supported by the Australian Research Council (ARC) Centre of

Excellence for Integrative Brain Function (ARC Centre Grant CE140100007) to JBM

and EA, and by an ARC Discovery Project (DP170100908) to EA. JBM was supported

by ARC Australian Laureate Fellowship (FL110100103) and by the Canadian Institute

for Advanced Research (CIFAR). We would like to thank Morgan McIntyre who helped

with data collection.

References

1.   Mattingley, J. B., Davis, G. & Driver, J. Preattentive filling-in of visual surfaces in
parietal extinction. *Science* **275,** 671–674 (1997).
2.   Cowey, A. & Stoerig, P. Blindsight in monkeys. *Nature* **373,** 247–249 (1995).
3.   Enns, J. T. & Di Lollo, V. What's new in visual masking? *Trends Cogn Sci* **4,** 345–
352 (2000).
4.   Broadbent, D. E. & Broadbent, M. H. P. From detection to identification:
Response to multiple targets in rapid serial visual presentation. *Percept*
Psychophys **42,** 105–113 (1987).
5.   Raymond, J. E., Shapiro, K. L. & Arnell, K. M. Temporary suppression of visual
processing in an RSVP task: an attentional blink? . *Journal of Experimental*
Psychology: Human Perception and Performance **18,** 849–860 (1992).
6.   Tong, F., Nakayama, K., Vaughan, J. T. & Kanwisher, N. Binocular Rivalry and
Visual Awareness in Human Extrastriate Cortex. *Neuron* **21,** 753–759 (1998).
7.   Ling, S. & Blake, R. Suppression During Binocular Rivalry Broadens Orientation
Tuning:. *Psychological Science* **20,** 1348–1355 (2009).
8.   Rees, G., Kreiman, G. & Koch, C. Neural correlates of consciousness in humans.
Nat Rev Neurosci **3,** 261 EP —270 (2002).

9.   Logothetis, N. K. & Schall, J. D. Neuronal correlates of subjective visual
perception. *Science* **245,** 761–763 (1989).
10. Leopold, D. A. & Logothetis, N. K. Activity changes in early visual cortex reflect
monkeys' percepts during binocular rivalry. *Nature* **379,** 379549a0–553 (1996).
11. Polonsky, A., Blake, R., Braun, J. & Heeger, D. J. Neuronal activity in human
primary visual cortex correlates with perception during binocular rivalry. *Nat
Neurosci* **3,** 1153–1159 (2000).
12. Srinivasan, R., Russell, D. P., Edelman, G. M. & Tononi, G. Increased
Synchronization of Neuromagnetic Responses during Conscious Perception. *J
Neurosci* **19,** 5435–5448 (1999).
13. Tononi, G., Srinivasan, R., Russell, D. P. & Edelman, G. M. Investigating neural
correlates of conscious perception by frequency-tagged neuromagnetic
responses. *Proc Natl Acad Sci USA* **95,** 3198–3203 (1998).
14. Brown, R. J. & Norcia, A. M. A method for investigating binocular rivalry in real-
time with the steady-state VEP. *Vis Res* **37,** 2401–2408 (1997).
15. Lansing, R. W. Electroencephalographic Correlates of Binocular Rivalry in Man.
Science **146,** 1325–1327 (1964).
16. Zhang, P., Jamison, K., Engel, S., He, B. & He, S. Binocular Rivalry Requires
Visual Attention. *Neuron* **71,** 362–369 (2011).
17. Haynes, J.-D., Deichmann, R. & Rees, G. Eye-specific effects of binocular rivalry
in the human lateral geniculate nucleus. *Nature* **438,** 496–499 (2005).
18. Tong, F. & Engel, S. A. Interocular rivalry revealed in the human cortical blind-
spot representation. *Nature* **411,** 195–199 (2001).
19. Pasley, B. N., Mayes, L. C. & Schultz, R. T. Subcortical Discrimination of
Unperceived Objects during Binocular Rivalry. *Neuron* **42,** 163–172 (2004).
20. Williams, M. A., Morris, A. P., McGlone, F., Abbott, D. F. & Mattingley, J. B.
Amygdala Responses to Fearful and Happy Facial Expressions under Conditions
of Binocular Suppression. *J Neurosci* **24,** 2898–2904 (2004).
21. Kullback, S. & Leibler, R. A. On Information and Sufficiency. *The Annals of
Mathematical Statistics* **22,** 79–86 (1951).
22. Lunghi, C. & Alais, D. Touch Interacts with Vision during Binocular Rivalry with a
Tight Orientation Tuning. *PLoS ONE* **8,** e58754 (2013).
23. Brouwer, G. J. & Heeger, D. J. Decoding and reconstructing color from
responses in human visual cortex. *J Neurosci* **29,** 13992–14003 (2009).
24. Garcia, J. O., Srinivasan, R. & Serences, J. T. Near-real-time feature-selective
modulations in human cortex. *Current Biology* **23,** 515–522 (2013).
25. Sprague, T. C. & Serences, J. T. Attention modulates spatial priority maps in the
human occipital, parietal and frontal cortices. **16,** 1879–1887 (2013).
26. Tang, M. F., Smout, C. A., Arabzadeh, E. & Mattingley, J. B. Prediction Error and
Repetition Suppression Have Distinct Effects on Neural Representations of Visual
Information. *bioRxiv* 213710 (2017). doi:10.1101/213710
27. Alais, D. & Blake, R. *Binocular rivalry*. (MIT Press, 2005).
28. Levelt, W. J. *On binocular rivalry*. (Perception Rvo-Tno., 1965).

29.   Myers, N. E. *et al.* Testing sensory evidence against mnemonic templates. *eLife*
Sciences **4,** e09000 (2015).
30.   Alais, D., Cass, J., O'Shea, R. P. & Blake, R. Visual Sensitivity Underlying
Changes in Visual Consciousness. *Current Biology* **20,** 1362–1367 (2010).
31.   Logothetis, N. K., Leopold, D. A. & Sheinberg, D. L. What is rivalling during
binocular rivalry? *Nature* **380,** 621–624 (1996).
32.   Dehaene, S. & Naccache, L. Towards a cognitive neuroscience of
consciousness: basic evidence and a workspace framework. *Cognition* (2001).
doi:10.1016/S0010-0277(00)00123-2
33.   Salti, M. *et al.* Distinct cortical codes and temporal dynamics for conscious and
unconscious percepts. *eLife Sciences* **4,** e05652 (2015).
34.   Driver, J. & Mattingley, J. B. Parietal neglect and visual awareness. **1,** 17–22
(1998).
35.   Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming
numbers into movies. *Spat Vis* **10,** 437–442 (1997).
36.   Brainard, D. H. The Psychophysics Toolbox. *Spat Vis* **10,** 433–436 (1997).
37.   O'Shea, R. P. Effects of orientation and spatial frequency on monocular and
binocular rivalry. in 67–70 (Springer Verlag, Singapore, 1998).
38.   Haynes, J.-D. & Rees, G. Predicting the orientation of invisible stimuli from
activity in human primary visual cortex. *Nature Publishing Group* **8,** 686–691
(2005).
39.   Keil, A. *et al.* Committee report: Publication guidelines and recommendations for
studies using electroencephalography and magnetoencephalography.
Psychophysiology **51,** 1–21 (2014).
40.   Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K.-M. & Robbins, K. A. The PREP
pipeline: standardized preprocessing for large-scale EEG analysis. *Front.*
Neuroinform. **9,** B153 (2015).
41.   Wolff, M. J., Jochim, J., Akyürek, E. G. & Stokes, M. G. Dynamic hidden states
underlying working-memory-guided behavior. **20,** 864–871 (2017).
42.   Kok, P., Mostert, P. & de Lange, F. P. Prior expectations induce prestimulus
sensory templates. *Proc Natl Acad Sci USA* **26,** 201705652 (2017).
43.   Blankertz, B., Lemm, S., Treder, M., Haufe, S. & Müller, K.-R. Single-trial analysis
and classification of ERP components — A tutorial. *NeuroImage* **56,** 814–825
(2011).
551