Non-lethal oxidative stress boosts bacterial survival and evolvability
under lethal exposure

Alexandro Rodríguez-Rojas[1]*, Josh Jay Kim[1], Paul Johnston[1,2,3], Olga Makarova[1],
Murat Eravci[4], Christoph Weise[4], Regine Hengge[5] and Jens Rolff[1]

[1]Institut für Biologie, Freie Universität Berlin, Königin-Luise-Strasse 1-3, 14195
Berlin, Germany.
[2]Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin,
Germany
[3]Berlin Center for Genomics in Biodiversity Research, Berlin, Germany
[4]Institut für Chemie und Biochemie, Freie Universität Berlin, Thielallee 63,
14195 Berlin, *Germany*.
[5]Institut für Biologie/Mikrobiologie, Humboldt-Universität zu Berlin, Philippstr.
13, 10115 Berlin, Germany.

*Corresponding author: a.rojas@fu-berlin.de

Abstract

20
When oxygen began to accumulate by photosynthesis and to be utilized for
respiration, reactive oxygen species such as hydrogen peroxide, a toxic by-
product of aerobic respiration, became a challenge. Later on, multicellular
organisms began to employ reactive oxygen species as potent weapons
against microbes. Although bacterial defences against lethal and sub-lethal
oxidative stress have been studied in model bacteria, the role of prevalent
fluctuating concentrations remain unexplored. It is known that sub-lethal
exposure of *E. coli* to hydrogen peroxide results in enhanced survival upon
further exposure. Here we show how pre-treated or 'primed' cells survive
better when they later face are subsequently exposed to otherwise lethal
concentrations concentrations. The basis and the duration of the response
(memory) was also determined by time-lapse quantitative proteomics. A low
level of peroxide induces several scavenging enzymes that are quite stable,

protecting cells from future exposure. We then asked if the phenotypic resistance against H_2O_2 alters resistance evolution against oxygen stress. Experimental evolution for H_2O_2 resistance revealed faster evolution and higher levels of resistance in primed cells. Several mutations were found to be associated with resistance in evolved populations affecting different loci but counterintuitively, none of them were implicated in scavenging systems. Our results have important implications for host colonisation and infections where microbes often encounter reactive oxygen species in gradients.

Author summary

Throughout evolutionary time bacteria were exposed to reactive oxygen species and evolved the ability to scavenge toxic oxygen radicals. Multicellular organisms evolved also the ability to produce such oxygen species against pathogens. Recent studies also suggest that reactive oxygen species such as hydrogen peroxide play an important role during host gut colonisation by its microbiota. Traditionally, experiments with different antimicrobials are carried out using fixed concentrations while in nature, including intra-host environment, microbes experience this type of stress in steps or gradients. Here we show that bacteria treated with sub-lethal concentrations of hydrogen peroxide (priming) survive far better than non-treated cells when they later encounter a lethal concentration. We also found that 'priming' response has a protective role from lethal mutagenesis. This protection is provided by long-living proteins that, upon priming, remains at a high level for some generations as we could determine by time-lapse LC-Mass spectrometry. Bacteria that were primed evolved resistance faster and to a higher level. Moreover, mutations that increase resistance to hydrogen peroxide, as determined by whole genome sequencing, do not occur in the scavenger codifying genes but in other different gene functions, at least in *E. coli*.

Introduction

The ability to solicit an improved stress response when encountering repeated stress relies on 'remembering' a similar event from the past (memory) and it is a common trait to many biological entities [1]. Bacteria encounter increasing and repeated stress imposed by the host immune system, during the course of an infection or the colonization of a host [2,3]. Obata et al, for example, recently demonstrated that low levels of hydrogen peroxide (H_2O_2) in the gut of *Drosophila melanogaster* shape the composition of the gut microbiota resulting in differential survival of the flies [3]. Here, we report how bacteria respond to low levels of ROS exposure and how this impacts bacterial fitness. We then investigate if the phenotypic response to a sub-lethal dose of H_2O_2 facilitates resistance evolution, hence providing a test of the phenotype-first hypothesis, which proposes that environmentally initiated phenotypic change precedes or even facilitates evolutionary adaptation [4].

In *Escherichia coli*, the defences against oxidative stress depend on transcriptional regulators such as OxyR or SoxR that detect different changes in the redox balance and induce the production of detoxifying enzymes, DNA repair and protection system and other proteins [5]. Oxidation of OxyR by hydrogen peroxide (H_2O_2) leads to the formation of an intra-molecular disulfide bond between cysteine residues 199 and 208 and OxyR is deactivated by enzymatic reduction with glutaredoxin I (Grx) or thioredoxin (Trx). OxyR positively regulates catalases and peroxidases. Because also the Grx/GorA system is transcriptionally regulated by OxyR, the whole response is autoregulated [6].

The OxyR-mediated oxidative stress response results in scavenging of H_2O_2 and hence mitigates the toxicity of this by-product of aerobic metabolism. It includes the induction of the Suf proteins that form a complex to supply apo-enzymes with iron-sulfur clusters. This system replaces the Isc system, which is disrupted by H_2O_2 stress [7,8]. The iron sulphur clusters of dehydratases are one of the most H_2O_2-sensitive systems. The repair of those clusters by Suf is

necessary to prevent the failure of the TCA cycle. The iron-sulfur clusters are
enzymes that employ ferrous iron as a co-factor which can increase the risk of
fuelling a Fenton reaction. The nucleoid-binding protein Dps binds intracellular
free iron to limit the generation of hydroxyl radical close to the vulnerable DNA
[9].

In a classic paper it has been shown that the pre-treatment with a low dose (60
µM) of hydrogen peroxide can increase the survival of *E. coli* upon subsequent
exposure to a lethal dose (30 mM) [10]. However, neither the duration of such
priming responses nor the molecular mechanisms of its maintenance, i.e. the
memory, have been studied. Our study has two main aims. First, we
investigated the main factors in the H_2O_2 priming response and for how long
the response is sustained. Second, we used this system to test the hypothesis
that inducible phenotypes accelerate adaptive evolution [4]. Therefore, we
experimentally evolved *Escherichia coli* under a growing level of Hydrogen
peroxide with and without priming . For H_2O_2-resistant populations evolved
after priming and non-priming regimes, genome re-sequencing analyses were
performed to identify mutations. We focus our study on growing bacteria
(exponential phase), as this better reflects the situations during an infection or
colonisation of host surfaces including the gut. Moreover, this allows us to
focus on low H_2O_2-primed resistance in the absence of the stationary phase-
induced RpoS-dependent general stress response, which also generates
pronounced phenotypic H_2O_2 resistance [11,12].

Results and Discussion

Priming by H_2O_2 results in higher bacterial survival

We find that the minimal inhibitory concentration for hydrogen peroxide is 1
mM and find a clear dose-effect in survival rate (figure 1a, ranging from 50 µM
to 1 mM, figure 1a, p=2.1×10^{-16}, DRC model fitted based on maximum
likelihood [13]). Based on these results we primed cells with H_2O_2 0.1 mM, the
maximum concentration that shows no difference in growth rate compared to

non-treated cells, and monitored survival at minimal inhibitory concentration, 1
mM, starting 30 min after removal of the treatment. Primed (pre-treated)
populations of *E. coli* showed higher survival than naïve cells by more than one
order of magnitude (Cox proportional hazard model, $p < 0.05$, figure 1b).

These results are in agreement with data previously reported by Imlay et al
[10], showing survival protection even to doses as high as 30 mM, a
concentration close to hydrogen peroxide usage as a disinfectant. Our
concentrations are also in the range of some *in vivo* situations. For example,
tailfin transection on zebrafish larvae induces a rapid increase in H_2O_2 levels
ranging from 100–200 µM in the wound margins [2]. In some cases, more than
100 µM of H_2O_2 have been reported in human and other animal aqueous and
vitreous humors of the eye [14].

Duration of the priming response

As the priming response protects the cells effectively from lethal exposure to
H_2O_2, an important question is for how long this response remains effective? To
address this question, we pre-treated *E. coli* again with H_2O_2 0.1 mM during 30
minutes, but we applied the trigger concentration (1 mM) at different times-
points (30, 60, 90, 120 and 150 minutes after the 30 min priming period, also
keeping cell density constant. We observed a significant decay of the priming
response from 120 minutes after H_2O_2 pre-treatment removal (figure 2),
roughly four divisions, suggesting that the priming effect is also trans-
generational. After 150 minutes, the survival rate of primed populations no
longer differed from naïve populations. Another study has shown long-term
memory based on an epigenetic switch able to control a bimodal virulence
alternation also in the scale of hours in *E. coli* [15].

How is the state of priming maintained for up to five generations? Bacteria can
store information about recent stress via stable transcripts or proteins [15,16].
We investigated the memory at the protein level and initially studied the
impact of the two concentrations used in most of our experiments (0.1 and 1

mM) on the proteome of *E. coli* by quantitative LC-mass spectrometry. From
the quantitative proteomic dataset, we detected many of the known enzymes
that are induced by hydrogen peroxide just five minutes after the addition of
H_2O_2 (figures 3, and tables S1 and S2). For both concentrations, we detected
and quantified many proteins belonging to OxyR regulon but many other
relevant genes such as *ahpC/F*, *xthA* and *suf* operon showed a weak difference
or no response at all. We should keep in mind that the samples in this
experiment were taken after only five minutes of treatment. The reason for
short exposure is that viability after the 1 mM treatment would be too at later
timepoints. For the dose of 0.1 mM it is possible to collect viable cells after a
longer period of exposure and these results are presented in the figure 4.

To explore the temporal dynamics of the proteome after a 30-minute stimulus
with H_2O_2 (0.1 mM), we followed the changes over almost three hours (five time
points: 30, 60, 90, 120 and 150 minutes). The decline in anti-H_2O_2 protein
levels correlates well with the decrease of the response (figure 4). Some
important proteins such as KatG, AhpCF and YaaA declined slowly after
removal of the H_2O_2, corresponding probably to a sustained production with a
minor contribution of dilution due to cell division and poor degradation rate.
These results indicate that many of these proteins are stable and show a
significantly higher abundance than in the control even at 150 minutes after
H_2O_2 treatment. Other proteins such as GrxA, YaaA and XthA, SufA, SufS, AcrA-
AcrB completely declined at this point indicating that these proteins may be
subject to proteolysis and have shorter half-lives (figure 4) but also that the
stressful situation is alleviated. The overall results of this proteomic experiment
strongly suggest that the memory of priming response in *E. coli* is mediated by
the scavenging proteins such as KatG and AhpCF.

An interesting approach to visualise the global impact of H_2O_2 on bacterial
physiology is a network analysis based on protein-protein and function [17]. If
we examine, for example, the proteome response to priming concentration
(H_2O_2 0.1 mM during 30 minutes, we observe a high degree of connectivity in
network of protein-protein interaction and functional relation of both, up- and

downregulated proteins (figure 5). If we compare our network with a large-
scale protein-protein interaction network of *E. coli* [18], we find a wide
perturbation including the most important nodes. This analysis also suggests
that there is a proteome-wide readjustment to cope with hydrogen peroxide
stress that does not only involve scavenging systems and shows the profound
impact of oxidative stress across the entire proteome in a dose that does not
change the growth rate in a rich medium.

Another contributing factor to memory could be based on long-life transcripts.
Thus, we sequenced the full transcriptome after exposure to H_2O_2. RNAseq
captured both, sRNA and mRNA, and we sampled just before the decline of the
response (120 minutes after the stimulus). The highest expressed transcript
was OxyS, a small regulatory RNA (sRNA) induced by active OxyR (tables S3
and S4). OxyS regulates several genes, and although several targets have been
identified, its function is not fully understood [19]. We did not detect other
transcripts related to the OxyR regulon.

OxyS is a potential candidate to explain the duration of priming because it is
relatively stable with a half-life between 18 to 20 minutes during the
exponential phase [20–22]. To explore this, we used an *oxyS*-deficient mutant
to test its susceptibility to H_2O_2 over 30 minutes. We did not find any significant
difference in susceptibility (figure 6) consistent with prior reports [21,23,24].
Although OxyS did not provide us with a mechanism to explain the duration of
the memory, its stability could play some role in alleviating DNA damage as
recently suggested [25]. However, such protection does not seem to have a
significant impact on survival, according to our results.

Based on the comparison of proteomic and transcriptomic data we can say that
the capacity of the cells 'to remember' the stimulus is mainly based on the
stability of scavenging proteins as documented in the proteomic dataset, that
after induction remain present at higher concentrations than in the control
samples as long as 120 minutes after removal of peroxide. This indicates that
these enzymes are not degraded but probably their production go on later after

the stimulus with a minor impact of proteolysis and dilution during cell divisions
which could explain the pattern that we observe in the decline of the response.
234

235 **Priming response is compromised by disrupting important H_2O_2-**
236 **scavenging genes**
237

The expression of important genes for H_2O_2 stress survival was also confirmed
by qPCR 30 minutes after the addition of H_2O_2 (0.1 mM). H_2O_2 induced a
significant up-regulation of representative genes such as *katG*, *ahpF/ahpC*, *dps*,
mntH and *sufA* (figure S1 and table S5). As previously described in the
literature, *oxyR* and *fur* genes do not change in expression level when cells are
treated with H_2O_2, since their activation relies on swapping between active or
inactive forms of the proteins depending on the intracellular level of H_2O_2 or
iron respectably [6,7].
246

To understand the priming response to H_2O_2 at molecular level, we constructed
a set of mutants for the coding gene of key proteins that we found in our
proteomic dataset. In proliferating *E. coli*, OxyR is the major regulator
controlling cellular response to H_2O_2 [26]. We explored the involvement of OxyR
in the priming response because many of the upregulated detected proteins
are under this regulator. We found that by disrupting OxyR, there was a
dramatic change in sensitivity with full loss of viability after 30 minutes (figure
6). The priming response is completely abolished (Figure 7), indicating that the
enhanced survival, due to pre-exposure to H_2O_2, depends on OxyR. OxyR is a
fundamental transcription factor, probably the most important one, that
protects *E. coli* against H_2O_2 during the exponential phase [26,27]. The active
form of OxyR positively regulates dozens of genes.
Next, we selected, from the most highly expressed proteins and taking into
account previous reports [7,9], many from the OxyR regulon, to determine
which ones contribute more significantly to survival (figure 6) and priming
(figure 7). Removal of KatG (catalase), AhpF (one subunit of the alkyl
peroxidase AhpCF) or RecA (DNA repair) significantly decreases survival in the
presence of hydrogen peroxide (1 mM). The double mutant for KatG/AhpF

showed even stronger sensitivity. However, the absence of other important proteins induced by H_2O_2 such as YaaA (decreases intracellular iron [28]), LipA (lipoate synthase, important for the repair of iron-sulfur clusters [29]), GltD (glutamate synthase subunit), GhxA or GhrA did not result in differential survival to 1 mM concentration. The OxyS sRNA was the transcript with the highest level of expression in the RNAseq data, but the mutant did not display any increased susceptibility to H_2O_2. Assuming that H_2O_2 sensitivity and priming response are closely linked, we included in the additional priming response experiment mostly those genes with increased susceptibility (figure 7). The removal of KatG indicates that catalase importantly contributes to priming response but it does not fully explain the protection observed for the WT strain (compare to Fig. 1B). In the case of AhpF, we also observe differences in priming response with the naïve state but not that pronounced as the case of KatG deficient strain. The double mutant defective in KatG and AhpF showed a dramatic decrease in priming response but still significantly different from the naïve. Another protein that illustrates an important influence on priming response is RecA, with the mutant showing a decreased priming response to H_2O_2. OxyS mutant is not suppressive for priming response, indicating that its presence contributes neither to the survival or priming. Overall, our data indicate that the priming by a low dose of H_2O_2 is multifactorial, with several OxyR-controlled proteins such as KatG, AhpCF or other factors related to DNA as RecA contributing to the priming response.

Priming response enhances the survival of evolving populations

To find out whether the priming response described above has an influence on the rate by which bacterial resistance against hydrogen peroxide evolves was the goal of the following experiments. We evolved bacterial cultures with a treatment protocol described in detail in M&M section. Parallel populations were evolved where one group was periodically exposed to a sub-lethal concentration of hydrogen peroxide, an activating signal that should protect the populations in comparison with naïve ones when later exposed to a higher

dose. We continued daily intermittent exposures to hydrogen peroxide doubling both, priming and triggering concentrations until extinction occured. We observed that the extinction rate was faster for the naïve populations in comparison to primed populations (Log-rank test, p< 0.05), figure 8). Naïve cells evolved resistance of up 8 mM H_2O_2, whereas primed cells evolved resistance to at least 16 mM H_2O_2. This first result indicates that priming potentially increases the evolvability of pre-treated populations.

Further analysis of the resistant populations by whole-genome sequencing revealed that all populations harboured different sets of mutations. Surprisingly, we did not find any modification in the enzymatic scavenger systems, such as catalases or any other proteins related to peroxide protection. We found several hundreds of single nucleotide polymorphisms (SNPs) and other type of mutations that were unique to each population for both regimes (figure 9). These mutations probably represent many of the neutral or non-lethal changes that populations accumulated during the exposure to H_2O_2, that is not simply a selective agent, but it also speeds up evolution by increasing the mutation rate. At the moment we cannot be certain about the contribution of each mutation to H_2O_2 stress adaptation and they will be subject of detailed studies in the future. However, it is possible to speculate about the possible role of many of the mutations we found. For example, a set of changes are located in genes coding for iron binding proteins or related with iron transport such as *iceT*, *feoA, yaaX/yaaA* or *rsxC*. The control of intracellular iron is crucial to decrease the adverse effects of Fenton chemistry [7].

There was also a set of more frequent changes that were common to both types of population, evolved under priming and non-priming conditions (figure 9). The most frequent mutations were *yodB* (a cytochrome b561 homolog), intergenic mutations between *insA* and *uspC* (universal stress protein C), intergenic mutations between *flhD* and *insB1*. Another frequent mutation was in the gene *yagH*, belonging to the CP4-6 prophage. CP4-6 is a cryptic prophage in *E. coli* that has been suggested could play a role in bacterial survival under adverse environmental conditions [30].

Priming alters the mutational spectrum and H₂O₂-induced mutagenesis in evolving populations

To assess if the evolved populations have a similar mutation spectrum, we analysed the total pool of mutations segregated by the treatments and determined the proportion of every type of found mutation. We used Monte Carlo hypergeometric test implemented by iMARS [31], to assess the overall differences between mutational spectra. Both groups, of the evolving populations under priming and non-priming conditions, differed significantly (p=0.00021, figure 6). ROS induces a particular type of mutations, with a characteristic signature in the DNA. The guanine base in genomic DNA is highly susceptible to oxidative stress due to having the lowest oxidation potential. Therefore, G·C→T·A and G·C→C·G transversion mutations frequently occur under oxidative conditions [32,33]. Thus, we investigated the proportion of C→A and C→G substitutions between the two types of evolving regimes, but we did not find significant differences (p=0.056 and p=0.11 respectively, two-tailed Fisher's exact-test, figure 10).

A possible explanation for these results is that the pre-activation of antioxidant defences helps to decrease the number of non-adaptive mutations and lethality by decreasing the extent to which the Fenton reaction occurs. We observed an increase in the frequency of substitutions G→A, T→C, G→C, T→G and A→G in naïve populations compared to primed populations, suggesting that the primed cells have DNA-damage control. Primed populations showed an increase in frameshift mutations by deletion of 1 bp, and additions of C or G; these last two were only present in primed populations that also showed higher number of A→C substitutions.

It is expected that the priming response would decrease DNA damage and hence also the rate of lethal mutations. To assess this possibility, we tested if priming is able to decrease the mutant frequency and hence lethality due to mutagenesis. Preserving the conditions of the previous experiments, we treated the bacterial cultures with the same priming and triggering concentrations at low cell density, the basal mutant frequency without H₂O₂

treatment was also determined. We found that the priming response decreases
H_2O_2-induced mutant frequency close to one order of magnitude if compared
with naïve cells ($p<0.001$ primed vs naïve, Welch's test, figure S2). In addition,
no difference was found between primed bacteria and the basal level of
mutagenesis ($p=0.03$ primed vs basal, Welch's test), indicating that priming is
able to supress most of the mutagenesis caused by H_2O_2. In our experiment,
naïve and primed populations showed different death rates (figure 1). We also
showed previously that different inoculum size on similar conditions to our
experiment has no influence on mutagenesis [34]. This experiment allows us to
deduce that one of the most important consequences of the priming response
to H_2O_2 is a drastic decrease in lethal mutagenesis, indicating that although
hydrogen peroxide damages most of the cellular components [7], damage to
DNA is likely a major contributor to lethality.
377
In principle, an increase of mutation rate is a parameter that increases the
evolvability of asexual populations [35–37]. How is it possible that naïve
populations show lower evolvability compared to primed populations despite a
higher mutation rate? A possible explanation is that evolvability is also
influenced by the population size and the mutation supply that is increased by
mutagenesis is balanced by the decrease in population size of the naïve
population. We also found that the mutational spectra of our evolving
populations are different. Mutational spectra are a qualitative property of
mutation rate that could enhance or hinder the access to beneficial mutations
[38–40]. It is possible that the observed changes in mutational spectra between
primed and naïve evolving populations could play a role in balancing the ratio
of deleterious/beneficial mutations but to confirm this fact in our experiments
is not a trivial task.
391
392 **Conclusions**
393
394 We found that priming by hydrogen peroxide in exponentially growing bacteria
395 is quantitatively driven by genes that are mostly under OxyR control. Some
396 other genes such as *recA* also seem to play an important role. The memory of

the priming response can last up to four generations upon first exposure to H_2O_2 taking into account the time after addition of the stressor and number of divisions. We also showed high stability of hydrogen peroxide detoxifying proteins which play a significant role in survival. Moreover, we found that priming plays a role in evolving populations by enhancing the survival under oxidative stress, which supports a link between the phenotype as a target of selection during evolutionary processes and provides a validation for the phenotype first hypothesis [4]. The type and number of mutations indicates that scavenger systems against oxidative stress are optimally evolved since no mutations affecting these systems were found under H_2O_2 stress selective pressure. Our work also contributes to understanding how the adaptation of early life with oxygen could have taken place.

Material and methods

Bacteria and growth conditions. The *E. coli* MG1655 was used as bacterial model for all experiments with hydrogen peroxide. For genetic manipulation, *Escherichia coli* DH5α strain was used and routinely cultured in Lysogeny Broth (LB medium), supplemented with antibiotics when appropriate. All bacterial strains were cultured in Lysogeny broth lenox (Carl Roth, Germany). For all experiments with hydrogen peroxide the LB was fresh prepared and kept in the dark until use.

Construction and verification of deletion mutants. All mutants used in this work (table S6) were generated in *E. coli* K-12 strain MG1655 following the a modified methodology described elsewhere [41]. Briefly, transformants carrying a red recombinase helper plasmid, pKD46, were grown in 5-ml SOB medium with ampicillin (100 μg/ml) and L-arabinose at 30°C to an O.D.600 of ~0.5 and then made electrocompetent. PCR products with homology regions were generated using specific primers (table S7) to amplified the region of interest from using the corresponding mutants of the Keio collection [42]. The PRC-generated fragments were purified (MinElute PCR Purification Kit, Qiagen). Competent cells in 50 μl aliquots were electroporated with 100 ng of PCR product. Cells were added immediately 0.9 ml of SOC, incubated 1 h at 37°C, and then 100 μl aliquots spread onto LB agar with kanamycin (30 μg/ml). The mutants were verified by PCR and the antibiotic resistance cassette was removed using the plasmid pCP20. The correct inactivation of genes was verified by PCR. To construct double mutants, singles mutants obtained in MG1655 were transduced using the P1vir phage procedure as previously described [43].

Hydrogen peroxide susceptibility testing. When working with hydrogen peroxide, we carried out all the experiments with fresh-made LB to avoid the accumulation of chemically formed H_2O_2 by joined the action of light and flavin present in the medium during storage. Susceptibility to H_2O_2 was determined by exposing exponentially phase bacteria at a density of ~ $2x10^7$ CFU/ml to different concentrations and times, taking as a reference previous works.

Priming experiments with H_2O_2. Starting from $2x10^7$ CFU/ml, where *E. coli* the bacteria were exposed (stimulus) to 0.1 mM of hydrogen peroxide during 30 minutes at at 37°C with shaking. Catalase was added to a final concentration of 4 μl/ml, centrifuged at 4000 g during 10 minutes and allow to recover during 90 minutes, keeping the cell density constant by removing the needed volume and replace it with appropriate amount of fresh LB every 30 minutes. The trigger (H_2O_2 1 mM) was added 90 minutes after removal the

stimulus. The challenge lasted for 30 minutes. At this point catalase (4 µl/ml) was added and cultures were diluted and plated to determine cell viability. Non treated cells were used as control. Five cultures per group were used.

Priming experiments in multi-well plates. The 96-multiwell plates were prepared under aseptic conditions. All experiments were carried out in a final volume of 100 µL of fresh LB. Bacteria were exposed to serially diluted concentrations of H_2O_2 from 256 mM by adding 50 µL of peroxide containing LB first 50 µL of mid-exponential phase bacteria rendering approximately 10^7 CFU/ml suspension. The plate was placed into a Synergy H1 plate reader (BioTek, USA) and kinetics of killing or growth were followed during two hours. Once those relevant non-lethal concentrations in these conditions were established, a second experiment was carried out. In a similar plate and conditions, bacteria were exposed to serial dilutions of H_2O_2 starting from 4 mM during 30 minutes. The plate was centrifuged, the supernatant removed and fresh LB transferred. The cells were allowed to recover for two hours. After this period, 50 µL were removed per well and replaced with 50 µL LB containing 32 mM of H_2O_2 for a final concentration of 16 mM. The killing and survival were followed for 8 hours by absorbance at 600 nm.

Memory of the priming response. After applying a stimulus (0.1 mM H_2O_2) during half an hour, the cells were allowed to recover for 30, 60, 90, 120 and 150 minutes before the addition of the trigger (1 mM H_2O_2). During the experiment, the D.O.600 of all cultures were kept around 0.05 by adding suitable volume and removal of equivalent quantity to maintain the population in exponential phase and the same population size. Bacteria remained exposed to the trigger during 30 minutes before starting the dilution in 1 ml of LB containing 4 µg of catalase (Sigma-Aldrich, Germany). Appropriated dilutions were made and plated onto LB agar to determine the survival rate.

Global proteomics by mass spectrometry. *E. coli* strain MG1655 was grown at 37°C in LB medium to OD600 0.5. The cultures were diluted ten times in fresh LB. H_2O_2 was added to tubes at 0.1 and 1 mM. Non treated samples

were used as control. All tubes were incubated during five minutes with
shaking at 37°C. Remaining hydrogen peroxide was removed by adding 4
µg/ml of catalase first and by centrifugation at 10000 g during 2 minutes. After
removal of the supernatant, equivalent amount of fresh LB was added. For the
memory decline experiment, the procedure was identical, except that only 0.1
mM of hydrogen peroxide was used and the samples were treated during 30
minutes. After removal of the treatment, time-point samples were taken after
30, 60, 90, 120 and 150 minutes. Each experimental condition consisted of six
replicas. Two millilitres per sample were pelleted by centrifugation at 10000 g
for 2 minutes. The cells were re-suspended in 50 ml of TE (10 mM Tris-HCl pH
8.0, 1 mM EDTA) that contained chicken lysozyme (1 mg/ml, Sigma) and DTT
(10 mM) and incubated at room temperature for five minutes with occasional
swirling. A volume of 250 µl of denaturation buffer (6M urea/2 M thiourea in 10
mM HEPES pH 8.0) was added into each sample. Then, 10 µl (which
approximately corresponds to 50 µg of protein) of the resulting lysate were
mixed with 150 µl of the denaturation buffer and used for in-solution protein
digestion as described previously [44]. Briefly, proteins, demobilised in
denaturation buffer, were reduced by addition of 1 ml of 10 mM DTT dissolved
in 50 mM ammonium bicarbonate (ABC) and incubation for 30 minutes,
followed by alkylation with 1 µl of 55 mM iodacetamid dissolved in 50 mM ABC
for 20 minutes, digestion with 1 µg Lysyl endopeptidase (LysC, Wako, Japan),
resuspended in 50 mM ABC, for 3 hours. After pre-digestion with LysC, protein
samples were diluted four times with 50 mM ABC and subjected to overnight
trypsin digestion using 1 µg of sequencing grade modified trypsin, Promega),
diluted in 50 mM ABC. All in-solution protein digestion steps were performed at
room temperature and after addition of iodacetamid the samples were
protected from the light until the digestion was stopped by adding 10 ml of
Buffer A* (5% acetonitrile, 3% trifluoric acid). After trypsin digestion overnight
at room temperature, samples were diluted with 200 µl of buffer A (5%
acetonitrile, 3% trifluoracetic acid) treated as described elsewhere and vacuum
dried. Re-dissolved samples were loaded on a ReprosilPur C18 reverse phase
column and peptides were analyzed using a nano-HPLC Dionex Ultimate 3000
system (Thermo scientific) coupled to an Orbitrap Velos mass spectrometer

(Thermo scientific). MS and MS/MS data from each LC/MS run were analysed
with MaxQuant software [45]. Identification of proteins was performed using
the MaxQuant implemented Andromeda peptide search engine.

Quantification of gene expression. To assess the quality of hydrogen
peroxide stress response related genes in *E. coli*, treated bacteria comparison
with 0.1 mM of hydrogen peroxide were compared to non-treated control.
Preparation of each of the six treatments and the control samples was carried
out by two operators three times on three consecutive days, thereby
comprising three biological replicates for each of the six experimental and one
control groups. Overnight (ON) cultures were diluted 1:100 in fresh LB,
subdivided into six 10 ml aliquots and incubated in 50 ml-Falcon tubes at 37°C
for 2.5 hours at 220 rpm. When the OD600 of bacterial cultures reached 0.4-
0.5, the cultures were diluted ten times with LB and hydrogen peroxide was
added to the final concentrations of 0.1 mM. These bacterial cultures were
incubated for 30 minutes at 37°C with shaking. Catalase was added to each
tube to a final concentration of 4 µg/ml. Ten millilitres of treated cultures and
from the control tubes were collected, immediately centrifuged at 10 000 g for
2 minutes and the supernatant was fully removed. The bacterial pellets were
resuspended in 1 ml LB, followed by addition of 1ml of RNAprotect Bacteria
Reagent (Qiagen, Germany), incubated during 2 minutes and centrifuged at
10000 g for 2 minutes at room temperature. The supernatant was discarded
and the bacterial pellet was immediately frozen and stored at -80°C until RNA
extraction.

RNA was isolated using RNeasy kit (Qiagen) according to the manufacturer's
instructions and eluted in 50 µl of RNase-free water. The nucleic acid yield and
purity were determined by measuring the optical density at A260/280 using
Nanodrop spectrophotometer (Thermo Scientific). RNA samples were treated
with TURBO DNase (Life Technologies). Briefly, 10 µg of RNA were used in a
total volume of 500 µl containing 20 units of TURBO DNase, incubated for 30
minutes at 37°C, immediately followed by RNeasy (Qiagen) clean-up and
elution in 30 µl of RNase-free water. Following DNase treatment, RNA integrity

was assessed using Agilent RNA 6000 Nano kit and 2100 Bioanalyzer
instrument (both Agilent Technologies). All samples had RIN values above 8.
564
For cDNA synthesis, total RNA (250 ng per reaction) and random primers were
used for cDNA synthesis using High-Capacity cDNA Reverse Transcription Kit
with RNase Inhibitor (Applied Biosystems). Initially, to ensure linear conversion
of the transcripts, the dynamic range of reverse transcription reaction was
tested by performing a standard curve with cDNA, synthesised using various
input amounts of pooled RNA. To obtain sufficient amount of cDNA, several
batches of 20 µl RT reactions were pooled, diluted 50-fold with RNase-free
water and stored in single-use aliquots at -80°C until further use. All 21
samples were tested for presence of contaminating genomic DNA by running
the *mdoG* assay with cDNA and the respective no reverse transcription (-RT)
controls. There was no amplification in the majority of –RT controls. In –RT
samples with detectable amplification, difference in Ct values when compared
with +RT varied between the samples, but was no less than 10 cycles for all of
them with the lowest Ct values in –RT control samples ≥ 30.
579
For primer design, *Escherichia coli* strain K-12 MG1655 complete genome
(accession U00096) sequence was downloaded from NCBI database
(www.ncbi.nlm.nih.gov) and used as reference. Target sequence accession
number, and primer sequences for each assay can be found in table S8.
Primers were designed using Primer Express software (Applied Biosystems) and
optimized for annealing temperature of 60°C. Each primer pair and amplicon
were checked for secondary structure formation using Oligo Tool (Integrated
DNA technologies, http://eu.idtdna.com/analyzer/Applications/OligoAnalyzer/).
588
Quantitative Real-time PCR reactions were prepared manually in a total volume
of 10 µl by mixing 5 µl 2x KAPA SYBR FAST ABI Prism master mix (KAPA
Biosystems), 0.2 µl forward and reverse primer mix (10 µM each primer), 2.8 µl
RNase-free water and 2 µl cDNA in MicroAmp Fast Optical 48-wells reaction
plates (Applied Biosystems). PCR reactions were run in the Fast mode using
StepOne thermocycler (Applied Biosystems) with the following cycling

conditions: 95°C 3' /40x(95°C 3''/60°C 20'')/melting curve analysis. Each assay

was run in duplicate. No template controls (NTC) were run each time. Presence

of a single specific product was verified by running a melt curve analysis

followed by visualisation of the qPCR product in a 2% agarose gel stained with

SYBR Safe DNA Gel stain (Life Tecnologies). Additionally, each PCR assay was

tested for reaction efficiency as follows: equimolar amounts of cDNA from all 21

samples were pooled together and used for the preparation of the standard

curve by serial dilution (1:3) over five dilution points and run in triplicate.

Expression of target genes was normalized to the expression levels of three

reference genes (*arcA*, *mdoG* and *tus*), selected based on the assessment of

the expression stability across all experimental conditions using BestKeeper

software. Reaction efficiency information inferred from the standard curve data

was used to correct for differences in amplification efficiencies in the REST

2009 software [46]. Default settings (2000 iterations) were used for

randomisation and bootstrapping analysis to test significance of gene

expression. Expression values with p-values ≤0.05 were assigned as

differentially expressed. We followed the Minimum Information for Publication

of Quantitative Real-Time PCR experiments (MIQE) guidelines-compliant check-

list [47].

For data analysis, qPCR Amplification curves were first visually examined in the

StepOne software (Applied Biosystems). No baseline and threshold line

adjustments were necessary. Ct values of the technical replicates were

averaged and used for relative gene expression analysis in the REST 2009

software (Qiagen, Germany) [46]. Expression of target genes was normalized

to the expression levels of three reference genes (*arcA*, *mdoG* and *tus*),

selected based on the assessment of the expression stability across all

experimental conditions using BestKeeper software. Reaction efficiency

information inferred from the standard curve data was used to correct for

differences in amplification efficiencies in the REST 2009 software. Default

settings (2000 iterations) were used for randomisation and bootstrapping

analysis to test significance of gene expression. Expression values with p-

values ≤0.05 were assigned as differentially expressed. We followed the

Minimum Information for Publication of Quantitative Real-Time PCR

experiments (MIQE) guidelines-compliant check-list [47].

Transcriptome sequencing. The transcriptome sequence was carried out on

samples treated with 0.1 mM of hydrogen peroxide during the experiment of

memory decline. The time point used corresponded to 120 minutes after

removal of the treatment. Bacterial cells that were kept in low density (0.05

O.D.600) were concentrated ten times and the pellets were resuspended in 1

ml of lysis buffer. Both, small RNA fraction and the large one were isolated

using the microRNA & small RNA Isolation kit (Thermo Fisher Scientific,

Germany). Traces of genomic DNA were removed from 10 µg of RNA by

digestion in a total volume of 500 µl containing 20 units of TURBO DNase,

incubated for 30 minutes at 37°C, immediately followed by RNeasy (Qiagen)

clean-up and elution in 30 µl of RNase-free water. Following DNase treatment,

RNA integrity was assessed using Agilent RNA 6000 Nano kit and 2100

Bioanalyzer instrument (both Agilent Technologies, USA). Both fractions were

depleted from ribosome RNA using the Ribo-Zero depletion Kit for Gram

negative bacteria (Illumina, USA). Libraries were prepared using a TruSeq

Stranded Total RNA library preparation kit (Illumina, USA) and were sequenced

on a MiSeq platform. All sequences are available are available from the NCBI

SRA under BioProject accession PRJNA485867.

Evolution experiment. Before the beginning of the experiment, five

independent clones of *E. coli* MG1655 were pre-adapted to the experimental

conditions such as culture medium, temperature and oxygen level and other

parameters. The pre-adaptation was carried out by diluting an overnight

culture of each clone 1:1000 and incubated at 37°C in the plate reader with the

same shaking conditions used during the rest of the experiment in 0.2 ml in

fresh-made Lysogeny Broth Lenox(LB). The bacteria we cultured by serial

passage every 24 hours using a bottleneck of 1% (1/100 dilution) to allow for

approximately fifty generations during ten days. Contamination checks by

plating out on LB and cryopreservation of culture aliquots at -80°C in LB

containing 15% glycerol solution were carried out periodically. All lines showed

similar fitness and five independent colonies (one per line) were selected for the evolution experiment. The experiment was performed at 37°C with periodical shaking in a microplate reader (Synergy 2, Biotek). We used flat bottom polypropylene non-binding 96-well plates from Greiner Bio-One GmbH (Germany). The plates were covered with polystyrene lids with anti-evaporation rings also from Greiner Bio-One GmbH. A final volume of 200 µl per well was used. Growth curves were generated by taking measurements of OD600 every 20 minutes (preceded by a brief shaking of 5 seconds). Every day, the experiment consisted of two stages. Every morning a V-bottom plate containing 190 µl of LB was inoculated with 2 µl of overnight culture and incubated for two hours to reach an OD between 0.2 to 0.6. At this point, 3 µl of LB containing the corresponding priming concentration of H_2O_2 (for populations to be primed) or only LB (control populations) were added to each well. The plate was then incubated for 30 minutes, centrifuged and the supernatant removed using an 8-channel vacuum line provided of disposable tips. The pellets were resuspended in 200 µl of fresh LB and 190 µl were transferred to a new flat-bottom plate, allowed the cell to recover for 30 minutes at 37 °C and challenged with a ten-fold higher concentration of the priming one. The dish was located in the plate reader and the growth was followed as described above during 20 hours. The serial passage started at 50µM of H_2O_2. Next day, the procedure was identical by both, priming and challenging concentrations were doubled to reach a final challenging concentration of 32 mM of H_2O_2 where all bacterial population went extinct. Besides, 10 µl of the remaining cultures were added to 180 µl of sterile 0.9% NaCl and then serially diluted, plated and inspected for contamination.

DNA isolation. Genomic DNA samples for whole genome sequencing were isolated using in house method based on fast phenol:chloroform extraction, removal of RNA and ethanol precipitation, capturing the DNA in paper filter as previously described. The DNA quantity and quality were estimated by measuring the optical density at A260/280 using the Nanodrop spectrophotometer (Thermo Scientific) and agarose gel electrophoresis.

Genome re-sequencing. We sequenced the total genomic DNA coming from
sixty populations from the evolution experiment, all from the previous passage
before extinction (20 evolved from priming regime, 20 from non-priming
regime and 20 control strains). TruSeq DNA PCR-free libraries were constructed
according to the manufacturer's instructions and sequenced for 600 cycles
using a MiSeq at the Berlin Center for Genomics in Biodiversity Research.
Sequence data are available from the NCBI SRA under BioProject accession
PRJNA485867.

The haploid variant calling pipeline snippy [48] was used to identify mutations
in the selection lines. Snippy uses bwa [49] to align reads to the reference
genome and identifies variants in the resulting alignments using FreeBayes
[50]. All variants were independently verified using a second computational
pipeline, breseq [51].

Determination of H_2O_2-induced mutagenesis. This procedure was carried
out following similar protocols to previous studies with some modifications
[52,53]. Five independent cultures (5 ml each one) of *E. coli* MG1655 was
grown in fresh-made LB medium to an OD of ~0.2. At this point, each culture
was diluted 10 times (volume was increased to 50 ml) with LB and divided into
two separate sets of 25 ml per culture each one. One set, consisting of five
cultures of 25 ml was treated with 0.1 mM of H_2O_2 (primed) while the second
set of culture remained untreated (naïve). After 30 minutes, both cultures were
treated with 1 mM of H_2O_2 during another 30 minutes. Then, H_2O_2 was
removed, first, by quickly adding 4 µl/ml of catalase and, second, by collecting
the cells by centrifugation and washing them with one ml of fresh LB. An
additional set of tubes received no treatment all and was used to determine
the basal frequency of mutants. For determination of H_2O_2-induced mutation
frequency, 4 ml of fresh LB was inoculated with 1 ml of washed bacteria from
each culture. The cultures were allowed to recover overnight 16 h at 37°C.
Serial dilutions of each culture were plated next day onto LB plates containing
100 ug/ml of rifampicin or without antibiotic to estimate the viability. The three
groups were compared among them using a Welch's test.

Statistical analysis. For point to point comparisons in sensitivity and priming killing curves a Welch's test was used. For survival analysis we used a log-rank test. Some additional analyses are mentioned elsewhere in the manuscript. All tests were performed with software package R [54].

Funding information. ARR and JR were supported SFB 973 (Deutsche Forschungsgemeinschaft, CRC973, project C5) and OM and JR by European Research Council grant 260986 to JR.

Acknowledgements. We would like to thank Heidrun Häweker and Elisa Bitterman for help with library preparations and technical assistance. We are grateful to Gisela Storz from the National Institutes of Health, Bethesda, USA for sharing with us the OxyS defective mutant.

References

1. Hilker M, Schwachtje J, Baier M, Balazadeh S, Bäurle I, Geiselhardt S, et al. Priming and memory of stress responses in organisms lacking a nervous system. Biol Rev Camb Philos Soc. 2015; doi:10.1111/brv.12215

2. Wittmann C, Chockley P, Singh SK, Pase L, Lieschke GJ, Grabher C, et al. Hydrogen Peroxide in Inflammation: Messenger, Guide, and Assassin. Adv Hematol. Hindawi Publishing Corporation; 2012;2012: 1–6. doi:10.1155/2012/541471

3. Obata F, Fons CO, Gould AP. Early-life exposure to low-dose oxidants can increase longevity via microbiome remodelling in Drosophila. Nat Commun. Nature Publishing Group; 2018;9: 975. doi:10.1038/s41467-018-

03070-w

4.   Levis NA, Pfennig DW. Evaluating 'Plasticity-First' Evolution in Nature: Key
Criteria and Empirical Approaches. Trends Ecol Evol. 2016;31: 563–574.
doi:10.1016/j.tree.2016.03.012

5.   Zheng M, Wang X, Templeton LJ, Smulski DR, LaRossa RA, Storz G. DNA
microarray-mediated transcriptional profiling of the Escherichia coli
response to hydrogen peroxide. J Bacteriol. 2001;183: 4562–70.
doi:10.1128/JB.183.15.4562-4570.2001

6.   Zheng M, Aslund F, Storz G. Activation of the OxyR transcription factor by
reversible disulfide bond formation. Science. 1998;279: 1718–21.
Available: http://www.ncbi.nlm.nih.gov/pubmed/9497290

7.   Imlay J a. Cellular defenses against superoxide and hydrogen peroxide.
Annu Rev Biochem. 2008;77: 755–76.
doi:10.1146/annurev.biochem.77.061606.161055

8.   Ezraty B, Vergnes A, Banzhaf M, Duverger Y, Huguenot A, Brochado AR, et
al. Fe-S cluster biosynthesis controls uptake of aminoglycosides in a ROS-
less death pathway. Science. 2013;340: 1583–7.
doi:10.1126/science.1238328

9.   Imlay JA. The molecular mechanisms and physiological consequences of
oxidative stress : lessons from a model bacterium. Nat Publ Gr. Nature
Publishing Group; 2013; doi:10.1038/nrmicro3032

10.   Imlay JA, Linn S. Bimodal pattern of killing of DNA-repair-defective or
anoxically grown Escherichia coli by hydrogen peroxide. J Bacteriol.
1986;166: 519–27. Available:
http://www.ncbi.nlm.nih.gov/pubmed/3516975

11.   Jenkins DE, Schultz JE, Matin A. Starvation-induced cross protection
against heat or H2O2 challenge in Escherichia coli. J Bacteriol. 1988;170:
3910–3914. doi:10.1128/jb.170.9.3910-3914.1988

12.   Lange R, Hengge-Aronis R. Identification of a central regulator of
stationary-phase gene expression in *Escherichia coli*. Mol Microbiol.
1991;5: 49–59. doi:10.1111/j.1365-2958.1991.tb01825.x

13.   Ritz C, Baty F, Streibig JC, Gerhard D. Dose-Response Analysis Using R.
Xia Y, editor. PLoS One. Public Library of Science; 2015;10: e0146021.

doi:10.1371/journal.pone.0146021

14.   Halliwell B, Clement MV, Long LH. Hydrogen peroxide in the human body.
FEBS Lett. No longer published by Elsevier; 2000;486: 10–13. doi:10.1016/
S0014-5793(00)02197-9

15.   Ronin I, Katsowich N, Rosenshine I, Balaban NQ. A long-term epigenetic
memory switch controls bacterial virulence bimodality. Elife. 2017;6.
doi:10.7554/eLife.19599

16.   Veening J-W, Smits WK, Kuipers OP. Bistability, Epigenetics, and Bet-
Hedging in Bacteria. Annu Rev Microbiol. Annual Reviews ; 2008;62: 193–
210. doi:10.1146/annurev.micro.62.081307.163002

17.   Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas
J, et al. STRING v10: protein–protein interaction networks, integrated over
the tree of life. Nucleic Acids Res. 2015;43: D447–D452.
doi:10.1093/nar/gku1003

18.   Arifuzzaman M, Maeda M, Itoh A, Nishikata K, Takita C, Saito R, et al.
Large-scale identification of protein-protein interaction of Escherichia coli
K-12. Genome Res. 2006;16: 686–691. doi:10.1101/gr.4527806

19.   Gottesman S. The Small RNA Regulators of *Escherichia coli* : Roles and
Mechanisms. Annu Rev Microbiol. 2004;58: 303–328.
doi:10.1146/annurev.micro.58.030603.123841

20.   Basineni SR, Madhugiri R, Kolmsee T, Hengge R, Klug G. The influence of
Hfq and ribonucleases on the stability of the small non-coding RNA OxyS
and its target rpoS in E. coli is growth phase dependent. RNA Biol.
2009/12/18. 2009;6: 584–594. doi:10082 [pii]

21.   González-Flecha B, Demple B. Role for the oxyS gene in regulation of
intracellular hydrogen peroxide in Escherichia coli. J Bacteriol. 1999;181:
3833–6. Available: http://www.pubmedcentral.nih.gov/articlerender.fcgi?
artid=93864&tool=pmcentrez&rendertype=abstract

22.   Zhang A, Altuvia S, Tiwari A, Argaman L, Hengge-Aronis R, Storz G. The
OxyS regulatory RNA represses rpoS translation and binds the Hfq (HF-I)
protein. EMBO J. 1998;17: 6061–6068. doi:10.1093/emboj/17.20.6061

23.   Altuvia S, Weinstein-Fischer D, Zhang A, Postow L, Storz G. A small, stable
RNA induced by oxidative stress: role as a pleiotropic regulator and

antimutator. Cell. 1997;90: 43–53. Available: http://www.ncbi.nlm.nih.gov/
pubmed/9230301

24.	Tjaden B, Goodwin SS, Opdyke JA, Guillier M, Fu DX, Gottesman S, et al.
Target prediction for small, noncoding RNAs in bacteria. Nucleic Acids
Res. 2006;34: 2791–2802. doi:10.1093/nar/gkl356

25.	Barshishat S, Elgrably-Weiss M, Edelstein J, Georg J, Govindarajan S, Haviv
M, et al. OxyS small RNA induces cell cycle arrest to allow DNA damage
repair. EMBO J. 2018;37: 413–426. doi:10.15252/embj.201797651

26.	Imlay JA. Transcription Factors That Defend Bacteria Against Reactive
Oxygen Species. Annu Rev Microbiol. Annual Reviews ; 2015;69: 93–108.
doi:10.1146/annurev-micro-091014-104322

27.	Aslund F, Zheng M, Beckwith J, Storz G. Regulation of the OxyR
transcription factor by hydrogen peroxide and the cellular thiol-disulfide
status. Proc Natl Acad Sci U S A. 1999;96: 6161–5. Available:
http://www.ncbi.nlm.nih.gov/pubmed/10339558

28.	Liu Y, Bauer SC, Imlay JA. The YaaA protein of the Escherichia coli OxyR
regulon lessens hydrogen peroxide toxicity by diminishing the amount of
intracellular unincorporated iron. J Bacteriol. American Society for
Microbiology (ASM); 2011;193: 2186–96. doi:10.1128/JB.00001-11

29.	Ollagnier-De Choudens S, Sanakis Y, Hewitson KS, Roach P, Baldwin JE,
Münck E, et al. Iron-sulfur center of biotin synthase and lipoate synthase.
Biochemistry. 2000;39: 4165–73. Available:
http://www.ncbi.nlm.nih.gov/pubmed/10747808

30.	Wang X, Kim Y, Ma Q, Hong SH, Pokusaeva K, Sturino JM, et al. Cryptic
prophages help bacteria cope with adverse environments. Nat Commun.
Nature Publishing Group; 2010;1: 147. doi:10.1038/ncomms1146

31.	Morgan C, Lewis PD. iMARS - Mutation analysis reporting software: An
analysis of spontaneous cII mutation spectra. Mutat Res - Genet Toxicol
Environ Mutagen. 2006;603: 15–26. doi:10.1016/j.mrgentox.2005.09.010

32.	Kino K, Sugiyama H. Possible cause of G·C→C·G transversion mutation by
guanine oxidation product, imidazolone. Chem Biol. Cell Press; 2001;8:
369–378. doi:10.1016/S1074-5521(01)00019-9

33.	Rodríguez-Rojas A, Makarova O, Müller U, Rolff J. Cationic Peptides

Facilitate Iron-induced Mutagenesis in Bacteria. Matic I, editor. PLOS
Genet. Public Library of Science; 2015;11: e1005546.
doi:10.1371/journal.pgen.1005546

34. Rodríguez-Rojas A, Makarova O, Rolff J. Antimicrobials, Stress and
Mutagenesis. PLoS Pathog. 2014;10. doi:10.1371/journal.ppat.1004445

35. Galhardo RS, Hastings PJ, Rosenberg SM. Mutation as a Stress Response
and the Regulation of Evolvability. Crit Rev Biochem Mol Biol. 2007;42:
399–435. doi:10.1080/10409230701648502

36. Blázquez J, Couce A, Rodríguez-Beltrán J, Rodríguez-Rojas A, Blazquez J,
Couce A, et al. Antimicrobials as promoters of genetic variation. Curr Opin
Microbiol. 2012/08/15. 2012;15: 561–569. doi:S1369-5274(12)00102-6
[pii]10.1016/j.mib.2012.07.007

37. Ram Y, Hadany L. The evolution of stress-induced hypermutation in
asexual populations. Evolution (N Y). 2012;66: 2315–2328.
doi:10.1111/j.1558-5646.2012.01576.x

38. Couce A, Guelfo JR, Blázquez J. Mutational spectrum drives the rise of
mutator bacteria. PLoS Genet. Public Library of Science; 2013;9:
e1003167. doi:10.1371/journal.pgen.1003167

39. Couce A, Rodríguez-Rojas A, Blázquez J. Bypass of genetic constraints
during mutator evolution to antibiotic resistance. Proc Biol Sci. 2015;282:
20142698. doi:10.1098/rspb.2014.2698

40. Agashe D. The road not taken: Could stress-specific mutations lead to
different evolutionary paths? PLOS Biol. 2017;15: e2002862.
doi:10.1371/journal.pbio.2002862

41. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A.
2000/06/01. 2000;97: 6640–6645. doi:10.1073/pnas.120163297
120163297 [pii]

42. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al.
Construction of Escherichia coli K-12 in-frame, single-gene knockout
mutants: the Keio collection. Mol Syst Biol. 2006/06/02. 2006;2: 2006
0008. doi:msb4100050 [pii] 10.1038/msb4100050

43. Thomason LC, Costantino N, Court DL. *E. coli* Genome Manipulation by P1

Transduction. Current Protocols in Molecular Biology. Hoboken, NJ, USA:

John Wiley & Sons, Inc.; 2007. pp. 1.17.1-1.17.8.

doi:10.1002/0471142727.mb0117s79

44. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification,

enrichment, pre-fractionation and storage of peptides for proteomics

using StageTips. Nat Protoc. 2007;2: 1896–906.

doi:10.1038/nprot.2007.261

45. Tyanova S, Temu T, Carlson A, Sinitcyn P, Mann M, Cox J. Visualization of

LC-MS/MS proteomics data in MaxQuant. Proteomics. 2018;15: 1453–

1456. doi:10.1002/pmic.201400449

46. Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool

(REST) for group-wise comparison and statistical analysis of relative

expression results in real-time PCR. Nucleic Acids Res. 2002;30: e36.

Available: http://www.pubmedcentral.nih.gov/articlerender.fcgi?

artid=113859&tool=pmcentrez&rendertype=abstract

47. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al.

The MIQE guidelines: minimum information for publication of quantitative

real-time PCR experiments. Clin Chem. 2009;55: 611–22.

doi:10.1373/clinchem.2008.112797

48. Seemann T. Snippy: fast bacterial variant calling from NGS reads

[Internet]. 2015. Available: https://github.com/tseemann/snippy

49. Li H, Durbin R. Fast and accurate short read alignment with Burrows-

Wheeler transform. Bioinformatics. 2009;25: 1754–1760.

doi:10.1093/bioinformatics/btp324

50. Garrison E, Marth G. Haplotype-based variant detection from short-read

sequencing. 2012; Available: https://arxiv.org/pdf/1207.3907.pdf

51. Deatherage DE, Barrick JE. Identification of Mutations in Laboratory-

Evolved Microbes from Next-Generation Sequencing Data Using breseq.

Methods in molecular biology (Clifton, NJ). 2014. pp. 165–188.

doi:10.1007/978-1-4939-0554-6_12

52. Sanders LH, Sudhakaran J, Sutton MD. The GO system prevents ROS-

induced mutagenesis and killing in Pseudomonas aeruginosa. FEMS

Microbiol Lett. 2009/03/18. 2009;294: 89–96. doi:FML1550 [pii]

10.1111/j.1574-6968.2009.01550.x

53.   Rodríguez-Rojas A, Blázquez J. The pseudomonas aeruginosa PfpI gene

plays an antimutator role and provides general stress protection. J

Bacteriol. 2009;191. doi:10.1128/JB.01081-08

54.   R Core Team. R: A language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing; 2015.

Legend to the figures

933
934

Figure 1. Hydrogen peroxide killing curves of *E. coli* MG1655. Panel A shows bacterial sensitivity to the 1 mM of hydrogen peroxide with a strong dose response effect ($p = 2.1 \times 10^{-16}$, DRC model fitted based on maximum likelihood). Panel B shows the priming effect or improved response when cells are treated in advanced with a non-lethal dose (0.1 mM). Curves represent the mean of five independent cultures per time point. Asterisks represents significant differences (Welch's test, one asterisk for $p < 0.05$ and two asterisks for $p < 0.01$). Only comparison between primed and non-primed groups are shown.

Figure 2. During memory decline experiment, the viability of the cells remains unaltered before the addition of the trigger (A). The priming memory declined over time after induction of priming response with H_2O_2 0.1 mM (trigger) (B). Asterisks indicate significant differences between each time-point pair (Welch's test, one asterisk for $p < 0.05$ and two asterisks for $p < 0.01$).

Figure 3. Heatmap of relative protein expression based on quantitative lable-free quantification by liquid chromatography/mass spectrometry (LC-MS). Hierarchical clustering of the intensities was performed by using Euclidean distances between means. Row indicate the fluctuation of protein (by gene names) level at 5 minutes after addition of 0.1 and 1 mM H_2O_2. Intensity ranges of the log2 fold-changes are given from highest intensity (green) to lowest (red). Only the 50 statistically most significant upregulated proteins are shown, taking as a reference the 0.1 mM concentration.

Figure 4. Heatmap of relative protein expression based on quantitative label-
free quantification by liquid chromatography/mass spectrometry (LC-MS).
Hierarchical clustering of the intensities was performed by using Euclidean
distances between means. Row indicate the fluctuation of protein (by gene
names) level at different time points. Intensity ranges of the log2 fold-changes
are given from highest intensity (green) to lowest (red). Only the most 50
statistically significant upregulated proteins at time point T1 (30 minutes after
priming) are represented to follow their fluctuation in the other time points with
30 minutes between them.

Figure 5. Projection of differentially translated proteins 30 minutes after
addition of H_2O_2 on protein-protein interaction network of *E. coli* K12. Pale
green nodes indicate upregulated proteins while pale red represents
downregulated ones. Note that the most critical nodes (DnaK and GloL) of the
protein-protein interaction network are affected. The interactions among nodes
show the wide-proteome impact of H_2O_2.

Figure 6. Sensitivity of a selected set of mutants in relevant proteins of *E. coli*
MG1655 that presented an increased level of expression being part of priming
response. Asterisks represents significant differences between the wild-type
(wt) strain and its derivatives mutants (Welch's test, one asterisk for $p < 0.05$
and two asterisks for $p < 0.01$)

Figure 7. Priming experiments of different mutants of *E. coli* MG1655 in
responsive genes to oxidative stress. Asterisks represents significant
differences (Welch's test, one asterisk for $p < 0.05$ and two asterisks for $p <$
0.01). Only comparison between primed and non-primed groups are shown.

Figure 8. Proportion of population extinctions (20 populations per treatment)
due to exposure to growing concentration of H_2O_2 during experimental
evolution of 40 individual populations. Both, priming and triggering doses were
increased twofold daily up to 32 mM, where total extinction was reached.

Evolvability differs between the two treatments, naïve and primed populations
(Log-rank test, p< 0.01).
993
994 **Figure 9**. Mapping of mutations that are unique to each H_2O_2-evolved
995 population shared ones. The positions indicate the approximate locations of
996 mutations in the *E. coli* MG1655 chromosome relative to the wild-type. Three
997 generic main type of changes are indicated.
998
Figure 10. Mutational spectra. Frequencies of different types of mutations
among evolved populations of both primed and no-primed conditions. The
number of mutations (nucleotide substitutions and indels) is plotted against
respective nucleotide positions within the gene fragment. In this analysis, the
mutations were taken by type regardless of the targets by generating two
frequency datasets to compare frequency by type of mutation of evolved
populations under priming and non-priming regimes. The mutational spectra
are significantly different between the two conditions according to the
spectrum analysis software iMARS [32].
1008
1009
1010 **Supporting Information Legends**
1011
1012 **Figure S1.** Relative gene expression for E. coli MG1655 OxyR regulon selected
1013 genes in treatments with 0.1 mM H_2O_2 vs. non-treated bacteria. Cells were
1014 collected 30 minutes after exposure. Error bars represent the standard error of
1015 the mean of three independent biological replicas, each one being the average
1016 of three technical repetitions.
1017
Figure S2. Boxplot showing the H_2O_2-induced mutant frequency. Naïve and
primed cells (pre-treated with 0.1 mM, 30 minutes in advanced) cultures
challenged with 1 mM, allowed torecover and plated in rifampicin (100 µg/ml).
The basal level of mutagenesis for non pre-treated, non-challenged cells is also
shown. Every sample consisted in five independent replications. Equal letter
indicates significant difference while same express no differences (Welch's

test, p=0.03 for basal level vs primed, p<0.01 for both basal vs naïve and
primed vs naïve).

1026

Table S1. Output table of the proteomic experiment reporting time-lapse
decline of the *E. coli* response to 1 mM H_2O_2 during 30 minutes. Bacteria were
sampled 30, 60, 90, 120 and 150 minutes after removal of the treatment. Each
treatment group consisted of six independent replications and bacteria before
treatment (T0) were used as control. Statistical analysis used student t-test and
false discovery rate for correction of the p-values (data analysis using
Maxquant and Perseus software for label-free quantification of proteins with LC-
MS).

1035

Table S2. Output table of the proteomic experiment reporting *E. coli* response
to H_2O_2 treatment of 0.1 and 1 mM during 5 minutes. Each treatment group
consisted of six independent replications and bacteria before treatment (T0)
were used as control. Statistical analysis used student t-test and false
discovery rate for correction of the p-values (data analysis using Maxquant and
Perseus software for label-free quantification of proteins with LC-MS).

1042

1043   **Table S3.** Transcripts differentially expressed (±2 log2) in the fraction of small
1044   of RNA (<200 nt) during the decay of H_2O_2 response (120 minutes after
1045   removal the treatment).

1046

1047   **Table S4.** Transcripts differentially expressed (± 2.5 log2) in the fraction of
1048   large RNA (>200 nt) during the decay of H_2O_2 response (120 minutes after
1049   removal the treatment).

1050

1051   **Table S5.** Relative gene expression (qPCR) for *E. coli* MG1655 selected
1052   responsive genes after a treatment with 0.1 mM H_2O_2 vs. non-treated bacteria
1053   during 30 minutes.

1054

1055   **Table S6.** Strains used in this work and their relevant phenotypes

1056

1057 **Table S7.** Primers used for amplification of kanamycin insertion of mutants
1058 from the Keio collection which PCR product was used to transfer the mutations
1059 to the *E. coli* MG1655 strain.

1060

1061 **Table S8.** Primers used for relative gene expression quantification by real time
1062 PCR. *E. coli* MG1655 was treated with 0.1 mM H_2O_2 during 30 minutes.

1063

1064

1065

1066

1067

1068

1069

1070

Changes in the 50 most significantly upregulated proteins (by gene names)

Changes in the 50 most significantly upregulated proteins (by gene names)

Non-priming evolved populations

Priming evolved populations

Mutations in common
Only in primed populations

non-synonymous amino acid change
intergenic mutations
frameshift