Supra-threshold perception and neural representation of tones presented in noise in conditions of masking release

Katharina Egger, Torsten Dau[1], and Bastian Epp[1*]

[1] Hearing Systems Section, Department of Health Technology, Technical University of Denmark, Ørsteds Plads, Building 352, 2800 Kgs. Lyngby, Denmark

* Corresponding author

bepp@dtu.dk

Abstract

The neural representation and perceptual salience of tonal signals presented in different noise maskers were investigated. The properties of the maskers and signals were varied such that they produced different amounts of either monaural masking release, binaural masking release, or a combination of both. The signals were then presented at different levels above their corresponding masked thresholds and auditory evoked potentials (AEPs) were measured. It was found that, independent of the masking condition, the amplitude of the P2 component of the AEP was similar for the same stimulus levels above masked threshold, suggesting that both monaural and binaural effects of masking release were represented at the level of P2 generation. The perceptual salience of the signal was evaluated at equal levels above masked threshold using a rating task. In contrast to the electrophysiological findings, the subjective ratings of the perceptual signal salience were less consistent with the signal level above masked threshold and varied strongly across listeners and masking conditions. Overall, the results from the present study suggest that the P2 amplitude of the AEP represents an objective indicator of the audibility of a target signal in the presence of complex acoustic maskers.

Introduction

One major task of the auditory system is to distinguish between different sound sources and to segregate single sources from the acoustic background. For sound segregation and identification, the auditory system utilizes monaural cues, such as spectro-temporal information contained in the sound, as well as binaural cues, such as interaural disparities occurring at the two ears (i.e., interaural level- and time differences). The ability of the auditory system to benefit from these cues is often assessed via the effect of masking of a tonal signal by a noise masker. Many aspects of auditory signal detection in noise can be accounted for by the power spectrum model of masking [1]. This model assumes that the frequency selectivity of the auditory system can be represented as a bank of overlapping band-pass filters and that the detection threshold of a tonal signal in the presence of a noise masker is mainly determined by the signal-to-masker energy ratio at the output of the band-pass filter centered at the signal frequency.

Some masking data, however, cannot be accounted for by the power spectrum model. For example, when the masker exhibits coherent intensity fluctuations across frequency, the masked thresholds of signals presented in noise are lower than predicted by the power spectrum model. This monaural phenomenon of improved signal detectability due to coherent intensity fluctuations of the masker across frequency has been referred to as comodulation masking release (CMR) (for a review, see [2] and [3]). There are two experimental CMR paradigms, the band-widening paradigm and the flanking-band paradigm. In the band-widening paradigm, the masker is spectrally centered at the signal frequency and is either unmodulated or comodulated. The bandwidth-dependent difference in masked threshold between the comodulated and the unmodulated masker conditions is then referred to as CMR, or modulated-unmodulated difference [4]. In the flanking-band paradigm, the masker consists of several narrow bands of noise whereby one masker band is centered at the signal frequency, the signal centered band (SCB), and one or more bands, the flanking bands (FBs), are spectrally separated from the signal frequency (e.g., [5-7]). In this paradigm, CMR has been defined

in two ways: 1) as the difference in masked threshold between the conditions with the SCB alone

and when the SCB and comodulated FBs are presented; or 2) as the difference in masked threshold

for unmodulated versus comodulated masker bands.

A release from masking can also be found when the signal and the masker are presented at

different spatial locations compared to a condition in which both sounds are presented at the same

location. Differences in the spatial location of the signal and the masker lead to differences in

binaural cues between the signal and the masker. These binaural cues can lead to an enhanced

signal detectability which is commonly referred to as binaural masking level difference (BMLD) [8-

9]. Effects of BMLD have typically been investigated in experiments with tonal signals masked by

noise where either the signal or the noise masker contained an interaural phase difference (IPD)

[10]. The BMLD has been defined as the difference in threshold between the diotic condition,

where the signal and the masker are presented in phase across ears, and a dichotic condition, where

either the signal or the masker is presented interaurally out of phase.

The effects of CMR, BMLD, and their combination have commonly been studied at signal levels

close to masked threshold (e.g., [11-16]), whereas their contribution to the perception of signals

above masked threshold (i.e., supra-threshold) has received much less attention. The benefit of

CMR at supra-threshold levels has mainly been discussed in connection to either speech detection

and recognition tasks (e.g., [17-19]) or regarding signal intensity and frequency discrimination tasks

(e.g., [20-21]). In terms of binaural processing, the perceived intensity of a signal, i.e., its loudness,

has primarily been studied in conditions of BMLD at different supra-threshold levels of the signal

in the presence of noise (e.g., [10,22-23]). More recently, Verhey and Heise [24] investigated the

magnitude of the tonal content as well as the partial loudness of a tonal signal in the presence of a

noise masker. In their study, conditions with and without monaural and binaural masking releases at

different signal levels above the respective masked thresholds were used. Verhey and Heise [24]

showed that the masking releases were strongest at low signal levels above masked threshold and

decreased towards higher levels. The above studies were all either focused on monaural processing,

regarding CMR, or on binaural processing, regarding BMLD, whereas the supra-threshold

perception of the combined effect of CMR and BMLD has not yet been investigated. It is therefore

still unknown to what extent the two effects contribute to supra-threshold perception.

Various physiological studies investigated the neural mechanisms underlying the processing and

representation of the auditory cues that lead to a perceptual masking release. A physiological

correlate of CMR in human listeners has been proposed at cortical levels using

electroencephalography (EEG) [25], magnetoencephalography (MEG) [26], as well as functional

magnetic resonance imaging (fMRI [27] [28]). Physiological correlates of BMLD in humans have

also been investigated at different levels along the auditory pathway using EEG [29-31], MEG [32-

33], and fMRI [34].

Billings et al. [35] compared late auditory potentials evoked by a speech syllable presented in

speech-shaped noise with the performance in a behavioral sentence identification task at different

signal-to-masker ratios. They identified the N1 component of the electrophysiological response as a

correlate of behavioral performance. However, while the N1 component was found to be mainly

dependent on the signal-to-masker ratio, the behavioral data showed a signal-level dependent

behavior, complicating the relation between behavioral performance and evoked potentials

amplitude.

Epp et al. [36] recorded evoked potentials of masked tonal signals in conditions of masking

release at different signal-to-masker ratios. At a fixed signal-to-masker ratio, the level of the signal

above masked threshold was altered by the introduction of monaural across-frequency cues to the

masker (yielding CMR), binaural cues to the tonal signal (yielding BMLD), or a combination of

both. Epp et al. [36] found a combined effect of CMR and BMLD reflected in the N1/P2 complex

of the evoked potential. The N1 component was sensitive to the introduction of binaural cues, but

did not show any sensitivity to monaural across-frequency cues. In contrast, the amplitude of the P2

component at a given signal-to-masker ratio was found to be larger in conditions with comodulated

noise than in conditions with uncorrelated noise. A larger P2 amplitude was also observed when

binaural cues were introduced to the tonal signal as well as when both cues were present. In fact, the P2 amplitude was of similar size in experimental conditions where the signal was presented at a similar level above its masked threshold. This was not the case for the N1 component, nor the P2-N1 difference amplitude. Epp et al. [36] therefore suggested that the P2 component might be a potential candidate for the neural representation of the partially masked signal at the corresponding level above masked threshold.

The hypothesis of the present study was that signals at the same level above masked threshold evoke the same P2 amplitudes in the corresponding evoked potential. It was further postulated that there exists a perceptual correlate of the signal level above masked threshold, the "perceptual salience", and that signals at the same level above masked threshold evoke the same perceptual salience. To test this, the present study extended the investigations of Epp et al. [36] and considered three experiments: In the first experiment, behavioral detection thresholds of masked tonal signals were obtained in conditions reflecting either purely monaural, purely binaural, or a combination of monaural and binaural masking releases. In a second experiment, AEPs for the masked signals were recorded in the same listeners at individually adjusted signal levels corresponding to equal levels above their respective masked thresholds. Finally, in a third experiment, the listeners rated the perceptual salience of the signal embedded in the noise masker at equal signal levels above their respective masked thresholds.

Materials and methods

Stimuli

The signal was a 700 Hz tone of 300 ms duration, including 20 ms raised-cosine on- and offset

ramps. The signal was presented either with the same interaural phase, i.e., an IPD of 0 degree

(diotic condition), or interaurally out of phase (dichotic condition) with an IPD of 150 degree[1].

The masker consisted of either five narrowband noise bands ("multi-band conditions") or one

broadband noise ("broadband condition"). In the multi-band conditions, each noise band was 24 Hz

wide and had a sound pressure level (SPL) of 50 dB. One noise band (SCB) was centered at the

signal frequency (700 Hz) and the other four bands, the FBs, were arranged symmetrically (on a

linear frequency scale) with respect to the SCB at 300, 400, 1000, and 1100 Hz. The remote spectral

distance between the SCB and the FBs was chosen in order to reduce within-channel cues

contributing to CMR (e.g., Cohen, 1991). The noise bands had either uncorrelated or comodulated

intensity fluctuations across frequency, subsequently denoted as UN and CM masker, respectively.

The broadband (BB) masker was an 824-Hz wide noise band centered at the signal frequency (700

Hz) and had an overall level of 60 dB SPL. This level was chosen since it produces approximately

the same amount of energy that passes the auditory filter centered at the signal frequency as the

SCB in the multi-band conditions [37]. Hence, a similar masking effect produced by the BB masker

was expected as for the UN masker.

All maskers were presented diotically and had a duration of 900 ms, including 20 ms raised-

cosine on- and offset ramps. The signal was presented in the last 300 ms of the masker. This was

done to avoid a temporal overlap of the masker onset response and the signal onset response in the

AEPs. The signal in both binaural conditions was combined with each of the maskers resulting in

six different stimulus conditions: the diotic signal presented in either uncorrelated noise (UN_0),

---

[1]   An IPD of 150 degrees was encouraged by the maximum IPD that human listeners can perceive due to the limitation

      of the circumference of the head, based on the average maximum, nonambiguous IPD calculated out of head-related

      transfer functions (HRTFs) of 68 listeners (HRTF database of the Acoustics Research Institute, Austrian Academy

      of Sciences)

comodulated noise (CM_0), or broadband noise (BB_0) and the dichotic signal presented in the same

masker types (UN_{150}, CM_{150}, and BB_{150}, respectively).

The noise bands were generated by multiplying a random-phase sinusoidal carrier at the desired

frequency with a low-pass noise without a DC component. The low-pass noise was generated in the

frequency domain by assigning numbers between ±0.5 from a uniformly distributed random process

to the real and imaginary parts of the respective frequency components. For the UN and BB

maskers, independent realizations of the low-pass noise were used for each masker band, while the

same low-pass noise was used for all five bands of the CM masker. The masking noise was newly

generated for each interval and each trial.

Apparatus

All stimuli were digitally generated in MATLAB with a sampling rate of 44100 Hz and a 16-bit

resolution, converted from digital to analogue (RME DIGI96/8 PAD, experiments 1 and 3; RME

Fireface UCX, experiment 2) and presented via circumaural headphones (Sennheiser HD580,

experiments 1 and 3) or via in-ear headphones (Etymotic ER-2, experiment 2). In experiment 2, the

Psychophysics Toolbox extensions for MATLAB [38-39] were used. The headphones were

calibrated and equalized at the signal frequency of the tone.

AEPs were recorded using a BioSemi ActiveTwo measurement system. The listeners wore an

elastic cap with plastic electrode holders for 64 sintered Ag/AgCl pin-type electrodes. The electrode

holders were filled with highly conductive, Signa electrode gel to reduce the contact impedance

between electrode and skin. The common reference was the electrode placed at the left mastoid (P9),

according to the extended 10/20 layout as standardized by the American Electroencephalographic

Society (e.g., [40-41]). The electrodes were connected to the ActiveTwo AD-box, which amplified

and performed A/D conversion of the measured potentials with a sampling rate of 1024 Hz and a

24-bit resolution. The potentials were recorded with the data acquisition software ActiView

(version 6.05), which streamed the continuous EEG to hard disk. During the recordings, an anti-

aliasing digital low-pass filter with a cut-off frequency of 200 Hz was used together with a digital high-pass filter with a cut-off frequency of 0.16 Hz to reduce the influence of slow, non-neural potentials, such as skin potentials [42].

Listeners

Eight listeners (three female, five male), aged between 22 and 28 years, participated in the experiments. None of them reported any history of hearing impairment. All listeners had pure-tone hearing thresholds within 15 dB HL for the standard audiometric frequencies from 125 to 4000 Hz. All listeners were paid an hourly wage for their participation. The listeners were the same in all three experiments. During the psychoacoustical experiments, the listeners were seated in a double-walled, sound-attenuating booth. The AEP experiment was carried out in a double-walled and electrically shielded, sound-attenuating booth.

Procedure

In the first experiment, detection thresholds of the masked signals were measured for each listener in all six stimulus conditions (UN_0, CM_0, BB_0, UN_{150}, CM_{150}, and BB_{150}). An adaptive, three interval, three-alternative forced-choice procedure with visual feedback was used. The intervals within a trial were separated by pauses of 500 ms. The listeners had to indicate the interval in which the signal was presented by pressing the corresponding key on the keyboard. The adaptive signal level adjustment followed a one-up two-down algorithm to estimate the 70.7% point of the psychometric function [43]. The initial step size was 8 dB. After each lower reversal, the step size was halved until it reached the minimum step size of 1 dB. This step size was kept constant for another six reversals of which the arithmetic mean was calculated and used as the estimated threshold for that run. Each listener performed four threshold measurements per stimulus condition whereof the arithmetic mean of the last three runs was taken as the final individual threshold estimate. The different stimulus conditions were presented in random order within blocks, where each condition occurred once.

In the second experiment, late AEPs for the masked signals were recorded for all stimulus

conditions. The signal levels were adjusted to 10, 15, 20, and 25 dB above the masked thresholds of

the individual listener (obtained in experiment 1). The stimuli were presented in random order,

evenly distributed over 400 sweeps (i.e., 400 repetitions per stimulus condition and level). The

presentations were separated by an inter-stimulus interval of 300 ms plus an additional jitter of 250

to 350 ms. During the EEG recordings, the listeners were presented a quiet movie of their choice

with subtitles on a low-radiation screen. The listeners were asked to relax but not to fall asleep and

to avoid moving as far as possible. The experiment was divided in six blocks of approximately 35

minutes each, whereof two blocks were recorded in one session. The listeners participated in three

sessions distributed over different days.

In the third experiment, the listeners rated the perceptual salience of the tone embedded in the

masking noise for all stimulus conditions by magnitude estimation. The signal levels were

individually-adjusted to -10, 0, 5, 10, 15, and 20 dB relative to the masked thresholds. The test

signals to be judged (i.e., the tones embedded in masking noise) were presented in paired trials with

a fixed reference which was a diotic 700-Hz tone presented at 70 dB SPL. In each trial, the

reference tone and the test signal could be played as often as the listeners desired and were accessed

either by pressing the corresponding keys or by clicking the corresponding buttons with the mouse.

The number of times the reference and the test signal were presented in each trial was recorded. The

listeners had to rate the salience of the tone in the test signal on an endpoint-anchored scale by using

a continuous slider. The scale was anchored by 'not audible' on the lower end and by 'reference' on

the upper end. These anchors were combined with a scale without labels in between, as suggested in

[44]. The slider's position was mapped onto a numerical scale between 0 (designating the lower

anchor 'not audible') and 10 (designating the upper anchor 'reference'). The listeners were not aware

of the mapping. Prior to the experiment, the listeners were told that the experiment contained the

same types of stimuli they had been listening to in the previously performed measurements.

To familiarize the listeners with the task, a short training session was provided before the initial

run of the experiment, using stimulus examples of varying perceptual saliences. Four different types

of stimulus examples were presented: (1) the reference tone; (2) the masking noise alone, randomly

representing either the UN, CM, or BB masker; (3) the masked signal in one stimulus condition

randomly selected out of all six (UN_0, CM_0, BB_0, UN_{150}, CM_{150}, or BB_{150}), at a supra-threshold

level, corresponding to 25 dB above masked threshold; and (4) randomly selected stimuli of all

stimulus conditions with signal levels as used in the experiment.

The stimuli were tested in random order within blocks, where each level for each stimulus

condition occurred once. Sixteen blocks were presented, whereof the very first block was

considered as additional training and excluded from further data analysis. The individual salience

ratings were therefore derived from fifteen responses per stimulus condition and level.

Data analysis

The continuous EEG data were processed off-line in MATLAB to extract the late AEPs. Single

sweeps of a time interval of 1200 ms (150 ms pre-stimulus period and 1050 ms following the

stimulus onset) were extracted corresponding to the different stimulus conditions and levels. For

each sweep, a linear fit was subtracted from the waveform to remove low-frequency potential drifts

(e.g., [45]). The fit was based on the intervals 150 ms before the signal onset (at 600 ms) and 150

ms after the stimulus. The data were then digitally filtered with a zero-phase, second order forward-

backward Butterworth low-pass filter with a cut-off frequency of 20 Hz. Each sweep was baseline

corrected by subtracting the arithmetic mean over the 150 ms pre-stimulus period. An iterative

weighted averaging method was used to average all single sweeps per stimulus condition and level

for each listener [45]. The weighted average was computed by weighting each single sweep by the

inverse power of its noise [46]. Single sweeps which exceeded the artefact rejection threshold of

±100 μV in any of the recorded channels were discarded and not included in the average. This

resulted in the average AEP including the response to the masker onset.

To obtain the neural response following the signal onset within the masker for each listener,

referred to as the AEP change complex, the AEP in the latency interval between 600 and 1050 ms

was extracted from the averaged sweeps. Baseline correction was applied considering a 150 ms pre-

stimulus period (with respect to the signal onset at 600 ms). The grand mean AEP change complex

was computed as an arithmetic mean over all individual AEPs.

In addition, the amplitude components of the AEP change complex were evaluated individually

for each listener. The change complex can be described by the response extrema N1, measured

between 90 and 190 ms, and P2, measured between 180 and 290 ms (with respect to the signal

onset). Peak amplitudes for the N1 and P2 components were extracted for the electrode placed at

the vertex (Cz), individually for each listener. Electrode Cz was used since it showed the largest

response averaged across all conditions. Peak amplitudes were extracted separately for each

stimulus condition and level, using a semi-automatic procedure including identification of N1 and

P2 components by a peak-scan-algorithm and subsequent verification by visual inspection of all

waveforms. The algorithm located minima and maxima in the time windows defined for N1 and P2,

respectively. The peaks were indicated automatically if the first derivative of the potential showed a

zero-crossing and had a change in sign within the time window. If multiple extrema were found, the

algorithm selected the ones with the largest amplitude. If an extremum with the wrong polarity or

no response maximum was found by the algorithm, peaks were selected manually by requiring a

clear peak within the time window with correct curvature.

The behavioral salience data were analyzed in MATLAB using a non-parametric Friedman test.

The effects of level and stimulus condition on the rated signal salience were tested. The analysis

was based on median salience ratings across all listeners. Since no interaction effects can be tested

using the Friedman test, two separate tests were conducted. To test the effects of level, the data

obtained at the different levels were arranged in columns and the stimulus conditions were

represented along the rows of the data matrix. To test the effects of stimulus condition, the matrix

was transposed before testing.

Results & Discussion

Experiment 1: Masked signal detection thresholds in noise

Masked thresholds for the eight individual listeners as a function of the signal IPD are shown in Fig 1 (A-H). The error bars in each panel indicate the standard deviations of the thresholds. The bottom right panel shows the grand average thresholds across the listeners. The results are shown for the UN masker (squares), the CM masker (triangles), and the BB masker (circles). For all masker types, the thresholds obtained in the dichotic conditions (150 degree IPD) were lower than those obtained in the diotic conditions (0 degree IPD), reflecting a BMLD. All listeners, except for the one represented in panel A, showed a decrease in threshold when the masker envelope across frequency was changed from being uncorrelated (UN masker) to comodulated (CM masker), reflecting a CMR effect. This was the case in both binaural conditions. For listener A, the difference in masked threshold between the UN and the CM maskers, i.e., the effect of CMR, nearly vanished in the dichotic condition. The thresholds obtained with the signal presented in the BB masker were similar to those obtained with the UN masker, except for listener A, who showed a considerably lower threshold for the BB masker in the diotic condition than all other listeners. The amount of BMLD obtained for the different masker types differed across the listeners. The BMLD ranged from 8.7 dB (listener H) to 20.4 dB (listener A) for the UN masker, from 7.4 dB (listener H) to 14.6 dB (listener F) for the CM masker, and from 8 dB (listener A) to 13 dB (listeners B and D) for the BB masker. The amount of CMR decreased with increasing IPD for half of the listeners (A, C, D, and E). Three listeners (B, G, and H) showed roughly the same amount of CMR for the two signal IPDs; listener F showed a larger CMR in the dichotic condition compared to the diotic condition. The diotic CMR varied from 9 dB (listener E) to 12 dB (listener C) and the dichotic CMR varied from 1.2 dB (listener A) to 12.4 dB (listener F).

Fig 1. Mean thresholds. Individual mean thresholds of the listeners (panels A-H) and grand

average thresholds across all listeners (bottom right panel) as a function of signal IPD. Thresholds

are plotted for UN (squares), CM (triangles), and BB (circles) maskers. Error bars indicate plus-

minus one standard deviation.

Regarding the average data, a benefit was observed by introducing masker comodulation

(yielding CMR), a signal IPD (yielding BMLD), or a combination of both. The thresholds were

largest for UN_0 (48.9 dB) and smallest for CM_{150} (26.7 dB). CMR reached a magnitude of 10.7 dB

in the diotic condition and decreased to 8.4 dB in the dichotic condition. Thresholds were lower in

the dichotic conditions than in the diotic conditions, demonstrating a BMLD of 13.9 dB in the case

of the UN masker and approximately 11.5 dB in the cases of the CM and BB maskers.

The data are consistent with findings reported in the literature [13,16,36]. Interestingly, also in

Epp and Verhey [13] and Hall et al. [16], a few listeners showed a strong reduction of CMR in the

dichotic relative to the diotic condition in combination with large BMLDs, whereas the majority of

the listeners showed a smaller BMLD and a comparable CMR in the diotic and dichotic conditions.

Experiment 2: Auditory-evoked potentials to tones in conditions of masking release

The grand mean AEPs across all listeners, for the latency interval of 550 to 950 ms relative to the

stimulus onset, i.e., the interval associated with the AEP change complex, is shown in Fig 2. The

three panels illustrate the results obtained with the diotic signals (solid lines) and the dichotic

signals (dashed lines) in the presence of the UN masker (left panel), the CM masker (middle panel),

and the BB masker (right panel), respectively. The ordinate represents different levels of the signal

above the individual listeners' masked thresholds. Independent of the masker condition (UN, CM,

and BB) and the binaural condition (0 versus 150 degree IPD), the magnitude of the AEP change

complex increased with increasing signal level relative to masked threshold (bottom to top in Fig 2).

This is consistent with the initial hypothesis of this study that a higher signal level above masked

threshold yields a stronger neural response. However, for a given level above masked threshold, the

grand mean AEP magnitude was larger in the dichotic (dashed) than in the diotic (solid) conditions.

Hence, the grand mean data, when ignoring possible individual differences in latency of the AEP, is

at odds with the initial hypothesis of this study that signals presented at equal levels above masked

threshold evoke the same AEP magnitude, independent of the stimulus condition.

Fig 2. Grand mean AEP. Grand mean evoked potential data for all listeners for a latency

interval of 550 to 950 ms relative to the stimulus onset. The three panels illustrate the averaged

AEPs measured for diotic (solid lines) and dichotic (dashed lines) signals in the presence of UN

(left), CM (middle), and BB (right) maskers. Waveforms are plotted for four different signal levels:

10, 15, 20, and 25 dB relative to masked thresholds (indicated on the y-axes on the left-hand side).

The potentials were baseline corrected considering a 150 ms pre-stimulus period (with respect to the

signal onset at 600 ms).

The results shown in Fig 2 are compatible with the data obtained in the study of [47] who

measured AEPs evoked by a diotic or antiphasic tone in a broadband masker. In their study, larger

responses were found for antiphasic than for diotic signals at a fixed level above masked threshold.

However, only the N1-P2 peak of the change complex was evaluated and only three listeners were

considered. The grand average AEP data show also a similar trend as the results obtained in Epp et

al. [36], where larger grand average potentials were obtained for dichotic than for diotic signals. In

the study by Epp et al. [36], the signal was however presented at equal SPLs rather than at equal

levels above masked threshold.

The grand average AEPs, however, do not account for individual differences in latency across

listeners which might challenge the interpretation of the grand mean AEP. To compensate for

individual differences in AEP latency, the peak amplitudes of the change complex were extracted

and analyzed individually for each listener[2]. The grand mean of the extracted N1 (left panel), P2

(middle panel), and P2-N1 (right panel) amplitudes as a function of the signal level relative to the

individual listeners' masked thresholds are shown in Fig 3. The open symbols represent the

stimulus conditions with the diotic signal in the presence of the UN (squares), CM (triangles), and

BB (circles) maskers, respectively. The filled symbols represent the corresponding stimulus

conditions with the dichotic signal. The magnitude of the N1 (negativity) increased with signal level.

For a given level, the mean N1 amplitude varied between 1.1 and 2.2 µV across the stimulus

conditions. The magnitude of the P2 (positivity) also increased with signal level but showed a

markedly smaller amount of variability (of 0.5 to 1.1 µV) across stimulus conditions than N1. The

difference amplitude, P2-N1, often considered in AEP studies, showed variability of up to 3 µV.

Fig 3. Mean amplitudes of AEP. Grand mean of amplitudes N1 (left panel), P2 (middle panel),

and the difference between P2 and N1 (right panel) across all listeners as a function of signal level

relative to masked threshold. Amplitudes are plotted for diotic (open symbols) and dichotic (filled

symbols) signals in the presence of UN (squares), CM (triangles), or BB (circles) maskers. Error

bars in the lower, left corner indicate plus-minus one standard error (averaged over all signal levels)

for each stimulus condition, respectively.

The P2 amplitudes for both binaural conditions were found to be similar at equal levels above

masked thresholds (Fig 3, middle panel), in agreement with the initial hypothesis of this study.

---

[2]   Even though the latency was shown to represent an informative dimension in the evaluation of AEPs (Billings et al.,
2009), this dimension is closely correlated to the amplitude of the change complex component and shows the same
or an even larger variability.

Hence, individual differences in latency resulted in a broadening of the grand mean AEP peaks and

therefore in different amplitudes for different conditions in Fig 2. The N1 amplitude (and, hence,

the P2-N1 difference) showed a large variability across stimulus conditions and, in contrast to the

P2 amplitude, no clear pattern between N1 amplitude and condition was found.

In summary, the present results support the hypothesis that the P2 component of the AEP change

complex correlates with the signal level above masked threshold. Specifically, P2 may represent an

objective indicator of the overall perceptual masking release, obtained here with monaural and

binaural signal-masker configurations.

Experiment 3: Perceptual salience ratings of tones in conditions of masking release

Panels A-H of Fig 4 show the individual salience ratings of the signal as a function of the signal

level relative to the masked threshold. Since the salience scale had fixed endpoints, the data are

represented by the median and the corresponding interquartile ranges. The open symbols represent

the results for the diotic conditions, whereas the filled symbols indicate the results for the dichotic

conditions obtained with the UN masker (squares), the CM masker (triangles), and the BB maskers

(circles), respectively. For all listeners, the salience ratings increased with increasing signal level in

each stimulus condition. The salience tended to be highest for stimulus conditions without a release

from masking (UN_0 and BB_0). Two listeners (F, G) rated the salience of the dichotic tone embedded

in the CM masker (CM_{150}) clearly lower than in all other stimulus conditions. Some listeners (A, C,

D, and H) showed a similar increase in salience with increasing signal level relative to masked

threshold, particularly for the CM masker conditions. Other listeners (B, E, F, and G) showed

ratings that are ordered according to their physical signal-to-masker level ratio rather than according

to the signal level relative to masked threshold. The salience for signals that were presented below

masked threshold (i.e., -10 dB SL) nearly vanished for all but listeners A and C.

A non-parametric Friedman test was used on the pooled data to investigate the effects of level

and stimulus condition on the rated salience. The test provided a significant main effect of level

[$chi^2(5) = 224.03$, $p < 0.001$] and of stimulus condition [$chi^2(5) = 71.03$, $p < 0.001$]. The significant

effect of stimulus condition is contrary to the initial hypothesis of this study that signals at the same

level above masked threshold evoke the same perceptual salience for all stimulus conditions.

Fig 4. Individual salience ratings. Individual median salience ratings of the individual listeners

(panels A-H) as a function of masked threshold. Salience is plotted for diotic (open symbols) and

dichotic (filled symbols) signals in the presence of UN (squares), CM (triangles), and BB (circles)

maskers. Error bars indicate the interquartile range (panels A-H). PC_r and PC_t refer to the average

number of times the reference and the test signal were presented in each trial, respectively.

The variability across listeners observed in these data suggests that different perceptual attributes

might have influenced the salience rating. The perception of a tone in the presence of a background

noise has been discussed mainly in connection to perceptual attributes like tonality, tonalness, pitch

strength, and partial loudness, or in relation to measures of annoyance (for a review, see [48]).

Effects of monaural and/or binaural masking releases on tone perception have not been considered

in this context yet. The order of the ratings at a given signal level (above threshold) in one of the

listener groups (B, E, F, G) suggests that the partial loudness of the signal might have contributed to

the judgments.

Relation between auditory evoked potential amplitude and salience rating

The salience ratings increased monotonically with increasing signal level relative to masked

threshold. However, for a given level above masked threshold, the rated salience differed

considerably across stimulus conditions and showed a large variability across listeners. These

results are not consistent with the AEP data for the P2 component and, thus, do not support the

initial hypothesis of this study that masked signals evoking similar P2 amplitudes result in the same perceptual salience of the signal. The statistical significance of stimulus condition further provides evidence against the validity of this hypothesis.

The observed mismatch between the salience ratings and the electrophysiological responses using the measurement paradigm chosen in the present study may be caused by the dominance of certain perceptual attributes in the salience rating. The individual contributing factors, such as partial loudness or tonalness, would need to be investigated further by isolating them separately in additional perceptual experiments.

An attempt to relate partial loudness to electrophysiological data was made by Rupp et al. [49]. They measured auditory evoked fields (AEFs) in response to stimulation with tone pulses in the presence of Schroeder-phase tone-complex maskers and compared the AEFs with patterns of a loudness experiment obtained with the same listeners. While the tone pulses were kept at a constant sound pressure level, the temporal position of the pulses within the maskers was varied, resulting in varying neuromagnetic response amplitudes and a varying partial loudness of the tone pulses. The behavioral measure in Rupp et al. [49] was a two-alternative-forced choice paired-comparison of the stimuli. The paired comparisons were then projected on a perceptual ratio scale using the Bradley-Terry-Luce (BTL) method [50]. Rupp et al. [49] found a correlation between the AEFs and the partial loudness patterns. In particular, in those conditions where the tone pulses were masked more effectively by the tone-complex maskers, the corresponding AEF showed a smaller amplitude than in the conditions where the signal was masked less effectively. The BTL method has also been used to measure other perceptual attributes of sounds like annoyance with subsequent analysis of other contributing factors [51]. Since loudness most likely was a contributing factor to the perceptual salience of the signal in the present study, a similar approach might help to clarify the observed disparity between AEP amplitude and salience rating.

Summary and conclusions

The present study investigated the neural representation and perceptual salience of tonal signals

presented in different noise maskers. The maskers and the target signals were chosen such that they

produced different amounts of either monaural masking release (CMR), binaural masking release

(BMLD), or a combination of both. The signals were then presented at different levels above their

respective masked thresholds. AEPs were used as an objective indicator of the signal's neural

representation in the maskers. It was found that signals presented at equal levels above the listeners'

individual masked threshold (but at different physical intensities) resulted in similar amplitudes of

the P2 component (but different amplitudes of the N1 component) across various conditions of

masking release. This suggests that both monaural and binaural effects of perceptual masking

release are represented at the level of P2 generation. In contrast to the electrophysiological findings,

the subjective ratings of the signal salience showed a significant effect of stimulus condition. Hence,

the perceptual salience as considered in the present study did not show a consistent correlation with

the corresponding P2 amplitude patterns. The reason for this might be the contribution of several

other perceptual cues to the salience ratings, with relative weights that may vary across listeners.

Possibly, the partial loudness and the magnitude of tonal content of the supra-threshold tone

embedded in the complex masker contributed to the perceived salience of the signal.

Overall, if the correlation of the P2 amplitude of the AEP change complex with the level above

masked threshold also holds true for other conditions of masking release and more complex signals

like speech, P2 might serve as an indicator of the supra-threshold signal audibility in the presence of

a complex masker in various conditions of perceptual masking.

Acknowledgments

463   We thank all our listeners for their participation in this study.

References

[1] H. Fletcher, Auditory Patterns, Rev Mod Phys, vol. 12, no. 1, pp. 47-65, 1 1940.

[2] J. W. Hall, M. P. Haggard and M. A. Fernandes, Detection in noise by spectro-temporal pattern

analysis, The Journal of the Acoustical Society of America, 2005.

[3] J. L. Verhey, D. Pressnitzer and I. M. Winter, The psychophysics and physiology of comodulation

masking release., Exp Brain Res, vol. 153, no. 4, pp. 405-17, 12 2003.

[4] R. P. Carlyon, S. Buus and M. Florentine, Comodulation masking release for three types of modulator

as a function of modulation rate, Hearing Research, 1989.

[5] J. W. Hall, M. P. Haggard and M. A. Fernandes, Detection in noise by spectro-temporal pattern

analysis, The Journal of the Acoustical Society of America, vol. 76, no. 1, pp. 50-56, 1984.

[6] J. W. Hall, J. H. Grose and M. P. Haggard, Effects of flanking band proximity, number, and modulation

pattern on comodulation masking release, The Journal of the Acoustical Society of America,

1990.

[7] G. P. Schooneveldt and B. C. J. Moore, Comodulation masking release (CMR): Effects of signal

frequency, flanking-band frequency, masker bandwidth, flanking-band level, and monotic

versus dichotic presentation of the flanking band, The Journal of the Acoustical Society of

America, 1987.

[8] L. A. Jeffress, H. C. Blodgett, T. T. Sandel and C. L. I. Wood, Masking of tonal signals, The Journal of

the Acoustical Society of America, vol. 28, no. 3, pp. 416-426, 1956.

[9] S. van de Par and A. Kohlrausch, Dependence of binaural masking level differences on center

frequency, masker bandwidth, and interaural parameters., The Journal of the Acoustical Society

of America, vol. 106, no. 4 Pt 1, pp. 1940-7, 10 1999.

[10] I. Hirsh and I. Pollack, The role of interaural phase in loudness, The Journal of the Acoustical Society of

America, vol. 20, no. 6, pp. 761-766, 1948.

[11] M. F. Cohen and E. D. Schubert, Comodulation masking release and the masking-level difference, The Journal of the Acoustical Society of America, vol. 89, no. 6, pp. 3007-3008, 1991.

[12] B. Epp and J. L. Verhey, Superposition of masking releases., J Comput Neurosci, vol. 26, no. 3, pp. 393-407, 6 2009.

[13] B. Epp and J. L. Verhey, Combination of masking releases for different center frequencies and masker amplitude statistics., The Journal of the Acoustical Society of America, vol. 126, no. 5, pp. 2479-89, 11 2009.

[14] J. W. Hall, J. A. Cokely and J. H. Grose, Combined monaural and binaural masking release, The Journal of the Acoustical Society of America, vol. 83, pp. 1839-1845, 1988.

[15] J. W. Hall, E. Buss and J. H. Grose, Binaural comodulation masking release: Effects of masker interaural correlation, The Journal of the Acoustical Society of America, 2006.

[16] J. W. Hall, E. Buss and J. H. Grose, Exploring the additivity of binaural and monaural masking release., The Journal of the Acoustical Society of America, vol. 129, no. 4, pp. 2080-7, 4 2011.

[17] J. M. Festen, Contributions of comodulation masking release and temporal resolution to the speech-reception threshold masked by an interfering voice, The Journal of the Acoustical Society of America, no. 94, pp. 1295-1300, 1993.

[18] J. H. Grose and J. W. Hall, Comodulation masking release for speech stimuli, The Journal of the Acoustical Society of America, no. 91, pp. 1042-1050, 1992.

[19] B. J. Kwon, Co-modulation masking release in consonant recognition, The Journal of the Acoustical Society of America, vol. 112, no. 2, pp. 634-41, 11 2002.

[20] E. Buss, J. H. Grose and J. W. Hall, Frequency discrimination under conditions of comodulation masking release (L), The Journal of the Acoustical Society of America, 2012.

[21] E. Buss and J. W. Hall, Effects of masker envelope coherence on intensity discrimination, The Journal of the Acoustical Society of America, 2009.

[22] R. D. Soderquist, D. R. and Shilling, Loudness and the binaural masking level difference, Bull. Psychon. Soc., no. 28, pp. 553-555, 1990.

[23] T. H. Townsend and D. P. Goldstein, Suprathreshold binaural unmasking, The Journal of the Acoustical Society of America, no. December, pp. 621-624, 1972.

[24] J. L. Verhey and S. J. Heise, Suprathreshold Perception of Tonal Components in Noise Under Conditions of Masking Release, Acta Acustica united with Acustica, pp. 1-24, 2012.

[25] A. G. Androulidakis and S. J. Jones, Detection of signals in modulated and unmodulated noise observed using auditory evoked potentials., Clin Neurophysiol, vol. 117, no. 8, pp. 1783-93, 8 2006.

[26] I. Rupp, Andre Las, Liora Nelken, Neuromagnetic representation of comodulation masking release in the human auditory cortex, in Hearing – From Sensory Processing to Perception, B. Kollmeier, G. Klump, V. Hohmann, U. Langemann, M. Mauermann, S. Uppenkamp and J. Verhey, Eds., Berlin, Heidelberg, Springer Berlin Heidelberg, 2007, pp. 125-132.

[27] S. M. A. Ernst, J. L. Verhey and S. Uppenkamp, Spatial dissociation of changes of level and signal-to-noise ratio in auditory cortex for tones in noise., NeuroImage, vol. 43, no. 2, pp. 321-8, 1 11 2008.

[28] S. M. A. Ernst, S. Uppenkamp and J. L. Verhey, Cortical representation of release from auditory masking., NeuroImage, vol. 49, no. 1, pp. 835-42, 1 1 2010.

[29] C. Fowler and C. Mikami, Phase effects on the middle and late auditory evoked potentials., Journal of the American Academy of …, vol. 7, pp. 23-30, 1996.

[30] I. M. Ishida and D. R. Stapells, Does the 40-Hz auditory steady-state response show the binaural masking level difference?, Ear and hearing, vol. 30, no. 6, pp. 713-5, 12 2009.

[31] W. Y. S. Wong and D. R. Stapells, Brain stem and cortical mechanisms underlying the binaural masking level difference in humans: an auditory steady-state response study., Ear and hearing, vol. 25, no. 1, pp. 57-67, 3 2004.

[32] B. Ross, T. Fujioka, K. L. Tremblay and T. W. Picton, Aging in binaural hearing begins in mid-life: Evidence from cortical auditory-evoked responses to changes in interaural phase., J Neurosci, vol. 27, no. 42, pp. 11172-8, 17 10 2007.

[33] T. Sasaki, T. Kawase, N. Nakasato, A. Kanno, M. Ogura, T. Tominaga and T. Kobayashi, Neuromagnetic evaluation of binaural unmasking., NeuroImage, vol. 25, no. 3, pp. 684-9, 15 4

2005.

[34] D. S. Wack, J. L. Cox, C. V. Schirda, C. R. Magnano, J. E. Sussman, D. Henderson and R. F. Burkard, Functional Anatomy of the Masking Level Difference, an fMRI Study, PLoS ONE, vol. 7, no. 7, p. e41263, 27 7 2012.

[35] C. J. Billings, G. P. McMillan, T. M. Penman and S. M. Gille, Predicting perception in noise using cortical auditory evoked potentials, JARO - Journal of the Association for Research in Otolaryngology, vol. 14, pp. 891-903, 2013.

[36] B. Epp, I. Yasin and J. L. Verhey, Objective measures of binaural masking level differences and comodulation masking release based on late auditory evoked potentials., Hearing research, vol. 306, pp. 21-8, 12 2013.

[37] B. R. Glasberg and B. C. Moore, Derivation of auditory filter shapes from notched-noise data., Hear Res, vol. 47, no. 1-2, pp. 103-38, 1 8 1990.

[38] D. H. Brainard, The Psychophysics Toolbox, Spatial Vision, 1997.

[39] D. Kleiner, M., Brainard, D., and Pelli, What's new in Psychtoolbox-3?, in Percept. ECVP Abstr. Suppl., 2007.

[40] H. H. Jasper, The ten twenty electrode system of the international federation., Electroen Clin Neuro, vol. 10 (append, pp. 371-375, 1957.

[41] F. Sharbrough, G. Chatrian, R. Lesser, H. Lüders, M. Nuwer and T. Picton, American Electroencephalographic Society guidelines for standard electrode position nomenclature., J Clin Neurophysiol, vol. 8, no. 2, pp. 200-202, 1991.

[42] S. J. (. J. Luck, An Introduction to the Event-Related Potential Technique, second edition, 2014.

[43] H. Levitt, Transformed up-down methods in psychoacoustics, The Journal of the Acoustical society of America, pp. 467-477, 1971.

[44] R. Guski, Psychological Methods for Evaluating Sound Quality and Assessing Acoustic Information, Acta Acustica united with Acustica, 1997.

[45] H. Riedel, M. Granzow and B. Kollmeier, Single-sweep-based methods to improve the quality of

auditory brain stem responses Part II: Averaging methods, Zeitschrift für Audiologie / Audiological Acoustics, vol. 40, no. 2, pp. 62-85, 2001.

[46] M. Hoke, B. Ross, R. Wickersberg and B. Lütkenhöner, Weighted averaging - theory and application to electric response audiometry, Electroen Clin Neuro, vol. 57, pp. 484-9, 1984.

[47] A. Yonovitz, C. L. Thompson and J. Lozar, Masking level differences: auditory evoked responses with homophasic and antiphasic signal and noise., Journal of speech and hearing research, vol. 22, no. 2, pp. 403-11, 6 1979.

[48] H. Hansen, J. L. Verhey and R. Weber, The Magnitude of Tonal Content. A Review, Acta Acustica united with Acustica, vol. 97, no. 3, pp. 355-363, 1 5 2011.

[49] A. Rupp, N. Sieroka, A. Gutschalk and T. Dau, Representation of Auditory-Filter Phase Characteristics in the Cortex of Human Listeners, Journal of Neurophysiology, vol. 99, no. 3, pp. 1152-1162, 3 2008.

[50] H. David, The Method of Paired Comparisons, New York: Oxford University Press, 1988.

[51] W. Ellermeier, M. Mader and P. Daniel, Scaling the Unpleasantness of Sounds According to the BTL Model: Ratio-Scale Representation and Psychoacoustical Analysis, Acta Acustica (Stuttgart), 2004.

[52] D. A. H. Thomas and H. A. David, The Method of Paired Comparisons., The Statistician, 2006.

[53] C. J. Billings, K. L. Tremblay, G. C. Stecker and W. M. Tolin, Human evoked cortical activity to signal-to-noise ratio and absolute signal level., Hearing research, vol. 254, no. 1-2, pp. 15-24, 8 2009.

Signal level (dB re masked threshold)

Salience (y-axis)

Signal level (dB re masked threshold) (x-axis)

Panel labels: A, B, C, D, E, F, G, H

A: PCr = 1.0, PCt = 2.4
B: PCr = 2.6, PCt = 4.3
C: PCr = 1.0, PCt = 3.8
D: PCr = 1.0, PCt = 2.4
E: PCr = 1.6, PCt = 3.1
F: PCr = 1.8, PCt = 4.7
G: PCr = 1.3, PCt = 2.4
H: PCr = 1.7, PCt = 4.3

Legend:
UN_0
UN_{150}
CM_0
CM_{150}
BB_0
BB_{150}