MicroRNA-transcriptome networks in whole blood and monocytes of women undergoing preterm labor

Alison G. Paquette, PhD[1] ; Oksana Shynlova, PhD[2] ; Xiaogang Wu, PhD[1] ; Mark Kibschull, PhD[2] ; Kai Wang, PhD[1] ; Nathan D. Price, PhD[1] ; Stephen J Lye, PhD[2*]

1. Institute for Systems Biology, Seattle WA 98119 2. Program in Development and Fetal Health, Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, Ontario, Canada

[*]Corresponding Author: Dr Stephen J Lye
Address: Lunenfeld-Tanenbaum Research Institute at Sinai Health System
6-1004 - 25 Orde Street, Toronto, Ontario, M5T 3H7.
Tel: 416-558-8640
Email: lye@lunenfeld.ca

ABSTRACT

Preterm birth is attributed to neonatal morbidity as well as cognitive and physiological challenges. We have previously identified significant differences in mRNA expression in whole blood and monocytes, as well as differences in miRNA concentration in blood plasma, extracellular vesicles (EV) and EV-depleted plasma in women undergoing spontaneous preterm labor (sPTL). The goal of this analysis was to identify differences in miRNA expression within whole blood (WB) and peripheral monocytes (PM) from the same population of women undergoing sPTL compared to nonlaboring controls matched by gestational age. We performed single end small RNA sequencing in whole blood and peripheral monocytes from women undergoing sPTL with active contractions(24-34 weeks of gestation, N=15) matched for gestational age to healthy pregnant non laboring controls (>37 weeks gestation, N=30) who later delivered at term as a part of the Ontario Birth Study (Toronto, Ontario CA). We identified significant differences in expression of 16 miRNAs in PMs and 9 miRNAs in WB in women undergoing sPTL. In PMs, these miRNAs were predicted targets of 541 genes, including 28 previously associated with sPTL. In WB, miRNAs were predicted to target 303 genes, including 9 previously associated with sPTL. These genes were involved in a variety of immune pathways, including interleukin 2 signaling. This study is the first to identify changes in miRNA expression in WB and PMs of women undergoing sPTL. Our results shed light on potential mechanisms by which miRNAs may play a role in mediating systemic inflammatory response in pregnant women that deliver prematurely.

INTRODUCTION

Premature birth, defined as delivery before 37 weeks of gestation, occurs in

11.1% of pregnancies worldwide and is associated with neonatal morbidity and

mortality[1] . Pregnancies characterized by pathological changes including placental

insufficiency, subclinical infections, disruptions in maternal immune-tolerance to

pregnancy, and decidual senescence often result in spontaneous preterm labor (sPTL).

There is a paucity of research examining transcriptomic changes that occur during

sPTL[2] , which have the potential to transform our understanding of the molecular

mechanisms underlying this heterogeneous syndrome [3] . Therefore, there is a crucial

research need for robust and multidimensional characterization of transcriptomic changes

relating to sPTL.

Throughout pregnancy, maternal blood circulates through the in-utero

environment and responds to fetal cues. At the beginning of gestation, the number of

monocytes in maternal blood drastically increase[4] , which regulate placental invasion,

angiogenesis, and tissue remodeling [5] . After infiltration into uterine tissues, monocytes

differentiate into macrophages characterized by an immunosuppressive (M2) phenotype

in normal pregnancies and an inflammatory (M1) state in complicated pregnancies[5] .

Maternal immune cells play a crucial role in normal pregnancy maintenance and provide

insight into changes that occur in pregnancies complicated by preterm birth.

Micro(mi)RNAs are a subtype of small, non-coding RNAs which are

transcriptional regulators of gene expression in all human organs. MiRNAs are expressed

within the placenta throughout pregnancy and are involved in fetal and maternal

signaling. Placental miRNA expression profiles have been associated with preeclampsia

[6] [7] and sPTL[8]. Like other organs, the placenta releases miRNAs into the circulation,

and unique placenta derived miRNAs (from the C19MC and C14MC miRAN cluster) are

detectable within maternal plasma[9–11]. Differential miRNA expression in plasma has

been detected in pregnancies complicated by fetal growth restriction preeclampsia[13],

and preterm birth[14, 15]. Circulating placenta enriched miRNAs are detectable in whole

blood, and have been associated with fetal hypoxia[16]. MiRNAs may also play a role in

monocyte differentiation throughout pregnancy, as there are differences in miRNA

concentration in different monocyte subpopulations (CD16+ vs. CD16-)[17]. miRNAs

have emerged as important transcriptional regulators and signaling molecules during

pregnancy with the potential to play a role in the underlying molecular perturbations that

occur during pregnancy-related complications such as sPTL.

Genome scale transcriptomic analysis of maternal blood provides a window into

the changes that occur throughout pregnancy and during parturition. We have previously

identified differences in mRNA expression between whole blood and peripheral

monocytes[18], as well as differences in miRNA concentration in whole plasma,

extracellular vesicles (EVs), and EV depleted plasma in women undergoing sPTL. The

goal for this analysis was to identify changes in miRNA expression in maternal whole

blood and monocytes in the same population (15 women undergoing sPTL and 30

pregnant women matched on gestational age not undergoing labor) and to integrate these

results with our previous findings. We hypothesize that there are changes in miRNAs in

whole blood and monocytes of women undergoing sPTL compared to controls, which are

related to the role of these miRNAs in transcriptional regulation of labor-associated

genes. Through this miRNA analysis and integration with next generation sequencing

mRNA data from the same individuals, we obtain a better understanding of the

transcriptional regulation that occurs in the context of PTL.

MATERIALS AND METHODS

Study Design

Participants were recruited within the Ontario Birth Study (OBS); a continuously

enrolling prospective cohort at Mount Sinai Hospital (MSH) in Toronto Canada.

Inclusion criteria for the OBS are: pregnancy diagnosed <17 weeks, maternal age >18,

English speaking, signed informed consent, and intent to deliver and receive antenatal

care at MSH. Exclusion criteria included non-viable neonate and/or inability to provide

consent. Peripheral blood was collected from 15 patients who were in preterm labor,

defined as cervical dilatation >4 cm and active uterine contractions, with no other

accompanying pathology; who delivered prematurely between 24-34 weeks of gestation

(Preterm labor or PTL). They were matched with 30 healthy asymptomatic pregnant

women whose blood was taken at the same time point during routine clinical

visits(gestational age 24-34 weeks), who later delivered at full term (TL). A full

description of this subpopulation is described in other manuscripts [18] [19]. This study

was approved by the Research Ethics Board of Mount Sinai Hospital, Toronto, Canada

(#04-0024-E). All patients provided written consent as part of the OBS at Sinai Health

System, Toronto, Canada.

Specimen Collection

In PTL patients, peripheral blood samples were collected prospectively at the point of

hospital admission, and in TL controls blood was collected during the regular antenatal

visit. Blood was collected into both PAXgene tubes (Qiagen; Hilden, Germany) for

whole blood isolation and EDTA blood collection tubes for monocyte separation.

Monocytes were separated through the Monocyte RosetteSep cocktail (Stemcell

Technologies, Vancouver CA), followed by high density gradient centrifugation to

generate a highly purified monocyte fraction for subsequent mRNA isolation.

Small RNA Sequencing and Quantification

Whole blood RNA was isolated using a PAXgene blood miRNA kit (Qiagen Hilden,

Germany), and from peripheral monocytes using TRIzol LS reagent (Thermo Fisher,

Waltham MA) following manufacturer's instructions. RNA quality was determined using

Experion analyzers (BioRad, Hercules CA) and sequenced at The Center for Applied

Genomics at the SickKids Hospital, Toronto, Canada. Library preparation was performed

following the New England Biolabs NEB Next multiplex small library preparation

protocol, with 400 ng of total RNA as the input. The 3' adapter was ligated to the small

RNA, followed by reverse transcriptase (RT) primer hybridization to the 3' adapter and

then the 5' adapter was ligated to the opposite end. Libraries were generated using first

stranded synthesis then enriched by PCR. Quality and size was determined using the

Bioanalyzer 2100 DNA High Sensitivity chip (Agilent Technologies, Palo Alto CA), and

libraries were quantified by qPCR using the Kapa Library Quantification Illumina/ABI

Prism Kit protocol (KAPA Biosystems, Wilmington MA) and sequenced on Illumina

HiSeq 2500 (Illumina, San Diego CA).

Sequencing data was preprocessed using the RNA analysis pipeline sRNAnalyzer

([20], http://srnanalyzer.systemsbiology.net/). In the data preprocessing, adaptor

sequences were trimmed and low-quality sequences were removed. Processed sequences

were aligned to all human miRNAs (miRBase Release 21,[21]) with no mismatches. We

removed miRNAs with 0 read counts in >50% of samples or a mean read count <20

producing a final dataset of 417 miRNAs in whole blood and 274 miRNAs in peripheral

monocytes, CPM (Count Per Million) values were calculated using RNA sequencing

analysis software "edgeR"[22], and log2 transformed. Data from this analysis is publicly

available within the Gene Expression Omnibus (GEO) as GSE108876 and GSE108877.

Statistical Analyses

Differentially expressed miRNAs were identified using edgeR [22]. miRNAs were

considering significantly associated with PTL if they exhibited a Benjamini Hochberg

(BH) adjusted q value of <0.05, and a log2 fold change of >1. In order to mitigate batch

effects, we eliminated differentially expressed miRNAs which were correlated with

order they were loaded onto the array (N=3 miRNAs in whole blood and monocytes),

identified using spearman correlation tests with a statistical cutoff of FDR adjusted q

value<0.05. We compared expression of miRNA in whole blood and peripheral

monocytes to previously published data from the same patient population, including

miRNA data from matched plasma and EVs(GSE106224)[19], as well as mRNA

expression from whole blood and peripheral monocytes (GSE96097) [18].

Putative miRNA targets were detected using the quantitative model TargetScan (V. 7.0,

targetscan.org), which characterizes canonical targeting of miRNAs based on 14 features,

and has the best predictive performance compared to comparable tools[23]. MiRNA-

target gene relationships were identified in whole blood and peripheral monocytes by

examining correlations between each miRNA and its proposed target genes. For each

miRNA, we used only mRNA targets with an absolute value of "context++" score (a

metric of miRNA target prediction accuracy used by TargetScan) which was higher than

the median, ensuring only highest quality relationships were validated. miRNA and target

relationships were considered statistically significant using a Spearman correlation

coefficient <-0.3 and a $p < 0.05$. These mRNAs were then matched to the list of

Differentially Expressed Genes (DEGs) identified in whole blood and monocytes from

the same individuals [18].

Gene set enrichment analysis was performed for mRNA targets using two-sided

hypergeometric tests conducted on Gene Ontology (GO) Biological Process Gene sets

using GO gene set visualization application "ClueGO" within the cytoscape environment

[24]. GO gene sets with more than 5 genes were included in the analysis, and were

considered significant with a Benjamini Hochberg adjusted $q < 0.05$. These GO gene sets

were clustered into ClueGO groups based on the similarity in the number of genes in the

gene sets calculated by a Kappa score within ClueGO. Data was analyzed and visualized

in R (Version 3.3.1) and Cytoscape (Version 3.6.0) [24].

RESULTS

Population Characteristics

Demographic characteristics of women undergoing sPTL (defined as active

uterine contractions with cervical dilation followed by delivery between 24-34 weeks)

were compared to healthy women not in labor that went on to deliver at full term (**Table**

1). There were no significant differences in delivery method, ethnicity, or fetal sex

between sPTL cases and controls ($p>0.05$, Fisher's exact test). Women who delivered

prematurely were more likely to be between 18 and 30 years old vs. older than 30,

although this relationship was borderline statistically significant (p=0.06).

Differences in miRNA expression in Whole Blood and Monocytes

We identified miRNAs with significantly different concentrations in peripheral

monocytes and whole blood of women undergoing PTL using generalized linear models

within edgeR [22]. In monocytes, 11 miRNAs were higher and 5 miRNAs were lower in

women undergoing PTL compared to women who delivered at term ***(Figure 1A-B,***

supplemental table 1). In whole blood, 6 miRNAs were higher and 3 miRNAs were

lower in women undergoing PTL compared to controls ***(Figure 1 C-D, supplemental***

table 1). Two of these miRNAs (mIR-495-3p and miR-381-3p) were from the C14MC

MC miRNA cluster; a region of chromosome 14 which encodes miRNAs which are

substantially enriched in placental tissue. Overall, there was a higher number of

significant miRNAs and larger concentration changes in relation to sPTL in monocytes

compared to whole blood.

MiR-1291-5p expression was significantly increased in whole blood of women

undergoing sPTL(Log fold change 1.12, FDR adjusted q=2.22x10^{-4}), but decreased in

monocytes of women undergoing sPTL (Log fold change -1.32, FDR adjusted

q=2.45x10^{4}). We observed a significant negative correlation between expression of this

miRNA in whole blood and in monocytes (p=0.02, ρ =-0.02, Spearman correlations,

Supplemental Figure 1). This higher level in whole blood suggests that other sub-

populations of blood cells (such as erythrocytes, platelets, lymphocytes or granulocytes)

may have the opposite expression profile compared to monocytes and also contribute to

the miRNA pool in women undergoing sPTL.

Comparison of expression differences across blood compartments

In our prior analysis of plasma from same patients from the OBS cohort, we

identified significant differences in concentration of 132 miRNAs in whole plasma, EVs

and EV depleted plasma [19]. Five of these miRNAs were also significantly different in

monocytes, and three miRNAs were significantly different in whole blood

(**supplemental table 2**). We examined correlations of these 8 miRNAs that were

significantly upregulated in plasma samples as well as in whole blood and monocytes

from the same patients (**supplemental figure 2,** N=44 samples with plasma and whole

blood RNAseq data, N=42 samples with monocyte and whole blood sRNA sequencing

data, and N=14 samples with RNAseq data in EVs and monocytes/whole blood).

Two miRNAs (miR-183-5p, and miR-331-3p) were increased and one miRNA

(miR-328-3p) was decreased in both plasma and monocytes of women undergoing sPTL

compared to TL controls. The concentration/expression of these miRNAs was not

significantly correlated between whole plasma and monocytes (p>0.05, Spearman

correlations). MiR-181b-5p was increased in monocytes and decreased in the plasma of

women undergoing sPTL, and the expression/concentration of these miRNAs was

inversely correlated in plasma and monocytes (ρ= -0.28, p=0.07). MiR-378c-5p was

increased in both monocytes and in extracellular vesicles of women undergoing sPTL.

We observed significant positive correlations (p<0.05) between whole blood miRNA

levels and the levels in plasma and EVs in the 3 miRNAs significantly associated with

sPTL in the previous studies: (1) MiR-374a-5p was decreased in both plasma and whole

blood of women undergoing sPTL, (2) miR-381-3p was decreased in plasma but

increased in whole blood of women undergoing sPTL, and (3) miR495-5p was decreased

in the plasma and extracellular vesicles but increased in the whole blood of women

undergoing sPTL compared to controls. This indicates that there is some overlap in signal

related to sPTL that is detectable across different components of the blood within the

same individuals.

Confirmation of target mRNAs of miRNAs associated with sPTL

We identified the putative mRNA targets of the differentially expressed miRNAs

using the TargetScan database. TargetScan incorporates 14 features to predict miRNA-

mRNA interactions and has shown to have the highest predictive value compared to

similar tools [23] . From the TargetScan database, we included only the top 50% of

predicted mRNA based on the "context ++" score (a metric of accuracy used by the

database). We then compared mRNA expression of these predicted targets to the miRNA

concentration within matched samples (GSE96097,[18]). We constrained miRNA-

mRNA predicted interactions to only those which had a Spearman correlation coefficient

$\rho < -0.3$ and a $p < 0.05$, which represented 7.29% of the predicted genes in monocytes and

4.26% in whole blood (**Supplemental table 3, Supplemental Figure 4**). In monocytes,

541 genes were associated with 16 miRNAs, and in whole blood 303 genes were

associated with 7 miRNAs. Based on this criteria, there were 2 miRNAs in whole blood

with no confirmed mRNA targets.

In previous transcriptomic analyses from the same patients, we identified 262

genes in monocytes and 181 genes in whole blood which were associated with sPTL[18].

Here, we compared the changes in mRNA expression related to sPTL and the confirmed

miRNA target genes we have identified in this current study (**Figure 2**). In whole blood,

9 of the 181 DEGs were confirmed miRNA targets of 3 differentially expressed miRNAs

in whole blood (Shown in **Figure 2A**). Among these 3 miRNAs in whole blood, miR-

4742-3p emerged as the strongest regulator, since it was negatively associated with 5

genes which were positively associated with sPTL, including *SPH*, *CD177*, *CYP1B1*,

ELOVL7 and *GRB10.* In monocytes, 28 of these 262 genes were confirmed mRNA

targets of 7 differentially expressed miRNAs. In monocytes, MiR-1291-5p emerged as

the strongest regulator, and was negatively associated with 11 of the 28 genes which are

positively associated with sPTL, including *IL1B*, *IL1R1* and *CD177* (**Figure 2B**). In both

whole blood and monocytes, the direction of the associations with PTL is congruent with

the directionality in mRNAs (i.e., if a miRNA is positively associated with PTL, the

miRNA is negatively correlated with mRNAs that are negatively associated with PTL).

We also examined mRNAs which were in the same genomic region as the miRNAs in

our study, but found that none of the associated mRNAs were statistically significantly

associated with sPTL in monocytes and whole blood leukocytes. This integrated analysis

suggests that the miRNAs identified here may play a role in the transcriptional regulation

of genes associated with sPTL in monocytes and whole blood leukocytes.

miRNA target mRNA networks in sPTL

We performed enrichment analysis to identify gene sets which were

overrepresented by the confirmed mRNA targets of the differentially expressed miRNAs

identified in whole blood and monocytes. Using CytoScape application "ClueGO", we

identified significantly enriched GO gene sets which were grouped together based on

intersecting common genes, which are shown in **Figure 3, and in supplemental table 4**.

In whole blood, 59 different GO gene sets were significantly enriched for the 303 genes

which were mRNA targets for the 9 miRNAs associated with sPTL in whole blood.

These gene sets were grouped into 19 ClueGO groups, including 8 distinct groups (i.e.

only had 1 GO gene set), and one ClueGO group which contained 22 GO gene sets

including the 5 most significant gene sets: "positive regulation of cytokine production" ,

"positive regulation of adaptive immune response" , "interleukin-2 production" ,

"positive regulation of adaptive immune response based on somatic recombination of

immune receptors built from immunoglobulin superfamily domains" , and" T cell

receptor signaling pathway". This diversity of GO gene sets and strong signal indicates

that these miRNAs associated with sPTL are involved in a wide variety of biological

pathways in whole blood; particularly immune pathways.

In monocytes, 30 different gene sets were significantly enriched for the 541 genes

which were targets for the 16 miRNAs associated with sPTL. These gene sets were

clustered into 11 groups, including 8 gene sets which were distinct (i.e only had 1 GO

term), and 1 gene set (Group 10), which had 13 different terms. The top 5 GO terms

based on q value were: "small GTPase mediated signal transduction"," negative

regulation of protein modification process", "positive regulation of interleukin 2

production", "positive regulation of mononuclear cell proliferation", and "negative

regulation of protein kinase activity". These terms are reflective of changes associated

with monocyte proliferation and immune function, indicating that miRNAs may play a

role in regulation of these pathways in monocytes.

The Gene Ontology gene set "positive regulation of interlukin-2 production" was

significantly enriched in both whole blood and monocytes. This gene set contains 31

genes, and the differentially expressed miRNAs in both whole blood and monocytes were

predicted negative regulators of 5 unique genes within this gene set, each representing

17.86% of the total genes in this pathway. In whole blood, 4 miRNAs were identified as

regulators of the 5 genes (*CCR2*, *CD4*, *MALT1*, *PDE4B*, and *CD28*). In monocytes, 3

miRNAs were identified as negative regulators of the 5 genes (*SASH3, CD83, IL1B,*

TNFSF4, and CD28) involved in interleukin signaling (**Table 2**). In this gene set, *IL1B*

was a target of miR-1291-5p, and *IL1B* expression was significantly increased in women

undergoing sPTL (FDR adjusted q=4.02x10-3, logFC=2.01). Through this analysis, we

have identified that miRNA expression within whole blood and peripheral monocytes

significantly alters genes involved in production of interleukin 2.

DISCUSSION

This study is the first comprehensive miRNA profiling of women undergoing

sPTL using small RNA sequencing within both whole blood and peripheral monocytes.

Our key findings included (A) identification of differentially expressed miRNAs in whole

blood and peripheral monocytes, which (B) were negative regulators of genes associated

with sPTL, which we identified using previously generated RNA sequencing data on

whole blood and monocytes from the same individuals and (C) identification of unique

and shared gene sets enriched for mRNA targets of these differential miRNAs in

monocytes and whole blood of women undergoing sPTL.

Overall, we identified a greater number of miRNAs with concentration changes,

and more genes related to miRNAs associated with sPTL. Additionally, we observed no

congruence in the differential miRNAs identified in monocytes and whole blood.

Maternal blood includes three major subsets of immune cells; monocytes, lymphocytes,

and granulocytes, which circulate through gestational tissues including the myometrium,

decidua and placenta, and are exposed and respond to signals from these tissues. RNA

sequencing data derived from monocytes exhibits less cellular heterogeneity and thus

likely has a cleaner signal. We suggest that the signaling we observe in monocytes may

be related to their functional role throughout pregnancy and parturition [4] . Altogether,

our work suggests that a stronger signal related to sPTL can be obtained from monocytes

compared to whole blood.

We observed a number of miRNAs which were differentially expressed in in

whole blood and monocytes of women undergoing sPTL which also exhibited significant

concentration differences previously identified in whole plasma, EV-depleted plasma and

EVs in the same population of individuals [19] . We identified fewer differential miRNAs

in immune cells compared to the plasma component, but overlap in the differentially

expressed miRNAs, as well as concurrence in the directionality of associations with

sPTL. This suggests functional signaling that occurs within blood in context of sPTL. In

independent populations, increased expression of miR-223 has been identified in plasma

of women at 20 weeks who went on to deliver prematurely[14]. Similarly, we observed

increased levels of this miRNA in whole blood of women who delivered prematurely

compared to TL controls, which supports the potential of this miRNA as having a

functional role in PTL.

In whole blood, 2 of the miRNAs (miR-495-3p and miR-381-3p) with

concentration differences in women undergoing sPTL were part of the C14MC region; a

group of imprinted miRNAs located on chromosome 14q32 which is highly produced by

the placenta and enters maternal circulation [25]. In plasma samples from the same

patients, these miRNA concentrations were negatively associated with sPTL[19], and

changes in average expression of C14MC miRNAs in plasma have also been associated

with preterm labor[15]. On average, C14MC miRNAs have higher expression and lower

variance in the fetal compartment and the placenta compared to the maternal plasma, thus

this signal may be reflecting changes in this compartment[26]. We hypothesize that the

changes observed within the whole blood and monocyte fraction of leukocytes are

reflective of functional transcriptional changes that occur within the blood or they may be

reflective of concentration changes related to miRNA signals from the developing fetus.

We identified mRNAs which were transcriptionally regulated by differentially

expressed miRNAs using predicted targets from TargetScan, which were filtered using

matched mRNA data derived through RNA sequencing based on negative correlations

with miRNA[18]. Of the targets identified by TargetScan, on average only 4.26% of

these predicted targets passed our filter whole blood and 7.29% in monocytes. These

target prediction algorithms are only designed to reveal gene expression regulation

potential, and expression can also be altered by post transcriptional modifications

including transcriptional regulation, histone acetylation and DNA methylation [27], and

are not cell type specific [23]. This analysis was designed to collect miRNA and mRNA

from the same patient, which allowed us to perform correlations and obtain a far more

accurate understanding of miRNA-mediated regulation of gene expression than by

relying solely on miRNA prediction algorithms alone.

We used our curated, confirmed miRNA target lists generated to perform an

enrichment analysis of gene sets related to our differentially expressed miRNAs. In

whole blood, we identified immune related gene sets. This may be reflective of the

inflammatory and immune response that occurs as a breakdown in maternal fetal

tolerance or inflammation related to preterm labor[28]. The target genes of the miRNAs

differentially expressed within monocytes were enriched for a total of 19 GO gene sets

including those related to interleukin production and activation of the immune system,

such as "regulation of interleukin production (Interleukins 2, 4 and 6), as well as T Cell,

lymphocyte, and monocyte proliferation. The monocyte population expands during

pregnancy and differentiates into a pro-inflammatory phenotype during labor [30]. Our

work indicates that part of this expansion in monocytes may be transcriptionally

regulated by a subset of miRNAs in monocytes. More work is required to understand if

these miRNAs induce monocyte expansion and proliferation *in vitro*, which is beyond the

scope of this analysis.

In monocytes, miR-1291-5p emerged as a master miRNA regulator of the highest

number of genes associated with sPTL, including *IL1B* and its receptor *IL1R1*, which

have previously been associated with sPTL. The positive regulation of interleukin 2 gene

set was significantly enriched for miRNA target genes in both whole blood and

monocytes, which contains *IL1B*. This gene set is involved in immune response and

activation of inflammatory cytokines [31]. Inflammatory cytokines increase at the end of

pregnancy as monocytes shift into an inflammatory M1 phenotype [30], and are

implicated in sPTL[32]. Our work suggests that this may be in part modulated through

changes in miRNAs expression in whole blood and monocytes.

We are limited in our ability to perform extensive characterization,

stratification/clustering or any predictive analyses of these miRNAs due to inadequate

statistical power resulting from a small sample size. Additionally, we cannot separate

miRNA signals related to PTL from those related to the labor process (uterine

contractions) or to the betamethasone treatment that women delivering prematurely

received prior to labor, as previously discussed [18] . Due this these limitations, this work

was not designed to interrogate miRNAs as a potential biomarker for sPTL. Further

functional analyses are required, including functional validation of miRNA/mRNA

targets, as well as the functionality of these miRNAs on key tissues involved in

pregnancy (i.e. placenta and myometrium). Additionally, our findings will need to be

validated in independent cohort of pregnant women. This work instead focuses on

potential mechanisms by which miRNAs may play a role in mediating systemic

inflammatory response in pregnant women that deliver prematurely, serves as an

important foundation to subsequent analyses, and highlights the importance of sample

source when performing blood related assessments.

Our analysis stands out among related studies because it is the first to generate

high throughput RNA sequencing data of both miRNA and mRNA from the same patient

source. We have carefully selected a reliable methodology to identify potential miRNA

targets based on experimental data [23], and have used novel alignment tools to obtain

comprehensive coverage of miRNA expression [20]. This robust, multi-dimensional data

has allowed us to confidently build mRNA-miRNA regulation networks related to sPTL

in a way that has not previously been done, which provides additional insight into

mechanisms of transcriptional regulation of human labor. This type of integrative

analysis has been suggested to overcome many challenges in performing pregnancy

research [3]. Our analysis has highlighted miRNA mediated transcriptional regulatory

networks of sPTL-associated genes in monocytes and whole blood, which are involved in

important biological pathways, including interleukin signaling. Further work is needed to

validate these data in independent populations and elucidate the miRNA mechanisms of

signaling in whole blood and in a subpopulation of peripheral monocytes.

ACKNOWLEDGEMENTS

The authors would like to thank the women who participated in this study as well as the

staff at Mount Sinai Hospital, in particular Drs Robin Thurman and Kellie Murphy. We

would like to thank members of the Price lab, particularly Ben Heavner and Rory

Donovan for their help with the data management and members of Lye lab, particularly

Mrs Elzbieta Matysiak-Zablocki and Mrs Anna Dorogin for their help with blood

processing. We would like to acknowledge Drs Lee Hood, Stephen Matthews, Sam

Mesiano, Leslie Myatt, Craig Pennell and Yoel Sadovsky on behalf of the GAPPS

"Systems Biology of Preterm Birth" pilot study, as well as Mary Brunkow for her

coordination of this study. We would also like to thank Beverly Anne Bessie and Theresa

Fitzgerald for their help with the coordination of this manuscript. This work is supported

by the Global Alliance to Prevent Prematurity and Stillbirth (GAPPS), Grant Title

"Systems Biology of Preterm Birth: A Pilot Study", as well as the Eunice Kennedy

418   Shriver National Institute of Child Health & Human Development of the National

419   Institutes of Health: Grant R01HD091527 (to N.P), and K99HD096112 (to A.P).

CONFLICTS OF INTEREST

421   The authors confirm that there are no conflicts of interest.

WORKS CITED

[1]   **Blencowe H, Cousens S, Oestergaard MZ, et al.** National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *The Lancet* n.d.; 379; 2162–72.

[2]   **Eidem HR, Ackerman WE, McGary KL, et al.** Gestational tissue transcriptomics in term and preterm human pregnancies: a systematic review and meta-analysis. *BMC Med. Genomics* 2015; 8; 27.

[3]   **Eidem HR, McGary KL, Capra JA, et al.** The transformative potential of an integrative approach to pregnancy. *Placenta* 2017; 57; 204–15.

[4]   **Tang M-X, Hu X-H, Liu Z-Z, et al.** What are the roles of macrophages and monocytes in human pregnancy? *J. Reprod. Immunol.* 2015; 112; 73–80.

[5]   **Brown MB, von Chamier M, Allam AB, et al.** M1/M2 Macrophage Polarity in Normal and Complicated Pregnancy. *Front. Immunol.* 2014; 5; 606.

[6]   **Hu Y, Li P, Hao S, et al.** Differential expression of microRNAs in the placentae of Chinese patients with severe pre-eclampsia. *Clin. Chem. Lab. Med.* 2009; 47; 923–9.

[7]   **Enquobahrie DA, Abetew DF, Sorensen TK, et al.** Placental microRNA expression in pregnancies complicated by preeclampsia. *Am. J. Obstet. Gynecol.* 2011; 204; 178.e12-178.e21.

[8]   **Mayor-Lynn K, Toloubeydokhti T, Cruz AC, et al.** Expression Profile of MicroRNAs and mRNAs in Human Placentas From Pregnancies Complicated by Preeclampsia and Preterm Labor. *Reprod. Sci.* 2011; 18; 46–56.

[9]   **Williams Z, Ben-Dov IZ, Elias R, et al.** Comprehensive profiling of circulating microRNA via small RNA sequencing of cDNA libraries reveals biomarker potential and limitations. *Proc. Natl. Acad. Sci.* 2013; 110; 4255–60.

[10]   **Chim SSC, Lee WS, Ting YH, et al.** Systematic identification of spontaneous preterm birth-associated RNA transcripts in maternal plasma. *PloS One* 2012; 7; e34328.

[11]   **Weber JA, Baxter DH, Zhang S, et al.** The microRNA spectrum in 12 body fluids. *Clin. Chem.* 2010; 56; 1733–41.

[12]   **Chim SSC, Shing TKF, Hung ECW, et al.** Detection and Characterization of Placental MicroRNAs in Maternal Plasma. *Clin. Chem.* 2008; 54; 482–90.

[13]   **Gunel T, Zeybek YG, Akcakaya P, et al.** Serum microRNA expression in pregnancies with preeclampsia. *Genet. Mol. Res. GMR* 2011; 10; 4034–40.

[14] **Gray C, McCowan LM, Patel R, et al.** Maternal plasma miRNAs as biomarkers during mid-pregnancy to predict later spontaneous preterm birth: a pilot study. *Sci. Rep.* 2017; 7; 815.

[15] **Wommack JC, Trzeciakowski JP, Miranda RC, et al.** Micro RNA clusters in maternal plasma are associated with preterm birth and infant outcomes. *PLOS ONE* 2018; 13; e0199029.

[16] **Whitehead CL, Teh WT, Walker SP, et al.** Circulating MicroRNAs in Maternal Blood as Potential Biomarkers for Fetal Hypoxia In-Utero. *PLoS ONE* 2013; 8; e78487.

[17] **Dang T-M, Wong W-C, Ong S-M, et al.** MicroRNA expression profiling of human blood monocyte subsets highlights functional differences. *Immunology* 2015; 145; 404–16.

[18] **Paquette AG, Shynlova O, Kibschull M, et al.** Comparative analysis of gene expression in maternal peripheral blood and monocytes during spontaneous preterm labor. *Am. J. Obstet. Gynecol.* 2018; 218; 345.e1-345.e30.

[19] **Fallen S, Baxter D, Wu X, et al.** Extracellular vesicle RNAs reflect placenta dysfunction and are a biomarker source for preterm labour. *J. Cell. Mol. Med.* 2018; DOI: 10.1111/jcmm.13570.

[20] **Wu X, Kim T-K, Baxter D, et al.** sRNAnalyzer-a flexible and customizable small RNA sequencing data analysis pipeline. *Nucleic Acids Res.* 2017; DOI: 10.1093/nar/gkx999.

[21] **Griffiths-Jones S, Saini HK, van Dongen S, et al.** miRBase: tools for microRNA genomics. *Nucleic Acids Res.* 2008; 36; D154–8.

[22] **Robinson MD, McCarthy DJ, Smyth GK**. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2010; 26; 139–40.

[23] **Agarwal V, Bell GW, Nam J-W, et al.** Predicting effective microRNA target sites in mammalian mRNAs. *ELife* 2015; 4; e05005.

[24] **Shannon P, Markiel A, Ozier O, et al.** Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 2003; 13; 2498–504.

[25] **Seitz H, Royo H, Bortolin M-L, et al.** A large imprinted microRNA gene cluster at the mouse Dlk1-Gtl2 domain. *Genome Res.* 2004; 14; 1741–8.

[26] **Paquette A, Chu, Tianjiao, Wu, Xiaogang, et al.** Distinct communication patterns of trophoblastic miRNA among the maternal-placental-fetal compartments. *Placenta* n.d.; 72–73; 28–35.

[27] **Allis CD, Jenuwein T**. The molecular hallmarks of epigenetic control. *Nat. Rev. Genet.* 2016; 17; 487.

[28] **Romero R, Dey SK, Fisher SJ**. Preterm labor: one syndrome, many causes. *Science* 2014; 345; 760–5.

[29] **Ball G, Srinivasan L, Aljabar P, et al.** Development of cortical microstructure in the preterm human brain. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110; 9541–6.

[30] **Zhang Y-H, He M, Wang Y, et al.** Modulators of the Balance between M1 and M2 Macrophages during Pregnancy. *Front. Immunol.* 2017; 8; 120.

[31] **Hoyer KK, Dooms H, Barron L, et al.** Interleukin-2 in the development and control of inflammatory disease. *Immunol. Rev.* 2008; 226; 19–28.

503 [32] **Nadeau-Vallee M, Obari D, Quiniou C, et al.** A critical role of interleukin-1 in
504   preterm labor. *Cytokine Growth Factor Rev.* 2016; 28; 37–51.
505
506 **TABLES**

507 **Table 1: Participant Characteristics**

A. Categorical Variables	N	%	N	%	P*
	Term (N=30)		Preterm (N=15)		
Delivery Method					0.09
C Section	12	27%	2	13%	
Vaginal	17	38%	13	87%	
Fetal Sex					1.00
Female	13	43%	6	40%	
Male	16	53%	9	60%	
Maternal Ethnicity					0.71
Nonwhite	8	27%	4	27%	
White	20	67%	7	47%	
Previous Preterm Delivery					1.00
No	26	87%	13	87%	
Yes	3	10%	2	13%	
Multiple Pregnancies					0.38
Yes	0	0%	10	67%	
No	29	97%	5	33%	
APH					0.59
No	28	93%	13	87%	
Yes	2	7%	2	13%	
Corticosteroids					0.04
No	30	100%	3	20%	
Yes	0	0%	12	80%	
Antibiotics					4.29E-04
No	23	77%	3	20%	
Yes	7	23%	12	80%	
MgSO4					0.04
No	30	100%	3	20%	
Yes	0	0%	12	80%	
Gestational Diabetes					1.00
No	26	87%	13	87%	
Yes	4	13%	2	13%	

	Min			Std	
Pregnancy Smoking Status					1.00
Yes	0	0%	0	0%	
No	30	100%	15	100%	

B. Continuous Variables	Min	Mean	Max	Std Deviation	P**
Maternal Age					0.06
Term	21	35.60	47	5.65	
Preterm	21	31.33	46	7.64	
Maternal Pre-Pregnancy BMI					0.53
Term	20	27.87	35	3.79	
Preterm	19	27.64	39	6.03	
Gestational Age at Blood Collection					0.19
Term	24.14	27.81	29.00	1.15	
Preterm	23.29	28.78	33.29	2.97	
Gestational Age at Delivery					6.39E-08
Term	36.29	39.33	41.71	1.32	
Preterm	23.29	28.78	33.29	2.97	
Gravidity					0.52
Term	1	2.13	5	1.07	
Preterm	1	2.07	6	1.44	
Parity					0.61
Term	0	0.63	2	0.72	
Preterm	0	0.53	2	0.74	

Table 2: mRNA targets of differentially expressed miRNAs involved in Positive regulation of Interleukin-2 signaling

miRNA	Gene targets in GO:0032743
Monocytes	

hsa-miR-1248-5p	*SASH3, CD83*
hsa-miR-1291-5p	*IL1B*
hsa-miR-378c-5p	*CD28,TNFSF4*
Whole Blood	
hsa-miR-4742-3p	*CCR2*
hsa-miR-1299-3p	*CD4,MALT1*
hsa-miR-582-3p	*PDE4B*
hsa-miR-1291-5p	*CD28*

FIGURE LEGENDS

Figure 1: MA plot of log2 fold changes vs. the mean of normalized counts in the regularized logarithmic distribution of RNA sequencing data in (**A**) monocytes, and (**C**) whole blood. mRNAs significantly different in linear models after adjustment for multiple comparisons are highlighted in red, and lines indicate log2 fold changes >1. Heatmap of the 16 differentially expressed genes identified in monocytes (**B**) or the 9 differentially expressed genes identified in whole blood (**D**), with blue indicating lower expression and red indicating higher expression. miRNA expression of women who underwent term labor is on the left and highlighted in grey, and miRNA expression of women in sPTL is highlighted in blue on the right.

Figure 2: A. Venn diagram in whole blood (**A**) and monocytes (**B**) of genes previously associated with sPTL and gene targets of the differentially expressed miRNAs identified in this study. In the overlapping genes, we show the log fold change and P values of the

miRNAs (from supplemental table 1), correlation with mRNA concentration, and log fold

change and correlation with sPTL in the previous study).

Figure 3: Biomolecular network of miRNA target genes in (**A**) monocytes and (**B**) whole

blood. Enrichment analysis was performed on gene ontology (GO) gene sets and

significant gene sets are shown (Benjamini Hoch adjusted q<0.05). Network node layout

is based on similarity between genes within GO gene sets, where nodes are colored based

on multiple occurrences within different go categories. GO terms which were significant

in both whole blood and monocytes have a red outer circle.

Supplemental Figure 1: miR-1291-5p expression in whole blood and monocytes.

miRNA expression values for women who delivered at term are highlighted in orange,

and values for women who delivered prematurely are in blue.

Supplemental Figure 2: Correlations in miRNA expression which were identified as

significant in (A) monocytes and plasma (B) monocytes and exosomes (C) whole blood

and plasma/exosomes. Blue dots indicate correlations involving monocytes, and red dots

indicate correlations involving whole blood. Orange dots reflect correlation between

whole blood and exosomes. Circled dots are expression values involving exosomes.

Correlations were evaluated using spearman correlations, with significance defined as

p<0.05. All correlations; including those which are insignificant are shown here.

Supplemental Figure 3: Number of miRNA targets identified in (A) whole blood and

(B) Monocytes. Blue lines indicates the number of targetscan candidates which had

higher than the median context ++ scores, which is a measure of prediction accuracy to

bind to target miRNA. Green bars indicate mRNAs with matched expression data in

GSE96097. Yellow bars indicate samples whose miRNA expression in whole blood

and/or monocytes was significantly correlated with the mRNA expression (defined as ρ

<-0.3, $p<0.05$, spearman correlation coefficients) in matched samples.

A. miRNA MA plot: Monocytes

B. Heatmap of significant miRNAs: Monocytes

C. miRNA MA plot: Whole Blood

D. Heatmap of significant miRNAs: Whole Blood

A. miRNA target genes associated with sPTL: Whole Blood

Genes associated with sPTL

Target genes of miRNAs associated with sPTL

172   9   294

	miRNA Association w/ sPTL		Correlation w/ mRNA			miRNA Association w/ sPTL	
	Log Fold Change	Q Value	Gene	P Value	Rho	Log Fold Change	Q Value
hsa-miR-1299-3p	2.06	1.38E-02	SMPD3	4.71E-02	-0.32	-1.12	3.59E-05
			ZNF749	3.86E-02	-0.34	-1.03	4.20E-04
hsa-miR-374a-5p	-1.02	2.46E-03	MMP14	1.27E-03	-0.50	1.13	5.35E-06
			SAP30	1.20E-02	-0.40	1.39	2.30E-07
hsa-miR-4742-3p	-1.07	5.85E-11	ASPH	1.55E-02	-0.39	1.04	1.12E-05
			CD177	9.44E-03	-0.42	2.37	5.92E-06
			CYP1B1	1.03E-02	-0.41	1.48	1.78E-08
			ELOVL7	3.51E-03	-0.46	1.08	9.72E-04
			GRB10	4.93E-02	-0.32	1.30	2.53E-05

B. miRNA target genes associated with sPTL: Monocytes

Genes associated with sPTL

Target genes of miRNAs associated with sPTL

233   28   513

	miRNA Association w/ sPTL		Correlation w/ mRNA			miRNA Association w/ sPTL	
	Log Fold Change	Q Value	Gene	P Value	Rho	Log Fold Change	Q Value
hsa-miR-1248-5p	-1.18	1.02E-02	CXCR1	9.08E-04	-0.52	3.47	4.47E-30
			DHCR24	1.13E-02	-0.41	1.08	1.95E-03
			GPR146	4.85E-02	-0.33	1.74	3.37E-02
			PROK2	4.51E-02	-0.33	2.23	5.93E-16
hsa-miR-1291-5p	-1.32	6.76E-05	ALPL	3.44E-04	-0.56	4.59	4.37E-16
			AMPH	7.59E-03	-0.43	2.91	7.46E-22
			CD177	4.79E-02	-0.33	2.44	1.12E-05
			DHRS13	5.47E-03	-0.45	1.17	4.92E-11
			IL1B	8.89E-03	-0.42	2.01	4.02E-03
			IL1R1	3.08E-02	-0.36	1.25	2.05E-07
			ITGA7	3.16E-02	-0.35	2.17	2.64E-02
			KREMEN1	5.96E-03	-0.44	1.81	1.16E-15
			LCK	2.98E-02	-0.36	2.03	9.19E-05
			OR2W3	1.91E-02	-0.38	1.02	4.14E-02
			UNC5A	1.62E-02	-0.39	1.01	4.68E-03
hsa-miR-21-3p	1.43	1.67E-09	AKAP2	7.72E-05	-0.60	-1.00	1.86E-02
			C15orf54	4.15E-04	-0.55	-1.39	7.17E-05
			GBP4	2.54E-02	-0.37	-1.06	3.13E-04
			SCUBE1	3.89E-02	-0.34	-1.32	3.93E-02
			SEZ6L	4.57E-02	-0.33	-1.43	7.21E-08
hsa-miR-3120-5p	-1.09	1.65E-02	GYPC	4.05E-03	-0.46	1.01	3.96E-02
			MAPK13	2.14E-02	-0.38	1.11	4.11E-08
			SLC1A3	1.47E-03	-0.50	1.71	1.42E-02
			SLC6A8	8.56E-03	-0.43	1.35	1.08E-02
hsa-miR-342-5p	-1.01	5.32E-04	MCTP2	2.37E-05	-0.64	1.57	5.56E-22
			PGLYRP1	2.04E-04	-0.57	1.56	7.27E-04
hsa-miR-4286-5p	1.03	9.28E-04	PPM1N	1.03E-02	-0.42	-1.58	1.33E-02
			SCAMP5	2.42E-02	-0.37	-1.36	4.14E-03

A. Pathway Enrichment: miRNA target Genes (Whole Blood)

positive regulation of NF-kappaB transcription factor activity

protein depolymerization

regulation of microtubule-based process

response to organophosphorus

B cell differentiation

water-soluble vitamin metabolic process

vitamin metabolic process

coenzyme biosynthetic process

cofactor biosynthetic process

thioester biosynthetic process

acyl-CoA biosynthetic process

thioester metabolic process

acyl-CoA metabolic process

antigen receptor-mediated signaling pathway

T cell receptor signaling pathway

regulation of proteasomal ubiquitin-dependent protein catabolic process

ovulation cycle

organelle transport along microtubule

positive regulation of leukocyte mediated immunity

positive regulation of monocyte interferon-gamma production

regulation of lymphocyte mediated immunity

regulation of production of molecular mediator of immune response

regulation of adaptive immune response

positive regulation of adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains

positive regulation of production of molecular mediator of immune response

regulation of B cell activation

positive regulation of adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains

regulation of interleukin-2 production

interferon-gamma production

interleukin-2 production

cytokine biosynthetic process

regulation of cytokine biosynthetic process

cytokine metabolic process

positive regulation of interleukin-2 production

organic hydroxy compound catabolic process

organophosphate catabolic process

negative regulation of viral process

response to hydrogen peroxide

cellular response to hydrogen peroxide

dendritic spine development

dendrite development

regulation of dendritic spine development

regulation of dendrite development

regulation of dendrite morphogenesis

positive regulation of cytokine production

regulation of myeloid leukocyte differentiation

osteoclast differentiation

ncRNA transcription

transcription elongation from RNA polymerase II promoter

DNA-templated transcription, elongation

type I interferon production

regulation of type I interferon production

mitochondrial membrane organization

apoptotic mitochondrial changes

positive regulation of hemopoiesis

B. Pathway Enrichment: miRNA target Genes (Monocytes)

regulation of interleukin-4 production

interleukin-6 production

regulation of interleukin-6 production

interleukin-4 production

lymphocyte activation

response to antibiotic

protein stabilization

protein ubiquitination

negative regulation of protein phosphorylation

positive regulation of alpha-beta T cell activation

regulation of leukocyte activation

positive regulation of interleukin-2 production

negative regulation of protein modification process

negative regulation of kinase activity

protein dephosphorylation

negative regulation of protein kinase activity

positive regulation of leukocyte proliferation

small GTPase mediated signal transduction

negative regulation of protein serine/threonine kinase activity

negative regulation of cyclin-dependent protein kinase activity

positive regulation of mononuclear cell proliferation

positive regulation of lymphocyte proliferation

oligosaccharide metabolic process

cellular response to mechanical stimulus

cell adhesion mediated by integrin

regulation of cell adhesion mediated by integrin