Publicly available transcriptomes provide the opportunity for dual RNA-Seq meta-analysis in *Plasmodium* infection

Parnika Mukherjee[1,2,3] , Gaétan Burgio[3] and Emanuel Heitlinger[1,2]

[1]Department of Molecular Parasitology, Humboldt University Berlin, Germany

[2]Research Group Ecology and Evolution of Molecular Parasite-Host Interactions, Leibniz-Institute for Zoo and Wildlife Research (IZW), Berlin, Germany

[3]Department of Immunology and Infectious Diseases, John Curtin School of Medical Research, Australian National University, Canberra, ACT, Australia

Abstract

Dual RNA-Seq is the simultaneous transcriptomic analysis of interacting symbionts, for example, in malaria. Potential cross-species interactions identified by correlated gene expression might highlight interlinked signaling, metabolic or gene regulatory pathways in addition to physically interacting proteins. Often, malaria studies address one of the interacting organisms – host or parasite – rendering the other "contamination". Here we perform a meta-analysis using such studies for cross-species expression analysis.

We screened experiments for gene expression from host and *Plasmodium.* Out of 171 studies in *Homo sapiens, Macaca mulatta* and *Mus musculus*, we identified 63 potential studies containing host and parasite data. While 16 studies (1950 samples) explicitly performed dual RNA-Seq, 47 (1398 samples) originally focused on one organism. We found 915 experimental replicates from 20 blood studies to be suitable for co-expression analysis and used orthologs for meta-analysis across different host-parasite systems. Centrality metrics from the derived gene expression networks correlated with gene essentiality in the parasites. We found indications of host immune response to elements of the *Plasmodium* protein degradation system, an antimalarial drug target. We identified well-studied immune responses in the host with our co-expression networks as our approach recovers known broad processes interlinked between hosts and parasites in addition to individual host and parasite protein associations.

The set of core interactions represents commonalities between human malaria and its model systems for prioritization in laboratory experiments. Our approach might also allow insights into the transferability of model systems for different pathways in malaria studies.

Importance

Malaria still causes about 400,000 deaths a year and is one the most studied infectious diseases. The disease is studied in mice and monkeys as lab models to derive potential therapeutic intervention in human malaria. Interactions between *Plasmodium* spp. and its hosts are either conserved across different host-parasite systems or idiosyncratic to those systems. Here we use correlation of gene expression from different RNA-Seq studies to infer common host-parasite interactions across human, mouse and monkey studies. We, firstly, find a set of very conserved interactors, worth further scrutiny in focussed laboratory experiments. Secondly, this work might help assess to which extent experiments and knowledge on different pathways can be transferred from models to humans for potential therapy.

Introduction

Transcriptomes are often analysed in a first attempt to understand molecular, cellular and organismic events. A comprehensive profile of RNA expression can be obtained using high-throughput sequencing of cDNA from reverse transcribed expressed RNA. Such RNA-sequencing (RNA-Seq) provides high technical accuracy at a reasonable cost, making it the current method of choice for transcriptomics [1,2].

In an infection experiment, RNA-Seq can assess host and pathogen transcriptomes simultaneously if RNA from both organisms is contained in a sample. On the one hand, it has been proposed to analyse transcriptomes of both organisms involved in an infection for a more complete understanding of the disease [3]–[5], such as virulence of a pathogen resulting from interlinked processes of both host and pathogen (host-pathogen interactions). This approach is called dual RNA-Seq. Some recent studies on malaria pathogenesis make use of dual RNA-Seq to study the host and the parasite simultaneously. On the other hand, researchers intending to study one of the two organisms, the target, might consider transcripts from the non-target organism, "contamination". Malaria research is indeed traditionally designed to target one organism, either the host or the parasite. Nevertheless, expression of "contaminant" transcripts potentially corresponds to response to stimuli.

Malaria is the most thoroughly investigated disease caused by a eukaryotic organism and the accumulation of these two types of studies, RNA-Seq with "contaminants" and intentional dual RNA-Seq, provides a rich resource for meta-analysis. In case of malaria, unlike in bacterial or viral

65 infections, both the parasite, *Plasmodium* spp*.* and the host are eukaryotic organisms with similar transcriptomes [4]. Their messenger RNA (mRNA) have a long poly-A tail at the 3'-end, therefore host and parasite mRNA is selected simultaneously when poly-dT priming is used to amplify polyadenylated mRNA transcripts [4,5]. This makes most malaria transcriptome datasets potentially suitable for dual RNA-Seq analysis.

70 Such a meta-analysis can use co-regulated gene expression to infer host-parasite interactions. In the mammalian intermediate host, *Plasmodium* spp. first multiply in the liver and then invade red blood cells (RBCs) for development and asexual expansion. While the nuclear machinery from both host and parasite cells produces mRNA in the liver, RBCs are enucleated and transcriptionally inactive in the mammalian host. In the blood stage of the infection, leukocytes or White Blood Cells

75 (WBCs) provide most of the transcriptomic response and are, thus, the source of host mRNA. Correlation of mRNA expression can be indicative of different types of biological interactions: protein products could be directly involved in the formation of complexes and might therefore be produced at quantities varying similarly under altered conditions. Alternatively, involvement in the same biological pathways can result in co-regulated gene expression without physical interactions.

80 This broad concept of interaction has long been exploited in single organisms (e.g. [6]-[8]). We (and others before [9,10]) propose to extrapolate this to interactions between the host and the parasite. It can be expected that a stimulus presented by the parasite to a host causes host immune response and the parasite in turn tries to evade this response, creating a cascade of genes co-regulated at different experimental conditions.

85 Here we showed that existing raw sequencing read datasets collectively present the potential to answer questions that have not been investigated by individual studies on malaria. We explored this potential by conducting a comparative meta-analysis of dual RNA-Seq transcriptomes of *Plasmodium* and its hosts in the blood stage of infection. Since rodent and simian malaria are often used as laboratory models for human malaria, we demonstrated the availability and suitability of

90 mRNA sequencing data from three evolutionarily close hosts - *Homo sapiens, Macaca mulatta* and

Mus musculus - and their respective *Plasmodium* species. We summarised available data and

conceived an approach to elucidate host-parasite interactions using orthologs across different

host-parasite pairs. We demonstrated that this approach provides meaningful results by cross-

validation of information content from networks from different host-parasite systems. We found that

95 these networks highlight known functional determinants of host-parasite interactions in broad

categories such as invasion and calcium ion homeostasis for the parasite and host immune

response and cadherin mediated cell-cell adhesion on the host side. We also found novel

interactions at a finer scale potentially worth further investigation.

Methods

Data review and curation of potentially suitable studies

Sequence data generated in biological experiments is submitted to one of the three mirroring

databases of the International Nucleotide Sequence Database Collaboration (INSDC): NCBI

Sequence Read Archive (SRA) [11], EBI European Nucleotide Archive (ENA) [12] and DDBJ

Sequence Read Archive (DRA) [13]. We used SRAdb v1.36.0 [14], a Bioconductor/R package [15],

105 to query SRA [11] for malaria RNA-Seq studies with the potential to provide host and *Plasmodium*

reads for our meta-analysis. We first selected studies with library_strategy "RNA-Seq" and with

"Plasmodium" in the study title, abstract or sample attributes fields using the function

dbGetQuery(). Then we used the getSRA() function with the query "(malaria OR Plasmodium) AND

RNA-Seq". This function searches all fields. We manually curated the combined results and added

110 studies based on a literature review using the terms described for the getSRA() function in PubMed

and Google Scholar. We disregarded studies on vectors, non-target hosts and those derived from

cell culture conditions devoid of transcriptionally active host cells. In these databases, all

experiments submitted under a single accession are given a single "study accession number" and

are collectively referred to as a "study" here onwards. We used prefetch and fastq-dump functions

115 from SRAtoolkit v2.8.2, to download all replicate samples (called runs in the databases) of the

selected studies. The curation and the download of the studies was performed on 21 January, 2019 and updated on 24 July, 2020.

Mapping and quantification of gene expression

We mapped sequencing reads onto concatenated host and parasite reference genomes using STAR v2.6.0c [16,17] with default parameters. Only runs with at least 70% uniquely mapping reads were considered for further analysis. Host and *Plasmodium* sp. genome assemblies and gene annotation files were downloaded from Ensembl version 43. Simultaneous mapping against both genomes should avoid non-specific mapping of reads in regions conserved between host and parasites. We quantified the sequencing reads mapped to exons using the countOverlaps function of the GenomicRanges package v3.7 [18].

Identification of co-expressed genes via correlation techniques

Reid and Berriman [9] recommended using empirical p-values for the analysis of gene co-expression. This allows not only to scrutinise housekeeping genes likely to show almost uniform expression under different experimental conditions but also requires fewer assumptions about the quality of the input data. We computed correlation indices for each gene-pair and obtained empirical p-values by comparison against null distributions computed using permutations of the given data, instead of assuming a theoretical null distribution. This is a robust way to estimate whether gene pairs are correlated because of specific events (treatment condition, time point) and not by chance (e.g., housekeeping genes) [19,20]. To obtain p-values corrected for multiple testing, since host and parasite genomes total at nearly 30,000 genes, the number of permutations would have to be around 10^{12} for a resolution of 0.1% FDR. As computational costs for these permutations are too high we used uncorrected p-values to rank genes as initially proposed by Reid and Berriman [9]. Here we only consider uncorrected p-values of 10^{-5} (in 100,000 permutations) as evidence of co-regulation.

Selection of runs for analysis

The construction of host-parasite gene co-expression networks require RNA-Seq runs to have both host and parasite transcript expression. To fulfil these criteria, we implemented thresholds based on host and parasite transcript expression, on the selection of runs. If a study had at least 5 runs with 50% detectable transcriptome expression from the host and from the parasite, we defined 2 thresholds for runs: 1) "intermediate" (int) with at least 50% detectable host and parasite transcriptome expression and 2) "stringent" (str) with 70% detectable host and parasite transcriptome expression. If a single study had less than 5 runs in the intermediate threshold, these runs were pooled with runs from other such studies.

To compare sub-datasets with runs selected at different thresholds and including all runs without thresholds, we calculated the Jaccard index for every pairwise combination of these datasets.

The Jaccard Index is defined as $\frac{|A \cap B|}{|A \cup B|}$ [21], where A and B are the set of bipartite edges in from two datasets. To include the maximum amount of host and parasite data and to use the best dataset from each study, we calculated the sum of all Jaccard Indices from each dataset. We chose the sub-dataset with the highest Jaccard index for a given study for further analysis and concatenated these representative studies into an "overall" dataset to compute possible conserved interactions from all host-parasite systems.

Identification of orthologs

Orthologs are genes in different species that have been derived from a single gene in the last common ancestor of those species by vertical descent. This means that orthologous genes from different species, in general, are likely to perform the same function [22].

Whole genome based predictions of the proteome for host species were downloaded from Ensembl release 43 and for *Plasmodium* from PlasmoDB release 37. Orthologs were identified using OrthoFinder v2.2.7 [23] wherein blastp [24] results for all-versus-all protein comparisons are

clustered using the MCL clustering algorithm [25] (tools bear default parameters for OrthoFinder).

165 One-to-one ortholog clusters, or orthogroups, were recovered for hosts and for parasites.

Network and functional analyses

Bipartite networks/graphs are graphs in which the nodes of the graph can be divided into two independent sets, and the nodes from one set connect only with nodes in the other set [26,27]. To visualise host-parasite interactions, bipartite networks were constructed with R package igraph

170 v1.2.5 [28] and Cytoscape v3.8.0 [29] where the nodes were the genes and their interaction were represented with an edge. To cluster nodes into network modules, we used the edge-betweenness algorithm (function edge_betweenness()) from igraph. The UpSetR package (v1.4.0) [30] in R was used to visualise intersection sizes of overlapping edges between networks.

For functional analysis of the genes found in host-parasite interactions from co-expression

175 analysis, their Gene Ontology [31] terms were found using the package topGO v2.36.0 and enriched using Kolmogorov-Smirnov test [32]. PlasmoDB [33] and Malaria.tools [34] databases were used for looking up gene functions and stage-associated transcript expression.

Statistical modeling

Using the R package fitdistrplus v1.1-1, we determined how Relative Growth Rate (RGR) [35] and

180 Mutagenesis Index Score (MIS) [36] are distributed. We used the betareg package in R to model RGR and MIS using centrality properties of our networks.
The centrality properties measured for our networks were node degree (DG), eigenvector centrality (EC) and betweenness (BW) using their dedicated functions in igraph package in R: "degree()", "betweenness()" and "eigen_centrality()". RGR and MIS (as response variables) were modelled

185 first with a single centrality measure as a predictor and then with combinations of two centrality measures: DG with EC and DG with BW. "Nested" models were compared based on likelihood-ratio tests and the more complex model was accepted when delta likelihood exceeded 2. In addition to this, the Akaike Information Criterion (AIC) was computed for each model to compare models and again models were considered differing in explanatory power at a delta-AIC of 2.

190　Models on different datasets (and different response variables) were compared without explicit

statistical testing discussing differences in p-values for variables in question.

In general, R (v3.4.3 - 4.0.0) [37] was used for analysis. All code is available at

https://doi.org/10.5281/zenodo.4535898 (as used in this manuscript) and github repository

https://github.com/parnika91/CompBio-Dual-RNAseq-Malaria (under further development).

Results

Potentially suitable studies for human, mouse and simian malaria

We found 63 potentially suitable studies [3,27,38-97] (Supplementary Table S1) on querying SRA

and performing a literature review. The host organism for 27 studies was *Homo sapiens*, for 26,

200　*Mus musculus* and for 10, *Macaca mulatta*. The infecting parasites were *P. falciparum, P. vivax* and

P. berghei in human studies (including artificial infections of human hepatocyte culture with *P.*

berghei), *P. yoelii, P. chabaudi* and *P. berghei* in mouse studies and *P. cynomolgi* and *P. coatneyi* in

macaque studies (Table 1). For 16 out of the 63 studies, the authors state that they intended to

simultaneously study host and parasite transcriptomes. This includes 8 studies from MaHPIC

205　(Malaria Host-Pathogen Interaction Center), based at Emory University, which made extensive

omics measurements in simian malaria. The original focus of the remaining 47 studies was on the

parasite in 23 and on the host in 24 cases. Our collection of studies comprises data derived from

blood and liver for all three host organisms human, mouse and macaque. Experiments performed

on mouse blood focus on the parasite instead of the host (12 vs. 0). Studies on human blood

210　infection focus more often on the host immune response than on the parasite (10 vs. 6). Liver and

spleen studies focus on host and parasite almost equally as often, with sources for host tissue in

this case being either mice (in vivo) or hepatoma cultures (in vitro).

We note that 21 of the 63 studies depleted or enriched specific classes of cells from their samples.

Seventeen blood-stage studies depleted or enriched host WBCs to focus expression analysis on

215　*Plasmodium* or host immunity, respectively. Assuming this depletion was imperfect, we tested

whether such samples present mRNA from both organisms. The physiologically asymptomatic liver

stage [72] is an appealing target for sporozoite-derived vaccines [98]-[100], but low parasite numbers make it difficult to study *Plasmodium* transcriptomes in liver. To reduce overwhelming host RNA levels, 10 out of the 14 liver studies sorted infected hepatoma cells from uninfected cells. Three out of the remaining 4 studies were among the ones interested in the host expression during the infection and 1 enriched for host cells.

Blood and liver samples from different studies and host parasite systems are potentially suitable for dual RNA-Seq analysis

A sample or run must provide sufficient gene expression from both host and parasite to be suitable for dual RNA-Seq co-expression analysis. Analysing the proportions of sequencing reads mapping to the host and parasite transcriptomes, we found that the original focus of the study was not always reflected (Fig 1). In many native samples (not enriched or depleted for host material), the number of host reads was found to be overwhelming. However, probably when parasitemia was very high, parasite transcriptomes were still recovered. Some examples are runs in the studies SRP032775, SRP029990 and ERP106769. Similarly, many studies using depletion or enrichment of a certain cell type prior to RNA sequencing show considerable expression of the non-target organism (enriched/depleted 'E' in Fig 1A). Examples are the studies ERP023892, ERP002273, ERP004598, ERP005730, ERP110375 and SRP112213. Studies depleting whole blood from leukocytes to focus on parasite transcriptomes still show considerable host gene expression and provide potentially suitable runs for the analysis of blood infection at lower intensities. Although, we note that this might come with the caveat that host expression might be biased by unequal depletion of particular cell types.

The number of sequencing reads mapping onto a transcriptome is a major determinant of the proportion of the organism's transcriptome detected as expressed (transcripts with at least one read mapped; Fig 1B and C). As expected, only runs that were sequenced deeply were able to capture the expression of a high proportion of the transcriptome. For both host and parasite, this plateaus at the total number of genes expressed in the transcriptome of the respective tissue in these organisms. Runs with an acceptable proportion of the transcriptome detected as expressed

were chosen in our analysis of blood stage malaria below. We propose that thresholds based on
the proportion of the transcriptome detected as expressed are most suitable for this selection of
runs. Table 2 provides an overview of the number of studies and runs available for each of these
analyses.

Evaluation of thresholds on transcriptome representation improve the analysis of co-regulated gene-expression

Deciding on the stringency of the thresholds applied to a dual RNA-Seq meta-analysis requires
additional analysis. Previous studies have reported that estimated proportions of the
transcriptionally active parts of the *P. falciparum* genome intraerythrocytic stages range from 60-
90% [101]-[103]. Here we test different thresholds that maximise the signal common between
blood studies - the number of edges shared between correlation networks for different studies (Fig
2).

36 studies investigated the blood stages of *Plasmodium.* 13 of them provided more than 5 runs at
all thresholds (schematic in Fig 2A) and were analysed as independent datasets. Similarly, 11
studies could be analysed separately with selection of runs at a "stringent" threshold of 70%
transcriptome coverage. To make use of runs not meeting thresholds from studies that couldn't be
analysed separately, 6 runs from 2 studies were pooled into a combined "humanPvivax" dataset
and 10 runs from 5 studies into a combined mouse dataset at intermediate thresholds. The
combined mouse dataset comprised 4 runs from 2 *P. berghei* studies, 5 from 2 *P. chabaudi* studies
and 1 from a *P. yoelii* study. 16 studies did not provide any runs meeting the thresholds and were
therefore excluded from the analysis. Based on the sum of Jaccard indices (see Methods) of the
datasets, we selected a total of 15 sub-datasets maximizing overlap between individual study
networks. We concatenated them to construct the "overall" dataset: 4 studies without thresholds
("all"), 7 at intermediate (50%; "int") and 4 at stringent (70% "str") thresholds (marked with an
asterix in Fig 2B,C,D). This means that a total of 915 runs were included in the "overall" dataset.

We can conclude from this analysis that the edges shared between different studies within and among host-parasite systems (see below for an exception regarding human-*P. falciparum* studies) outnumber random expectations and are highly significantly ($p < 0.001$, Fisher's exact test; between all selected studies), indicating a biological signal shared between datasets (Fig 2D). We suggest analysing the optimal version (selection of runs) of each study instead of setting a fixed (e.g. intermediate) threshold. Fig 2 indicates which different thresholds we deem most suitable for different studies and our selection for downstream analyses.

Across different studies, across different host-parasite systems

Knowledge of one-to-one orthologs between different hosts and different parasite species can be used in the next steps to integrate across different host-parasite systems (Fig 3). Humans share 18179 1:1 orthologous genes with macaques and 17089 with mice. 13986 genes are 1:1:1 orthologs among the three host species. Similarly, 4005 one-to-one groups of orthologous genes were found among the *Plasmodium* species (Fig 3A).

We have identified two different but interlinked workflows to reconstruct a consensus network of expression correlation. A first approach integrates data from different studies of one host-parasite system by simply appending expression profiles of their runs. Expression data across host-parasite systems is combined in orthogroups and correlations of gene expression are computed for all samples simultaneously. We refer to the results of this approach as the "overall" network (Fig 3C). It has the benefits that it uses and weighs all the information sources in one concise process.

Alternatively, we search for consensus networks by comparison of individual networks from different studies (Fig 3C). Similar to the approach of optimising expression thresholds (Fig 2, 3B), we compare overlapping edges in a multilayer network analysis. Sets of overlapping edges from multiple studies and host-parasite systems offer more control when querying for similar correlation in different layers representing different host-parasite systems. Similarly, correlations from different types of tissues (blood and liver) could be combined as multilayer networks in future work.

Correlating expression on the "overall" dataset resulted in a larger network (3.64 million edges, 12652 host genes, 3996 parasite genes) than on 13 out of 15 individual study datasets - two macaque datasets were exceptions (Fig 4). 528,883 bipartite edges are found only in the "overall" dataset and not in any individual dataset. 13 - 43% of edges from the "overall" dataset are recovered in each individual dataset (median 18.5%), indicating a substantial contribution from each individual dataset to the "overall" network. The most dominant study (presenting 43% of the edges in the "overall" network in it's own network) was a mouse study (study ID ERP004598 [91]) in which parasite gene expression was the primary focus. 91% of this study's bipartite edges are shared with at least one other individual study in addition to the "overall" network. The second most influential study was a dual RNA-Seq study (on monkey, study ID SRP118827 [27]) that shares 89% of its bipartite edges with at least one other study (Fig 4A). This not only illustrates the contributions of studies that initially focused on one organism ("with contaminants") in dual RNA-Seq analysis, but also that information can be pooled across different studies and host-parasite systems with our approach. Surprisingly, there were no common edges found between two human studies ([38] on Thai patients and [3] on Gambian patients, both infected with *P. falciparum*), possibly indicating differences in sampling from patients. Alternatively, this might hint towards idiosyncrasies in human blood studies using native samples with high parasitemia. The potential difficulty to apply RNA-Seq meta-analysis to these human samples highlights the need to transfer such data across model systems while controlling for concordance with the human system.

A "core" network of evolutionarily conserved interactions

The "overall" network is a highly connected graph with a total of ~3.64 million edges (Supplementary Table S2A). Even though this network provides some resolution relative to the ~56 million edges possible in a network of 13986 host and 4005 parasite genes, the resolution of this network might still be improved. We therefore defined a "core" network: using the "overall" network as a scaffold, we extracted edges that were recovered in at least one human study and at least one model organism. Using this definition, the resultant "core" network has 1876 host genes and 2050 parasite genes connected by 15324 edges (Supplementary Table S2B). A list of GO terms enriched or depleted in the genes of the "core" network are provided in Supplementary Table S3.

As expected, many GO terms were the same as in the "overall" network. Most GO terms in the "overall" network were enriched more strongly because of the higher number of genes in that network. However, many GO terms in the "overall" network were broader than in the "core" network. The recovery of more specific functions in the "core" network indicates higher resolution in this network (Table S3).

Gene co-expression explains gene essentiality

A node is important in a network if it is central, i.e., highly connected to other nodes in the network. The metrics node degree (DG), betweenness (BW) and eigenvector centrality (EC) are network centrality measures that can quantify this. Similarly, an essential gene is classically described as a gene performing a function necessary for the viability of a cell or organism. We thus hypothesized central nodes in our parasite networks would tend to be more essential genes of *Plasmodium*.

In quantitative terms, more essential genes are more important for cell growth and thus, the disruption of those genes leads to larger growth defects. Such data is available for *P. berghei* in mice, for which [35] reported "relative growth rates" (RGR) of mutant parasites reflecting the essentiality of a gene. A second study [36] reported *P. falciparum* growth rates impacted by genome wide mutagenesis covering 5,399 protein-coding genes. Here, the Mutagenesis Index Score (MIS) quantifies essentiality of a gene.

First, we tested the different network metrics for predicting gene essentiality (RGR and MIS) as response variables in beta-regression models. EC generally resulted in the best models and combination of the centrality measures did not improve the models in most cases (Supplementary Table S4). This might be explained by DG and EC being tightly correlated. We therefore report models with EC as a single predictor.

We then compared how well EC from different networks explained gene essentiality (Table 3, Fig 5): EC in the network derived from human - *P. falciparum* studies did significantly predict *P. falciparum* MIS and *P. berghei* RGR. EC from *P. berghei* networks, was also a significant predictor for both *P. falciparum* MIS and *P. berghei* RGR. Surprisingly, *P. berghei* network centrality was a better predictor of MIS in *P. falciparum* than the centrality in *P. falciparum* expression networks. EC

from both the "overall" network and the "core" network was a significant predictor for gene

350   essentiality. Considering effect sizes, p-values and AIC, we conclude that centrality in the *P. berghei* network best explains both RGR and MIS, followed by the metric from the "core" network. Given the much smaller size of the "core" network, the predictive power of centrality within this network is striking.

We can conclude from this analysis, that a) our co-expression networks capture biological

355   characteristics independently measured for the parasite and b) that parasite genes with a central position in the "core" network are important for the parasite's growth and survival. Highly connected genes in the intra-parasitic "core" network, however, might control the reaction to host signals without necessarily directly interacting with host genes.

Interacting parasite processes and host immune response

360   Based on functional annotation of our networks, we show that the correlation between host and parasite transcript expression can highlight known processes important in host-parasite interactions (Fig 6, Table S3). Biological processes for hosts consistent among almost all networks include "Cell adhesion by cadherin" and "Calcium-dependent cell adhesion". Cadherins are cell-adhesion proteins that depend on calcium. *Plasmodium* infection causes systemic endothelial

365   activation of host blood vessels when the infected RBCs (iRBCs) sequester on the lining of blood vessels. This is accompanied with an increased interaction of endothelial cells with WBCs, as cell adhesion molecules on the vessel lining direct WBC trafficking to infected RBC sequestration sites [104]. In addition, immune cells like B cells and monocytes express adhesion-related genes found in this GO term, like Fer (Tyrosine-protein kinase) and ICAM-1 (Intercellular adhesion molecule 1),

370   respectively [105]-[107]. To connect co-occurring host and parasite biological processes, we found associations between enriched (p-value <= 0.05) host and parasite GO terms based on the interactors of the genes in these enriched GO terms. From the "core" network, we identified a GO network of 617 host GO terms and 464 parasite GO terms. This analysis suggests that besides the detailed co-expression of gene products, broader enriched host and parasite processes or

375   pathways are likely interacting in malaria infection (Fig 7; Supplementary Table 5).

Another host biological process commonly found in our interaction networks is the response to oxidative stress combined with response to hydrogen peroxide and nitric oxide mediated signalling.

It has been found that iRBCs produce twice as many free OH^- radicals and H_2O_2 than uninfected RBCs [91,108]. This mechanism is believed to be a defence mechanism to abate the infection,

380 although the exact mechanism of parasite killing is still unclear. Nitric oxide (NO) production by the host is also believed to promote parasite death in malaria [109]. An interesting GO term observed in the results was "Killing by host of symbiont cells". This term included genes that were all involved in defence response towards microbes by invoking immune cells (eg, Neutrophil cytosol factor 1 and Cathepsin G), respiratory burst (Ncf1) and blood coagulation (eg, prothrombin).

385 Host-parasite interaction networks from all infection systems were consistently enriched for the parasite processes, "Calcium ion homeostasis" and "Late endosome to vacuole transport". Calcium ion homeostasis regulated by calmodulins and Ca^{2+}-dependent kinases (CDPKs) play roles in complex signalling pathways and are important for apicomplexan parasite virulence [110,111]. Vacuolar protein sorting-associated proteins-1, -2 and -46 (VPS1, 2 and 46) along with

390 serine/threonine protein kinase (VPS15) are the underlying signal for the enrichment of "Late endosome to vacuole transport" process and are interlinked with host immune reaction genes in our interaction networks. The endomembrane system is important for the parasite in order to invade the host cell, to establish the parasitophorous vacuole and to obtain nutrients. The Endosomal Sorting Complex Required for Transport (ESCRT) is involved in late endosome

395 formation as endosomes exists at different stages of formation - from early to late endosome, before finally fusing with the lysosome or vacuole (reviewed in [112]). The interrelation of the immune system and endosomal formation and sorting events should be a future focus of research including our deeper studies of correlated gene expression networks.

To test whether our networks (gene clusters) are indicative of correlated gene expression

400 originating from specific immune cells, we used immune cell marker genes established in [113]. These marker genes are specific for 6 types of immune cells in the event of a *Plasmodium* infection in the blood: neutrophils and monocytes (innate immunity), T cells and B cells (adaptive

immunity), and myeloid dendritic cells and NK cells (innate and adaptive). We found

(Supplementary Table S6) that specific markers from neutrophils were significantly

405   overrepresented (Fisher's exact test, p-value = 0.0089) in our "overall" network. Specific markers

from monocytes were not significant but overrepresented (Fisher's exact test, p-value = 0.19). This

likely indicates phagocytosis and killing of *Plasmodium* via respiratory burst in which neutrophils

and monocytes have a central role [114,115]. Specific markers for myeloid dendritic cells had

significant underrepresentation (Fisher's exact test, p-value = 0.0027). Marker genes for all other

410   cell types were underrepresented (Fisher's exact test, p-value > 0.05). In the "core" network,

marker genes were not significantly enriched, but those specific for monocytes, neutrophils and NK

cells were tentatively overrepresented. We then used the presence of these specific cell type

markers to annotate network clusters with the cell-type indicated (Fig 8). This resulted in 68

clusters in the "overall" network and 14 clusters in the "core" network annotated with a single cell

415   type and additional 49 and 5 clusters annotated with multiple cell types, respectively. Among the

most densely interconnected cluster in the "core" network with strong evidence of specific cell

types was a set of 203 host genes (three are specific for T cells, two for neutrophils) correlating in

their expression with only two parasite genes. One of them, a ubiquitin regulatory protein

(PBANKA_1222400), is connected to 269 host genes in total in the "core" network. It appears to be

420   uniformly expressed in all stages of the *Plasmodium* life cycle [34]. WLL-vs (Leu-Leu-Trp vinyl

sulfone) and WLW-vs (Trp-Leu-Trp vinyl sulfone) are proteasome inhibitors and likely drug

candidates [116]. In an experiment to study genetic changes mediating parasite recrudescence in

WLL-vs and WLW-vs resistant mutants, this protein's level was amplified in recrudescent lines

[117]. Thus it was suggested to confer low-grade resistance to such proteasome inhibitors. The

425   second *Plasmodium* gene is a putative dynamin-like protein (DrpC, PBANKA_1434100),

connected to 11 host genes. Dynamins are mechanochemical enzymes with a GTPase domain

and one of the functions of dynamin-related proteins (Drp) is to traffic vesicles [118]. DrpC is

conserved in apicomplexans and was shown to be localized at the base and periphery of the

mitochondrion in *Toxoplasma gondii*, implying a probable role in mitochondrial fission [119].

430   Another module of three host genes included a NK cell marker correlated with two parasite genes -

a 26S proteasome subunit (PBANKA_1206600) and ubiquitin fusion degradation protein UFD1

(PBANKA_1024700). The parasite 26S protease subunit [120]-[122] was reported as an essential

protein in *P. berghei* but not in *P. falciparum* [35,36]. In general, the ubiquitin-proteasome system

(UPS) is essential for quality control of proteins in eukaryotes. In *Plasmodium*, the UPS is

435 expressed across all life cycle stages and is speculated to be a promising drug target. It helps

adapt the parasite to stress, such as changes in oxidative environment and temperature

differences, ensuring the survival and virulence of the parasite. The *P. falciparum* ortholog of UFD1

(PF3D7_1418000) was reported to be a pathogenesis-related protein in an *in silico* module-based

subnetwork alignment approach using protein-protein interactions of *E. coli* as references [123].

440 One module had genes specific to three different immune cells - monocytes, myeloid DC and NK

cells - the highest number of distinct immune cell types represented in a "core" network cluster.

Among the 46 co-expressed parasite genes in this module were genes related to apicoplast

biogenesis, mitochondrial fission, transcription, vacuolar transport from Golgi apparatus to the

endoplasmic reticulum and Fe-S cluster assembly proteins.

445 Overall, these results (Supplementary Table S6) give an indication that certain gene expression

clusters in our networks might be associated with specific cells of the innate immune response

(neutrophils and monocytes). In addition, we show that parasite processes likely invoking this

immune response include the expression of genes involved in drug resistance and in vesicular

transport.

450 We next looked at overlaps between networks in our multilayer network analysis. Twenty

interactions of specific gene pairs were conserved across 6 individual networks and the "overall"

network (Fig 4A): this includes i) negatively correlated (Pearson's rho = -0.26) Kelch13 in

Plasmodium with Laminin subunit beta-2 (LAMB2) in the host and ii) negatively correlated

(Pearson's rho = -0.33) parasite 26S protease subunit and host LAMB2, both recovered only in

455 mouse and monkey studies. LAMB2 is an extracellular high-affinity receptor and is associated with

GO term "substrate adhesion-dependent cell spreading" to which infected erythrocytes (IE) have

been reported to bind, along with to other adhesion molecules such as ICAM, VECAM, etc. [124,

125] found that laminin levels in serum were increased in severe malaria. [126] later contradicted

this, finding that the binding of infected erythrocytes to endothelial receptors including laminin was

460 the same in severe and non-severe malaria. Kelch13 is a well-studied protein in which a C580Y

mutation in *P. falciparum* confers resistance to artemisinin, a frontline anti-malarial drug. It was found to be essential in both *P. falciparum* and *P. berghei* intraerythrocytic stages. Associated proteins and Kelch13 form an endocytic compartment and are essential for feeding on host hemoglobin. Kelch13 is also suggested to be a ubiquitin ligase with a role in the ubiquitin-proteasome system by labeling proteins for degradation [127]. Artemisinin and its derivatives (ART) are known to be activated by the products of hemoglobin degradation and promote cell death by increasing ER stress facilitated by the accumulation of polyubiquitinated proteins. Resistance-conferring mutations on Kelch13 reduce host cell and hemoglobin endocytosis and along with 26S proteasome system (which includes the 26S protease subunit, our other parasite gene in discussion here), maintain the normal proteasomal degradation pathway preventing cell death, and thus, resulting in resistance [128-132]. Our work here is, to our knowledge, the first suggestion that Kelch13 and LAMB2 might be involved in interacting host-parasite processes.

The gene pairs discussed above were recovered in 7 datasets altogether (6 individual + "overall"), but were not recovered by any of the human - *Plasmodium* studies and are therefore not present in the "core" network. Of the 20 interactions that were found in 7 datasets, 5 were also found in the "core" network and are indicated as such in the Supplementary Table 2C. Of these 5, one was an interaction between host protein Odorant-receptor ODR4 homolog (ENSG00000157181) and parasite Thioredoxin-like associated protein 2 (TLAP2; PBANKA_0518100). TLAP2 is a microtubule-associated complex. It was found to be dispensable in both *P. berghei* and *P. falciparum* [35,36]. TLAP2 is conserved in *Toxoplasma gondii* and *Plasmodium* spp. [133]. Along with other TLAP proteins, it is associated with protein TrxL1 (Thioredoxin-Like protein 1) and as a complex, coats cortical microtubules [133]. These coating proteins, as an ensemble, stabilize the cortical microtubules [134]. Host ODR4 is involved in trafficking of GPCR proteins [135] and in protein localisation [136].

A second interaction was between host biotinidase (ENSG00000169814) and parasite with protein kinase (PBANKA_1016200). In the host, biotinidase makes biotin available from dietary sources. Host biotin was found to be essential for *Plasmodium* survival in the liver stages but not in the blood stages [137]. In general, several mutations in biotinidase reduce biotin availability without

causing severe diseases. There have been suggestions that biotinidase mutations could be used

490   as antimalarial therapies, following evidence that there are biotinidase mutations in blood spots of

Somali populations [138]. Putative parasite protein kinase PBANKA_1016200 is a membrane

protein involved in protein phosphorylation and ATP binding. Even though it is dispensable in both

P. berghei and *P. falciparum*, in general, protein kinases have been an attractive subset of proteins

to be used as antimalarial targets (reviewed in [138]).

495   We propose that interactions between these proteins (physical protein-protein interaction) or

interlinkage of associated pathways may be worth further scrutiny in mechanistic investigations.

Conclusion

Our analysis recovers both very broad and well-known processes involved in host-parasite

500   interactions as well as strong interaction signals for narrower processes, including specific

interacting gene pairs. This indicates that our analysis of interaction networks might uncover novel

links between host and parasite processes helping in focus further research into detailed

mechanisms of host-parasite interactions. Further analyses building on the methodology and

concepts conceived here will involve networks derived from liver studies curated here, in which

505   host and parasite cells interact more directly than in the blood. Methodologically, we show that

analyses of expression from different host-parasite systems as multilayer networks can link host

and parasite network modules. Meta-analysis of co-expression networks can thus assess whether

host-parasite interactions are shared among different studies and treatments, different tissues and

different host-parasite systems. This allows the identification of processes at different scales of

510   molecular pathway organisation as depicted above.

Moreover, the sharing of pathways among co-expression networks from different host-parasite

systems might indicate how easily results on those pathways can be translated between different

host-parasite systems - here presented in a "core" network. Our results can thus provide insights

into how easily and for which pathways observations made in malaria models can be translated to

515   human malaria.

Conflict of Interest: The authors declare that there is no conflict of interest.

Author Contribution: EH and PM designed the study. PM obtained and analysed the data. EH, GB and PM wrote the manuscript. All authors contributed to the final version of the manuscript.

Acknowledgement: This research project was undertaken with the assistance of resources and services from the National Computational Infrastructure (NCI), which is supported by the Australian Government. We thank the statistics user group of the Leibniz-IZW for very helpful comments on our analysis.

Funding: This work was supported by the Alliance Berlin Canberra "Crossing Boundaries: Molecular Interactions in Malaria", which is co-funded by a grant from the Deutsche Forschungsgemeinschaft (DFG) for the International Research Training Group (IRTG) 2290 and the Australian National University. G.B.'s work is supported by the National Health and Medical Research Council of Australia (Grant APP1143008), the Australian Research Council (Grant DP180101494) and the National Natural Science Foundation of China (Grant 81772214)

References:

[1] Z. Wang, M. Gerstein, and M. Snyder, "RNA-Seq: a revolutionary tool for transcriptomics," Nat. Rev. | Genet., vol. VOLUME 10, no. january 2009, pp. 57–63, 2009.

[2] R. Stark, M. Grzelak, and J. Hadfield, "RNA sequencing: the teenage years," Nat. Rev. Genet., vol. 20, no. 11, pp. 631–656, 2019.

[3] Lee, Hyun Jae, Athina Georgiadou, Michael Walther, Davis Nwakanma, Lindsay B. Stewart, Michael Levin, Thomas D. Otto, David J. Conway, Lachlan J. Coin, and Aubrey J. Cunnington. 2018. "Integrated Pathogen Load and Dual Transcriptome Analysis of Systemic Host-Pathogen Interactions in Severe Malaria." *Science Translational Medicine* 10(447):eaar3619.

[4] A. J. Westermann, S. A. Gorski, and J. Vogel, "Dual RNA-seq of pathogen and host," Nat. Rev. Microbiol., vol. 10, no. 9,pp. 618–630, Sep. 2012.

[5] A. J. Westermann, L. Barquist, and J. Vogel, "Resolving host–pathogen interactions by dual RNA-seq," PLOS Pathog., vol. 13, no. 2, p. e1006033, Feb. 2017.

[6] J. M. Stuart, E. Segal, D. Koller, and S. K. Kim, "A Gene-Coexpression Network for Global Discovery of Conserved Genetic Modules," Science (80-.)., vol. 302, no. 5643, pp. 249 LP – 255, Oct. 2003.

[7] H. Bono and Y. Okazaki, "Functional transcriptomes: comparative analysis of biological pathways and processes in eukaryotes to infer genetic networks among transcripts," Curr. Opin. Struct. Biol.,vol. 12, no. 3, pp. 355–361, Jun. 2002.

[8] H. K. Lee, A. K. Hsu, J. Sajdak, J. Qin, and P. Pavlidis, "Coexpression analysis of human genes across many microarray data sets," Genome Res., vol. 14, no. 6, pp. 1085–1094, Jun. 2004.

[9] A. J. Reid and M. Berriman, "Genes involved in host–parasite interactions can be revealed by their correlated expression," Nucleic Acids Res., vol. 41, no. 3, pp. 1508–1518, Dec. 2012.

[10] Musungu, Bryan M., Deepak Bhatnagar, Robert L. Brown, Gary A. Payne, Greg OBrian, Ahmad M. Fakhoury, and Matt Geisler. 2016. "A Network Approach of Gene Co-Expression in the Zea Mays/Aspergillus Flavus Pathosystem to Map Host/Pathogen Interaction Pathways." *Frontiers in Genetics* 7(NOV):1–14.

[11] R. Leinonen, H. Sugawara, M. Shumway, and on behalf of the I. N. S. D. Collaboration, "The Sequence Read Archive," Nucleic Acids Res., vol. 39, no. suppl_1, pp. D19–D21, Nov. 2010.

[12] Leinonen, Rasko, Ruth Akhtar, Ewan Birney, Lawrence Bower, Ana Cerdeno-Tárraga, Ying Cheng, Iain Cleland, Nadeem Faruque, Neil Goodgame, Richard Gibson, Gemma Hoad, Mikyung Jang, Nima Pakseresht, Sheila Plaister, Rajesh Radhakrishnan, Kethi Reddy, Siamak Sobhany, Petra Ten Hoopen, Robert Vaughan, Vadim Zalunin, and Guy Cochrane. 2011. "The European Nucleotide Archive." *Nucleic Acids Research* 39(Database issue):D28–31.

[13] O. Ogasawara, Y. Kodama, J. Mashima, T. Kosuge, and T. Fujisawa, "DDBJ Database updates and computational infrastructure enhancement," Nucleic Acids Res., vol. 48, no. D1, pp. D45–D50, Jan. 2020.

[14] Y. Zhu, R. M. Stephens, P. S. Meltzer, and S. R. Davis, "SRAdb: query and use public next-generation sequencing data from within R," BMC Bioinformatics, vol. 14, no. 1, p. 19, 2013.

[15] Huber, Wolfgang, Vincent J. Carey, Robert Gentleman, Simon Anders, Marc Carlson, Benilton S. Carvalho, Hector Corrada Bravo, Sean Davis, Laurent Gatto, Thomas Girke, Raphael Gottardo, Florian Hahne, Kasper D. Hansen, Rafael A. Irizarry, Michael Lawrence, Michael I. Love, James MacDonald, Valerie Obenchain, Andrzej K. Oleś, Hervé Pagès, Alejandro Reyes, Paul Shannon, Gordon K. Smyth, Dan Tenenbaum, Levi Waldron, and Martin Morgan. 2015. "Orchestrating High-Throughput Genomic Analysis with Bioconductor." *Nature Methods* 12(2):115–21.

[16] Dobin, Alexander, Carrie A. Davis, Felix Schlesinger, Jorg Drenkow, Chris Zaleski, Sonali Jha, Philippe Batut, Mark Chaisson, and Thomas R. Gingeras. 2012. "STAR: Ultrafast Universal RNA-Seq Aligner." *Bioinformatics* 29(1):15–21.

[17] A. Dobin and T. R. Gingeras, "Mapping RNA-seq Reads with STAR," Curr. Protoc. Bioinforma., vol. 51, pp. 11.14.1-11.14.19, Sep. 2015.

[18] Lawrence, Michael, Wolfgang Huber, Hervé Pagès, Patrick Aboyoun, Marc Carlson, Robert Gentleman, Martin T. Morgan, and Vincent J. Carey. 2013. "Software for Computing and Annotating Genomic Ranges." *PLOS Computational Biology* 9(8):e1003118.

[19] T. A. Knijnenburg, L. F. A. Wessels, M. J. T. Reinders, and I. Shmulevich, "Fewer permutations, more accurate P-values," Bioinformatics, vol. 25, no. 12, pp. 161–168, 2009.

[20] B. Phipson and G. K. Smyth, "Permutation P-values should never be zero: Calculating exact P-values when permutations are randomly drawn," Stat. Appl. Genet. Mol. Biol., vol. 9, no. 1, pp.1–12, 2010.

[21] Jaccard, Paul (February 1912). "THE DISTRIBUTION OF THE FLORA IN THE ALPINE ZONE.1". New Phytologist. 11 (2): 37–50. doi:10.1111/j.1469-8137.1912.tb05611.x

[22] E. V Koonin, "An apology for orthologs - or brave new memes," Genome Biol., vol. 2, no. 4, p. comment1005.1, 2001.

[23] D. M. Emms and S. Kelly, "OrthoFinder: phylogenetic orthology inference for comparative genomics," Genome Biol., vol. 20, no. 1, p. 238, 2019.

[24] S. F. Altschul, W. Gish, W. Miller, E. W. Myers, and D. J. Lipman, "Basic local alignment search tool," J. Mol. Biol., vol. 215, no. 3, pp. 403–410, 1990.

[25] S. van Dongen and C. Abreu-Goodger, "Using MCL to Extract Clusters from Networks BT - Bacterial Molecular Networks: Methods and Protocols," J. van Helden, A. Toussaint, and D. Thieffry, Eds. New York, NY: Springer New York, 2012, pp. 281–295.

[26] J. Li, S. Wang, Z. Chen, and Y. Wang, "A Bipartite Network Module-Based Project to Predict Pathogen–Host Association," Front. Genet., vol. 10, no. January, pp. 1–9, 2020.

[27] Tang, Yan, Anuj Gupta, Swetha Garimalla, Mary R. Galinski, Mark P. Styczynski, Luis L. Fonseca, and Eberhard O. Voit. 2018. "Metabolic Modeling Helps Interpret Transcriptomic Changes during Malaria." *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 1864(6, Part B):2329–40.

[28] G. Csardi & T. Nepusz (2006). "The Igraph Software Package for Complex Network Research." InterJournal 2006, Complex Systems, 1695.

[29] Shannon, Paul, Andrew Markiel, Owen Ozier, Nitin S. Baliga, Jonathan T. Wang, Daniel Ramage, Nada Amin, Benno Schwikowski, and Trey Ideker. 2003. "Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks." Genome Research 13(11):2498–2504.

[30] A. Lex, N. Gehlenborg, H. Strobelt, R. Vuillemot, and H. Pfister, "UpSet: Visualization of Intersecting Sets," IEEE Trans. Vis. Comput. Graph., vol. 20, no. 12, pp. 1983–1992, Dec. 2014.

[31] P. Gaudet, N. Škunca, J. C. Hu, and C. Dessimoz, "Primer on the Gene Ontology," Methods Mol. Biol., vol. 1446, pp. 25–37, 2017.

[32] A. Alexa and J. Rahnenfuhrer, "Gene set enrichment analysis with topGO," Jan. 2007.

[33] Bahl, Amit, Brian Brunk, Jonathan Crabtree, Martin J. Fraunholz, Bindu Gajria, Gregory R. Grant, Hagai Ginsburg, Dinesh Gupta, Jessica C. Kissinger, Philip Labo, Li Li, Matthew D. Mailman, Arthur J. Milgram, David S. Pearson, David S. Roos, Jonathan Schug,

Christian J. Stoeckert Jr, and Patricia Whetzel. 2003. "PlasmoDB: The Plasmodium Genome Resource. A Database Integrating Experimental and Computational Data." *Nucleic Acids Research* 31(1):212–15.

[34] Q. W. Tan and M. Mutwil, "Malaria.tools—comparative genomic and transcriptomic database for Plasmodium species," Nucleic Acids Res., vol. 48, no. D1, pp. D768–D775, Aug. 2019.

[35] Bushell, Ellen, Ana Rita Gomes, Theo Sanderson, Burcu Anar, Gareth Girling, Colin Herd, Tom Metcalf, Katarzyna Modrzynska, Frank Schwach, Rowena E. Martin, Michael W. Mather, Geoffrey I. McFadden, Leopold Parts, Gavin G. Rutledge, Akhil B. Vaidya, Kai Wengelnik, Julian C. Rayner, and Oliver Billker. 2017. "Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes." Cell 170(2):260-272.e8.

[36] Zhang, Min, Chengqi Wang, Thomas D. Otto, Jenna Oberstaller, Xiangyun Liao, Swamy R. Adapa, Kenneth Udenze, Iraad F. Bronner, Deborah Casandra, Matthew Mayho, Jacqueline Brown, Suzanne Li, Justin Swanson, Julian C. Rayner, Rays H. Y. Jiang, and John H. Adams. 2018. "Uncovering the Essential Genes of the Human Malaria Parasite Plasmodium Falciparum by Saturation Mutagenesis." Science 360(6388).

[37] R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/.

[38] Yamagishi, Junya, Anna Natori, Mohammed E. M. Tolba, Arthur E. Mongan, Chihiro Sugimoto, Toshiaki Katayama, Shuichi Kawashima, Wojciech Makalowski, Ryuichiro Maeda, Yuki Eshita, Josef Tuda, and Yutaka Suzuki. 2014. "Interactive Transcriptome Analysis of Malaria Patients and Infecting Plasmodium Falciparum." *Genome Research* 24(9):1433–44. **2 SRA studies are associated with this paper: DRP000987 and DRP001953.**

[39] SRP059851. https://www.ncbi.nlm.nih.gov/sra/?term=SRP059851. Accessed: 21-01-2019.

[40] ERP017542. https://www.ncbi.nlm.nih.gov/sra/?term=ERP017542 Accessed: 21-01-2019.

[41] ERP023982. https://www.ncbi.nlm.nih.gov/sra/?term=ERP023982 Accessed: 21-01-2019.

[42] ERP105548 https://www.ncbi.nlm.nih.gov/sra/?term=ERP105548 Accessed: 21-01-2019.

[43] ERP107298 https://www.ncbi.nlm.nih.gov/sra/?term=ERP107298 Accessed: 21-01-2019.

[44] Tran, T. M., Jones, M. B., Ongoiba, A., Bijker, E. M., Schats, R., Venepally, P., Skinner, J., Doumbo, S., Quinten, E., Visser, L. G., Whalen, E., Presnell, S., O'connell, E. M., Kayentao, K., Chaussabel, D., Lorenzi, H., Nutman, T. B., Ottenhoff, T. H. M., Haks, M. C., Traore, B., Kirkness, E. F., Sauer-Wein, R. W., and Crompton, P. D. Transcriptomic evidence for modulation of host inflammatory responses during febrile Plasmodium falciparum malaria. 6 (2016). doi: https://doi.org/10.1038/srep31291.

[45] Jaijyan, D. K., Singh, H., and Singh, A. P. A sporozoite-and liver stage-expressed tryptophan-rich protein plays an auxiliary role in Plasmodium liver stage development and is a potential vaccine candidate. 290 (2015). doi: https://doi.org/10.1074/jbc.M114.588129.

[46] Rojas-Peña, M. L., Vallejo, A., Herrera, S., Gibson, G., and Arévalo-Herrera, M. Transcription profiling of malaria-nïive and semi-immune colombian volunteers in a Plasmodium vivax sporozoite challenge. 9, 8 (08 2015), 1–18. doi: https://doi.org/10.1371/journal.pntd.0003978.

[47] SRP059851 https://www.ncbi.nlm.nih.gov/sra/?term=SRP059851 Accessed: 21-01-2019.

[48] SRP071199 https://www.ncbi.nlm.nih.gov/sra/?term=SRP071199 Accessed: 21-01-2019.

[49] Rojas-Pena, M. C., Arafat, D. C., Velasquez, J. M., Garimalla, S. C., Arevalo-Herrera, M., Herrera, S., and Gibson, G. Profiling gene expression of the host response to a Plasmodium vivax irradiated sporozoite immunization and infectious challenge. (2018). doi: https://doi.org/10.1101/501957.

[50] SRP083918 https://www.ncbi.nlm.nih.gov/sra/?term=SRP083918 Accessed: 21-01-2019.

[51] Götz, A., Tang, M. S., Ty, M. C., Arama, C., Ongoiba, A., Doumtabe, D., Traore, B., Crompton, P. D., Loke, P., and Rodriguez, A. Atypical activation of dendritic cells by Plasmodium falciparum. Proceedings of the National Academy of Sciences 114, 49 (2017), E10568–E10577. doi: https://doi.org/10.1073/pnas.1708383114.

[52 Rothen, J., Murie, C., Carnes, J., Anupama, A., Abdulla, S., Chemba, M., Mpina, M., Tanner, M., Lee Sim, B. K., Hoffman, S. L., Gottardo, R., Daubenberger, C., and Stuart, K. Whole blood transcriptome changes following controlled human malaria infection in malaria pre-exposed volunteers correlate with parasite prepatent period. 13, 6 (06 2018), 1–18. doi: https://doi.org/10.1371/journal.pone.0199392

[53] Kim, A., Popovici, J., Vantaux, A., Samreth, R., Bin, S., Kim, S., Roesch, C., Liang, L., Davies, H., Felgner, P., Herrera, S., Arévalo-Herrera, M., MÉUARD, D., and Serre, D. Characterization of P. vivax blood stage transcriptomes from field isolates reveals similarities among infections and complex gene isoforms. 7 (2017). doi: https://doi.org/10.1038/s41598-017-07275-9.

[54] LaMonte, G., Orjuela-Sanchez, P., Wang, L., Li, S., Swann, J., Cowell, A., Zou, B. Y., Abel-Haleem Mohamed, A., Villa-Galarce, Z., Moreno, M., Tong-Rios, C., Vinetz, J., Lewis, N., and Winzeler, E. A. Dual RNAseq identifies human mucosal immunity protein muc13 as a hallmark of Plasmodium exoerythrocytic infection. 10 (2019). doi: https://doi.org/10.1038/s41467-019-08349-0.

[55] Quin, J. E., Bujila, I., Chérif, M., Sanou, G. S., Qu, Y., Vafa Homann, M., Rolicka, A., Sirima, S. B., O'connell, M. A., Lennartsson, A., Troye-Blomberg, M., Nebie, I., and Östlund Farrants, A.-K. Major transcriptional changes observed in the fulani, an ethnic group less susceptible to malaria. 6 (sep 2017), e29156. doi: https://doi.org/10.7554/eLife.29156.

[56] Gural, N., Mancio-Silva, L., Miller, A. B., Galstian, A., Butty, V. L., Levine, S. S., Patrapuvich, R., Desai, S. P., Mikolajczak, S. A., Kappe, S. H., Fleming, H. E., March, S., Sattabongkot, J., and Bhatia, S. N. In vitro culture, drug sensitivity, and transcriptome of Plasmodium vivax hypnozoites. 23, 3 (2018), 395–406.e4. doi: https://doi.org/10.1016/j.chom.2018.01.002.

[57] SRP143371 https://www.ncbi.nlm.nih.gov/sra/?term=SRP143373 Accessed: 21-01-2019

[58] Cubi, R., Vembar, S. S., Biton, A., Franetich, J.-F., Bordessoulles, M., Sossau, D., Zanghi, G., Bosson-Vanga, H., Benard, M., Moreno, A., Dereuddre-Bosquet, N., Le Grand, R., Scherf, A., and Mazier, D. Laser capture microdissection enables transcriptomic analysis of dividing and quiescent liver stages of Plasmodium relapsing species. 19, 8 (2017), e12735. doi: https://doi.org/10.1111/cmi.12735.

[59] Van Der Wel, A. V., Roma, G., Gupta, D. K., Schuierer, S., Nigsch, F., Carbone, W., Zeeman, A.-M., Lee, B. H., Hofman, S. O., Faber, B. W., Knehr, J., Pasini, E., Kinzel, B., Bifani, P., Bonamy, G. M. C., Bouwmeester, T., Kocken, C. H. M., and Diagana, T. T. A comparative transcriptomic analysis of replicating and dormant liver stages of the relapsing malaria parasite Plasmodium cynomolgi. 6 (2017). doi: https://doi.org/10.7554/eLife.29605.

[60] Joyner, C., Moreno, A., Meyer, E. V. S., Cabrera-Mora, M., The MAHPIC CONSORTIUM, Kong, A., Ibegbu, C., Arafat, D., Jones, D., Nace, D. P., Voit, E. O., Pohl, J., Humphrey, J., Debarry, J., Sullivan, J., Gutierrez, J. B., Kevin, K. J., Fonseca, L. L., Styczynski, M., Nural, M., Pruett, S. T., Lapp, S. A., Pakala, S., Lamb, T. J., Williams, T., Uppal, K., Tran, V., Yin, W., Yan, Y., Kissinger, J. C., Barnwell, J. W., and Galinski, M. R. Plasmodium cynomolgi infections in rhesus macaques display clinical and parasitological features pertinent to modelling vivax malaria pathology and relapse infections. 15, 1 (Sep 2016), 451. doi: https://doi.org/10.1186/s12936-016-1480-6.

[61] SRP116117 https://www.ncbi.nlm.nih.gov/sra/?term=SRP116117 Accessed: 21-01-2019

[62] SRP116593 https://www.ncbi.nlm.nih.gov/sra/?term=SRP116593 Accessed: 21-01-2019

[63] SRP116793 https://www.ncbi.nlm.nih.gov/sra/?term=SRP116793 Accessed: 21-01-2019

[64] SRP118503 https://www.ncbi.nlm.nih.gov/sra/?term=SRP118503 Accessed: 21-01-2019

[65] SRP118827 https://www.ncbi.nlm.nih.gov/sra/?term=SRP118827 Accessed: 21-01-2019

[66] SRP118996 https://www.ncbi.nlm.nih.gov/sra/?term=SRP118996 Accessed: 21-01-2019

[67] Ogun, S. A., Tewari, R., Otto, T. D., Howell, S. A., Knuepfer, E., Cunningham, D. A., Xu, Z., Pain, A., and Holder, A. A. Targeted disruption of py235ebp-1: Invasion of erythrocytes by Plasmodium yoelii using an alternative Py235 erythrocyte binding protein. 7, 2 (02 2011), 1–14. doi: https://doi.org/10.1371/journal.ppat.1001288.

[68] Lawton, J., Brugat, T., Yan, Y. X., Reid, A. J., Böhme, U., Otto, T. D., Pain, A., Jackson, A., Berriman, M., Cunningham, D., Preiser, P., and Langhorne, J.Characterization and gene expression analysis of the cir multi-gene family of Plasmodium chabaudi chabaudi (AS). 13, 1 (Mar 2012), 125. doi: https://doi.org/10.1186/1471-2164-13-125.

[69] Spence, P. J., Jarra, W., Lévy, P., Reid, A. J., Chappell, L., Brugat, T., Sanders, M., Berriman, M., and Langhorne, J. Vector transmission regulates immune control of Plasmodium virulence. 498 (2013). doi: https://doi.org/10.1038/nature12231

[70] ERP002116 https://www.ncbi.nlm.nih.gov/sra/?term=ERP002116 Accessed: 21-01-2019

[71] ERP002273 https://www.ncbi.nlm.nih.gov/sra/?term=ERP002273 Accessed: 21-01-2019

[72] ERP004042 https://www.ncbi.nlm.nih.gov/sra/?term=ERP004042 Accessed: 21-01-2019

[73] Guttery, D. S., Poulin, B., Ramaprasad, A., Wall, R. J., Ferguson, D. J., Brady, D., Patzewitz, E.- M., Whipple, S., Straschil, U., Wright, M. H., Mohamed, A. M., Radhakrishnan, A., Arold, S. T., Tate, E. W., Holder, A. A., Wickstead, B., Pain, A., and Tewari, R. Genome-wide functional analysis of Plasmodium protein phosphatases reveals key regulators of parasite development and differentiation. 16, 1 (2014), 128–140. doi: https://doi.org/10.1016/j.chom.2014.05.020.

[74] ERP005730 https://www.ncbi.nlm.nih.gov/sra/?term=ERP005730 Accessed: 21-01-2019

[75] ERP106769 https://www.ncbi.nlm.nih.gov/sra/?term=ERP106769 Accessed: 21-01-2019

[76] Lin, J. W., Reid, A. J., Cunningham, D., Böhme, U., Tumwine, I., Keller-Mclaughlin, S., Sanders, M., Berriman, M., & Langhorne, J. (2018). Genomic and transcriptomic comparisons of closely related malaria parasites differing in virulence and sequestration pattern. *Wellcome open research*, *3*, 142. https://doi.org/10.12688/wellcomeopenres.14797.2

[77] Miller, J. L., Sack, B. K., Baldwin, M., Vaughan, A. M., and Kappe, S. H. Interferon-mediated innate immune responses against malaria parasite liver stages. 7, 2 (2014), 436–447. doi: https://doi.org/10.1016/j.celrep.2014.03.018.

[78] Li, J., Cai, B., Qi, Y., Zhao, W., Liu, J., Xu, R., Pang, Q., Tao, Z., Hong, L., Liu, S., Leerkes, M., Quiñones, M., and Su, X.-Z. UTR introns, antisense rna and differentially spliced transcripts between Plasmodium yoelii subspecies. 15, 1 (Jan 2016), 30. doi: https://doi.org/10.1186/s12936-015-1081-9.

[79] Torre, S., Polyak, M. J., Langlais, D., Fodil, N., Kennedy, J. M., Radovanovic, I., Berghout, J., Leiva-Torres, G. A., Krawczyk, C. M., Ilangumaran, S., Mossman, K., Liang, C., Knobeloch, K.-P., Healy, L. M., Antel, J., Arbour, N., Prat, A., Majewski, J., Lathrop, M., Vidal, S. M., and Gros, P. USP15 regulates type I interferon response and is required for pathogenesis of neuroinflammation. 18, 54 (2016). doi: https://doi.org/10.1038/ni.3581.

[80] Yeoh, L. M., Goodman, C. D., Mollard, V., McFadden, G. I., and Ralph, S. A. Comparative transcriptomics of female and male gametocytes in Plasmodium berghei and the evolution of sex in alveolates. 18, 1 (Sep 2017), 734. doi: https://doi.org/10.1186/s12864-017-4100-0.

[81] SRP109709 https://www.ncbi.nlm.nih.gov/sra/?term=SRP109709 Accessed: 21-01-2019

[82] RP112213 https://www.ncbi.nlm.nih.gov/sra/?term=SRP112213 Accessed: 21-01-2019

[83] Gabryšová, L., Alvarez-Martinez, M., Luisier, R., Cox, L. S., Sodenkamp, J., Hosking, C., Pérez-Mazliah, D., Whicher, C., Kannan, Y., Potempa, K., Wu, X., Bhaw, L., Wende, H., Sieweke, M. H., Elgar, G., Wilson, M., Briscoe, J., Metzis, V., Langhorne, J., Luscombe, N. M., and O'garra, A. c-Maf controls immune responses by regulating disease-specific gene networks and repressing IL-2 in CD4+ T cells. 19 (2018). doi: https://doi.org/10.1038/s41590-018-0083-5.

[84] Mamedov, M. R., Scholzen, A., Nair, R. V., Cumnock, K., Kenkel, J. A., Oliveira, J. H. M., Trujillo, D. L., Saligrama, N., Zhang, Y., Rubelt, F., Schneider, D. S., Hsiu Chien, Y., Sauerwein, R. W., and Davis, M. M. A macrophage colony-stimulating-factor-producing Tcell subset prevents malarial parasitemic recurrence. 48, 2 (2018), 350–363.e7. doi: https://doi.org/10.1016/j.immuni.2018.01.009.

[85] Burrack, K. S., Huggins, M. A., Taras, E., Dougherty, P., Henzler, C. M., Yang, R., Alter, S., Jeng, E. K., Wong, H. C., Felices, M., Cichocki, F., Miller, J. S., Hart, G. T., Johnson, A. J., Jameson, S. C., and Hamilton, S. E. Interleukin-15 complex treatment protects mice from cerebral malaria by inducing interleukin-10-producing natural killer cells. 48, 4 (2018), 760–772.e4. doi: https://doi.org/10.1016/j.immuni.2018.03.012.

[86] Hirako, I. C., Assis, P. A., Hojo-Souza, N. S., Reed, G., Nakaya, H., Golenbock, D. T., Coimbra, R. S., and Gazzinelli, R. T. Daily rhythms of tnf expression and food intake regulate synchrony of Plasmodium stages with the host circadian cycle. 23, 6 (2018), 796–808.e6. doi: https://doi.org/10.1016/j.chom.2018.04.016.

[87] Lai, S. M., Sheng, J., Gupta, P., Renia, L., Duan, K., Zolezzi, F., Karjalainen, K., Newell, E. W., and Ruedl, C. Organ-specific fate, recruitment, and refilling dynamics of tissue-resident macrophages during blood-stage malaria. 25, 11 (2018), 3099–3109.e3. doi: https://doi.org/10.1016/j.celrep.2018.11.059.

[88] SRP164767 https://www.ncbi.nlm.nih.gov/sra/?term=SRP164767 Accessed: 21-01-2019

[89] SRP164768 https://www.ncbi.nlm.nih.gov/sra/?term=SRP164767 Accessed: 21-01-2019

[90] SRP261098 https://www.ncbi.nlm.nih.gov/sra/?term=srp261098 Accessed 24-07-2020

[91] ERP108490 https://www.ncbi.nlm.nih.gov/sra/?term=ERP108490 Accessed 24-07-2020

[92] ERP114892 https://www.ncbi.nlm.nih.gov/sra/?term=ERP114892 Accessed 24-07-2020

[93] M. Toro-Moreno, K. Sylvester, T. Srivastava, D. Posfai, and E. R. Derbyshire, "RNA-Seq Analysis Illuminates the Early Stages of Plasmodium& Liver Infection," *MBio*, vol. 11, no. 1, pp. e03234-19, Feb. 2020.

[94] Tweedell, Rebecca E., Dingyin Tao, Timothy Hamerly, Tanisha M. Robinson, Simon Larsen, Alexander G. B. Grønning, Alessandra M. Norris, Jonas G. King, Henry Chun Hin Law, Jan Baumbach, Elke S. Bergmann-Leitner, and Rhoel R. Dinglasan. 2019. "The Selection of a Hepatocyte Cell Line Susceptible to Plasmodium Falciparum Sporozoite Invasion That Is Associated With Expression of Glypican-3 ." *Frontiers in Microbiology* 10:127.

[95] Bradwell, Katie R., Drissa Coulibaly, Abdoulaye K. Koné, Matthew B. Laurens, Ahmadou Dembélé, Youssouf Tolo, Karim Traoré, Amadou Niangaly, Andrea A. Berry, Bourema Kouriba, Christopher V Plowe, Ogobara K. Doumbo, Kirsten E. Lyke, Shannon Takala-Harrison, Mahamadou A. Thera, Mark A. Travassos, and David Serre. 2020. "Host and Parasite Transcriptomic Changes upon Successive ≪Span Class="Named-Content Genus-Species" Id="Named-Content-1">Plasmodium Falciparum≪/Span> Infections in Early Childhood" edited by P. Flórez de Sessions. *MSystems* 5(4):e00116-20.

[96] Kurup, S. P., Anthony, S. M., Hancox, L. S., Vijay, R., Pewe, L. L., Mioffer, S. J., Sompallae, R., Janse, C. J., Khan, S. M., & Harty, J. T. (2019). Monocyte-Derived CD11c+ Cells Acquire Plasmodium from Hepatocytes to Prime CD8 T Cell Immunity to Liver-Stage Malaria. *Cell host & microbe*, 25(4), 565–577.e6. https://doi.org/10.1016/j.chom.2019.02.014

[97] Subudhi, A. K., O'Donnell, A. J., Ramaprasad, A., Abkallo, H. M., Kaushik, A., Ansari, H. R., Abdel-Haleem, A. M., Ben Rached, F., Kaneko, O., Culleton, R., Reece, S. E., & Pain, A. (2020). Malaria parasites regulate intra-erythrocytic development duration via serpentine receptor 10 to coordinate with host rhythms. *Nature communications*, 11(1), 2763. https://doi.org/10.1038/s41467-020-16593-y

[98] M. R. Hollingdale and M. Sedegah, "Development of whole sporozoite malaria vaccines," Expert Rev. Vaccines, vol. 16, no. 1, pp. 45–54, Jan. 2017.

[99] Seder, Robert A., Lee-Jah Chang, Mary E. Enama, Kathryn L. Zephir, Uzma N. Sarwar, Ingelise J. Gordon, LaSonji A. Holman, Eric R. James, Peter F. Billingsley, Anusha Gunasekera, Adam Richman, Sumana Chakravarty, Anita Manoj, Soundarapandian Velmurugan, MingLin Li, Adam J. Ruben, Tao Li, Abraham G. Eappen, Richard E. Stafford, Sarah H. Plummer, Cynthia S. Hendel, Laura Novik, Pamela J. M. Costner, Floreliz H. Mendoza, Jamie G. Saunders, Martha C. Nason, Jason H. Richardson, Jittawadee Murphy, Silas A. Davidson, Thomas L. Richie, Martha Sedegah, Awalludin Sutamihardja, Gary A. Fahle, Kirsten E. Lyke, Matthew B. Laurens, Mario Roederer, Kavita

Tewari, Judith E. Epstein, B. Kim Lee Sim, Julie E. Ledgerwood, Barney S. Graham, and Stephen L. Hoffman. 2013. "Protection Against Malaria by Intravenous Immunization with a Nonreplicating Sporozoite Vaccine." *Science* 341(6152):1359 LP – 1365.

[100] E. M. Bijker, S. Borrmann, S. H. Kappe, B. Mordmüller, B. K. Sack, and S. M. Khan, "Novel approaches to whole sporozoite vaccination against malaria," Vaccine, vol. 33, no. 52, pp. 7462–7468, Dec. 2015.

[101] Chappell, Lia, Philipp Ross, Lindsey Orchard, Timothy J. Russell, Thomas D. Otto, Matthew Berriman, Julian C. Rayner, and Manuel Llinás. 2020. "Refining the Transcriptome of the Human Malaria Parasite Plasmodium Falciparum Using Amplification-Free RNA-Seq." *BMC Genomics* 21(1):395.

[102] Z. Bozdech, M. Llinás, B. L. Pulliam, E. D. Wong, J. Zhu, and J. L. DeRisi, "The Transcriptome of the Intraerythrocytic Developmental Cycle of Plasmodium falciparum," PLOS Biol., vol. 1, no. 1, p. e5, Aug. 2003.

[103] Otto, Thomas D., Daniel Wilinski, Sammy Assefa, Thomas M. Keane, Louis R. Sarry, Ulrike Böhme, Jacob Lemieux, Bart Barrell, Arnab Pain, Matthew Berriman, Chris Newbold, and Manuel Llinás. 2010. "New Insights into the Blood-Stage Transcriptome of Plasmodium Falciparum Using RNA-Seq." *Molecular Microbiology* 76(1):12–24.

[104] Turner, G. D., V. C. Ly, T. H. Nguyen, T. H. Tran, H. P. Nguyen, D. Bethell, S. Wyllie, K. Louwrier, S. B. Fox, K. C. Gatter, N. P. Day, T. H. Tran, N. J. White, and A. R. Berendt. 1998. "Systemic Endothelial Activation Occurs in Both Mild and Severe Malaria. Correlating Dermal Microvascular Endothelial Cell Phenotype and Soluble Cell Adhesion Molecules with Disease Severity." *The American Journal of Pathology* 152(6):1477–87.

[105] Möst, J., W. Schwaeble, J. Drach, A. Sommerauer, and M. P. Dierich. 1992. "Regulation of the Expression of ICAM-1 on Human Monocytes and Monocytic Tumor Cell Lines." The Journal of Immunology 148(6):1635 LP – 1642.

[106] Uhlén, Mathias, Linn Fagerberg, Björn M. Hallström, Cecilia Lindskog, Per Oksvold, Adil Mardinoglu, Åsa Sivertsson, Caroline Kampf, Evelina Sjöstedt, Anna Asplund, IngMarie Olsson, Karolina Edlund, Emma Lundberg, Sanjay Navani, Cristina Al-Khalili Szigyarto, Jacob Odeberg, Dijana Djureinovic, Jenny Ottosson Takanen, Sophia Hober, Tove Alm, Per-Henrik Edqvist, Holger Berling, Hanna Tegel, Jan Mulder, Johan Rockberg, Peter Nilsson, Jochen M. Schwenk, Marica Hamsten, Kalle von Feilitzen, Mattias Forsberg, Lukas Persson, Fredric Johansson, Martin Zwahlen, Gunnar von Heijne, Jens Nielsen, and Fredrik Pontén. 2015. "Tissue-Based Map of the Human Proteome." Science 347(6220):1260419.

[107] Aguilar, Stephanie Vargas, Oscar Aguilar, Rhys Allan, El Ad David Amir, Veronique Angeli, Maxim N. Artyomov, Natasha Asinovski, Jilian Astarita, K. Frank Austen, Geetika Bajpai, et al., and The Immunological Genome Project. 2020. "ImmGen at 15." Nature Immunology 21(7):700–703.

[108] H. Atamna and H. Ginsburg, "Origin of reactive oxygen species in erythrocytes infected with Plasmodium falciparum," Mol. Biochem. Parasitol., vol. 61, no. 2, pp. 231–241, 1993.

[109] Percário, Sandro, Danilo R. Moreira, Bruno A. Q. Gomes, Michelli E. S. Ferreira, Ana Carolina M. Gonçalves, Paula S. O. C. Laurindo, Thyago C. Vilhena, Maria F. Dolabela, and Michael D. Green. 2012. "Oxidative Stress in Malaria." *International Journal of Molecular Sciences* 13(12):16346–72.

[110] C. R. S. Garcia, "Calcium Homeostasis and Signaling in the Blood-stage Malaria Parasite," Parasitol. Today, vol. 15, no. 12, pp. 488–491, Dec. 1999.

[111] S. Lourido and S. N. J. Moreno, "The calcium signaling toolkit of the Apicomplexan parasites Toxoplasma gondii and Plasmodium spp," Cell Calcium, vol. 57, no. 3, pp. 186–193, 2015.

[112] E. Jimenez-Ruiz, J. Morlon-Guyot, W. Daher, and M. Meissner, "Vacuolar protein sorting mechanisms in apicomplexan parasites," Mol. Biochem. Parasitol., vol. 209, no. 1–2, pp. 18–25, 2016.

[113] Vallejo, Andres F., Robert C. Read, Myriam Arevalo-Herrera, Sócrates Herrera, Tim Elliott, and Marta E. Polak. 2018. "Malaria Systems Immunology: Plasmodium Vivax Induces Tolerance during Primary Infection through Dysregulation of Neutrophils and Dendritic Cells." Journal of Infection 77(5):440–47.

[114] Aitken, Elizabeth H., Agersew Alemu, and Stephen J. Rogerson. 2018. "Neutrophils and Malaria." Frontiers in Immunology 9:3005.

[115] Ortega-Pajares, Amaya and Stephen J. Rogerson. 2018. "The Rough Guide to Monocytes in Malaria Infection." Frontiers in Immunology 9:2888.

[116] Li, Hao, Anthony J. O'Donoghue, Wouter A. van der Linden, Stanley C. Xie, Euna Yoo, Ian T. Foe, Leann Tilley, Charles S. Craik, Paula C. A. da Fonseca, and Matthew Bogyo. 2016. "Structure- and Function-Based Design of Plasmodium-Selective Proteasome Inhibitors." Nature 530(7589):233–36.

[117] Stokes, Barbara H., Euna Yoo, James M. Murithi, Madeline R. Luth, Pavel Afanasyev, Paula C. A. da Fonseca, Elizabeth A. Winzeler, Caroline L. Ng, Matthew Bogyo, and David A. Fidock. 2019. "Covalent Plasmodium Falciparum-Selective Proteasome Inhibitors Exhibit a Low Propensity for Generating Resistance in Vitro and Synergize with Multiple Antimalarial Agents." PLoS Pathogens 15(6):e1007722–e1007722.

[118] Breinich, Manuela S., David J. P. Ferguson, Bernardo J. Foth, Giel G. van Dooren, Maryse Lebrun, Doris V Quon, Boris Striepen, Peter J. Bradley, Friedrich Frischknecht, Vern B. Carruthers, and Markus Meissner. 2009. "A Dynamin Is Required for the Biogenesis of Secretory Organelles in Toxoplasma Gondii." *Current Biology* 19(4):277–86.

[119] Melatti, Carmen, Manuela Pieperhoff, Leandro Lemgruber, Ehmke Pohl, Lilach Sheiner, and Markus Meissner. 2019. "A Unique Dynamin-Related Protein Is Essential for Mitochondrial Fission in Toxoplasma Gondii." PLOS Pathogens 15(4):e1007512.

[120] Wang, Lihui, Claire Delahunty, Karin Fritz-Wolf, Stefan Rahlfs, Judith Helena Prieto, John R. Yates, and Katja Becker. 2015. "Characterization of the 26S Proteasome Network in Plasmodium Falciparum." Scientific Reports 5(1):17818.

[121] Aminake, Makoah Nigel, Hans-Dieter Arndt, and Gabriele Pradel. 2012. "The Proteasome of Malaria Parasites: A Multi-Stage Drug Target for Chemotherapeutic Intervention?" International Journal for Parasitology: Drugs and Drug Resistance 2:1–10.

[122] Wu, Yang, Vesela Encheva, Judith L. Green, Edwin Lasonder, Adchara Prommaban, Simone Kunzelmann, Evangelos Christodoulou, Munira Grainger, Ngoc Truongvan, Sebastian Bothe, Vikram Sharma, Wei Song, Irene Pinzuti, Chairat Uthaipibull, Somdet Srichairatanakool, Veronique Barault, Gordon Langsley, Hermann Schindelin, Benjamin Stieglitz, Ambrosius P. Snijders, and Anthony A. Holder. 2019. "Protein Ubiquitylation Is Essential for the Schizont to Merozoite Transition in Plasmodium Falciparum; Blood-Stage Development." BioRxiv 755439.

[123] Cai, Hong, Timothy G. Lilburn, Changjin Hong, Jianying Gu, Rui Kuang, and Yufeng Wang. 2015. "Predicting and Exploring Network Components Involved in Pathogenesis in the Malaria Parasite via Novel Subnetwork Alignments." BMC Systems Biology 9(4):S1.

[124] Burgmann, H., S. Looareesuwan, C. Viravan, S. Vanijanonta, K. Zedwitz-Liebenstein, H. Vorbach, and W. Graninger. 1996. "Serum Laminin and Basic Fibroblast Growth Factor Concentrations in Patients with ComplicatedPlasmodium Falciparum Malaria." Journal of Clinical Immunology 16(5):278–82.

[125] Wenisch, C., W. Graninger, C. Viravan, S. Looareesuwan, B. Parschalk, and W. Wernsdorfer. 1994. "Serum Laminin in Malaria." Journal of Clinical Pathology 47(9):787 LP – 789.

[126] Mahamar, Almahamoudou, Oumar Attaher, Bruce Swihart, Amadou Barry, Bacary S. Diarra, Moussa B. Kanoute, Kadidia B. Cisse, Adama B. Dembele, Sekouba Keita, Benoît Gamain, Santara Gaoussou, Djibrilla Issiaka, Alassane Dicko, Patrick E. Duffy, and Michal Fried. 2017. "Host Factors That Modify Plasmodium Falciparum Adhesion to Endothelial Receptors." Scientific Reports 7(1):13872.

[127] Gnädig, Nina F., Barbara H. Stokes, Rachel L. Edwards, Gavreel F. Kalantarov, Kim C. Heimsch, Michal Kuderjavy, Audrey Crane, Marcus C. S. Lee, Judith Straimer, Katja Becker, Ilya N. Trakht, Audrey R. Odom John, Sachel Mok, and David A. Fidock. 2020. "Insights into the Intracellular Localization, Protein Associations and Artemisinin Resistance Properties of Plasmodium Falciparum K13." PLOS Pathogens 16(4):e1008482.

[128] Birnbaum, Jakob, Sarah Scharf, Sabine Schmidt, Ernst Jonscher, Wieteke Anna Maria Hoeijmakers, Sven Flemming, Christa Geeke Toenhake, Marius Schmitt, Ricarda Sabitzki, Bärbel Bergmann, Ulrike Fröhlke, Paolo Mesén-Ramírez, Alexandra Blancke Soares, Hendrik Herrmann, Richárd Bártfai, and Tobias Spielmann. 2020. "A Kelch13-Defined Endocytosis Pathway Mediates Artemisinin Resistance in Malaria Parasites." Science 367(6473):51 LP – 59.

[129] Bridgford, Jessica L., Stanley C. Xie, Simon A. Cobbold, Charisse Flerida A. Pasaje, Susann Herrmann, Tuo Yang, David L. Gillett, Lawrence R. Dick, Stuart A. Ralph, Con Dogovski, Natalie J. Spillman, and Leann Tilley. 2018. "Artemisinin Kills Malaria Parasites by Damaging Proteins and Inhibiting the Proteasome." Nature Communications 9(1):3801.

[130] Tilley, Leann, Judith Straimer, Nina F. Gnädig, Stuart A. Ralph, and David A. Fidock. 2016. "Artemisinin Action and Resistance in Plasmodium Falciparum." Trends in Parasitology 32(9):682–96.

[131] Ng, Caroline L., David A. Fidock, and Matthew Bogyo. 2017. "Protein Degradation Systems as Antimalarial Therapeutic Targets." Trends in Parasitology 33(9):731–43.

[132] Dogovski, Con, Stanley C. Xie, Gaetan Burgio, Jess Bridgford, Sachel Mok, James M. McCaw, Kesinee Chotivanich, Shannon Kenny, Nina Gnädig, Judith Straimer, Zbynek Bozdech, David A. Fidock, Julie A. Simpson, Arjen M. Dondorp, Simon Foote, Nectarios Klonis, and Leann Tilley. 2015. "Targeting the Cell Stress Response of Plasmodium Falciparum to Overcome Artemisinin Resistance." PLOS Biology 13(4):e1002132.

[133] Liu, Jun, Laura Wetzel, Ying Zhang, Eiji Nagayasu, Stephanie Ems-McClung, Laurence Florens, and Ke Hu. 2013. "Novel Thioredoxin-Like Proteins Are Components of a Protein Complex Coating the Cortical Microtubules of <Span Class="Named-Content Genus-Species" Id="Named-Content-1">Toxoplasma Gondii</Span>" Eukaryotic Cell 12(12):1588 LP – 1599.

[134] Liu, Jun, Yudou He, Imaan Benmerzouga, William J. Sullivan Jr, Naomi S. Morrissette, John M. Murray, and Ke Hu. 2016. "An Ensemble of Specifically Targeted Proteins Stabilizes Cortical Microtubules in the Human Parasite Toxoplasma Gondii." Molecular Biology of the Cell 27(3):549–71.

[135] Chen, Changchun, Eisuke Itakura, Katherine P. Weber, Ramanujan S. Hegde, and Mario de Bono. 2014. "An ER Complex of ODR-4 and ODR-8/Ufm1 Specific Protease 2 Promotes GPCR Maturation by a Ufm1-Independent Mechanism." PLOS Genetics 10(3):e1004082.

[136] Dwyer, Noelle D., Emily R. Troemel, Piali Sengupta, and Cornelia I. Bargmann. 1998. "Odorant Receptor Localization to Olfactory Cilia Is Mediated by ODR-4, a Novel Membrane-Associated Protein." Cell 93(3):455–66.

[137] Dellibovi-Ragheb, Teegan A., Hugo Jhun, Christopher D. Goodman, Maroya S. Walters, Daniel R. T. Ragheb, Krista A. Matthews, Krithika Rajaram, Satish Mishra, Geoffrey I. McFadden, Photini Sinnis, and Sean T. Prigge. 2018. "Host Biotin Is Required for Liver Stage Development in Malaria Parasites." Proceedings of the National Academy of Sciences 115(11):E2604 LP-E2613.

[138] Mustière, Romain, Patrice Vanelle, and Nicolas Primas. 2020. "Plasmodial Kinase Inhibitors Targeting Malaria: Recent Developments." Molecules (Basel, Switzerland) 25(24):5949.

[138] Sarafoglou, K., K. Bentler, A. Gaviglio, K. Redlinger-Grosse, C. Anderson, M. McCann, B. Bloom, D. Babovic-Vuksanovic, D.
920   Gavrilov, and S. A. Berry. 2009. "High Incidence of Profound Biotinidase Deficiency Detected in Newborn Screening Blood Spots in the
Somalian Population in Minnesota." *Journal of Inherited Metabolic Disease* 32(S1):169–73.

Table 1. Potentially available host-parasite systems, number of studies and runs in different tissues after database querying. Numbers are arranged as the number of studies / number of runs.

	Blood	Spleen	Liver	Brain	Bone marrow	Lungs
Human - *P. falciparum*	14 / 855	0 / 0	0 / 0	0 / 0	0 / 0	0 / 0
Human - *P. vivax*	6 / 141	0 / 0	1 / 4	0 / 0	0 / 0	0 / 0
Human - *P. berghei*	0 / 0	0 / 0	6 / 145	0 / 0	0 / 0	0 / 0
Mouse - *P. yoelii*	3 / 19	2 / 10	3 / 20	0 / 0	1 / 4	1 / 6
Mouse - *P. chabaudi*	4 / 204	5 / 795	2 / 46	0 / 0	0 / 0	0 / 0
Mouse - *P. berghei*	3 / 44	0 / 0	0 / 0	3 / 83	0 / 0	0 / 0
Monkey - *P. cynomolgi*	5 / 811	0 / 0	2 / 59	0 / 0	0 / 0	0 / 0
Monkey - *P. coatneyi*	1 / 35	0 / 0	0 / 0	0 / 0	2 / 66	0 / 0

Table 2. Number of blood studies for each host-parasite pair and suitability analysis of their experimental replicates (runs). [* = after applying a quality threshold of a unique read mapping
930   proportion of 70%, yellow: intermediate threshold, green: stringent threshold]

Host-parasite	# studies*	# runs*	Threshold: % transcriptome expressed	# studies with runs meeting threshold	# runs meeting threshold
Human – *P. falciparum*	13	621	30	8	180
			40	8	152
			50	6	131
			60	5	112
			70	4	101
Human – *P. vivax*	6	105	30	3	11
			40	3	10
			50	2	6
			60	2	5
			70	1	2
Monkey – *P. cynomolgi*	5	811	30	5	535
			40	5	500
			50	5	478
			60	5	370

			70	5	237
Monkey – *P. coatneyi*	1	34	30	1	21
			40	1	20
			50	1	18
			60	1	16
			70	1	13
Mouse – *P. berghei*	3	44	30	3	28
			40	3	23
			50	3	20
			60	1	13
			70	1	1
Mouse – *P. chabaudi*	4	91	30	4	31
			40	4	20
			50	3	8
			60	1	2
			70	0	0
Mouse – *P. yoelii*	3	14	30	1	5
			40	1	4
			50	1	1

Table 3. Overview of beta regression models to explain Relative Growth rate (RGR) and Mutagenesis Index Score (MIS) using eigenvector centrality measures of 5 networks: from 1) a single study on *P. berghei* infection in mice, from 2) overall, 3) core, 4) a single study on human *P.falciparum* infection in Indonesia (I) and 5) another singly study on human *P. falciparum* infection in Gambia (G).

	Dependent variable:									
	RGR					MIS				
	1	2	3	4	5	1	2	3	4	5
P. berghei EC	-1.445*** (0.089)					-0.754*** (0.087)				
Overall EC		-1.279*** (0.128)					-0.418*** (0.128)			
Core EC			-0.984*** (0.073)					-0.511*** (0.071)		

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
P. falciparum(I) EC				-0.833***					-0.952***	
				(0.267)					(0.272)	
P. falciparum(G) EC					-0.746***					-0.454***
					(0.070)					(0.069)
Constant	1.092***	1.505***	0.880***	0.486***	0.767***	1.146***	1.159***	1.032***	0.836***	1.001***
	(0.048)	(0.107)	(0.042)	(0.030)	(0.041)	(0.049)	(0.108)	(0.043)	(0.032)	(0.042)
Observations	2,169	2,169	2,169	2,169	2,169	2,169	2,169	2,169	2,169	2,169
R2	0.083	0.038	0.053	0.003	0.033	0.03	0.004	0.018	0.004	0.016
Log Likelihood	2,847.969	2,766.102	2,789.414	2,718.350	2,718.52	6,065.9	6,032.86	6,050.41	6,033.30	6,019.49

Note: Values are effect sizes with error in brackets; *p<0.1; **p<0.05; ***p<0.01

Figure 1. Proportion and number of sequencing reads and expressed genes from parasite and host in selected malaria RNA-Seq studies. We mapped sequencing reads from studies selected for their potential to provide both host and parasite gene expression data (total studies=63, total runs=3351) against

appropriate host and parasite genomes. (A) The percentage of parasite reads (x-axis) is plotted for runs in each study (host and parasite add to 100%). The studies are categorised according to the host organisms studied and "enriched/depleted" to indicate enrichment of infected hepatocytes or depletion of leukocytes from blood. Studies labeled "dual" were originally intended to simultaneously assess host and parasite transcriptomes. We also plot the number of reads mapped against the number of expressed genes for host (B) and parasite (C). The proportion of transcriptome detected as expressed increases with sequencing depth towards the maximum of all expressed in the transcriptome.

Figure 2. Overlapping bipartite edges between datasets across host-parasite systems

Based on the proportion of transcriptome expressed in each study, three thresholds were implemented on the selection of runs for co-expression analysis. Without thresholds "all", includes all runs. The intermediate ("int") and stringent ("str")

thresholds, include runs in which 50% ("int") and 70% ("str") of the transcriptome of both host and the parasite are detected as expressed (covered), respectively. (A) Schematic to show how samples were selected for coexpression analysis. (B) For each dataset, the median host and parasite transcriptome coverage is indicated. The heatmap shows the log10 transformed number of bipartite edges for each study on the diagonal. The set size of the overlapping bipartite edges between corresponding datasets are displayed in the remaining fields. The datasets are clustered based on euclidean distances between these set sizes. The size of the overlapping set of bipartite edges is not determined by the median parasite or host transcriptome coverage, even studies with low coverage of the parasite transcriptome provide datasets leading to high number of bipartite edges found in common with other datasets. The optimal overlap set size does not suggest a single threshold for the selection of sequencing runs - all three thresholds are found to have a high number of overlapping edges with other datasets. (C) Jaccard indices (size of intersection as a ratio of the size of union) for each dataset pair are displayed as (-)log10 transformed indices. The versions of the datasets used for further analysis are marked with an asterix. (D) The size of the intersection was tested for significance using Fisher's exact test. The heatmap shows that all datasets led to correlation networks that overlapped with other networks from other datasets more than expected by chance. We can conclude that they contain biological signals suitable for combined analysis.

Figure 3. Alternative and interlinked strategies to reconstruct host-parasite interaction networks.

We designed two interlinked approaches to obtain a consensus network involving multiple host-parasite systems. (A) An implementation of permutation tests derives "empirical p-values" to test correlation coefficients for significance and infers interactions. (B) The methodology used to compare transcriptome coverage thresholds (50%, intermediate, "int"; 70%, stringent, "str") is using different subsets of runs from each individual study ("sub-datasets" S1, S2 and S3 in this illustration). As in fig 2, without thresholds "all", includes all runs. Each subset is analysed separately to infer interactions,

 allowing the subset (and threshold) leading to highest information overlap with other studies to be selected. (C) shows the inference of common interactions across host-parasite systems using the knowledge of single-copy orthologs across the hosts and the parasites. Here, we can take two approaches: the first approach uses intersections in a multilayer network approach to come to a consensus of common interactions. In the second approach, we concatenate runs and correlate gene expression on those as a single "overall" dataset. In the analysis for our final results we combined these approaches mapping networks from each study onto the common network from an "overall" dataset.

Figure 4. Multi-layer networks of host-parasite interactions across host-parasite systems increase resolution over neworks from single studies

We performed pairwise gene correlation tests on individual studies for gene expression in malaria and on an "overall" dataset constructed by concatenating data from 15 datasets. (A) A small part of the "overall" network shows those overlaps. We selected this part of the network starting from a set of "core edges" found in most networks (the "overall" network and networks from six individual studies). Edges in the neighbourhood to this were randomly selected in equal numbers, where available, based on how many networks, out of 15, they were found in. The number of networks an edge is found in is represented by the edge colour. Several edges were found common to multiple datasets across host-parasite systems, a result later used to derive a "core" network. (B) The (log 10 transformed) number of edges in networks from individual studies (horizontal bars) and shared (vertical bars) among networks shows the large overlaps between individual studies but also large differences in the sizes for those overlaps. (C) shows the effect of performing an increasing number of permutations on the p-value of an edge within a single network (datasets ERP004598_all,

SRP118827_int, DRP000987_str and "overall" as examples). As the number of permutations increases, the increase in
resolution (decrease in the number of significantly correlated edges recovered) slows, warranting a different approach to
increase resolution, such as the multi-layer analysis depicted in previous panels. (D) shows the distribution of p-values
across the range of correlation coefficients. The same studies as in fig (C) are used. While individual datasets might
represent different aspects of malaria infection and have a reduced number of interactions in common, we here show the
benefit of analysing the "overall" dataset: the "overall" analysis identifies both positive and negative correlations. In the
results of the "overall" dataset, we reach a consensus of all the datasets included.

Figure 5. Explanation of gene essentiality with gene co-expression networks

Eigenvector centrality in a "core" network of correlated parasite-parasite gene expression is plotted against Relative
Growth Rate (RGR, in A) and Mutagenesis Index Score (MIS, in B), respectively. Lines depict predictions of the
essentiality scores from the centrality measure in a beta-regression: the higher the centrality, the lower is the essentiality
score (RGR and MIS), meaning that genes central in the network are more essential to the parasites growth and survival.
Color for both figures indicates phenotypes as categorised in [35].

(A)

DRP000987_str

Response to toxic substance
Positive regulation of humoral immune response
Positive regulation of acute inflammatory response
Cellular response to hypoxia
Cellular response to oxidative stress
Immunoglobulin secretion
Positive regulation of IL6 secretion
Macrophage differentiation
Positive regulation of nitric oxide biosynthesis

SRP116793_all

G protein -coupled receptor signalling pathway
Regulation of MAPK cascade
Nitric oxide mediated signal transduction
Positive regulation of cAMP-mediated signalling
Immature B cell differentiation
Calcium ion transport into cytosol

Response to oxygen levels
Cellular response to liposaccharide
Response to wounding
T-cell lineage commitment
Cell-cell adhesion mediated by cadherin

Response to oxygen levels
Calcium-depndent cell-cell adhesion
Cellular response to decreased oxygen levels
T cell lineage commitment

ERP106451_int

Ammonium transport
Positive regulation of B cell differentiation
Response to drug
Positive regulation of G-protein coupled receptors
Regulation of IL2 biosynthesis
Erythrocyte development
Cellular response to oxidative stress
Regulaion of cAMP-mediated signalling pathway
Response to starvation

SRP118827_int

Response to oxygen levels
Calcium-dependent cell-cell adhesion
Lymphocyte chemotaxis
Positive regulation of activated T cell proliferation
Negative regulation of alpha-beta T cell differentiation
Response to hydrogen peroxide

G protein-coupled receptor signalling pathway
Cell adhesion
Regulation of cGMP-mediated signaling
Nitric oxide mediated signal transduction
Antimicrobial humoral immune response
Response to toxc substance
Xenobiotic metabolic process
Activation of MAPK activity
Hydrogen peroxide biosynthesis process
Positive regulation of IL6 production
Neutrophil chemotaxis

Cell-cell adhesion mediated by cadherin
Regulation of reactive oxygen species
Regulation of macrophage chemotaxis
Fat cell differentiation
Response to hydrogen peroxide

Hematopoietic stem cell differentiation
Positive regulation of IFNg
B cell apoptotic process
T cell lineage commitment
Pre-B cell differentiation
Fat cell differentiation
Negative regulation of macrophage chemotaxis
Positive regulation of chemokine
Killing by host of symbiont cells
Drug transport

Response to bacterium
Response to oxygen levels
Regulation of IFNg biosynthesis
B cell differentiation
Cytokine production
Regulatory T cell differentiation
Regulation of isotype switching to IgG
Cell-cell adhesion mediated by cadherin

Response to oxygen levels
Cellular response to decreased oxygen levels
Cell-cell adhesion by cadherin
Negative regulation of alpha-beta T cell
Cellular response to oxygen levels

G protein-coupled receptor signaling pathway
Positive regulation of cAMP-mediated signaling
Immune response
Positive regulation of G protein-coupled receptor
Regulation of MAPK cascade
Response to hydrogen peroxide
Response to drug
Cell-cell adhesion by integrin
Cellular response to xenobiotic stimulus

G protein-coupled receptor signalling pathway
Positive regulation of cAMP-mediated signalling
Transmembrane transport
Modification by host of symbiont morphology
Nitric oxide mediated signal transduction
NK T cell activation
IL13 production
cGMP metabolic process
Phagocytosis
Cellular response to xenobiotic stimulus

ERP004598_all   **ERP110375_int**

(B)

DRP000987_str

Cell gliding
Entry into host cell
Mitochondrial electron transport
Late endosome to vacuole transport

Entry into host
Pathogenesis
Cell-cell adhesion
Cellular calcium ion homeostasis

SRP116793_all

Entry into host cell
Late endosome to vacuole transport
Transmembrane transport

Cellular calcium ion homeostasis
Cell-cell adhesion
Selenocysteine incorporation
Pathogenesis
De novo UMP biosynthesis

ERP106451_int

Outer dynein arm assembly
Carbohydrate transport
Late endosome to vacuole transport

Selenocysteine incorporation
De novo UMP biosynthesis
Cellular calcium ion homeostasis

Activation of MAPK activity

Cellular oxidant detoxification

Cellular response to drug

Ribosome disassemlby

Ribosomal large subunit biogenesis

Cellular calcium ion homeostasis
Selenocysteine incorporation
Pathogenesis
Cell-cell adhesion
Do novo UMP biosynthesis

SRP118827_int

Entry into host cell
Late endosome to vacuole transport
Cellular response to drug
Endonucleolytic cleavage in ITS1
Transmembrane transport

Celular calcium ion homeostasis
Selenocysteine incorporation
Pathogenesis
De novo UMP biosynthesis

Late endosome to vacuole transport
Carbohydrate transport

Cellular calcium ion homepstasis
Cell-cell adhesion
Pathogenesis
Entry into host
De novo UMP biosynthesis

Entry into host cell
Late endosome to vacuole transport

ERP004598_all   **ERP110375_int**

1015

Figure 6. Biological Processes (BP) Gene Ontology (GO) terms shared between datasets across host-parasite systems.

We performed GO term enrichment analysis for the "overall" dataset and its 15 constituent datasets for (A) host and (B) parasite genes. Here we show GO terms for the 6 largest studies for each host species. Enriched GO terms (p-value >= 0.05) are presented in sets coloured in red, green and blue for human, mouse and macaque studies. The GO terms of the "overall" dataset are displayed in the central yellow set. GO terms for the six individual datasets with the highest contribution to the "overall" dataset are shown in the perimeter of this. Overlapping between the respective areas shows shared GO terms found in either dataset. Set overlaps are not illustrated between the six individual datasets but respective terms are underlined (shared between 4-5 studies) or bold (shared between all 6 studies).

Figure 7. An interaction network of host-parasite GO terms.

Host and parasite genes in the "overall" and "core" gene expression correlation networks were analysed for GO term enrichment. For each individual enriched host GO term interacting genes from the other species were then analysed for enrichment more specifically. This gives and understanding of what GO biological processes in one organism, say, the host, are associated with specific processes in the parasite. In this figure, we show all parasite GO terms associated with host GO terms related to "adhesion", in the "core" network.

Figure 8. A "core" network highlights parasite interaction with specific immune cells

We derived a "core" network containing interacting genes supported by correlated gene expression in at least one human and one model organism study, A) this complete "core" network is displayed with grey nodes for host and red nodes for parasite genes. The size of a node is based on its connectivity - the higher its degree, the larger is the node. (B) illustrates the identification of node clusters in the "core" network using an edge betweenness algorithm from igraph. Clusters with immune cells specific marker genes are coloured (see legend, including a count for the number of marker genes in brackets). The remaining clusters are left grey.