1

PINK1-parkin-mediated mitophagy generates somatic mosaicism of the

mitochondrial genome

Arnaud Ahier[1], Nadia Cummins[1], Chuan-Yang Dai[1], Jürgen Götz[1] and Steven Zuryn[1]*.

[1] *Clem Jones Centre for Ageing Dementia Research, Queensland Brain Institute, The University*

of Queensland, Brisbane, Australia, 4072.

Corresponding author.

21 Running title: PINK-1-parkin determines mtDNA mosaicism

23 Keywords: Mitochondria, mtDNA, mitophagy, PINK1, parkin, heteroplasmy, proteotoxicity

Abstract

The composition of the mitochondrial DNA (mtDNA) in each cell is in constant flux through processes of mutation, replication, and degradation. In multiple species, certain tissues consistently harbour greater loads of mtDNA mutations relative to others. However, the mechanisms that drive this stereotyped distribution of mtDNA mutations within individuals are unknown. Using *C. elegans* as a model system, we found that mitochondrial autophagy (mitophagy), mediated through the conserved PTEN-induced putative kinase (PINK1) and the E3 ubiquitin-protein ligase parkin, selectively removes mtDNA mutations from specific tissue types. Moreover, neurons in particular were prone to accumulating mtDNA mutations when mitophagy was perturbed genetically, or through the ectopic expression of proteotoxic species that are associated with neurodegenerative diseases. These results suggest that mitophagy deterministically drives differences in heteroplasmy between tissues, and points at the existence of a causal link between proteotoxicity, mitophagy, and mtDNA mutation levels in neurons.

Introduction

Mitochondria act as central hubs for cellular bioenergetics, macromolecule precursor synthesis, redox balance, Ca^{2+} handling, apoptosis, and immunity [1, 2]. These organelles house their own genome (mtDNA), as well as RNA and protein-synthesizing systems, which together code and coordinate the assembly of core subunits of oxidative phosphorylation (OXPHOS), a process that generates adenosine triphosphate, the primary energy substrate of the cell. Although mtDNA lesions that perturb the assembly of OXPHOS enzymes can cause devastating metabolic disorders [3], mutations are buffered by the polyploidy nature of the mitochondrial genome; tens to hundreds of mitochondria can populate each cell, with each

organelle containing multiple copies of mtDNA. As a result, a mixture of multiple mtDNA

variants can co-exist within individual cells [4]. This state is termed heteroplasmy, and the

factors that influence its dynamics are critical determinants of the pathogenesis of

mitochondrial diseases, and possibly a number of age-related conditions, including

neurodegeneration, cardiomyopathy, diabetes, and cancer [5-7].

At any given time, an interplay between stochastic and deterministic processes influences

cellular heteroplasmy [5]. Mitotic segregation, genetic drift, and the competing effects of

homeostatic [8-12] and retrograde signalling pathways, such as the mitochondrial unfolded

protein response [9, 10], may contribute to the divergence of heteroplasmy levels between

cells, generating heteroplasmy mosaicism. However, there is evidence that specific organs

and tissues are prone to accumulating certain mtDNA mutations, leading to a stereotyped

pattern of heteroplasmy mosaicism across individuals [13-17]. For example, the caudate,

putamen and substantia nigra regions of the human brain show an age-dependent increase

in the heteroplasmy levels of the mtDNA^{4977} deletion (a mutant genome harbouring a 4,977bp

deletion), relative to the rest of the brain [15]. Stereotyped heteroplasmy differences

between tissues have also been observed in *C. elegans*. Here, mtDNA molecules harbouring

a large deletion accumulate at different heteroplasmy levels in distinct somatic and germ

lineages over time [16]. However, how these non-random patterns of differences in mtDNA

heteroplasmy evolve between tissues is still unknown.

Here, we provide evidence that mitochondrial autophagy (mitophagy) – orchestrated through

the PINK1/parkin signalling pathway – selectively removes mutant copies of mtDNA in a

subset of somatic tissue-types in *C. elegans*, resulting in the formation of mosaic and

stereotyped heteroplasmy differences between tissues of individuals. Whereas mitochondria

within the nervous system, epidermis and intestine use PINK1 and parkin to maintain low

levels of a mitochondrial genome harbouring a large deletion (*uaDf5,* also called ΔmtDNA),

mitochondria in the body wall muscles do not. Instead PINK1 and parkin act unselectively

against ΔmtDNA in muscles to control total mitochondrial volume. As a result, removing the

PINK1/parkin axis equalises heteroplasmy across all tissue types, abolishing mosaicism

between organs. Moreover, we found that neurons were most dependent upon PINK1 and

parkin for maintaining low heteroplasmy levels of mtDNA, and that proteotoxic threats

associated with neurodegenerative disease and mitophagy inhibition enhanced ΔmtDNA

heteroplasmy levels in neurons. Together, these results suggest that PINK1-parkin-mediated

mitophagy acts deterministically to drive the formation of a mosaic and stereotyped

heteroplasmic landscape in individual organisms, and that genetic or proteotoxic

interruptions to this process can increase mtDNA mutation levels in the nervous system.

Results

In *C. elegans*, somatic tissues display stereotyped heteroplasmy, whereby body wall muscles,

epidermis, neurons, and intestine consistently harbour decreasing heteroplasmy levels of

mtDNA mutations relative to each other [16]. To determine how this distinct pattern of

heteroplasmy is formed, we assessed whether each tissue responded differently to the same

mtDNA mutation, a 3.1 kb deletion (*uaDf5*, also called ΔmtDNA; Figure S1A) [18]. The *uaDf5*

deletion exists in heteroplasmy with wild-type copies of the mtDNA at a level of

approximately 50-60% across populations of animals (Figure S1B) [9, 18]. This deletion

disrupts four genes that encode core subunits of the electron transport chain (Figure S1A). At

50-60% heteroplasmy, animals display reductions in basal oxygen consumption as well as

total respiratory capacity [9], suggesting that they experience mitochondrial dysfunction. In

order to determine how ΔmtDNA altered mitochondrial morphology in each tissue, we

introduced the ΔmtDNA genome into animals expressing a fluorescent fusion protein

(*Mito::mKate2*) that localizes to mitochondria and is controlled by either a pan-neuronal

promoter (*rgef-1p*), an intestinal-specific promoter (*ges-1p*), an epidermis-specific promoter

(*dpy-7p*), or a body wall muscle-specific promoter (*myo-3p*). While gross morphology

remained unaltered, three-dimensional reconstruction of fluorescence images acquired from

each strain revealed that the mitochondrial volume changed in response to the presence of

ΔmtDNA in a tissue-dependent manner (Figure 1A-H and Figure S2). The volume of the

mitochondrial network was highly increased in the epidermis, only mildly increased in

neurons and intestine (Figure 1A-C and 1E-G), and unchanged in body wall muscles (Figure 1D

and H). This revealed that when threatened with a common mtDNA defect, distinct tissue

types responded uniquely and independently of each other, hinting at the possibility that

tissue-specific differences in mitochondrial physiology could generate disparity in ΔmtDNA

heteroplasmy.

Because heteroplasmy is directly determined by the net effect of the propagation and

degradation of mutations, we next investigated how mitochondria within each tissue type

altered mtDNA copy numbers in response to the presence of ΔmtDNA. To this end, we

performed cell-type-specific mitochondrial affinity purification (Figure S3) to separate and

isolate intact mitochondria from each tissue type of animals that harboured either wild-type

or both mutant and wild-type genomes. Mitochondria from each of the four tissues were

purified from transgenic strains carrying *TOMM-20::mKate2::HA* in which the outer surface

of the mitochondria was labelled with a hemagglutinin (HA) epitope in a tissue-specific

manner. Using anti-HA antibody-coated beads, mitochondria from each tissue type were

affinity-purified from homogenates and the number of individual mitochondria within each

sample was quantified before their DNA extracted to determine the copy number of genomes

per organelle using a fluorescent probe-based qPCR assay. We found that as a result of the

presence of ΔmtDNA the mtDNA copy number per mitochondrion increased in all tissues

except epidermis, with the largest absolute increase in copy number per organelle observed

in mitochondria isolated from body wall muscles (Figure 2A-D). This suggests that muscle

cells increase mtDNA replication in the presence of ΔmtDNA more than any other somatic

tissue investigated here. Because the ΔmtDNA genome is smaller in size than its wild-type

counterpart, it has an inherent replicative advantage [17, 19]. As such, any enhanced rates of

mtDNA replication in muscle cells may disproportionately increase heteroplasmy over time

and contribute to greater heteroplasmy loads.

The heightened mtDNA copy numbers in body wall muscles may reflect the comparatively

demanding energetic function that this tissue performs. Body wall muscles include four

quadrants of somatic muscle bundles, each of which consists of 23-24 cells arranged into

staggered pairs that stretch longitudinally from the head to the tail of the animal [20]. The

coordinated contractile activities of these bundles produce powerful sinusoidal body bends

that drive locomotion. Increasing the total mtDNA copy number in body wall muscles may

help to functionally compensate for ΔmtDNA, in a similar fashion as recently shown in mouse

sperm harbouring high levels of mtDNA mutations [21]. Indeed, under strenuous conditions

(performing the thrashing assay, see Materials and Methods), muscle function was

unperturbed in the presence of ΔmtDNA, even when we forced heteroplasmy to very high

loads (93.39 ± 0.34%) by genetically bottlenecking and selecting individual animals with high

ΔmtDNA levels across several generations (Figure 2E-H and Figure S4). Furthermore, body

wall muscle function was still unperturbed in animals with high ΔmtDNA heteroplasmy at

older ages (Figure 2G and H and Figure S4). Therefore, increasing mtDNA copy number may

functionally compensate for ΔmtDNA, particularly in energetically demanding cells such as

muscle, but it could also act to promote the propagation of mutant genomes with a replicative

advantage leading to the development of tissue disparity in heteroplasmy over time.

In addition to DNA replication, ΔmtDNA heteroplasmy can be modified through selective

degradation processes. Mitophagy contributes to mitochondrial turnover by selectively

destroying depolarized mitochondria harbouring mtDNA mutations [9, 10, 18], and it can be

triggered by a diverse array of cellular events, including cellular differentiation, and in

response to physiological cues, such as cellular activity and nutrient levels [22]. As such,

mitophagy may operate under distinct parameters in different cell types, which, in the

presence of ΔmtDNA mutations, could contribute to determining the heteroplasmy levels of

specific tissues and organs.

To test this idea, we introduced the ΔmtDNA mutation into double-mutant strains with null

deletions in the genes *pink-1* (PINK1) and *pdr-1* (parkin)[23], key mediators of mitophagy in

multiple species including *C. elegans* [24-27]. PINK1 normally induces mitophagy by

accumulating on the outer mitochondrial membrane of defective organelles, where it recruits

the E3 ubiquitin ligase parkin [28]. Parkin decorates the surface of these mitochondria with

polyubiquitin chains, thereby inducing engulfment by autophagosomes and destruction

following fusion with lysosomes [29]. The *pink-1(tm1779);pdr-1(gk448)* mitophagy-deficient

mutant animals harbouring ΔmtDNA were crossed to transgenic animals carrying the *TOMM-*

20::mKate2::HA transgene expressed under the individual tissue-specific promoters (muscle,

intestine, epidermis and neurons), and intact mitochondria were isolated from each tissue

using affinity purification (Figure 3A) [16]. To eliminate the possibility of any population

variability affecting our analyses, we compared mitochondria purified from each tissue to the

total mitochondria (homogenate) from the same samples. In addition, we performed the

same experiments in animals in which mitochondria from all cell types were purified using a

strain expressing *TOMM-20::mKate2::HA* under the ubiquitous promoter *eft-3p*. As expected,

ΔmtDNA heteroplasmy levels were not different between total mitochondria (homogenate)

and mitochondria affinity purified from all cell types (ubiquitous) in both wild-type and *pink-*

*1;pdr-1*backgrounds (Figure 3B), confirming that deficiencies in PINK-1 and PDR-1 did not

influence the mitochondrial isolation procedure. Deleting *pink-1* and *pdr-1*, however, resulted

in a significant increase in ΔmtDNA heteroplasmy levels in mitochondria purified from

neurons, intestine, and epidermis (Figure 3C-E), suggesting that PINK-1 and PDR-1 operate

within these tissues to reduce ΔmtDNA heteroplasmy. Neither *pink-1* nor *pdr-1* appeared to

affect mtDNA replication, as mtDNA copy number was unchanged in *pink-1;pdr-1* double

mutant backgrounds (Figure 3H). Instead, and as previously demonstrated [8-10], one

plausible explanation for our finding is that mitophagy preferentially targets and destroys

mitochondria that harbour the highest loads of ΔmtDNA. Interestingly, this activity was not

conserved in the body wall muscles. Removal of *pink-1* and *pdr-1* had no effect on ΔmtDNA

heteroplasmy levels in this tissue (Figure 3F), suggesting that mitophagy has distinct roles,

including in mtDNA quality control, in the different tissue types. The lack of mitophagy-driven

selectivity against ΔmtDNA in muscle cells may lead to the formation of heteroplasmy

disparity between this tissue and other somatic tissues. In support of this notion, differences

in ΔmtDNA heteroplasmy between muscle and other tissues was abolished when *pink-1* and

pdr-1 were deleted (Figure 3G, G', and G'').

It is possible that PINK-1-PDR-1-mediated mitophagy is inactive in body wall muscle cells, and

that deleting the *pink-1* and *pdr-*1 would therefore be inconsequential to ΔmtDNA

heteroplasmy in this tissue. However, we found that a PINK-1::GFP reporter driven by its

native promoter [30] was expressed in the muscle cells, in addition to the other cell types

studied here (Figure 4A-D). A translational GFP::PDR-1 reporter is also expressed in body wall

muscles in *C. elegans*, suggesting that PDR-1 is normally present in this tissue [23]. Moreover,

removal of *pink-1* and *pdr-1* dramatically increased mitochondrial volume in the muscle cells,

suggesting that mitophagy is used to constrain the mitochondrial network size in this cell type

(Figure S5). Indeed, previous studies have found that RNA inactivation (RNAi) of *pink-1* also

increased mitochondrial volume in the body wall muscle cells of *C. elegans* [25]. This is in

contrast to neurons, intestine, and epidermis, in which *pink-1* and *pdr-1* removal had no effect

on mitochondrial volume (Figure S5). Taken together, these results suggest that PINK-1-PDR-

1-mediated mitophagy performs different roles in each tissue type, which leads to the

development of distinct and stereotyped heteroplasmy levels in different organs.

Our results indicate that the activity of PINK-1-PDR-1-mediated mitophagy can determine

ΔmtDNA heteroplasmy in neurons, intestinal cells, and epidermal cells. We recently

demonstrated that proteotoxic tau, a protein that forms aggregates in tauopathies such as

Alzheimer's disease [31], perturbs parkin translocation to the mitochondria, and that human

tau expressed in the *C. elegans* nervous system inhibits mitophagy [24]. Because inactivating

pink-1 and *pdr-1* increased ΔmtDNA heteroplasmy levels in neurons more than in any other

investigated tissue type, we next determined whether the expression of tau could also

increase heteroplasmy levels in the nervous system. We constructed strains harbouring

ΔmtDNA and expressing human tau (hTau) in the nervous system (*aex-3p::hTau*) [24] as well

as the neuron-specific TOMM-20::mKate2::HA tag. We then performed affinity purification of

neuronal mitochondria in these strains and found that hTau significantly increased

heteroplasmy (Fig. 4E), suggesting that the expression of mitophagy-inhibiting proteins can

increase ΔmtDNA levels in neurons. Other proteotoxic species have also recently been found

to inhibit mitophagy in neurons. The expanded polyglutamine (polyQ) tract in mutant forms

of the Huntingtin protein (Htt) that cause Huntington disease were recently reported to

impede mitophagy in differentiated striatal neurons [32]. Indeed, we found that the

expression of a polyglutamine (polyQ) tract (*rgef-1p::Q40::yfp*) [33] in the *C. elegans* nervous

recapitulated the effect of hTau and increased ΔmtDNA heteroplasmy in neurons (Fig. 4E).

These results suggest that diverse proteotoxic threats associated with a broad range of

neurodegenerative diseases may promote the accumulation of mtDNA deletions in the

nervous system.

Discussion

Polyploidy of the mitochondrial genome gives rise to intercellular and intertissue differences

in heteroplasmy, whereby adjacent cells or tissues within the same individual develop

different mtDNA mutation loads. Our results suggest that the tissue-specific activities of *pink-*

1 and *pdr-1*, conserved genes that mediate key steps of mitophagy in *C. elegans* [24-27], give

rise to stereotyped differences in heteroplasmy between major somatic tissues (Fig 4F).

Furthermore, we have demonstrated that proteotoxic threats that can perturb mitophagy in

neurons can increase heteroplasmy levels in the nervous system. Taken together, our results

suggest that individual cellular contexts, genetic perturbations, and disease-related

proteostatic stresses that can each modify the activity of mitophagy processes, lead to

changes in the distribution and abundance of mtDNA mutations throughout an organism.

Previously, others have shown that mtDNA mutations can alter mitochondrial morphology

and mtDNA copy number [18, 34]. Here we have assessed these changes with cell-type-

specific resolution and have found that the ΔmtDNA mutation invoked distinct alterations in

mitochondrial volume and mtDNA copy number in individual tissues. The fact that each tissue

type displays a unique response to an identical mtDNA perturbation suggests that specific

changes in mitochondrial physiology are activated under distinct cellular contexts. For

example, in the presence of ΔmtDNA, mitochondrial volume in the epidermis increased

significantly, but did not change in the body wall muscles. Oppositely, mtDNA copy number

per mitochondrion was unchanged in the epidermis, but increased the greatest in muscle

cells. These distinct responses may be derived from tissue function and hence energetic

requirements. Relative to the epidermal tissue, the energetically strenuous function of the

body wall muscle during locomotion may require compensatory increases in mtDNA

biogenesis to overcome deficiencies caused by the ΔmtDNA mutation. Indeed, others have

found the same to be true in energetically demanding cells. In mice harbouring high levels of

mtDNA mutations [35, 36], an increase in total mtDNA copy number in spermatocytes

rescued the fertility phenotype and normalized spermatocyte proteome changes, in spite of

unaltered mtDNA mutation load [21]. Moreover, increasing mtDNA copy number in mice

harbouring a heteroplasmic mtDNA mutation (m.5024C>T) ameliorated pathology in a

number of energetically demanding tissues, such as the heart [37]. Even when heteroplasmy

was forced to very high values through genetic bottlenecking, we found that muscle function

was unperturbed in ΔmtDNA animals, despite the mutational load being the greatest in the

body wall muscle cells. Therefore, certain cell types with energetically demanding functions,

such as muscle, may preferentially compensate for the effect of deleterious mtDNA mutations

by increasing total mtDNA copy number, rather than by clearing the mtDNA mutation through

processes such as mitophagy. An unintended consequence of such a strategy may be to

propagate those mtDNA mutations that have a replicative advantage, such as large deletions.

Although enhanced mtDNA propagation may help to contribute to heightened ΔmtDNA

heteroplasmy levels in body wall muscles, our results indicate that the removal of *pink-1* and

pdr-1 was sufficient to equalise ΔmtDNA heteroplasmy levels across all tissue types. This

suggests that the tissue-specific activities of PINK-1-PDR-1-mediated mitophagy alone can

drive the formation of heteroplasmy disparity between tissues. While in neurons, intestinal

cells, and epidermal cells, deleting *pink-1* and *pdr-1* resulted in an increase in ΔmtDNA

heteroplasmy, levels remained unchanged in muscle cells. This suggests that whereas PINK-

1-PDR-1-mediated mitophagy selects against the ΔmtDNA mutation in neurons, intestine, and

epidermis, it does not do so in body wall muscle. Instead, mitophagy acts to constrain the

volume of the mitochondrial network in the body wall muscles, rather than selectively

reducing the heteroplasmy levels of ΔmtDNA mutations. This indicates that PINK-1-PDR-1-

mediated mitophagy performs distinct roles in different cell types, which predisposes certain

tissues and organs to the accumulation of ΔmtDNA.

In humans, large mtDNA deletions have been shown to accumulate in specific regions of the

brain during ageing [15]. Although there is strong evidence that PINK1-parkin-mediated

mitophagy occurs in mammalian neurons [38, 39], it is not known whether differences in the

activity of mitophagy plays a role in specifying the heteroplasmy levels of mtDNA deletions

within distinct regions of the brain. In mice homozygous for a proofreading deficiency in DNA

polymerase γ, which induces a high level of mtDNA single nucleotide mutations [35, 36],

parkin knockout significantly increases the levels of pathogenic mtDNA mutations in the

striatum [40]. Moreover, mutations in Pink1 [41] and parkin [42] can lead to early onset

Parkinson's disease, and substantia nigra neurons in Parkinson's disease patients have high

levels of mtDNA deletions [43, 44]. Although we did not examine subsets of neurons in this

study, our results suggest that neurons in general are the most vulnerable to increases in

heteroplasmy when mitophagy is perturbed. Therefore, our work demonstrates that

mutations in Pink1 and parkin have a widely conserved effect across phyla of enhancing the

accumulation of mtDNA deletions in neurons, which may perturb their function and cause

disease.

In addition to Pink1 and parkin mutations, we found that toxic proteins associated with

neurodegenerative diseases other than Parkinson's disease also increase heteroplasmy.

Overexpression in neurons of hTau, as well as a polyQ tract, significantly increased ΔmtDNA

heteroplasmy levels in the *C. elegans* nervous system. To our knowledge, this represents the

first reported incidence of proteotoxicity increasing the cellular load of mtDNA mutations.

Interestingly, similar proteotoxic species have been recently reported to inhibit mitophagy in

neurons. Via its projection domain, hTau can associate with Parkin in the cytoplasm and

specifically impair its translocation to the surface of defective mitochondria in human

neuroblastoma cells [24]. In addition, a polyQ tract within expanded mutant versions of the

Huntingting protein can prevent the formation of the Beclin1-Vps34 initiation complex,

negatively affecting the efficiency of mitophagy in post-mitotic striatal neurons [32]. These

results demonstrate that perturbing mitophagy indirectly with proteotoxicity has the same

effect as perturbing mitophagy genetically through mutations in *pink-1* and *pdr-1*. They also

demonstrate that toxic proteins associated with diverse neurodegenerative diseases can

modify the genetic landscape of the mitochondrial genome in neurons and provide evidence

that two of the most common hallmarks of neurodegeneration - proteotoxicity and

mitochondrial dysfunction - may converge at the level of the mitochondrial genome.

Materials and Methods

C. elegans **strains and culture**. LB138 *him-8(e1489) IV; uaDf5/+*; EG8078 *oxTi185 I; unc-*

119(ed3) III; BR4006 *pink-1(tm1779) II; byEx655 [pink-1p::pink-1::GFP; myo-2p::mCherry;*

herring sperm DNA]; VC1024 *pdr-1(gk448) III*; AM101 *rmIs110 [rgef-1p::Q40::YFP]* and wild-

type (N2) were acquired from the *Caenorhabditis* Genetics Center (CGC), University of

Minnesota, USA, which is funded by the NIH Office of Research Infrastructure Programs (P40

OD010440). APD178 *apdIs11 [aex-3p::tau4RIN + coel::gfp]* were kindly provided by Dr H.

Nicholas (The University of Sydney, Sydney, Australia). *C. elegans* culture and maintenance

were performed according to standard protocols [45]. The transgenic strains used for cell-

specific mitochondrial purification experiments were generated as previously described [16].

Cell-specific mitochondrial affinity purification (CS-MAP). CS-MAP was performed as

previously described [16]. Heteroplasmy in mitochondria derived from different cell types

was investigated across very large populations of pooled animals (>10,000 individuals). This

approach enabled population-scale trends to be resolved by overcoming confounding

influences that can be introduced by the presence of stochastic inter-individual variations.

Briefly, 10,000 to 20,000 L4 staged worms were grown on 2 x 150 mm NGM plates seeded

with *E. coli* OP50 bacteria. Worms were homogenized into a hypotonic buffer [50 mM KCl, 110 mM mannitol, 70 mM sucrose, 0.1 mM EDTA (pH 8.0), 5 mM Tris-HCl (pH 7.4)] with a dounce homogenizer and the subsequent crude mitochondrial fraction was enriched by differential centrifugations. The mitochondria were then isolated from this fraction using anti-HA (influenza hemagglutinin) magnetic beads (Pierce). The beads were washed and resuspended in 20 µl of hypotonic buffer. All CS-MAP experiments were repeated in at least three independent experiments and data for each genetic background were obtained at similar times to those previously reported for control wild-type backgrounds [16].

mtDNA extraction and quantitative PCR of heteroplasmy. For all mtDNA analyses, total input and affinity-purified mitochondria were lysed and extracted using phenol-chloroform-isoamyl alcohol, followed by an ethanol precipitation. Extracted DNA was resuspended in 10 mM Tris-HCl (pH 8), 0.1 mM EDTA. Multiplex quantitative PCR was then performed using the Luna Universal Probe qPCR Master Mix (ref M3004, NEB), a Roche LightCycler or Rotor Gene Q real-time PCR (Qiagen) machine and Rotor Gene Q pure detection software (V2.3.1). The primers 5'-cgtaagaaaatcaaaatatggtataattgg-3', 5'-aaaatgttacgatccttattaataaagc-3', 5'-gcagagatgtttattgaagctgac-3' and the probes 5'-HEX/tgaggccag/ZEN/ttcatattgttccaga gtg/IABKFQ-3' (Iowa Black FQ) and 5'-6-FAM/ccatccgtg/ZEN/ctagaagacaaagaatttc/IABKFQ-3' were used to quantify the wild-type mtDNA and ΔmtDNA levels. Calculations of mtDNA copies per mitochondrion were derived from the quantification of purified organelles in individual samples, as previously performed [16]. Control wild-type values have been previously reported by Ahier et al., 2018 [16].

Quantification of mtDNA copy number

Ten (10) L4 worms from each genetic background were lysed in 10 μl of standard lysis buffer

supplemented with 200 ng ml−1 proteinase K and incubated at 65°C for 2 h, followed by heat-

inactivation at 95 °C for 15 min. A 1/50 dilution of the lysate was used for subsequent

quantitative PCR. The Sensifast SYBR no-rox Kit (ref. BIO-98005, Bioline) and a Rotor Gene Q

Real-Time PCR (Qiagen) Machine, with Rotor Gene Q pure detection software (V2.3.1), were

used for quantitative PCR (qPCR). The primers SZ35 (5'-AGGCTAAGCCGGGGTAAGTT-3') and

SZ36 (5'-GCCAAAAGCTTAAACTGCGG-3') were used to quantify the nuclear DNA and the

primers   AA178   (5'-CATACCGAATAAACATCAGGGTAATCT-3'),   AA179   (5'-

ACGGGTGTTACACTATGATGAAGA-3') were used to quantify the mtDNA levels. The cycle

conditions were 95°C for 10 min, followed by 45 cycles of 95°C for 15 s and 58°C for 40 s.

Primer efficiencies were determined by amplifying a series of exponential dilutions (1 to

10^{-8} ng) of cloned wild-type mtDNA (pSZ116) and *ges-1* promoter (pSZ57) fragments,

confirming an equivalent efficiency of both sets of primers (91.2% for SZ35 and SZ36 and

90.6% for AA178 and AA179).

Image acquisition and processing. Imaging was performed on live animals mounted on a 2%

agarose pad on glass slides with 1 mM levamisole (Sigma). For quantification of purified

mitochondria, we visualized fluorescence using a Zeiss Z2 imager microscope with a Zeiss

Axiocam 506 mono camera and Zen2 (version 2.0.0.0) software. Mitochondria were counted

with the aid of the Image J Grid plugin, and comparisons were made on the same

immunoprecipitation volume for each sample. For 3D reconstruction, imaging/analysis was

performed at the Queensland Brain Institute's Advanced Microscopy Facility using a spinning-

disk confocal system (Marianas; 3I, Inc.) consisting of an Aanio Observer Z1 (Carl Zeiss)

equipped with a CSU-W1 spinning-disk head (Yokogawa Corporation of America), ORCA-

Flash4.0 v2 sCMOS camera (Hamamatsu Photonics), and 20x 0.8 NA PlanApo and 40x 1.2 NA

C-Apo objectives. Image acquisition was performed using SlideBook 6.0 (3I, Inc). 3D

reconstruction and mitochondrial volume calculations were performed using 3D rendering in

Imaris software (version 8.4.1, Bitplane).

Thrashing assays. Thrashing assays were performed on worms suspended in 40 µl of M9

buffer on a hydrophobic slide (ref G350308BK, ProSciTech). After an initial 3 min period, a 30

second video sequence of the thrashing animals was recorded using a Nikon SMZ745T

stereomicroscope and a TrueChrome IIS camera (Tucsen Photonics). The number of body

bends per minute, as well as other mobility metrics, was determined using CeleST Computer

Vision Software as described previously [46].

Statistics and reproducibility. The statistical analyses are described for each figure. Generally,

an unpaired parametric two-tailed *t*-test was used to determine the significance of

comparisons made between two data sets, unless otherwise noted in the figure legend. One-

way analysis of variance (ANOVA) was performed for comparisons across multiple

independent samples, using Tukey's multiple comparisons correction. All experiments were

reproduced at least 3 times with similar results. All data and biological reagents that support

the conclusions of this manuscript are available from the corresponding author on request.

Acknowledgements

We are grateful to M. Hilliard and R. Tweedale for comments on the manuscript and members

of the Zuryn laboratory for discussions and comments. We also thank A. Gaudin for support

with microscopy. Some strains were provided by the CGC, which is funded by the NIH Office

of Research Infrastructure Programs (P40 OD010440). This work was supported by NHMRC

Project Grants GNT1128381 and GNT1162553, a Stafford Fox Senior Research Fellowship to

S.Z., ARC Discovery grant DP160103812 to J.G. and a University of Queensland International

Scholarship to C.Y.D.

413 Author contributions

414 A.A. carried out most experiments. N.C., C.Y.D., S.Z. contributed some experiments. J.G.

415 supervised N.C. and contributed to the interpretation of experiments. A.A., N.C., C.Y.D and

416 S.Z. designed and interpreted experiments. A.A. and S.Z. wrote the paper. All authors

417 approved the final manuscript.

419 Conflict of interest

420 The authors declare that they do not have any financial or non-financial competing interests.

428 Figures

Figure 1. The presence of ΔmtDNA alters mitochondrial volume in a tissue-dependent manner. (**A-D**) Representative three-dimensional reconstructions of *C. elegans'* (**A**) complete nervous system mitochondria, (**B**) intestinal mitochondria, (**C**) epidermal mitochondria, and (**D**) body wall muscle mitochondria in (**a**) wild-type animals and (**b**) animals harbouring ΔmtDNA. Scale bar, 50 μm. (**E-H**) Quantification of mitochondrial volumes from each tissue type. Columns are means ± s.e.m. of individual animals (*n* = 5). ** = *p*< 0.01. *P* values were determined using one-way ANOVA and Tukey's correction.

Figure 2. The presence of ΔmtDNA alters mtDNA copy number in a tissue-specific manner.

(**A-D**) Quantification of mtDNA molecules per mitochondrion. Each dot represents the average number of mtDNA molecules per mitochondrion from at least 1,000 mitochondria analysed from (**A**) neurons, (**B**) intestinal cells, (**C**) epidermal cells, and (**D**) body wall muscle cells. Bars are means of three biologically independent experiments ± s.e.m. * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$. *P* values were determined using one-way ANOVA and Tukey's correction. (**E-H**) Animals harbouring wild-type mtDNA or ΔmtDNA have equivalent levels of body wall muscle activity as determined by the thrashing assay. (**E** and **G**) Representative heatmaps of an individual animal's body angle over time as it thrashes in liquid. (**F** and **H**) Quantification of body bends per minute. BWM, body wall muscle. Columns are means ± s.e.m. of individual animals (dots). Two-sided *t*-test. L4, larval stage 4; 1 DOA, 1-day-old adult; 5 DOA, 5-day-old adult.

Figure 3. Deletion of *pink-1* and *pdr-1* increases heteroplasmy in a tissue-specific manner

and removes ΔmtDNA mosaicism. (A) Schematic of *C. elegans* and mitochondria originating

from major somatic tissue types in which ΔmtDNA (*uaDf5*) heteroplasmy was determined. (**B-**

F) Percentage of ΔmtDNA in mitochondria purified from each tissue type in L4 animals,

relative to total homogenate. Bars are means ± s.e.m. of three independent biological

replicates (dots) consisting of ~10,000 animals each. * = *p*< 0.05; ** = *p*< 0.01. *P* values were

determined using one-way ANOVA and Tukey's correction. Comparison of heteroplasmy

between tissues in (**G**) wild-type animals and (**G'**) *pink-1;pdr-1* double mutants indicates that

inter-tissue differences in ΔmtDNA heteroplasmy is lost upon deletion of *pink1* and *pdr-1*.

(**G''**) Graphical representation of the loss of stereotyped inter-tissue variation in ΔmtDNA

heteroplasmy in *pink-1;pdr-1* double mutant animals. Each tissue type is coloured according

to ΔHeteroplasmy, which is the difference in *uaDf5* heteroplasmy relative to BWM. The

significance of heteroplasmy variation between tissues was calculated using one-way ANOVA.

** = *p*< 0.01; *** = *p*< 0.001. *P* values were determined using one-way ANOVA and Tukey's

correction. (**H**) Quantitative PCR analysis of mtDNA copy number in L4 wild type and *pink-*

1;pdr-1 double mutants. Columns are means $\pm$ s.e.m. of three independent biological

replicates (dots), each consisting of 10 animals, analysed with a two-sided *t*-test.

Figure 4. Expression of proteotoxic proteins in neurons increases ΔmtDNA heteroplasmy.

(**A**) A *pink-1* translation fluorescent reporter is expressed in most somatic tissues in *C. elegans*, including in the (**A**) intestine, (**B**), epidermis, (**C**) neurons, and (**D**) body wall muscle. Yellow arrowheads indicate the axon, while the white arrowhead indicates the cell body of a neuron. White scale bar, 50μm; red scale bar, 5μm. (**E**) Expression of either human tau (hTau) or a polyglutamine tract (Q40) in neurons increases the heteroplasmy levels of ΔmtDNA (*uaDf5*) in the nervous system. Bars are means ± s.e.m. of at least three independent biological replicates (dots) consisting of ~10,000 animals each performed at day 1 of adulthood. *** = *p*< 0.001. *P* values were determined using one-way ANOVA and Tukey's correction. (**F**) Proposed model that leads to mosaicism of mtDNA. PINK-1-PDR-1-mediated mitophagy displays distinct activities in different tissue types which leads to the formation of a stereotyped pattern of ΔmtDNA heteroplasmy. In neurons, the expression of proteotoxic proteins can inhibit mitophagy, leading to an increase in heteroplasmy in the nervous system.

Supplemental Figure Legends

Figure S1.

(**A**) Map of *C. elegans* mtDNA showing the location of the 3.1kb *uadF5* deletion. (**B**)

Quantification of *uaDf5* heteroplasmy in whole animal extracts. Heteroplasmy was stable at

around 50-60% for each of the strains used. Bar represents the mean value of three

independent experiments ± s.e.m.

Figure S2. Changes in mitochondrial volume in response to ΔmtDNA and *pink-1/pdr-1*

mutation in different cell types. (**A-D**) Representative microimages corresponding to the

three-dimensional reconstructions shown in Figure 2. (**A**) complete nervous system

mitochondria, (**B**) intestinal mitochondria, (**C**) epidermal mitochondria, and (**D**) body wall

muscle mitochondria. Images presented in Figure 2 were reconstructed using Imaris software

and straightened using ImageJ for presentation. Scale bar, 50μm.

Figure S3.

(**A**) Schematic of the CS-MAP (cell-specific mitochondrial affinity purification) process to

isolate mitochondria from specific tissue types in *C. elegans*. (**B**) Microscope image of purified

mitochondria isolated from body wall muscle cells (white arrowheads). Scale bar, 10μm.

Figure S4.

Advanced behavioural analyses measurement show that *uaDf5* does not compromise body

wall muscle (BWM) function. Behavioural analyses were performed in animals harbouring the

uaDf5 mtDNA mutation at different heteroplasmy levels (~60 and ~93%) and at various ages.

(**A**) Attenuation measures how well the depth of a body-wave is maintained as it propagates

from the head to the tail of the animal as it swims in liquid. It will detect any deformation in

BWM contraction along the length of the body. (**B**) Stretch measures the maximum

differences in curvature of the body. (**C**) Asymmetry evaluates how balanced the swim

posture is and whether the animal preferentially bends toward one side. (**D**) Brush stroke

evaluates the full extent of movement and flexibility of the animal. Bars and columns are

means ± s.e.m from individual animals (dots).

Figure S5. Deletion of *pink-1* and *pdr-1* results in an increase in mitochondrial volume in body wall muscles. (**A-D**) Representative microimages corresponding to the three-dimensional reconstructions shown in E-H. (**A**) complete nervous system mitochondria, (**B**) intestinal mitochondria, (**C**) epidermal mitochondria, and (**D**) body wall muscle mitochondria. Scale bar, 50μm. (**E-H**) Representative three-dimensional reconstructions of *C. elegans'* (**A**) complete nervous system mitochondria, (**B**) intestinal mitochondria, (**C**) epidermal mitochondria, and (**D**) body wall muscle mitochondria in (**a**) animals harbouring ΔmtDNA, and (**b**) *pink-1(tm1779);pdr-1(gk448)* animals harbouring ΔmtDNA. For (**H**), dashed boxes are magnified in (**a'-b'**). (**I-L**) Quantification of mitochondrial volumes from each tissue type. Columns are means ± s.e.m. of individual animals (*n* = 5). ** = *p*< 0.01. *P* values were determined using one-way ANOVA and Tukey's correction.

References

1.   West AP, Shadel GS (2017) Mitochondrial DNA in innate immune responses and
inflammatory pathology. *Nat Rev Immunol* **17**: 363-375

2.   Spinelli JB, Haigis MC (2018) The multifaceted contributions of mitochondria to
cellular metabolism. *Nat Cell Biol* **20**: 745-754

3.   Gorman GS, Chinnery PF, DiMauro S, Hirano M, Koga Y, McFarland R, Suomalainen A,
Thorburn DR, Zeviani M, Turnbull DM (2016) Mitochondrial diseases. *Nat Rev Dis Primers* **2**:
16080

4.   Morris J, Na YJ, Zhu H, Lee JH, Giang H, Ulyanova AV, Baltuch GH, Brem S, Chen HI,
Kung DK*, et al.* (2017) Pervasive within-Mitochondrion Single-Nucleotide Variant
Heteroplasmy as Revealed by Single-Mitochondrion Sequencing. *Cell Reports* **21**: 2706-2713

5.   Hahn A, Zuryn S (2019) The Cellular Mitochondrial Genome Landscape in Disease.
Trends Cell Biol **29**: 227-240

6.   Stewart JB, Chinnery PF (2015) The dynamics of mitochondrial DNA heteroplasmy:
implications for human health and disease. *Nat Rev Genet* **16**: 530-42

7.   McManus MJ, Picard M, Chen HW, De Haas HJ, Potluri P, Leipzig J, Towheed A,
Angelin A, Sengupta P, Morrow RM*, et al.* (2019) Mitochondrial DNA Variation Dictates
Expressivity and Progression of Nuclear DNA Mutations Causing Cardiomyopathy. *Cell
Metab* **29**: 78-90 e5

8.   Valenci I, Yonai L, Bar-Yaacov D, Mishmar D, Ben-Zvi A (2015) Parkin modulates
heteroplasmy of truncated mtDNA in Caenorhabditis elegans. *Mitochondrion* **20**: 64-70

9.   Lin YF, Schulz AM, Pellegrino MW, Lu Y, Shaham S, Haynes CM (2016) Maintenance
and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded
protein response. *Nature* **533**: 416-9

10.   Gitschlag BL, Kirby CS, Samuels DC, Gangula RD, Mallal SA, Patel MR (2016)
Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in C. elegans. *Cell
Metab* **24**: 91-103

11.   Suen DF, Narendra DP, Tanaka A, Manfredi G, Youle RJ (2010) Parkin overexpression
selects against a deleterious mtDNA mutation in heteroplasmic cybrid cells. *Proc Natl Acad
Sci U S A* **107**: 11835-40

12.   Kandul NP, Zhang T, Hay BA, Guo M (2016) Selective removal of deletion-bearing
mitochondrial DNA in heteroplasmic Drosophila. *Nat Commun* **7**: 13100

13.   Wang Y, Michikawa Y, Mallidis C, Bai Y, Woodhouse L, Yarasheski KE, Miller CA,
Askanas V, Engel WK, Bhasin S*, et al.* (2001) Muscle-specific mutations accumulate with
aging in critical human mtDNA control sites for replication. *Proc Natl Acad Sci U S A* **98**:
4022-7

14.   Samuels DC, Li C, Li B, Song Z, Torstenson E, Boyd Clay H, Rokas A, Thornton-Wells
TA, Moore JH, Hughes TM*, et al.* (2013) Recurrent tissue-specific mtDNA mutations are
common in humans. *PLoS Genet* **9**: e1003929

15.   Soong NW, Hinton DR, Cortopassi G, Arnheim N (1992) Mosaicism for a Specific
Somatic Mitochondrial-DNA Mutation in Adult Human Brain. *Nature Genetics* **2**: 318-323

16.   Ahier A, Dai C-Y, Tweedie A, Bezawork-Geleta A, Kirmes I, Zuryn S (2018) Affinity
purification of cell-specific mitochondria from whole animals resolves patterns of genetic
mosaicism. *Nat Cell Biol* **20**: 352-360

17.   Cortopassi GA, Shibata D, Soong NW, Arnheim N (1992) A pattern of accumulation of
a somatic deletion of mitochondrial DNA in aging human tissues. *Proc Natl Acad Sci U S A*
89: 7370-4

18.   Tsang WY, Lemire BD (2002) Stable heteroplasmy but differential inheritance of a
large mitochondrial DNA deletion in nematodes. *Biochem Cell Biol* **80**: 645-54

19.   Diaz F, Bayona-Bafaluy MP, Rana M, Mora M, Hao H, Moraes CT (2002) Human
mitochondrial DNA with large deletions repopulates organelles faster than full-length
genomes under relaxed copy number control. *Nucleic Acids Research* **30**: 4626-4633

20.   Gieseler K, Qadota H, Benian GM (2017) Development, structure, and maintenance
of C. elegans body wall muscle. *WormBook* **2017**: 1-59

21.   Jiang M, Kauppila TES, Motori E, Li X, Atanassov I, Folz-Donahue K, Bonekamp NA,
Albarran-Gutierrez S, Stewart JB, Larsson NG (2017) Increased Total mtDNA Copy Number
Cures Male Infertility Despite Unaltered mtDNA Mutation Load. *Cell Metab* **26**: 429-436 e4

22.   Pickles S, Vigie P, Youle RJ (2018) Mitophagy and Quality Control Mechanisms in
Mitochondrial Maintenance. *Curr Biol* **28**: R170-R185

23.   Springer W, Hoppe T, Schmidt E, Baumeister R (2005) A Caenorhabditis elegans
Parkin mutant with altered solubility couples alpha-synuclein aggregation to proteotoxic
stress. *Hum Mol Genet* **14**: 3407-23

24.   Cummins N, Tweedie A, Zuryn S, Bertran-Gonzalez J, Götz J (2019) Disease-
associated tau impairs mitophagy by inhibiting Parkin translocation to mitochondria. *EMBO*
J **38**

25.   Palikaras K, Lionaki E, Tavernarakis N (2015) Coordination of mitophagy and
mitochondrial biogenesis during ageing in C. elegans. *Nature* **521**: 525-8

26.   Kirienko NV, Ausubel FM, Ruvkun G (2015) Mitophagy confers resistance to
siderophore-mediated killing by Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **112**:
1821-6

27.   Ryu D, Mouchiroud L, Andreux PA, Katsyuba E, Moullan N, Nicolet-Dit-Felix AA,
Williams EG, Jha P, Lo Sasso G, Huzard D*, et al.* (2016) Urolithin A induces mitophagy and

prolongs lifespan in C. elegans and increases muscle function in rodents. *Nat Med* **22**: 879-
88

28.   Palikaras K, Lionaki E, Tavernarakis N (2018) Mechanisms of mitophagy in cellular
homeostasis, physiology and pathology. *Nat Cell Biol* **20**: 1013-1022

29.   Harper JW, Ordureau A, Heo JM (2018) Building and decoding ubiquitin chains for
mitophagy. *Nat Rev Mol Cell Biol* **19**: 93-108

30.   Samann J, Hegermann J, von Gromoff E, Eimer S, Baumeister R, Schmidt E (2009)
Caenorhabditis elegans LRK-1 and PINK-1 act antagonistically in stress response and neurite
outgrowth. *J Biol Chem* **284**: 16482-91

31.   Polanco JC, Li C, Bodea LG, Martinez-Marmol R, Meunier FA, Götz J (2018) Amyloid-
beta and tau complexity - towards improved biomarkers and targeted therapies. *Nat Rev*
Neurol **14**: 22-39

32.   Franco-Iborra S, Plaza-Zabala A, Sebastian D, Vila M, Martinez-Vicente M (2019)
Impairment of neuronal mitophagy by mutant huntingtin. *bioRxiv*

33.   Gidalevitz T, Ben-Zvi A, Ho KH, Brignull HR, Morimoto RI (2006) Progressive
disruption of cellular protein folding in models of polyglutamine diseases. *Science* **311**:
1471-4

34.   Picard M, Shirihai OS, Gentil BJ, Burelle Y (2013) Mitochondrial morphology
transitions and functions: implications for retrograde signaling? *Am J Physiol Regul Integr*
Comp Physiol **304**: R393-406

35.   Trifunovic A, Wredenberg A, Falkenberg M, Spelbrink JN, Rovio AT, Bruder CE,
Bohlooly YM, Gidlof S, Oldfors A, Wibom R, *et al.* (2004) Premature ageing in mice
expressing defective mitochondrial DNA polymerase. *Nature* **429**: 417-23

36.   Kujoth GC, Hiona A, Pugh TD, Someya S, Panzer K, Wohlgemuth SE, Hofer T, Seo AY,
Sullivan R, Jobling WA, *et al.* (2005) Mitochondrial DNA mutations, oxidative stress, and
apoptosis in mammalian aging. *Science* **309**: 481-4

37.   Filograna R, Koolmeister C, Upadhyay M, Pajak A, Clemente P, Wibom R, Simard ML,
Wredenberg A, Freyer C, Stewart JB, *et al.* (2019) Modulation of mtDNA copy number
ameliorates the pathological consequences of a heteroplasmic mtDNA mutation in the
mouse. *Sci Adv* **5**: eaav9824

38.   Ashrafi G, Schlehe JS, LaVoie MJ, Schwarz TL (2014) Mitophagy of damaged
mitochondria occurs locally in distal neuronal axons and requires PINK1 and Parkin. *J Cell*
Biol **206**: 655-70

39.   Bingol B, Tea JS, Phu L, Reichelt M, Bakalarski CE, Song Q, Foreman O, Kirkpatrick DS,
Sheng M (2014) The mitochondrial deubiquitinase USP30 opposes parkin-mediated
mitophagy. *Nature* **510**: 370-5

40.   Pickrell AM, Huang CH, Kennedy SR, Ordureau A, Sideris DP, Hoekstra JG, Harper JW,
Youle RJ (2015) Endogenous Parkin Preserves Dopaminergic Substantia Nigral Neurons
following Mitochondrial DNA Mutagenic Stress. *Neuron* **87**: 371-81

41.   Valente EM, Abou-Sleiman PM, Caputo V, Muqit MM, Harvey K, Gispert S, Ali Z, Del
Turco D, Bentivoglio AR, Healy DG*, et al.* (2004) Hereditary early-onset Parkinson's disease
caused by mutations in PINK1. *Science* **304**: 1158-60

42.   Kitada T, Asakawa S, Hattori N, Matsumine H, Yamamura Y, Minoshima S, Yokochi M,
Mizuno Y, Shimizu N (1998) Mutations in the parkin gene cause autosomal recessive juvenile
parkinsonism. *Nature* **392**: 605-8

43.   Kraytsberg Y, Kudryavtseva E, McKee AC, Geula C, Kowall NW, Khrapko K (2006)
Mitochondrial DNA deletions are abundant and cause functional impairment in aged human
substantia nigra neurons. *Nat Genet* **38**: 518-20

44.   Bender A, Krishnan KJ, Morris CM, Taylor GA, Reeve AK, Perry RH, Jaros E, Hersheson
JS, Betts J, Klopstock T*, et al.* (2006) High levels of mitochondrial DNA deletions in substantia
nigra neurons in aging and Parkinson disease. *Nat Genet* **38**: 515-7

45.   Brenner S (1974) The genetics of Caenorhabditis elegans. *Genetics* **77**: 71-94

46.   Restif C, Ibanez-Ventoso C, Vora MM, Guo SZ, Metaxas D, Driscoll M (2014) CeleST:
Computer vision software for quantitative analysis of C. elegans swim behavior reveals
novel features of locomotion. *Plos Comp Biol* **10**
665