PINK1-parkin-mediated mitophagy generates stereotyped somatic mosaicism

of the mitochondrial genome

Arnaud Ahier, Chuan-Yang Dai, and Steven Zuryn*.

The University of Queensland, Queensland Brain Institute, Clem Jones Centre for Ageing
Dementia Research, Brisbane, Australia.
Corresponding author.

Abstract

Mitochondria are critical for complex life and are characterized by the presence of their own

genome (mtDNA). The mtDNA makeup within each cell is in a constant state of flux through

processes of mutation, replication, and degradation, resulting in a mosaic mtDNA landscape

that inevitably varies between cells, tissues, and organs within individuals. However, despite

the stochastic nature of these processes, mosaic patterns of mtDNA mutations can become

stereotyped across the tissues of individuals in both invertebrate and vertebrate species. The

mechanisms that determine the non-random spatiotemporal distribution of mtDNA

mutations are unknown. We find that PTEN induced putative kinase (PINK1) and the E3

ubiquitin-protein ligase parkin drive the formation of mtDNA heteroplasmy disparity between

the major somatic tissue types of *C. elegans*, generating a stereotyped genetic mosaicism of

the mitochondrial genomic landscape. PINK1 and parkin are conserved mediators of

mitochondrial autophagy (mitophagy), but while PINK1/parkin preferentially direct the

removal of mtDNA mutations in neurons, intestinal cells, and hypodermal cells, they act non-

selectively in muscle cells to reduce mitochondrial network volume. These data suggest that

different cell types use alternative strategies to cope with mtDNA mutations and implicate

the nuclear genome encoded PINK1/parkin signalling axis in shaping stereotyped

mitochondrial genomic mosaicism across individuals.

Introduction

Mitochondria act as central hubs for cellular bioenergetics, co-factor and macromolecule

precursor synthesis, redox balance, Ca^{2+} handling, apoptosis, and immunity (Spinelli and

Haigis, 2018; West and Shadel, 2017). These organelles house their own genome (mtDNA), as

well as RNA and protein-synthesizing systems that together code and coordinate the

assembly of core subunits of the oxidative phosphorylation (OXPHOS) machinery used to generate the cell's primary energy substrate, ATP. Although mtDNA lesions that perturb OXPHOS assembly can cause devastating metabolic disorders (Gorman et al., 2016), mutations are buffered by the polyploidy of the mitochondrial genome. Hundreds to thousands of mtDNA copies can populate each cell, which can result in a mixture of multiple mtDNA variants within individual cells and organelles (Morris et al., 2017). This state, termed heteroplasmy, and the factors that influence its dynamics are critical determinants for the pathogenesis of rare mitochondrial diseases, and possibly a wide range of common age-onset inflictions, including neurodegeneration, diabetes, and cancer (Hahn and Zuryn, 2018; Stewart and Chinnery, 2015).

At any given time, an interplay between stochastic and deterministic processes influence cellular heteroplasmy (Hahn and Zuryn, 2018). Mitotic segregation, genetic drift, and the competing effects of homeostatic (Gitschlag et al., 2016; Kandul et al., 2016; Lin et al., 2016; Suen et al., 2010; Valenci et al., 2015) and retrograde signalling pathways (Gitschlag et al., 2016; Lin et al., 2016) may contribute to the divergence of heteroplasmy levels between the cells within a tissue, creating heteroplasmy mosaicism. However, there is evidence that specific organs and tissues are prone to accumulating certain mtDNA mutations more than others, leading to a stereotyped pattern of heteroplasmy mosaicism (Ahier et al., 2018; Samuels et al., 2013; Soong et al., 1992; Wang et al., 2001). For example, the caudate, putamen and substantia nigra regions of the brain show an age-dependent increase in the heteroplasmy levels of the mtDNA^{4977} deletion (a mutant genome harbouring a 4,977bp deletion), relative to the rest of the brain (Soong et al., 1992). In the invertebrate *C. elegans*, stereotyped heteroplasmy differences have also been observed. Here, mtDNA molecules harbouring a large deletion accumulate at different heteroplasmy levels in distinct somatic and germ lineages in a reproducible manner (Ahier et al., 2018). However, it is unknown how these non-random patterns mtDNA heteroplasmy between tissues evolves within individuals.

Here, we have determined that cell-type-specific roles for PTEN-induced putative kinase 1 (PINK1) and the E3 ubiquitin-protein ligase parkin generate stereotyped patterns of mtDNA mosaicism across major somatic tissues in *C. elegans*. PINK1 and parkin are conserved mediators of mitochondrial autophagy (mitophagy), a selective form of autophagy that

removes depolarized mitochondria (Palikaras et al., 2018). The defective organelles are

decorated with polyubiquitin chains on their outer surface via PINK1 recruitment of parkin,

and are subsequently captured by autophagosomes and destroyed following fusion with

lysosomes (Harper et al., 2018). We show that mutations in PINK1 and parkin reverse the

stereotyped divergence of mtDNA heteroplasmy observed between neuronal, muscle,

intestinal, and hypodermal tissue types. Moreover, we reveal that distinct tissue types use

unique strategies to manage mtDNA mutations, which may be dependent upon metabolic

requirements. Our results provide a mechanistic rationale for stereotyped mtDNA mosaicism

in the soma and provide insights into the processes that contribute to the evolving mtDNA

landscape within individuals in both healthy and diseased states.

Results and Discussion

In order to determine whether conserved homeostatic mechanisms such as mitophagy shape

stereotyped mosaic patterns of somatic mtDNA distribution, we introduced a heteroplasmic

variant of mtDNA that harbours a 3.1 kb deletion (*uaDf5*, also called ΔmtDNA) (Tsang and

Lemire, 2002) into double-mutant strains with null deletions in the genes PINK1 (*pink-1*) and

parkin (*pdr-1*). The *uaDf5* deletion exists in heteroplasmy with wild-type copies of the mtDNA

at a level of approximately 60% across populations of animals (Lin et al., 2016; Tsang and

Lemire, 2002). We next crossed these triple mutant strains to transgenic animals carrying a

mos1-mediated single-copy insertion (mosSCI) of *TOMM-20::mKate2::HA* controlled by either

a pan-neuronal promoter (*rgef-1p*), intestinal-specific promoter (*ges-1p*), hypodermis

(epidermis)-specific promoter (*dpy-7p*), or body wall muscle (BWM)-specific promoter (*myo-*

3p). Using this panel of strains, we performed cell-type-specific mitochondrial affinity

purification (CS-MAP) (Ahier et al., 2018), allowing comparisons of mtDNA heteroplasmy

across major tissue types in animals lacking PINK-1-PDR-1-mediated mitophagy (Figure 1A).

Heteroplasmy in cell-type-specific mitochondria was investigated across very large

populations of pooled animals (>10,000 individuals, see methods). This approach enabled

population-scale trends to be resolved by overcoming confounding influences that can be

introduced by the presence of stochastic inter-individual variations in mosaicism.

Mitochondria were purified from both *pink-1(tm1779);pdr-1(gk448)* double-mutant and wild-

type backgrounds carrying ΔmtDNA. To accurately quantify ΔmtDNA heteroplasmy levels

from each tissue type, we used a fluorescence multiplex quantitative PCR (qPCR) assay (Ahier

et al., 2018). To eliminate the influence of any inter-population variability affecting our

analyses, we compared mitochondria purified from each cell type to the total mitochondria

(homogenate) from the same samples. In addition, we performed the same experiments in

animals in which mitochondria from all cell types were purified using the same CS-MAP

technique. In this case, the ubiquitous promoter *etf-3p* was used to drive the expression of a

TOMM-20::mKate2::HA transgene integrated into the same genomic location as the other

transgenes. As expected, ΔmtDNA heteroplasmy levels did not change between total input

mitochondria and mitochondria purified from all cells of wild-type and *pink-1(tm1779);pdr-*

1(gk448) backgrounds (Figure 1B). This confirmed that deficiencies in PINK-1 and PDR-1 did

not influence the CS-MAP procedure.

However, we found that removal of the PINK-1-PDR-1 mitophagy pathway in mitochondria

purified from neurons, intestine, and hypodermis resulted in a significant increase in ΔmtDNA

heteroplasmy levels (Figure 1B). This suggests that mitophagy operates within these tissues

to selectively remove ΔmtDNA molecules. In wild-type animals, mitochondria isolated from

neurons and intestinal cells displayed lower levels of heteroplasmy relative to the

homogenate. However, inactivating PINK-1-PDR-1-mediated mitophagy completely reversed

this trend, suggesting that PINK-1-PDR-1-dependent mitophagy is particularly important in

these cell types for maintaining heteroplasmy levels below the average for the whole animal

(Figure 1B). Interestingly, we found that removal of *pink-1* and *pdr-1* had no effect on ΔmtDNA

heteroplasmy levels in BWM cells, indicating that PINK-1-PDR-1-mediated mitophagy was not

utilised in this tissue type for mitochondrial genome quality control (Figure 1B and 1C).

Moreover, removal of *pink-1* and *pdr-1* abolished any differences in ΔmtDNA heteroplasmy

levels between all of the major tissue types studied (Figure 1D), suggesting that PINK-1-PDR-

1-mediated mitophagy acts to drive stereotyped mosaicism of the mitochondrial genome.

Different cell types have distinct functions and therefore have varying energetic demands.

The BWM is composed of four quadrants of somatic muscle bundles, each of which consists

of 23-24 cells arranged into staggered pairs that stretch longitudinally from the head to the

tail of the animal (Gieseler et al., 2017). The coordinated contractile activities of these bundles

produce powerful and energetically demanding sinusoidal body bends that drive locomotion.

We therefore hypothesised that BWM cells might avoid mitophagy in combating mtDNA
mutations so as to prevent reductions of the mitochondrial network. To test this idea, we
analysed the volume of mitochondria in all tissues using three-dimensional reconstruction of
fluorescence images acquired from strains carrying *TOMM-20::mKate2::HA* (Figure 2). We
found that the presence of ΔmtDNA tended to increase the volume of the mitochondrial
network in neurons, intestine, and hypodermis, although significant increases were only
observed in hypodermal tissue (Figure 2A-C and 2E-G). Mitochondrial volume remained
unchanged in BWM tissue in the presence of ΔmtDNA (Figure 2D and H). However, we found
that removal of *pink-1* and *pdr-1* in this background almost tripled the volume of the
mitochondrial network in BWM cells (Figure 2Dc and H). This suggests, as previously
demonstrated during ageing (Palikaras et al., 2015), that PINK-1-PDR-1-mediated mitophagy
controls the mitochondrial density in *C. elegans* BWM cells. However, unlike in other major
tissue types, mitophagy does not counteract the presence of deleterious mitochondrial
genomes in the muscle. In contrast, whereas PINK-1-PDR-1-mediated mitophagy appeared to
preferentially coordinate the removal of ΔmtDNA in the nervous system, intestine, and
hypodermis, it did not appear to play a role in shaping mitochondrial volume, which remained
unchanged when both *pink-1* and *pdr-1* were removed in ΔmtDNA backgrounds (Figure 2A-C
and 2E-G). One possible explanation for this observation is that cells with high energy
demand, such as muscle cells, stimulate high amounts of mitochondrial biogenesis that is
balanced through mitophagy-mediated removal. As such, under high mitochondrial turnover
conditions, mitophagy may fulfil a different primary role and therefore lose selectivity
towards the removal of genetically faulty mitochondria. Indeed, we found that mitophagy
restricted the mitochondrial networks to almost one third the volume they would otherwise
be in BWM cells, a workload that may saturate the mitophagy machinery. A similar role was
not observed in the other major tissue types. Moreover, in wild-type backgrounds, BWM cells
had the highest ΔmtDNA heteroplasmy levels of any somatic tissue type (Figure 1B),
suggesting that mtDNA quality control is, overall, less efficient in muscle.

The *uaDf5* deletion disrupts four genes that encode core subunits of the electron transport
chain and as such, ΔmtDNA is insufficient for the assembly of a functional OXPHOS system.
Despite this, we found that BWM activity was unperturbed by the presence of ΔmtDNA.
Animals carrying ΔmtDNA could still produce sinusoidal body bends under strenuous

conditions (thrashing assay, see materials and methods) at rates comparable to those of their
wild-type counterparts, even when genetically bottlenecked to very high heteroplasmy levels
(93.39 ± 0.34%) and tested at various ages up to 5-day old adults (Figure 3A). This suggested
that BWM cells could effectively compensate for the presence of deleterious mtDNA in order
to maintain function over their lifetimes. Recently demonstrated in mouse sperm, one
potential mechanism to compensate for deleterious heteroplasmic mtDNA mutations is to
increase the copy number of total mitochondrial genomes (Jiang et al., 2017). Although not
helping to remove or repair the mutations themselves, energy supplies might be restored if
the expression level of mtDNA-encoded components is increased through an indiscriminate
elevation in both mutant and wild-type genome copy number. We therefore investigated
whether BWM cells responded to the presence of ΔmtDNA by enhancing mtDNA replication.
Indeed, we found that in all tissues except hypodermis, the mtDNA copy number per
mitochondrion increased in the presence of ΔmtDNA (Figure 3B). However, the largest
absolute increase was observed in BWM mitochondria, with almost 2 additional copies of
mtDNA in each mitochondrion (Figure 3C). This suggests that BWM mitochondria have a
greater capacity to respond to ΔmtDNA by perpetuating a higher overall copy number of
mtDNA per mitochondrion, which may act as an alternative mechanism for maintaining cell
function in the absence of mitophagy-mediated mtDNA quality control.

Together, our results suggest that major and distinct tissue systems comprising the three
primary germ layers manage deleterious mtDNA mutations by differentially adopting
mitophagy and mtDNA biogenesis processes (Figure 3D). Moreover, we have determined that
stereotyped distribution patterns of mitochondrial genomes harbouring deletions are caused
by disparate roles of PINK-1-PDR-1-mediated mitophagy in different cell types. Interestingly,
large mtDNA deletions have been shown to follow stereotyped patterns of accumulation in
specific regions of the human brain during ageing (Soong et al., 1992). In the context of our
findings, it would be interesting to determine whether such changes are the result of
alterations in mitophagy processes in these cells. Indeed, the increase in heteroplasmy levels
observed following removal of *pink-1* and *pdr-1* was the greatest in the nervous system. This
suggests that neurons are heavily reliant upon mitophagy for the surveillance and removal of
mtDNA deletions. Accumulating evidence suggests that mtDNA mutations contribute to the
progressive nature of age-related neurodegenerative diseases (Stewart and Chinnery, 2015)

and mitophagy itself has also been shown to be disrupted in models of neurodegeneration
(Cummins et al., 2018; Ye et al., 2015). Indeed, mutations in both PINK1 (Valente et al., 2004)
and parkin (Kitada et al., 1998) cause early-onset Parkinson's disease, suggesting that
perturbations to mitophagy promote neurodegeneration, which in light of our results, may
be attributable to the nervous system's reliance on mitophagy for quality control of the
neuronal mtDNA landscape.

Figures

Fig.1 *pink-1* and *pdr-1* determine stereotyped patterns of ΔmtDNA somatic mosaicism. (A)
Schematic of *C. elegans* and mitochondria originating from major somatic tissue types.
TOMM-20::mKate2::HA transgenes driven by tissue-specific promoters were integrated into
the *oxTi185* site on chromosome 1. In these transgenic animals, cell-specific mitochondrial
affinity purification (CS-MAP) was used to purify mitochondria of each cell type from whole
animals. Mitochondria harbour both wild-type and ΔmtDNA (*uaDf5*) genomes, a situation
known as heteroplasmy. The percentages of each genome were quantified from each sub-
population of mitochondria obtained from each cell type. (**B**) Percentage of ΔmtDNA in
mitochondria purified from each tissue type, relative to total homogenate. +, animals with
functional *pink-1* and *pdr-1* genes; -, animals with deletions in *pink-1* and *pdr-1*. Bars are
means ± s.e.m. *P* values were determined using one-way ANOVA and Tukey's correction. (**C**)
Summary of the swing in ΔmtDNA % caused by deletion of *pink-1* and *pdr-1* genes, relative to

wild-type animals. (**D**) Graphical representation of the loss of stereotyped inter-tissue

variation in ΔmtDNA heteroplasmy in *pink-1;pdr-1* double mutant animals. Each tissue type is

coloured according to ΔHeteroplasmy, which is the difference in *uaDf5* heteroplasmy relative

to BWM. The significance of heteroplasmy variation between tissues was calculated using

one-way ANOVA.

Fig. 2 PINK-1-PDR-1 mediated mitophagy controls mitochondrial network volume in muscle

cells. (**A-D**) Representative three-dimensional reconstructions of *C. elegans* (**A**) Complete

nervous system mitochondria, (**B**) intestinal mitochondria, (**C**) hypodermal mitochondria, and

(**D**) body wall muscle (BWM) mitochondria in (**a**) wild-type animals, (**b**) animals harbouring

ΔmtDNA, and (**c**) animals carrying ΔmtDNA in the absence of *pink-1* and *pdr-1* genes (**A-D**,

scale bar, 50 μm). For (**D**), dashed boxes are magnified in (**a'-c'**) (scale bar, 20 μm). (**E-H**)

Quantification of mitochondrial volumes from each tissue type. Columns are means ± s.e.m.

of individual animals (dots). *P* values were determined using one-way ANOVA and Tukey's

correction.

Fig. 3 Tissue-specific mtDNA replication in response to the presence of ΔmtDNA. (**A**)

Quantification of body wall muscle (BWM) function using the thrashing assay. ΔmtDNA

heteroplasmy levels are shown above each graph. Columns are means ± s.e.m. of individual

animals (dots). Two-sided *t*-test. L4, larval stage 4; 1 DOA, 1-day old adult; 5 DOA, 5-day old

adult. (**B**) Quantification of mtDNA molecules per mitochondrion (see materials and

methods). -, animals not carrying ΔmtDNA; +, animals carrying ΔmtDNA. Bars are means ±

s.e.m. *P* values were determined using one-way ANOVA and Tukey's correction. (**C**) Summary

of change in mtDNA copy number per mitochondrion caused by the presence of ΔmtDNA,

relative to wild-type animals. (**D**) Model summarizing the distinct mechanisms by which

different cell types manage mtDNA deletions. Neurons, intestinal cells, and hypodermal cells

use PINK-1-Parkin to reduce ΔmtDNA heteroplasmy levels, whereas muscle cells do not. Only

neurons, intestinal cells, and muscle cells induce mtDNA replication in response to the

presence of ΔmtDNA, with muscle cell mitochondria showing the greatest induction of this

response.

Materials and Methods

C. elegans **strains and culture**. LB138 *him-8(e1489) IV; uaDf5/+*; EG8078 *oxTi185 I; unc-*

119(ed3) III; BR4006 *pink-1(tm1779) II; byEx655 [pink-1p::pink-1::GFP; myo-2p::mCherry;*

herring sperm DNA]; VC1024 *pdr-1(gk448) III* and wild-type (N2) were acquired from the

Caenorhabditis Genetics Center (CGC), University of Minnesota, USA, which is funded by the

NIH Office of Research Infrastructure Programs (P40 OD010440). *C. elegans* culture and

maintenance were performed according to standard protocols (Brenner, 1974). The

transgenic strains used for cell-specific mitochondrial purification experiments were

generated as previously described (Ahier et al., 2018).

Cell-specific Mitochondrial Affinity Purification (CS-MAP). CS-MAP was performed as

previously described (Ahier et al., 2018). Briefly, 10,000 to 20,000 L4 staged worms were

grown on 2 x 150 mm NGM plates seeded with *E. coli* OP50 bacteria. Worms were

homogenised into a hypotonic buffer [50 mM KCl, 110 mM mannitol, 70 mM sucrose, 0.1 mM

EDTA (pH 8.0), 5 mM Tris-HCl (pH 7.4)] with a dounce homogenizer and the subsequent crude

mitochondrial fraction was enriched by differential centrifugations and mitochondria were

isolated from this fraction using anti-HA (Influenza Hemagglutinin) magnetic beads (Pierce).

Beads were washed and resuspended in 20 µl of hypotonic buffer. All CS-MAP experiments

were repeated in at least three independent experiments and data for each genetic

background were obtained at similar times as previously reported in Ahier et al., 2018 for

control wild-type backgrounds.

mtDNA extraction and Quantitative PCR. For all mtDNA analyses, total input and affinity-

purified mitochondria were lysed and extracted using phenol-chloroform-isoamyl alcohol,

followed by an ethanol precipitation. Extracted DNA was resuspended in 10 mM Tris-HCl (pH

8), 0.1 mM EDTA. Then, multiplex quantitative PCR was performed using the Luna Universal

Probe qPCR Master Mix (ref M3004, NEB), a Rotor Gene Q Real-Time PCR (Qiagen) Machine

and Rotor Gene Q pure detection software (V2.3.1). The primers 5'-

cgtaagaaaatcaaaatatggtataattgg-3', 5'-aaaatgttacgatccttattaataaagc-3', 5'-

gcagagatgtttattgaagctgac-3' and the probes 5'-HEX/tgaggccag/ZEN/ttcatattgttccaga

gtg/IABKFQ-3' (Iowa Black FQ) and 5'-6-FAM/ccatccgtg/ZEN/ctagaagacaaagaatttc/IABKFQ-3'

were used to quantify the wild-type mtDNA and ΔmtDNA levels. Calculations of mtDNA copies

per mitochondrion were derived from the quantification of purified organelles in individual

samples, as previously performed (Ahier et al., 2018). Control wild-type values are reported

in Ahier et al., 2018.

Image acquisition and processing. Imaging was performed on live animals mounted on a 2% agarose pad on glass slides with 1 mM levamisole (Sigma). For quantification of purified mitochondria, we visualized fluorescence using a Zeiss Z2 imager microscope with a Zeiss Axiocam 506 mono camera and Zen2 (version 2.0.0.0) software. Mitochondria were counted with the aid of the Image J Grid plugin, and comparisons were made on the same immunoprecipitation volume for each sample. For 3D reconstruction, images were acquired using a spinning-disk confocal system (Marianas; 3I, Inc.) consisting of an Aanio Observer Z1 (Carl Zeiss) equipped with a CSU-W1 spinning-disk head (Yokogawa Corporation of America), ORCA-Flash4.0 v2 sCMOS camera (Hamamatsu Photonics), 20x 0.8 NA PlanApo and 40x 1.2 NA C-Apo objectives. Image acquisition was performed using SlideBook 6.0 (3I, Inc). 3D reconstruction and mitochondrial volume calculations were performed using 3D rendering in Imaris software (version 8.4.1, Bitplane). Imaging was performed at the Queensland Brain Institute's Advanced Microscopy Facility.

Thrashing assays. Thrashing assays were performed on worms suspended in 40 µl of M9 buffer on a hydrophobic slide (ref G350308BK, ProSciTech). After an initial 3 min period, a 30 second video sequence of the thrashing animals was recorded using a Nikon SMZ745T stereomicroscope and a TrueChrome IIS camera (Tucsen Photonics). Body bends per minutes were determined using CeleST Computer Vision Software as described previously (Restif et al., 2014).

Statistics and Reproducibility. Statistical analyses performed are described for each figure. One-way analysis of variance (ANOVA) was performed for comparisons across multiple independent samples, using Tukey's multiple comparisons correction. All experiments were reproduced at least 3 times with similar results.

Data and reagent availability. All data and biological reagents that support the conclusions of this manuscript are available from the corresponding author on request.

Acknowledgements

We are grateful to R. Tweedale for comments on the manuscript and members of the Zuryn laboratory for discussions and comments. We also thank A. Gaudin for support with

microscopy. Some strains were provided by the CGC, which is funded by the NIH Office of

Research Infrastructure Programs (P40 OD010440). This work was supported by NHMRC

Project Grants GNT1128381 and GNT1162553, a Stafford Fox Senior Research Fellowship to

S.Z., and a University of Queensland International Scholarship to C.Y.D.

Contributions

A.A. carried out most experiments. C.Y.D., S.Z. contributed some experiments. A.A., C.Y.D and

S.Z. designed and interpreted experiments and wrote the paper.

Competing Interest

The authors declare that they do not have any financial or non-financial competing interests.

References

Ahier, A., Dai, C.-Y., Tweedie, A., Bezawork-Geleta, A., Kirmes, I., and Zuryn, S. (2018).
Affinity purification of cell-specific mitochondria from whole animals resolves patterns of
genetic mosaicism. Nat Cell Biol *20*, 352-360.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.

Cummins, N., Tweedie, A., Zuryn, S., Bertran-Gonzalez, J., and Gotz, J. (2018). Disease-
associated tau impairs mitophagy by inhibiting Parkin translocation to mitochondria. EMBO
J.

Gieseler, K., Qadota, H., and Benian, G.M. (2017). Development, structure, and maintenance
of C. elegans body wall muscle. WormBook *2017*, 1-59.

Gitschlag, B.L., Kirby, C.S., Samuels, D.C., Gangula, R.D., Mallal, S.A., and Patel, M.R. (2016).
Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in C. elegans. Cell
Metab *24*, 91-103.

Gorman, G.S., Chinnery, P.F., DiMauro, S., Hirano, M., Koga, Y., McFarland, R., Suomalainen,
A., Thorburn, D.R., Zeviani, M., and Turnbull, D.M. (2016). Mitochondrial diseases. Nat Rev
Dis Primers *2*, 16080.

Hahn, A., and Zuryn, S. (2018). The Cellular Mitochondrial Genome Landscape in Disease.
Trends Cell Biol.

Harper, J.W., Ordureau, A., and Heo, J.M. (2018). Building and decoding ubiquitin chains for
mitophagy. Nat Rev Mol Cell Biol *19*, 93-108.

Jiang, M., Kauppila, T.E.S., Motori, E., Li, X., Atanassov, I., Folz-Donahue, K., Bonekamp, N.A.,
Albarran-Gutierrez, S., Stewart, J.B., and Larsson, N.G. (2017). Increased Total mtDNA Copy

Number Cures Male Infertility Despite Unaltered mtDNA Mutation Load. Cell Metab *26*, 429-
436 e424.

Kandul, N.P., Zhang, T., Hay, B.A., and Guo, M. (2016). Selective removal of deletion-bearing
mitochondrial DNA in heteroplasmic Drosophila. Nat Commun *7*, 13100.

Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S., Yokochi,
M., Mizuno, Y., and Shimizu, N. (1998). Mutations in the parkin gene cause autosomal
recessive juvenile parkinsonism. Nature *392*, 605-608.

Lin, Y.F., Schulz, A.M., Pellegrino, M.W., Lu, Y., Shaham, S., and Haynes, C.M. (2016).
Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial
unfolded protein response. Nature *533*, 416-419.

Morris, J., Na, Y.J., Zhu, H., Lee, J.H., Giang, H., Ulyanova, A.V., Baltuch, G.H., Brem, S., Chen,
H.I., Kung, D.K.*, et al.* (2017). Pervasive within-Mitochondrion Single-Nucleotide Variant
Heteroplasmy as Revealed by Single-Mitochondrion Sequencing. Cell Reports *21*, 2706-
2713.

Palikaras, K., Lionaki, E., and Tavernarakis, N. (2015). Coordination of mitophagy and
mitochondrial biogenesis during ageing in C. elegans. Nature *521*, 525-528.

Palikaras, K., Lionaki, E., and Tavernarakis, N. (2018). Mechanisms of mitophagy in cellular
homeostasis, physiology and pathology. Nature Cell Biology *20*, 1013-1022.

Restif, C., Ibanez-Ventoso, C., Vora, M.M., Guo, S.Z., Metaxas, D., and Driscoll, M. (2014).
CeleST: Computer vision software for quantitative analysis of C. elegans swim behavior
reveals novel features of locomotion. Plos Comp Biol *10*.

Samuels, D.C., Li, C., Li, B., Song, Z., Torstenson, E., Boyd Clay, H., Rokas, A., Thornton-Wells,
T.A., Moore, J.H., Hughes, T.M.*, et al.* (2013). Recurrent tissue-specific mtDNA mutations are
common in humans. PLoS Genet *9*, e1003929.

Soong, N.W., Hinton, D.R., Cortopassi, G., and Arnheim, N. (1992). Mosaicism for a Specific
Somatic Mitochondrial-DNA Mutation in Adult Human Brain. Nature Genetics *2*, 318-323.

Spinelli, J.B., and Haigis, M.C. (2018). The multifaceted contributions of mitochondria to
cellular metabolism. Nat Cell Biol *20*, 745-754.

Stewart, J.B., and Chinnery, P.F. (2015). The dynamics of mitochondrial DNA heteroplasmy:
implications for human health and disease. Nat Rev Genet *16*, 530-542.

Suen, D.F., Narendra, D.P., Tanaka, A., Manfredi, G., and Youle, R.J. (2010). Parkin
overexpression selects against a deleterious mtDNA mutation in heteroplasmic cybrid cells.
Proc Natl Acad Sci U S A *107*, 11835-11840.

Tsang, W.Y., and Lemire, B.D. (2002). Stable heteroplasmy but differential inheritance of a
large mitochondrial DNA deletion in nematodes. Biochem Cell Biol *80*, 645-654.

Valenci, I., Yonai, L., Bar-Yaacov, D., Mishmar, D., and Ben-Zvi, A. (2015). Parkin modulates
heteroplasmy of truncated mtDNA in Caenorhabditis elegans. Mitochondrion *20*, 64-70.

Valente, E.M., Abou-Sleiman, P.M., Caputo, V., Muqit, M.M., Harvey, K., Gispert, S., Ali, Z.,
Del Turco, D., Bentivoglio, A.R., Healy, D.G*., et al.* (2004). Hereditary early-onset Parkinson's
disease caused by mutations in PINK1. Science *304*, 1158-1160.

Wang, Y., Michikawa, Y., Mallidis, C., Bai, Y., Woodhouse, L., Yarasheski, K.E., Miller, C.A.,
Askanas, V., Engel, W.K., Bhasin, S*., et al.* (2001). Muscle-specific mutations accumulate with
aging in critical human mtDNA control sites for replication. Proc Natl Acad Sci U S A *98*,
4022-4027.

West, A.P., and Shadel, G.S. (2017). Mitochondrial DNA in innate immune responses and
inflammatory pathology. Nat Rev Immunol *17*, 363-375.

Ye, X., Sun, X., Starovoytov, V., and Cai, Q. (2015). Parkin-mediated mitophagy in mutant
hAPP neurons and Alzheimer's disease patient brains. Hum Mol Genet *24*, 2938-2951.
398