PINK1 and parkin shape the organism-wide distribution of a deleterious mitochondrial genome

Arnaud Ahier[1], Nadia Cummins[1], Chuan-Yang Dai[1], Jürgen Götz[1] and Steven Zuryn[1]*.

[1] *Clem Jones Centre for Ageing Dementia Research, Queensland Brain Institute, The University of Queensland, Brisbane, Australia, 4072.*

Corresponding author.

Running title: PINK-1-parkin shape mutant mtDNA distribution

Keywords: Mitochondria, mtDNA, mitophagy, PINK1, parkin, heteroplasmy, proteotoxicity

Abstract

In multiple species, certain tissue types are prone to acquiring greater loads of

mitochondrial genome (mtDNA) mutations relative to others, however the

mechanisms that drive these heteroplasmy differences are unknown. We found that

the conserved PTEN-induced putative kinase (PINK1/PINK-1) and the E3 ubiquitin-

protein ligase parkin (PDR-1), which are required for mitochondrial autophagy

(mitophagy), underlie stereotyped differences in heteroplasmy of a deleterious

mitochondrial genome mutation (ΔmtDNA) between major somatic tissues types in

Caenorhabditis elegans. We demonstrate that tissues prone to accumulating

ΔmtDNA have lower mitophagy responses than those with low mutation levels, such

as neurons. Moreover, we show that ΔmtDNA heteroplasmy increases when

proteotoxic species that are associated with neurodegenerative disease and

mitophagy inhibition are overexpressed in the nervous system. Together, these

results suggest that PINK1 and parkin drive organism-wide patterns of heteroplasmy

and provide evidence of a causal link between proteotoxicity, mitophagy, and mtDNA

mutation levels in neurons.

Introduction

Mitochondria act as central hubs for cellular bioenergetics, macromolecule precursor

synthesis, redox balance, Ca^{2+} handling, apoptosis, and immunity [1, 2]. They house

their own genome (mtDNA), as well as RNA and protein-synthesizing systems,

which together code and coordinate the assembly of core subunits of oxidative

phosphorylation (OXPHOS). Mutations in the mtDNA that perturb the assembly of

OXPHOS enzymes can cause devastating metabolic disorders [3]. The level of

heteroplasmy of a pathogenic mtDNA mutation correlates with the severity of the

clinical phenotype and mosaic distributions of mutations across an individual have

been reported to lead to organ-specific dysfunction [4]. Heteroplasmy and mosaicism

are therefore important determinants of mitochondrial disease pathophysiology.

The composition of the mitochondrial DNA (mtDNA) in each cell is in constant

flux through processes of mutation, replication, and degradation. Mitotic segregation

of mitochondria may result in asymmetric proportions of heteroplasmic allelic

variants in descendant cellular lineages [5]. Alternatively, due to the high rate and

relaxed replication of mtDNA, mosaicism can arise through clonal expansion and

subsequent genetic drift between adjacent cells and tissues over time [6]. Indeed,

there is continuous replication of mtDNA in all tissues, allowing variations in

heteroplasmy to even develop between post-mitotic cells. Although there is evidence

of selectivity [7], it is generally assumed that mitotic segregation and genetic drift are

largely stochastic processes and therefore lead to random variation in mtDNA

heteroplasmy and mosaicism within individuals [4]. Evidence in several species,

however, suggests that differences in heteroplasmy between certain cell types is not

random and may instead be deterministic. For example, relative to the rest of the

human brain, the caudate, putamen and substantia nigra regions show an age-

dependent increase in the heteroplasmy levels of the mtDNA[4977] deletion (a mutant

genome harbouring a 4,977bp deletion) [8]. Heteroplasmy differences between

tissues have also been observed in *C. elegans*. Here, mtDNA molecules harbouring

a 3.1 kb deletion (ΔmtDNA) accumulate at different heteroplasmy levels in distinct

somatic and germ lineages over time [9, 10]. How different cell, tissue, or organ

types reproducibly develop higher heteroplasmy levels than others is unclear.

By purifying and analysing mitochondria from different cell types, we show

that PINK-1 and PDR-1 are required to set heteroplasmy differences between major

somatic tissue groups in *C. elegans* and that removal of these genes equalises

ΔmtDNA heteroplasmy across the organism. Mechanistically, our evidence indicates

that heteroplasmy is lowered by PINK-1 and PDR-1 through the selective removal of

ΔmtDNA, but to varying degrees of efficiency in different cell types with neurons,

intestinal cells, and epidermal cells being more amenable to ΔmtDNA clearance than

body wall muscle cells. As a result, body wall muscle cells accumulate ΔmtDNA at

higher levels than the other tissues. Interestingly, we find that neurons, which have

the greatest dependency upon PINK-1 and PDR-1 for maintaining low ΔmtDNA

heteroplasmy levels, are susceptible to increases in heteroplasmy caused by

overexpressing either tau or polyglutamate repeats in the nervous system,

proteotoxic species associated with neurodegenerative disease and mitophagy

inhibition. Together, our results uncover a nuclear-encoded mechanism that

determines organism-wide patterns of mtDNA heteroplasmy and establishes links

between proteotoxicity, mitophagy, and mtDNA mutation levels in neurons.

Results

To understand the mechanisms that underlie the divergence in heteroplasmy levels

between different tissue types within individuals, we took advantage of a strain of *C.*

elegans that harbours a 3.1 kb deletion in its mitochondrial genome (*uaDf5*, also

called ΔmtDNA). This mutation exists in stable heteroplasmy of 50-60% (Figure S1A)

and is deleterious for mitochondrial function as it disrupts four mtDNA genes that

encode core subunits of the respiratory chain (Figure S1B). Worms harbouring the

uaDf5 allele at around 60% heteroplasmy display reductions in basal oxygen

consumption and total respiratory capacity as well as constitutive induction of the

mitochondrial unfolded protein response (UPR^{mt}) [11, 12]. Because cellular

heteroplasmy is determined by the net result of the propagation and degradation of

mtDNA mutations, we investigated whether mitophagy, a process that can

selectively destroy depolarized mitochondria harbouring mtDNA mutations, such as

ΔmtDNA [11-13], may underlie tissue-specific heteroplasmy levels. To test this idea,

we introduced the ΔmtDNA mutation into double-mutant strains with null deletions in

the genes *pink-1* and *pdr-1* [14], key mediators of mitophagy in multiple species

including *C. elegans* [15-18]. PINK1/PINK-1 can induce mitophagy by accumulating

on the outer mitochondrial membrane of defective organelles, where it recruits the

E3 ubiquitin ligase parkin/PDR-1 [19]. Parkin/PDR-1 decorates the surface of these

mitochondria with polyubiquitin chains, thereby inducing engulfment by

autophagosomes and subsequent destruction following fusion with lysosomes [20].

In some invertebrate species such as *Drosophila*, parkin can act independently of

PINK1 during mitophagy [21], prompting us to use *pink-1(tm1779);pdr-1(gk448)*

double mutants to unambiguously disrupt mitophagy in *C. elegans*. *Pink-*

1(tm1779);pdr-1(gk448) double mutants harbouring heteroplasmic ΔmtDNA were

crossed to a series of transgenic strains carrying a *TOMM-20::mKate2::HA*

transgene under the control of either the promoter of the gene *myo-3, ges-1, rgef-1*,

or *dpy-7* which selectively drive body wall muscle, intestinal, panneuronal, and

epidermal expression, respectively (Figure 1A). We also generated a similar strain

which carried the transgene *TOMM-20::mKate2::HA* under the ubiquitous control of

the promoter of *eft-3*. Intact mitochondria were isolated from each of the strains

above, as well as wild-type animals in which mitophagy was unperturbed. This was

performed by using immunoprecipitation against the HA epitope, which labelled the

outer surface of the mitochondria because of its attachment to the C-terminus of

TOMM-20 [9]. In doing so, we were able to selectively purify mitochondria from each

tissue type in which the *TOMM-20::mKate2::HA* transgene was expressed.

Heteroplasmy   comparisons   in   animals   expressing   *eft-3p::TOMM-*

20::mKate2::HA revealed that there was no relative difference in ΔmtDNA levels

between immunoprecipitated mitochondria and total (non-affinity purified input

sample) mitochondria (Figure 1B). This was as expected and indicated that the

mitochondrial isolation procedure used did not artificially alter heteroplasmy level

measurements   made   in   either   wild-type   or   *pink-1(tm1779);pdr-1(gk448)*

backgrounds. Comparisons in relative ΔmtDNA levels made between mitochondria

purified from each tissue type in wild-type worms revealed that neurons, intestinal

cells, epidermal cells, and body wall muscle cells displayed distinct and reproducible

differences in ΔmtDNA heteroplasmy relative to each other (Figures 1C-F), with

muscle cells always harbouring the highest load of mutations of any somatic tissue

type   studied   (Figure   1G)[9].   Interestingly,   these   intertissue   differences   in

heteroplasmy were abolished in *pink-1(tm1779);pdr-1(gk448)* mutants (Figure 1H

and 1I), suggesting that the activities of *pink-1* and *pdr-1*, two nuclear-encoded

genes, were required for setting distinct and stereotyped patterns of heteroplasmy in

major somatic tissues of *C. elegans*. This equalisation effect wasn't caused by global

changes in mtDNA copy number, as *pink-1(tm1779);pdr-1(gk48)* mutants had normal

levels of mtDNA (Figure 1J). At the level of individual tissues in *pink-1(tm1779);pdr-*

1(gk488) double mutants, we observed increases in heteroplasmy in neurons,

intestinal cells, and epidermal cells (Figure 1C-E), but no alteration in ∆mtDNA levels

in body wall muscle cells (Figure 1F). This suggested that *pink-1* and *pdr-1* acted

only in select cell types to remove ∆mtDNA, leading to stereotyped differences in

heteroplasmy between tissues over time.

158   Because these results indicated that *pink-1* and *pdr-1* did not influence

heteroplasmy in muscle cells, we hypothesised that the genes were either inactive or

that they had different functions that did not include ∆mtDNA clearance in this tissue

type. However, we found that *pink-1* was expressed in body wall muscle cells, as

indicated by a *pink-1p::PINK-1::GFP* [22] transgene in live animals (Figure 2A).

Because a translational GFP::PDR-1 reporter is also expressed in body wall muscles

in *C. elegans* [14], it is likely that PINK-1-PDR-1-mediated mitophagy is active in this

tissue.   In *C. elegans*, mitophagy can be monitored through the colocalization of

mitochondria with autophagosomes, which defines one of the terminal stages of

mitophagy prior to lysomal fusion and the destruction of encapsulated mitochondria

[16, 23]. By quantifying the colocalization of *myo-3p::*TOMM-20::mKate2::HA and

lgg-1p::LGG-1::GFP [24], which label muscle mitochondria and autophagosomes,

respectively, we determined that mitophagy was inhibited in the body wall muscle

cells of *pink-1(tm1779);pdr-1(gk488)* worms (Figure 2B). Indeed, in these mutants,

using multiple fluorescent markers (*myo-3p::TOMM-20::mKate2::HA* and *myo-*

3p::^{MTS}RFP) and three-dimensional reconstruction of high magnification images, we

also observed a dramatic increase in total mitochondrial muscle volume (Figure 2C,

D and Figure S3), suggesting that the mitochondrial network increased in size in the

absence of mitophagy-mediated turnover. Together, these results indicate that *pink-*

1 and *pdr-1* are active and required for mitophagy in muscle cells in *C. elegans*,

consistent with previous findings [16].

Mitophagy is initiated by the stabilisation of PINK1 on the surface of

depolarised mitochondria. However, although mitochondria are ubiquitous, they can

exhibit cell-type specific morphologies, dynamics, and functions and as such,

mitochondria from distinct cell-types may respond physiologically differently to the

presence of ΔmtDNA in a manner that impinges on mitophagy and therefore

ΔmtDNA clearance. For example, if the mitochondrial membrane potential (MMP) is

unaffected by ΔmtDNA in muscle cells, it is possible that mitochondria harbouring

ΔmtDNA may not be recognised and removed via mitophagy. To investigate this

possibility further, we compared the effects of ΔmtDNA in body wall muscle cells and

neurons, two cell types with opposite heteroplasmy trends (Figure 1C and F).

Although gross mitochondrial morphology and volume appeared to be unaffected by

ΔmtDNA (Figure S3), we found that ΔmtDNA induced a decrease in the MMP in both

muscle and neural mitochondria to the same degree (Figure 3A and B). Furthermore,

we found that total mtDNA copy number increased in mitochondria purified from

muscle cells and neurons, which is likely to be a compensatory response to

mitochondrial dysfunction caused by the presence of ΔmtDNA (Figure 3C). This

suggests that ΔmtDNA invokes comparable levels of mitochondrial dysfunction and

depolarisation in muscle cells and neurons and that tissue-specific differences in

heteroplasmy, as determined by *pink-1* and *pdr-1* activities, are not attributable to

intrinsic differences in mitochondrial physiology between cell types.

Although *pink-1* and *pdr-1* are expressed and are required for mitophagy in

both body wall muscle (Figure 2B) and neurons, as we showed previously in *C.*

elegans [15], it is possible that mitophagy responds differently to the presence of

ΔmtDNA in each cell type, which may affect its clearance. We compared tissue-

specific levels of mitophagy in the nervous system and body wall muscle in wild-type

animals and animals harbouring ΔmtDNA by determining the percentage overlap

(volume:volume) of mitochondria (muscle, *myo-3p::TOMM-20::mKate2::HA*; neural

rgef-1p::TOMM-20::mKate2::HA) to autophagosomes (*lgg-1p::LGG-1::GFP*).

Interesting, we found that mitophagy was induced by the presence of ΔmtDNA in

both cell types, suggesting that mtDNA mutations can directly activate mitochondrial

clearance pathways *in vivo* (Figure 3D). More importantly, we found that relative to

mitochondrial volume, there was much more basal mitophagy in neurons compared

to muscle cells. Moreover, although both tissue types responded to the presence of

ΔmtDNA by increasing mitophagy, the neural response was much greater than that

in muscle (Figure 3D). This suggests that mitophagy has a greater capacity to

respond to ΔmtDNA in the nervous system than in muscle and could therefore

explain why neurons have lower heteroplasmy levels than muscle and why

mutations in *pink-1* and *pdr-1(gk488)* have a greater effect on heteroplasmy in the

nervous system.

Because our results indicated that ΔmtDNA strongly induces mitophagy in the

neurons and that the activity of PINK-1-PDR-1-mediated mitophagy is important in

the nervous system for maintaining low levels of heteroplasmy, we next determined

whether other cellular conditions that perturbed mitophagy could raise ΔmtDNA

levels in these cells types. We recently demonstrated that proteotoxic tau, a protein

that forms aggregates in tauopathies such as Alzheimer's disease [25], perturbs

parkin translocation to the mitochondria, and that human tau expressed in the *C.*

elegans nervous system inhibits mitophagy [15]. To determine whether this

increased heteroplasmy levels in neurons, we constructed strains harbouring

∆mtDNA and expressing human tau (hTau) in the nervous system (*aex-3p::hTau*)

[15] as well as the neuron-specific *rgef-1p::TOMM-20::mKate2::HA* transgene

(Figure 4A). We then immunoprecipitated neuronal mitochondria in these strains and

found that hTau significantly increased heteroplasmy (Figure 4B), suggesting that

the expression of mitophagy-inhibiting proteins can increase ∆mtDNA levels in

neurons. Other proteotoxic species have also recently been found to inhibit

mitophagy in neurons. The expanded polyglutamine (polyQ) tract in mutant forms of

the Huntingtin protein (Htt) that cause Huntington disease were recently reported to

impede mitophagy in differentiated striatal neurons [26]. Indeed, we found that the

expression of a polyglutamine (polyQ) tract (*rgef-1p::Q40::YFP*) [27] in the *C.*

elegans nervous system recapitulated the effect of hTau and increased ∆mtDNA

heteroplasmy in neurons (Figure 4B). These results suggest that diverse proteotoxic

threats associated with a broad range of neurodegenerative diseases may promote

the accumulation of mtDNA deletions in the nervous system.

Discussion

Stochastic processes such as mitotic segregation and genetic drift influence cellular

heteroplasmy [4]. However, certain tissues are prone to accumulating mtDNA

mutations, implying that cell-type specific events deterministically set heteroplasmy

levels. Our results suggest that cell-type differences in the activity and

responsiveness of mitophagy, mediated through *pink-1* and *pdr-1*, contribute to the

intertissue divergence in heteroplasmy of a deleterious mitochondrial genome. We

showed that deletion of *pink-1* and *pdr-1* equalised heteroplasmy levels across

tissues that normally display stereotypical differences in ΔmtDNA heteroplasmy.

Comparisons between body wall muscle cells and neurons revealed that although

active in both tissues, mitophagy responded to ΔmtDNA at greater levels relative to

total mitochondrial volume in neurons than muscle, leading to higher ΔmtDNA

heteroplasmy in muscle. Furthermore, we have demonstrated that proteotoxic

threats that can perturb mitophagy in neurons can increase heteroplasmy levels in

the nervous system. Taken together, our results suggest a model (Figure 4C)

whereby organism-wide patterns of heteroplasmy of mtDNA deletions are

determined by the activity of PINK-1-PDR-1-mediated mitophagy and that genetic

perturbations and disease-related proteotoxic threats that perturb mitophagy can

increase the abundance of mtDNA mutations in the nervous system of an organism.

In humans, large mtDNA deletions have been shown to accumulate in specific

regions of the brain during ageing [8]. Although there is strong evidence that PINK1-

parkin-mediated mitophagy occurs in mammalian neurons [28, 29], it is not known

whether differences in the activity of mitophagy plays a role in specifying the

heteroplasmy levels of mtDNA deletions within distinct regions of the brain. In mice

homozygous for a proofreading deficiency in DNA polymerase γ, which induces a

high level of mtDNA single nucleotide mutations [30, 31], parkin knockout

significantly increases the levels of pathogenic mtDNA mutations in the striatum [32].

Moreover, mutations in Pink1 [33] and parkin [34] can lead to early onset Parkinson's

disease, and substantia nigra neurons in Parkinson's disease patients have high

levels of mtDNA deletions [35, 36]. Although we did not examine subsets of neurons

in this study, our results suggest that neurons in general are the most vulnerable to

increases in heteroplasmy when mitophagy is perturbed. Therefore, our work

demonstrates that mutations in Pink1 and parkin have a widely conserved effect

across phyla of enhancing the accumulation of mtDNA deletions in neurons, which

may perturb their function and cause disease. It will be interesting to determine

whether these principles are specific to mtDNA deletions or other types of mtDNA

mutations as well.

In addition to Pink1 and parkin mutations, we found that toxic proteins

associated with neurodegenerative diseases other than Parkinson's disease also

increase heteroplasmy. Overexpression in neurons of hTau, as well as a polyQ tract,

significantly increased ΔmtDNA heteroplasmy levels in the *C. elegans* nervous

system. To our knowledge, this represents the first reported incidence of

proteotoxicity increasing the cellular load of mtDNA mutations. Interestingly, similar

proteotoxic species have been recently reported to inhibit mitophagy in neurons. Via

its projection domain, hTau can associate with Parkin in the cytoplasm and

specifically impair its translocation to the surface of defective mitochondria in human

neuroblastoma cells [15]. In addition, a polyQ tract within expanded mutant versions

of the Huntingting protein can prevent the formation of the Beclin1-Vps34 initiation

complex, negatively affecting the efficiency of mitophagy in post-mitotic striatal

neurons [26]. These results demonstrate that perturbing mitophagy indirectly with

proteotoxicity has the same effect as perturbing mitophagy genetically through

mutations in *pink-1* and *pdr-1*. They also demonstrate that toxic proteins associated

with diverse neurodegenerative diseases can modify the genetic landscape of the

mitochondrial genome in neurons and provide evidence that two of the most

common hallmarks of neurodegeneration - proteotoxicity and mitochondrial

dysfunction - may converge at the level of the mitochondrial genome.

Materials and Methods

***C. elegans* strains and culture**. LB138 *him-8(e1489) IV; uaDf5/+*; EG8078 *oxTi185*

I; unc-119(ed3) III; BR4006 *pink-1(tm1779) II; byEx655 [pink-1p::pink-1::GFP; myo-*

2p::mCherry; herring sperm DNA]; VC1024 *pdr-1(gk448) III*; AM101 *rmIs110 [rgef-*

1p::Q40::YFP], SJ4103 *zcIs14 [myo-3p::GFP^{mt}]*, MAH235 *sqIs13[lgg-1p::gfp::lgg-1 +*

odr-1p::RFP] [24] and wild-type (N2) were acquired from the *Caenorhabditis*

Genetics Center (CGC), University of Minnesota, USA, which is funded by the NIH

Office of Research Infrastructure Programs (P40 OD010440). APD178 *apdIs11 [aex-*

3p::tau4RIN + coel::gfp] were kindly provided by Dr H. Nicholas (The University of

Sydney, Sydney, Australia). QH5425 *vdIs30 [unc-25p::MTS::mRFP + odr-1p::dsRed]*

was kindly provided as a gift by Dr M. Hilliard and J. Chaplin (The University of

Queensland, Queensland Brain Institute, Brisbane, Australia). *C. elegans* culture and

maintenance were performed according to standard protocols [37]. The transgenic

strains used for cell-specific mitochondrial purification experiments, *foxSi16[myo-*

3p::TOMM-20::mKate2::HA], *foxSi37 [ges-1p::TOMM-20::mKate2::HA]*, *foxSi41[dpy-*

7p::TOMM-20::mKate2::HA], *foxSi44[rgef-1p::TOMM-20::mKate2::HA], foxSi75[eft-*

3p::TOMM-20::mKate2::HA] and *foxSi2[myo-3p:: ^{MTS}RFP]* were used were

generated as previously described [9].

Cell-specific mitochondrial affinity purification (CS-MAP). CS-MAP was

performed as previously described [9]. Heteroplasmy in mitochondria derived from

different cell types was investigated across very large populations of pooled animals

(>10,000 individuals). This approach enabled population-scale trends to be resolved

by overcoming confounding influences that can be introduced by the presence of

stochastic inter-individual variations. Briefly, 10,000 to 20,000 L4 staged worms were

grown on 2 x 150 mm NGM plates seeded with *E. coli* OP50 bacteria. We chose L4

animals as it represents a stage in which all somatic tissues are fully formed (e.g.

nervous system and epidermal seam cells) and developing embryos are not yet

present in the gonads of animals. Worms were homogenized into a hypotonic buffer

[50 mM KCl, 110 mM mannitol, 70 mM sucrose, 0.1 mM EDTA (pH 8.0), 5 mM Tris-

HCl (pH 7.4)] with a dounce homogenizer and the subsequent crude mitochondrial

fraction was enriched by differential centrifugations. The mitochondria were then

isolated from this fraction using anti-HA (influenza hemagglutinin) magnetic beads

(Pierce). The beads were washed and resuspended in 20 µl of hypotonic buffer. All

CS-MAP experiments were repeated in at least three independent experiments and

data for each genetic background were obtained at similar times to those previously

reported for control wild-type backgrounds [9]. To eliminate the possibility of any

population variability affecting our analyses, we compared mitochondria purified from

each tissue to the total mitochondria (homogenate) from the same samples.

mtDNA extraction and quantitative PCR of heteroplasmy. For all mtDNA

analyses, total input and affinity-purified mitochondria were lysed and extracted

using phenol-chloroform-isoamyl alcohol, followed by an ethanol precipitation.

Extracted DNA was resuspended in 10 mM Tris-HCl (pH 8), 0.1 mM EDTA. Multiplex

quantitative PCR was then performed using the Luna Universal Probe qPCR Master

Mix (ref M3004, NEB), a Roche LightCycler or Rotor Gene Q real-time PCR (Qiagen)

machine and Rotor Gene Q pure detection software (V2.3.1). The primers 5'-

cgtaagaaaatcaaaatatggtataattgg-3',   5'-aaaatgttacgatccttattaataaagc-3',   5'-

gcagagatgtttattgaagctgac-3' and the probes 5'-HEX/tgaggccag/ZEN/ttcatattgttccaga

gtg/IABKFQ-3'   (Iowa   Black   FQ)   and   5'-6-

FAM/ccatccgtg/ZEN/ctagaagacaaagaatttc/IABKFQ-3' were used to quantify the wild-

type mtDNA and ΔmtDNA levels. Calculations of mtDNA copies per mitochondrion

were derived from the quantification of purified organelles in individual samples, as

previously performed [9]. Control wild-type values have been previously reported by

Ahier et al., 2018 [9].

Quantification of mtDNA copy number

Ten (10) L4 worms from each genetic background were lysed in 10 μl of standard

lysis buffer supplemented with 200 ng ml–1 proteinase K and incubated at 65°C

for 2 h, followed by heat-inactivation at 95 °C for 15 min. A 1/50 dilution of the

lysate was used for subsequent quantitative PCR. The Sensifast SYBR no-rox Kit

(ref. BIO-98005, Bioline) and a Rotor Gene Q Real-Time PCR (Qiagen) Machine,

with Rotor Gene Q pure detection software (V2.3.1), were used for quantitative PCR

(qPCR). The primers SZ35 (5'-AGGCTAAGCCGGGGTAAGTT-3') and SZ36 (5'-

GCCAAAAGCTTAAACTGCGG-3') were used to quantify the nuclear DNA and the

primers   AA178   (5'-CATACCGAATAAACATCAGGGTAATCT-3'),   AA179   (5'-

ACGGGTGTTACACTATGATGAAGA-3') were used to quantify the mtDNA levels.

The cycle conditions were 95°C for 10 min, followed by 45 cycles of 95°C for 15 s

and 58°C for 40 s. Primer efficiencies were determined by amplifying a series of

exponential dilutions (1 to 10^{-8} ng) of cloned wild-type mtDNA (pSZ116) and *ges-1*

promoter (pSZ57) fragments, confirming an equivalent efficiency of both sets of

primers (91.2% for SZ35 and SZ36 and 90.6% for AA178 and AA179).

Image acquisition and processing. Imaging was performed on live animals

mounted on a 2% agarose pad on glass slides with 1 mM levamisole (Sigma). For

quantification of purified mitochondria and mitochondrial membrane potential, we

visualized fluorescence using a Zeiss Z2 imager microscope with a Zeiss Axiocam

506 mono camera and Zen2 (version 2.0.0.0) software. Mitochondria were counted

with the aid of the Image J Grid plugin, and comparisons were made on the same

immunoprecipitation volume for each sample. For 3D reconstruction,

imaging/analysis was performed at the Queensland Brain Institute's Advanced

Microscopy Facility using a spinning-disk confocal system (Marianas; 3I, Inc.)

consisting of an Aanio Observer Z1 (Carl Zeiss) equipped with a CSU-W1 spinning-

disk head (Yokogawa Corporation of America), ORCA-Flash4.0 v2 sCMOS camera

(Hamamatsu Photonics), and 20x 0.8 NA PlanApo and 40x 1.2 NA C-Apo objectives.

Image acquisition was performed using SlideBook 6.0 (3I, Inc). 3D reconstruction

and mitochondrial volume calculations were performed using 3D rendering in Imaris

software (version 8.4.1, Bitplane). For all mitochondrial and mitophagy volume

measurements, 100x magnification was used and representative images depict

multiple images stitched together and straightened using image J.

Mitochondrial membrane potential. Mitochondrial membrane potential was

measured by observing mitochondrial accumulation of the fluorescent dye

tetramethylrhodamine ethyl ester (TMRE; Sigma), as previously described [38, 39].

Briefly, 1-day old adult animals were transferred to growth plates seeded with 200 μl

of M9 buffer containing 1 mM of TMRE. Animals were allowed to grow for 15 hours

on the plates before being transferred to growth plates without TMRE for 5 hours to

allow excess dye to be cleared from the intestinal cavity. Live animals were then

prepared for fluorescence microscopy, as described above. To quantify the amount

of dye accumulation in mitochondria from each tissue type, a region of interest was

selected from images (40x magnification) containing mitochondria labelled with GFP

fused to an MTS and driven under promoters that express in either body wall muscle

(*myo-3*p::GFP^{mt}) [40], or the PLM neuron (*mec-4*p::^{MTS}GFP) [41]. Mean TMRE

intensity was calculated for mitochondria of individual animals using Image J

software.

Statistics and reproducibility. The statistical analyses are described for each

figure. Generally, an unpaired parametric two-tailed *t*-test was used to determine the

significance of comparisons made between two data sets, unless otherwise noted in

the figure legend. One-way analysis of variance (ANOVA) was performed for

comparisons across multiple independent samples, using Tukey's multiple

comparisons correction. All experiments were reproduced at least 3 times with

similar results. All data and biological reagents that support the conclusions of this

manuscript are available from the corresponding author on request.

Acknowledgements

We are grateful to M. Hilliard and R. Tweedale for comments on the manuscript and

members of the Zuryn laboratory for discussions and comments. We also thank A.

Gaudin for support with microscopy. Some strains were provided by the CGC, which

is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

This work was supported by NHMRC Project Grants GNT1128381 (S.Z.) and

GNT1162553 (S.Z.), ARC Discovery grants DP200101630 (S.Z.) and DP160103812

(J.G.), a Clem Jones Centre for Ageing Dementia Flagship grant (S.Z.), a Stafford

Fox Senior Research Fellowship to S.Z., and a University of Queensland

International Scholarship to C.Y.D.

Author contributions

433   A.A. carried out most experiments. N.C., C.Y.D., S.Z. contributed some experiments.

434   J.G. supervised N.C. and contributed to the interpretation of experiments. A.A., N.C.,

435   C.Y.D and S.Z. designed and interpreted experiments. A.A. and S.Z. wrote the

436   paper. All authors approved the final manuscript.

Conflict of interest

439   The authors declare that they do not have any financial or non-financial competing

440   interests.

References

References

1.   West AP, Shadel GS (2017) Mitochondrial DNA in innate immune responses and
inflammatory pathology. *Nat Rev Immunol* **17**: 363-375

2.   Spinelli JB, Haigis MC (2018) The multifaceted contributions of mitochondria to
cellular metabolism. *Nat Cell Biol* **20**: 745-754

3.   Gorman GS, Chinnery PF, DiMauro S, Hirano M, Koga Y, McFarland R, Suomalainen A,
Thorburn DR, Zeviani M, Turnbull DM (2016) Mitochondrial diseases. *Nat Rev Dis Primers* **2**:
16080

4.   Hahn A, Zuryn S (2019) The Cellular Mitochondrial Genome Landscape in Disease.
Trends Cell Biol **29**: 227-240

5.   Wilton PR, Zaidi A, Makova K, Nielsen R (2018) A Population Phylogenetic View of
Mitochondrial Heteroplasmy. *Genetics* **208**: 1261-1274

6.   Carelli V, Maresca A, Caporali L, Trifunov S, Zanna C, Rugolo M (2015) Mitochondria:
Biogenesis and mitophagy balance in segregation and clonal expansion of mitochondrial
DNA mutations. *Int J Biochem Cell Biol* **63**: 21-4

7.   Katajisto P, Dohla J, Chaffer CL, Pentinmikko N, Marjanovic N, Iqbal S, Zoncu R, Chen
W, Weinberg RA, Sabatini DM (2015) Asymmetric apportioning of aged mitochondria
between daughter cells is required for stemness. *Science* **348**: 340-343

8.   Soong NW, Hinton DR, Cortopassi G, Arnheim N (1992) Mosaicism for a Specific
Somatic Mitochondrial-DNA Mutation in Adult Human Brain. *Nature Genetics* **2**: 318-323

9.   Ahier A, Dai C-Y, Tweedie A, Bezawork-Geleta A, Kirmes I, Zuryn S (2018) Affinity
purification of cell-specific mitochondria from whole animals resolves patterns of genetic
mosaicism. *Nat Cell Biol* **20**: 352-360

10.   Liau WS, Gonzalez-Serricchio AS, Deshommes C, Chin K, LaMunyon CW (2007) A
persistent mitochondrial deletion reduces fitness and sperm performance in heteroplasmic
populations of C. elegans. *BMC Genet* **8**: 8

11.   Lin YF, Schulz AM, Pellegrino MW, Lu Y, Shaham S, Haynes CM (2016) Maintenance
and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded
protein response. *Nature* **533**: 416-9

12.   Gitschlag BL, Kirby CS, Samuels DC, Gangula RD, Mallal SA, Patel MR (2016)
Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in C. elegans. *Cell*
Metab **24**: 91-103

13.   Tsang WY, Lemire BD (2002) Stable heteroplasmy but differential inheritance of a
large mitochondrial DNA deletion in nematodes. *Biochem Cell Biol* **80**: 645-54

14.   Springer W, Hoppe T, Schmidt E, Baumeister R (2005) A Caenorhabditis elegans
Parkin mutant with altered solubility couples alpha-synuclein aggregation to proteotoxic
stress. *Hum Mol Genet* **14**: 3407-23

15.   Cummins N, Tweedie A, Zuryn S, Bertran-Gonzalez J, Götz J (2019) Disease-
associated tau impairs mitophagy by inhibiting Parkin translocation to mitochondria. *EMBO
J* **38**

16.   Palikaras K, Lionaki E, Tavernarakis N (2015) Coordination of mitophagy and
mitochondrial biogenesis during ageing in C. elegans. *Nature* **521**: 525-8

17.   Kirienko NV, Ausubel FM, Ruvkun G (2015) Mitophagy confers resistance to
siderophore-mediated killing by Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **112**:
1821-6

18.   Ryu D, Mouchiroud L, Andreux PA, Katsyuba E, Moullan N, Nicolet-Dit-Felix AA,
Williams EG, Jha P, Lo Sasso G, Huzard D, *et al.* (2016) Urolithin A induces mitophagy and
prolongs lifespan in C. elegans and increases muscle function in rodents. *Nat Med* **22**: 879-
88

19.   Palikaras K, Lionaki E, Tavernarakis N (2018) Mechanisms of mitophagy in cellular
homeostasis, physiology and pathology. *Nat Cell Biol* **20**: 1013-1022

20.   Harper JW, Ordureau A, Heo JM (2018) Building and decoding ubiquitin chains for
mitophagy. *Nat Rev Mol Cell Biol* **19**: 93-108

21.   Park J, Lee SB, Lee S, Kim Y, Song S, Kim S, Bae E, Kim J, Shong MH, Kim JM, *et al.*
(2006) Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin.
Nature **441**: 1157-1161

22.   Samann J, Hegermann J, von Gromoff E, Eimer S, Baumeister R, Schmidt E (2009)
Caenorhabdititis elegans LRK-1 and PINK-1 act antagonistically in stress response and neurite
outgrowth. *J Biol Chem* **284**: 16482-91

23.   Palikaras K, Tavernarakis N (2017) Assessing Mitochondrial Selective Autophagy in
the Nematode Caenorhabditis elegans. *Methods Mol Biol* **1567**: 349-361

24.   Lapierre LR, De Magalhaes Filho CD, McQuary PR, Chu CC, Visvikis O, Chang JT,
Gelino S, Ong B, Davis AE, Irazoqui JE, *et al.* (2013) The TFEB orthologue HLH-30 regulates
autophagy and modulates longevity in Caenorhabditis elegans. *Nat Commun* **4**: 2267

25.   Polanco JC, Li C, Bodea LG, Martinez-Marmol R, Meunier FA, Götz J (2018) Amyloid-
beta and tau complexity - towards improved biomarkers and targeted therapies. *Nat Rev
Neurol* **14**: 22-39

26.   Franco-Iborra S, Plaza-Zabala A, Sebastian D, Vila M, Martinez-Vicente M (2019)
Impairment of neuronal mitophagy by mutant huntingtin. *bioRxiv*

27.   Gidalevitz T, Ben-Zvi A, Ho KH, Brignull HR, Morimoto RI (2006) Progressive
disruption of cellular protein folding in models of polyglutamine diseases. *Science* **311**:
1471-4

28.   Ashrafi G, Schlehe JS, LaVoie MJ, Schwarz TL (2014) Mitophagy of damaged
mitochondria occurs locally in distal neuronal axons and requires PINK1 and Parkin. *J Cell
Biol* **206**: 655-70

29.   Bingol B, Tea JS, Phu L, Reichelt M, Bakalarski CE, Song Q, Foreman O, Kirkpatrick DS,
Sheng M (2014) The mitochondrial deubiquitinase USP30 opposes parkin-mediated
mitophagy. *Nature* **510**: 370-5

30.   Trifunovic A, Wredenberg A, Falkenberg M, Spelbrink JN, Rovio AT, Bruder CE,
Bohlooly YM, Gidlof S, Oldfors A, Wibom R, *et al.* (2004) Premature ageing in mice
expressing defective mitochondrial DNA polymerase. *Nature* **429**: 417-23

31.   Kujoth GC, Hiona A, Pugh TD, Someya S, Panzer K, Wohlgemuth SE, Hofer T, Seo AY,
Sullivan R, Jobling WA, *et al.* (2005) Mitochondrial DNA mutations, oxidative stress, and
apoptosis in mammalian aging. *Science* **309**: 481-4

32.   Pickrell AM, Huang CH, Kennedy SR, Ordureau A, Sideris DP, Hoekstra JG, Harper JW,
Youle RJ (2015) Endogenous Parkin Preserves Dopaminergic Substantia Nigral Neurons
following Mitochondrial DNA Mutagenic Stress. *Neuron* **87**: 371-81

33.   Valente EM, Abou-Sleiman PM, Caputo V, Muqit MM, Harvey K, Gispert S, Ali Z, Del
Turco D, Bentivoglio AR, Healy DG, *et al.* (2004) Hereditary early-onset Parkinson's disease
caused by mutations in PINK1. *Science* **304**: 1158-60

34.   Kitada T, Asakawa S, Hattori N, Matsumine H, Yamamura Y, Minoshima S, Yokochi M,
Mizuno Y, Shimizu N (1998) Mutations in the parkin gene cause autosomal recessive juvenile
parkinsonism. *Nature* **392**: 605-8

35.   Kraytsberg Y, Kudryavtseva E, McKee AC, Geula C, Kowall NW, Khrapko K (2006)
Mitochondrial DNA deletions are abundant and cause functional impairment in aged human
substantia nigra neurons. *Nat Genet* **38**: 518-20

36.   Bender A, Krishnan KJ, Morris CM, Taylor GA, Reeve AK, Perry RH, Jaros E, Hersheson
JS, Betts J, Klopstock T, *et al.* (2006) High levels of mitochondrial DNA deletions in substantia
nigra neurons in aging and Parkinson disease. *Nat Genet* **38**: 515-7

37.   Brenner S (1974) The genetics of Caenorhabditis elegans. *Genetics* **77**: 71-94

38.   Valmas N, Zuryn S, Ebert PR (2008) Mitochondrial uncouplers act synergistically with
the fumigant phosphine to disrupt mitochondrial membrane potential and cause cell death.
Toxicology **252**: 33-39

39.   Zuryn S, Kuang J, Ebert P (2008) Mitochondrial modulation of phosphine toxicity and
resistance in Caenorhabditis elegans. *Toxicological Sciences* **102**: 179-186

40.   Benedetti C, Haynes CM, Yang Y, Harding HP, Ron D (2006) Ubiquitin-like protein 5
positively regulates chaperone gene expression in the mitochondrial unfolded protein
response. *Genetics* **174**: 229-39

41.   Neumann B, Coakley S, Giordano-Santini R, Linton C, Lee ES, Nakagawa A, Xue D,
Hilliard MA (2015) EFF-1-mediated regenerative axonal fusion requires components of the
apoptotic pathway. *Nature* **517**: 219-22
555

Figure Legends

Figure 1. P*ink-1* and *pdr-1* are required for establishing differences in ΔmtDNA heteroplasmy between major tissue types. (**A**) Overview of mitochondrial immunoprecipitation from specific tissues in *pink-1(tm1779);pdr-1(gk488)* double mutants harbouring ΔmtDNA. (**B-F**) Relative heteroplasmy differences in mitochondria isolated from each cell type in wild type and *pink-1(tm1779);pdr-1(gk488)* backgrounds. Bars are means ± s.e.m. of three independent biological replicates (dots) consisting of ~10,000 animals each. * = $p < 0.05$; ** = $p < 0.01$. *P* values were determined using one-way ANOVA and Tukey's correction. Comparison of heteroplasmy between tissues in (**G**) wild-type animals and (**H**) *pink-1(tm1779);pdr-1(gk488)* double mutants indicates that inter-tissue differences in ΔmtDNA heteroplasmy is abolished by removing *pink1* and *pdr-1*. (**I**) Graphical representation of heteroplasmy divergence (relative to body wall muscle) between tissues in each background. The significance of heteroplasmy variation between tissues was calculated using one-way ANOVA. ** = $p < 0.01$; *** = $p < 0.001$. *P* values were determined using one-way ANOVA and Tukey's correction. (**J**) Quantitative PCR analysis of mtDNA copy number in L4 wild type and *pink-1(tm1779);pdr-1(gk488)* double mutants. Columns are means ± s.e.m. of three independent biological replicates (dots), each consisting of 10 animals, analysed with Student's t-test.

Figure 2. *Pink-1* and *pdr-1* are required for mitophagy in body wall muscle cells. (**A**) PINK-1 translational reporter is expressed in most somatic tissues in *C.*

elegans, including the body wall muscle. White box is magnified in the panels on the

left and body muscle cells are outlined in white dashed lines. White scale bar, 50μm;

red scale bar, 5μm. (**B**) Quantification of mitophagy in body wall muscle cells.

Columns are means ± s.e.m. of independent animals (grey and black dots). ** = *p*<

0.01. *P* values were determined using Student's t-test. (**C**) Representative images

and three-dimensional reconstructions of the *C. elegans* body wall muscle

mitochondrial network. These representative figures have been constructed from

high magnification (40x) z-stack images of live animals stitched together and three-

dimensionally reconstructed. (**D**) Quantification of body wall muscle mitochondrial

volume. Columns are means ± s.e.m. of individual animals (black dots). ** = *p*< 0.01,

P values were determined using Student's t-test.

Figure 3. ΔmtDNA induces mitochondrial dysfunction and mitophagy in

muscle cells and neurons. (**A**) Representative micrographs taken of live animals

expressing tissue-specific GFP reporters (*myo-3*p::GFP^{mt}, body wall muscle and

*mec-4*p::^{MTS}GFP, mechanosensory neurons) localised to the mitochondria and

stained with the mitochondrial membrane potential sensitive dye, TMRE (pseudo-

coloured magenta). Scale bar, 10μm. (**B**) Quantification of mean TMRE intensity

from mitochondria of each tissue type. a.u., arbitrary units. Columns are means ±

s.e.m., each dot is the mean TMRE intensity calculated from an independent animal.

** = *p*< 0.01. *P* values were determined using Student's t-test. (**C**) Quantification of

mtDNA molecules per mitochondrion. Each dot represents an independent

experiment in which the average number of mtDNA molecules per mitochondrion (n

> 1,000 mitochondria per sample) were calculated. Bars are means of three

biologically independent experiments ± s.e.m. ** = *p*< 0.01; *** = *p*< 0.001. *P* values

were determined using one-way ANOVA and Tukey's correction. (**D**) Quantification

of mitophagy in body wall muscle cells and neurons. Columns are means ± s.e.m.,

each dot represents an independent animal analysed. * = $p < 0.05$; ** = $p < 0.01$; *** =

$p < 0.001$; **** = $p < 0.0001$. *P* values were determined using one-way ANOVA and

Tukey's correction.

Figure 4. Proteotoxicity increases ΔmtDNA heteroplasmy in neurons. (**A**)

Overview of the genetic strains used to isolate mitochondria from neurons

overexpressing proteotoxic species. (**B**) Expression of either human tau (hTau) or a

polyglutamine tract (Q40) in neurons increases ΔmtDNA levels in the nervous

system. Bars are means ± s.e.m. of at least three independent biological replicates

(dots). *** = $p < 0.001$. *P* values were determined using one-way ANOVA and

Tukey's correction. (**C**) Proposed model.

Supplemental Figure Legends

Figure S1. ΔmtDNA heteroplasmy and genetic overview. (**A**) Quantification of

uaDf5 heteroplasmy in whole animal extracts. Heteroplasmy was stable at around

50-60% for each of the strains used. Bar represents the mean value of three

independent experiments ± s.e.m. (**B**) Schematic of *C. elegans* mtDNA showing the

location of the 3.1kb *uadF5* deletion.

Figure S2. Perturbing mitophagy increases mitochondrial volume in body wall

muscle cells. (**A**) Representative microimages of body wall muscle mitochondria.

White scale bar, 50μm; red scale bar, 5μm. All images are stitched together from

multiple high magnification (40x) z-stack images of live animals (see materials and

methods).   (**B**) Quantification of mitochondrial. Columns are means $\pm$ s.e.m. of

individual animals (black dots). ** = *p*< 0.01. *P* values were determined using

Student's t-test.

Figure S3. ΔmtDNA does not alter mitochondrial morphology or volume in

body wall muscle cells and neurons. Representative microimages of body wall

muscle (**A**) and (**D**) motor neuron mitochondria. White scale bar, 50µm; red scale

bar, 5µm. All images are stitched together from multiple high magnification (40x) z-

stack images of live animals (see materials and methods).   Quantification of

mitochondrial volumes from (**B-C**) body wall muscle cells and (**E-F**) motor neurons.

Columns are means $\pm$ s.e.m. of individual animals (black dots).

A

pink-1(tm1779); pdr-1(gk488)

○ Ubiquitous (*myo-3p::TOMM-20::mKate2::HA*)
○ Neurons (*rgef-1p::TOMM-20::mKate2::HA*)
○ Intestine (*ges-1p::TOMM-20::mKate2::HA*)
○ Epidermis (*dpy-7p::TOMM-20::mKate2::HA*)
○ Body wall muscles (*myo-3p::TOMM-20::mKate2::HA*)

○ Wild-type mtDNA
○ ΔmtDNA (*uaDf5*)

αHA   IP

Neuron mitochondria

B **Ubiquitous**

ΔmtDNA(uaDf5) % relative to input

ns

wild type
pink-1(tm1779); pdr-1(gk488)

C **Neurons**

ΔmtDNA(uaDf5) % relative to input

**

wild type
pink-1(tm1779); pdr-1(gk488)

D **Intestine**

ΔmtDNA(uaDf5) % relative to input

*

wild type
pink-1(tm1779); pdr-1(gk488)

E **Epidermis**

ΔmtDNA(uaDf5) % relative to input

**

wild type
pink-1(tm1779); pdr-1(gk488)

F **Body wall muscles**

ΔmtDNA(uaDf5) % relative to input

ns

wild type
pink-1(tm1779); pdr-1(gk488)

G **Wild type**

ΔmtDNA(uaDf5) % relative to input

***
**

neurons
intestine
epidermis
body wall muscles

H **pink-1(tm1779); pdr-1(gk448)**

ΔmtDNA(uaDf5) % relative to input

ns

neurons
intestine
epidermis
body wall muscles

I **Heteroplasmy divergence(%)**

0   2   4   6   8

Wild type

p=0.002

pink-1(tm1779); pdr-1(gk448)

ns

J **mtDNA**

copies of mtDNA/nuDNA

ns

wild type
pink-1(tm1779); pdr-1(gk48)

A *pink-1*p::PINK-1::GFP

DIC   GFP   Merge

B

mitoautophagosomes/
μm³ mitochondria x10⁻⁴

$$\text{mitoautophagosomes/} \mu m^3 \text{ mitochondria} \times 10^{-4}$$

**

wild type

*pink-1(tm1779);
pdr-1(gk488)*

*myo-3p::TOMM-20::mKate2::HA;
lgg-1p::LGG-1::GFP*

C myo-3p::TOMM-20::mKate2::HA

Wild type   *pink-1(tm1779);pdr-1(gk48)*

ΔmtDNA   ΔmtDNA

ΔmtDNA

pink-1(tm1779);pdr-1(gk448);ΔmtDNA

ΔmtDNA   *pink-1(tm1779);pdr-1(gk448);
ΔmtDNA*

20 μm

D

Volume (μm³)

**

wild type

*pink-1(tm1779);
pdr-1(gk448)*

myo-3p::TOMM-20::mKate2::HA

A

aex-3p::hTau
rgef-1p::Q40::YFP

Wild-type mtDNA
ΔmtDNA (*uaDf5*)

αHA

rgef-1p::TOMM-20::mKate2::HA

B

neuron-specific
mitochondria

ΔmtDNA(uaDf5) %
relative to input

7.5
5.0
2.5
0.0
-2.5
-5.0
-7.5
-10.0

*** ***

wild type
aex-3p::hTau
rgef-1p::Q40::YFP

C

PINK-1-PDR-1-
mediated
mitophagy

Proteotoxicity
hTau
polyQ40

Less mitophagy
per mitochondria

More mitophagy
per mitochondria

Divergence in heteroplasmy
between tissues

Tissue-type:
Muscle Neurons

mutant
mtDNA %

Time

Muscle Neurons

Time

Muscle Neurons

Epidermis

Body wall
muscles

Neurons

Intestine