1 **Iron-sulfur cluster repair contributes to *Y. pseudotuberculosis* survival within deep tissues.**

4 Running title: Iron-sulfur cluster repair during *Y. pseudotuberculosis* infection

8 Kimberly M. Davis[1,#a,*], Joanna Krupp[1], Stacie Clark[1,2], Ralph R. Isberg[1]

9
10
11
12 [1]Department of Molecular Biology and Microbiology
13 [2]Graduate Program in Molecular Microbiology, Sackler School of Graduate Biomedical Science
14 Tufts University School of Medicine, Boston, MA, USA
15 [#a]Current Address: W. Harry Feinstone Department of Molecular Microbiology and Immunology
16 Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

22 *Corresponding author:

23 E-mail: kdavi140@jhu.edu (KMD)
24

Abstract

To successfully colonize host tissues, bacteria must respond to and detoxify many different host-

derived antimicrobial compounds, such as nitric oxide (NO). NO can directly kill bacteria,

primarily through attack on iron-sulfur (Fe-S) cluster-containing proteins. NO detoxification

plays an important role in promoting bacterial survival, but it remains unclear if repair of Fe-S

clusters is also important for bacterial survival within host tissues. Here we show that the Fe-S

cluster repair protein, YtfE, contributes to the survival of *Y. pseudotuberculosis* within the spleen

following nitrosative stress. *Y. pseudotuberculosis* forms clustered centers of replicating bacteria

within deep tissues, where peripheral bacteria express the NO-detoxifying gene, *hmp*. *ytfE*

expression also occurred specifically within peripheral cells at the edges of microcolonies. In the

absence of *ytfE*, the area of microcolonies was significantly smaller than WT, consistent with

ytfE contributing to the survival of peripheral cells. The loss of *ytfE* did not alter the ability of

cells to detoxify NO, which occurred within peripheral cells in both WT and Δ*ytfE*

microcolonies. In the absence of NO-detoxifying activity by *hmp*, NO diffused across Δ*ytfE*

microcolonies, and there was a significant decrease in the area of microcolonies lacking *ytfE*,

indicating that *ytfE* also contributes to bacterial survival in the absence of NO detoxification.

These results indicate a role for Fe-S cluster repair in the survival of *Y. pseudotuberculosis*

within the spleen, and suggest that extracellular bacteria may rely on this pathway for survival

within host tissues.

Introduction

Nitric oxide (NO) is a diffusible gas that has a wide range of physiological functions within mammals (1, 2). The effects of NO are tissue concentration-dependent, as it promotes vasodilation, cell proliferation and cell differentiation at low concentrations (3, 4), while high concentrations drive apoptosis and defense against bacterial, fungi, and parasites (5-8). NO is produced by three different nitric oxide synthase (NOS) isoforms within mammalian tissues: the neuronal NOS (nNOS), endothelial NOS (eNOS), and inducible NOS (iNOS). nNOS and eNOS are expressed at low levels by endothelial and neuronal cells respectively (2). In contrast, iNOS is expressed by a wide range of cell types, specifically in response to NF-κB-dependent sensing of, and is responsible for the high levels of NO produced during infection (1, 9).

NO has direct bactericidal activity and can also react with reactive oxygen species to produce additional toxic compounds. NO antibacterial activity occurs primarily through nitrosylation of iron-sulfur (Fe-S) cluster-containing proteins, which play critical roles in cellular respiration, DNA synthesis, and gene regulation. One of the global regulators in *E. coli* that has been shown to be inactivated by NO is NsrR (10), which regulates the response to nitrosative stress and is also associated with the oxidative stress response (11-13). Nitrosylation of the NsrR-associated Fe-S cluster relieves repression of at least 60 genes in *E. coli* (14-17). Included in this regulon is the *hmp* gene, which encodes a flavohemoglobin that detoxifies NO (18, 19), and YtfE (also known as Repair of Iron Centers (RIC) in *E. coli*), which functions to repair Fe-S clusters following NO damage (12, 16). Repair of Fe-S cluster proteins by YtfE can eliminate the need for new Fe-S cluster biogenesis when Fe availability is limited (12, 20, 21). A variety of studies argue that YtfE contributes to bacterial survival within host cells, but it remains unclear if

YtfE contributes to the survival of extracellular pathogens replicating within host tissues, and if

YtfE contributes to survival following exposure to NO (22-24).

Yersinia pseudotuberculosis is an oral pathogen that is typically contained within

intestinal tissues and gut-associated lymphoid tissues, but has the capacity to spread systemically

in susceptible individuals (25-28). Following bloodstream access, *Y. pseudotuberculosis*

colonizes deep tissue sites where individual bacteria replicate to form clonal microcolonies (29-

31). Neutrophils, monocytes, and macrophages are recruited to sites of bacterial replication,

which are kept at bay by *Yersinia*, primarily via substrates of the bacterial type III secretion

system (32-34). Recruited monocytes and macrophages produce NO, which diffuses across a

layer of neutrophils and is inactivated at the periphery of the microcolony by bacteria expressing

Hmp, preventing diffusion of NO into the interior of the microcolony (31). The consequences of

the selective attack of NO on the peripheral bacteria are unclear, although it is likely that

nitrosative stress may slow their growth. Additional members of the nitrosative stress response,

such as YtfE, may also be required in peripheral cells to ensure their survival.

Bacteria responding to RNS appear to recover and remain viable, consistent with

members of the NsrR regulon cooperating to repair NO-mediated damage in peripheral bacteria

within microcolonies. Two of the members of the *E. coli* NsrR regulon, *hmp* and *ytfE,* are

upregulated in the bubonic plague model of *Yersinia pestis* infection (35), so *ytfE* is likely a

member of the NsrR regulon in *Yersinia* and could contribute to microbial fitness during

microcolony growth. Additionally, very few studies have explored the role of Fe-S cluster repair

during infection with extracellular bacteria. Here, we show that *ytfE* contributes to the survival

of extracellular bacteria, specifically through upregulation of *ytfE* within peripheral cells of *Y.*

pseudotuberculosis microcolonies.

Results.

***Y. pseudotuberculosis ytfE* expression is regulated by NsrR and occurs within peripheral**

cells during growth in the spleen.

The *ytfE* gene is known to be a member of the NsrR regulon in a number of bacterial

species (12, 13, 36), so *ytfE* is expected to be repressed by NsrR in *Y. pseudotuberculosis*. We

also expected that *ytfE* is transcribed during *Y. pseudotuberculosis* growth within the mouse

spleen as dthe NsrR-regulated *hmp* gene is expressed in this tissue (22-24, 31, 35). To determine

if *Y. pseudotuberculosis ytfE* is expressed during splenic growth, C57BL/6 mice were

intravenously challenged with bacteria and bacterial RNA was isolated at day 3 post-inoculation

(PI). Based on qRT-PCR analysis, *ytfE* transcript levels increased within the mouse spleen

relative to the inoculum culture grown in the absence of an NO-generating system, with marked

mouse-to-mouse variation (Figure 1A) (31).   To determine if *ytfE* expression is NsrR-dependent,

we compared *ytfE* transcription levels in WT and Δ*nsrR* strains in the presence (+) and absence (-)

of nitrogen stress imparted by acidified nitrite (NO_2). Transcription of *ytfE* increased by over

100X in the WT strain with the addition of nitrogen stress (Figure 1B). As expected, transcription

of *ytfE* was high in the Δ*nsrR* strain in either the presence or absence of nitrogen stress, indicating

that NsrR negatively regulates *ytfE* transcription.

Hmp, another member of the NsrR regulon, is specifically expressed on the periphery of

Y. pseudotuberculosis microcolonies during splenic growth (31). At the transcriptional level,

this is associated with considerable variation in expression levels between individual mice. The

variable *ytfE* expression levels in the mouse and NsrR-dependence is reminiscent of *hmp*

expression, indicating that there could be a link between spatial expression and inter-mouse

variation. To determine if *ytfE* and *hmp* have similar expression patterns during growth in the

spleen, mice were intravenously inoculated with a WT *Y. pseudotuberculosis* strain containing

hmp::mCherry (chromosomal integration of *mCherry* downstream of *hmp*) and *ytfE::gfp*

(chromosomal integration of *gfp* downstream of *ytfE*). Microcolonies were visualized within the

spleen using fluorescence microscopy, and *ytfE* and *hmp* reporter signals were quantified within

the same cells at the center and periphery of the microcolonies (Materials and Methods). The

hmp reporter signal increased within individual cells at the periphery relative to cells at the

centroid of microcolonies, which generated a ratio value greater than 1, consistent with *hmp*

peripheral expression (Figure 1C). The *ytfE* signal was dim relative to *hmp*, but also increased at

the periphery of microcolonies relative to the centroid. The Periphery/Centroid signal intensity

ratio value for *ytfE* was similar to *hmp*, indicating that *ytfE* was expressed at the periphery of

microcolonies (Figure 1C). There was significant overlap in the *hmp* and *ytfE* signals within

these images, and based on NsrR-dependent regulation of both genes, it is likely that *hmp* and

ytfE were expressed in the same cells (Figure 1D).

ytfE contributes to the virulence of _Y. pseudotuberculosis_ in the spleen.

YtfE repairs Fe-S clusters damaged by NO, and there appears to be sufficient NO at the

periphery of splenic microcolonies to allow synthesis of this protein and to promote repair within

this subpopulation of bacteria. To determine if loss of *yftE* alters the overall fitness of *Y.*

pseudotuberculosis during growth within the spleen, we constructed a *Y. pseudotuberculosis*

strain that lacks the *ytfE* gene (Δ*ytfE*) and harbors a constitutive *gfp*-expressing plasmid, to

visualize growth within the spleen. To compare differences in relative fitness, mice were

infected intravenously with equal amounts of WT mCherry$^+$ and Δ*ytfE* GFP$^+$ strains, and spleens

were harvested at day 3 PI to determine the competitive index by colony forming units (CFUs)

and quantify microcolony areas within the same animals. At day 3 PI, the median competitive

index was less than 1, indicating lowered fitness of the ΔytfE strain relative to the WT strain

(Figure 2A). The areas of individual microcolonies within these tissues were also visualized and

quantified by fluorescence microscopy. In addition, the ΔytfE microcolonies were significantly

smaller than WT microcolonies within the same organs, indicating that ytfE contributes to the

survival of *Y. pseudotuberculosis* in the spleen, presumably due to lowered fitness of the

bacterial population located at the periphery of the microcolonies (Figure 2B).

Since YtfE could directly repair the Fe-S cluster of NsrR, the absence of ytfE could alter

expression of the NsrR regulon, by resulting in heightened expression within peripheral cells. To

determine if microcolonies from the *Y. pseudotuberculosis* ΔytfE strain have sustained

expression of the NsrR regulon relative to the WT strain, we infected mice intravenously with

WT GFP$^+$ *hmp::mCherry* or ΔytfE GFP$^+$ *hmp::mCherry* integrated *hmp* reporter strains, and

spleens were harvested at day 3 PI to visualize reporter expression by fluorescence microscopy.

Similar to Fig. 1D, *hmp* reporter expression was significantly higher at the periphery relative to

the centroid in the WT strain (Figure 2C). The reporter expression pattern was very similar in

ΔytfE microcolonies, indicating that there was still a gradient of NO exposure in the mutant, in

which Hmp activity in the peripheral population protects the central population of bacteria, and

that loss of *ytfE* did not significantly alter expression of the NsrR regulon (Figure 2D).

Loss of *ytfE* and *hmp* reduces *Y. pseudotuberculosis* fitness.

YftE contributed to the growth of *Y. pseudotuberculosis* microcolonies in the spleen

despite expression limited to the microcolony periphery. We were then interested in determining

if YtfE-mediated repair played an important role in the context of a Δ*hmp* strain, where all

bacteria in a microcolony are exposed to NO (31). We first compared the virulence of WT and

Δ*hmp* Δ*ytfE* microcolonies during single strain infection. Mice were infected intravenously with

WT GFP$^+$ or Δ*hmp* Δ*ytfE* GFP$^+$ strains, and spleens were harvested at day 3 PI to visualize

microcolony areas by fluorescence microscopy. Individual Δ*hmp* Δ*ytfE* microcolonies were

significantly smaller than WT, consistent with impaired growth of the Δ*hmp* Δ*ytfE* strain (Figure

3A). We also determined the relative fitness of the WT and Δ*hmp* Δ*ytfE* strains by comparing

growth within the same tissue during co-infection. Mice were infected intravenously with WT

mCherry$^+$ and Δ*hmp* Δ*ytfE* GFP$^+$ strains, and spleens were harvested at day 3 PI to quantify

CFUs and visualize microcolony areas by fluorescence microscopy. The median competitive

index value was below 1, indicating that the Δ*hmp* Δ*ytfE* strain had reduced fitness relative to the

WT strain (Figure 3B). The microcolony areas of the Δ*hmp* Δ*ytfE* were also significantly smaller

than WT microcolonies, indicating the Δ*hmp* Δ*ytfE* had reduced fitness in the spleen relative to

the WT strain (Figure 3C).

***ytfE* contributes to bacterial survival in the absence of *hmp*.**

As the fitness of the Δ*ytfE* or Δ*hmp* Δ*ytfE* strains showed similar defects when compared

to WT (Figure 3), it is not clear if YftE contributes to *Y. pseudotuberculosis* colonization of the

spleen in the absence of Hmp function. To address this point, we challenged mice intravenously

with Δ*hmp* GFP$^+$ P$_{hmp}$::*mCherry* and Δ*hmp* Δ*ytfE* GFP$^+$ P$_{hmp}$::*mCherry* strains, and spleens were

harvested at day 3 PI to quantify CFUs, visualize microcolony areas, and detect reporter signals

by fluorescence microscopy. The median competitive index (CI) value for the double mutant

was below 1, indicating that Δ*hmp* Δ*ytfE* strain was less fit than Δ*hmp* (Figure 4A).

Interestingly, in 4 mice the CI was at least 1, indicating that the Δ*hmp* Δ*ytfE* strain could compete

efficiently with Δ*hmp* in these animals. We then compared the total CFU in the organs in mice

in which the Δ*hmp* strain outcompeted Δ*hmp* Δ*ytfE*, or in mice in which no such outcompetition

took place (Figure 4B). The total CFUs were significantly lower in organs in which the Δ*hmp*

did not outcompete the $\Delta hmp\Delta ytfE$ strain, indicating that the fitness differences were suppressed

in animals showing increased restriction of bacterial growth. The microcolony areas were

quantified within all the spleens depicted in Figure 4A, and the areas of Δhmp $\Delta ytfE$

microcolonies were significantly smaller than Δhmp (Figure 4C). Δhmp $\Delta ytfE$ microcolonies had

similar P_{hmp} reporter expression at the centroid and periphery, indicating NO diffused across

these centers (Figure 4D). Together, these results confirm that $ytfE$ contributes to bacterial

survival in the absence of Hmp detoxifying activity.

$ytfE$ has limited effects on NO sensitivity in the absence of hmp

The $\Delta ytfE$ single mutant strain had reduced fitness relative to the WT strain (Figure 2),

however the single $\Delta ytfE$ mutant was not significantly more sensitive than the WT strain to the

presence of acidified nitrite during growth in culture.:. Similarly, despite the lowered fitness of

the Δhmp $\Delta ytfE$ strain in spleens relative to Δhmp, the Δhmp $\Delta ytfE$ strain was not significantly

more sensitive than the Δhmp mutant strain to the presence of acidified nitrite during growth in

culture (Figure 5A; compare $+NO_2$ samples). This is consistent with previous reports in other

organisms (36, 37), perhaps due to the presence of other backup repair pathways that are active

in the absence of Hmp or because YftE plays a role in protection from other stress species.

We then compared hmp reporter expression in $\Delta ytfE$ and $\Delta hmp\Delta ytfE$ strains to confirm

that NO diffusion occurred across Δhmp $\Delta ytfE$ microcolonies using plasmid-borne reporters.

Mice were infected intravenously with $\Delta ytfE$ GFP$^+$ $P_{hmp}::mCherry$ or Δhmp $\Delta ytfE$ GFP$^+$

$P_{hmp}::mCherry$ strains, and spleens were harvested 3 days PI to visualize reporter expression by

fluorescence microscopy. The $\Delta ytfE$ strain had peripheral P_{hmp} reporter expression, as seen with

the chromosomally-integrated hmp reporter, indicating that NO diffusion across the microcolony

is inhibited by peripheral cells in $\Delta ytfE$ microcolonies (Figure 5B, 5C). Δhmp $\Delta ytfE$

microcolonies showed no such preference for the periphery, indicating NO diffused across these

centers, as expected based on the loss of *hmp*.

Rescue of *ytfE* restores microcolony size.

To show that the loss of *ytfE* was responsible for decreased microcolony size, we rescued

the Δ*ytfE* strain with a WT copy of *ytfE*, and transformed the *ytfE* rescued strain with the

constitutive *gfp*-expressing plasmid. Mice were infected intravenously with WT mCherry[+] and

ytfE rescued GFP[+] strains, and spleens were harvested at day 3 PI to quantify CFUs and visualize

microcolony areas by fluorescence microscopy. The median competitive index for the *ytfE*

rescued strain was close to a value of 1, indicating the fitness of this strain was roughly

equivalent to the WT strain (Figure 6A). The microcolony areas were quantified within all the

spleens depicted in Figure 6A, and the areas of *ytfE* rescued microcolonies were very similar to

WT microcolonies (Figure 6B, 6C). These results indicate that the Δ*ytfE* strain was rescued by

the WT copy of *ytfE*, which confirms that the reduced fitness of the Δ*ytfE* strain was specifically

due to a loss of *ytfE*.

Discussion

The detoxification of NO and other reactive nitrogen species is critical for bacterial

survival within host tissues (38-41). Bacterial proteins involved in NO detoxification, however,

are not synthesized until NO accumulates and damages the Fe-S clusters of NsrR, resulting in

inactivation of this repressor (14, 15). Many additional Fe-S cluster-containing proteins are

simultaneously damaged, so bacteria need to either repair damaged proteins or synthesize

replacement proteins, while simultaneously replenishing proteins that detoxify NO and prevent

further damage (16, 36). Although Fe-S cluster repair is likely required for bacterial survival, it

has been unclear whether or not this plays an important role within host tissues. We have chosen

to ask this question in a mouse model of *Y. pseudotuberculosis* infection, where it is known that

bacteria respond to RNS and that Fe concentrations are limiting (23, 24, 31).   Here, we have

shown that Fe-S cluster repair contributes to successful *Y. pseudotuberculosis* replication within

the spleen.

We found that a place where Fe-S cluster repair in *Y. pseudotuberculosis* likely occurs is

in the peripheral subpopulation of bacteria that responds to NO assault within microcolonies.

This is consistent with a role for YftE in supporting survival of the peripheral subpopulation. In

the absence of *ytfE*, we would expect bacteria on the periphery to be exposed to stress associated

with NO exposure, leading to sequential loss of the peripheral population and progressively

smaller microcolonies. Consistent with this hypothesis, we observed Δ*ytfE* microcolonies were

smaller than those established by the WT strain. It is expected that the difference between WT

and Δ*ytfE* microcolony areas will become progressively more pronounced as the infection

proceeds, because Δ*ytfE* microcolonies should continuously lose their peripheral subpopulation.

This prediction is based on our previous observation that NO-sensitive microcolonies are

progressively reduced during the course of disease, with elimination of Δ*hmp* bacteria by NO

most pronounced at late timepoints post-inoculation, concurrent with a timepoint in which the

animals inoculated with the WT strain are moribund (PI).

NO alone was not sufficient to limit the growth of Δ*ytfE* bacteria in bacteriological

media, consistent with previous studies that indicated the uropathogenic *E. coli* (UPEC) Δ*ytfE*

strain had reduced intracellular survival within host cells, but was not sensitive to exogenous NO

alone (37). Presumably, the Δ*ytfE* strain is sensitive to other antimicrobial compounds generated

within host tissues, possibly by the generation of a variety of reactive nitrogen species (RNS) as

a consequence of NO reaction with reactive oxygen species (ROS) or other compounds.

Upregulation of *ytfE* following hydrogen peroxide-mediated damage of *Staphylococcus aureus*

indicates that YtfE plays a broad role in repair, instead of just a response to nitrogen stress (42).

An additional regulatory signal for the *yftE* gene is iron limitation, which depresses

synthesis of new Fe-S cluster proteins, thus requiring cellular YftE function (13, 20, 43). The

spleen is expected to be an iron-limiting environment requiring Fe-S cluster repair, but we found

no evidence for NO-independent induction of the *ytfE* gene within the center of microcolonies,

using our fluorescent promoter constructions. This contrasts with data arguing that *Y.*

pseudotuberculosis expresses many Fe acquisition genes during growth in host tissues, consistent

with Fe-limiting conditions within host tissues (24, 44). It is possible that the center of the

microcolony represents a protected environment with low exposure to stresses such as reactive

nitrogen and oxygen species as well as limited damage to Fe-S centers. Additionally, *ytfE*

expression may be induced at very low NO concentrations, as seen in *Salmonella enterica*

serovar Typhimurium (36), but may require severe iron limitation for expression in the absence

of NO.

The loss of YftE function also has the potential to alter metabolite levels due to

disruption of protein functions that are Fe-S center-related. In the presence of RNS, a Δ*ytfE*

strain may have reduced aconitase and fumarase activity due to the role of YtfE in Fe-S cluster

repair specifically for these proteins (45). The Fe-S cluster of the NsrR repressor is also repaired

by YtfE, which could lead to prolonged expression of the NsrR regulon in the presence of NO

(12). Our results indicate that expression of the NsrR regulon is similar in the presence or

absence of *ytfE*, although it remains possible that prolonged expression of the NsrR regulon

could be detected at later timepoints post-inoculation. Future work will investigate these issues,

and determine the interplay between iron regulation, NO-induced damage and repair of critical

Fe-S centers.

Materials and Methods

Bacterial strains & growth conditions. The WT *Y. pseudotuberculosis* strain, YPIII, was

used throughout. For all mouse infection experiments, bacteria were grown overnight to post-

exponential phase in 2xYT broth (LB, with 2x yeast extract and tryptone) at 26° C with rotation.

Exponential phase cultures were sub-cultured 1:100 from overnight cultures, and grown at 26° C

with rotation for an additional 2 hours. Sodium nitrite (2.5mM) was added to LB pH5.5 to

induce the nitrogen stress response.

Generation of ytfE mutant strains. The *hmp* and *nsrR* deletion strains were previously

described (31). Deletion derivative strains were generated for *ytfE* by amplifying the start codon

+ 3 downstream codons, the 3' terminal 3 codons + the stop codon and fusing these fragments to

generate a start + 6 aa + stop construct. Deletion constructions were amplified with 500 base

pairs flanking sequence on each side, cloned into the suicide vector, pSR47S, and transformed

into *Y. pseudotuberculosis*. Sucrose selection was used to select for bacteria that had

incorporated the desired mutation after a second cycle of recombination (30). PCR, sequencing,

and qRT-PCR were used to confirm deletion strains.

Integrated ytfE reporter construction (ytfE^{+} rescue strain). The *ytfE::gfp* reporter was

generated by cloning *gfp* immediately downstream of the *ytfE* gene (in between the *ytfE* stop

codon and terminator sequence) by overlap extension PCR. The *ytfE* start codon was amplified

with 500 base pairs upstream flanking sequence, while the stop codon of *gfp* was amplified with

500 base pairs downstream flanking sequence. This fragment was cloned into the suicide vector,

pSR47S, and transferred by conjugation into WT *Y. pseudotuberculosis hmp::mCherry*

(chromosomally integrated), selecting for kanamycin resistance. For the *ytfE^{+}* rescue strain, a

WT *ytfE* gene product, including 500 base pairs upstream and downstream of *ytfE*, was amplified

from genomic DNA and cloned into pSR47S. This vector was transferred by conjugation into

$\Delta ytfE$ *Y. pseudotuberculosis*, selecting for kanamycin resistance. A second round of

recombination was selected on sucrose-containing media to isolate strains that had recombined

each *ytfE* construct. PCR and sequencing were used to confirm integration of *gfp* or rescue of

the *ytfE* deletion.

Generation of plasmid-based reporter strains. Several of the *Y. pseudotuberculosis*

reporter strains in this study have been previously described: WT GFP[+], WT mCherry[+]

(*yopE::mCherry)*, WT *hmp::mCherry*, WT GFP[+] *hmp::mCherry*, and Δhmp GFP[+] (31). For this

study, GFP[+] strains were constructed by transforming deletion strains with the constitutive GFP

plasmid, which expresses GFP from an unrepressed P_{tet} of pACYC184. The $P_{hmp}::mCherry$ was

also transformed into GFP[+] strains. The $P_{hmp}::mCherry$ transcriptional fusion was previously

constructed by fusing the *hmp* promoter to *mCherry* using overlap extension PCR, and cloned

into the pMMB67EH plasmid (31).

Murine model of systemic infection. Six to 8-week old female C57BL/6 mice were

obtained from Jackson Laboratories (Bar Harbor, ME). All animal studies were approved by the

Institutional Animal Care and Use Committee of Tufts University. Mice were injected

intravenously with 10^3 bacteria for all experiments. For co-infection experiments, mice were

inoculated with 5×10^2 CFU of each strain, for a total of 10^3 CFUs. At the indicated timepoints

post-inoculation (PI) (3 days), spleens were removed and processed.

qRT-PCR to detect bacterial transcripts in broth-grown cultures. Bacterial cells were

grown in broth to $A_{600} = 0.3$ under appropriate stress conditions, pelleted, resuspended in Buffer

RLT (QIAGEN) + ß-mercaptoethanol, and RNA was isolated using the RNeasy kit (QIAGEN)

according to manufacturer's protocol. DNA contamination was eliminated using the DNA-free

kit (Ambion) according to manufacturer's protocol. RNA was reverse transcribed using M-MLV

reverse transcriptase (Invitrogen), in the presence of the RNase inhibitor, RnaseOUT

(Invitrogen), according to manufacturer's protocol. Approximately 30 ng cDNA was used as a

template in reactions with 0.5 µM of forward and reverse primers and SYBR Green (Applied

Biosystems) according to manufacturer's protocol. Control samples were prepared that lacked

M-MLV, to confirm DNA was eliminated from samples and was not amplified by qRT-PCR.

Reactions were carried out using the StepOnePlus Real-Time PCR system, and relative

comparisons were obtained using the $\Delta\Delta C_T$ or $2^{-\Delta Ct}$ method (Applied Biosystems).

qRT-PCR to detect bacterial transcripts from mouse tissues. Mice were inoculated

intravenously with the WT strain, and at day 3 PI spleens were harvested, and immediately

submerged in RNALater solution (QIAGEN). Tissue was homogenized in Buffer RLT + ß-

mercaptoethanol, and RNA was isolated using the RNeasy kit (QIAGEN) according to

manufacturer's protocol. Bacterial RNA was enriched following depletion of host mRNA and

rRNA from total RNA samples, using the MICROB*Enrich* kit (Ambion) according to

manufacturer's protocol. DNA digestion, reverse transcription, and qRT-PCR were performed

as described above.

Fluorescence microscopy. C57BL/6 mice were inoculated intravenously with the *Y.*

pseudotuberculosis WT strain, at day 3 PI spleens were harvested and immediately fixed in 4%

paraformaldehyde in PBS for 3 hours. Tissues were frozen-embedded in Sub Xero freezing

media (Mercedes Medical) and cut by cryostat microtome into 10µm sections. To visualize

reporters, sections were thawed in PBS, stained with Hoechst at a 1:10,000 dilution, washed in

PBS, and coverslips were mounted using ProLong Gold (Life Technologies). Tissue was imaged

with 20x or 63x objectives, using a Zeiss Axio Observer.Z1 (Zeiss) fluorescent microscope with

Colibri.2 LED light source, an Apotome.2 (Zeiss) for optical sectioning, and an ORCA-R^2 digital

CCD camera (Hamamatsu).

Image analysis. Volocity image analysis software was used to quantify microcolony

areas. Image J was used to quantify the signal intensity of each channel at the centroid and

periphery of each microcolony, to generate relative signal intensities of fluorescent reporters.

Thresholding was used to define the area of each microcolony, the centroid was calculated, and

0.01 pixel^2 squares were selected to calculate values at the centroid. Peripheral measurements

depict bacteria in contact with host cells.

Author Contributions. Designed and performed experiments: KMD, JK, SC.

Intellectual/conceptual contribution: KMD, RRI. Analyzed the data: KMD, JK, SC, RRI. Wrote

the paper: KMD, RRI.

Acknowledgments. We thank the members of the Isberg lab, who provided valuable

advice and feedback throughout this project. We also thank the members of the Davis lab, who

provided feedback and suggestions during the final steps of manuscript preparation. The authors

of this manuscript declare no conflicts of interest. This work was supported by NIAID award

R01 AI110684 as well as by an American Cancer Society-Ellison Foundation Postdoctoral

Fellowship (PF-13-360-01-MPC), and a NIAID K22 Career Transition Award (1K22AI123465-

01).

Figure Legends:

Figure 1: *ytfE* **expression is regulated by NsrR and occurs within peripheral cells.** A)

C57BL/6 mice were inoculated intravenously (i.v.) with 10^3 WT *Y. pseudotuberculosis*, and

spleens were harvested at day 3 post-inoculation (PI). Bacterial transcripts were isolated from

splenic tissue, transcript levels were quantified by qRT-PCR relative to 16S, and the fold

increase in *ytfE* transcript levels is shown relative to the inoculum. Each dot represents an

individual mouse. B) Nitrogen stress was induced (NO_2, +) in cultures of WT and Δ*nsrR* strains,

and compared to untreated cultures (NO_2, -). The *ytfE* transcript levels are expressed relative to

16S, fold increase is relative to the average level in untreated WT cultures. Each column: 5

biological replicates, mean and SEM. C) C57BL/6 mice were inoculated i.v. with the WT

hmp::mCherry ytfE::gfp strain, and spleens were harvested at day 3 PI for fluorescence

microscopy. Fluorescent signal intensity was determined within the same peripheral and centroid

cells for *hmp* and *ytfE* reporters, and divided to generate the periphery/centroid ratio (4

mice/group). Dots: individual microcolonies. D) Representative image of *hmp::mCherry* and

ytfE::gfp reporters; merge and single channel images are shown. Statistics: Wilcoxon matched

pairs, **p<0.01, n.s.: not significant.

Figure 2: *ytfE* **contributes to growth in the spleen.** A) Co-infection with the WT and Δ*ytfE*

strains. C57BL/6 mice were inoculated i.v. with an equal mixture of mCherry$^+$ (*yopE::mCherry*)

WT and GFP$^+$ Δ*ytfE* strains, and spleens were harvested at day 3 PI. Competitive index: ratio of

Δ*ytfE*/WT CFUs in spleen at day 3, divided by ratio of Δ*ytfE*/WT CFUs in the inoculum. Dots:

individual mice. Dotted line: represents equal fitness. B) WT and Δ*ytfE* microcolony areas

(μm^2) from the co-infection quantified (Experimental Procedures) in 6 mice. C) C57BL/6 mice

were inoculated i.v. with the WT GFP$^+$ *hmp::mCherry* or Δ*ytfE* GFP$^+$ *hmp::mCherry* strains, and

spleens were harvested at day 3 PI for fluorescence microscopy. Reporter signals were

quantified within peripheral and centroid cells, *hmp* reporter signal was divided by GFP. D)

Representative image of a Δ*ytfE* GFP$^+$ *hmp::mCherry* microcolony, stained with Hoechst to

detect host nuclei; merge and *hmp* single channel images are shown Statistics: B): Mann-

Whitney, C): Wilcoxon matched pairs, ***$p<0.001$.

<u>**Figure 3:**</u> **Δ*hmp* Δ*ytfE* is attenuated for growth in the spleen.** A) C57BL/6 mice inoculated i.v.

with the WT GFP$^+$ or Δ*hmp* Δ*ytfE* GFP$^+$ strains, and spleens were harvested at day 3 PI to

quantify microcolony areas (μm^2) from single strain infections. Dots: individual microcolonies.

B) C57BL/6 mice were inoculated i.v. with an equal mixture of WT mCherry$^+$ (*yopE::mCherry*)

and Δ*hmp* Δ*ytfE* GFP$^+$ strains, and spleens were harvested at day 3 PI. Competitive index: ratio

of Δ*hmp* Δ*ytfE*/WT CFUs in spleen at day 3, divided by ratio of Δ*hmp* Δ*ytfE*/WT CFUs in the

inoculum. Dots: individual mice. Dotted line: represents equal fitness. C) WT and Δ*hmp* Δ*ytfE*

microcolony areas (μm^2) from the co-infection quantified (Experimental Procedures) in 7 mice.

Statistics: Mann-Whitney, *$p<0.05$.

<u>**Figure 4:**</u> **Δ*ytfE* contributes to survival in the absence of *hmp*.** A) C57BL/6 mice were

inoculated i.v. with an equal mixture of *hmp* GFP$^+$ and Δ*hmp* Δ*ytfE* GFP$^+$ P_{hmp}::*mCherry* strains,

and spleens were harvested at day 3 PI. Competitive index: ratio of Δ*hmp* Δ*ytfE*/Δ*hmp* CFUs in

spleen at day 3, divided by ratio of Δ*hmp* Δ*ytfE*/Δ*hmp* CFUs in the inoculum. Dots: individual

mice. Dotted line: represents equal fitness. B) Total CFUs in the spleen during co-infection,

when CI was less than 1 (Δ*hmp* wins) or above or equal to 1 (Δ*hmp* Δ*ytfE* wins). Dots: individual

mice. C) Δhmp and Δhmp $\Delta ytfE$ microcolony areas (μm^2) from the co-infection quantified

(Experimental Procedures) in 10 mice. Dots: individual microcolonies. D) Reporter signals were

quantified within peripheral and centroid cells in the Δhmp $\Delta ytfE$ strain during co-infection, *hmp*

reporter signal was divided by GFP. Dots: individual microcolonies. Statistics: B) and C): Mann-

Whitney, D): Wilcoxon matched pairs, *p<0.05, **p<0.01, n.s.: not significant.

Figure 5: YtfE activity is required in the absence of NO detoxification (Δ*hmp*). A) Optical

density (A_{600nm}) was measured every 2 hours (time, hrs) within cultures of the indicated strains

during incubation in LB pH5.5 in the presence ($+NO_2$) and absence of $NaNO_2$. Data represents

two independent experiments. B) C57BL/6 mice were inoculated i.v. with the $\Delta ytfE$ GFP[+]

P_{hmp}::mCherry or Δhmp $\Delta ytfE$ GFP[+] *P_{hmp}::mCherry* strains, and spleens were harvested at day 3

PI for fluorescence microscopy. Reporter signals were quantified within peripheral and centroid

cells, *hmp* reporter signal was divided by GFP. Dots: individual microcolonies. C)

Representative images of $\Delta ytfE$ and Δhmp $\Delta ytfE$ microcolonies; merge and *hmp* single channel

images are shown. Statistics: Wilcoxon matched pairs, *p<0.05, n.s.: not significant.

Figure 6: Rescue of *ytfE* restores fitness of the Δ*ytfE* strain. A) C57BL/6 mice were

inoculated i.v. with an equal mixture of *ytfE* rescued GFP[+] and WT mCherry[+] strains, and spleens

were harvested at day 3 PI. Competitive index: ratio of *ytfE* rescue/WT CFUs in spleen at day 3,

divided by ratio of *ytfE* rescue/WT CFUs in the inoculum. Dots: individual mice. Dotted line:

represents equal fitness. B) WT and *ytfE* rescue microcolony areas (μm^2) from the co-infection

quantified (Experimental Procedures) in 10 mice. Dots: individual microcolonies. C)

Representative images of WT mCherry[+] and *ytfE* rescued GFP[+] microcolonies. Scale bar: 20μm.

Statistics: Mann-Whitney, n.s.: not significant.

<u>References</u>

1.   Bogdan C. Nitric oxide and the immune response. Nat Immunol. 2001;2(10):907-16.

2.   Nathan C. Nitric oxide as a secretory product of mammalian cells. FASEB J.
1992;6(12):3051-64.

3.   Pervin S, Singh R, Hernandez E, Wu G, Chaudhuri G. Nitric oxide in physiologic
concentrations targets the translational machinery to increase the proliferation of human breast
cancer cells: involvement of mammalian target of rapamycin/eIF4E pathway. Cancer Res.
2007;67(1):289-99.

4.   Pervin S, Singh R, Chaudhuri G. Nitric oxide-induced cytostasis and cell cycle arrest of a
human breast cancer cell line (MDA-MB-231): potential role of cyclin D1. Proc Natl Acad Sci.
2001;98(6):3583-8.

5.   Dubey M, Nagarkoti S, Awasthi D, Singh A, Chandra T, Kumaravelu J, et al. Nitric
oxide-mediated apoptosis of neutrophils through caspase-8 and caspase-3-dependent mechanism.
Cell Death Dis. 2016;7(9):e2348.

6.   Kumar S, Barthwal M, Dikshit M. Cdk2 nitrosylation and loss of mitochondrial potential
mediate NO-dependent biphasic effect on HL-60 cell cycle. Free Radic Biol Med.
2010;48(6):851-61.

7.   Kobayashi Y. The regulatory role of nitric oxide in proinflammatory cytokine expression
during the induction and resolution of inflammation. J Leukoc Biol. 2010;88(6):1157-62.

8.   Zeigler M, Doseff A, Galloway M, Opalek J, Nowicki P, Zweier J, et al. Presentation of
nitric oxide regulates monocyte survival through effects on caspase-9 and caspase-3 activation. J
Biol Chem. 2003;278(15):12894-902.

9.   Pautz A, Art J, Hahn S, Nowag S, Voss C, Kleinert H. Regulation of the expression of
inducible nitric oxide synthase. Nitric Oxide. 2010;23(2):75-93.

10.   Pullan S, Gidley M, Jones R, Barrett J, Stevanin T, Read R, et al. Nitric oxide in
chemostat-cultured *Escherichia col*i is sensed by Fnr and other global regulators: unaltered
methionine biosynthesis indicates lack of S nitrosation. J Bacteriol. 2007;189(5):1845-55.

11.   Gilberthorpe N, Lee M, Stevanin T, Read R, Poole R. NsrR: a key regulator
circumventing *Salmonella enterica* serovar Typhimurium oxidative and nitrosative stress *in vitro*
and in IFN-gamma-stimulated J774.2 macrophages. Microbiology. 2007;153(6):1756-71.

12.   Overton T, Justino M, Li Y, Baptista J, Melo A, Cole J, et al. Widespread distribution in
pathogenic bacteria of di-iron proteins that repair oxidative and nitrosative damage to iron-sulfur
clusters. J Bacteriol. 2008;190(6):2004-13.

13.   Bodenmiller D, Spiro S. The *yjeB* (*nsrR*) gene of *Escherichia coli* encodes a nitric oxide-
sensitive transcriptional regulator. J Bacteriol. 2006;188(3):874-81.

14.   Tucker N, Hicks M, Clarke T, Crack J, Chandra G, Le Brun N, et al. The transcriptional
repressor protein NsrR senses nitric oxide directly via a [2Fe-2S] cluster. PLoS ONE.
2008;3(11):e3623.

15.   Volbeda A, Dodd E, Darnault C, Crack J, Renoux O, Hutchings M, et al. Crystal
structures of the NO sensor NsrR reveal how its iron-sulfur cluster modulates DNA binding. Nat
Commun. 2017;8:15052.

16.   Filenko N, Spiro S, Browning D, Squire D, Overton T, Cole J, et al. The NsrR regulon of
Escherichia coli K-12 includes genes encoding the hybrid cluster protein and the periplasmic,
respiratory nitrite reductase. J Bacteriol. 2007;189(12):4410-7.

17.   Partridge J, Bodenmiller D, Humphrys M, Spiro S. NsrR targets in the *Escherichia coli*
genome: new insights into DNA sequence requirements for binding and a role for NsrR in the
regulation of motility. Mol Microbiol. 2009;73(4):680-94.

18.   Robinson J, Brynildsen M. A kinetic platform to determine the fate of nitric oxide in
Escherichia coli. PLoS Comput Biol. 2013;9(5):e103049.

19.   Poole R, Anjum M, Membrillo-Hernández J, Kim S, Hughes M, Stewart V. Nitric oxide,
nitrite, and Fnr regulation of *hmp* (flavohemoglobin) gene expression in *Escherichia coli* K-12. J
Bacteriol. 1996;178(18):5487-92.

20.   Justino M, Almeida C, Gonçalves V, Teixeira M, Saraiva L. *Escherichia coli* YtfE is a
di-iron protein with an important function in assembly of iron-sulphur clusters. FEMS Microbiol
Lett. 2006;257(2):278-84.

21.   Todorovic S, Justino M, Wellenreuther G, Hildebrandt P, Murgida D, Meyer-Klaucke W,
et al. Iron-sulfur repair YtfE protein from *Escherichia coli*: structural characterization of the di-
iron center. J Biol Inorg Chem. 2008;13(5):765-70.

22.   Miller H, Kwuan L, Schwiesow L, Bernick D, Mettert E, Ramirez H, et al. IscR is
essential for *Yersinia pseudotuberculosis* type III secretion and virulence. PLoS Pathog.
2014;10(6):e1004194.

23.   Miller H, Schwiesow L, Au-Yeung W, Auerbuch V. Hereditary hemochromatosis
predisposes mice to *Yersinia pseudotuberculosis* infection even in the absence of the type III
secretion system. Front Cell Infect Microbiol. 2016;6:69.

24.   Schwiesow L, Mettert E, Wei Y, Miller H, Herrera N, Balderas D, et al. Control of *hmu*
heme uptake genes in *Yersinia pseudotuberculosis* in response to iron sources. Front Cell Infect
Microbiol. 2018;8:47.

25.   Hubbert W, Petenyi C, Glasgow L, Uyeda C, Creighton S. *Yersinia pseudotuberculosis*
infection in the United States. Septicemia, appendicitis, and mesenteric lymphadenitis. Am J
Trop Med Hyg. 1971;20(5):679-84.

26.   Gurry J. Acute terminal ileitis and *Yersinia* infection. Br Med J. 1974;2(5913):264-6.

27.   Capron J, Delamarre J, Delcenserie R, Gineston J, Dupas J, Lorriaux A. Liver absess
complicating *Yersinia pseudotuberculosis* ileitis. Gastroenterology. 1981;81(1):150-2.

28.   Paff J, Triplett D, Saari T. Clinical and laboratory aspects of *Yersinia pseudotuberculosis*
infections, with a report of two cases. Am J Clin Pathol. 1976;66(1):101-10.

29.   Barnes P, Bergman M, Mecsas J, Isberg R. *Yersinia pseudotuberculosis* disseminates
directly from a replicating bacterial pool in the intestine. J Exp Med. 2006;203(6):1591-601.

30.   Crimmins G, Mohammadi S, Green E, Bergman M, Isberg R, Mecsas J. Identification of
MrtAB, an ABC transporter specifically required for *Yersinia pseudotuberculosis* to colonize the
mesenteric lymph nodes. PLoS Pathog. 2012;8(8):e1002828.

31.   Davis K, Mohammadi S, Isberg R. Community behavior and spatial regulation within a
bacterial microcolony in deep tissue sites serves to protect against host attack. Cell Host
Microbe. 2015;17(1):21-31.

32.   Zhang Y, Khairallah C, Sheridan B, van der Velden A, Bliska J. CCR2+ inflammatory
monocytes are recruited to *Yersinia pseudotuberculosis* pyogranulomas and dictate adaptive
responses at the expense of innate immunity during oral infection. Infect Immun.
2018;86(3):e00782-17.

33.   Peterson L, Philip N, DeLaney A, Wynosky-Dolfi M, Asklof K, Gray F, et al. RIPK1-
dependent apoptosis bypasses pathogen blockage of innate signaling to promote immune
defense. J Exp Med. 2017;214(11):3171-82.

34.   Durand E, Maldonado-Arocho F, Castillo C, Walsh R, Mecsas J. The presence of
professional phagocytes dictates the number of host cells targeted for Yop translocation during
infection. Cell Microbiol. 2010;12(8):1064-82.

35.   Sebbane F, Lemaître N, Sturdevant D, Rebeil R, Virtaneva K, Porcella S, et al. Adaptive
response of *Yersinia pestis* to extracellular effectors of innate immunity during bubonic plague.
Proc Natl Acad Sci. 2006;103(31):11766-71.

36.   Karlinsey J, Bang I, Becker L, Frawley E, Porwollik S, Robbins H, et al. The NsrR
regulon in nitrosative stress resistance of *Salmonella enterica* serovar Typhimurium. Mol
Microbiol. 2012;85(6):1179-93.

37.   Shepherd M, Achard M, Idris A, Totsika M, Phan M, Peters K, et al. The cytochrome bd-
I respiratory oxidase augments survival of multidrug-resistant *Escherichia coli* during infection.
Sci Rep. 2016;6:35285.

38.   Shiloh M, MacMicking J, Nicholson S, Brause J, Potter S, Marino M, et al. Phenotype of
mice and macrophages deficient in both phagocyte oxidase and inducible nitric oxide synthase.
Immunity. 1999;10(1):29-38.

39.   Vasquez-Torres A, Jones-Carson J, Mastroeni P, Ischiropoulos H, Fang F. Antimicrobial
actions of the NADPH phagocyte oxidase and inducible nitric oxide synthase in experimental
salmonellosis. I. Effects on microbial killing by activated peritoneal macrophages *in vitro*. J Exp
Med. 2000;192(2):227-36.

40.   Shiloh M, Nathan C. Reactive nitrogen intermediates and the pathogenesis of *Salmonella*
and *Mycobacteria*. Curr Opin Microbiol. 2000;3(1):35-42.

41.   Bang I, Liu L, Vasquez-Torres A, Crouch M, Stamler J, Fang F. Maintenance of nitric
oxide and redox homeostasis by the *Salmonella* flavohemoglobin *hmp*. J Biol Chem.
2006;281(38):28039-47.

42.   Mashruwala A, Boyd J. The *Staphylococcus aureus* SrrAB regulatory system modulates
hydrogen peroxide resistance factors, which imparts protection to aconitase during aerobic
growth. PLoS ONE. 2017;12(1):e0170283.

43.   Djaman O, Outten F, Imlay J. Repair of oxidized iron-sulfur clusters in *Escherchia coli*. J
Biol Chem. 2004;279(43):44590-99.

44.   Nuss A, Beckstette M, Pimenova M, Schmühl C, Opitz W, Pisano F, et al. Tissue dual
RNA-seq allows fast discovery of infection-specific functions and riboregulators shaping host-
pathogen transcriptomes. Proc Natl Acad Sci. 2017;114(5):E791-E800.

45.   Justino M, Almeida C, Teixeira M, Saraiva L. *Escherichia coli* di-iron YtfE protein is
necessary for the repair of stress-damaged iron-sulfur clusters. J Biol Chem.
2007;282(14):10352-9.
558
559