1 **Iron-sulfur cluster repair contributes to *Y. pseudotuberculosis* survival within deep tissues.**

4 Running title: Iron-sulfur cluster repair during *Y. pseudotuberculosis* infection

8 Kimberly M. Davis[1,#a,*], Joanna Krupp[1], Stacie Clark[1,2], Ralph R. Isberg[1]

9
10
11
12 [1]Department of Molecular Biology and Microbiology
13 [2]Graduate Program in Molecular Microbiology
14 Tufts University School of Medicine, Boston, MA, USA
15 [#a]Current Address: W. Harry Feinstone Department of Molecular Microbiology and Immunology
16 Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

22 *Corresponding author:

23 E-mail: kdavi140@jhu.edu (KMD)
24

Abstract

To successfully colonize host tissues, bacteria must respond to and detoxify many different host-

derived antimicrobial compounds, such as nitric oxide (NO). NO has direct antimicrobial

activity through attack on iron-sulfur (Fe-S) cluster-containing proteins. NO detoxification plays

an important role in promoting bacterial survival, but it remains unclear if repair of Fe-S clusters

is also important for bacterial survival within host tissues. Here we show that the Fe-S cluster

repair protein, YtfE, contributes to the survival of *Y. pseudotuberculosis* within the spleen

following nitrosative stress. *Y. pseudotuberculosis* forms clustered centers of replicating bacteria

within deep tissues, where peripheral bacteria express the NO-detoxifying gene, *hmp*. *ytfE*

expression also occurred specifically within peripheral cells at the edges of microcolonies. In the

absence of *ytfE*, the area of microcolonies was significantly smaller than WT, consistent with

ytfE contributing to the survival of peripheral cells. The loss of *ytfE* did not alter the ability of

cells to detoxify NO, which occurred within peripheral cells in both WT and Δ*ytfE*

microcolonies. In the absence of NO-detoxifying activity by *hmp*, NO diffused across Δ*ytfE*

microcolonies, and there was a significant decrease in the area of microcolonies lacking *ytfE*,

indicating that *ytfE* also contributes to bacterial survival in the absence of NO detoxification.

These results indicate a role for Fe-S cluster repair in the survival of *Y. pseudotuberculosis*

within the spleen, and suggest that extracellular bacteria may rely on this pathway for survival

within host tissues.

Introduction

Nitric oxide (NO) is a diffusible gas that has a wide range of physiological functions

within mammals (1, 2). The effects of NO are tissue concentration-dependent, as it promotes

vasodilation, cell proliferation and cell differentiation at low concentrations (3, 4), while high

concentrations drive apoptosis and defense against bacterial, fungi, and parasites (5-8). NO is

produced by three different nitric oxide synthase (NOS) isoforms within mammalian tissues: the

neuronal NOS (nNOS), endothelial NOS (eNOS), and inducible NOS (iNOS). nNOS and eNOS

are expressed at low levels by endothelial and neuronal cells respectively (2). In contrast, iNOS

is expressed by a wide range of cell types, specifically in response to NF-κB-dependent sensing

of, and is responsible for the high levels of NO produced during infection (1, 9).

NO has direct antimicrobial activity and can also react with reactive oxygen species to

produce additional toxic compounds, such as peroxynitrite. NO may be bacteriostatic, while

peroxynitrite is known to have direct bactericidal activity (10). NO antibacterial activity occurs

through nitrosylation of iron-sulfur (Fe-S) cluster-containing proteins, which play critical roles in

cellular respiration, DNA synthesis, and gene regulation. NO also targets heme groups, reactive

thiols, tyrosyl radicals, and can cause DNA damage (1, 10). One of the global regulators in *E.*

coli that has been shown to be inactivated by NO is NsrR (11), which regulates the response to

nitrosative stress and is also associated with the oxidative stress response (12-14). Nitrosylation

of the NsrR-associated Fe-S cluster relieves repression of at least 60 genes in *E. coli* (15-18).

Included in this regulon is the *hmp* gene, which encodes a flavohemoglobin that detoxifies NO

(19, 20), and YtfE (also known as Repair of Iron Centers (RIC) in *E. coli*), which functions to

repair Fe-S clusters following NO damage (13, 17). Repair of Fe-S cluster proteins by YtfE can

eliminate the need for new Fe-S cluster biogenesis when Fe availability is limited (13, 21, 22). A

variety of studies argue that YtfE contributes to bacterial survival within host cells, but it remains

unclear if YtfE contributes to the survival of extracellular pathogens replicating within host

tissues, and if YtfE contributes to survival following exposure to NO (23-25).

Yersinia pseudotuberculosis is an oral pathogen that is typically contained within

intestinal tissues and gut-associated lymphoid tissues, but has the capacity to spread systemically

in susceptible individuals (26-29). Following bloodstream access, *Y. pseudotuberculosis*

colonizes deep tissue sites where individual bacteria replicate to form clonal microcolonies (30-

32). Neutrophils, monocytes, and macrophages are recruited to sites of bacterial replication,

which are kept at bay by *Yersinia*, primarily via substrates of the bacterial type III secretion

system (33-35). Recruited monocytes and macrophages produce NO, which diffuses across a

layer of neutrophils and is inactivated at the periphery of the microcolony by bacteria expressing

Hmp, preventing diffusion of NO into the interior of the microcolony (32). The consequences of

the selective attack of NO on the peripheral bacteria are unclear, although it is likely that

nitrosative stress may slow their growth. Additional members of the nitrosative stress response,

such as YtfE, may also be required in peripheral cells to ensure their survival.

Bacteria responding to RNS appear to recover and remain viable, consistent with

members of the NsrR regulon cooperating to repair NO-mediated damage in peripheral bacteria

within microcolonies. Two of the members of the *E. coli* NsrR regulon, *hmp* and *ytfE,* are

upregulated in the bubonic plague model of *Yersinia pestis* infection and during Peyer's patch

colonization by *Y. pseudotuberculosis* (36, 37). The similar expression patterns of *hmp* and *ytfE*

may suggest that both genes are also members of the NsrR regulon in *Yersinia,* and could both

contribute to microbial fitness during microcolony growth. Additionally, very few studies have

explored the role of Fe-S cluster repair during infection with extracellular bacteria. Here, we

show that *ytfE* contributes to the survival of extracellular bacteria, specifically through

upregulation of *ytfE* within peripheral cells of *Y. pseudotuberculosis* microcolonies.

Results.

Y. pseudotuberculosis ytfE **expression is regulated by NsrR and occurs within peripheral**

cells during growth in the spleen.

The *ytfE* gene is known to be a member of the NsrR regulon in a number of bacterial

species (13, 14, 38), so *ytfE* is expected to be repressed by NsrR in *Y. pseudotuberculosis*. We

also expected that *ytfE* is transcribed during *Y. pseudotuberculosis* growth within the mouse

spleen as the NsrR-regulated *hmp* gene is expressed in this tissue (23-25, 32, 36). To determine if

Y. pseudotuberculosis ytfE is expressed during splenic growth, C57BL/6 mice were intravenously

challenged with bacteria and bacterial RNA was isolated at day 3 post-inoculation (PI). Based on

qRT-PCR analysis, *ytfE* transcript levels increased within the mouse spleen relative to the

inoculum culture grown in the absence of an NO-generating system, with marked mouse-to-

mouse variation (Figure 1A) (32). To determine if *ytfE* expression is NsrR-dependent, we

compared *ytfE* transcription levels in WT and Δ*nsrR* strains in the presence (+) and absence (-) of

nitrogen stress imparted by acidified nitrite (NO_2). Transcription of *ytfE* increased by over 100X

in the WT strain with the addition of nitrogen stress (Figure 1B). Transcription of *ytfE* was high

in the Δ*nsrR* strain in either the presence or absence of nitrogen stress, indicating that NsrR

negatively regulates *ytfE* transcription. A slight increase in *ytfE* expression in the Δ*nsrR* strain

with nitrogen stress suggests additional pathways may regulate *ytfE* expression.

Hmp, another member of the NsrR regulon, is specifically expressed on the periphery of

Y. pseudotuberculosis microcolonies during splenic growth (32). At the transcriptional level,

this is associated with considerable variation in expression levels between individual mice.

Variation is likely due to the presence of different size microcolonies within different organs, as

distinct peripheral expression of *hmp* is visualized in large microcolonies, and smaller

microcolonies are more homogenous. The variable *ytfE* expression levels in the mouse and

NsrR-dependence is reminiscent of *hmp* expression, indicating that there could be a link between

spatial expression and inter-mouse variation. To determine if *ytfE* and *hmp* have similar

expression patterns during growth in the spleen, mice were intravenously inoculated with a WT

Y. pseudotuberculosis strain containing *hmp::mCherry* (chromosomal integration of *mCherry*

downstream of *hmp*) and *ytfE::gfp* (chromosomal integration of *gfp* downstream of *ytfE*).

Microcolonies were visualized within the spleen using fluorescence microscopy, and *ytfE* and

hmp reporter signals were quantified within the same cells at the center and periphery of the

microcolonies (Materials and Methods). The *hmp* reporter signal increased within individual

cells at the periphery relative to cells at the centroid of microcolonies, which generated a ratio

value greater than 1, consistent with *hmp* peripheral expression (Figure 1C). The *ytfE* signal was

dim relative to *hmp*, but also increased at the periphery of microcolonies relative to the centroid.

The Periphery/Centroid signal intensity ratio value for *ytfE* was similar to *hmp*, indicating that

ytfE was expressed at the periphery of microcolonies (Figure 1C). To confirm the dim *ytfE*

signal was not due to mCherry fluorescence detected in the gfp channel, experiments were also

performed in a WT strain containing *yopE::mCherry* and *ytfE::gfp*. *yopE* and *ytfE* exhibited

distinct reporter patterns indicating the detected gfp signal was due to *ytfE* reporter expression

(Figure 1C, 1D). There was significant overlap in the *hmp* and *ytfE* signals within these images,

and based on NsrR-dependent regulation of both genes, it is likely that *hmp* and *ytfE* were

expressed in the same cells (Figure 1D).

ytfE contributes to the virulence of Y. pseudotuberculosis in the spleen.

YtfE repairs Fe-S clusters damaged by NO, and there appears to be sufficient NO at the

periphery of splenic microcolonies to allow synthesis of this protein and to promote repair within

this subpopulation of bacteria. To determine if loss of *yftE* alters the overall fitness of *Y.*

pseudotuberculosis during growth within the spleen, we constructed a *Y. pseudotuberculosis*

strain that lacks the *ytfE* gene (Δ*ytfE*) and harbors a constitutive *gfp*-expressing plasmid, to

visualize growth within the spleen. To compare differences in relative fitness, mice were

infected intravenously with equal amounts of WT mCherry$^+$ and Δ*ytfE* GFP$^+$ strains, and spleens

were harvested at day 3 PI, a late stage of infection in this model, to determine the competitive

index by colony forming units (CFUs) and quantify microcolony areas within the same animals.

At day 3 PI, the median competitive index was significantly less than 1, indicating lowered

fitness of the Δ*ytfE* strain relative to the WT strain (Figure 2A; p=0.0115). The areas of

individual microcolonies within these co-infected tissues were also visualized and quantified by

fluorescence microscopy in 11 mice. The Δ*ytfE* microcolonies were significantly smaller than

WT microcolonies within the same organs, indicating that *ytfE* contributes to the survival of *Y.*

pseudotuberculosis in the spleen, presumably due to lowered fitness of the bacterial population

located at the periphery of the microcolonies (Figure 2B).

Since YtfE could directly repair the Fe-S cluster of NsrR, the absence of *ytfE* could alter

expression of the NsrR regulon, by resulting in heightened expression within peripheral cells. To

determine if microcolonies from the *Y. pseudotuberculosis* Δ*ytfE* strain have sustained

expression of the NsrR regulon relative to the WT strain, we infected mice intravenously with

WT GFP$^+$ *hmp::mCherry* or Δ*ytfE* GFP$^+$ *hmp::mCherry* integrated *hmp* reporter strains, and

spleens were harvested at day 3 PI to visualize reporter expression by fluorescence microscopy.

Similar to Fig. 1D, *hmp* reporter expression was significantly higher at the periphery relative to

the centroid in the WT strain (Figure 2C). The reporter expression pattern was very similar in

Δ*ytfE* microcolonies, indicating that there was still a gradient of NO exposure in the mutant, in

which Hmp activity in the peripheral population protects the central population of bacteria, and

that loss of *ytfE* did not significantly alter expression of the NsrR regulon (Figure 2D).

***ytfE* contributes to bacterial survival in the absence of *hmp*.**

YftE contributed to the growth of *Y. pseudotuberculosis* microcolonies in the spleen

despite expression limited to the microcolony periphery. We were then interested in determining

if YtfE-mediated repair played an important role in the context of a Δhmp strain, where all

bacteria in a microcolony are exposed to NO (32). To address this point, we challenged mice

intravenously with Δhmp GFP$^+$ $P_{hmp}::mCherry$ and Δhmp $\Delta ytfE$ GFP$^+$ $P_{hmp}::mCherry$ strains, and

spleens were harvested at day 3 PI to quantify CFUs, visualize microcolony areas, and detect

reporter signals by fluorescence microscopy. The Δhmp $\Delta ytfE$ strain showed no potentiation of

the defect in single strain infections, but this defect can be observed during co-infection. The

median competitive index (CI) value for the double mutant was below 1, but was not

significantly less than 1, which suggests that Δhmp $\Delta ytfE$ strain may not be less fit than Δhmp

(Figure 3A). Interestingly, in 4 mice the CI was at least 1, indicating that the Δhmp $\Delta ytfE$ strain

could compete efficiently with Δhmp in these animals. We then compared the total CFU in the

organs in mice in which the Δhmp strain outcompeted Δhmp $\Delta ytfE$, or in mice in which no such

outcompetition took place (Figure 3B). The total CFUs were significantly lower in organs in

which the Δhmp did not outcompete the $\Delta hmp\Delta ytfE$ strain, indicating that the fitness differences

were suppressed in animals showing increased restriction of bacterial growth. It is also possible

that the $\Delta hmp\Delta ytfE$ strain had reduced levels of initial seeding within tissues. The microcolony

areas were quantified within all the spleens depicted in Figure 3A, and the areas of Δhmp $\Delta ytfE$

microcolonies were significantly smaller than Δhmp (Figure 3C). Δhmp $\Delta ytfE$ microcolonies had

similar P_{hmp} reporter expression at the centroid and periphery, indicating NO diffused across

these centers (Figure 3D). Together, these results confirm that *ytfE* contributes to bacterial

survival in the absence of Hmp detoxifying activity.

ytfE has limited effects on NO sensitivity in the absence of _hmp_

187   The Δ*ytfE* single mutant strain had reduced fitness relative to the WT strain (Figure 2),

however the single Δ*ytfE* mutant was not significantly more sensitive than the WT strain to the

presence of acidified nitrite during growth in culture. Similarly, despite the lowered fitness of the

Δ*hmp* Δ*ytfE* strain in spleens relative to Δ*hmp*, the Δ*hmp* Δ*ytfE* strain was not significantly more

sensitive than the Δ*hmp* mutant strain to the presence of acidified nitrite during growth in culture

(Figure 4A; compare +NO_2 samples). The absence of *ytfE* also did not significantly alter the

sensitivity of strains to the NO donor compound, DETANONOate, during growth in minimal

media (Figure 4B). This is consistent with previous reports in other organisms (38, 39), perhaps

due to the presence of other backup repair pathways that are active in the absence of Hmp or

because YtfE plays a role in protection from other stress species.

197   We then compared *hmp* reporter expression in Δ*ytfE* and Δ*hmp*Δ*ytfE* strains to confirm

that NO diffusion occurred across Δ*hmp* Δ*ytfE* microcolonies using plasmid-borne reporters.

Mice were infected intravenously with Δ*ytfE* GFP$^+$ P$_{hmp}$::*mCherry* or Δ*hmp* Δ*ytfE* GFP$^+$

P$_{hmp}$::*mCherry* strains, and spleens were harvested 3 days PI to visualize reporter expression by

fluorescence microscopy. The Δ*ytfE* strain had peripheral P$_{hmp}$ reporter expression, as seen with

the chromosomally-integrated *hmp* reporter, indicating that NO diffusion across the microcolony

is inhibited by peripheral cells in Δ*ytfE* microcolonies (Figure 4C, 4D). Δ*hmp* Δ*ytfE*

microcolonies showed no such preference for the periphery, indicating NO diffused across these

centers, as expected based on the loss of *hmp*.

Rescue of _ytfE_ restores microcolony size.

To show that the loss of *ytfE* was responsible for decreased microcolony size, we rescued

the Δ*ytfE* strain with a WT copy of *ytfE*, and transformed the *ytfE* rescued strain with the

constitutive *gfp*-expressing plasmid. Mice were infected intravenously with WT mCherry$^+$ and

ytfE rescued GFP$^+$ strains, and spleens were harvested at day 3 PI to quantify CFUs and visualize

microcolony areas by fluorescence microscopy. The median competitive index for the *ytfE*

rescued strain was close to a value of 1, indicating the fitness of this strain was roughly

equivalent to the WT strain (Figure 5A). The microcolony areas were quantified within all the

spleens depicted in Figure 5A, and the areas of *ytfE* rescued microcolonies were very similar to

WT microcolonies (Figure 5B, 5C). These results indicate that the Δ*ytfE* strain was rescued by

the WT copy of *ytfE*, which confirms that the reduced fitness of the Δ*ytfE* strain was specifically

due to a loss of *ytfE.*

Discussion

The detoxification of NO and other reactive nitrogen species is critical for bacterial survival within host tissues (40-43). Bacterial proteins involved in NO detoxification, however, are not synthesized until NO accumulates and damages the Fe-S clusters of NsrR, resulting in inactivation of this repressor (15, 16). Many additional Fe-S cluster-containing proteins are simultaneously damaged, so bacteria need to either repair damaged proteins or synthesize replacement proteins, while simultaneously replenishing proteins that detoxify NO and prevent further damage (17, 38). Although Fe-S cluster repair is likely required for bacterial survival, it has been unclear whether or not this plays an important role within host tissues. We have chosen to ask this question in a mouse model of *Y. pseudotuberculosis* infection, where it is known that bacteria respond to RNS and that Fe concentrations are limiting (24, 25, 32). *ytfE* expression had been detected in *Yersinia* species replicating within host tissues, however it was unclear if Fe-S cluster repair or assembly play a significant role during infection (36, 37). Here, we have shown that Fe-S cluster repair contributes to successful *Y. pseudotuberculosis* replication within the spleen.

We found that a place where Fe-S cluster repair in *Y. pseudotuberculosis* likely occurs is in the peripheral subpopulation of bacteria that responds to NO assault within microcolonies. This is consistent with a role for YftE in supporting survival of the peripheral subpopulation. In the absence of *ytfE*, we would expect bacteria on the periphery to be exposed to stress associated with NO exposure, leading to sequential loss of the peripheral population and progressively smaller microcolonies. Consistent with this hypothesis, we observed Δ*ytfE* microcolonies were smaller than those established by the WT strain. It is expected that the difference between WT and Δ*ytfE* microcolony areas will become progressively more pronounced as the infection

241   proceeds, because Δ*ytfE* microcolonies should continuously lose their peripheral subpopulation.

242   This prediction is based on our previous observation that NO-sensitive microcolonies are

243   progressively reduced during the course of disease, with elimination of Δ*hmp* bacteria by NO

244   most pronounced at late timepoints post-inoculation, concurrent with a timepoint in which the

245   animals inoculated with the WT strain are moribund (32).

246   NO alone was not sufficient to limit the growth of Δ*ytfE* bacteria in bacteriological

media, consistent with previous studies that indicated the uropathogenic *E. coli* (UPEC) Δ*ytfE*

strain had reduced intracellular survival within host cells, but was not sensitive to exogenous NO

alone (39). Presumably, the Δ*ytfE* strain is sensitive to other antimicrobial compounds generated

within host tissues, possibly by the generation of a variety of reactive nitrogen species (RNS) as

a consequence of NO reaction with reactive oxygen species (ROS) or other compounds.

Upregulation of *ytfE* following hydrogen peroxide-mediated damage of *Staphylococcus aureus*

indicates that YtfE may play a broad role in repair, instead of just a response to nitrogen stress

(44, 45). Additional studies in *E. coli* also suggest that YtfE repairs Fe-S cluster proteins

damaged by hydrogen peroxide (13, 46-49).

256   An additional regulatory signal for the *yftE* gene is iron limitation, which depresses

synthesis of new Fe-S cluster proteins, thus requiring cellular YftE function (14, 21, 50). The

spleen is expected to be an iron-limiting environment requiring Fe-S cluster repair, but we found

no evidence for NO-independent induction of the *ytfE* gene within the center of microcolonies,

using our fluorescent promoter constructions. This contrasts with data arguing that *Y.*

pseudotuberculosis expresses many Fe acquisition genes during growth in host tissues, consistent

with Fe-limiting conditions within host tissues (25, 37). It is possible that the center of the

microcolony represents a protected environment with low exposure to stresses such as reactive

nitrogen and oxygen species as well as limited damage to Fe-S centers. Additionally, *ytfE*

expression may be induced at very low NO concentrations, as seen in *Salmonella enterica*

serovar Typhimurium (38), but may require severe iron limitation for expression in the absence

of NO.

The loss of YftE function also has the potential to alter metabolite levels due to

disruption of protein functions that are Fe-S center-related. In the presence of RNS, a Δ*ytfE*

strain may have reduced aconitase and fumarase activity due to the role of YtfE in Fe-S cluster

repair specifically for these proteins (47). The Fe-S cluster of the NsrR repressor is also repaired

by YtfE, which could lead to prolonged expression of the NsrR regulon in the presence of NO

(13). Our results indicate that expression of the NsrR regulon is similar in the presence or

absence of *ytfE*, although it remains possible that prolonged expression of the NsrR regulon

could be detected at later timepoints post-inoculation. Future work will investigate these issues,

and determine the interplay between iron regulation, NO-induced damage and repair of critical

Fe-S centers.

Materials and Methods

Bacterial strains & growth conditions. The WT *Y. pseudotuberculosis* strain, YPIII, was used throughout. For all mouse infection experiments, bacteria were grown overnight to post-exponential phase in 2xYT broth (LB, with 2x yeast extract and tryptone) at 26º C with rotation. Exponential phase cultures were sub-cultured 1:100 from overnight cultures, and grown at 26º C with rotation for an additional 2 hours. Sodium nitrite (2.5mM) was added to LB pH5.5 to induce the nitrogen stress response. DETA-NONOate (2.5mM) NO donor compound (Cayman Chemicals) was added to M9 minimal media to test NO sensitivity.

Generation of ytfE mutant strains. The *hmp* and *nsrR* deletion strains were previously described (32). Deletion derivative strains were generated for *ytfE* by amplifying the start codon + 3 downstream codons, the 3' terminal 3 codons + the stop codon and fusing these fragments to generate a start + 6 aa + stop construct. Deletion constructions were amplified with 500 base pairs flanking sequence on each side, cloned into the suicide vector, pSR47S, and transformed into *Y. pseudotuberculosis*. Sucrose selection was used to select for bacteria that had incorporated the desired mutation after a second cycle of recombination (31). PCR, sequencing, and qRT-PCR were used to confirm deletion strains.

Integrated ytfE reporter construction (ytfE⁺rescue strain). The *ytfE::gfp* reporter was generated by cloning *gfp* immediately downstream of the *ytfE* gene (in between the *ytfE* stop codon and terminator sequence) by overlap extension PCR. The *ytfE* start codon was amplified with 500 base pairs upstream flanking sequence, while the stop codon of *gfp* was amplified with 500 base pairs downstream flanking sequence. This fragment was cloned into the suicide vector, pSR47S, and transferred by conjugation into WT *Y. pseudotuberculosis hmp::mCherry* (chromosomally integrated), selecting for kanamycin resistance. For the *ytfE⁺* rescue strain, a

WT *ytfE* gene product, including 500 base pairs upstream and downstream of *ytfE*, was amplified

from genomic DNA and cloned into pSR47S. This vector was transferred by conjugation into

Δ*ytfE Y. pseudotuberculosis*, selecting for kanamycin resistance. A second round of

recombination was selected on sucrose-containing media to isolate strains that had recombined

each *ytfE* construct. PCR and sequencing were used to confirm integration of *gfp* or rescue of

the *ytfE* deletion.

Generation of plasmid-based reporter strains. Several of the *Y. pseudotuberculosis*

reporter strains in this study have been previously described: WT GFP$^+$, WT mCherry$^+$

(*yopE::mCherry)*, WT *hmp::mCherry*, WT GFP$^+$ *hmp::mCherry*, and Δ*hmp* GFP$^+$ (32). For this

study, GFP$^+$ strains were constructed by transforming deletion strains with the constitutive GFP

plasmid, which expresses GFP from an unrepressed P_{tet} of pACYC184. The P_{hmp}*::mCherry* was

also transformed into GFP$^+$ strains. The P_{hmp}*::mCherry* transcriptional fusion was previously

constructed by fusing the *hmp* promoter to *mCherry* using overlap extension PCR, and cloned

into the pMMB67EH plasmid (32).

Murine model of systemic infection. Six to 8-week old female C57BL/6 mice were

obtained from Jackson Laboratories (Bar Harbor, ME). All animal studies were approved by the

Institutional Animal Care and Use Committee of Tufts University. Mice were injected

intravenously with 10^3 bacteria for all experiments. For co-infection experiments, mice were

inoculated with 5 x 10^2 CFU of each strain, for a total of 10^3 CFUs. At the indicated timepoints

post-inoculation (PI) (3 days), spleens were removed and processed.

qRT-PCR to detect bacterial transcripts in broth-grown cultures. Bacterial cells were

grown in broth to $A_{600} = 0.3$ under appropriate stress conditions, pelleted, resuspended in Buffer

RLT (QIAGEN) + ß-mercaptoethanol, and RNA was isolated using the RNeasy kit (QIAGEN)

according to manufacturer's protocol. DNA contamination was eliminated using the DNA-free

kit (Ambion) according to manufacturer's protocol. RNA was reverse transcribed using M-MLV

reverse transcriptase (Invitrogen), in the presence of the RNase inhibitor, RnaseOUT

(Invitrogen), according to manufacturer's protocol. Approximately 30 ng cDNA was used as a

template in reactions with 0.5 µM of forward and reverse primers and SYBR Green (Applied

Biosystems) according to manufacturer's protocol. Control samples were prepared that lacked

M-MLV, to confirm DNA was eliminated from samples and was not amplified by qRT-PCR.

Reactions were carried out using the StepOnePlus Real-Time PCR system, and relative

comparisons were obtained using the $\Delta\Delta C_T$ or $2^{-\Delta Ct}$ method (Applied Biosystems).

qRT-PCR to detect bacterial transcripts from mouse tissues. Mice were inoculated

intravenously with the WT strain, and at day 3 PI spleens were harvested, and immediately

submerged in RNALater solution (QIAGEN). Tissue was homogenized in Buffer RLT + ß-

mercaptoethanol, and RNA was isolated using the RNeasy kit (QIAGEN) according to

manufacturer's protocol. Bacterial RNA was enriched following depletion of host mRNA and

rRNA from total RNA samples, using the MICROB*Enrich* kit (Ambion) according to

manufacturer's protocol. DNA digestion, reverse transcription, and qRT-PCR were performed

as described above.

Fluorescence microscopy. C57BL/6 mice were inoculated intravenously with the *Y.*

pseudotuberculosis WT strain, at day 3 PI spleens were harvested and immediately fixed in 4%

paraformaldehyde in PBS for 3 hours. Tissues were frozen-embedded in Sub Xero freezing

media (Mercedes Medical) and cut by cryostat microtome into 10µm sections. To visualize

reporters, sections were thawed in PBS, stained with Hoechst at a 1:10,000 dilution, washed in

PBS, and coverslips were mounted using ProLong Gold (Life Technologies). Tissue was imaged

with 20x or 63x objectives, using a Zeiss Axio Observer.Z1 (Zeiss) fluorescent microscope with

Colibri.2 LED light source, an Apotome.2 (Zeiss) for optical sectioning, and an ORCA-R^2 digital

CCD camera (Hamamatsu).

Image analysis. Volocity image analysis software was used to quantify microcolony

areas. Image J was used to quantify the signal intensity of each channel at the centroid and

periphery of each microcolony, to generate relative signal intensities of fluorescent reporters.

Thresholding was used to define the area of each microcolony, the centroid was calculated, and

0.01 pixel^2 squares were selected to calculate values at the centroid. Peripheral measurements

depict bacteria in contact with host cells.

Author Contributions. Designed and performed experiments: KMD, JK, SC.

Intellectual/conceptual contribution: KMD, RRI. Analyzed the data: KMD, JK, SC, RRI. Wrote

the paper: KMD, RRI.

Acknowledgments. We thank the members of the Isberg lab, who provided valuable

advice and feedback throughout this project. We also thank the members of the Davis lab, who

provided feedback and suggestions during the final steps of manuscript preparation. The authors

of this manuscript declare no conflicts of interest. This work was supported by NIAID award

R01 AI110684 as well as by an American Cancer Society-Ellison Foundation Postdoctoral

Fellowship (PF-13-360-01-MPC), and a NIAID K22 Career Transition Award (1K22AI123465-

01).

Figure Legends:

Figure 1: *ytfE* expression is regulated by NsrR and occurs within peripheral cells. A)

C57BL/6 mice were inoculated intravenously (i.v.) with 10^3 WT *Y. pseudotuberculosis*, and

spleens were harvested at day 3 post-inoculation (PI). Bacterial transcripts were isolated from

splenic tissue, transcript levels were quantified by qRT-PCR relative to 16S, and the fold

increase in *ytfE* transcript levels is shown relative to the inoculum. Each dot represents an

individual mouse. B) Nitrogen stress was induced (NO_2, +) in cultures of WT and Δ*nsrR* strains,

and compared to untreated cultures (NO_2, -). The *ytfE* transcript levels are expressed relative to

16S, fold increase is relative to the average level in untreated WT cultures. Each column: 5

biological replicates, mean and SEM. C) C57BL/6 mice were inoculated i.v. with the WT

hmp::mCherry ytfE::gfp strain or the WT *yopE::mCherry ytfE;:gfp* strain, and spleens were

harvested at day 3 PI for fluorescence microscopy. Fluorescent signal intensity was determined

within the same peripheral and centroid cells for *hmp* and *ytfE* reporters or *yopE* and *ytfE*

reporters, and divided to generate the periphery/centroid ratio (4 mice/group). Dots: individual

microcolonies. D) Representative image of *hmp::mCherry* and *ytfE::gfp* reporters (top panels) or

yopE::mCherry and *ytfE::gfp* reporters (bottom panels); merge and single channel images are

shown. Scale bars: 20μm. Statistics: Wilcoxon matched pairs, **$p<0.01$, n.s.: not significant.

Figure 2: *ytfE* contributes to growth in the spleen. A) Co-infection with the WT and Δ*ytfE*

strains. C57BL/6 mice were inoculated i.v. with an equal mixture of mCherry⁺ (*yopE::mCherry*)

WT and GFP⁺ Δ*ytfE* strains, and spleens were harvested at day 3 PI. Competitive index: ratio of

Δ*ytfE*/WT CFUs in spleen at day 3, divided by ratio of Δ*ytfE*/WT CFUs in the inoculum. Dots:

individual mice. Dotted line: equal fitness, replicates completed on two separate days. B) WT

and $\Delta ytfE$ microcolony areas (μm^2) from the co-infection in panel 2A quantified (Experimental

Procedures) in 11 mice. C) C57BL/6 mice were inoculated i.v. with the WT GFP$^+$

hmp::mCherry or $\Delta ytfE$ GFP$^+$ *hmp::mCherry* strains, and spleens were harvested at day 3 PI for

fluorescence microscopy. Reporter signals were quantified within peripheral and centroid cells,

hmp reporter signal was divided by GFP. D) Representative image of a $\Delta ytfE$ GFP$^+$

hmp::mCherry microcolony, stained with Hoechst to detect host nuclei; merge and *hmp* single

channel images are shown Statistics: A): Wilcoxon Signed Rank Test, compared to a value of 1,

B): Mann-Whitney, C): Wilcoxon matched pairs, *p<0.05, ***p<0.001.

Figure 3: $\Delta ytfE$ contributes to survival in the absence of *hmp*. A) C57BL/6 mice were

inoculated i.v. with an equal mixture of *hmp* GFP$^+$ and Δhmp $\Delta ytfE$ GFP$^+$ P_{hmp}::*mCherry* strains,

and spleens were harvested at day 3 PI. Competitive index: ratio of Δhmp $\Delta ytfE$/Δhmp CFUs in

spleen at day 3, divided by ratio of Δhmp $\Delta ytfE$/Δhmp CFUs in the inoculum. Dots: individual

mice. Dotted line: represents equal fitness. Statistics: Wilcoxon Signed Rank Test, compared to

a value of 1, n.s.: not significant. B) Total CFUs in the spleen during co-infection, when CI was

less than 1 (Δhmp wins) or above or equal to 1 (Δhmp $\Delta ytfE$ wins). Dots: individual mice. C)

Δhmp and Δhmp $\Delta ytfE$ microcolony areas (μm^2) from the co-infection quantified (Experimental

Procedures) in 10 mice. Dots: individual microcolonies. D) Reporter signals were quantified

within peripheral and centroid cells in the Δhmp $\Delta ytfE$ strain during co-infection, *hmp* reporter

signal was divided by GFP. Dots: individual microcolonies. Statistics: B) and C): Mann-

Whitney, D): Wilcoxon matched pairs, *p<0.05, **p<0.01, n.s.: not significant.

414 **Figure 4**: ***ytfE* has limited effects on NO sensitivity in the absence of *hmp*.** A) Optical density

415 (A_{600nm}) was measured every 2 hours (time, hrs) within cultures of the indicated strains during

416 incubation in LB pH5.5 in the presence ($+NO_2$) and absence of $NaNO_2$. Data represents two

417 independent experiments. B) Optical density (A_{600nm}) was measured every 2 hours (time, hrs)

418 within cultures of the indicated strains during incubation in M9 minimal media in the presence

419 ($+NO$) and absence of the DETANONOate NO donor. Data represents three independent

420 experiments. C) C57BL/6 mice were inoculated i.v. with the $\Delta ytfE$ GFP^+ $P_{hmp}::mCherry$ or

421 Δhmp $\Delta ytfE$ GFP^+ $P_{hmp}::mCherry$ strains, and spleens were harvested at day 3 PI for fluorescence

422 microscopy. Reporter signals were quantified within peripheral and centroid cells, *hmp* reporter

423 signal was divided by GFP. Dots: individual microcolonies. D) Representative images of $\Delta ytfE$

424 and Δhmp $\Delta ytfE$ microcolonies; merge and *hmp* single channel images are shown. Statistics:

425 Wilcoxon matched pairs, *p<0.05, n.s.: not significant.

427 **Figure 5: Rescue of *ytfE* restores fitness of the $\Delta ytfE$ strain.** A) C57BL/6 mice were

428 inoculated i.v. with an equal mixture of *ytfE* rescued GFP^+ and WT $mCherry^+$ strains, and spleens

429 were harvested at day 3 PI. Competitive index: ratio of *ytfE* rescue/WT CFUs in spleen at day 3,

430 divided by ratio of *ytfE* rescue/WT CFUs in the inoculum. Dots: individual mice. Dotted line:

431 equal fitness, replicates completed on two separate days. Statistics: Wilcoxon Signed Rank Test,

432 compared to a value of 1, n.s.: not significant. B) WT and *ytfE* rescue microcolony areas (μm^2)

433 from the co-infection quantified (Experimental Procedures) in 10 mice. Dots: individual

434 microcolonies. C) Representative images of WT $mCherry^+$ and *ytfE* rescued GFP^+ microcolonies.

435 Scale bar: 20μm. Statistics: Mann-Whitney, n.s.: not significant.

References

1.   Bogdan C. Nitric oxide and the immune response. Nat Immunol. 2001;2(10):907-16.

2.   Nathan C. Nitric oxide as a secretory product of mammalian cells. FASEB J.
1992;6(12):3051-64.

3.   Pervin S, Singh R, Hernandez E, Wu G, Chaudhuri G. Nitric oxide in physiologic
concentrations targets the translational machinery to increase the proliferation of human breast
cancer cells: involvement of mammalian target of rapamycin/eIF4E pathway. Cancer Res.
2007;67(1):289-99.

4.   Pervin S, Singh R, Chaudhuri G. Nitric oxide-induced cytostasis and cell cycle arrest of a
human breast cancer cell line (MDA-MB-231): potential role of cyclin D1. Proc Natl Acad Sci.
2001;98(6):3583-8.

5.   Dubey M, Nagarkoti S, Awasthi D, Singh A, Chandra T, Kumaravelu J, et al. Nitric
oxide-mediated apoptosis of neutrophils through caspase-8 and caspase-3-dependent mechanism.
Cell Death Dis. 2016;7(9):e2348.

6.   Kumar S, Barthwal M, Dikshit M. Cdk2 nitrosylation and loss of mitochondrial potential
mediate NO-dependent biphasic effect on HL-60 cell cycle. Free Radic Biol Med.
2010;48(6):851-61.

7.   Kobayashi Y. The regulatory role of nitric oxide in proinflammatory cytokine expression
during the induction and resolution of inflammation. J Leukoc Biol. 2010;88(6):1157-62.

8.   Zeigler M, Doseff A, Galloway M, Opalek J, Nowicki P, Zweier J, et al. Presentation of
nitric oxide regulates monocyte survival through effects on caspase-9 and caspase-3 activation. J
Biol Chem. 2003;278(15):12894-902.

9.   Pautz A, Art J, Hahn S, Nowag S, Voss C, Kleinert H. Regulation of the expression of
inducible nitric oxide synthase. Nitric Oxide. 2010;23(2):75-93.

10.   Fang F. Perspective series: host/pathogen interactions. Mechanisms of nitric oxide-
related antimicrobial activity. J Clin Invest. 1997;99(12):2818-25.

11.   Pullan S, Gidley M, Jones R, Barrett J, Stevanin T, Read R, et al. Nitric oxide in
chemostat-cultured *Escherichia coli* is sensed by Fnr and other global regulators: unaltered
methionine biosynthesis indicates lack of S nitrosation. J Bacteriol. 2007;189(5):1845-55.

12.   Gilberthorpe N, Lee M, Stevanin T, Read R, Poole R. NsrR: a key regulator
circumventing *Salmonella enterica* serovar Typhimurium oxidative and nitrosative stress *in vitro*
and in IFN-gamma-stimulated J774.2 macrophages. Microbiology. 2007;153(6):1756-71.

13.   Overton T, Justino M, Li Y, Baptista J, Melo A, Cole J, et al. Widespread distribution in
pathogenic bacteria of di-iron proteins that repair oxidative and nitrosative damage to iron-sulfur
clusters. J Bacteriol. 2008;190(6):2004-13.

14.   Bodenmiller D, Spiro S. The yjeB (nsrR) gene of *Escherichia coli* encodes a nitric oxide-
sensitive transcriptional regulator. J Bacteriol. 2006;188(3):874-81.

15.   Tucker N, Hicks M, Clarke T, Crack J, Chandra G, Le Brun N, et al. The transcriptional
repressor protein NsrR senses nitric oxide directly via a [2Fe-2S] cluster. PLoS ONE.
2008;3(11):e3623.

16.   Volbeda A, Dodd E, Darnault C, Crack J, Renoux O, Hutchings M, et al. Crystal
structures of the NO sensor NsrR reveal how its iron-sulfur cluster modulates DNA binding. Nat
Commun. 2017;8:15052.

17.   Filenko N, Spiro S, Browning D, Squire D, Overton T, Cole J, et al. The NsrR regulon of
Escherichia coli K-12 includes genes encoding the hybrid cluster protein and the periplasmic,
respiratory nitrite reductase. J Bacteriol. 2007;189(12):4410-7.

18.   Partridge J, Bodenmiller D, Humphrys M, Spiro S. NsrR targets in the *Escherichia coli*
genome: new insights into DNA sequence requirements for binding and a role for NsrR in the
regulation of motility. Mol Microbiol. 2009;73(4):680-94.

19.   Robinson J, Brynildsen M. A kinetic platform to determine the fate of nitric oxide in
Escherichia coli. PLoS Comput Biol. 2013;9(5):e103049.

20.   Poole R, Anjum M, Membrillo-Hernández J, Kim S, Hughes M, Stewart V. Nitric oxide,
nitrite, and Fnr regulation of *hmp* (flavohemoglobin) gene expression in *Escherichia coli* K-12. J
Bacteriol. 1996;178(18):5487-92.

21.   Justino M, Almeida C, Gonçalves V, Teixeira M, Saraiva L. *Escherichia coli* YtfE is a
di-iron protein with an important function in assembly of iron-sulphur clusters. FEMS Microbiol
Lett. 2006;257(2):278-84.

22.   Todorovic S, Justino M, Wellenreuther G, Hildebrandt P, Murgida D, Meyer-Klaucke W,
et al. Iron-sulfur repair YtfE protein from *Escherichia coli*: structural characterization of the di-
iron center. J Biol Inorg Chem. 2008;13(5):765-70.

23.   Miller H, Kwuan L, Schwiesow L, Bernick D, Mettert E, Ramirez H, et al. IscR is
essential for *Yersinia pseudotuberculosis* type III secretion and virulence. PLoS Pathog.
2014;10(6):e1004194.

24.   Miller H, Schwiesow L, Au-Yeung W, Auerbuch V. Hereditary hemochromatosis
predisposes mice to *Yersinia pseudotuberculosis* infection even in the absence of the type III
secretion system. Front Cell Infect Microbiol. 2016;6:69.

25.   Schwiesow L, Mettert E, Wei Y, Miller H, Herrera N, Balderas D, et al. Control of *hmu*
heme uptake genes in *Yersinia pseudotuberculosis* in response to iron sources. Front Cell Infect
Microbiol. 2018;8:47.

26.   Hubbert W, Petenyi C, Glasgow L, Uyeda C, Creighton S. *Yersinia pseudotuberculosis*
infection in the United States. Septicemia, appendicitis, and mesenteric lymphadenitis. Am J
Trop Med Hyg. 1971;20(5):679-84.

27.   Gurry J. Acute terminal ileitis and yersinia infection. Br Med J. 1974;2(5913):264-6.

28.   Capron J, Delamarre J, Delcenserie R, Gineston J, Dupas J, Lorriaux A. Liver absess
complicating *Yersinia pseudotuberculosis* ileitis. Gastroenterology. 1981;81(1):150-2.

29.   Paff J, Triplett D, Saari T. Clinical and laboratory aspects of *Yersinia pseudotuberculosis*
infections, with a report of two cases. Am J Clin Pathol. 1976;66(1):101-10.

30.   Barnes P, Bergman M, Mecsas J, Isberg R. *Yersinia pseudotuberculosis* disseminates
directly from a replicating bacterial pool in the intestine. J Exp Med. 2006;203(6):1591-601.

31.   Crimmins G, Mohammadi S, Green E, Bergman M, Isberg R, Mecsas J. Identification of
MrtAB, an ABC transporter specifically required for *Yersinia pseudotuberculosis* to colonize the
mesenteric lymph nodes. PLoS Pathog. 2012;8(8):e1002828.

32.   Davis K, Mohammadi S, Isberg R. Community behavior and spatial regulation within a
bacterial microcolony in deep tissue sites serves to protect against host attack. Cell Host
Microbe. 2015;17(1):21-31.

33.   Zhang Y, Khairallah C, Sheridan B, van der Velden A, Bliska J. CCR2+ inflammatory
monocytes are recruited to *Yersinia pseudotuberculosis* pyogranulomas and dictate adaptive
responses at the expense of innate immunity during oral infection. Infect Immun.
2018;86(3):e00782-17.

34.   Peterson L, Philip N, DeLaney A, Wynosky-Dolfi M, Asklof K, Gray F, et al. RIPK1-
dependent apoptosis bypasses pathogen blockage of innate signaling to promote immune
defense. J Exp Med. 2017;214(11):3171-82.

35.   Durand E, Maldonado-Arocho F, Castillo C, Walsh R, Mecsas J. The presence of
professional phagocytes dictates the number of host cells targeted for Yop translocation during
infection. Cell Microbiol. 2010;12(8):1064-82.

36.   Sebbane F, Lemaître N, Sturdevant D, Rebeil R, Virtaneva K, Porcella S, et al. Adaptive
response of *Yersinia pestis* to extracellular effectors of innate immunity during bubonic plague.
Proc Natl Acad Sci. 2006;103(31):11766-71.

37.   Nuss A, Beckstette M, Pimenova M, Schmühl C, Opitz W, Pisano F, et al. Tissue dual
RNA-seq allows fast discovery of infection-specific functions and riboregulators shaping host-
pathogen transcriptomes. Proc Natl Acad Sci. 2017;114(5):E791-E800.

38.   Karlinsey J, Bang I, Becker L, Frawley E, Porwollik S, Robbins H, et al. The NsrR
regulon in nitrosative stress resistance of *Salmonella enterica* serovar Typhimurium. Mol
Microbiol. 2012;85(6):1179-93.

39.   Shepherd M, Achard M, Idris A, Totsika M, Phan M, Peters K, et al. The cytochrome bd-
I respiratory oxidase augments survival of multidrug-resistant *Escherichia coli* during infection.
Sci Rep. 2016;6:35285.

40.   Shiloh M, MacMicking J, Nicholson S, Brause J, Potter S, Marino M, et al. Phenotype of
mice and macrophages deficient in both phagocyte oxidase and inducible nitric oxide synthase.
Immunity. 1999;10(1):29-38.

41.   Vasquez-Torres A, Jones-Carson J, Mastroeni P, Ischiropoulos H, Fang F. Antimicrobial
actions of the NADPH phagocyte oxidase and inducible nitric oxide synthase in experimental

salmonellosis. I. Effects on microbial killing by activated peritoneal macrophages in vitro. J Exp
Med. 2000;192(2):227-36.

42.   Shiloh M, Nathan C. Reactive nitrogen intermediates and the pathogenesis of *Salmonella*
and *Mycobacteria*. Curr Opin Microbiol. 2000;3(1):35-42.

43.   Bang I, Liu L, Vasquez-Torres A, Crouch M, Stamler J, Fang F. Maintenance of nitric
oxide and redox homeostasis by the *Salmonella* flavohemoglobin *hmp*. J Biol Chem.
2006;281(38):28039-47.

44.   Mashruwala A, Boyd J. The *Staphylococcus aureus* SrrAB regulatory system modulates
hydrogen peroxide resistance factors, which imparts protection to aconitase during aerobic
growth. PLoS ONE. 2017;12(1):e0170283.

45.   Kinkel T, Roux C, Dunman P, Fang F. The *Staphylococcus aureus* SrrAB two-
component system promotes resistance to nitrosative stress and hypoxia. MBio.
2013;4(6):e00696-13.

46.   Jang S, Imlay J. Micromolar intracellular hydrogen peroxide disrupts metabolism by
damaging iron-sulfur enzymes. J Biol Chem. 2007;282(2):929-37.

47.   Justino M, Almeida C, Teixeira M, Saraiva L. *Escherichia coli* di-iron YtfE protein is
necessary for the repair of stress-damaged iron-sulfur clusters. J Biol Chem.
2007;282(14):10352-9.

48.   Velayudhan J, Karlinsey J, Frawley E, Becker L, Nartea M, Fang F. Distinct roles of the
Salmonella enterica serovar Typhimurium CyaY and YggX proteins in the biosynthesis and
repair of iron-sulfur clusters. Infect Immun. 2014;82(4):1390-401.

49.   Silva L, Baptista J, Batley C, Andrews S, Saraiva L. The di-iron RIC protein (YtfE) of
Escherichia coli interacts with the DNA-binding protein from starved cells (Dps) to diminish
RIC protein-mediated redox stress. J Bacteriol. 2018;200(24):e00527-18.

50.   Djaman O, Outten F, Imlay J. Repair of oxidized iron-sulfur clusters in *Escherchia coli*. J
Biol Chem. 2004;279(43):44590-99.
574

A

Competitive Index

* p=0.0117

WT + Δ*ytfE*

B

Microcolony area (μm²)

*

WT   Δ*ytfE*

C

hmp (mCherry)/GFP ratios

***   ***

centroid periphery centroid periphery

WT   Δ*ytfE*

D

Δ*ytfE*
gfp
hmp::mCherry
Hoechst

20 μm

Δ*ytfE*
hmp::mCherry

A

Competitive Index

WT + *ytfE* rescue

n.s. (p=0.922)

B

Microcolony areas (µm²)

n.s.

WT   *ytfE* rescue

C

WT

ytfE rescue