Characterization of coral-associated microbial aggregates (CAMAs) within tissues

of the coral *Acropora hyacinthus*

Naohisa Wada[1,2,7], Mizuki Ishimochi[2,3], Taeko Matsui[2], F. Joseph Pollock[4,7,8], Sen-Lin Tang[1],

Tracy D. Ainsworth[5], Bette L. Willis[6,7], Nobuhiro Mano[2, †], David G. Bourne[6,7,9 †]

[1]Biodiversity Research Center, Academia Sinica, Nangang, Taipei 11529, Taiwan

[2]Department of Marine Science and Resources, College of Bioresource Science, Nihon

University, Fujisawa, Kanagawa 252-0813, Japan

[3]Faculty of Science, University of the Ryukyus, Nishihara, Okinawa 903-0213, Japan

[4]The Nature Conservancy, Caribbean Division, Coral Gables, FL 33134

[5]School of Biological, Earth and Environmental Science, University of New South Wales,

Sydney, NSW 2052, Australia

[6]College of Science and Engineering, James Cook University, Townsville, QLD 4811, Australia

[7]AIMS@JCU, Townsville, QLD 4811, Australia

[8]Department of Biology, The Pennsylvania State University, State College, PA 16802, USA

[9]Australian Institute of Marine Science, Townsville, QLD 4810, Australia

†Equal correspond authors

Dr. Nobuhiro Mano (mano.nobuhiro@nihon-u.ac.jp) Department of Marine Science and

Resources, College of Bioresource Science, Nihon University, Fujisawa, Kanagawa, 252-

0813 Japan. Phone +81 466 84 3357

Dr. David G. Bourne (david.bourne@jcu.edu.au) College of Science and Engineering, James

Cook University, Townsville, QLD, 4811 Australia. Phone +61 7 47814790

Abstract

Bacterial diversity associated with corals has been studied extensively,

however, localization of bacterial associations within the holobiont is still poorly

resolved. Here we provide novel insight into the localization of coral-associated

microbial aggregates (CAMAs) within tissues of the coral *Acropora hyacinthus*. In

total, 318 and 308 CAMAs were characterized via histological and fluorescent *in situ*

hybridization (FISH) approaches respectively, and shown to be distributed extensively

throughout coral tissues collected from five sites in Japan and Australia. The densities

of CAMAs within the tissues were negatively correlated with the distance from the

coastline (i.e. lowest densities at offshore sites). CAMAs were randomly distributed

across the six coral tissue regions investigated. Within each CAMA, bacterial cells had

similar morphological characteristics, but bacterial morphologies varied among

CAMAs, with at least five distinct types identified. Identifying the location of

microorganisms associated with the coral host is a prerequisite for understanding their

contributions to fitness. Localization of tissue-specific communities housed within

CAMAs is particularly important, as these communities are potentially important

contributors to vital metabolic functions of the holobiont.

Introduction

Scleractinian corals associate with broad consortium of microorganisms, including endosymbiont dinoflagellates (Symbiodiniaceae), protozoa, fungi, bacteria, archaea and viruses, which collectively are termed the coral holobiont [1–3]. The importance of symbiotic dinoflagellates in provisioning the coral host with essential nutrients through translocated photosynthates has been established [e.g. 4,5], however the roles of other microorganisms within the holobiont are less well understood (reviewed in[3]). Some of the functions attributed to coral-associated microbiota include supply of essential nutrients and vitamins through processes such as nitrogen fixation [6–9] and metabolizing dimethylsulfoniopropionate (DMSP) to produce biologically important byproducts like dimethylsulfide [10]. The coral microbiota is also likely important for directly facilitating disease resistance through production of antimicrobials [11,12] or indirectly by preventing colonization of opportunistic or pathogenic organisms [13,14].

Corals are considered simple metazoans, but despite their basal phylogenetic position, they nevertheless form complex three-dimensional structures. Anatomically, the coral consists of (1) a surface mucus layer, (2) polyps consisting of feeding tentacles, actinopharynx, mesenteries (including their filaments), and walls (surface and basal), (3) a gastrovascular system that includes the gastrovascular cavity (formerly coelenteron) and connecting canals and (4) an external calcium carbonate skeleton [15,72]. The coral tissue layers are composed of epithelia, calicodermis (formerly calicoblastic epithelium) and the gastrodermis, which also contains the Symbiodinaceae. Within all these microhabitat niches, bacteria can reside as either transient communities or established symbionts with functional roles that may be positive, neutral or negative to the coral holobiont [16]. A multitude of studies have reported on the diversity of the coral-associated microbial communities, in some cases finding conserved microbial communities associated with some coral species, and in others finding shifting microbial profiles that reflect varying geographic, temporal or health status patterns [17–20]. To understand the significance of coral-microbial associations, care must be

exercised so that diversity patterns reflect the specific ecological niche that these

communities inhabit, such as the surface mucus layer, tissue layers, and/or the skeleton

[21–28]. Defining the locations of specific microorganisms is essential for elucidating the

importance of their role within the holobiont. For example, mucus bacteria are more

likely to have a loose association with the coral host, being sloughed off as the mucus is

exuded from the corals [29]. Conversely tissue-associated microorganisms are potentially

more integrated in shared metabolic pathways and may reside in specific associations

with host coral as a consequence of potential host selection [6,20,30].

To date, few studies have precisely localized bacterial communities within

coral tissues. Studies that have focused on localization often find that bacterial

communities within coral cell layers (i.e., epidermal and gastrodermal epithelia) form

aggregations termed coral-associated microbial aggregates (CAMAs) [31–34]. CAMAs

were first reported as potential pathogens when observed within healthy tissues of

Caribbean corals displaying signs of white-band disease [35,36]. Further studies

subsequently reported that CAMAs are widespread in tissues of healthy corals sampled

from geographically dispersed areas [32,36,37]. To date, CAMAs have been reported from 5

species of corals in the Caribbean [35,36] and 24 species from the Indo-Pacific [32], although

their frequency varies among coral genera, with the genera *Acropora*, *Porites*, and

Pocillopora most commonly hosting bacterial aggregates [32]. Identification of the

microorganisms that constitute these CAMAs has been poorly resolved. Neave et al.

(2016) visualized aggregates (i.e., "cyst-like aggregations") of *Endozoicomonas* within

tissues of *Stylophora pistillata* at the interface of the epidermis and gastrodermis and

confirmed their distribution in samples taken from widely-separated biogeographic

regions [38,39]. However, the fine-scale spatial distributions of microorganisms and

potential microhabitat-associated structure of the microbiome still have not been

clarified. To facilitate a more comprehensive understanding of the roles bacterial

communities may play in the coral holobiont, an improved understanding of the

localization of these microbial communities is essential, particularly tissue-associated

bacterial communities housed within structures termed coral-associated microbial

aggregates (CAMAs). Here, we visualize the localization, distribution and morphology

of CAMAs associated with the coral *Acropora hyacinthus* sampled from Sesoko Island,

Okinawa, Japan and sites in the northern and central regions of the Great Barrier Reef

(GBR) Australia, located along an inshore to offshore gradient.

Results and Discussion

Comparison of CAMA prevalence among geographic locations

At the time of field collection, all 48 colonies of *A. hyacinthus* sampled (one

fragment from each colony), from the 5 geographic locations (see **Fig. 1**), appeared

visually healthy. This was confirmed by subsequent histological analyses, which found

that all tissues displayed normal cell morphology, including no signs of fragmentation,

wound repair or necrosis (as per criteria in [40]). In total, 318 CAMAs were characterized

via histology (**Fig. 2a**) with CAMAs detected in fragments from 27 of the 48 sampled

colonies (~56%). The vast majority stained basophilic (95.9%), compared to only 4.1 %

staining eosinophilic with the hematoxylin and eosin staining procedure. At the Sesoko

Island site, tissues derived from the fragments of all 10 sampled colonies contained

CAMAs (**Fig. 2c**), whereas at the other four sites, CAMAs were observed only in

tissues of some of the coral fragments. For example, CAMAs were clearly visible in

80% of Inner Shelf samples, 30% of Lizard Island, 25% of Outer Shelf, and 40% of

Orpheus Island samples (n=10 samples at all sites, except at the Outer Shelf site where

n=8 samples). In general, the prevalence of CAMAs was significantly higher at the

Sesoko Island site than at the Lizard Island, Outer Shelf, and Orpheus Island sites

(Fisher's exact   test: $p = 0.00099$, **Fig. 2c,** and see more detail of the pairwise test in

Suppl. table S1).

Fig. 1 Map showing the locations of five study sites in two countries: (a, b) Sesoko Island (SI) in Okinawa, Japan; (a, c) Inner Shelf (IS), Lizard Island (LI) and Outer Shelf (OS) sites in the Northern Great Barrier Reef; and (a, d) Orpheus Island (OI) in the central Great Barrier Reef, Australia.

Fig. 2 Appearance, occurrence of CAMAs in the coral *Acropora hyacinthus*. (**a**) Numerous CAMAs (indicated by arrows) are visible in a histological section stained by hematoxylin and eosin of a colony from Sesoko, Japan. In the panel, small photo shows close-up of a CAMA located in a mesentery of a polyp. (**b**) Right panel shows two of CAMAs (red) distributed with Symbiodiniaceae (green) in the tentacle (coral tissue: blue) of a sample from the Inner Shelf, GBR using FISH. (**c, d**) Pie diagrams showing the proportion of colonies sampled that contained CAMAs at five sites in Japan and Australia based on the detections of (**c**) HE stain and (**d**) FISH. (**c**) A significantly higher proportion of Sesoko Is. samples contained CAMAs (including basophilic and eosinophilic) than samples collected from other sites (Lizard Is., Outer Shelf northern GBR site, and Orpheus Is.). (**d**) In the FISH experiment, both proportions of CAMA in samples from Sesoko Is. and Inner Shelf were also significantly higher than in samples from other three sites (** $p<0001$; Fisher's exact test followed by a Benjamini–Hochberg false discovery rate correction). Scale bars indicate 600 µm (**a**), 50 µm (**small panel in a**) and 100 µm (**b**).

A total of 308 CAMAs were also identified by FISH in the serial-sectioned

tissues derived from the sampled colonies (again observed in 27 fragments from 48

sampled colonies) (**Fig. 2b**). CAMAs were observed from all samples derived from

Sesoko Island and the Inner Shelf (**Fig. 2d**). In contrast CAMAs were observed in 30%

of Lizard Island, 25% of Outer Shelf, and 20% of Orpheus Island samples. The

prevalence of CAMAs detected by FISH was significantly higher at both Sesoko Island

and Inner Shelf than at the other sites (Lizard Island, Outer Shelf, and Orpheus Island

sites; Fisher's exact test: $p = 0.0000019$, **Fig. 2d** and see more detail of the pairwise

test in **Suppl. table S1**).

The density of basophilic (i.e. stained with hematoxylin) CAMAs in tissues

was highest in the Sesoko Island samples (n=10) with 20.13 ± 17.1 per cm^2 (average

and S.E.), compared to 6.78 ± 9.3 per cm^2 for the Inner Shelf (n=8) samples, 0.92 ± 0.1

per cm^2 at Lizard Island (n=3) and 3.55 ± 3.3 per cm^2 in Orpheus Island (n=3) samples

(see more detail in **Suppl. table S2**). Basophilic CAMAs were not detected in samples

of the Outer Shelf corals. Eosinophilic (i.e., stained by eosin) CAMAs were only

detected in coral fragments derived from three sites, with 2.32, 5.39 ± 6.7, and $1.28 \pm$

2.8 per cm^2 in Inner Shelf (n=1), Outer Shelf (n=3) and Orpheus Island (n=2) samples,

respectively (see more detail in **Suppl. table S2**). The densities of CAMAs detected by

FISH were reflective of the histological patterns with 18.72 ± 12.7 per cm^2 in Sesoko

Island (n=10), 7.90 ± 8.2 per cm^2 in Inner Shelf (n=10), 1.26 ± 0.6 per cm^2 in Lizard

Island (n=3), 0.90 ± 0.5 per cm^2 in Outer Shelf (n=2), and 4.48 ± 4.2 per cm^2 in

Orpheus Island (n=2) samples (see more detail in **Suppl. table S2**). In one sample from

Sesoko Island, 48.9 basophilic CAMAS were detected per cm^2 of tissue (41.1 detected

by FISH in the serial section), the greatest density of CAMAs observed across all 48

samples investigated. The maximum number observed in a sample from the Outer Shelf

region was 10.10 eosinophilic CAMAS per $cm^{2.}$

156   The patterns of CAMAs prevalence derived from histological and FISH

analysis confirm that CAMAs commonly occur in healthy tissues of the coral *Acropora*

hyacinthus collected from sites in Japan and Australia separated by more than 40

degrees of latitude. Although CAMAs were not detected in all tissue samples collected,

this may reflect limitations in the area of coral tissue that can be surveyed via

histological approaches. The presence of CAMAs in a histological section will depend

on the tissue sectioned (i.e., the location of the fragment on the colony and on the

section from that fragment), and on the scale and orientation of the section. Indeed, our

results showed variation of both presence and density of CAMAs, even from samples

derived from the same site (see more detail in **Suppl. table. S2**). Therefore, although

not all coral tissue samples (and therefore not all colonies) were found to host CAMAs,

we cannot exclude the possibility that other tissue areas of the same colony had CAMAs

present. Other studies have also reported that CAMAs are common in tissues of many

coral species in the Caribbean [35,36], Indo-Pacific and Red Sea [32,33,37,38]. In particular,

CAMAs were common in species of *Acropora*, *Porites*, and *Pocillopora*, although often

their presence was patchy within a population sample [32].

172   Interestingly the distance from the coast negatively correlated with the

proportion and density of CAMAs within coral tissues across the sampling sites (**Table**

1). A predictive model was constructed which yielded an ordinary negative binomial

regression which was found preferable to a zero-inflated model (Vuong test; Voung z-

statistic= 22.57413, *p*-value < 0.001 in the basophilic CAMAs, and Voung z-statistic=

3.120e-6, *p*<0.001 in CAMAs detected by FISH). Therefore an increased distance from

the coastline influenced negatively the proportion and density of both basophilic

CAMAs and CAMAs detected by FISH (**Fig. 3a, c**). The density of eosinophilic

CAMAs did not follow the negative binominal regression model (**Table 1, Fig. 3b**).

Although our study provides only a snapshot of five sites sampled at one time point, it

suggests that inshore reef environments may promote the development of CAMAs

within coral tissues. The inshore site at Sesoko Island has high nutrient influxes,

especially phosphates [41,42]. Similarly, nearshore GBR sites are influenced by influxes of

dissolved nutrients from terrestrial runoff, with higher concentrations typically found at

inshore compared with offshore sites [43]. Temperature fluctuations may also influence

CAMA abundance, although differences in seasonal temperature fluctuations are

minimal across the northern GBR sites [44]. The coral microbiome community has been

shown to shift in response to environmental stressors [29,45–49], thus water quality

parameters influencing microbiological composition and function [50] could stimulate

CAMA abundance. Further studies, particularly of potential links between nutrient

levels and CAMA development, are needed to understand what might drive the

increased prevalence and density of CAMAs in coral tissues at inshore sites, and to

determine if hosting more CAMAs is beneficial to corals or is an indicator of negative

impacts on the coral holobiont.

Table 1 Results from the generalized linear regression models with negative binominal distribution testing the effect of the distance

from closest coastline on the densities of CAMAs

		Estimate	SE	Wald chi-square (z-value)	p-value (z)	AIC	LR (X^2)	p-value (LR)
Basophilic CAMAS	Intercept	3.53176	0.399923	8.846	<2e-16 ***	187.52	67.427	2.2e-16 ***
	Distance[†]	-0.17477	0.02769	-6.312	2.76e-10 ***			
Eosinophilic CAMAS	Intercept	-3.508	1.3253	-2.647	0.00812**	46.991		
	Distance[†]	0.077	0.0459	1.678	0.09643			
CAMAs detected by FISH	Intercept	3.19428	0.35556	8.984	<2e-16 ***	212.09	51.662	6.594e-13***
	Distance[†]	-0.12714	0.02092	-6.077	1.22e-09 ***			

198 [†]The distance indicates distance to closest coastline.

199 P values are **$p < 0.01$ and ***$p < 0.001$.

200 AIC and LR indicates Akaike's Information Criterion and likelihood-ratio.

Fig. 3 Relationship between densities of CAMAs and distance to the coast based on a negative binominal regression model. (a) Density of basophilic CAMAs was negatively correlated with the distance from the coast to the offshore sampling sites. **(b)** No correlation of eosinophilic CAMAs were detected in relation to the distance to the coast, although higher density (max. 10.1 cm^2) was found in the Outer Shelf (offshore). **(c)** Density of the CAMAs detected by FISH was also negatively associated with the distance. Model prediction and associated 95% confidence intervals were obtained from a negative binominal regression model between the densities of CAMAs and the distance to the coastline **(a, c)**. Scale bars indicate 10 μm.

Distribution of CAMAs within anatomical regions of the coral polyp

CAMAs were localized (by FISH) within the six anatomical regions of the

coral polyp: the tentacle, actinopharynx, mesenterymesenterial filament, coenenchyme,

and calicodermis (see **Fig. 4a**). The CAMAs appeared to be randomly distributed across

the anatomical regions investigated, though because the numbers of CAMAs

characterized for the Lizard Island, Outer Shelf and Orpheus Island samples were low

(**Fig. 4b**), meaningful comparisons can only be made between the Sesoko Island and

Northern Inner Shelf samples. For the Sesoko Island samples, CAMAs were found

predominantly in the tentacles (36.5%), mesenterial filaments (34.7%) and

coenenchyme (16.5%) regions; in Inner Shelf samples from Northern GBR sites,

CAMAs were mostly located in the calicodermis (46.8%), tentacles (17.5%) and

coenenchyme (16.7%) (**Fig. 4b**). An additional 31 CAMA-like shaped structures were

observed from three samples derived from Sesoko Island, but these were discounted as

bacterial aggregates due to non-probe binding signals and excluded from the analysis.

CAMAs spanned a wide size range, from 23 to 6,761 μm^2 across tissue

samples from all sites (**Fig. 4c**). Measurements likely underestimated the size of

CAMAs, given they are dependent on the orientation of sectioning and it is unlikely that

most CAMAs were sectioned through their greatest diameter. Acknowledging

constraints associated with sectioning, the average size of CAMAs was 1,304 μm^2.

Sesoko Island samples contained generally larger average and median-sized aggregates

(1,507.7±1,522.4 μm^2, 921.3 μm^2 respectively) than samples from the GBR region

(Inner Shelf: 637.4±734 μm^2, 321.6 μm^2). However, given the large range in sizes

measured, the fact that only two locations had sufficient sample sizes for comparison,

the issue of sectioning orientation potentially biasing size measurements, and high

variation among the individual colony level (**Suppl. fig. S1**), such patterns require

further validation. No patterns in the size of CAMAs across different anatomical regions

were detected (**Fig. 4c**).

Fig. 4 Distribution of CAMAs within six anatomical regions of the coral *Acropora hyacinthus* collected from five sites in Japan and Australia. (a) Schematic drawing of a coral polyp showing the six anatomical regions examined microscopically (same color coding used in **(b)** and **(c)**. **(b)** Pie charts showing the distribution of CAMAs among six anatomical regions in samples from the five sites. **(c)** Dot plots comparing the size of CAMAs among anatomical regions for each location. N=308 CAMAs detected in tissues treated with FISH. In the dot plot, the lines indicate median (thick line), 3[rd] (upper) and 1[st] (lower) quartiles (long thin lines), whiskers (short thin lines; upper value indicates largest value within 1.5* interquartile range from 3[rd] quartile, lower: min value).

Morphology of CAMAs within coral tissues

High resolution imaging was used to partially characterize and compare the

morphology of bacteria across the CAMAs detected (**Fig. 5**). Interestingly, each CAMA

appeared to be composed of a single morphological type of bacteria, although

morphological types varied among CAMAs. Overall, five different morphological types

of bacteria were identified: rod-shaped (length 2.5±0.1 µm, width 0.6±0.0 µm, **Fig. 5a,**

e), an atypical coccus (length 4.8±0.3 µm, width 3.1±0.2 µm, **Fig. 5b, f**), a longer rod

morphology (length 8.0±0.0 µm, width 0.8±0.0 µm, **Fig. 5c, g**), filamentous-like

bacteria (length N.D., width 0.4±0.0 µm, **Fig. 5d, h**), and a rod-shaped morphology but

with spore-like structures (**Fig. 5i–j**, **m–n**). While the consistency of morphological

characteristics within each CAMA may indicate that CAMAs are hosting single

bacterial types, it is also possible that they host multiple bacterial species with similar

morphologies. Interestingly, the fluorescent signal detected for some CAMAs was not

uniform over the entire aggregation. The lack of signal within some CAMAs (see **Fig.**

5d for example) may be due to the probe not targeting microbial cells that inhabit that

space. Defining a specific morphological shape for the bacterial cells observed was

sometimes difficult, with patchy probe hybridization producing images of structures that

potentially protruded from the tissue sections and seemed amorphous (**Fig. 5k–l, o–p**).

However, this again could be the result of mixed microbial communities within the

CAMAs, with some cells targeted by the probes, but others not hybridizing to the

probe-fluorochrome conjugate.

Fig. 5 Morphological variation of bacteria housed within different CAMAs. Note that bacteria within each CAMA are morphologically similar. Dotted lines (in **a–d** and **i–l**) delineate regions magnified in close-up images (**e–h** and **m–p**). Overall, five bacterial morphologies were detected: rod-like (**a**, **e**), pleomorphic (**b**, **f**), long rods (**c**, **g**), filamentous-like (**d**, **h**), rod-shaped with spore-like structures (**i–j** and **m–n**), and putative amorphous masses (k–l and o–p). Scale bars indicate 10 µm (**a–d** and **i–l**) and 5 µm (**e–h** and **m–p**).

The variation in bacterial cell morphology among CAMAs may contribute to

the different HE staining properties observed, with the majority being basophilic

(95.9%), but 4.1% staining eosinophilic (n=318 CAMAs detected in total). Variability

in the staining of CAMAs has been reported previously [32,51] and attributed to varying

degrees of protein and DNA production or local tissue pH conditions [32]. However, these

staining patterns may also be influence by different bacteria housed within the CAMAs.

Previous studies have reported that the abundant coral-associated bacterial genera

Endozoicomonas forms aggregates within tissues of the corals *Stylophora pistillata* and

Pocillopora verrucosa [38,39]. Indeed, extensive sequence-based phylogenetic surveys of

coral microbiomes have revealed that several dominant bacterial groups are common [20],

including the *Proteobacteria* (particularly *Alpha-* and *Gammaproteobacteria* inclusive

of *Endozoicomonas*), as well as *Actinobacteria*, *Bacteroidetes* (especially

Flavobacteria), and *Cyanobacteria* [3]. Hence some of the different cellular morphologies

we detected within CAMAs may represent common coral-associated bacterial groups

profiled in microbiome diversity studies. We note, however, that bacterial morphology

can be plastic and dependent on numerous variables, such as division stage,

colonization, chemical environment, physical constraints and nutrient availability [52].

Future work targeting CAMAs with taxa-specific probes and gene-sequencing

approaches would help to resolve both the taxonomy, composition and potential

functional roles of bacteria within these structures of the coral holobiont.

FISH imaging showed that for some of the CAMAs, patchy probe-specific

labeling occurred, resulting in dull or dark areas within the structures. There are a

number of potential methodological reasons for this observation, including: 1)

insufficient probe sensitivity due to low ribosomal rRNA content in target cells [53,54], 2)

methodological and environmental factors that prevent probes from accessing target

cellular rRNA at these sites [55], 3) the bacterial community penetrating and proliferating

within the epidermis of coral tissues [37], or 4) lipid or fat solvents deposited through

dehydration and dewaxing steps showing empty spaces in the tissues [56]. Alternatively,

little to no signal in central regions of some of the CAMAs could indicate that the probe

EUB338mix did not target the taxonomic group of microbes within these regions. The

EUB338mix is estimated to cover 96% of the *Eubacteria* domain, but taxa outside this

coverage, including archaeal lineages, may be present [57]. We speculate that some of the

CAMAs may be mixed communities containing bacteria not targeted by the probes or

even *Archaea,* which have been identified to associate with corals in microbiome

diversity studies [58]. A recent coral metagenomic study recovered *Thermarchaeota*

genome bins from a *Porites* sp. that was potentially metabolically linked through

nitrogen cycling to other coral microbial-associated taxa, including *Nitrospira* [6]. Co-

aggregation of ammonia-oxidizing archaea with nitrite-oxidizing bacteria is common in

other organisms, such as sponges [59,60].

Further three-dimensional reconstructions of z-stacked FISH images of

select 100 µm stained sections of tentacles visualized the CAMAs as typically spheroid

or ellipsoid-shaped structures (**Fig. 6a**). In one example, a single CAMA was located

independently in the epidermis of a tentacle (**Fig. 6b**). In a different tentacle, multiple

smaller CAMAs were localized close to Symbiodiniaceae cells in the gastrodermis (**Fig.**

6c). Sizes of the large single CAMA and the multiple smaller CAMAs were 33,400 µm^3

(**Fig. 6d**, surface area 7,729 µm^2) and 1,978 ± 141.2 µm^3 (**Fig. 6e**, 809.5 ± 59.9 µm^2,

n=4 CAMAs), respectively. Even though the CAMAs were located in the same polyp,

the size of the single large CAMA was approximately 40-fold greater than the multiple

smaller CAMAs, demonstrating inherent size variability for these structures. The

bacteria within these CAMAs displayed a similar rod-shaped morphology (see **Fig.5a**

and e), with the average cross-sectional area of each bacterium being 175.7 µm^2 and

22.6 ± 4.2 µm^2, respectively (**Suppl. fig. S2**). Based on cell size, the number of

individual rod-shaped bacterial cells within the 3D rendered images of these CAMAs

was estimated as ~ 47,275 cells for the large single CAMA located in the epithelium of

the tentacle (**Fig. 6d** and **Suppl. fig. S2a**), and 2,799 ± 200 cells (**Fig. 6e** and **Suppl. fig.**

S2b) for each of the smaller CAMAs localized close to Symbiodiniaceae cells.

Fig. 6 Three-dimensional (3D) images of CAMAs (red) within a tentacle of the coral *Acropora hyacinthus*, as visualized using FISH. (**a**) Section of tentacle showing localization of two types of CAMAs (10x magnification). (**b**) Single aggregation of bacteria in a large structure within the epithelium (40x magnification; composed of 92 z-stack images). (**c**) Multiple aggregations of bacteria in smaller structures within the gastrodermis (40x magnification; composed of 56 z-stack images). 3D rendering of CAMAs (**d**, **e**) reconstructed from 3D images in **b** and **c**. Coral tissue and Symbiodiniaceae appear as blue and green structures, respectively.

Using these estimates of cell numbers within CAMAs we further

extrapolated that bacterial densities in tissues from Sesoko Island and Inner Shelf GBR

corals are approximately 5.6×10^4 and 1.5×10^4 cells per cm^2 (along a linear cross-

section), respectively. Very few studies have accurately determined bacterial cell

densities associated with corals. Counts for bacteria in the coral mucus layer can be as

high as 10^7 cells ml^{-1} [61]. Estimates of bacteria on tissue surfaces range from 1×10^5 to

10^6 cells per cm^2 for the coral *Pocillopora damicornis* [62] and 8.3×10^6 to 6.2×10^7 cells

per cm^2 for the coral *Oculina patagonica* [24]. However, these counts are based on

bacteria external to coral tissues and therefore not directly comparable to estimates from

our 3D reconstructions of CAMAs visualized within coral tissues. Our results highlight

that tissue-associated microbial communities are present across the different anatomical

regions of the coral polyp and that differentiating these communities from external and

mucus-associated microbial communities is important for accurate appraisals of the

coral microbiome and for identifying their role(s) within the coral holobiont.

Conclusion

Localization of microorganisms associated with corals is vital to understand

their symbiotic relationships and reveal their function(s) within the holobiont. Here we

provide novel insight into the distributions and densities of bacteria within tissues of the

coral *Acropora hyacinthus* sampled from five different locations. CAMAs were

common in coral tissues sampled, although their prevalence were higher in colonies

sampled close to shore compared to offshore geographic sites, potentially linked to

water quality parameters. While each CAMA appeared to be dominated by a single

bacterial morphological type, different CAMA hosted different bacterial morphotypes.

CAMAs have been defined as facultative symbionts, not necessary for host fitness [32],

however their high prevalence and abundance in coral tissues may indicate they are

integrated into shared metabolic pathways and central to maintaining coral fitness

through provisioning benefits. For example, findings that CAMAs are often co-

localized near Symbiodiniaceae cells highlights the potential metabolic integrated links

between bacterial and Symbiodiniaceae symbionts. Such propositions require testing by

tracing metabolic pathways, as well as improved taxonomic and functional assessment

of the microorganisms housed within these CAMAs.

Methods

Sample collections

Fragments (~ 3 cm x 3 cm; n=48 colonies) of the tabulate coral *Acropora*

hyacinthus (Dana, 1846) were collected from five geographic locations across two

countries (Australia and Japan). Samples from Japan (n=10) were collected from

Sesoko Island, Okinawa (26˚37'40.3"N, 127˚51'36.4"E, depth 1.5–3.0 m, distance from

coastline of Okinawa main Island 2.4 Km, average colony size 62.8 ± 41.7 cm) in July

2015. On the Great Barrier Reef (GBR) Australia, samples were collected from the

Northern sector along an inshore to offshore gradient inclusive of Lizard Island (see

Fig. 1; n=10 from an Inner Shelf site (14˚47'29.4"S, 145˚20'18.8"E, depth 3–5 m,

distance from coastline 12.5 Km, average colony size 77.69 ± 43.0 cm), n=10 from

Lizard Island (14˚41'48.75"S, 145˚27'49.22"E, depth 2–4 m, distance from coastline

28.6 Km, average colony size 61.67±17.5 cm) and n=8 from an Outer Shelf site

(14˚38'30.0"S, 145˚38'22.5"E, depth 2–5 m, distance from coastline 46.2 Km, average

colony size 41±20.5 cm) in January 2013. Samples (n =10) were also collected from

reefs around Orpheus Island (18˚35'55.4"S, 146˚29'33.8"E, depth 2–3 m, distance from

coastline 16.3 Km, colony size N.D.) in the inner-central region of the Great Barrier

Reef in March 2013.

Histological preparations

Following collection, samples were immediately rinsed with sterile seawater

and then fixed in 4% paraformaldehyde (Electron Microscopy, USA and Wako, Japan)

in 10mM phosphate buffered saline (PBS; pH 7.4) for 8–10 hours maintained at 4 ˚C.

Samples were subsequently rinsed twice with 70% ethanol and stored in 70% ethanol at

4 ˚C prior to decalcification. After samples were rinsed by PBS twice for 30 min each,

samples were decalcified at 4 ˚C in a 10% EDTA (sigma-Aldrich, USA) solution (w/v;

pH 8.0 adjusted by sodium hydroxide [Wako, Japan]), which was exchanged

approximately every two days until no coral skeleton remained (approx. two weeks).

The decalcified samples were rinsed in PBS, and dehydrated sequentially through 70%,

90%, and two changes in abs. 100% ethanol series (60 min each), then processed

through a 1:1 solution of abs. 100% ethanol and toluene, and two changes in toluene (30

min each), and embedded in paraffin. Nine sections (three serial sections x three sets;

interval = 100 µm between each set) of each coral fragment, each 4 µm thick, were cut

from each paraffin-embedded sample (**Suppl. fig. S3**). In addition, thicker 100 µm

sections were cut from coral tissues to allow reconstructions of three-dimensional

configurations of the CAMAs (see below). Serial sections were mounted on one slide

coated with egg-white glycerin and on two slides with poly-L-lysine solution (Sigma-

Aldrich, USA) for hematoxylin and eosin (HE) staining and fluorescence *in situ*

hybridization (FISH), respectively. We analyzed a total 144 sets (432 sections) of the

serial sections for HE staining and FISH among five location sites.

Hematoxylin and Eosin (HE) staining

For hematoxylin and eosin (HE) staining, one serial section from each set was

dewaxed in xylene (2 x 15 min), rehydrated through ethanol series with abs. 100%,

99%, 90% and 70% (5 min each) and rehydrated completely in sterile water. Hydrated

sections were stained in Mayer's hematoxylin (Wako, Japan) for 10 min, rinsed in water

for 5 min, then stained with eosin Y (Merck, Germany) for 5 min, and further rinsed in

water for 30 sec. The stained sections were dehydrated through the same ethanol series

in reverse with agitation (few sec each), cleared by xylene (2 x 5 min) and finally

mounted in Entellan mounting medium (Merck, Germany). HE stained sections were

observed and recorded using an ECLIPSE Ni microscope (Nikon, Japan) and

BIOREVO BZ–9000 microscope (KEYENCE, Japan). All sampled corals appeared

visually healthy at the time of collection and tissues displayed normal cell morphology,

including no signs of fragmentation, wound repair or necrosis (as per criteria in [40]). The

number of CAMA were counted into two categories, basophilic and eosinophilic

CAMAs. The area of whole tissue (including skeleton region) was measured by tracing

the outer edge of the tissue section using Fiji software [63].

Fluorescence in situ hybridization (FISH)

405   The other two serial sections from each set were subjected to FISH according

to the protocol detailed in Wada et al. (2016) [64]. Briefly, sections were dewaxed in

xylene (2 x 15 min), dehydrated briefly once in 100% ethanol and dried completely.

The dried sections were immersed in a 0.2 M HCl solution for 12 min, followed by a 20

mM Tris-HCl solution (pH 8.0) for 10 min at room temperature. The sections were

mounted with proteinase K (50 µg ml^{-1}) in 20 mM Tris-HCl solution at 37 ˚C for 5 min

for bacterial cell wall permeabilization, and washed in a 20 mM Tris-HCl solution at

room temperature for 10 min. Oligonucleotide probes, including a probe targeting the

16S rRNA gene (EUB338mix: 5'-GCWGCCWCCCGTAGGWGT-3') and a nonsense,

negative control probe (Non338: 5'- ACATCCTACGGGAGGC -3'), were labeled with

the Cy3 fluochrome (eurofins, USA) [65,66]. Tissue sections were covered with

hybridization buffer (30% v/v formamide, 0.9 M NaCl, 20 mM Tris-HCl [pH 8.0],

0.01% SDS), then each oligonucleotide probe was added to a final concentration of 25

ng µl^{-1} to each serial section. The slides were incubated at 46 ˚C for 1.5 hour. After

incubation, sections were washed in 50-ml falcon tubes containing preheated wash

buffer (0.112 M NaCl, 20 mM Tris HCl [pH 8.0], 0.01% SDS, 5 mM EDTA [pH 8.0])

in a water bath at 48 ˚C for 10 min, then soaked immediately with agitation in cold

water, and air dried completely. The dried sections were mounted in an antifade

mounting medium Fluoromount/Plus (Diagnostic BioSystems, USA). Sections were

examined and recorded using a FV1000-D confocal microscope (Olympus, Japan) with

two channels, using the following settings: (1) laser: 405 nm and 559 nm; (2) excitation

dichroic mirror: DM405/473/559; (3) emission dichroic mirror: SDM560 and mirror;

(4) band-pass filter: None and BA575–620 for detecting autofluorescence of coral tissue

(blue) and Cy3 signal (red), respectively. Non-specific probe binding in tissue sections

was identified as detailed by Wada et al. (2016)[64]. Specific detection of bacteria by

fluorescent *in situ* hybridization (FISH) can be problematic for coral samples due to

nonspecific binding of probes and background autoflorescence of granular cells and

nematocysts [64]. Therefore, CAMAs were identified by their characteristic shapes, in

addition to fluorescent signals bound by the general bacteria-targeted probe set

EUB338mix and comparisons to background autofluorescence and non-specific binding

(using Non338 probe). The number of CAMAs that hybridized with the EUB338mix

probe in each section was counted and the area of whole tissue (including skeleton

region) was measured by tracing the outer edge of the tissue section. We also traced the

edges of each CAMA to calculate the area (the size of CAMA was dependent on the

orientation of sectioning). Image processing was conducted in Fiji software [63].

Reconstruction of three-dimensional images of CAMAs

Three-dimensional images (3D) of CAMAs were reconstructed from the thick

sections (100 µm) which were cut carefully from two samples (S6 from the Sesoko

Island and LO8 from the Outer Shelf GBR site). Sections were visualized via FISH

according to methods described above, and examined using a LSM 880 (ZEISS,

Germany) with two tracks and the following settings: (1) laser: 405 nm and 561 nm; (2)

beam splitter: MBS -405 and MBS 488/561; (3) filter: 371–479 nm for auto-

fluorescence of coral tissue (blue) and 627–758 nm for Symbiodiniaceae (green) in

track 1 and 565–588 nm for Cy3 signals (red) in track 2, respectively. For

reconstructing the 3D images, the sections consisted of z-stack images at 3.0 µm

intervals for 10x and 0.7 µm each and 40x magnifications using the Z-stack function in

LSM 880. The z-stack images were processed and reconstructed with surface rending of

the Cy3 signals in Imaris software ver. 8.0.2 (BitplaneAG, USA).

Statistical analysis

All statistical analyses were conducted using R Stats ver. 3.5.1 [67] with the following

packages: rcompanion ver. 2.0.0 [68], MASS 7.3-51.4 [69], pscl 1.5.2 [70] and car 3.0-2 [71]. To

compare the abundances of CAMAs at the colony level among sites, Fisher's exact

test followed by post hoc Benjamini–Hochberg false discovery rate correction were

used. A generalized linear model was used to evaluate the relationship between the

distance of samples from the coastline (explanatory variable) with the distribution and

density of CAMAs (count data of response variables: basophilic, eosinophilic and FISH

detected CAMAs; offset variable: tissue area) among the five sites. The tests were

performed to fit using the 'glm.nb' function (negative binomial distribution) in the

MASS package, due to over dispersions among the data. We also performed zero-

inflated negative binomial regression using the 'zeroinf' function in the pscl package,

since the count data included excessive zeros. We then performed a Vuong test that

compares an ordinary negative binomial regression model against the zero-inflated

model using the 'vuong' function in the pscl package, and adopted the model's fitness

based on AIC (Akaike's Information Criterion), and yielded the model, as appropriate.

Finally, the likelihood-ratio (LR) chi-square was calculated for each model using the car

package.

Author contribution

NW, NM and DGB conceived the study. NW, FJP and DGB designed the sampling and

FISH experiments. NW and FJP conducted the field sampling. NW, MI and TM

performed the histological work and all observations and data collection. NW and DGB

had major contribution in the manuscript writing and the figure making. FJP, S-LT,

TDA, BLW and NM contributed to writing and editing the manuscript. All authors

critically reviewed, revised and ultimately approved this final version. Finally the

authors confirm that there are no competing interests and no non-financial competing

interests that impact this manuscript

Acknowledgements

Dr. Joleah Lamb (University of California, Irvine), Mr. Sefano Katz (Pacific Blue

Foundation), Dr. Gergely Torda (ARC Center of Excellence for Coral Reef Studies,

James Cook University), and Dr. Melanie Trapon (CSIRO) are thanked for helping to

collect samples. We thank A/Prof Esther Peters from George Mason University for her

advice and help in updating coral anatomy terminology. We are also grateful to Mr.

Yasuhiko Sato from Zeiss Microscopy Japan, and Mr. Ji-Ying Huang and IPMB Live-

Cell-Imaging Core Lab in the Institute Academia Sinica for supporting the microscopy

and the Imaris software used.

References

1. Rohwer, F., Seguritan, V., Azam, F. & Knowlton, N. Diversity and distribution of coral-associated bacteria. *Mar. Ecol. Prog. Ser.* **243**, 1–10 (2002).

2. Rosenberg, E., Koren, O., Reshef, L., Efrony, R. & Zilber-Rosenberg, I. The role of microorganisms in coral health, disease and evolution. *Nat. Rev. Microbiol.* **5**, 355–62 (2007).

3. Bourne, D. G., Morrow, K. M. & Webster, N. S. Insights into the Coral Microbiome: Underpinning the Health and Resilience of Reef Ecosystems. *Annu. Rev. Microbiol.* **70**, 317–340 (2016).

4. Yellowlees, D., Rees, T. A. V. & Leggat, W. Metabolic interactions between algal symbionts and invertebrate hosts. *Plant Cell Environ.* **31**, 679–694 (2008).

5. Petrou, K., Nielsen, D. A. & Heraud, P. Single-Cell Biomolecular Analysis of Coral Algal Symbionts Reveals Opposing Metabolic Responses to Heat Stress and Expulsion. *Front. Mar. Sci.* **5**, (2018).

6. Robbins, S. J. *et al.* A genomic view of the coral holobiont. *Nat. Microbiol.* (in press).

7. Lema, K. A., Willis, B. L. & Bourne, D. G. Corals Form Characteristic Associations with Symbiotic Nitrogen-Fixing Bacteria. *Appl. Environ. Microbiol.* **78**, 3136–3144 (2012).

8. Pogoreutz, C. *et al.* Nitrogen Fixation Aligns with nifH Abundance and Expression in Two Coral Trophic Functional Groups. *Front. Microbiol.* **8**, (2017).

9. Yang, S.-H. *et al.* Metagenomic, phylogenetic, and functional characterization of predominant endolithic green sulfur bacteria in the coral Isopora palifera. *Microbiome* **7**, 3 (2019).

10. Raina, J.-B. *et al.* DMSP biosynthesis by an animal and its role in coral thermal stress response. *Nature* **502**, 677–680 (2013).

11. Raina, J.-B. *et al.* Isolation of an antimicrobial compound produced by bacteria associated with reef-building corals. *PeerJ* **4**, e2275 (2016).

12. Gochfeld, D. J. & Aeby, G. S. Antibacterial chemical defenses in Hawaiian corals

provide possible protection from disease. *Mar. Ecol. Prog. Ser.* **362**, 119–128

(2008).

13. Ritchie, K. B. Regulation of microbial populations by coral surface mucus and

mucus-associated bacteria. *Mar. Ecol. Prog. Ser.* **322**, 1–14 (2006).

14. Shnit-Orland, M. & Kushmaro, A. Coral mucus-associated bacteria: a possible first

line of defense. *FEMS Microbiol. Ecol.* **67**, 371–380 (2009).

15. Galloway, S. B. *et al. Coral Disease and Health Workshop: Coral Histopathology*

II. (NOAA Technical Memorandum NOS NCCOS 56 and NOAA Technical

Memorandum CRCP 4, National Oceanic and Atmospheric Administration, Silver

spring, 2007).

16. Peixoto, R. S., Rosado, P. M., Leite, D. C. de A., Rosado, A. S. & Bourne, D. G.

Beneficial Microorganisms for Corals (BMC): Proposed Mechanisms for Coral

Health and Resilience. *Front. Microbiol.* **8**, (2017).

17. Yang, S.-H. *et al.* Long-Term Survey Is Necessary to Reveal Various Shifts of

Microbial Composition in Corals. *Front. Microbiol.* **8**, (2017).

18. Shiu, J.-H. *et al.* A Newly Designed Primer Revealed High Phylogenetic Diversity

of Endozoicomonas in Coral Reefs. *Microbes Environ.* **33**, 172–185 (2018).

19. Hernandez-Agreda, A., Leggat, W., Bongaerts, P., Herrera, C. & Ainsworth, T. D.

Rethinking the Coral Microbiome: Simplicity Exists within a Diverse Microbial

Biosphere. *mBio* **9**, e00812-18 (2018).

20. Pollock, F. J. *et al.* Coral-associated bacteria demonstrate phylosymbiosis and

cophylogeny. *Nat. Commun.* **9**, 4921 (2018).

21. Knowlton, N. & Rohwer, F. Multispecies Microbial Mutualisms on Coral Reefs:

The Host as a Habitat. *Am. Nat.* **162**, S51–S62 (2003).

22. Ritchie, K. B. & Smith, G. W. Microbial Communities of Coral Surface

Mucopolysaccharide Layers. in *Coral Health and Disease* (eds. Rosenberg, P. E. &

Loya, P. Y.) 259–264 (Springer Berlin Heidelberg, 2004).

23. Bourne, D. G. & Munn, C. B. Diversity of bacteria associated with the coral

Pocillopora damicornis from the Great Barrier Reef. *Environ. Microbiol.* **7**, 1162–

1174 (2005).

24. Koren, O. & Rosenberg, E. Bacteria Associated with Mucus and Tissues of the

Coral Oculina patagonica in Summer and Winter. *Appl. Environ. Microbiol.* **72**,

5254–5259 (2006).

25. Johnston, I. S. & Rohwer, F. Microbial landscapes on the outer tissue surfaces of

the reef-building coral Porites compressa. *Coral Reefs* **26**, 375–383 (2007).

26. Chiu, H.-H., Mette, A., Shiu, J.-H. & Tang, S.-L. Bacterial Distribution in the

Epidermis and Mucus of the Coral Euphyllia glabrescens by CARD-FISH. *Zool.*

Stud. 11 (2012).

27. Ainsworth, T. D. *et al.* The coral core microbiome identifies rare bacterial taxa as

ubiquitous endosymbionts. *ISME J.* **9**, 2261–2274 (2015).

28. Yang, S.-H. *et al.* Prevalence of potential nitrogen-fixing, green sulfur bacteria in

the skeleton of reef-building coral Isopora palifera. *Limnol. Oceanogr.* **61**, 1078–

1086 (2016).

29. Glasl, B., Herndl, G. J. & Frade, P. R. The microbiome of coral surface mucus has

a key role in mediating holobiont health and survival upon disturbance. *ISME J.* **10**,

2280–2292 (2016).

30. van de Water, J. A. J. M., Allemand, D. & Ferrier-Pagès, C. Host-microbe

interactions in octocoral holobionts - recent advances and perspectives. *Microbiome*

6, 64 (2018).

31. Ainsworth, T. D., Fine, M., Roff, G. & Hoegh-Guldberg, O. Bacteria are not the

primary cause of bleaching in the Mediterranean coral Oculina patagonica. *ISME J.*

2, 67–73 (2007).

32. Work, T. & Aeby, G. Microbial aggregates within tissues infect a diversity of

corals throughout the Indo-Pacific. *Mar. Ecol. Prog. Ser.* **500**, 1–9 (2014).

33.   Water, J. A. J. M. van de *et al.* The coral immune response facilitates protection
against microbes during tissue regeneration. *Mol. Ecol.* **24**, 3390–3404 (2015).

34.   La Rivière, M., Garel, M. & Bally, M. Localization of endobacteria in the
gastrodermis of a Mediterranean gorgonian coral, Paramuricea clavata, using
fluorescence in situ hybridization. *Mar. Biol.* **163**, 206 (2016).

35.   Peters, E., Oprandy, J. & Yevich, P. Possible causal agent of" white band disease"
in Caribbean acroporid corals. *J. Invertebr. Pathol.* **396**, 394–396 (1983).

36.   Peters, E. A survey of cellular reactions to environmental stress and disease in
Caribbean scleractinian corals. *Helgol Meeresunters* **137**, 113–137 (1984).

37.   Ainsworth, T. & Hoegh-Guldberg, O. Bacterial communities closely associated
with coral tissues vary under experimental and natural reef conditions and thermal
stress. *Aquat. Biol.* **4**, 289–296 (2009).

38.   Neave, M. J. *et al.* Differential specificity between closely related corals and
abundant Endozoicomonas endosymbionts across global scales. *ISME J.* (2016).
doi:10.1038/ismej.2016.95

39.   Bayer, T. *et al.* The Microbiome of the Red Sea Coral Stylophora pistillata Is
Dominated by Tissue-Associated Endozoicomonas Bacteria. *Appl. Environ.*
Microbiol. **79**, 4759–4762 (2013).

40.   Work, T. M. & Aeby, G. S. Pathology of tissue loss (white syndrome) in Acropora
sp. corals from the Central Pacific. *J. Invertebr. Pathol.* **107**, 127–31 (2011).

41.   Tada, K., Sakai, K., Nakano, Y., Takemura, A. & Montani, S. Size-fractionated
phytoplankton biomass in coral reef waters off Sesoko Island, Okinawa, Japan. *J.*
Plankton Res. **25**, 991–997 (2003).

42.   Tanaka, Y. Nutrient uptake by reef building corals and the ecophysiological effects.
Oceanogr. Jpn. 101-117 (in Japanese) (2012).

43.   Brodie, J. E., Devlin, M., Haynes, D. & Waterhouse, J. Assessment of the
eutrophication status of the Great Barrier Reef lagoon (Australia). *Biogeochemistry*
106, 281–302 (2011).

44. Berkelmans, R., Weeks, S. J. & Steinberga, C. R. Upwelling linked to warm
summers and bleaching on the Great Barrier Reef. *Limnol. Oceanogr.* **55**, 2634–
2644 (2010).

45. Bourne, D., Iida, Y., Uthicke, S. & Smith-Keune, C. Changes in coral-associated
microbial communities during a bleaching event. *ISME J.* **2**, 350–363 (2007).

46. Vega Thurber, R. L. *et al.* Metagenomic analysis of stressed coral holobionts.
Environ. Microbiol. **11**, 2148–2163 (2009).

47. Littman, R., Willis, B. L. & Bourne, D. G. Metagenomic analysis of the coral
holobiont during a natural bleaching event on the Great Barrier Reef. *Environ.*
Microbiol. Rep. **3**, 651–660 (2011).

48. Kelly, L. W. *et al.* Local genomic adaptation of coral reef-associated microbiomes
to gradients of natural variability and anthropogenic stressors. *Proc. Natl. Acad.*
Sci. **111**, 10227–10232 (2014).

49. Shiu, J.-H. *et al.* Dynamics of coral-associated bacterial communities acclimated to
temperature stress based on recent thermal history. *Sci. Rep.* **7**, 14933 (2017).

50. Glasl, B., Webster, N. S. & Bourne, D. G. Microbial indicators as a diagnostic tool
for assessing water quality and climate stress in coral reef ecosystems. *Mar. Biol.*
164, 91 (2017).

51. Sudek, M., Work, T. M., Aeby, G. S. & Davy, S. K. Histological observations in
the Hawaiian reef coral, Porites compressa, affected by Porites bleaching with
tissue loss. *J. Invertebr. Pathol.* **111**, 121–125 (2012).

52. van Teeseling, M. C. F., de Pedro, M. A. & Cava, F. Determinants of Bacterial
Morphology: From Fundamentals to Possibilities for Antimicrobial Targeting.
Front. Microbiol. **8**, (2017).

53. Amann, R. I., Ludwig, W. & Schleifer, K. H. Phylogenetic identification and in situ
detection of individual microbial cells without cultivation. *Microbiol Mol Biol Rev*
59, 143–169 (1995).

54. Wagner, M., Horn, M. & Daims, H. Fluorescence in situ hybridisation for the
identification and characterisation of prokaryotes. *Curr. Opin. Microbiol.* **6**, 302–
309 (2003).

55. Bouvier, T. & del Giorgio, P. A. Factors influencing the detection of bacterial cells
using fluorescence in situ hybridization (FISH): A quantitative review of published
reports. *FEMS Microbiol. Ecol.* **44**, 3–15 (2003).

56. Prophet, E. B. *Laboratory methods in histotechnology*. (Amer Registry of
Pathology, 1992).

57. Riou, V., Périot, M. & Biegala, I. C. Specificity Re-evaluation of Oligonucleotide
Probes for the Detection of Marine Picoplankton by Tyramide Signal
Amplification-Fluorescent In Situ Hybridization. *Front. Microbiol.* **8**, (2017).

58. Siboni, N., Ben-Dov, E., Sivan, A. & Kushmaro, A. Global distribution and
diversity of coral-associated Archaea and their possible role in the coral holobiont
nitrogen cycle. *Environ. Microbiol.* **10**, 2979–2990 (2008).

59. Arp, D. J. & Bottomley, P. J. Nitrifers: more than 100 years from isolation to
genome sequences. *Microbe-Am. Soc. Microbiol.* **1**, 229 (2006).

60. Feng, G., Sun, W., Zhang, F., Karthik, L. & Li, Z. Inhabitancy of active
Nitrosopumilus-like ammonia-oxidizing archaea and Nitrospira nitrite-oxidizing
bacteria in the sponge Theonella swinhoei. *Sci. Rep.* **6**, 24966 (2016).

61. Garren, M. & Azam, F. New Method for Counting Bacteria Associated with Coral
Mucus. *Appl. Environ. Microbiol.* **76**, 6128–6128 (2010).

62. Garren, M. & Azam, F. Corals shed bacteria as a potential mechanism of resilience
to organic matter enrichment. *ISME J.* **6**, 1159–1165 (2012).

63. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
Nat. Methods **9**, 676–682 (2012).

64. Wada, N. *et al. In situ* visualization of bacterial populations in coral tissues: pitfalls
and solutions. *PeerJ* **4**, e2424 (2016).

65. Wallner, G., Amann, R. & Beisker, W. Optimizing fluorescent in situ hybridization

with rRNA-targeted oligonucleotide probes for flow cytometric identification of

microorganisms. *Cytometry* **14**, 136–43 (1993).

66. Daims, H., Brühl, A., Amann, R., Schleifer, K. H. & Wagner, M. The domain-

specific probe EUB338 is insufficient for the detection of all Bacteria: development

and evaluation of a more comprehensive probe set. *Syst. Appl. Microbiol.* **22**, 434–

44 (1999).

67. R Core Team. R: A Language and Environment for Statistical Computing. *Vienna,*

Austria: R Foundations for Statistical Computing. (2013).

68. Mangiafico, S. rcompanion: Functions to Support Extension Education Program

Evaluation (R package version 2.0.0). *The Comprehensive R Archive Network*

(2018).

69. Venables, W. N. & Ripley, B. D. *Modern Applied Statistics with S*. (Springer-

Verlag, 2002).

70. Zeileis, A., Kleiber, C. & Jackman, S. Regression Models for Count Data in R. *J.*

Stat. Softw. **27**, 1–25 (2008).

71. Fox, J. & Weisberg, S. *An R companion to applied regression*. (Sage Publications,

2018).

72. Peters, E. C. Anatomy. in Diseases of Coral 85–107 (John Wiley & Sons, Ltd,

2015). doi:10.1002/9781118828502.ch6