How pirate phage interferes with helper phage: Comparison of the two distinct strategies

Namiko Mitarai*

The Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, Copenhagen, 2100-DK, Denmark.

Abstract

Pirate phages use the structural proteins encoded by unrelated helper phages to propagate. The best-studied example is the pirate P4 and helper P2 of coliphages, and it has been known that the *Staphylococcus aureus* pathogenicity islands (SaPIs) that can encode virulence factors act as pirate phages, too. When alone in the host, the pirate phages act as a prophage, but when the helper phage gene is also in the same host cell, the pirate phage has ability to exploit the helper phages structural proteins to produce pirate phage particles and spread, interfering with the helper phage production. The known helper phages in these systems are temperate phage. Interestingly, the interference of the pirate phage to the helper phage occurs in different manner between the SaPI-helper system and the P4-P2 system. The SaPIs cannot lyse the helper lysogen upon infection, while when the helper phage lyse the SaPI lysogen, most of the phage particles produced are SaPI particles. On the contrary, in the P4-P2 system, the pirate P4 can lyse helper P2 lysogens to produce mostly the P4 particles, while when P2 phage lyses a P4 lysogen, most of the produces phages are P2 particles. Here, the consequences of these different strategies in the pirate and helper phage spreading among uninfected host was analyzed by using mathematical model. Both in the well-mixed environment and the spatially structured environment, SaPI's strategy interfere with the helper phage

---

*Corresponding author
Email address: mitarai@nbi.dk (Namiko Mitarai)

spreading significantly more than the P4's strategy, because SaPI interfere with the helper phage's main reproduction step, while P4 interfere only by forcing the helper lysogens to lyse. However, the interference is weaker in the spatially structured environment, because the system self-organize so that the helper phages take over the front of propagation due to the need of helper phage for the pirate phage spreading.

Competing interests: The author declares no competing interest.

Keywords: temperate phage, popuplation dynamics, Lotoka-Volterra equation

---

Introduction

Spreading of pathogenic genes including antibiotic-resistance genes and toxin genes among bacteria is a real threat to the human society [1, 2]. One of the well-known examples of the mobile genetic elements that contribute the spreading

5   process is the *Staphylococcus aureus* pathogenicity islands (SaPIs). SaPIs spread among the host bacteria by using bacteriophages as their "helper" [3, 4, 5, 1, 2]. Interestingly, the mechanism of SaPIs spreading has many aspects in common with that of the well-studied pirate phage P4, that uses the helper phage P2 to spread among the host *Escherichia coli* [3, 6, 5].

10   The pirate phages and helper phages we consider here all belong to the order *Caudovirales* or the tailed bacteriophages [7, 5], hence a phage particle consists of a head that contains the double stranded DNA genomes and a tail. However, SaPI or pirate phage P4 does not have the structural genes that encode head and tail proteins, thus it cannot produce its progeny when infecting a host cell alone,

15   and typically it becomes a prophage (Fig. 1a). Helper phages for them (e.g., ϕ11 for SaPIbov1, P2 for P4) are temperate phages. When infecting a sensitive host cell, it can enter either the lysogenic pathway with probability α_H where the phage genome is integrated to the host chromosome as a prophage, provide immunity to the same phage, and replicate with the host, or the lytic pathway

20   with probability $1 - \alpha_H$ to produce multiple copies of the phage particles and come out by lysing the host cell (Fig. 1b). Interestingly, pirate phages encode

genes that allow them to interfere with the helper phage burst if it happens in the same host cell, including a factor to modify the size of helper head capsid so that it becomes too small to carry the helper genome. As a result, pirate phages take over the structural proteins produced from helper genes and produce pirate phages. This phenomenon was termed as "molecular piracy" [5].

The fascinating mechanisms of the molecular piracy have been intensively studied, first for P4 which was found in 1963 [8]. SaPIs have attracted attention as a cause of the outbreak of toxic shock syndrome [9] and on-going studies are revealing the importance of them in *S. aureus* [2]. The pirate phages are ubiquitous; it has been found that the P4-type sequences are widespread among *E. coli* strains [10], while on average one SaPI was found per natural *S. aureus* strain [2]. These findings demonstrate the importance of the pirate phages as a source of horizontal gene transfer in bacteria.

When looking closer, however, SaPIs and P4 appear to have chosen very different strategies in their behavior. In the case of SaPI, when it infect a helper lysogen, it simply integrates its genome to the host genome to form a double lysogen [11]. However, if a helper phage infect a SaPI lysogen and goes lytic (or the helper prophage in a double lysogen is induced), majority (fraction $f_P^{(H)} \geq 0.9$) of the produced phage particles are SaPI particles [12]. The interference is so severe that helper phages are normally not able to make a visible plaque on a pirate lysogen lawn [4] (Fig. 1c). On the contrary, when infecting a helper P2 lysogen, a pirate P4 phage chooses lysogenic pathway with probability $\alpha_P \sim 0.4$, otherwise it chooses lysis [13]. Upon bust, almost all the produced phages are P4 (fraction $f_P^{(P)} \sim 1$) [14], i.e., P4 phage can spread on a lawn of P2 lysogen. When a P2 phage infects a P4 lysogen, the interference is weak, and upon lysis only small part (fraction $f_P^{(H)} \sim 0.001$) of the produced phages are P4 [15], enabling P2 phage to spread on a lawn of P4 lysogens (Fig. 1d). The key pirate phage parameters that illustrate these differences are summarized in Table 1.

Since both SaPIs and P4 are widespread, both choices seems to be viable. However, at least when looking at spreading on lysogens, the SaPI-helper system

is inhibiting each other, while the P4-P2 system is supporting each other. A nontrivial question is how they spread when both pirate phage and helper phage

encounter a population of host cells without any prophage. Can both the pirate and helper phages spread among the host, or does the interference prevent the spreading? What is the difference in the outcome between the two different strategies?

In order to answer these questions, we study the spreading of pirate phage

and helper phage among uninfected host cells at population level by using a mathematical model. Previous studies on phage-bacteria systems have shown that the Lotka-Volterra type population dynamics model can reproduce many of the important phage-bacteria dynamics reasonably well in various length and time scales [16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. As a simple and

well-defined set up, we here analyze the following two cases: Overnight growth in a culture of uninfected cells well-mixed with the pirate and helper phages, and the plaque formation in a lawn of uninfected cells initially spotted by a droplet of pirate-helper phage mixture. For the well-mixed case, we using the Lotka-Volterra type equations, and for the plaque simulation we let bacteria

grow locally while let phage and nutrient diffuse, describing the system using a reaction-diffusion type model [19, 26]. In both cases, we consider the growth of an overnight, which is the time scale where effect of the induction and the mutation are negligible. We compare the two phage strategies reflected in the values of $f_P^{(H)}$, α_P, and $f_P^{(P)}$, to reveal their impact on the phage spreading.

Model equations

Growth in a well-mixed culture

We use Lotka-Volterra type ordinary differential equation to model the population dynamics of the system, which has been used to describe temperate phage and bacteria population dynamics [18, 23, 29, 26]. To extend the model

to the pirate-helper phage system described in Fig. 1, we consider the following variables time t; uninfected bacteria concentration $B(t)$, the pirate phage

lysogen concentration $L_P(t)$, the helper phage lysogen concentration $L_H(t)$, the pirate phage concentration $P(t)$, the helper phage concentration $H(t)$, intermediate states concentration I with subscripts that express each infection pathway, and the nutrient concentration $n(t)$. Note that, in the case of P4, 99% of the case the genome is integrated to the chromosome, but 1% go into the plasmid state [13]. We here ignore these distinctions for simplicity and treat them as lysogens. We express these populations in the unit of number per volume.

When the lytic pathway is chosen upon infection, the free phages are produced after intermediate steps. Naturally, when a helper phage infect a uninfected bacterium and choose lysis, it will produce helper phages only as long as there is no super infection by a pirate phage in early part of lytic pathway. The production of the pirate phage (or mixture of the helper and the pirate phages) can happen in different ways: (i) When a pirate phage infects a helper lysogen and force the lytic pathway with the probability $1 - \alpha_P$. (ii) When a helper phage infects the pirate lysogen and choose the lytic pathway with probability $1 - \alpha_H$. Further, we assume that some pirate phages can be produced (iii) when a pirate phage infects a bacterium in an early stage of the helper lytic pathway, to model the (almost) simultaneous co-infection. The model reflects these different possibilities for production of phages. The resulting model equations are as follows:

$$B_{tot} = B + L_P + L_{PH} + L_H + \sum_{i=1}^{M} \left(I_H^{(i)} + I_{HP}^{(i)} + I_P^{(i)} \right), \tag{1}$$

$$g_w(n) = g_{max} \frac{n}{k_s + n}, \tag{2}$$

$$\frac{dn}{dt} = -g_w(n)B_{tot}, \tag{3}$$

$$\frac{dB}{dt} = g_w(n)B - \eta_P \cdot P \cdot B - \eta_H H \cdot B, \tag{4}$$

$$\frac{dL_P}{dt} = g_w(n)L_P + \eta_P \cdot P \cdot B - \eta_H H \cdot L_P, \tag{5}$$

$$\frac{dL_H}{dt} = g_w(n)L_H + \alpha_H \cdot \eta_H \cdot H \cdot B - \eta_P P \cdot L_H, \tag{6}$$

$$\frac{dL_{PH}}{dt} = g_w(n)L_{PH} + \alpha_H \cdot \eta_H \cdot H \cdot L_P + \alpha_P \cdot \eta_P \cdot P \cdot L_H \tag{7}$$

$$\frac{dI_H^{(1)}}{dt} = (1 - \alpha_H) \cdot \eta_H \cdot H \cdot B - \eta_P \cdot P \cdot I_H^{(1)} - \frac{g_w(n)}{g_{max}} \frac{M}{\tau} I_H^{(1)}, \tag{8}$$

$$\frac{dI_H^{(i)}}{dt} = \frac{g_w(n)}{g_{max}} \frac{M}{\tau} (I_H^{(i-)1} - I_H^{(i)}) \qquad (\text{for } i = 2, \cdots, M), \tag{9}$$

$$\frac{dI_{HP}^{(1)}}{dt} = (1 - \alpha_H) \cdot \eta_H \cdot H \cdot L_P + \eta_P \cdot P \cdot I_H^{(1)} - \frac{g_w(n)}{g_{max}} \frac{M}{\tau} I_{HP}^{(1)}, \tag{10}$$

$$\frac{dI_{HP}^{(i)}}{dt} = \frac{g_w(n)}{g_{max}} \frac{M}{\tau} (I_{HP}^{(i-)1} - I_{HP}^{(i)}) \qquad (\text{for } i = 2, \cdots, M), \tag{11}$$

$$\frac{dI_P^{(1)}}{dt} = (1 - \alpha_P) \cdot \eta_P \cdot P \cdot L_H - \frac{g_w(n)}{g_{max}} \frac{M}{\tau} I_P^{(1)}, \tag{12}$$

$$\frac{dI_P^{(i)}}{dt} = \frac{g_w(n)}{g_{max}} \frac{M}{\tau} (I_P^{(i-)1} - I_P^{(i)}) \qquad (\text{for } i = 2, \cdots, M). \tag{13}$$

$$\frac{dH}{dt} = \beta \frac{M}{\tau} \left(I_H^{(M)} + (1 - f_P^{(H)}) I_{HP}^{(M)} + (1 - f_P^{(P)}) I_P^{(M)} \right) - \eta_H \cdot H \cdot B_{tot}$$
$$\qquad - \delta H, \tag{14}$$

$$\frac{dP}{dt} = \beta \frac{M}{\tau} \left(f_P^{(H)} I_{HP}^{(M)} + f_P^{(P)} I_P^{(M)} \right) - \eta_P \cdot P \cdot B_{tot} - \delta P. \tag{15}$$

Here, we assume Monod's growth law [30] with the maximum growth rate g_{max} and the Michaels constant k_s. Nutrient is measured in a unit where the yield for bacteria growth is one. We assume that the bacteria that are in intermediate steps of lysis consume the nutrient at the same rate as the growing bacteria. The latency time τ and the total number of phage particles produced per burst β are assumed to be the same for all infection pathways for simplicity. The intermediate infection steps are divided into $M = 10$ steps, that gives about 30% fluctuation in the latency time [26]. We assume that the latency time becomes longer proportional to the doubling time, ensuring that the phage production stops when the bacteria growth stops due to the lack of nutrient. Free phages decay at the rate δ, but it is in general small [31] that does not affect the present result.

Initial condition and Numerical integration

We consider the population dynamics when the uninfected cells are mixed with the pirate-helper phage lysate in a rich medium. This is represented by initially having finite uninfected bacteria and nutrient as $B(0) = b_0$ and $n(0) =$

n_0. The average phage inputs at time zero of helper phage API_H and pirate phage API_P determine the initial phage concentrations as $H(0) = b_0 \cdot API_H$ and and $P(0) = b_0 \cdot API_P$.

The integration was done by the 4-th order Runge-Kutta method, with time step 10^{-6}/h.

Plaque formation in a spatially structured environment

We model a plaque formation experiment setup called the spot assay [32], where large number of bacteria in a soft agar is cast in a thin layer on a hard agar plate that contains nutrient, a droplet of liquid that contains phages is placed, and subsequently incubated overnight. With this method, it is easy to co-infect the bacteria with both helper and pirate phages in a well-controlled manner. Bacteria cannot swim visibly in the high viscosity of soft agar, hence each initially casted bacterium grow and divide locally to form a microcolony [33, 34]. As time goes, phages produced upon lysis of infected bacteria reach neighbor bacteria by diffusion to infect them and continue spreading. When the nutrients run out, both bacteria and phage growth ceases, leaving the plaque in the final configuration. When the phage is temperate, lysogens can grow in the infected area, resulting in a turbid plaque.

The clear plaque formation of virulent phage has been modeled by using reaction-diffusion type models [19, 35] and extended for turbid plaque formation by the temperate phage [26]. Here, we modify the turbid plaque formation model proposed in [26], where the infection of bacteria by phage is modeled by using Lotka-Volterra type equation, to include the pirate phages.The latency time for the phage burst is taken into account by considering the intermediate states, and bacteria grows locally while the phage and the nutrient diffuse in space. Even though the local growth of bacteria into microcolonies may provide extra protection against phage attack [34, 27], for simplicity we here assume interaction in a locally mixed population. We use the same symbols as the well-mixed case to express the different populations, but now the they are expressed in the unit of number per area by summing up the population over the thin

soft-agar thickness Δa, and they are function of position $\mathbf{r}$ and time t. Note that this changes the unit of the variables from per volume in the well-mixed model. Furthermore, the diffusion terms are added for the nutrient and phages. The resulting model equations are parallel to the ones in eqs. (1)-(15), but the total derivative by time in the left hand side of eqs. (4)-(15) should be replaced with partial derivative by time, and the diffusion terms $D_n\nabla^2 n$, $D_H\nabla^2 H$, and $D_P\nabla^2 P$ should be added to the right hand side of eqs. (2), (14), and (15), respectively. Further more, he pirate (helper) phage adsorption rate η_P (η_H) needs to be divided by Δa reflecting that we are summing up the populations over this depth. Finally, to reflect the change of unit of the nutrient from per ml to per $\mu\mathrm{m}^2$, the symbol for the Michaels constant is changed from k_s in the well mixed equation (2) to K_s for the spatial model (used in the parameter table in Table 2).

Initial condition and Numerical integration

We solved the model partial differential equations in polar coordinates with the origin at the center of the initially placed droplet. We assume the symmetry in circular direction, and the spatial variation is only considered along the radial direction r. At a distant outer boundary $r = R_{max}$, which is set to be 9 mm, we impose reflecting boundary condition for both phage and nutrient concentrations.

We analyze the plaque spreading in a lawn of uninfected bacteria. This is represented by initially distributing uninfected bacteria and nutrient uniformly as $B(r,0) = B_0$ and $n(r,0) = N_0$, with the rest of the variables set to zero, except for the phages placed in the middle where the phages are spotted. The average phage inputs in the initial spot of helper phage API_H and pirate phage API_P determine the initial phage concentrations. The initial spot of radius $R_s = 1$mm is given by setting $H(r,0) = B_0 \cdot API_H \cdot \Theta(R_s - r)$ and $P(r,0) = B_0 \cdot API_P \cdot \Theta(R_s - r)$, where $\Theta(x)$ is the Heaviside step function. We set $B_0 = 1/(18\mu\mathrm{m})^2$ throughout this paper [26].

The integration was done by the finite difference method, and time integra-

tion was done by the 4-th order Runge-Kutta method. The size of the spatial discretization, Δr, was set to $18\mu m$, hence at the initial condition there is one bacterium per Δr^2 ($B_0 = 1/\Delta r^2$). Time step of integration was set to $\Delta t = 10^{-4} h$.

Parameters

Table 2 shows the list of the default parameters used, with the references for each value when available. Most of the values are chosen from the well-studied P4-P2 system, but when known the value for SaPI and a relatively well-studied helper phage for SaPIs, $\phi 11$, are also stated with references. The difference in the strategies are reflected in the values of $f_P^{(H)}$, α_P, and $f_P^{(P)}$, and the rest of the parameters are kept same between the two system.

Results

Growth in a well-mixed culture

In order to compare the SaPI-helper strategy and the P4-P2 strategy, we set most of the parameters same for both systems as described in Table 2. The difference in strategy is reflected by choice of the following parameters as summarized in Table 1: the pirate frequency of lysogeny upon infection of helper lysogen α_P, the fraction of the pirate phage production upon a pirate phage lysing a helper lysogen $f_P^{(P)}$, and the fraction of the pirate phage production upon a helper phage lysing a pirate lysogen $f_P^{(H)}$. The detail of the model equations are given in the Model section.

We first simulate the case where uninfected cells of concentration $b_0 = 10^5$/ml are mixed with the pirate phage and the helper phage lysate. We vary the initial average phage input for the pirate phage API_P and that for the helper phage API_H, by mixing $API_P \cdot b_0$ pirate phages and $API_H \cdot b_0$ helper phages to the culture at time zero. As the bacteria cells grow, the nutrient in the culture is consumed. The initial nutrient concentration was set so that the population would reach 10^9/ml if there were no killing by phage. When

the nutrient concentrations become zero, both the bacteria and the phage stop growing, though free phage adsorption to the bacteria still happens.

Figures 2 show the time course of the population concentrations for the SaPI-helper strategy (Figs. 2a) and the P4-P2 strategy (Figs. 2b), with $API_P = API_H = 100$.

In the case of the SaPI-helper system, we can see that the pirate phage concentration P rises following closely with the helper phage concentration H increase. The delay is because the pirate phage can replicate only if the pirate lysogens are infected by the helper phage, while helper phage can just replicate upon single infection. At first, the initially added pirate phages typically form lysogens and they grow, as reflected in the increase of L_P. The replicated helper phage starts to infect these pirate lysogens, and they produce mainly the pirate phages. This pirate phage replication severely interfere with the helper phage replication, hence overall phage replication rate is low (compare with the P4-P2 strategy Figs. 2b). As a result, significant number of bacteria are left uninfected (B) when the nutrient is depleted.

This is in contrast to the P4-P2 system, where the pirate phage can take over the helper phage only when it infects a helper lysogen. For the increase of the pirate phage, the increase of the helper lysogen L_H needs to happen first, which allows the helper phages to grow without feeling the interference by the pirate phage for quite some time. The pirate phage P starts to increase only about 6h after the start of the simulation, but the system soon run out of the nutrient and thus pirate phages does not increase much. At the same time, the lack of interference to the helper phage replication allows the helper phages to infect all the uninfected cells B and the pirate phage L_P, converting them to either helper lysogens L_H or the double lysogens L_{PH}.

Initial phage input dependence of the prophage spreading in well-mixed environment

The degree of interference between the helper phage and the pirate phage should depend on the initial phage inputs. This is studied in the Figs. 3 for

each strategy. Since the free phages are eventually adsorbed in the bacteria, we consider the number of the lysogens with helper prophage $L_H + L_{PH}$ as a measure of the helper phage spreading, while the lysogens with pirate prophage $L_P + L_{PH}$ as a measure of the pirate phage spreading.

Clearly, in the SaPI-helper system, a larger amount of the pirate phage in initial culture (large API_P) interferes more severe with the helper phage spreading (Fig. 3a left), especially when helper phage input API_H is relatively small. However, when API_P is high, the pirate phage spreading is not compromised much from the lack of helper phage growth (Fig. 3a right), since the initial amount of pirate phage is already high enough and the formed pirate phage lysogens can grow over time. At the same time, the pirate phage clearly benefit from taking over the helper phage for replication when API_P is low but API_H is high, reflected in the increased amount of $L_P + L_{PH}$ when increasing API_H with keeping $API_P \sim 0.1$.

On the contrary, in the P4-P2 system, the final concentration of bacteria with helper prophage $L_H + L_{PH}$ is almost independent of API_P (Fig. 3a right), indicating that the helper phage spreading is not interfered by the pirate phage much. Interestingly, the pirate phage spreading at a fixed API_P showed a non-monotonic dependence on API_H. This is because the lysis of the pirate lysogen by helper infection does not contribute to the pirate phage production, and the pirate phage needs to lyse helper lysogen for pirate phage production; increasing API_H for a fixed API_P contribute both processes while the latter become dominant only when API_H is high enough so that initially added pirate phages infect the helper lysogens and not the uninfected host. When API_H value is moderate, it can lyse the pirate lysogens formed due to initially added pirate phages, which reduces the amount of pirate lysogens.

When comparing the two strategies, SaPIs are better at spreading than P4 in a well-mixed system, as depicted by the larger population of $L_P + L_{PH}$ in all the studied parameter regions. At the same time, the helper phages are interfered strongly by SaPI but not by P4.

Plaque formation in a lawn of uninfected bacteria

The analysis of the well-mixed system showed that SaPI significantly interfere with the helper spreading, while P4 does not. The strong interference by SaPI could make the spreading of SaPI in the spatially structured harder, since if the helper phage is locally depleted by interference, the phage cannot spatially propagate even if there uninfected cells outside of the plaque.

In order to see if local depletion can be a problem, we next analyzed the plaque spreading in a lawn of uninfected bacteria. The simulation mimics the plaque formation by the spot assay [32], where soft agar with bacteria is cast over hard agar with nutrient, a droplet with mixture of phages is placed in the center, and incubated overnight. In the setup, bacteria cannot swim visibly and the local growth results in formation of a microcolony at the position where a bacterium was initially dispersed [33, 34]. Phages grow and diffuse to infect new bacteria, resulting in a circular zone of killing around the initial spotted area. When the phage is temperate, lysogens will survive and grow, making the plaque turbid. The simulation was started by placing a droplet of mixture of pirate and helper phage with radius 1mm. The concentration of phages in the droplet is defined by the average phage input per initially placed bacterium for the helper phage, API_H, and that for the pirate phage, API_P. The detail of the simulation is given in the method section.

The time courses of the plaque formation with $API_H = API_P = 1$ (i.e., on average there is one of each phage per initially placed bacterium under the initial spot) are shown in Fig. 4a-f for the SaPI-helper system and in Fig. 4g-l for the P4-P2 system, respectively. The populations at each location are shown as a function of distance from the infection origin, r. The unit of the population density is normalized to initial bacteria density B_0.

The profile for the uninfected bacteria and the helper phage at the front of the plaque spreading looks very similar between the two system (Fig 4ab vs. gh). This is because, the helper phage properties are set to be identical when there is no pirate phage, and at the front of the plaque propagation, the helper phages does not feel the interference by the pirate phage due to their low

density.

The difference can be seen is in how the pirate phage can propagate. We see that pirate phage can come up much later and close to the center in the P4-P2 system (Fig. 4i), where pirate phages can be produced only when the helper lysogen L_H is lysed by the pirate phage infection. In the SaPI-helper case (Fig. 4c), the wave of pirate phage follows closer to the front propagation of helper phage by taking over the part of the helper phage lysis, making it possible for the pirate phage to spread a lot faster and further away, making larger amount of the double lysogen in the final state (Fig. 4f vs. l).

These results depict that the SaPI-helper strategy is indeed beneficial for the spreading of the pirate phage prophage spreading in space, because the strong interference does not inhibit helper spreading as long as the helper phage takes over the front of the plaque propagation. The P4-P2 strategy, on the other hand, does allow the helper phages to spread, but the pirate phage gene spread is delayed more because pirate phage can grow only later, after the helper lysogens are established.

Initial phage input dependence of the prophage spreading in spatially structured environment

The spreading of the phage gene should depends on the initial concentration of the each phage, API_H and API_P, in the spot. When the initial pirate phage input API_P is too high compared to API_H, the strong interference of SaPI-helper strategy may inhibit spreading of both of the phages, while when $API_H \gg API_P$, the P4-P2 strategy may better to spread since it can lyse the helper lysogens.

For systematic understanding, we investigated how the outcome changes when varying the initial concentration of the phages in the spot, API_H and API_P. The results are summarized in Fig. 5 for (a) the SaPI-helper strategy and (b) the P4-P2 strategy by the final population with the helper phage gene $L_H + L_{PH}$ and the pirate phage gene $L_P + L_{PH}$ integrated over space. The integration is done outside of the initial spot radius, in order to focus on the

effect of the spreading.

For the helper phage spreading, we observed that the interference by SaPI is relatively weak (Fig. 5a top left). This is because the system tend to self-organize so that the helper take over the spreading front. Once the helper take over the front, it is the pirate phage cannot lyse the helper lysogen, hence there will be large amount of helper prophages left. An example of such self-organization is depicted in the case of $API_H = 1$ and $API_p = 100$ by showing the time course of the pirate phage and helper phage profile over time (Fig. 5a bottom). At the start, the pirate phage is in the front of propagation, but it goes down over time since the pirate phages simply diffuse and adsorbed. The helper phage can reproduce in small amount by infecting uninfected host, and at about 3h the helper phage catches up the propagation front, and start to replicate fast by lysing the uninfected cells at the propagation front, producing a peak of helper phage at 4h. This peak propagates with pirate phage peak closely following behind, because some of the helper phages lyses the pirate phage lysogens that were produced by the initially added pirate phages. Of course, this self-organization cannot happen when API_H is too small and API_P is too large, depicted by the significantly low level of helper phage spreading at $API_H \sim 0.01$ and $API_P \sim 100$. Overall, the pirate phage spread fairly well (Fig. 5a top right), because if the helper phage take over the propagating front, pirate phage can follow closely, while when the initial pirate phage level is so high to completely inhibit helper phage propagation, the abundant initial pirate phages diffuses and produce the pirate lysogens.

In the P4-P2 system, again the helper phage spreading is not interfered by the pirate phage P4 (Fig. 5b top left), because even when the $API_P \gg API_H$ at first, the helper phages simply lyse the pirate lysogens to produce helper phage and propagate, as depicted in the very rapid take over of the front by helper phages in the $API_H = 1$ and $API_P = 100$ case shown in Fig. 5b bottom. Overall, the pirate phage spreading increase with (Fig. 5b top right) increasing API_P and API_H, but the absolute level of final spreading of the pirate phage gene $L_P + L_{PH}$ is significantly less than the SaPI case.

Discussion

The pirate-helper phage system demonstrates the fascinating richness of the
interaction between phages in nature. At the same time, given the increased
number of possible infection scenarios, it becomes significantly complex to an-
alyze the population dynamics. Still, the SaPI-helper system and the P4-P2
system pose interesting examples, in that their spreading among the lysogen
populations are completely opposite (Fig. 1). The present analysis of spreading
among initially uninfected population showed that, in both well-mixed and spa-
tially structured cases, SaPI is spreading better than P4. At the same time, SaPI
strongly interferes with the helper phage spreading, while P4 does not interfere
much with the P2 spreading. This is natural, because SaPI takes over most of
the phage production when the helper phage trigger the lysis (i.e., $f_P^{(H)} \sim 1$),
which is the major part of the helper phage production. On the contrary, even
if the pirate phages can lyse the helper lysogens and produce mostly the pirate
phage ($\alpha_P < 1$, $f_P^{(P)} \sim 1$) as P4 does, for helper phage that is some reduc-
tion of the existing helper lysogens but not the interference to the helper phage
reproduction, hence effect is weak.

It is an interesting and unsolved question why SaPI-helper system and P4-
P2 system has such different strategies. Here, it is worth noting that $f_P^{(H)}$ of
P4-P2 system changes close to 1 upon mutation of P2 gene A or *rep* gene of the
host bacterium [15]; in other word, P4-P2 system is only a few mutations away
from the SaPI-helper type strategy. Further consideration of the co-evolution
of the whole system is needed to reveal what is keeping the P4-P2 system in the
present parameter space.

Another point that the present analysis clarified is the weaker trade-off be-
tween the helper phage spreading and the pirate phage spreading in the spatially
structured lawn of uninfected cells. Even if the interference by pirate phage upon
helper burst is strong as $f_P^{(H)} \sim 1$, as long as the helper phages come to the
front of spreading this does not inhibit the helper phage spreading, and pirate
phage can spread as well by following the front. Because the pirate phage can-

not spread by themselves, it is relatively easy for the helper phage to take over the front. The pirate-behind-helper configuration makes the most of the pirate phage lysogens be double lysogens with the helper. However, this contradicts somewhat to the observation that very few SaPIs are found as a double lysogen with its own helper phage [2]. The scarcity of SaPI-helper double lysogen could be due to strong selection pressure against being a helper phage because SaPI is inhibiting the helper phage so much upon induction of the double lysogen [2]. This possibility can be analyzed by extending the present population dynamics model to include inductions and mutations and simulate longer time scale. Also, the present results relies on the assumption that bacteria does not migrate, hence the propagation of the phages relies only on the diffusion on top of the replication. The virulent phage propagation on the migrating population of bacteria has been found to show very rich patterns [36], and it will be interesting to study similar setup with pirate and helper phages.

Naturally, the pirate-helper interaction parameters vary depending on the combination. The frequency of lysogenization α_H for phage 80α was found to be quite low ($10^{-5} \sim 10^{-3}$) [37], which will change the outcome even if other parameters were identical. The frequency of lysogenization for helper phage α_H may be selected mainly by the frequency of the host population collapse [38] or by competition with other temperate phages [29], independent of the interaction with the pirate phage, even though the value of α_H do affect the outcome of the pirate-helper system strongly. More interestingly, the P4 phage cannot lyse the phage 186 lysogen, i.e., $\alpha_P = 1$, while interference upon the phage 186 bursting P4 lysogen is moderately strong ($f_P^{(H)} \sim 0.6$ to 0.9)[39], which appear to be closer to the SaPI-helper strategy. It is difficult to tell how often a given pirate phage uses the found helper phage in natural environments, even though that could affect how much the system had co-evolved. It should also be noted that using multiple helper phages is beneficial by itself, since it greatly increases the host range of the pirate phages. However, at the same time, the selection for the ability for a pirate phage to interact with multiple helper phages may put some limit in optimizing the interaction with a specific helper phage due to the

molecular differences of helper phages. In addition, the pirate phage may deliver the gene beneficial to the host bacteria as SaPIs carry pathogenic genes, and this can significantly contribute to spreading of the pirate phage gene, too.

Finally, the similarity and the differences of the pirate phages from the defective interfering particles (DIPs) in viral systems should be commented. DIPs are the defective viral particles that cannot replicate by itself upon infection of the host, but when co-infected the host with intact virus, it will interfere with the production of intact viruses and produce DIPs instead [40, 41]. The known naturally occurring examples of DIPs such as those found in influenza virus are produced as the original virus replicates [40], but there are several effort to engineer DIPs to interfere the spreading of pathogenic virus, including the HIV virus [42]. The noticeable difference between DIPs and pirate phage discussed in this paper is that the "helper" viruses for DIPs typically shows chronic infection, i.e., it lives with the host cells for relatively long time and keep producing the viruses, and does not have dormant mode that corresponds to lysogeny in the helper temperate phage. As a consequence, there is no corresponding difference of the strategies of pirate phage replication that depends on the which phage determines to "go lytic". Nevertheless, some of the population dynamics discussed here has some similarity to those of DIP's. For example, an recent theoretical paper that analyzed spreading of the DIPs in spatially structured environment pointed out that the spreading speed of the virus is not affected by the DIPs if the virus takes the front of propagation, even if the wave of DIPs is propagating just behind. This is consistent with the present finding that once the helper phage takes over the front that is stably maintained (Fig. 5), and it is expected that also in DIP-virus system, there is a tendency to self-organize so that the intact virus take over the front. It is worth mentioning that experiment on the spreading of DIPs and viruses mixture in space reported highly heterogeneous pattern in some conditions [43], demonstrating the importance of stochasticity. It is an interesting future work to extend the model to include stochastic population fluctuations in pirate-helper phage system, as well as analyzing it experimentally.

Overall, this paper sheds light on one of the many facets of the art of war among bacteria and phages. Considering the ubiquity of the pirate phage, their impact on the ecology and evolution of the microbial system should be significant. Clearly, further study is required to understand the rich population dynamics of pirate-helper systems.

Acknowledgments

NM thanks Kim Sneppen for fruitful discussions. This work is funded by Danish National Research Foundation (BASP: DNRF120).

References

[1] J. R. Penadés, G. E. Christie, The phage-inducible chromosomal islands: a family of highly evolved molecular parasites, Annual review of virology 2 (2015) 181–201.

[2] R. P. Novick, G. Ram, The floating (pathogenicity) island: a genomic dessert, Trends in Genetics 32 (2) (2016) 114–126.

[3] B. H. Lindqvist, G. Dehò, R. Calendar, Mechanisms of genome propagation and helper exploitation by satellite phage p4., Microbiological reviews 57 (3) (1993) 683–702.

[4] G. Ram, J. Chen, K. Kumar, H. F. Ross, C. Ubeda, P. K. Damle, K. D. Lane, J. R. Penadés, G. E. Christie, R. P. Novick, Staphylococcal pathogenicity island interference with helper phage reproduction is a paradigm of molecular parasitism, Proceedings of the National Academy of Sciences 109 (40) (2012) 16300–16305.

[5] G. E. Christie, T. Dokland, Pirates of the caudovirales, Virology 434 (2) (2012) 210–221.

[6] G. E. Christie, R. Calendar, Interactions between satellite bacteriophage p4 and its helpers, Annual review of genetics 24 (1) (1990) 465–490.

[7] H.-W. Ackermann, Tailed bacteriophages: the order caudovirales, in: Advances in virus research, Vol. 51, Elsevier, 1998, pp. 135–201.

[8] E. Six, A defective phage depending on phage p2, in: Bacteriol. Proc, Vol. 138, 1963.

[9] J. A. Lindsay, A. Ruzin, H. F. Ross, N. Kurepina, R. P. Novick, The gene for toxic shock toxin is carried by a family of mobile pathogenicity islands in staphylococcus aureus, Molecular microbiology 29 (2) (1998) 527–543.

[10] F. W. Studier, P. Daegelen, R. E. Lenski, S. Maslov, J. F. Kim, Understanding the differences between genome sequences of escherichia coli b strains rel606 and bl21 (de3) and comparison of the e. coli b and k-12 genomes, Journal of molecular biology 394 (4) (2009) 653–680.

[11] R. P. Novick, G. E. Christie, J. R. Penadés, The phage-related chromosomal islands of gram-positive bacteria, Nature reviews. Microbiology 8 (8) (2010) 541.

[12] N. Quiles-Puchalt, R. Martínez-Rubio, G. Ram, Í. Lasa, J. R. Penadés, Unravelling bacteriophage ϕ11 requirements for packaging and transfer of mobile genetic elements in s taphylococcus aureus, Molecular microbiology 91 (3) (2014) 423–437.

[13] F. Briani, G. Dehò, F. Forti, D. Ghisotti, The plasmid status of satellite bacteriophage p4, Plasmid 45 (1) (2001) 1–17.

[14] E. W. Six, C. A. C. Klug, Bacteriophage p4: a satellite virus depending on a helper such as prophage p2, Virology 51 (2) (1973) 327–344.

[15] E. W. Six, B. H. Lindqvist, Mutual derepression in the p2–p4 bacteriophage system, Virology 87 (2) (1978) 217–230.

[16] A. Campbell, Conditions for the existence of bacteriophage, Evolution (1961) 153–165.

[17] B. R. Levin, F. M. Stewart, L. Chao, Resource-limited growth, competition, and predation: a model and experimental studies with bacteria and bacteriophage, The American Naturalist 111 (977) (1977) 3–24.

[18] F. M. Stewart, B. R. Levin, The population biology of bacterial viruses: why be temperate, Theoretical population biology 26 (1) (1984) 93–117.

[19] J. Yin, J. McCaskill, Replication of viruses in a growing plaque: a reaction-diffusion model, Biophysical journal 61 (6) (1992) 1540–1549.

[20] T. F. Thingstad, Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems, Limnology and Oceanography 45 (6) (2000) 1320–1328.

[21] J. S. Weitz, J. Dushoff, Alternative stable states in host–phage dynamics, Theoretical Ecology 1 (1) (2008) 13–19.

[22] Z. Wang, N. Goldenfeld, Fixed points and limit cycles in the population dynamics of lysogenic viruses and their hosts, Physical Review E 82 (1) (2010) 011918.

[23] N. D. Maynard, E. W. Birch, J. C. Sanghvi, L. Chen, M. V. Gutschow, M. W. Covert, A forward-genetic screen and dynamic analysis of lambda phage host-dependencies reveals an extensive interaction network and a new anti-viral strategy, PLoS genetics 6 (7) (2010) e1001017.

[24] L. F. Jover, M. H. Cortez, J. S. Weitz, Mechanisms of multi-strain coexistence in host–phage systems with nested infection networks, Journal of theoretical biology 332 (2013) 65–77.

[25] J. O. Haerter, N. Mitarai, K. Sneppen, Phage and bacteria support mutual diversity in a narrowing staircase of coexistence, The ISME journal 8 (11) (2014) 2317–2326.

[26] N. Mitarai, S. Brown, K. Sneppen, Population dynamics of phage and bacteria in spatially structured habitats using phage λ and escherichia coli, Journal of bacteriology 198 (12) (2016) 1783–1793.

[27] R. S. Eriksen, N. Mitarai, K. Sneppen, Sustainability of spatially distributed bacteria-phage systems, bioRxiv (2018) 495671.

[28] J. S. Weitz, G. Li, H. Gulbudak, M. H. Cortez, R. J. Whitaker, Viral invasion fitness across a continuum from lysis to latency, bioRxiv (2019) 296897.

[29] V. Sinha, A. Goyal, S. L. Svenningsen, S. Semsey, S. Krishna, In silico evolution of lysis-lysogeny strategies reproduces observed lysogeny propensities in temperate bacteriophages, Frontiers in microbiology 8 (2017) 1386.

[30] J. Monod, The growth of bacterial cultures, Annual Reviews in Microbiology 3 (1) (1949) 371–394.

[31] M. De Paepe, F. Taddei, Viruses' life history: towards a mechanistic basis of a trade-off between survival and reproduction among phages, PLoS Biol 4 (7) (2006) e193.

[32] A. D. Kaiser, Mutations in a temperate bacteriophage affecting its ability to lysogenize escherichia coli, Virology 3 (1) (1957) 42–61.

[33] X. Shao, A. Mugler, J. Kim, H. J. Jeong, B. Levin, I. Nemenman, Growth of bacteria in 3-d colonies, arXiv preprint arXiv:1605.01098.

[34] R. S. Eriksen, S. L. Svenningsen, K. Sneppen, N. Mitarai, A growing microcolony can survive and support persistent propagation of virulent phages, Proceedings of the National Academy of Sciences 115 (2) (2018) 337–342.

[35] J. Fort, V. Méndez, Time-delayed spread of viruses in growing plaques, Physical review letters 89 (17) (2002) 178101.

[36] D. Ping, T. Wang, D. Fraebel, S. Maslov, K. Sneppen, S. Kuehn, Collapse and contingency in phage infections of migrating bacterial populations, in: APS Meeting Abstracts, 2019.

[37] K. A. Manning, N. Quiles-Puchalt, J. R. Penadés, T. Dokland, A novel ejection protein from bacteriophage 80α that promotes lytic growth, Virology 525 (2018) 237–247.

[38] S. Maslov, K. Sneppen, Well-temperate phage: optimal bet-hedging against local environmental collapses, Scientific Reports 5 (2015) 10523.

[39] B. Sauer, R. Calendar, E. Ljungquist, E. Six, M. G. Sunshine, Interaction of satellite phage p4 with phage 186 helper, Virology 116 (2) (1982) 523–534.

[40] V. V. Rezelj, L. I. Levi, M. Vignuzzi, The defective component of viral populations, Current opinion in virology 33 (2018) 74–80.

[41] K. A. S. Thompson, J. Yin, Population dynamics of an rna virus and its defective interfering particles in passage cultures, Virology journal 7 (1) (2010) 257.

[42] L. I. Rast, I. M. Rouzine, G. Rozhnova, L. Bishop, A. D. Weinberger, L. S. Weinberger, Conflicting selection pressures will constrain viral escape from interfering particles: principles for designing resistance-proof antivirals, PLoS computational biology 12 (5) (2016) e1004799.

[43] A. Baltes, F. Akpinar, B. Inankur, J. Yin, Inhibition of infection spread by co-transmitted defective interfering particles, PloS one 12 (9) (2017) e0184029.

[44] L. E. Bertani, The effect of the inhibition of protein synthesis on the establishment of lysogeny, Virology 4 (1) (1957) 53–71.

[45] C. Lee, J. Iandolo, Structural analysis of staphylococcal bacteriophage phi 11 attachment sites., Journal of bacteriology 170 (5) (1988) 2409–2411.

[46] L. Rudin, J.-E. Sjöström, M. Lindberg, L. Philipson, Factors affecting competence for transformation in staphylococcus aureus, Journal of bacteriology 118 (1) (1974) 155–164.

[47] R. W. Hendrix, R. L. Duda, Bacteriophage lambda papa: not the mother of all lambda phages, Science 258 (5085) (1992) 1145–1148.

Figures

Figure 1: **Strategies of SaPI-helper system and P4-P2 system.** (a) Pirate phage can only become lysogen when infecting a host alone, hence it cannot propagate on a lawn of uninfected host. (b) Helper phage is temperate, with frequency of lysogenization α_H. Typically helper phage forms a turbid plaque on a lawn of uninfected host. (c) In the SaPI-helper system (e.g. SaPIbov1-ϕ11), infection of pirate phage SaPI particle to a helper lysogen results in formation of double lysogen (frequency of lysogenization of pirate phage upon infection of helper lysogen α_P is one). Hence SaPI cannot propagate on a lawn of helper lysogen. When a pirate phage infects a SaPI lysogen and choose the lytic pathway with probability $1-\alpha_H$, the majority of the produced particles are pirate phages (fraction of pirate phage upon helper lysis decision $f_P^{(H)} \geq 0.9$), which inhibits the helper phage propagation on a pirate lysogen lawn. (d) In the P4-P2 system, infection of P4 phage to a P2 lysogen can results in the lysis with probability $1 - \alpha_P \sim 0.6$, and upon burst almost all of the phages produced are P4 (fraction of pirate phage upon pirate lysis decision $f_P \sim 1$), allowing pirate phage propagation on a helper lysogen lawn. On the other hand, P4 cannot take over much of P2 when P2 infection lyses a P4 lysogen ($f_P^{(H)} \sim 0.001$), which allows helper phage propagation on a pirate lysogen lawn.

Figure 2: **Simulated well-mixed population dynamics with (a) SaPI-helper (b) P4-P2 system.** The initial condition is set so that $API_P = API_H = 100$. The concentrations of sensitive bacteria B, the pirate lysogen L_p, the helper lysogen L_H, the double lysogen L_{PH}, the helper phage H, and the pirate phage P are shown by solid line, coarse dashed line, fine dashed line, dash-dotted line, dash-double dotted line, and long dashed line, respectively.

Figure 3: **The initial condition dependence of the helper and pirate phage gene spreading in well mixed population.** In the initial condition, the sensitive bacteria of 10^5/ml is mixed with the phage particles of various average phage input API_H and API_P and the population dynamics was simulated for until the system reaches the final state by using (a) the SaPI-helper system's parameter set ($f_P^{(H)} = 0.9$, $\alpha_P = 1$) or (b) the P4-P2 system's parameter set ($f_P^{(H)} = 0.001$, $\alpha_P = 0.4$, $f_P^{(P)} = 1$). The number of lysogen cells with helper phage prophage $L_H + L_{PH}$ (pirate phage prophage $L_P + L_{PH}$) in the final state normalized to the initial nutrient concentration 10^9/ml are visualized by color code in figure left (right) panel, respectively, where the higher value is shown with lighter color in log scale.

Figure 4: **Simulated plaque formation by (a-f) SaPI-helper and (g-l) P4-P2 system.** (a-f) and (g-l) represents the time evolution of the populations. The unit is the number of the cells / phage particles per $18\mu m^2$, corresponding to the number per microcolony. The profiles at 1h, 3h, 5h, and 7h are shown by solid line, coarse dashed line, fine dashed line, and dash-dotted line, respectively.

Figure 5: **The initial condition dependence of the helper and pirate phage gene spreading.** The average phage input in the initial spot, API_H and API_P is varied, and the plaque formation is simulated by using (a) the SaPI-helper system's parameter set ($f_P^{(H)} = 0.9$, $\alpha_P = 1$) or (b) the P4-P2 system's parameter set ($f_P^{(H)} = 0.001$, $\alpha_P = 0.4$, $f_P^{(P)} = 1$). The number of lysogen cells with helper phage prophage $L_H + L_{PH}$ (pirate phage prophage $L_P + L_{PH}$) integrated over the space out side of the initial spot ($r > 1$mm) and normalized to the maximum observed value 1×10^8 are visualized by color code in figure left panels (right panels), respectively, where the higher value is shown with lighter color.

Tables

Table 1: **Key parameters for pirate phage strategies**

	Meaning	SaPI-helper	P4-P2
α_P	Lysogen frequency upon pirate phage infection	1	0.4
$f_P^{(P)}$	Fraction of pirate phage when the pirate phage lysing helper lysogen	N.A.	1
$f_P^{(H)}$	Fraction of pirate phage when the helper phage lysing pirate lysogen	0.9	0.001

Table 2: **Default parameters used**

	Meaning	Default value	References, comments
η_H	Helper phage adsorption rate	$3.3 \times 10^3 \mu\mathrm{m}^3/\mathrm{h}$	P2 [31]
η_P	Pirate phage adsorption rate	$13 \times 10^3 \mu\mathrm{m}^3/\mathrm{h}$	P4 [31]
τ	latency time at fast growth	0.75 h	P2 latency 30min[14]$\sim$ 48min [31] P4 latency on P2 lysogen 60 min [14, 31]
β	Total burst size	100	P2 on uninfected host: 60[44]$\sim$ 160[31] P4 on P2 lysogen: 100[15]$\sim$300[31] ϕ11 on uninfected host: $\sim$100 [45]
α_H	Lysogenization frequency for a helper phage	0.15 for both systems	P2: $0.15 \sim 0.2$ [44] ϕ11: ~ 0.14 [46]
α_P	Lysogenization frequency when a pirate phage infects a helper lysogen	0.4 for P4-P2 1 for SaPI -helper	$0.3 \sim 0.5$ [13] 1 [5]
$f_P^{(H)}$	fraction of pirate phage from a helper phage lysing a pirate lysogen	0.001 for P4-P2 0.9 for SaPI-helper	Yield is less than one per cell [15] 0.9-0.99 for SaPI-helper systems [12]
$f_P^{(P)}$	fraction of pirate phage from a pirate phage lysing a helper lysogen	1 (N.A. for SaPI)	$10^{-5} \sim 10^{-3}$ P2 and 100 P4 [14] P2 level possibly due to the induction
δ	Phage decay rate	0.001/h	Small in the time scale of interest.
g_{max}	Fastest growth rate	2/h	Typical *E. coli* growth in rich medium
M	number of intermediate steps	10	30% variation in latency time [26].
n_0	Initial nutrient level	10^9 /ml	well-mixed case
k_s	Constant for Monod growth in the well mixed simulation	$n_0/5$	Depends on the nutrient.
N_0	Initial nutrient level	$30/\mu\mathrm{m}^2$	spacially structured case [26]
K_s	Constant for Monod growth in the plaque simulation	$N_0/5$	Depends on the nutrient.
D_H	Helper phage diffusion constant	$10^4 \mu\mathrm{m}^2/\mathrm{h}$	4-fold less than the value used for λ [26]. λ similar size to P2 [31] but λ was selected for bigger plaque [47].
D_P	Pirate phage diffusion constant	$D_H \times \eta_P/\eta_H$	The difference in the adsorption likely due to the difference in diffusion.
D_n	Nutrient diffusion constant	$4 \times 10^5 \mu m^2/h$	[26]