Effects of dietary taurine level on visual function in European sea bass

(*Dicentrarchus labrax*)

Richard W. Brill*, Andrij Z. Horodysky[2], Allen R. Place[3], Mary E.M. Larkin[4], Renate

Reimschuessel[3]

[1] Department of Fisheries Science, Virginia Institute of Marine Science, Gloucester Point,
Virginia, United State of America
11
12   [2] Department of Marine and Environmental Science, Hampton University, Hampton, Virginia,
13   United State of America
14
15   [3] Institute of Marine and Environmental Technology, University of Maryland Center for
16   Environmental Science, Baltimore, Maryland, United State of America
17
18   [4] Center for Veterinary Medicine, U.S. Food and Drug Administration, Laurel, Maryland, United
19   State of America
20

23   * Corresponding author

24   E-mail: rbrill@vims.edu (RB)

Abstract

Dietary insufficiencies have been well documented to decrease growth rates and survival

(and therefore overall production) in fish aquaculture. By contrast, the effects of dietary

insufficiencies on the sensory biology of cultured fish remains largely unstudied. Diets based

solely on plant protein sources could have advantages over fish-based diets, because of the cost

and ecological effects of the latter, but lack the amino acid taurine. Adequate levels of taurine

are, however, necessary for the development of a fully functional visual system in mammals. As

part of ongoing studies to determine the suitability of plant-based diets, we investigated the

effects of normal and reduced taurine dietary levels on retinal anatomy and function in European

sea bass (*Dicentrarchus labrax*). We could not demonstrate any effects of dietary taurine level

on retinal anatomy, nor the functional properties of luminous sensitivity or temporal resolution

(measured as flicker fusion frequency). We did, however, find an effect on spectral sensitivity.

The peak of spectral sensitivity of individuals fed a 5% taurine diet was rightward shifted (i.e.,

towards longer wavelengths) relative to that of fish fed a 0% or 1.5 % taurine diet. This

difference in in spectral sensitivity was due to a relatively lower level of middle wavelength

pigment (maximum absorbance $\approx$ 500 nm) in fish fed a 5% taurine diet. Changes in spectral

sensitivity resulting from diets containing different taurine levels are unlikely to be detrimental

to fish destined for market but could be in fishes that are being reared for stock enhancement

programs.

Introduction

Carnivorous fishes cannot synthesize many essential amino and fatty acids and must receive them through their diet [1,2]. Consequently, the aquaculture of these species requires diets based on, or at least supplemented with, protein sources from wild-caught fish. Up to two to five times more fish is, however, required to culture a product than is provided by it [3]. This increases the cost of production and can have ecological consequences as capture of wild-caught fish may impact forage fish populations and the higher trophic level species that feed on them [4,5]. An equally important concern is the potential for fishmeal to contain xenobiotic compounds such as polychlorinated biphenyls (PCB's) and mercury. These undergo biomagnification leading to elevated levels of contaminates in the endproduct. In brief, fishmeal diets for the aquaculture of carnivorous fishes are costly, and potentially unsustainable as well as unsafe. Purely plant-based diets circumvent these issues, but generally lack essential and semi-essential amino acids (e.g., taurine, methionine, lysine), as well as many vitamins and minerals needed in microquantities [6]. Plant-based diets are therefore currently supplemented with fishmeal or fish oil [7,8,9]. But we argue, as have others [10)], that there remains an exigent need to determine if purely plant-based diets can support survival rates, growth rates, and feeding efficiencies (i.e., the ratio of the mass of fish produced per mass of feed) necessary for the successful aquaculture of carnivorous fishes.

Omnivorous species have been the easiest to convert to low or no fishmeal diets; whereas marine carnivorous species have been the most difficult [11]. But it is the latter that are produced by aquaculture for restocking programs (i.e., to augment wild populations [12]). Diets differing in fatty acid composition can have significant impacts on the growth performance, energetics, cardiorespiratory physiology, hypoxia tolerances, and exercise and recovery performance of

fishes [13-17]. It is therefore plausible that the metabolic performance and hypoxia tolerances of

fishes fed plant-based diets may be significantly different – with major implications for the

suitability of fishes for re-stocking programs.

More specific to our project, plant protein sources lack taurine. Taurine is a sulfur

containing amino acid that is found in higher concentrations than any other free amino acid (i.e.,

amino acids not incorporated into any known proteins) and its roles in the proper development

and function in a variety of vertebrate tissues have received considerable attention [18-21].

Taurine is considered a conditionally indispensable amino acid for humans and non-human

primates and an essential amino acid in some mammalian carnivores (e.g., felines) [20], but little

attention has been paid to the required levels of this amino acid or its roles in fishes. We

hypothesize that most marine carnivorous fishes lack the little ability to synthesize taurine (due

to the large quantities found in their natural prey items) and therefore require it to be supplied in

the diet. We also hypothesize that plant-based diets may not allow for development of fully

functional visual system as a low taurine diet, or treatments with agonists of taurine uptake, have

been documented to impede the embryonic development of the retina and maintenance of normal

retinal function in mammals; the latter because of taurine's role as an antioxidant and osmolyte

(i.e., a compound maintaining intracellular osmotic balance) [20-24]. High concentrations of

taurine have, moreover, been found in the photoreceptor cells (i.e., rod and cone cells) and retinal

pigment epithelium in several teleost fish species [25-30]. Although it has been suggested that

taurine is only an osmolyte in retinal cells of fishes [26], other investigators have concluded that

the role of taurine in development and maintenance of retinal function is conserved throughout

the vertebrate order [21,31]. We therefore posit that diets containing inadequate levels of taurine

could result in diminished visual system function in carnivorous fishes. We recognize that less

than fully functional visual systems are unlikely to be detrimental to fish destined for market, yet

we contend that a diminished functionality of the visual system of fishes being cultured for stock

enhancement programs would decrease their survival and fitness (e.g., growth and reproduction)

relative to wild individuals. If this is the case, programs rearing fish for restocking would be less

able to meet their ultimate objective. Therefore, in conjunction with our ongoing study

examining the overall efficacy of formulations of our plant-based diet for generating acceptable

growth rates in European sea bass, we expanded our efforts to examine the effects of dietary

taurine level on visual function.

Materials and Methods

Our study was carried out under protocols approved by the Institutional Animal Care and

Use Committees of the University of Maryland Baltimore Medical School and the College of

William and Mary and followed all applicable laws and regulations. European sea bass were

obtained from the Aquaculture Research Center at the Institute of Marine and Environmental

Technology (IMET, Baltimore, MD). The average starting weight was ~15 g for fish

subsequently reared on the 5% taurine diet and ~25 g for fish subsequently reared on the 0 or

1.5% taurine diet. Fish were divided by diet and housed in eight-foot diameter, four cubic meter

recirculating systems with shared mechanical and life support systems. The latter included a

protein skimmer, ozonation, mechanical filtration (in the form of bubble-bead filters), and

biological filtration. Water quality (measured two to three times per week) was not significantly

different between systems (ANOVA, $p>0.05$). Mean ($\pm$ SEM) water quality values in the tanks

were: dissolved oxygen 5.7 ± 1.6 mg L^{-1}, temperature 27 ± 2 ^{o}C, pH 7.6 ± 0.3, total ammonia

nitrogen (NH_3) 0.06 ± 0.06 mg L^{-1}, nitrite (NO_2^-) 0.12 ± 0.08 mg L^{-1}, nitrate (NO_3^-) 49 ± 9 mg

L^{-1}, alkalinity 96 ± 23 meq L^{-1}, and salinity 25 ± 2 ppt. Fish were fed 3.5% of their body

weight per day and maintained on each specific diets for five to six months.

Diet preparation

122   The three diets formulations (Table 1) were prepared by Zeigler Bros. (Gardners, PA,

123 USA) and analysis of their proximate composition (Table 2) was performed by New Jersey Feed

124 Laboratory, Inc. (Ewing Township, NJ, USA).

125
126 **Table 1. Formulations for the three experimental diets.**
127

Constituent	0% taurine	1.5% taurine	5% taurine
Profine VF	28.75	28.75	28.75
Soybean meal, 47.5%	23.33	23.33	23.33
Wheat flour, bagged	16.54	15.04	11.54
Corn gluten, 60%	15.34	15.34	15.34
Menhaden gold oil, top-dressed	5.96	5.96	5.96
Monocalcium phosphate FG	3.95	3.95	3.95
Lecithin FG	3	3	3
L-Lysine, 98.5%	0.75	0.75	0.75
Choline chloride, 70%	0.6	0.6	0.6
Potassium chloride FG	0.56	0.56	0.56
DL-Methionine, 99%	0.45	0.45	0.45
Sodium chloride	0.28	0.28	0.28
Vitamin C	0.2	0.2	0.2
Premix AquaVit	0.12	0.12	0.12
Premix Aquamin Fish	0.12	0.12	0.12
Magnesium oxide FG	0.05	0.05	0.05

Taurine FG	0	1.5	5

128   FG = food grade

129
130   **Table 2. Proximate composition (%) of three experimental diets.**
131

Constituent	0% taurine	1.5% taurine	5% taurine
Moisture	8.05	6.97	9.5
Protein	43.82	45.02	46.65
Fat	5.94	6.19	6.94
Fiber	3.06	1.95	2.25
Ash	7.08	7.75	7.71
Taurine	0.07	1.48	4.85

Tissue sampling for analysis retinal anatomy

Eyes were harvested from fish not used in the ERG experiments. Food was withheld for

24 hours and 10-12 fish from each diet were anesthetized in a bath containing 25 mg L^{-1} MS-222

(Syndel, Ferndale, WA, USA) buffered with 50 mg L^{-1} sodium bicarbonate (Sigma-Aldrich, St.

Louis, MO, USA). Upon removal from the anesthesia bath, the spinal cord was immediately

severed, and eyes harvested. Retinal tissue was subsequently embedded, sectioned, mounted on

glass slides, and stained with Hematoxylin-Eosin using standard histological procedures. Images

were captured using a BX53 microscope outfitted with a DP73 camera and visualized with the

cellsSens software (version 510) (Olympus, Center Valley, PA, USA).

Retinal responses to light stimuli

Fish were transferred to the Virginia Institute of Marine Science (Gloucester Point, VA)

where whole-animal corneal electroretinography (ERG) was used to assess three standard

metrics of retinal function: (1) luminous sensitivity (V log I response), (2) flicker fusion

frequency (FFF, a measure of temporal resolution), and (3) spectral sensitivity (32-36).

Anesthesia and handling methodologies were as described previously (33-35). Teflon-coated,

silver – silver chloride, 0.5mm wire electrodes were used to measure ERG potentials: the active

electrode was placed on the corneal surface and a reference electrode in the nasal cavity. All

subjects were dark-adapted for a minimum of 60 minutes prior to visual trials. Electrode

placements, as well as any further modifications to the experimental setup, were conducted under

a dim red LED light source (peak wavelength of 660 nm) that is beyond the spectral sensitivity

of European sea bass.

Luminous sensitivity was assessed using stimulus intensities covering six orders of

magnitude using a collimated white LED source and neutral density filters progressing from

subthreshold to saturation intensity levels in 0.2 log unit steps. FFF was assessed by measuring

the ability of the retinal responses to track sinusoidally modulated white light stimuli ranging in

frequency from 1 Hz (0 log units) to 100 Hz (2.0 log units), presented in increments of 0.2 log

unit frequency steps. FFF was measured at stimulus intensities of 25%, 50% and 100% of the

maximum response, as well as fixed light levels (log I = 1.9, 2.7, and 3.7; with I in units of

candela per m^2). Spectral sensitivity was assessed using stimuli over wavelengths from the

ultraviolet (300 nm) to the near infrared (700 nm) presented sequentially in 10 nm steps.

Monochromatic light flashes (50% bandwidth = 5 nm) were made approximately equally quantal

through a series of neutral density filters and subsequently corrected to predict isoquantal

responses, as described previously (33-36). To form hypotheses regarding the number and

spectral distribution of visual pigments present, and the effects of dietary taurine levels on the

distribution of pigments contributing to spectral sensitivity, we fitted the SSH [37] and GFRKD

[38] vitamin A1 rhodopsin absorbance templates separately to the photopic spectral sensitivity

data. Estimates of the unknown model parameters (λ_{max} values and their respective weighting

proportions) were derived by fitting the summed curves to the ERG data using maximum

likelihood. We objectively selected the appropriate template (SSH or GFRKD) and number of

contributing pigments using an information theoretic approach following Akaike's information

criterion that is a parsimonious measure that strikes a balance between model simplicity and

complex overparameterization (39). All parameter optimization, template fitting and model

selection was conducted using the software package R version 3.2.2 (R Development Core

Team).

Statistical tests comparing the effects of dietary taurine levels on the magnitude of

responses to various light levels were done using Sigmaplot (version 11.2, Systat Software, San

Jose, CA). Analysis of data were performed using a one-way analysis of variance (ANOVA) test

on means when the data were normally distributed, and a Kruskal-Wallis one-way analysis of

variance on ranks when the data were not. Comparisons the magnitude of responses to various

light levels (i.e., luminous sensitivity) tests were limited to the effects of diet within a given light

levels. Statistical tests comparing the effects of dietary taurine levels on FFF were preformed

using the two-way repeated measures ANOVA procedure in Sigmaplot, with the Holm-Sidak

method to conduct all pairwise multiple comparisons.

Results and discussion

Retinal morphology

There were no obvious effects of dietary taurine level on the morphology of identifiable

layers in European sea bass retina (Fig 1) not retinal cell layer thickness ratios (Fig 2) equivalent

to the massive disruption seen in retinal tissue of domestic cats fed low taurine diets [40].

Fig 1. Identifiable layers in European sea bass retinas. Panel A is a representative retina from

fish fed a 0% taurine diet, panel B is a representative retina from fish fed a 5% taurine diet.

G=ganglion cell layer, IP=inner plexiform layer, IN=inner nuclear layer, OP=outer plexiform

layer; ON=outer nuclear layer, C=cone photoreceptors, OS=outer segments of the photoreceptor

layer, PE=pigmented epithelium.

Fig 2. Retinal cell layer thickness ratios in European sea bass fed diets with 0 or 5% taurine

(open and cross hatched boxes, respectively). The boundary of the box closest to zero indicates

the 25[th] percentile, the line within the box marks the median, and the boundary of the box

farthest from zero indicates the 75[th] percentile. Whiskers (error bars) below and above the box

indicate the 10[th] and 90[th] percentiles, respectively. Data points above and below the whiskers are

considered outliers. T= total retina thickness, C= cone photoreceptors layer thickness, ON=outer

nuclear layer thickness, OS= thickness of the outer segments of photoreceptor layer, IN=inner

nuclear layer thickness (µm).

Retinal function

Retinal responses to increasing light levels showed the expected steep increases up to

those levels producing maximum response (Fig 3). When light levels are expressed in log units,

retinal response curves were the expected sigmoidal shape (Fig 3 insert). There was only one

significant effect of dietary taurine levels on luminous sensitivity in European sea bass, and only

at light levels needed to produce a response 75% of maximum (Fig 4). In other words,

significantly higher light levels were required to achieve responses above approximately 50% of

maximum in fish fed a diet lacking taurine. Flicker fusion frequencies generally showed the

expected increases with increasing light intensities, and dietary taurine level had no influence on

FFF when comparisons are made at the same light level (Fig 5).

Fig 3. Luminous sensitivity (i.e., intensity–response curves) of European sea bass. Data are

mean values (± SEM). Response values for individuals were normalized to 0-100%, the values

averaged, and the mean values rescaled to 0-100%. The inset shows the same data, but with light

intensities expressed in log units. Increases in light intensity were in 0.2 log unit steps.

Fig 4. Light levels needed to produce a given response relative to that of the maximin

response in intensity–response curves (Fig 3). The boundary of the box closest to zero

indicates the 25[th] percentile, the line within the box marks the median, and the boundary of the

236 box farthest from zero indicates the 75th percentile. Whiskers (error bars) above and below the

237 box indicate the 90th and 10th percentiles. Data points above and below the 90th and 10th

238 percentiles are considered outliers and are not used to determine median values.

241 **Fig 5. Flicker fusion frequencies (i.e., the highest frequency of sinusoidal light stimulus**

242 **detectable) with increasing light intensities of European seabass.** The light levels used

243 correspond to approximately those needed to produce responses approximately 25%, 50% and

244 75% of the maximum response in the intensity–response curves (Fig 3). The symbols #, *, and ^

245 indicate differences in flicker fusion frequencies between log light levels 1.9 and 2.7, between

246 log light levels 1.9 and 3.7, and between log light levels 2.7 and 3.7, respectively. The boundary

247 of the box closest to zero indicates the 25th percentile, the line within the box marks the median,

248 and the boundary of the box farthest from zero indicates the 75th percentile. Whiskers (error bars)

above and below the box indicate the 90[th] and 10[th] percentiles. Data points above and below the

90[th] and 10[th] percentiles are considered outliers and are not used to determine median values.

The spectral sensitivity of fish fed diets with reduced taurine levels (i.e., 0 and 1.5%)

were not significantly different from each other (Fig 6, panels A and B) and these data were

subsequently combined. The results from fitting the data to both SSH and GFRKD rhodopsin

templates suggested that European seabass are trichromats (i.e., have three visual pigments)

(Tables 3 and 4). The SSH rhodopsin template was clearly best fitting (i.e., had the lowest AIC

value) for the combined data from fish fed diets with reduced taurine levels (Table 3). In the case

of data from fish fed 5% taurine diet (Table 4), the AIC values (i.e., goodness of fits) for SSH

rhodopsin templates for dichromats (i.e., two retinal pigments) with secondary absorbing peaks

(i.e., β bands) on both the visual pigments, and the SSH and GFRKD rhodopsin templates for

trichromats, were indistinguishable (ΔAIC < 5). The SSH template for a dichromat with

secondary absorbing peaks on both pigments predicted, however, the short wavelength pigment

to have maximum absorbance in the UV wavelength (376 nm) which we consider unreasonable

given the results from ERG data fitted similarly from a variety of inshore fishes (34-35, 41). We

therefore based our subsequent conclusions on the effects diet on SSH rhodopsin templates for

trichromats.

Table 3. The spectral sensitivity of European seabass fed diets with reduced taurine levels
(i.e., 0 and 1.5%) where the data from fish in these two groups have been combined.
272

Condition	Template	$\lambda_{max,1}$	$\lambda_{max,2}$	$\lambda_{max,3}$	-log(L)	p	AIC	ΔAIC	Pigment weights
Mono	GFRKD	574			-32.1	8	-58	169	
	SSH	575			-26.5	8	-47	180	
Di, α	GFRKD	410	510		-91.3	8	-172	55	
	SSH	411	510		-91.0	8	-173	55	
Di, β, S	GFRKD	498	603		-79.7	8	-147	80	
	SSH	504	604		-97.7	8	-183	44	
Di, β, L	GFRKD	439	513		-79.5	8	-147	80	
	SSH	461	513		-96.5	8	-181	46	
Di, β, B	GFRKD	509	601		-67.2	8	-120	107	
	SSH	527	605		-82.0	8	-150	77	
Tri, α	GFRKD	388	501	603	-97.4	8	-181	47	
	SSH	389	512	611	120.7	8	-227	0	0.24, 0.91, 1

274   **Table 4. The spectral sensitivity of European seabass fed a 5% taurine diet.**
275

Condition	Template	$\lambda_{max,1}$	$\lambda_{max,2}$	$\lambda_{max,3}$	-log(L)	p	AIC	ΔAIC	Pigment weights
Mono	GFRKD	586			-52.2	8	-98	92	
	SSH	588			-44.6	8	-83	108	
Di, α	GFRKD	485	597		-83.7	8	-155	35	
	SSH	496	598		-96.9	8	-182	9	
Di, β, S	GFRKD	485	597		-83.7	8	-155	35	
	SSH	496	598		-96.9	8	-182	9	
Di, β, L	GFRKD	439	513		-79.5	8	-147	44	
	SSH	461	513		-96.5	8	-181	10	
Di, β, B	GFRKD	506	597		-76.6	8	-139	52	
	SSH	376	599		-102.4	8	-191	0	
Tri, α	GFRKD	382	492	597	-101.4	8	-189	2	
	SSH	380	497	598	-101.3	8	-189	2	0.17, 0.32, 1

276
277

Fig 6. Spectral sensitivity (i.e., responses to monochromatic light) of European sea bass.

Data are mean values (± SEM). Response values for individuals were normalized to 0-100%, the

values averaged, and the mean values rescaled to 0-100%. Panel A are the differences between

mean values of responses of fish fed the three different diets. Panel B are the mean responses of

fish feed diets containing 0%, 1.5% and 5% taurine. In both panels, fish feed diets containing

0%, 1.5% and 5% taurine are indicated by filled circles, open circles and filled triangles,

respectively. Because the spectral sensitivities of fish fed diets containing 0% or 1.5% taurine

were largely not different, the data from these individuals were combined and the mean relative

responses shown in panels B and C. Wavelengths at maximal absorptions (λ_{max} values) and

pigment-specific weight values (to the right of each λ_{max} value) are shown. Black lines are the

summed curves of the visual pigment curves multiplied by their respective weighting factors.

Color coded lines indicate the absorptive characteristics of individual visual pigments. The color

bar at the top of the figure shows the approximate human visual spectrum.

Our results suggested that the visual pigments of European seabass fed diets containing 0

or 1.4% taurine have maximal absorbance peaks (λ_{max}) of 389, 512, 611 nm (P$_1$, P$_2$, P$_3$,

respectively; Fig 6, panel C). The maximal absorbance peaks of the visual pigments in fish fed a

diet containing 5% taurine had maximal absorbance peaks (λ_{max}) of 380, 497, 598 nm (P$_1$, P$_2$, P$_3$,

respectively; Fig 6, panel D). The most significant differences were a significant effect of dietary

taurine level on the relative abundance of the three visual pigments (Fig 6, panel C). More

specifically, in fish fed 0 and 1.5% taurine diets, the ratio of the middle and long wavelength

visual pigments (P_2 and P_3, respectively) were approximately equal (0.91 to 1,) with a reduced

abundance of the short wavelength pigment (P_1) relative to the most abundant longwave length

pigment (P_3) (0.24 to 1) (Fig 6, panel C). By contrast, in fish fed a 5% taurine diet, there was a

reduction in the abundance of the middle wavelength pigment (P_2) relative to the long

wavelength pigment (P_3) (0.32 to 1) (Fig 6, panel D), while the ratio of the abundance of the

short wavelength pigment (P_1) relative to the most abundant longwave length pigment (P_3) was

relatively unchanged (0.17 to 1) (Fig 6, panel D). The overall effects of the changes in

abundance of visual pigments was that the peak in overall retinal spectral sensitivity was shifted

to longer wavelengths, from ≈ 530 nm in fish fed the 0% or 1.5% taurine diets (Fig 6, panel C), to

≈ 600 nm in fish fed the 5% taurine diet (Fig 6, panel D).

Replacing fishmeal with plant-based diets is a priority for the aquaculture industry, and

the development of optimized diets is underway [10,42]. Plant-based diets have a high protein

content [43], but commercial feed formulations often vary based on the availability and cost of

ingredients, with different batches containing significantly different proportions or levels of

quality of ingredients, or different ingredients all together. For this reason, the assessment of

multiple ingredients and ingredient combinations is necessary. To address this issue, we

formulated a diet based on available and cost-effective plant ingredients which are effective

fishmeal replacements in rainbow trout (*Oncorhyncus mykiss*) [44-45] and are highly digestible

by European sea bass [46]. But, as with other plant-based diets, our formulation lacks taurine.

There is evidence that, although that taurine can be recycled through the taurine transporter and

biliary recycling pathways, constant dietary supply is required to maintain proper function

throughout multiple tissue types [20]. We assumed that taurine's role as a photoreceptor

protectant would most likely be as an antioxidant, similar to the role it plays in the liver of

Atlantic salmon challenged with pro-oxidants like cadmium chloride [21,47]. We therefore

hypothesized that visual disparities between European seabass fed diets containing 0%, 1.5% and

5% taurine would be due to differences in photoreceptor and pigment repair.

We did not, however, see a reduction of luminous sensitivity resulting from reduced

dietary taurine (Fig 2). In contrast, other investigators have demonstrated a large progressive

decrease in ERG amplitudes to a range of light intensities (12-197 cd m^{-2}, log light level 1.1 to

2.3) in rats fed a soybean-based diet (containing a negligible amount of taurine) and treated with

guanidinoethyl sulfonate (a compound that depletes of taurine) over 15 weeks [23]. Similar

decreases in ERG amplitudes have been recorded in cats fed a taurine deficient (casein-based)

diet (40). These results (in comparison to ours) imply a loss of luminous sensitivity resulting

from a low taurine diet is related to a reduction in the cellular defense mechanisms countering

light- and oxygen induced damage in retinal receptor cells [24], and that taurine deficiency has

different effects in mammals and teleost fishes.

Effective visual function at low light levels requires both spatial and temporal summation

within the retina [50-51]. The former results from the convergence of photoreceptors onto

bipolar cells, as seen in fishes occupying dimly lit environments [52-54]. Spatial summation

therefore reduces visual acuity (i.e., the ability to detect the details of an object). Demonstrating

anatomical changes in spatial summation (i.e., changes in the convergence of photoreceptors

onto bipolar cells) was, however beyond the scope of this study. Temporal summation refers to

photoreceptors responding slowly to flashing light as evinced by temporal resolutions (i.e.,

reducsed FFF). Low FFF implies a blurring of fast-moving objects or the inability to detect them

at all [50-51, 55-57]. We did specifically investigate the effects of dietary taurine level on flicker

fusion frequency (Fig 5) but could we could not demonstrate any effects of dietary taurine level.

We therefore conclude that there was no effect of dietary taurine level on retinal anatomy in

terms of the degree of the convergence of receptor cells onto bipolar cells.

In terms of spectral sensitivity, our results imply that the European seabass visual system

has three visual pigments and, more importantly, that spectral sensitivity can be impacted by diet

taurine content (Fig 6). To the best of our knowledge, we are the first to demonstrate this

phenomenon. Our results could be due to: (1) changes in the relative abundance of the three

visual pigments equivalent to the seasonal shifts seen in other teleost fishes [58-60], or (2)

changes in the amino acid composition of opsin proteins (which influence peak wavelength

absorbancies of visual pigments) that result from changes in amino acid composition of opsin

proteins. Changes in amino acid composition of opsin proteins (due to changes in gene

expression) have been shown to occur over ontogeny, through adaptation of individuals to

different light spectra, or changes diurnal lighting patterns [61-67]. Our data do not allow us to

differentiate between changes in the relative abundance of visual pigments or changes in amino

acid composition of opsin proteins. Our results do, however, demonstrate that there is a threshold

effect of dietary taurine in European seabass, in that there were no discernable alterations in

spectral sensitivity of fish fed 0% and 1.5% taurine diets, but a clear difference spectral

sensitivity of fish fed a 5% taurine diet. The net effect of reduced taurine diets was a broader

overall spectral sensitivity, with the peak in spectral sensitivity being leftward-shifted (a greater

sensitivity to shorter wave lengths).

We note, however, that one of reasons for undertaking this study was to determine if

fishes being cultured for stock enhancement programs fed a plant-based diet lacking sufficient

taurine could decrease survival and fitness following release. If so, programs rearing fish for

restocking would be less able to meet their ultimate objective of enhancing population

abundance. Our demonstration that reduced dietary taurine levels influence spectral sensitivity

(Fig 6) implies that this is the case. This conclusion is congruent with both the visual pigment

sensitivity hypothesis and the contrast sensitivity hypothesis. The former posits that for optimal

visual function the absorbance of visual pigments must correspond to the spectral distribution of

the light environment [68-70], whereas the latter contends that aquatic animals maximize

contrast sensitivity to acquire information effectively in a turbid (i.e., low contrast) environment

[71-73]. Under both situations, however, changes in the relative abundance of visual pigment is

likely to reduce fitness in cultured fish released for restocking. It unknown, however, if these

deficits would remain following release of cultured, fish or if the visual properties of fish fed 0%

and 1.5% taurine diets we observed would converge onto those of fish fed a 5% taurine diet after

the switch to a natural prey diet. These questions are worth investigating.

380
381
382　# Conclusions
383
We could not demonstrate any effects of dietary taurine level on retinal anatomy, the

functional properties of luminous sensitivity nor temporal resolution (i.e., FFF) in European sea

bass. We did, however, find an effect on spectral sensitivity. The peak of spectral sensitivity of

individuals fed a 5% taurine diet was rightward shifted (i.e., towards longer wavelengths)

relative to that of fish fed a 0% or 1.5 % taurine diet. This difference in spectral sensitivity was

caused by a relatively lower level of middle wavelength pigment (maximum absorbance $\approx$500

nm) in fish fed a 5% taurine diet. Changes in spectral sensitivity resulting from diets containing

different taurine levels are unlikely to be detrimental to fish destined for market, but could be in

fishes that are being reared for stock enhancement programs.

393
394

Acknowledgements

The authors would like to thank the staff of the Aquaculture Research Center at the Institute of

Marine and Environmental Technology, including Steve Rodgers and Chris Tollini. This is

contribution XXXX from the Virginia Institute of Marine Science. This publication was made

possible by the National Oceanic and Atmospheric Administration, Office of Education

Educational Partnership Program award number (NA11SEC4810002). Its contents are solely the

responsibility of the award recipient and do not necessarily represent the official views of the

U.S. Department of Commerce, National Oceanic and Atmospheric Administration. The views

expressed in this article are those of the authors and may not reflect the official policy of the

Department of Health and Human Services, the U.S. Food and Drug Administration, or the U.S.

Government.

References

1. Silva SS, Anderson TA. Fish nutrition in aquaculture. London: Chapman & Hall; 1995.

2. Lunger AN, Craig SR, McLean E. Replacement of fish meal in cobia (*Rachycentron*

canadum) diets using an organically certified protein. Aquaculture 2006;257:393-399

3. Tacon A, Metian M. Global overview on the use of fish meal and fish oil in   industrially

compounded aquafeeds: Trends and future prospects. Aquaculture 2008;285: 146-158.

4. Pauly D, Christensen V, Dalsgaard J, Froese R, Torres F Jr. Fishing down marine food webs.

Science 1998;279:860-863.

5. Naylor RL, Goldburg RJ, Primavera JH, Kautsky N, Beveridge MC, Clay J, et al. Effect of

aquaculture on world fish supplies. Nature 2000;405:1017-1024.

6. Gatlin DM, Barrows FT, Brown P, Dabrowski K, Gaylord TG, Hardy RW et al. Expanding

the utilization of sustainable plant products in aquafeeds: a review. Aquac Res. 200;738:551-

579.

7. Barrows FT, Gaylord TG, Sealey, W.M., Smith, C.E., Porter, L. Supplementation of plant-

based diets for rainbow trout (*Oncorhynchus mykiss*) with macro-minerals and inositol.

Aquac Nutr. 2010;16:654-661.

8. Gaylord TG, Barrows FT, Teague AM, Johansen KA., Overturf KE, Shepherd B.

Supplementation of taurine and methionine to all-plant protein diets for rainbow trout

(*Oncorhynchus mykiss*). Aquaculture 2007;269:514-524.

9. Zhou QC, Wu Z, Chi S, Yang Q. Dietary lysine requirement of juvenile cobia (*Rachycentron*

canadum). Aquaculture 2007;273:634-640.

10. Turchini GM, Trushenski JT, Glencross BD. Thoughts for the future of aquaculture nutrition:

realigning perspectives to reflect contemporary issues related to judicious use of marine

resources in aquafeeds. N Am J Aquacult. 2018;DOI:10.1002/naaq.10067.

11. Fernandez-Jover D, Arechavala-Lopez P, Martinez-Rubio L, Tocher DR, Bayle-Sempere JT,

Lopez-Jimenez JA, et al. Monitoring the influence of marine aquaculture on wild fish

communities: benefits and limitations of fatty acid profiles. Aquacult Environ Interact.

2001;2:39-47.

12. Lorenzen K, Agnalt A-LA, Blankenship HL, Hines AH, Leber KM, Loneragan NR, Taylor

MD. Evolving context and maturing science: aquaculture-based enhancement and restoration

enter the marine fisheries management toolbox. Rev Fish Sci. 2013;21:213-221.

13. Agnisola C, McKenzie DJ, Taylor EW, Bolis CL, Tota B. Cardiac performance in relation to

oxygen supply varies with dietary lipid composition in sturgeon. Am J Physiol.

1996;271:R417-425.

14. McKenzie DJ. Effects of dietary fatty acids on the respiratory and cardiovascular physiology

of fish. Comp. Biochem Physiol A Mol Integr Physiol. 2001;128:607-621.

15. McKenzie DJ, Piraccini G, Steffensen JF, Bolis CL, Bronzi P, Taylor EW. Effects of diet on

spontaneous locomotor activity and oxygen consumption in Adriatic sturgeon (*Acipenser*

naccarii). Fish Physiol Biochem. 1995:14;341-55.

16. Randall DJ, Mckenzie DJ, Abrami G, Bondiolotti GP, Natiello F, Bronzi P, Bolis L, Agradi

E. Effects of Diet on Response to Hypoxia in Sturgeon (*Acipenser naccarii*). J Exp Biol.

1992:125;113-125.

17. Wagner GN, Balfry SK, Higgs DA, Lall SP, Farrell AP. Dietary fatty acid composition

affects the repeat swimming performance of Atlantic salmon in seawater. Comp Biochem

Physiol A Mol Integr Physiol. 2004:137;567-76.

18. Ripps H, Shen WK. Review: Taurine: A "very essential" amino acid. Mol Vis 2012;18:

2673-2686.

19. Schaffer SW, Jong CJ, Ramila KC, Azuma J. Physiological roles of taurine in heart and

muscle. J Biomed Sci. 2010:S2;1-2.

20. Schuller-Levis GB, Park E. Taurine: new implications for an old amino acid. FEMS

Microbiol Lett. 2003:226;195-202.

21. L'Amoreaux W. The roles of taurine in the retina. In: Taurine in Health and Disease, El

Idrissi A, L'Amoreaux W, editors. Kerala, India: Transworld Research Network; 2015. p.

215-254. ISBN: 978-81-7895-520-9.

22. Hillenkamp J, Hussain AA, Jackson TL, Constable PA, Cunningham JR, Marshall J.

Compartmental analysis of taurine transport to the outer retina in the bovine eye. Invest

Ophth Vis Sci. 2004:45;4099-4105.

23. Lake N. Electroretinographic deficits in rate treated with Guanidinoethyl sulfonate, a

depletory of taurine. Exp Eye Res. 1986:42;87-91.

24. Lake N, Malik N. Retinal morphology in rates treated with a taurine transport antagonist.

Exp Eye Res. 1987:44;331-346.

25. Gisselmann G, Galler A, Friedrich F, Hatt H, Bormann J. Cloning and functional

characterization of two glycine receptor alpha-subunits from the perch retina. Eur J Neurosci.

2002:16;69-80.

26. Marc RE, Cameron D. A molecular phenotype atlas of the zebrafish retina. J Neurocytol.

2001:30;593-654.

27. Negishi K, Drujan BD. Effects of some amino acids on horizontal cells in the fish retina. J

Neurosci Res. 1979:4;351-263.

28. Nusetti S, Obregón F, Quintal M, Benzo Z, Lima L. Taurine and zinc modulate outgrowth

from goldfish retinal explants. Neurochem Res. 2005:30:1483-1492.

29. Omura Y, Yoshimura R. Immunocytochemical localization of taurine in the developing

retina of the lefteye flounder *Paralichthys olivaceus*. Arch Histol Cytol. 1999:62;441-446.

30. Omura Y, Hach A, Furukawa E, Ueck M, Lake N. Immunocytochemical localization of

taurine in the pineal organ and retina of an anadromous fish, *Plecoglossus altivelis*. Arch

Histol Cytol. 1997:60;153-162.

31. Lombardini JB. Taurine: retinal function. Brain Res Rev. 1991:16;151-169.

32. Fritsches KA, Brill RW, Warrant EJ. Warm eyes provide superior vision in swordfishes. Curr

Biol. 2005:15;55-58.

33. Brill R. Magel C, Davis M, Hannah R, Rankin P. Effects of rapid decompression and

exposure to bright light (simulated capture) on the visual function of black rockfish (*Sebastes*

melanops) and Pacific halibut (*Hippoglossus stenolepsis*). Fish Bulletin. 2008:106;427-437.

34. Horodysky AZ, Brill RW, Warrant EJ, Musick JA, Latour RJ. Comparative visual function in

five sciaenid fishes inhabiting Chesapeake Bay. J Exp Biol 2008:211;3601-3612.

35. Horodysky AZ, Brill RW, Warrant EJ, Musick JA, Latour RJ. Comparative visual function in

four piscivorous fishes inhabiting Chesapeake Bay. J Exp Biol. 2010:213; 1751-1761.

36. Kalonoski M, Hirons A, Horodysky A, Brill R. Spectral sensitivity, light sensitivity, and

temporal resolution of the visual systems in three sympatric coastal shark species: sandbar

shark (*Carcharhinus plumbeus*), smooth dogfish (*Mustelus canis canis*), and spiny dogfish

(*Squalus acanthias*). J Comp Physiol B. 2014:200;997-1013.

37. Stavenga DG, Smiths RP, Hoenders BJ. Simple exponential functions describing the

absorbance bands of visual pigment spectra. Vis Res. 1993:33;1011-1017.

38. Govardovskii VI, Fyhrquist N, Reuter T, Kuzmin DG, Donner K. In search of the visual

pigment template. Vis Neurosci. 2000:17;509-528.

39. Burnham KP, Anderson DR. Model selection and multimodel inference: a practical

information-theoretic approach. New York: Springer; 2002.

40. Schmidt SY, Berson EL, Watson G, Huang C. Retinal degeneration in cats fed casein III.

Taurine deficiency and ERG amplitudes. Invest Ophthalmol Vis Sci. 1977:16;673-678.

41. Horodysky, A. Z., R. W. Brill, K. C. Crawford, E. S. Seagroves, and A. K. Johnson..

Comparative visual ecophysiology of mid‒Atlantic temperate reef fishes. Biol. Open.

2013:2;1371-1381.

42. Rust MB, Barrows FT, Hardy RW, Lazur AL, Naughten K, Silverstein J. Alternative feeds

initiative. The future of aquafeeds. National Oceanic and Atmospheric Administration /

United State Department of Agriculture 2011:NOAA Tech. Memo. NMFS–F/SPO–124.

43. Hardy RW. New developments in aquatic feed ingredients, and potential of enzyme

supplements. In: Cruz-Saurez LE, Ricque-Marie D, Tapia-Salazar M, Olvera-Novoa MA,

Civera-Cerecedo R, editors. Avances En Nutrition Acuicola V. Memorias Del V Simposium

Internacional de Nutricion Acuicola. 2000 Merida, Yucutan, Mexico, p. 216-226.

44. Barrows FT, Gaylord TG, Sealey WM, Porter L, Smith CE. The effect of vitamin premix in

extruded plant-based and fish meal based diets on growth efficiency and health of rainbow

trout, *Oncorhynchus mykiss*. Aquaculture. 2008:283;148-155.

45. Palti Y, Silverstein JT, Wieman H, Phillips JG, Barrows FT, Parsons JE. Evaluation of

family growth response to fishmeal and gluten-based diets in rainbow trout (*Oncorhynchus*

mykiss). Aquaculture. 2006:255;548-556.

46. Watson AM, Kissil GW, Barrows FT, Place AR. Developing a plant-based diet for cobia

Rachycentron canadum. Int. Aquafeed. 2012:15;34-38.

47. Espe M, Holen E. Taurine attenuates apoptosis in primary liver cells isolated from Atlantic

salmon (*Salmo salar*). British J Nutritio. 2013:110;20-28.

48. Hayes KC, Rabin AR, Berson EL. An ultrastructural study of nutritionally induced and

reversed retinal degeneration in cats. Am J Pathol. 1975a:78;504-524.

49. Hayes KC, Carey RC, Schmidt SY. Retinal degeneration associated with taurine deficiency

in the cat. Science 1975b:188;949-951.

50. Warrant EJ. Seeing better at night: life style, eye design and the optimum strategy of spatial

and temporal summation. Vis Res. 1999:39;1611-1630.

51. Warrant E. Vision in the dimmest habitats on earth. J Comp Physiol A. 2004:190;765-789.

52. Fritsches KA, Marshall NJ, Warrant EJ. Retinal specializations in the blue marlin: Eyes

designed for sensitivity to low light levels. Mar Freshwater Res. 2003:54;333-341.

53. Lisney TJ, Collin SP. Retinal ganglion cell distribution and spatial resolving power in

elasmobranchs. Brain Behav Evol. 2008:72;59-77.

54. Litherland L, Collin SP. Comparative visual function in elasmobranchs: spatial arrangement

and ecological correlates of photoreceptor and ganglion cell distributions. Vis Neurosci.

2008:25:549-561.

55. Frank TM. Comparative study of temporal resolution in the visual systems of mesopelagic

crustaceans. Biol Bull. 1999:196:137-144.

56. Frank TM. Temporal resolution in mesopelagic crustaceans. Philos Trans R Soc.

2000:355;1195-1198.

57. Frank TM. Effects of light adaptation on the temporal resolution of deep-sea crustaceans.

Integr Comp Biol. 2003:43;559-570.

58. Wald G. The significance of vertebrate metamorphosis. Science 1958:128;1481-1490.

59. Dartnall HJA, Lythgoe JN. The spectral clustering of visual pigments. Vision Res.

1965:5;81-100.

60. McFarland WN, Allen DM. The effect of extrinsic factors on two distinctive rhodopsin-

porphyropsin systems. Can J Zool. 1977:55;1000-1009.

61. Cheng CL, Flamarique IN. Opsin expression: new mechanism for modulating colour vision.

Nature 2004:428;279.

62. Bowmaker JK. Evolution of vertebrate visual pigments. Vis Res. 2008:48;2022-2041.

63. Schweikert LE, Grace MS. Altered environmental light drives retinal change in the Atlantic

Tarpon (*Megalops atlanticus*) over timescales relevant to marine environmental disturbance.

BMC Ecol. 2018:18;1, https://doi.org/10.1186/s12898-018-0157-0.

64. Schweikert LE, Grace MS. Spectral sensitivity change may precede habitat shift in the

developing retina of the Atlantic tarpon (*Megalops atlanticus*) Physiol Biochem Zool.

2017:90;553-563.

65. Johnson AM, Stanis S, Fuller RC. Diurnal lighting patterns and habitat alter opsin expression

and colour preferences in a killifish. Proc R Soc B 2013:280;20130796.

http://dx.doi.org/10.1098/rspb.2013.0796.

66. Shand J, Davies WL, Thomas N, Balmer L, Cowing JA, Pointer M. The influence of

ontogeny and light environment on the expression of visual pigment opsins in the retina of

the black bream, *Acanthopagrus butcheri*. J Exp Biol. 2008:211;1495-1503.

67. Taylor SM, Loew ER, Grace MR.. Ontogenic retinal changes in three ecologically distinct

elopmorph fishes (Elopomorpha: Teleostei) correlate with light environment and behavior.

Vis Neurosci. 2015:32;1-15.

68. Munz FW. Vision: visual pigments. In: Hoar WS, Randall DJ, editors. Fish Physiology vol 5.

New York: Academic Press; 1971: p 1-32.

69. Munz FW, McFarland WN. The significance of spectral position in the rhodopsins of tropical

marine fishes. Vision Res. 1973:13;1829-1874.

70. Crescitelli F, McFall-Ngai M, Horwitz, J. The visual pigment sensitivity hypothesis: further

evidence from fishes of varying habitats J Comp Physiol A, 1985:157;323-333.

71. Lythgoe JN Vision in fishes: ecological adaptations. In: Ali MA, editor. Environmental

physiology of fishes, NATO Advanced Study Institutes Series (Series A: Life Sciences) vol

35. Boston: Springer; 1980: p 431-445.

72. Lythgoe JN (1988) Light and vision in the aquatic environment. In: Atema J, Fay RR, Popper

AN, Tavolga WN, editors. Sensory biology of aquatic animals. New York: Springer; p 57-82.

73. McFarland W Light in the sea: the optical world of elasmobranchs. J Exp Zool. 1990:256;3-

12.

584
585