The model of local axon homeostasis - explaining the role and regulation of microtubule bundles in axon maintenance and pathology

Ines Hahn, André Voelzmann, Yu-Ting Liew, Beatriz Costa-Gomes, Andreas Prokop[#]

The University of Manchester, Manchester Academic Health Science Centre, Faculty of Biology, Medicine and Health, School of Biology, Manchester, UK

<u>Running title</u>**:** The role of microtubules in axon pathology

Key words: *Drosophila*, neurodegeneration, axons, actin, cytoskeleton, microtubules

[#]   author for correspondence:
       The University of Manchester
       Faculty of Life Sciences
       Oxford Road
       Manchester M13 9PT
       Tel: +44-(0)161-27-51556
       Fax: +44-(0)161-27-51505
       Andreas.Prokop@manchester.ac.uk

Abstract

Axons are the slender, cable-like, up to meter-long projections of neurons that electrically wire our brain and body. In spite of their challenging morphology, they usually need to be maintained for an organism's lifetime. This makes them key lesion sites in pathological processes of ageing, injury and neurodegeneration. To better understand how axons are formed and maintained long-term, we focus here on the parallel bundles of microtubules (MTs) which form their indispensable structural backbones and highways for life-sustaining cargo transport and organelle dynamics. Many MT-binding and -regulating proteins in axons have prominent hereditary links to axon degeneration, but knowing their molecular roles is usually insufficient to explain their roles during axon morphogenesis, maintenance or pathology. Such understanding requires deciphering how these proteins interact in regulatory networks to implement observed cellular phenomena. Here we propose the model of local axon homeostasis as a conceptual framework that attempts to combine current knowledge into one coherent interactome. According to this model, each area of an axon is self-sustaining through local auto-regulatory networks; these networks maintain a balance between (1) the enormous mechanical challenges posed by the life-sustaining intra-axonal motor dynamics and (2) the maintenance activities required to sustain MT bundles as the highways needed for those dynamics. This model offers a new level of explanation, and we hope that it will influence experimental approaches leading to more MT-related experimental data across the spectrum of axonopathies - which are urgently required to decipher the important contributions of MTs to axon biology and pathology.

Introduction

Axons are the slender, cable-like extensions of nerve cells which form the nerves and nerve tracts that wire our brain and body, sending neuronal messages in highly regulated manners. With diameters of only 0.1-15µm (Hoffman, 1995), they extend over distances of up to a meter in humans. To adopt such a unique morphology and physiology, axons display many specialised features (Fig.1).

Axons are indispensable for nervous system function, as illustrated by paralysis in spinal cord injury caused by the interruption of ascending and descending axon tracts (Bichenback, 2013; Tedeschi and Bradke, 2016). Axons are key lesion sites in injury-induced trauma and coma (Gaetz, 2004; Medana and Esiri, 2003; Smith et al., 2000; Tang-Schomer et al., 2012), and axon decay is believed to be the trigger for neuronal loss in ageing and many neurodegenerative disorders (Adalbert and Coleman, 2012; Salvadores et al., 2017). Notably, most neurons cannot be replaced, and compensation of lost axons through collateral branching of intact neighbouring axons has obvious limitations (Adalbert and Coleman, 2012; Sturrock, 1987).

This means that most axons have to be maintained for an organism's life time, i.e. up to a century in humans; unsurprisingly, mammals tend to lose almost half their axon mass towards high age (Calkins, 2013; Marner et al., 2003). This trend is severely enhanced in neurodegenerative disorders, as illustrated by gradually increasing paralysis in spastic paraplegia or motorneuron disease (Blackstone et al., 2011; Riancho et al., 2019).

Research into neurodegenerative disorders typically approaches the problem by describing observed phenotypes and unravelling the molecular mechanisms performed by proteins linked to the disease. However, it seems that this approach rarely leads to satisfactory explanations of the pathology (Aguzzi, 2019). We believe that more profound understanding will arise when widening the scope from molecular to cellular mechanisms, by studying how proteins work within regulatory networks to underpin observable processes of axon biology - thus performing investigations at the level of complexity at which pathology becomes manifest. Here we will illustrate this approach by focussing on the axonal cytoskeleton.

The importance of microtubule bundles for axon biology

As illustrated in Fig. 1, the cytoskeleton of the axon shaft consists of straight parallel bundles of MTs, which are interspersed with intermediate filaments (not shown) and longitudinal actin fibres called 'actin trails' - all running through evenly spaced periodic rings of cortical actin (Qu et al., 2017; Xu et al., 2013); significant deviations from this organisation, that will not be addressed in this review, exist at axon initial segments (not shown), growth cones and synapses (Dent et al., 2011; Leterrier, 2018; Leterrier et al., 2017; Prokop, 2013).

Of the three cytoskeleton classes, intermediate filaments were suggested by anatomical, developmental and genetic studies to regulate axon diameter, and their axonal aggregation is a hallmark of many neurodegenerative diseases (Friede and Samorajski, 1970; Hoffman, 1995; Perrot et al., 2008; Rao et al., 2003; Sakaguchi et al., 1993). However, intermediate filament accumulations are not necessarily the cause, but can be the consequence of axon decay (Eyer et al., 1998; Nguyen et al., 2000; Perrot et al., 2008). Notably, *Neurofilament-H-lacZ* mutant mice or *Quiver* mutant quail completely lacking axonal intermediate filaments, develop and

breed fairly normally (Eyer and Peterson, 1994; Yamasaki et al., 1991), and various arthropods form axons of defined diameters in the absence of any axonal intermediate filaments (Allen et al., 2006; Hirokawa, 1986; Voelzmann et al., 2016a). These examples suggest that intermediate filaments play no key roles in axon growth and maintenance. In contrast, the actin and microtubule (MT) cytoskeleton are essential for all stages of neuronal development and maintenance (Sakakibara et al., 2013; Voelzmann et al., 2016a) and this review will focus on the role and regulation of MTs.

MTs in axons are arranged into bundles which run all along axon shafts and are essential for their biology in at least three ways. Firstly, they serve as structural backbones, comparable to a vertebral column of a snake; since MTs in these bundles are discontinuous and expected to be interlinked via flexible connections (see section on cross-linkers), they are ideally suited to respond to longitudinal stretch and compression (similar to a half-extended telescope ladder), but also to torsion and flexure (Fig.2).

Secondly, MT bundles provide the highways for life-sustaining axonal transport between cell bodies and the axonal compartment. This transport is driven anterogradely by kinesins and retrogradely by the dynein/Dynactin complex; the cargoes include mRNAs, cytoplasmic proteins including signalling factors, vesicles delivering synaptic proteins, adhesion factors, neuropeptides and/or membrane lipids, as well as entire organelles including mitochondria (Fig. 3A-D; Goldstein et al., 2008; Gondre-Lewis et al., 2012; Gonzalez and Couve, 2014; Hirokawa et al., 2010; Pfenninger, 2009). Furthermore, local dynamics of organelles, such as fission or fusion of mitochondria, can be expected to require forces generated by MT-associated motor proteins (Fig. 3E; Saxton and Hollenbeck, 2012).

Third, axonal bundles provide a source for readily available MTs that can be used for other purposes (curved arrows in Fig.1); for example, splaying MTs can trigger axon extension processes in growth cones (Dent et al., 2011; Prokop et al., 2013), induce collateral branch formation along the axon shaft (Kalil and Dent, 2014), or support physiological changes at synapses (Bodaleo and Gonzalez-Billault, 2016).

Maintaining MT bundles is therefore crucial for axon longevity. Accordingly, there are prominent and numerous genetic links from MT regulators to hereditary neurodegenerative disorders (Suppl. Mat. in Prokop et al., 2013), and axon decay is a frequent side effect of MT-targeting chemotherapies (Prior et al., 2017; Wozniak et al., 2018; Wu et al., 2014). Of particular interest for this review are reports of pathological axon swellings where MT bundles have disintegrated into loops or waves (bottom of Fig.3), occurring in ageing, after injury and in certain axonopathies (Adalbert et al., 2009; Bernier and Kothary, 1998; Dalpe et al., 1998; Denton et al., 2014; Fassier et al., 2013; Havlicek et al., 2014; Sorbara et al., 2014; Tang-Schomer et al., 2012; Tarrade et al., 2006; Yamasaki et al., 1991; Yin et al., 2016). Notably, one study suggests that MT aberration upon ageing could cause swellings that trap and damage mitochondria, thus triggering axon degeneration (Fiala et al., 2007). However, in the existing literature too little emphasis is given to MTs and there are simply not enough data to deduce meaningful correlations between axon degeneration and MT bundle decay.

Even if there were a close correlation, this still does not exclude that, depending on the pathological condition, MT bundle deterioration may be a mere consequence rather than cause of axon decay (details in Fig.4). Ultimate clarification will only arise from developing a better understanding of MT bundle-forming and -maintaining machinery. Here we propose a conceptual framework that may facilitate such developments.

The integrated model of local axon homeostasis

The foundations for this conceptual framework were laid when we took the decision to use the
fruit fly *Drosophila melanogaster* as a means to study how cytoskeletal regulators collaborate in
orchestrating the morphogenetic changes that drive axon growth (Sánchez-Soriano et al.,
2007). *Drosophila* is not a miniature human, but it has many advantages and provides powerful
means to uncover the regulatory concepts behind the roles and regulation of axonal MTs, which
then often apply to higher organisms (Box 1; Aguzzi, 2019; Bellen et al., 2010; Elden et al.,
2010; Prokop, 2018). Through using *Drosophila* neurons as a consistent standardised cell
system, our group alone performed functional analyses of over 50 actin- and/or MT-binding or -
associating regulators (Prokop et al., 2013); these studies form an unprecedented pool of data
on the basis of which to develop novel concepts (Alves-Silva et al., 2012; Beaven et al., 2015;
Gonçalves-Pimentel et al., 2011; Qu et al., 2018; Qu et al., 2017; Voelzmann et al., 2016b).

Box1 Why using *Drosophila*?

The use of *Drosophila* neurons to study the neuronal cytoskeleton has a number of advantages that were detailed elsewhere (Prokop et al., 2013). Key aspects are the high degree of conservation of cytoskeletal proteins, regulators and dynamics, the experimental amenability of neurons in primary cell culture and *in vivo* (Prokop et al., 2013; Prokop et al., 2012; Sánchez-Soriano et al., 2010), and the relative ease of genetic manipulation based on available resources and efficient combinatorial genetics (Hahn et al., 2016). The power of combinatorial genetics is rooted in the relative ease, speed and cost effectiveness with which genes can be manipulated and functionally analysed, facilitating also combined analyses of multiple factors in the same animals/cells (Prokop, 2018; Prokop et al., 2013; Roote and Prokop, 2013). Combinatorial genetics has been extremely successful in overcoming problems of redundancy, and generating new conceptual understanding of co-operative networks of MT regulation (see main text). This can hardly be achieved through isolated work on individual factors.

Our loss-of-function analyses of 24 MT-binding or -associating (2nd order) proteins, revealed that
more than half displayed significant MT disorganisation. Interestingly, the MT disorganisation
found in these various conditions appears to display certain common characteristics: axons
display areas in which their bundles are dissolved into chaotic, intertwined, crisscrossing
arrangements of curled MTs (see examples in Fig.5). Notably, when using the same genetic
conditions *in vivo*, comparable phenotypes were observed in the fly brain (Qu et al., 2018).
Such *in vivo* phenotypes in the fly remind of the curled MT conformations in pathological axon
swellings of mammalian models mentioned in the previous section. Potential evolutionary
conservation of this phenomenon is also supported by the occurrence of MT curling and
disorganisation in mouse and rat primary neurons (Ahmad et al., 2006; Sánchez-Soriano et al.,
2009).

As an attempt to explain this surprising phenotype across mutant conditions and animal groups,
we developed the model of 'local axon homeostasis' (Prokop, 2016; Voelzmann et al., 2016a).
The model states that MTs, which usually behave like rigid rods (Hawkins et al., 2010), are
challenged to buckle ('d' in Fig.3) and/or curl up by the force-enriched axonal environment ('A-E'
in Fig.3; detailed in the next two sections). Through this bias towards MT curling, the parallel
axon bundles are at risk of becoming disorganised and turning into pathological swellings. The

model therefore proposes that this destructive tendency of axonal MTs is contained through the action of different classes of MT-associating and -regulating proteins, which co-operate and complement each other to form robust machinery that 'tames' MTs into bundles ('1-16' in Fig.3).

In this model, each axon segment uses local mechanisms to maintain its bundled MT organisation (hence 'local axon homeostasis'). Hereditary or acquired loss of single regulators would therefore be expected to weaken this machinery and increase the statistical risk of MT disorganisation. Such heightened probability might explain why many axonopathies affect primarily long axons (Prior et al., 2017), and why certain disorders linked to MT regulators display late onset axon decay (Voelzmann et al., 2017).

In the next two sections, we describe potential mechanisms underlying deteriorating destructive MT behaviours in axons and their relationship to molecular motor proteins. We will then summarise experimentally demonstrated MT maintaining and 'taming' mechanisms, and speculate about further potential maintenance mechanisms based on existing knowledge of known classes of axonal MT-regulating proteins.

Understanding the unusual curling behaviour of MTs in axons

MTs are polar polymers composed of α/ß-tubulin heterodimers which are arranged in a head-to-tail fashion into linear protofilaments; mostly 13 of these protofilaments are laterally aligned forming a straight tube of roughly 25 nm diameter (Fig.6A, C). But MTs can deviate from this norm: for example, axonal MTs were reported to contain 13 protofilaments in frog olfactory or goldfish brain axons, 11 or 15 in *C. elegans*, and 12 in *Drosophila*, crayfish and lobster (Benshalom and Reese, 1985; Burton et al., 1975; Savage et al., 1989). Deviation from the straight 13 protofilament conformation equips MTs with distinct, functionally relevant physical properties (Chaaban and Brouhard, 2017; Chalfie and Thomson, 1982). Importantly, protofilaments in these deviating MTs are skewed and cause a supertwist of the tubule (Fig.6D) (Chrétien and Fuller, 2000; Chrétien et al., 1996; Chrétien and Wade, 1991); this supertwist forces motor proteins to rotate around MTs (Ray et al., 1993) and is the likely explanation for supercoil of entire axons observed upon MT bundle destabilisation (Krieg et al., 2017; Shaw and Bray, 1977).

MTs are structurally active, and their properties can change upon protein binding (see interactions with kinesins below) or when altering the 'tubulin code'; the tubulin code is determined by the incorporation of different existing isotypes of α- and ß-tubulin into the MT lattice, and the addition of a range of distinct post-translational modifications (Fig.6B; Janke and Kneussel, 2010; Park and Roll-Mecak, 2018; Ti et al., 2018; Vemu et al., 2017). Some modifications influence the interaction with MT-binding proteins (e.g. spastin; Valenstein and Roll-Mecak, 2016), others are believed to structurally protect MTs from damage or depolymerisation, such as poly-aminations on various residues (Song et al., 2013) or acetylation of luminal lysine 40 (Fig.6B; Baas et al., 2016; Howes et al., 2014; Soppina et al., 2012; Xu et al., 2017). Notably, mutation of lysine 40 in *Drosophila* α1-tubulin caused mild but physiologically relevant *in vivo* phenotypes (Jenkins et al., 2017; Yan et al., 2018). Furthermore, the MT lumen may contain MIPs (MT inner proteins) that likely also contribute to MT stability (Ichikawa and Bui, 2018).

Although curvature is a key driver of MT plus end dynamics during de-/polymerisation (Brouhard and Rice, 2018; van Haren and Wittmann, 2019), MT lattices *in vitro* usually behave as rigid rods with a persistence length of 1-10 mm (as compared to ~12 µm measured for actin

filaments; Fletcher and Mullins, 2010; Hawkins et al., 2010; Howard, 2001). However, in so-
called gliding assays where MTs are moved around (minus-end-leading) on a carpet of active
kinesins, they can undergo fishtailing or form micron-sized arcs or loops (Amos and Amos,
1991; Lam et al., 2016; Weiss et al., 1991). Interestingly, this phenomenon seems not to occur
upon plus-end-leading movement on (axonemal) dynein carpets (although collisions are far
more frequent; Sumino et al., 2012).

On kinesin carpets, single MT loops form through motor-driven buckling upon substrate pinning
or collision, favoured by high MT and/or kinesin densities or exposure to non-polar conditions
(n-heptane, air; Tab.1); loops can be astonishingly stable (frequently >5 mins, as reported in Liu
et al., 2011). Their structural longevity is significantly enhanced when MTs are reversibly cross-
linked with biotin-streptavidin so that MTs bundle up (analogous to Fig.2C), thus forming spools
composed of up to dozens of MTs; spools appear to reproduce most behaviours observed for
single MTs (Tab.1): (1) the smallest inner diameters for loops and spools usually lie in the range
of 1-3 µm (with diameters of curvature below ~1 µm believed to break MTs; Odde et al., 1999;
Waterman-Storer and Salmon, 1997); (2) the direction of loop rotation is a function of the left-
versus right-handed supertwist of MTs (Fig.6D), and spools rotate according to the supertwist of
their constituent MTs (Kawamura et al., 2008; Liu et al., 2008); (3) spool diameters increase
with the degree of rigidity of its MTs (Wada et al., 2015). Interestingly, like off-track MTs in
axons, MTs can escape the bundled conformation of spools which can sometimes trigger spool
disassembly (Hess et al., 2005; Liu et al., 2008; VanDelinder et al., 2016b).

Notably, key parameters promoting MT loops in gliding assays can also be found in the narrow
axonal tube: it is force-enriched due to the high density of MT-associated motor proteins, and
growing/gliding MTs have a large probability to collide or get obstructed by the crowded
organelle or protein content of axons (Fig.1). In line with this argument, the observed loop
diameters *in vitro* roughly approximate observed diameters of disorganised MTs in axons
(Tab.1, Fig.5), and even spool shapes can resemble MT conformations observed in growth
cones of fly or mammalian neurons (Dent and Kalil, 2001; Hess et al., 2005; Sánchez-Soriano
et al., 2010). Therefore curled MT conformations in axons might find explanations from *in vitro*
work, and a number of mathematical models were put forward to describe loop or spool
dynamics in gliding assays (Crenshaw et al., 2011; Gosselin et al., 2016; Luria et al., 2011;
Pearce et al., 2018; Ziebert et al., 2015). Of these, two models provide mechanistic ideas for
how bending through external forces can result in semi-stable loops.

The first model builds on known conformational changes of tubulin. Thus, tubulins in non-
hydrolysed GMPCPP-MTs are 1-3% longer than hydrolysed GDP-tubulin, and taxol added after
(but not during) polymerisation achieves a similar elongation (Fig.6F; Alushin et al., 2014; Amos
and Löwe, 1999; Arnal and Wade, 1995; Castle et al., 2017; Hyman et al., 1995). Notably, this
conformational change seems physiologically relevant, as its suppression by the T238A
mutation in yeast ß-tubulin stabilises MTs *in vivo* and causes mitotic defects (Geyer et al., 2015;
Machin et al., 1995). Since virtually all gliding assays use taxol-treated (hence extended) MTs
(Tab.1), the first model proposes that kinesin-mediated MT bending switches tubulins located
on the concave side of the tube into the shorter conformation. This conformation can be
maintained as an energetically favoured state, which could be further assisted by MT-binding
proteins (Fig.6F; Ziebert et al., 2015).

The second model is based on work showing that kinesin-1 has a preference for convex MT
surfaces and can stabilise MT curvature by extending their lattice to similar degrees as taxol
(translating into a diameter of curvature of 3.2 µm; Peet et al., 2018). The conformational

changes imposed by kinesins include compaction of tubulin that goes beyond taxol- or
GMPCPP-induced effects (Krebs et al., 2004; Morikawa et al., 2015), and kinesins were found
to bind cooperatively to MTs potentially causing a snowball effect (Cross, 2019; Muto et al.,
2005). The model proposes therefore that binding of kinesin-1 to the convex side can drive a
bias towards curvature (Pearce et al., 2018). This model is particularly attractive for the non-
extended GDP-MTs in axons, and it could apply to further axonal lattice binding proteins such
as tau and doublecortin, which also seem to bind differently to straight and curved MTs
(Balabanian et al., 2017; Bechstedt et al., 2014; Ettinger et al., 2016; Samsonov et al., 2004).

Naturally, these models are in their infancy and will have to be refined by gradually
incorporating further reported findings. For example, MTs behave as elastic cylinders
(comparable to a garden hose) and can undergo softening through cross-sectional flattening
when strongly bent (Fig.6E; Kononova et al., 2014; Memet et al., 2018). In this same vein,
conformational changes of MTs upon kinesin-1 binding were reported to soften MTs locally
rather than increase their rigidity (Kabir et al., 2014). If confirmed, this would have important
implications for any existing models; together with the kinesin-induced tubulin compaction
(yellow asterisks in Fig.6F), it might be a mechanism to absorb energy and reduce the shear
force load on MTs.

In conclusion, formations of axonal areas of disorganised curled MTs could be seen as
processes of 'active self-organisation', for which insights from gliding assays provide attractive
explanations (Lam et al., 2016): extrapolation from *in vitro* work suggests that MT curling could
be caused through an interdependent relationship between the force-enriched environment and
the responses of MTs as a function of their intrinsic properties. It is now important to challenge
this view and perform thorough analyses across genetic conditions and animal models, by
obtaining neutral parametric descriptions of MT disorganisations observed in the different
conditions (Fig.5), and of the dynamics through which they are initiated and maintained. If such
studies revealed comparable parameters across different conditions, this would support the idea
of a common concept behind the unusual MT behaviours in axons - as proposed by the model
of local axon homeostasis. In the next section we will summarise roles of axonal MT-associated
motors during axon pathology and explore whether they are key drivers of MT disorganisations.

<u>The intricate relationship between MTs and their associated motor proteins</u>

MT-associated motors comprise the minus end-directed dynein/Dynactin complex and the
mostly plus-end directed proteins of the kinesin family (Hirokawa et al., 2010). Several kinesins
display direct roles in MT regulation (Sturgill and Ohi, 2013). These include active MT
depolymerisation (kinesin-8, -13) (Walczak et al., 2013) as well as MT polymerisation (kinesin-2,
-5; Chen and Hancock, 2015; Gumy et al., 2013; Guzik-Lendrum et al., 2017), MT-cross-linkage
(kinesin-5, -6, -12; see section on bundling), and roles in promoting MT orientation as a feature
of neuronal polarity (Tas et al., 2017; Zheng et al., 2008).

However, most attention is given to the active cargo and organelle transport and dynamics in
axons (Fig.3A-E; see section on axonal cytoskeleton), which is driven retrogradely by
dynein/Dynactin (Allan, 2011) and anterogradely by kinesins (primarily kinesin-1, -2, and -3)
(Hirokawa et al., 2010). The forces imposed by these dynamics and/or the size of cargoes
moved poses an obvious challenge to MT bundles (Appert-Rolland et al., 2015) and might be
the main correlate of the bending forces generated by kinesin carpets in gliding assays (see
previous section).

Clearly, there is an intricate mutual regulatory relationship and finely tuned balance between the
amount of transport, and the structural properties of the transport highways (Appert-Rolland et
al., 2015; Prokop, 2013). For example, MT density is higher in small calibre axons than in large
axons (~15 *versus* ~150 MTs/μm^2), and mathematical modelling suggests that this is required to
achieve the same transport efficiency as in large axons (Wortman et al., 2014 ; and references
within). The tubulin isotype composition of MTs or their posttranslational modifications, as well
as physical presence of other MT-binding proteins influence motor protein dynamics ('a' in Fig.3;
Balabanian et al., 2017; Monroy et al., 2018; Sirajuddin et al., 2014; Subramaniyan Parimalam
et al., 2016). *Vice versa*, it has been reported that motor proteins cause damage to the MTs
they walk on (Dumont et al., 2015; Peet et al., 2018; Triclin et al., 2018; VanDelinder et al.,
2016a), which likely triggers maintenance responses including MT repair (Akhmanova, 2018) or
potentially even replacement ('14' in Fig.3).

Tipping the balance in this mutual relationship can easily be imagined to cause reciprocal
deficiencies in transport rate and MT bundle organisation. For example, disorganisation or
partial breakage of MTs has been reported to cause pathological transport deficits (Fiala et al.,
2007; Tang-Schomer et al., 2012). *Vice versa*, immunological lesioning experiments initially
caused transport defects, which were then followed by MT disorganisation (Sorbara et al.,
2014). Analogously, we observe severe MT disorganisation in *Drosophila* primary neurons upon
loss of kinesin-1 or -3 (kinesin-1 shown in Fig.5E).

How loss of these kinesins can cause MT disorganisation can currently only be hypothesised.
For example, it has been reported for dendrites that kinesin-1 migrates on acetylated and
kinesin-3 on tyrosinated MTs (Tas et al., 2017). Provided the same is true in axons, the loss of
kinesin-1 would relieve acetylated MTs, but tyrosinated MTs would still bear their full transport
load - and *vice versa*. Such imbalances in transport distribution across MTs could lead to shear
forces that buckle MTs and seed MT disorganisation. In the same vein, MT disorganisation was
reported to be triggered by directional changes in motor traffic upon deficiency of the dynein
regulator NDEL1 at the axon initial segment (Kuijpers et al., 2016). Furthermore, the movement
of large cargoes likely induces dynamic rearrangements of local MT-MT crosslinking networks
(see section on cross-linkage); in this scenario, violating the balanced proportion between
cross-linkers and transport load may become a path to bundle aberration.

Alternatively, transport defects might affect MTs through biochemical routes, simply caused by
the fact that the bundle-maintaining machinery runs out of supply. There would be expected
distribution gaps (a) of tubulin heterodimers as building blocks, (b) of the proteins required to
execute MT bundle maintenance work ('b' in Fig.3), and (c) of organelles. Organelle deficiencies
can trigger systemic changes that would likely affect MT maintenance, as detailed in Box 2.

Functional interdependencies of organelles with MTs may explain why different types of
Charcot-Marie-Tooth disease or hereditary spastic paraplegias can be caused through motor
proteins as well as regulators of membranous compartments (Blackstone, 2018; Bucci et al.,
2012). In agreement with this line of argumentation, MT stabilising drugs have been beneficial in
animal models of neurodegeneration, including SPG4 (Box 2) and Alzheimer's disease
(Brunden et al., 2014). *Vice versa*, axonal swellings induced by senile plaques in the *Tg-*
swAPP^{Prp} mouse model of Alzheimer's disease were strongly enhanced when removing one
copy of the KLC1 gene (a linker required for kinesin-1 mediated transport), and this effect was
found to be conserved in *Drosophila* (Stokin et al., 2005).

Box 2. The intricate relationship between MTs and axonal organelles

Mitochondria are the main source for ATP (Sheng, 2017), required to fuel multiple processes relevant for MT regulation (red arrows in Fig.3); these include actin assembly and dynamics (Krendel and Mooseker, 2005; Skruber et al., 2018), protein phosphorylation (Bogoyevitch and Fairlie, 2007), GTP production required for MT polymerisation and signalling (Berg et al., 2002; Hall and Lalli, 2010; Voelzmann et al., 2016a), MT severing (McNally and Roll-Mecak, 2018), and MT-motor dynamics (Hirokawa et al., 2010)(; although vesicular transport uses local glycolysis to generate its own ATP; yellow star in Fig.3A;)(Hinckelmann et al., 2016; Zala et al., 2013). Secondly, the mitochondrial surface is an important signalling platform and could be used to orchestrate MT regulation locally (not shown in Fig.3;)(McBride et al., 2006). Thirdly, mitochondria cooperate with ER in the regulation of intracellular free calcium (yellow cloud in Fig.3;)(Rieusset, 2017; Wu et al., 2017) which has direct impact on MT regulators (e.g. spectraplakins, tau, kinesins;)(Kapur et al., 2012; McVicker et al., 2015) or even on MTs themselves (O'Brien et al., 1997). Fourthly, mitochondria collaborate with peroxisomes in the regulation of reactive oxygen species ('ROS' in Fig.3;)(Fransen et al., 2017; Pascual-Ahuir et al., 2017), which have known effects on MT regulation (Wilson and Gonzalez-Billault, 2015).

Furthermore, aberrations of axonal transport or MT-bundle organisation can cause mitochondrial damage or dysregulation of the mitochondria-peroxisome system, both leading to oxidative stress as a major path to axon pathology (Fiala et al., 2007; Liu et al., 2017; Pascual-Ahuir et al., 2017). Such causative relationships between MTs and oxidative stress can be experimentally demonstrated: for example the MT-stabilising drug epothilone B rescues oxidative stress caused by peroxisome transport deficiencies in a human iPSC model of SPG4 (spastin-linked spastic paraplegia 4;)(Wali et al., 2016).

Similar interdependencies would apply to other important organelles or membrane compartments that likewise depend on MT-binding motor proteins to undergo meaningful dynamics (Fig.3D); of particular importance are the ER with its multiple roles in calcium homeostasis, protein synthesis and lipidogenesis (Gonzalez and Couve, 2014), or the endolysosomal system required for proteostasis (Winckler et al., 2018). For example, drug-induced inhibition of the proteasome-ubiquitination system has been shown to induce alteration in MTs and axonal transport (Staff et al., 2013).

Naturally, the argumentative framework presented here is highly speculative, given the enormous complexity of the relationships between MT bundle organisation, motor protein activity and systemic factors. But we hope that these reflections will motivate experimenters to have a closer look at MTs in future studies of axon biology and pathology. More data are urgently needed, which does often not require more than analysing neuronal morphology with antisera against MTs (rather than restricting to intermediate filaments), or increasing the magnification in ongoing ultrastructural studies to have a closer look at MTs. In the following sections we will explore the mechanisms that are potentially used to prevent motor-induced MT bundle aberrations.

<u>MT polymerisation as a fundamental requirement for bundle maintenance</u>

As mentioned before, the numbers of axonal MTs have to be well adapted to the transport load (Wortman et al., 2014), and this requires a well-regulated machinery of MT polymerisation and

disassembly (blue stippled arrows in Fig.3). On the one hand, MT volume has to be generated
de novo during axon growth ('8' in Fig.3), and thereafter this volume has to be maintained at
steady-state and prevented from MT senescence, which requires MT repair but likely also MT
turn-over ('14' in Fig.1; Akhmanova, 2018; Triclin et al., 2018; Voelzmann et al., 2016a).

As we detailed in a previous review (Voelzmann et al., 2016a), the machinery of MT de-
/polymerisation requires three sub-machineries: (1) dynamic protein complexes at the MT plus
end (blue balls, 'Eb1' in Fig.3); (2) the supply of α/β-tubulin heterodimers as building blocks
which occurs through a complex regulatory network in close co-regulation with MT dynamics ('c'
in Fig.3; Gasic and Mitchison, 2018; Preitner et al., 2014); (3) proteins which bind or post-
translationally modify the MT lattice, for example through stabilising MTs against
depolymerisation ('7' in 'Fig.3).

Such complex machinery has to be orchestrated in tune with the wider systemic context of
axons. This is illustrated by our recent work in *Drosophila* neurons, showing that loss of cortical
actin rings in the axon shaft (Fig.1) causes a reduction in MT polymerisation speed, eventually
affecting MT bundle integrity; simultaneous application of MT-destabilising drugs or removal of
the MT-stabilising spectraplakin Short stop (Shot) exacerbated these effects, frequently even
eliminating entire axons (Qu et al., 2017). Similar co-dependencies are suggested by other
reports: (1) parallel loss of spectrin and tau causes axonal MT loss in *C. elegans* (Krieg et al.,
2017); (2) axon-shortening induced by the MT-stabiliser taxol can be ameliorated through co-
application of actin-destabilising drugs (in both chick and *Drosophila* neurons; Letourneau et al.,
1987; Sánchez-Soriano et al., 2010); (3) application of actin-destabilising drugs to PC12 cells
changes the tubulin to microtubule ratio (Dennerll et al., 1988). Co-dependencies of MT
polymerisation on MT stabilisation and actin networks are currently best explained by
biomechanical models as detailed in Box 3.

<u>Cortical guidance and elimination of polymerising MTs</u>

Whilst well-equilibrated MT polymerisation is a requirement for axonal maintenance, it also
poses a risk, in that extending MTs can accidentally project out of the bundle and seed MT
disorganisation ('4' in Fig.3), potentially similar to loop formation upon MT obstruction in gliding
assays (see section on curling). A key factor preventing this from happening is Eb1 (Alves-Silva
et al., 2012; Figs.3 and 5B). Eb1 directly binds at extending MT plus ends where it promotes
polymerisation (Zanic et al., 2013) and serves as a scaffold for many other proteins (Gupta et
al., 2014).

One mechanism through which Eb1 maintains extending MTs in bundled configuration, is MT
guidance mediated by Short stop (Shot). Shot is a well-conserved spectraplakin, able to cross-
link cortical actin, MTs and Eb1 ('5' in Fig.3), thus guiding polymerising MTs in parallel to the
axonal surface and laying them out into parallel bundles (Alves-Silva et al., 2012). Similar to
Eb1 deficiency, also loss of Shot causes severe MT disorganisation in axons - and the same is
true for its two mammalian homologues ACF7 and dystonin (Bernier and Kothary, 1998; Dalpe
et al., 1998; Sánchez-Soriano et al., 2009; Voelzmann et al., 2017) - of which the latter links to
the axonopathy HSAN6 (type 6 hereditary sensory and autonomic neuropathy; Edvardson et al.,
2012).

Such cortical guidance is complemented by at least one control mechanism: if MTs
(accidentally) leave their bundled arrangements and extend towards the cortex, they get
eliminated by Efa6 ('4' in Fig.3), a cortical collapse factor that associates with the axonal

membrane via its C-terminal plekstrin homology domain; consistent with the model of local axon homeostasis, loss of Efa6 causes three neuronal phenotypes, in culture as well as in fly brains: significant MT disorganisation (Fig.5D), longer axons, and more axonal branches (Qu et al., 2018); all three phenotypes are based on morphogenetic processes to which 'off-track' MTs can contribute (curved arrows in Fig.1). Our model would predict that mutant phenotypes caused by loss of Shot and Efa6 should enhance each other because they are caused through complementary mechanisms of MT bundle regulation. Accordingly, we found a clear increase in MT disorganisation when removing both Shot and Efa6 from the same neurons (Qu et al., 2018). We propose therefore that, Shot and Eb1 keep MTs away from the membrane, and Efa6 acts as a finely tuned quality control factor eliminating accidental off-track MTs, whilst still permitting enough MTs to get through to perform intended functions in axon growth and branching.

Box 3. Biomechanical models of axon growth

The regulation of axonal growth dynamics has been explained with the concept of tensegrity (tensional integrity), an architectural principle based on structural nets that are under continuous tension whilst containing isolated components under compression (Buckminster Fuller, 1961; Ingber and Folkman, 1989). Thus, in axons "*actin is under tension supported in part by microtubules under compression*" (Heidemann and Buxbaum, 1990). Tension is provided by pulling acto-myosin networks in growth cones (Fass and Odde, 2003; Heidemann et al., 1990), the rigid but contractile properties of cortical actin in the axon shaft (Fan et al., 2017; Heidemann and Buxbaum, 1990; Krieg et al., 2017; Xu et al., 2013) (; Fig.1), and the stiff nature of cross-linked MT bundles is well suited to oppose compressive forces up to a certain threshold (Fig.2).

In such a balanced system, destabilisation of MTs increases tension at the expense of compression, and a decrease in acto-myosin networks or contractility reduces tension in favour of compression; force-generating MT polymerisation is one component responding to and regulating this balance with ultimate impact on axon length (Buxbaum and Heidemann, 1988; Dennerll et al., 1988; Heidemann et al., 1990; Letourneau et al., 1987).

In further agreement with this hypothesis, pulling the tips of axons enhances their growth rate (Bray, 1984; Lamoureux et al., 2010; Zheng et al., 1991); even single MTs polymerise faster when being pulled on *in vitro* (Brouhard and Rice, 2018). The necessary translation of the mechanical stimulus into changes of MT polymerisation rates might therefore occur at the level of MT polymerases (Brouhard and Rice, 2018) and/or be mediated by mechano-sensitive calcium channels in the axonal membrane (Franze et al., 2009; He et al., 2019; Heidemann and Buxbaum, 1990).

Interestingly, the cortical collapse function of Efa6 is not conserved in vertebrates (Qu et al., 2018). But the concepts derived from Efa6 studies still appears relevant, because loss of the unrelated neuronal cortical collapse factor KIF21A causes analogous phenotypes in mammalian neurons: KIF21A mutations linked to the neurodevelopmental disorder CFEOM1 (type 1 congenital fibrosis of the extraocular muscles) affect axon growth and axonal branching just like Efa6 (Qu et al., 2018; van der Vaart et al., 2013), and we would also predict potential increases in MT disorganisation (no data available).

Guidance along cortical actin seems not the only mechanism through which Eb1 and Shot keep
MTs on track. This is illustrated by the simple fact that MT disorganisation observed upon loss
of Shot or Eb1 does not occur when removing actin from axon shafts (Alves-Silva et al., 2012;
Qu et al., 2017; Sánchez-Soriano et al., 2010). This suggests that both factors perform
additional, actin-independent functions or interactions to promote MT bundles.

For example, the unusual Shot-PH isoform, which is highly enriched in the nervous system and
harbours a plakin repeat region (PRR; conserved in mammalian spectraplakins), is a likely
candidate for such roles, but still has to be investigated ('11' in Fig.3) (Hahn et al., 2016;
Voelzmann et al., 2017). Eb1 has a long list of protein interactors besides Shot, and some of
them might attract extending MT plus ends to the axonal core ('9' in Fig.3); for example, they
could be proteins, such as APC-like or GAS2-LIKE family members (Pickled eggs/Pigs in
Drosophila), known to bind both MTs and Eb1 in *Drosophila* and mammals (Beaven et al., 2015;
Pines et al., 2010; Stroud et al., 2014).

<u>Role of severing proteins and MT-destabilising kinesins</u>

As indicated in the previous section, MT disorganisation, axon growth and collateral axon
branch formation are driven by MTs leaving the bundled conformation (curved arrows in Fig.1;
Kalil and Dent, 2014; Tint et al., 2009; Tymanskyj et al., 2017). Consequently, MT-
disassembling factors should have the principal potential to negatively regulate either process.
This is true for cortical MT collapse factors (*Drosophila* Efa6, mammalian Kif21A; see previous
section) which can down-regulate axonal growth, branching and MT disorganisation. Also the
MT-depolymerising kinesin-13 family member Kif2A (Homma et al., 2003) and MT severing
proteins (spastin, katanin and fidgetin) were reported to inhibit neurite growth and/or branching
(Leo et al., 2015; Mao et al., 2014; Tao et al., 2016). Even more, MT disorganisation is
observed upon the losses of *Drosophila* katanin (our unpublished results) or mammalian spastin
(Denton et al., 2014; Fassier et al., 2013; Havlicek et al., 2014; Tarrade et al., 2006).

However, other studies of spastin, katanin and fidgetin led to contradictory findings, describing
them as promoters rather than inhibitors of neurite growth and branching (Ahmad et al., 1999;
Butler et al., 2010; Havlicek et al., 2014; Karabay et al., 2004; Riano et al., 2009; Stewart et al.,
2012; Stone et al., 2012; Wood et al., 2006; Yu et al., 2008). Such stark, potentially context-
dependent deviations seem to reflect the complex regulation of severing proteins. Thus, spastin,
katanin and fidgetin are all members of the superfamily of AAA proteins (ATPases associated
with diverse cellular activities) (McNally and Roll-Mecak, 2018; Sharp and Ross, 2012; Zhang et
al., 2007), but their severing activity is differentially regulated through their individual responses
to (a) posttranslational MT modifications (in particular acetylation and poly-glutamylation; Bailey
et al., 2015; Lacroix et al., 2010; Leo et al., 2015; Shin et al., 2019; Sudo and Baas, 2010;
Valenstein and Roll-Mecak, 2016), (b) antagonistic MT shaft-binding proteins such as tau
(Qiang et al., 2018; Qiang et al., 2006; Yu et al., 2008), or (c) recruitment through specifically
localised proteins such as CAMSAP (Jiang et al., 2018).

Through this precise spatiotemporal regulation of their activity, severing proteins are believed to
either eliminate MTs or to break them up into stable fragments that serve as seeds for MT
amplification (Baas et al., 2016; McNally and Roll-Mecak, 2018; Vemu et al., 2018). These
properties of severing proteins could be used in different ways to prevent MT disorganisation:
First, by targeting disorganised MTs for elimination, MT severing proteins could serve as quality
control factors ('6' in Fig.3) that complement roles of cortical collapse factors ('4' in Fig.3). In

agreement with this idea, katanin in plant cells was reported to localise and sever preferentially at MT cross-points, which can be used to take out non-aligned MTs (McNally and Roll-Mecak, 2018).

Second, MT elimination functions could prevent MT bundle senescence. Thus, MTs suffer from damage through tear-and-wear (Dumont et al., 2015; Peet et al., 2018; Triclin et al., 2018; VanDelinder et al., 2016a), which might cause bundle aberration by abrogating interactions with MT-binding proteins (red cross at '16' in Fig.3). On the one hand, this is addressed by immediate lattice repair involving katanin or spastin (Davis et al., 2002; Diaz-Valencia et al., 2011; Gasic and Mitchison, 2018; Triclin et al., 2018; Vemu et al., 2018). On the other hand, spastin and katanin could prevent senescence through selective elimination of aged MTs (as similarly suggested for kinesin-8 or -13; Gardner et al., 2011), followed by compensatory polymerisation ('14' in Fig.3). Such a mechanism might explain why spastin deficiency in the *Sp$^\Delta$* mouse model causes a drop in MT polymerisation whilst triggering MT disorganisation (Fassier et al., 2013).

Third, MT shortening functions of katanin are required at MT minus ends. Thus, in both mammals and *Drosophila*, the minus-end capper CAMSAP/Patronin protects against MT disassembly, and recruits katanin to counterbalance against uncontrolled minus-end extension ('13' in Fig.3; Goodwin and Vale, 2010; Jiang et al., 2018; Nashchekin et al., 2016); such uncontrolled extension may cause MTs to go off-track or to buckle through extra forces produced.

Fourth, in the absence of MT multiplication through severing proteins, MT numbers might gradually decline, thus causing transport interruptions with additional knock-on effects on MT bundle organisation and eventually systemic pathology (see section on motor proteins; Wali et al., 2018; Wali et al., 2016). Such a scenario could provide an alternative interpretation also for the above mentioned *Sp$^\Delta$* mouse model, in that insufficient MT amplification might lie at the root of the problem rather than prevention of senescence; curiously, axon swellings in this model were reduced with low doses of MT-stabilising or -destabilising drugs (Fassier et al., 2013), not favouring either of the explanations.

Understanding spastin is important because it is by far the most prominent factor linking to spastic paraplegias worldwide (Koh et al., 2018; Schüle et al., 2016), and axonal swellings are a hallmark of the disease (Blackstone, 2018; Zempel and Mandelkow, 2015). Most SPG4-linked mutations lie within the AAA-ATPase domain (Shoukier et al., 2009), suggesting that MT severing is key to the disease pathology. However, point mutations might generate versions of spastin, which either act as dominant negative alleles (forming dysfunctional complexes that titrate out other spastin-interacting factors), or acquire gain-of-function qualities by diffusing away to perform other roles. One such MT-independent role of spastin is the isoform-specific regulation of the endoplasmic reticulum ('15' in Fig.3), including its shape, interaction with the endosome or production of lipid droplets (Allison et al., 2017; Papadopoulos et al., 2015; Park et al., 2010; Solowska and Baas, 2015). Therefore, it is difficult to exclude that at least part of those mutations causes systemic rather than MT aberrations as the key trigger of decay ('3' in Fig.4).

Structural support through MT-MT cross-linkage

MT-MT cross-linkage ('12' in Fig.3) is likely the oldest mechanistic concept put forward by neurobiologists to explain MT bundles (Lee and Brandt, 1992) and appears an obvious means

of suppressing MT disorganisation. In physical terms, axons have been described as a "*stiff spring in series with a viscoelastic (Voight) element composed of a less stiff spring in parallel with a fluid dashpot*" (Heidemann et al., 1990), meaning that axons are under rest tension and combine elastic and viscous properties. A central structural component underpinning such properties is likely provided by networks of MT-MT cross-linkers (Fig.2), where each linker is able to detach upon super-threshold pull or compression, and re-attach thereafter (slip-bonds). Physical cross-linking strands between axonal MTs were observed decades ago (Hirokawa, 1982; Hirokawa, 1986), and mathematical models support MT-MT cross-linkage as an important structural feature of axons (e.g. de Rooij and Kuhl, 2018; Lazarus et al., 2015; Li et al., 2018; Peter and Mofrad, 2012).

However, the underlying genetic factors remain surprisingly controversial to this day - to a degree that one cannot even fully exclude a model view where linkers do not attach but rather separate MTs, whilst the corset of contractile cortical actin rings (Fig.1) constrains them into a dense bundle (Fan et al., 2017). This uncertainty stems from the low number of publications reporting structural bundle defects upon loss of putative MT-MT cross-linkers (see below). Further doubt comes from the recognition that MT bundling observed upon neuronal linker expression in non-neuronal cells, might represent artefacts, because MT bundling can even be achieved through expression of isolated MT-binding domains, or the application of the MT-stabilising drug taxol which causes bundles with ultrastructural cross-bridges that are indistinguishable from those induced by tau or MAP2 (Chapin et al., 1991; DeBonis et al., 2015; Goriounov et al., 2003; Kader et al., 2017; Lee and Brandt, 1992). Another example is dynamin, which is linked to Charcot-Marie-Tooth disease, has been shown to bundle MTs *in vitro*, but *in vivo* seems to bind membranes instead (Scaife and Margolis, 1990; Shpetner and Vallee, 1989; Züchner et al., 2005).

For example, MTLC1 and MAP1B (Futsch in *Drosophila*) appear ideal cross-linkers, because they both possess MT-binding domains at their N- and C-termini, and induce bundles upon expression in non-neuronal cells (although MAP1B appears a weak bundler) (Kader et al., 2017; Penazzi et al., 2016; Satake et al., 2017). However, there are only isolated reports that axonal bundle defects occur in their absence (Bettencourt da Cruz et al., 2005; Satake et al., 2017). Instead, the long history of MAP1B research is mostly dedicated to aspects of axon development (Villarroel-Campos and Gonzalez-Billault, 2014). Its fly homologue Futsch promotes MT spools at synaptic terminals (Roos et al., 2000), but its axonal defects comprise merely growth aberrations but no MT disorganisation (Hummel et al., 2000; our unpublished results). Therefore, existing data for MAP1B/Futsch only vaguely support roles in axonal MT bundling.

The other conserved linker candidate tau, has only one central MT-binding region and achieves physical MT-MT linkage *in vitro* through N-terminal dimerisation (Chung et al., 2016; Méphon-Gaspard et al., 2016; Rosenberg et al., 2008). However, its dwell time on MTs seems very short (Janning et al., 2014; Samsonov et al., 2004), and reported tau-deficient phenotypes hardly ever comprise bundle aberration but rather developmental defects of neurons (Krieg et al., 2017; Penazzi et al., 2016).

Pinpointing MT-MT cross-linkage through tau is further complicated by the fact that normal tau has been associated with a whole array of further molecular functions relevant for MT dynamics. For example, tau can protect MTs from severing by katanin (Qiang et al., 2006), bind tubulin hetero-dimers (Shin et al., 2018), switch between bundled and single MT states (Prezel et al., 2018), cross-link MTs with actin or the cortex (Biswas and Kalil, 2018; Cabrales Fontela et al.,

2017; Maas et al., 2000), stabilise MTs during axon initiation (Brandt, 1998), maintain labile domains along MT shafts (Qiang et al., 2018), regulate End-binding proteins (Sayas et al., 2015), compete with kinesins (Trinczek et al., 1999), and promote MT nucleation and polymerisation (Penazzi et al., 2016). A similarly broad pleiotropy has been reported for MAP1B (Villarroel-Campos and Gonzalez-Billault, 2014).

A further complicating factor for pinpointing relevant MT-MT cross-linking activities are the obvious functional redundancies in the system. Thus, *MAP1B* and *tau* have enhanced growth phenotypes when both mutant conditions are combined in double-mutant mouse neurons (Takei et al., 2000). Complementary to this, co-expression of Futsch and Tau causes enhanced phenotypes in the *Drosophila* CNS (Hummel et al., 2000). Such functional redundancy likely extends to further potential cross-linkers. For example, Kinesin-5 (KIF11), kinesin-6 (KIF23, Pavarotti in *Drosophila*) and kinesin-12 (KIF15) are best known for their ability to slide anti-parallel MTs in the mitotic spindle (Baas, 1999); however, in axons where MTs are arranged in parallel these kinesins seem to inhibit sliding (Dong et al., 2019; Lin et al., 2012; Liu et al., 2010; Lu et al., 2013; Myers and Baas, 2007; Nadar et al., 2012) - and this can be considered a form of MT-MT cross-linkage. Notably, we observe that loss of *Drosophila* Pavarotti causes axonal MT disorganisation (our unpublished data), providing a readout for studying its potential linker functions. By incorporating redundancies redundancy of linker candidates into experimental approaches, there might be new opportunities to decipher the true molecular nature of MT-MT cross-linkage in axons.

Are MT bundles anchored to the axonal cortex?

Apart from cross-linkage between MTs, another bundle-stabilising aspect could be cross-linkage or anchorage to the wall of the axonal tube. For example, polymerisation occurs all along axon shafts, and it has been proposed from studies in developing vertebrate and *Drosophila* neurons, that the MT mass generated in the axonal shaft during axon growth gradually shifts anterogradely (Miller and Sheetz, 2006; O'Toole et al., 2008; Reinsch et al., 1991; Roossien et al., 2013). Contributing forces were suggested to be generated by pulling forces in the rear of growth cones (O'Toole et al., 2015), by MT polymerisation, by thermal motion of MT-MT cross-linkers (Lansky et al., 2015), by kinesins actively sliding MTs along other MTs ('B' in Fig.3; Lu and Gelfand, 2017) or by dyneins sliding MTs along cortical F-actin ('10' in Fig.3; Ahmad et al., 2006; He et al., 2005; Myers et al., 2006; Roossien et al., 2014).

Potential MT sliding along cortical actin would involve tethering MT bundles to the axolemma and raises the fundamental question as to whether MT bundles are free-flowing within the axonal tube, or anchored to it. Potential anchorage is suggested by observed co-drift of the axolemma with the axon core (Lamoureux et al., 2010; Popov et al., 1993; Zheng et al., 1991). Such anchorage would not have to be static, but might involve an interface of slip-bonds, as similarly suggested for back-flowing actin networks at stable focal adhesion sites (Case and Waterman, 2015). Furthermore, anchorage of MTs would not have to be restricted to the rigid cortical actin networks (Fig.1; '2' in Fig.3; Xu, 2013 #6895}, but could also involve links to adhesion factors ('3' in Fig.3), thus forming mechano-sensing modules (Yap et al., 2018) responding to local shear forces generated between MT bundles and the axonal environment. Such adhesion-mediated local mechano-sensing could explain local regulation phenomena: for example, net rates of fast mitochondrial transport are regulated in a way that they gradually decrease towards distal axon segments, indirectly proportional to the local MT drift rate along the axon (which increased towards the distal end; Miller and Sheetz, 2006). This proximo-distal

differences of fast mitochondrial transport could therefore be determined by mechano-sensing as a function of the local MT drift rate relative to the outer axonal environment.

Spectraplakins are candidate anchoring factors, as suggested by *Drosophila* neurons lacking the spectraplakin Shot and treated with the MT-stabilising drug taxol: in these neurons, axonal MTs shifted distally, often leaving tubulin-free zones in the proximal axon shaft (Voelzmann et al., 2017). However, this same effect was not observed when applying taxol whilst also removing actin (Sánchez-Soriano et al., 2010), suggesting that Shot does more than simply anchoring to cortical actin in this experimental context.

This conclusion is in agreement with the breadth of axonal functions proposed for spectraplakins, making them the most conspicuous regulators of the axonal cytoskeleton in any species investigated so far (Voelzmann et al., 2017). For example, the role of Shot in preventing MT shift may involve a combination of molecular functions including MT-MT cross-linkage ('11' in Fig.3), actin-MT cross-linkage ('2' and '5' in Fig.3), as well as two other potential anchoring mechanisms: Firstly, Shot is able to compartmentalise the localisation of the cell adhesion molecule Fasciclin 2 (Bottenberg et al., 2009; Prokop et al., 1998), suggesting that it might be able to link to membrane-associated proteins like its mammalian homologue dystonin ('3' in Fig.3; Voelzmann et al., 2017). Secondly, work on non-neuronal cells of fly and mammals revealed that spectraplakins can interact with the MT minus end-stabilising factor CAMSAP/Patronin (a factor known to be relevant for neuronal morphology; Yau et al., 2014), thus anchoring MT minus ends to the cell cortex ('1' in Fig.3; Nashchekin et al., 2016; Ning et al., 2016; Noordstra et al., 2016).

Therefore, tethering of axonal MT bundles to the cortex might occur through dynein and/or spectraplakins. Further MT-binding and/or -cross-linking proteins were similarly reported to also bind to actin or the cortex, and these include tau, MAP1B, APC and dynamin (Biswas and Kalil, 2018; Blanchoin and Michelot, 2012; Brandt et al., 1995; Elie et al., 2015; Gu et al., 2010; Maas et al., 2000; Mohan and John, 2015; Villarroel-Campos and Gonzalez-Billault, 2014). They could cross-link to actin at the cortex ('2' in Fig.3) or to longitudinal actin trails located within MT bundles (Fig.1; Leterrier et al., 2017).

Taken together, there are clear indications for an intricate cross-linking network that can hold MT bundles together whilst tethering them to the axonal surface. Holding cross-linked MT bundles in space would constitute a powerful means to prevent MT buckling and bundle deformation, caused by the enormous forces imposed by axonal cargo transport.

Conclusions and future perspectives

Here we have presented a conceptual view by describing a functional interactome that integrates the enormous complexity of cross-regulatory networks acting at the local level in axons. We propose that there has to be a fine balance between damaging effects inflicted by life-sustaining 'associated' motor movements ('A-E' in Fig.3) and the activities that maintain the highways required for this movement (MT-'taming' mechanisms; '1-15' in Fig.3), fine-tuned through a number of cross-regulatory mechanisms ('a-e' in Fig.3).

Our model view includes a broad range of reports from the literature and speculative conclusions drawn. But its original foundations are derived from our own work in *Drosophila* neurons as one consistent model. For this, *Drosophila* (like other genetic invertebrate models) provides a cost-effective and fast system to unravel the functional overlap and interface of

different genetic factors - ideal to dissect complex machinery and deliver data that then often apply to axons of higher animals (Beaven et al., 2015; Prokop, 2018; Prokop et al., 2013).

This strategy offers a path towards solving the daunting task of disentangling the enormous regulatory complexity; the model of local axon homeostasis could provide the basis on which to formulate helpful working hypotheses. A good starting point might be to break down the local axon homeostasis machinery into classifiable sub-machineries, such as those discussed in the different sections of this review.

In our view, we urgently require a shift away from prioritising molecular mechanisms and should rather dedicate to exploring their integration into wider regulatory networks - a type of research that is too often disqualified as 'incremental science' (Cohen, 2017). In our opinion, this would be a much needed paradigm shift with a realistic chance of providing us with far better understanding of axonal cell biology at the organisational level at which axonopathies become manifest.

Such an integrative approach has to take into consideration that knowledge derived from non-neuronal cells might not apply in neurons (Beaven et al., 2015). Furthermore, the interactome shown in Fig.3 makes clear that we will need quantitative and computational modelling approaches: we know increasingly well how factors bind to MTs and partly understand how they might compete with each other. But how crowded can a single MT be, how many molecules are there at any time in its surrounding, and how much dynamic exchange is taking place?

Integrated understanding of axon biology will also improve our knowledge of the next higher level of complexity, i.e. the mechanisms that orchestrate axon homeostasis and can help to maintain balance even during phases of change: when switching from growth to differentiation, or during stress, injury, degeneration or regeneration. Obviously, signalling networks or systemic factors such as second messengers or the 'tubulin code' will be key players to this end (Baas et al., 2016; Park and Roll-Mecak, 2018; Schelski and Bradke, 2017; Wilson and Gonzalez-Billault, 2015), and glial cells will likely act as important external influencers of such processes (Pan and Chan, 2017).

Finally, MTs have been recognised as promising therapeutic targets (Baas and Ahmad, 2013; Eira et al., 2016; Zempel and Mandelkow, 2015), and urgently needed advance on this translational path will be facilitated by a better understanding of the axonal MT homeostasis system.

Acknowledgements

Work underpinning this article was made possible through support by the BBSRC to A.P (BB/I002448/1, BB/P020151/1, BB/L000717/1, BB/M007553/1), by parents to Y.-T.L., by the Leverhulme Trust to I.H. (ECF-2017-247). The Manchester Bioimaging Facility microscopes used in this study were purchased with grants from the BBSRC, The Wellcome Trust and The University of Manchester Strategic Fund. The Fly Facility has been supported by funds from The University of Manchester and the Wellcome Trust (087742/Z/08/Z). We thank Anthony Brown for introducing A.P. to the arguments that support a role of intermediate filaments in axon diameter.

References

Adalbert, R., Coleman, M. P. (2012). Axon pathology in age-related neurodegenerative disorders. *Neuropathol Appl Neurobiol* 39, 90–108 -- *http://www.ncbi.nlm.nih.gov/pubmed/23046254*

Adalbert, R., Nogradi, A., Babetto, E., Janeckova, L., Walker, S. A., Kerschensteiner, M., Misgeld, T., Coleman, M. P. (2009). Severely dystrophic axons at amyloid plaques remain continuous and connected to viable cell bodies. *Brain* 132, 402-16 -- *http://www.ncbi.nlm.nih.gov/pubmed/19059977*

Aguzzi, A. (2019). 'Forward genetics' and the causes of ALS. *Nature Reviews Molecular Cell Biology* 20, 67-67 -- *https://doi.org/10.1038/s41580-018-0062-6*

Ahmad, F. J., He, Y., Myers, K. A., Hasaka, T. P., Francis, F., Black, M. M., Baas, P. W. (2006). Effects of dynactin disruption and dynein depletion on axonal microtubules. *Traffic* 7, 524-37 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids= 16643276*

Ahmad, F. J., Yu, W., McNally, F. J., Baas, P. W. (1999). An essential role for katanin in severing microtubules in the neuron. *J Cell Biol* 145, 305-15 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids= 10209026*

Akhmanova, A. (2018). Strengthening Microtubules by Cuts that Heal. *Dev Cell* 47, 400-401 -- *http://www.sciencedirect.com/science/article/pii/S1534580718309201*

Al-Bassam, J. (2017). Revisiting the tubulin cofactors and Arl2 in the regulation of soluble αβ-tubulin pools and their effect on microtubule dynamics. *Molecular Biology of the Cell* 28, 359-363 -- *http://www.molbiolcell.org/content/28/3/359.abstract*

Allan, V. J. (2011). Cytoplasmic dynein. *Biochem Soc Trans* 39, 1169-78 -- *http://www.ncbi.nlm.nih.gov/pubmed/21936784*

Allen, M. J., Godenschwege, T. A., Tanouye, M. A., Phelan, P. (2006). Making an escape: development and function of the *Drosophila* giant fibre system. *Semin Cell Dev Biol* 17, 31-41 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids= 16378740*

Allison, R., Edgar, J. R., Pearson, G., Rizo, T., Newton, T., Gunther, S., Berner, F., Hague, J., Connell, J. W., Winkler, J., Lippincott-Schwartz, J., Beetz, C., Winner, B., Reid, E. (2017). Defects in ER-endosome contacts impact lysosome function in hereditary spastic paraplegia. *J Cell Biol* 216, 1337-1355 -- *http://www.ncbi.nlm.nih.gov/pubmed/28389476*

Alushin, G. M., Lander, G. C., Kellogg, E. H., Zhang, R., Baker, D., Nogales, E. (2014). High-resolution microtubule structures reveal the structural transitions in alphabeta-tubulin upon GTP hydrolysis. *Cell* 157, 1117-29 -- *http://www.ncbi.nlm.nih.gov/pubmed/24855948*

Alves-Silva, J., Sánchez-Soriano, N., Beaven, R., Klein, M., Parkin, J., Millard, T., Bellen, H., Venken, K. J. T., Ballestrem, C., Kammerer, R. A., Prokop, A. (2012). Spectraplakins promote microtubule-mediated axonal growth by functioning as structural microtubule-associated proteins and EB1-dependent +TIPs (Tip Interacting Proteins). *J. Neurosci* 32, 9143-58 -- *http://www.jneurosci.org/content/32/27/9143.full*

Amos, L., Amos, W. (1991). The bending of sliding microtubules imaged by confocal light microscopy and negative stain electron microscopy. *J Cell Sci* 1991, 95-101 -- *http://jcs.biologists.org/content/joces/1991/Supplement_14/95.full.pdf*

Amos, L. A., Löwe, J. (1999). How taxol stabilises microtubule structure. *Chem Biol* 6, R65-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/10074470*

Appert-Rolland, C., Ebbinghaus, M., Santen, L. (2015). Intracellular transport driven by cytoskeletal motors: General mechanisms and defects. *Physics Reports* 593, 1-59 -- *http://www.sciencedirect.com/science/article/pii/S037015731500335X*

Arnal, I., Wade, R. H. (1995). How does taxol stabilize microtubules? *Curr Biol* 5, 900-8 -- *http://www.ncbi.nlm.nih.gov/pubmed/7583148*

Baas, P. W. (1999). Microtubules and neuronal polarity: lessons from mitosis. *Neuron* 22, 23-31 --

Baas, P. W., Ahmad, F. J. (2013). Beyond taxol: microtubule-based treatment of disease and injury of the nervous system. *Brain* -- *http://www.ncbi.nlm.nih.gov/pubmed/23811322*

Baas, P. W., Rao, A. N., Matamoros, A. J., Leo, L. (2016). Stability properties of neuronal microtubules. *Cytoskeleton (Hoboken)* 73, 442-60 -- *http://www.ncbi.nlm.nih.gov/pubmed/26887570*

Bailey, M. E., Sackett, D. L., Ross, J. L. (2015). Katanin Severing and Binding Microtubules Are Inhibited by Tubulin Carboxy Tails. *Biophys J* 109, 2546-61 -- *https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4699919/?otool=igbumllib*

Balabanian, L., Berger, C. L., Hendricks, A. G. (2017). Acetylated microtubules are preferentially bundled leading to enhanced kinesin-1 motility. *Biophys J* 113, 1551-1560 -- *http://www.sciencedirect.com/science/article/pii/S0006349517308664*

Beaven, R., Dzhindzhev, N. S., Qu, Y., Hahn, I., Dajas-Bailador, F., Ohkura, H., Prokop, A. (2015). *Drosophila* CLIP-190 and mammalian CLIP-170 display reduced microtubule plus end association in the nervous system. *Mol Biol Cell* 26, 1491-1508 -- *http://www.molbiolcell.org/content/26/8/1491.abstract*

Bechstedt, S., Lu, K., Brouhard, Gary J. (2014). Doublecortin recognizes the longitudinal curvature of the microtubule end and lattice. *Curr Biol* 24, 2366-2375 -- *http://www.sciencedirect.com/science/article/pii/S0960982214010525*

Bellen, H. J., Tong, C., Tsuda, H. (2010). 100 years of *Drosophila* research and its impact on vertebrate neuroscience: a history lesson for the future. *Nat Rev Neurosci* 11, 514-522 -- *http://www.nature.com/nrn/journal/v11/n7/full/nrn2839.html*

Benshalom, G., Reese, T. S. (1985). Ultrastructural observations on the cytoarchitecture of axons processed by rapid-freezing and freeze-substitution. *J Neurocytol* 14, 943-60 -- *http://www.ncbi.nlm.nih.gov/pubmed/2420942*

Berg, J. M., Tymoczko, J. L., Stryer, L. (2002). "Biochemistry (5th edition)." W H Freeman, New York

Bernier, G., Kothary, R. (1998). Prenatal onset of axonopathy in Dystonia musculorum mice. *Dev Genet* 22, 160-8 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9581287*

Bettencourt da Cruz, A., Schwarzel, M., Schulze, S., Niyyati, M., Heisenberg, M., Kretzschmar, D. (2005). Disruption of the MAP1B-related protein FUTSCH leads to changes in the neuronal cytoskeleton, axonal transport defects, and progressive neurodegeneration in *Drosophila*. *Mol Biol Cell* 16, 2433-42 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15772149*

Bichenback, J. (2013). "International perspectives on spinal cord injury." WHO, ISCOS, Switzerland

Biswas, S., Kalil, K. (2018). The microtubule-associated protein tau mediates the organization of microtubules and their dynamic exploration of actin-rich lamellipodia and filopodia of cortical growth cones. *J Neurosci* 38, 291-307 -- *http://www.ncbi.nlm.nih.gov/pubmed/29167405*

Blackstone, C. (2018). Hereditary spastic paraplegia. *Handb Clin Neurol* 148, 633-652 -- *http://www.ncbi.nlm.nih.gov/pubmed/29478605*

Blackstone, C., O'Kane, C. J., Reid, E. (2011). Hereditary spastic paraplegias: membrane traffic and the motor pathway. *Nat Rev Neurosci* 12, 31-42 -- *http://www.ncbi.nlm.nih.gov/pubmed/21139634*

Blanchoin, L., Michelot, A. (2012). Actin Cytoskeleton: A Team Effort during Actin Assembly. *Curr Biol* 22, R643-5 -- *http://www.ncbi.nlm.nih.gov/pubmed/22917514*

Bodaleo, F. J., Gonzalez-Billault, C. (2016). The presynaptic microtubule cytoskeleton in physiological and pathological conditions: lessons from *Drosophila* Fragile X Syndrome and Hereditary Spastic Paraplegias. *Frontiers in Molecular Neuroscience* 9, 60-60 -- *https://www.ncbi.nlm.nih.gov/pubmed/27504085*

Bogoyevitch, M. A., Fairlie, D. P. (2007). A new paradigm for protein kinase inhibition: blocking phosphorylation without directly targeting ATP binding. *Drug Discov Today* 12, 622-33 -- *http://www.ncbi.nlm.nih.gov/pubmed/17706543*

Bottenberg, W., Sánchez-Soriano, N., Alves-Silva, J., Hahn, I., Mende, M., Prokop, A. (2009). Context-specific requirements of functional domains of the Spectraplakin Short stop *in vivo*. *Mech Dev* 126, 489-502 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19409984*

Bradke, F., Fawcett, J. W., Spira, M. E. (2012). Assembly of a new growth cone after axotomy: the precursor to axon regeneration. *Nat Rev Neurosci* 13, 183-93 -- *http://www.ncbi.nlm.nih.gov/pubmed/22334213*

Brandt, R. (1998). Cytoskeletal mechanisms of axon outgrowth and pathfinding. *Cell Tissue Res.* 292, 181-189 -- *https://link-springer-com.manchester.idm.oclc.org/article/10.1007%2Fs004410051049*

Brandt, R., Bakota, L. (2017). Microtubule dynamics and the neurodegenerative triad of Alzheimer's disease: The hidden connection. *J Neurochem* 173, 409-17 -- *http://dx.doi.org/10.1111/jnc.14011*

Brandt, R., Leger, J., Lee, G. (1995). Interaction of tau with the neural plasma membrane mediated by tau's amino-terminal projection domain. *J Cell Biol* 131, 1327-40 -- *http://www.ncbi.nlm.nih.gov/pubmed/8522593*

Brangwynne, C. P., MacKintosh, F. C., Kumar, S., Geisse, N. A., Talbot, J., Mahadevan, L., Parker, K. K., Ingber, D. E., Weitz, D. A. (2006). Microtubules can bear enhanced compressive loads in living cells because of lateral reinforcement. *J Cell Biol* 173, 733-41 -- *http://www.ncbi.nlm.nih.gov/pubmed/16754957*

Brangwynne, C. P., MacKintosh, F. C., Weitz, D. A. (2007). Force fluctuations and polymerization dynamics of intracellular microtubules. *Proc Natl Acad Sci U S A* 104, 16128-33 -- *http://www.ncbi.nlm.nih.gov/pubmed/17911265*

Bray, D. (1984). Axonal growth in response to experimentally applied mechanical tension. *Dev Biol* 102, 379-89. -- *https://doi.org/10.1016/0012-1606(84)90202-1*

Bridge, K. E., Berg, N., Adalbert, R., Babetto, E., Dias, T., Spillantini, M. G., Ribchester, R. R., Coleman, M. P. (2009). Late onset distal axonal swelling in YFP-H transgenic mice. *Neurobiol Aging* 30, 309-21 -- *http://www.ncbi.nlm.nih.gov/pubmed/17658198*

Brouhard, G. J., Rice, L. M. (2018). Microtubule dynamics: an interplay of biochemistry and mechanics. *Nat Rev Mol Cell Biol* -- *https://doi.org/10.1038/s41580-018-0009-y*

Brunden, K. R., Trojanowski, J. Q., Smith, A. B., 3rd, Lee, V. M., Ballatore, C. (2014). Microtubule-stabilizing agents as potential therapeutics for neurodegenerative disease. *Bioorg Med Chem* 22, 5040-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/24433963*

Bucci, C., Bakke, O., Progida, C. (2012). Charcot-Marie-Tooth disease and intracellular traffic. *Prog Neurobiol* 99, 191-225 -- *http://www.ncbi.nlm.nih.gov/pubmed/22465036*

Buckminster Fuller, R. (1961). Tensegrity. *Portfolio and Art News Annual* 4, 112-127, 144, 148 -- *http://www.rwgrayprojects.com/rbfnotes/fpapers/tensegrity/tenseg01.html*

Burton, P. R., Hinkley, R. E., Pierson, G. B. (1975). Tannic acid-stained microtubules with 12, 13, and 15 protofilaments. *J Cell Biol* 65, 227-33 -- *http://www.ncbi.nlm.nih.gov/pubmed/47861*

Butler, R., Wood, J. D., Landers, J. A., Cunliffe, V. T. (2010). Genetic and chemical modulation of spastin-dependent axon outgrowth in zebrafish embryos indicates a role for impaired microtubule dynamics in hereditary spastic paraplegia. *Dis Model Mech* 3, 743-51 -- *http://dmm.biologists.org/content/3/11-12/743.long*

Buxbaum, R. E., Heidemann, S. R. (1988). A thermodynamic model for force integration and microtubule assembly during axonal elongation. *J Theor Biol* 134, 379-90 -- *http://www.ncbi.nlm.nih.gov/pubmed/3254435*

Cabrales Fontela, Y., Kadavath, H., Biernat, J., Riedel, D., Mandelkow, E., Zweckstetter, M. (2017). Multivalent cross-linking of actin filaments and microtubules through the microtubule-associated protein Tau. *Nature Communications* 8, 1981 -- *https://doi.org/10.1038/s41467-017-02230-8*

Calkins, D. J. (2013). Age-Related Changes in the Visual Pathways: Blame It on the AxonAge-Related Changes in the Visual Pathways. *Investigative Ophthalmology & Visual Science* 54, ORSF37-ORSF41 -- *https://dx.doi.org/10.1167/iovs.13-12784*

Case, L. B., Waterman, C. M. (2015). Integration of actin dynamics and cell adhesion by a three-dimensional, mechanosensitive molecular clutch. *Nat Cell Biol* -- *http://www.ncbi.nlm.nih.gov/pubmed/26121555*

Castle, B. T., McCubbin, S., Prahl, L. S., Bernens, J. N., Sept, D., Odde, D. J. (2017). Mechanisms of kinetic stabilization by the drugs paclitaxel and vinblastine. *Molecular Biology of the Cell* 28, 1238-1257 -- *https://www.molbiolcell.org/doi/abs/10.1091/mbc.e16-08-0567*

Chaaban, S., Brouhard, G. J. (2017). A microtubule bestiary: structural diversity in tubulin polymers. *Mol Biol Cell* 28, 2924-2931 -- *http://www.molbiolcell.org/content/28/22/2924.abstract*

Chalfie, M., Thomson, J. N. (1982). Structural and functional diversity in the neuronal microtubules of *Caenorhabditis elegans*. *J Cell Biol* 93, 15-23 -- *http://www.ncbi.nlm.nih.gov/pubmed/7068753*

Chapin, S. J., Bulinski, J. C., Gundersen, G. G. (1991). Microtubule bundling in cells. *Nature* 349, 24 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=1670738*

Chen, Y., Hancock, W. O. (2015). Kinesin-5 is a microtubule polymerase. *Nat Commun* 6, 8160 -- *http://www.ncbi.nlm.nih.gov/pubmed/26437877*

Chrétien, D., Fuller, S. D. (2000). Microtubules switch occasionally into unfavorable configurations during elongation. *J Mol Biol* 298, 663-76 -- *http://www.ncbi.nlm.nih.gov/pubmed/10788328*

Chrétien, D., Kenney, J. M., Fuller, S. D., Wade, R. H. (1996). Determination of microtubule polarity by cryo-electron microscopy. *Structure* 4, 1031-40 -- *http://www.ncbi.nlm.nih.gov/pubmed/8805589*

Chrétien, D., Wade, R. H. (1991). New data on the microtubule surface lattice. *Biol Cell* 71, 161-74 -- *http://www.ncbi.nlm.nih.gov/pubmed/1912942*

Chung, P. J., Song, C., Deek, J., Miller, H. P., Li, Y., Choi, M. C., Wilson, L., Feinstein, S. C., Safinya, C. R. (2016). Tau mediates microtubule bundle architectures mimicking fascicles of microtubules found in the axon initial segment. *Nat Commun* 7, 12278 -- *http://dx.doi.org/10.1038/ncomms12278*

Cioni, J. M., Koppers, M., Holt, C. E. (2018). Molecular control of local translation in axon development and maintenance. *Curr Opin Neurobiol* 51, 86-94 -- *http://www.ncbi.nlm.nih.gov/pubmed/29549711*

Cohen, B. A. (2017). How should novelty be valued in science? *Elife* 6, e28699 -- *https://doi.org/10.7554/eLife.28699*

Court, F. A., Midha, R., Cisterna, B. A., Grochmal, J., Shakhbazau, A., Hendriks, W. T., Van Minnen, J. (2011). Morphological evidence for a transport of ribosomes from Schwann cells to regenerating axons. *Glia* 59, 1529-1539 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/glia.21196*

Crenshaw, J. D., Liang, T., Hess, H., Phillpot, S. R. (2011). A cellular automation approach to the simulation of active self-assembly of kinesin-powered molecular shuttles. *J Comp Theoret Nanosci* 8, 1999-2005 -- *https://doi-org.manchester.idm.oclc.org/10.1166/jctn.2011.1916*

Cross, R. A. (2019). Microtubule lattice plasticity. *Curr Opin Cell Biol* 56, 88-93 -- *http://www.sciencedirect.com/science/article/pii/S0955067418301418*

Dalpe, G., Leclerc, N., Vallee, A., Messer, A., Mathieu, M., De Repentigny, Y., Kothary, R. (1998). Dystonin is essential for maintaining neuronal cytoskeleton organization. *Mol Cell Neurosci* 10, 243-57 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9604204*

Davis, L. J., Odde, D. J., Block, S. M., Gross, S. P. (2002). The importance of lattice defects in katanin-mediated microtubule mevering *in vitro*. *Biophysical Journal* 82, 2916-2927 -- *https://doi.org/10.1016/S0006-3495(02)75632-4*

de Rooij, R., Kuhl, E. (2018). Microtubule polymerization and cross-link dynamics explain axonal stiffness and damage. *Biophys J* 114, 201-212 -- *https://www.sciencedirect.com/science/article/pii/S0006349517312390*

DeBonis, S., Neumann, E., Skoufias, D. A. (2015). Self protein-protein interactions are involved in TPPP/p25 mediated microtubule bundling. *Sci Rep* 5, 13242 -- *http://www.ncbi.nlm.nih.gov/pubmed/26289831*

Dennerll, T. J., Joshi, H. C., Steel, V. L., Buxbaum, R. E., Heidemann, S. R. (1988). Tension and compression in the cytoskeleton of PC-12 neurites. II: Quantitative measurements. *J Cell Biol* 107, 665-74 -- *http://www.ncbi.nlm.nih.gov/pubmed/3417767*

Dent, E. W., Gupton, S. L., Gertler, F. B. (2011). The growth cone cytoskeleton in axon outgrowth and guidance. *Cold Spring Harb Perspect Biol* 3, a001800 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=21106647*

Dent, E. W., Kalil, K. (2001). Axon branching requires interactions between dynamic microtubules and actin filaments. *J Neurosci* 21, 9757-69 -- *http://www.jneurosci.org.manchester.idm.oclc.org/content/21/24/9757.long*

Denton, K. R., Lei, L., Grenier, J., Rodionov, V., Blackstone, C., Li, X. J. (2014). Loss of spastin function results in disease-specific axonal defects in human pluripotent stem cell-based models of hereditary spastic paraplegia. *Stem Cells* 32, 414-23 -- *http://www.ncbi.nlm.nih.gov/pubmed/24123785*

Diaz-Valencia, J. D., Morelli, M. M., Bailey, M., Zhang, D., Sharp, D. J., Ross, J. L. (2011). *Drosophila* katanin-60 depolymerizes and severs at microtubule defects. *Biophys J* 100, 2440-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/21575578*

Dogterom, M., Koenderink, G. H. (2019). Actin–microtubule crosstalk in cell biology. *Nat Rev Mol Cell Biol* 20, 38-54 -- *https://doi.org/10.1038/s41580-018-0067-1*

Dong, Z., Wu, S., Zhu, C., Wang, X., Li, Y., Chen, X., Liu, D., Qiang, L., Baas, P. W., Liu, M. (2019). Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)/Cas9-mediated kif15 mutations accelerate axonal outgrowth during neuronal development and regeneration in zebrafish. *Traffic* 20, 71-81 -- *http://www.ncbi.nlm.nih.gov/pubmed/30411440*

Dumont, E., L. P., Do, C., Hess, H. (2015). Molecular wear of microtubules propelled by surface-adhered kinesins. *Nat Nano* 10, 166-169 -- *http://dx.doi.org/10.1038/nnano.2014.334*

Edvardson, S., Cinnamon, Y., Jalas, C., Shaag, A., Maayan, C., Axelrod, F. B., Elpeleg, O. (2012). Hereditary sensory autonomic neuropathy caused by a mutation in dystonin. *Ann Neurol* 71, 569-72 -- *http://www.ncbi.nlm.nih.gov/pubmed/22522446*

Eira, J., Silva, C. S., Sousa, M. M., Liz, M. A. (2016). The cytoskeleton as a novel therapeutic target for old neurodegenerative disorders. *Progress in Neurobiology* -- *http://www.sciencedirect.com/science/article/pii/S0301008215300800*

Elbaum, M., Kuchnir Fygenson, D., Libchaber, A. (1996). Buckling microtubules in vesicles. *Phys Rev Lett* 76, 4078-4081 -- *http://www.ncbi.nlm.nih.gov/pubmed/10061186*

Elden, A. C., Kim, H.-J., Hart, M. P., Chen-Plotkin, A. S., Johnson, B. S., Fang, X., Armakola, M., Geser, F., Greene, R., Lu, M. M., Padmanabhan, A., Clay-Falcone, D., McCluskey, L., Elman, L., Juhr, D., Gruber, P. J., Rüb, U., Auburger, G., Trojanowski, J. Q., Lee, V. M. Y., Van Deerlin, V. M., Bonini, N. M., Gitler, A. D. (2010). Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* 466, 1069 -- *https://doi.org/10.1038/nature09320*

Elie, A., Prezel, E., Guerin, C., Denarier, E., Ramirez-Rios, S., Serre, L., Andrieux, A., Fourest-Lieuvin, A., Blanchoin, L., Arnal, I. (2015). Tau co-organizes dynamic microtubule and actin networks. *Sci Rep* 5, 9964 -- *http://www.ncbi.nlm.nih.gov/pubmed/25944224*

Ettinger, A., van Haren, J., Ribeiro, S. A., Wittmann, T. (2016). Doublecortin is excluded from growing microtubule ends and recognizes the GDP-microtubule lattice. *Curr Biol* 26, 1549-1555 -- *http://www.ncbi.nlm.nih.gov/pubmed/27238282*

Eyer, J., Cleveland, D. W., Wong, P. C., Peterson, A. C. (1998). Pathogenesis of two axonopathies does not require axonal neurofilaments. *Nature* 391, 584-7 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9468135*

Eyer, J., Peterson, A. (1994). Neurofilament-deficient axons and perikaryal aggregates in viable transgenic mice expressing a neurofilament-beta-galactosidase fusion protein. *Neuron* 12, 389-405 -- *http://www.ncbi.nlm.nih.gov/pubmed/8110465*

Fan, A., Tofangchi, A., Kandel, M., Popescu, G., Saif, T. (2017). Coupled circumferential and axial tension driven by actin and myosin influences in vivo axon diameter. *Sci Rep* 7, 14188 -- *http://www.ncbi.nlm.nih.gov/pubmed/29079766*

Farah, C. A., Nguyen, M. D., Julien, J. P., Leclerc, N. (2003). Altered levels and distribution of microtubule-associated proteins before disease onset in a mouse model of amyotrophic lateral sclerosis. *J Neurochem* 84, 77-86 -- *http://www.ncbi.nlm.nih.gov/pubmed/12485403*

Fass, J. N., Odde, D. J. (2003). Tensile force-dependent neurite elicitation via anti-beta1 integrin antibody-coated magnetic beads. *Biophys J* 85, 623-36 -- *http://www.ncbi.nlm.nih.gov/pubmed/12829516*

Fassier, C., Tarrade, A., Peris, L., Courageot, S., Mailly, P., Dalard, C., Delga, S., Roblot, N., Lefevre, J., Job, D., Hazan, J., Curmi, P. A., Melki, J. (2013). Microtubule-targeting drugs rescue axonal swellings in cortical neurons from spastin knockout mice. *Dis Model Mech* 6, 72-83 -- *http://www.ncbi.nlm.nih.gov/pubmed/22773755*

Ferrier, A., Boyer, J. G., Kothary, R. (2013). Cellular and molecular biology of neuronal Dystonin. *Int Rev Cell Mol Biol* 300, 85-120 -- *http://www.ncbi.nlm.nih.gov/pubmed/23273860*

Fiala, J. C., Feinberg, M., Peters, A., Barbas, H. (2007). Mitochondrial degeneration in dystrophic neurites of senile plaques may lead to extracellular deposition of fine filaments. *Brain Struct Funct* 212, 195-207 --

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17717688

Fletcher, D. A., Mullins, R. D. (2010). Cell mechanics and the cytoskeleton. *Nature* 463, 485-92 -- *http://www.ncbi.nlm.nih.gov/pubmed/20110992*

Fransen, M., Lismont, C., Walton, P. (2017). The Peroxisome-Mitochondria Connection: How and Why? *Int J Mol Sci* 18 -- *http://www.ncbi.nlm.nih.gov/pubmed/28538669*

Franze, K., Gerdelmann, J., Weick, M., Betz, T., Pawlizak, S., Lakadamyali, M., Bayer, J., Rillich, K., Gogler, M., Lu, Y. B., Reichenbach, A., Janmey, P., Kas, J. (2009). Neurite branch retraction is caused by a threshold-dependent mechanical impact. *Biophys J* 97, 1883-90 -- *http://www.ncbi.nlm.nih.gov/pubmed/19804718*

Friede, R. L., Samorajski, T. (1970). Axon caliber related to neurofilaments and microtubules in sciatic nerve fibers of rats and mice. *Anat Rec* 167, 379-87 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=5454590*

Frühbeis, C., Fröhlich, D., Kuo, W. P., Amphornrat, J., Thilemann, S., Saab, A. S., Kirchhoff, F., Möbius, W., Goebbels, S., Nave, K. A., Schneider, A., Simons, M., Klugmann, M., Trotter, J., Krämer-Albers, E. M. (2013). Neurotransmitter-triggered transfer of exosomes mediates oligodendrocyte-neuron communication. *PLoS Biol* 11, e1001604 -- *http://www.ncbi.nlm.nih.gov/pubmed/23874151*

Gaetz, M. (2004). The neurophysiology of brain injury. *Clin Neurophysiol* 115, 4-18 -- *http://www.ncbi.nlm.nih.gov/pubmed/14706464*

Gardner, M. K., Zanic, M., Gell, C., Bormuth, V., Howard, J. (2011). Depolymerizing kinesins Kip3 and MCAK shape cellular microtubule architecture by differential control of catastrophe. *Cell* 147, 1092-103 -- *http://www.ncbi.nlm.nih.gov/pubmed/22118464*

Gasic, I., Mitchison, T. J. (2018). Autoregulation and repair in microtubule homeostasis. *Curr Opin Cell Biol* 56, 80-87 -- *http://www.ncbi.nlm.nih.gov/pubmed/30415186*

Geyer, E. A., Burns, A., Lalonde, B. A., Ye, X., Piedra, F.-A., Huffaker, T. C., Rice, L. M. (2015). A mutation uncouples the tubulin conformational and GTPase cycles, revealing allosteric control of microtubule dynamics. *Elife* 4, e10113 -- *https://doi.org/10.7554/eLife.10113*

Giuditta, A., Eyman, M., Kaplan, B. B. (2002a). Gene expression in the squid giant axon: neurotransmitter modulation of RNA transfer from periaxonal glia to the axon. *Biol Bull* 203, 189-90 -- *https://www.journals.uchicago.edu/doi/10.2307/1543389*

Giuditta, A., Kaplan, B. B., van Minnen, J., Alvarez, J., Koenig, E. (2002b). Axonal and presynaptic protein synthesis: new insights into the biology of the neuron. *Trends Neurosci* 25, 400-4 -- *http://www.ncbi.nlm.nih.gov/pubmed/12127756*

Goldstein, A. Y., Wang, X., Schwarz, T. L. (2008). Axonal transport and the delivery of pre-synaptic components. *Curr Opin Neurobiol* 18, 495-503 -- *http://www.ncbi.nlm.nih.gov/pubmed/18950710*

Gonçalves-Pimentel, C., Gombos, R., Mihály, J., Sánchez-Soriano, N., Prokop, A. (2011). Dissecting regulatory networks of filopodia formation in a *Drosophila* growth cone model. *PLoS ONE* 6, e18340 -- *http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0018340*

Gondre-Lewis, M. C., Park, J. J., Loh, Y. P. (2012). Cellular mechanisms for the biogenesis and transport of synaptic and dense-core vesicles. *Int Rev Cell Mol Biol* 299, 27-115 -- *http://www.ncbi.nlm.nih.gov/pubmed/22959301*

Gonzalez, C., Couve, A. (2014). The axonal endoplasmic reticulum and protein trafficking: Cellular bootlegging south of the soma. *Semin Cell Dev Biol* 27, 23-31 -- *http://www.ncbi.nlm.nih.gov/pubmed/24361785*

Goodwin, S. S., Vale, R. D. (2010). Patronin regulates the microtubule network by protecting microtubule minus ends. *Cell* 143, 263-74 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20946984*

Goriounov, D., Leung, C. L., Liem, R. K. (2003). Protein products of human Gas2-related genes on chromosomes 17 and 22 (hGAR17 and hGAR22) associate with both microfilaments and microtubules. *J Cell Sci* 116, 1045-58 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12584248*

Gosselin, P., Mohrbach, H., Kulić, I. M., Ziebert, F. (2016). On complex, curved trajectories in microtubule gliding. *Physica D: Nonlinear Phenomena* 318–319, 105-111 -- *http://www.sciencedirect.com/science/article/pii/S0167278915002183*

Gu, C., Yaddanapudi, S., Weins, A., Osborn, T., Reiser, J., Pollak, M., Hartwig, J., Sever, S. (2010). Direct dynamin–actin interactions regulate the actin cytoskeleton. *The EMBO Journal* 29, 3593-3606 -- *http://emboj.embopress.org/content/embojnl/29/21/3593.full.pdf*

Gumy, L. F., Chew, D. J., Tortosa, E., Katrukha, E. A., Kapitein, L. C., Tolkovsky, A. M., Hoogenraad, C. C., Fawcett, J. W. (2013). The kinesin-2 family member KIF3C regulates microtubule dynamics and is required for axon growth and regeneration. *J Neurosci* 33, 11329-11345 -- *http://www.jneurosci.org/content/jneuro/33/28/11329.full.pdf*

Gupta, K. K., Alberico, E. O., Nathke, I. S., Goodson, H. V. (2014). Promoting microtubule assembly: A hypothesis for the functional significance of the +TIP network. *Bioessays* 36, 818-26 -- *http://www.ncbi.nlm.nih.gov/pubmed/24943963*

Guzik-Lendrum, S., Rayment, I., Gilbert, S. P. (2017). Homodimeric kinesin-2 KIF3CC promotes microtubule dynamics. *Biophys J* 113, 1845-1857 -- *http://www.ncbi.nlm.nih.gov/pubmed/29045878*

Gyoeva, F. K. (2014). The role of motor proteins in signal propagation. *Biochemistry (Moscow)* 79, 849-855 -- *http://dx.doi.org/10.1134/S0006297914090028*

Hahn, I., Ronshaugen, M., Sánchez-Soriano, N., Prokop, A. (2016). Functional and genetic analysis of spectraplakins in *Drosophila*. *Methods Enzymol* 569, 373-405 -- *https://tinyurl.com/y4vhld56*

Hall, A., Lalli, G. (2010). Rho and Ras GTPases in axon growth, guidance, and branching. *Cold Spring Harb Perspect Biol* 2, a001818 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20182621*

Havlicek, S., Kohl, Z., Mishra, H. K., Prots, I., Eberhardt, E., Denguir, N., Wend, H., Plotz, S., Boyer, L., Marchetto, M. C., Aigner, S., Sticht, H., Groemer, T. W., Hehr, U., Lampert, A., Schlotzer-Schrehardt, U., Winkler, J., Gage, F. H., Winner, B. (2014). Gene dosage-dependent rescue of HSP neurite defects in SPG4 patients' neurons. *Hum Mol Genet* 23, 2527-41 -- *http://www.ncbi.nlm.nih.gov/pubmed/24381312*

Hawkins, T., Mirigian, M., Selcuk Yasar, M., Ross, J. L. (2010). Mechanics of microtubules. *J Biomech* 43, 23-30 -- *http://www.ncbi.nlm.nih.gov/pubmed/19815217*

He, L., Ahmad, M., Perrimon, N. (2019). Mechanosensitive channels and their functions in stem cell differentiation. *Exp Cell Res* 374, 259-265 -- *http://www.ncbi.nlm.nih.gov/pubmed/30500393*

He, Y., Francis, F., Myers, K. A., Yu, W., Black, M. M., Baas, P. W. (2005). Role of cytoplasmic dynein in the axonal transport of microtubules and neurofilaments. *J Cell Biol* 168, 697-703 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15728192*

Heidemann, S. R., Buxbaum, R. E. (1990). Tension as a regulator and integrator of axonal growth. *Cell Motil Cytoskeleton* 17, 6-10 -- *http://www.ncbi.nlm.nih.gov/pubmed/2225090*

Heidemann, S. R., Lamoureux, P., Buxbaum, R. E. (1990). Growth cone behavior and production of traction force. *J Cell Biol* 111, 1949-57 -- *http://jcb.rupress.org.manchester.idm.oclc.org/content/111/5/1949.long*

Hess, H., Clemmens, J., Brunner, C., Doot, R., Luna, S., Ernst, K. H., Vogel, V. (2005). Molecular self-assembly of "nanowires"and "nanospools" using active transport. *Nano Lett* 5, 629-33 -- *http://www.ncbi.nlm.nih.gov/pubmed/15826099*

Hinckelmann, M. V., Virlogeux, A., Niehage, C., Poujol, C., Choquet, D., Hoflack, B., Zala, D., Saudou, F. (2016). Self-propelling vesicles define glycolysis as the minimal energy machinery for neuronal transport. *Nat Commun* 7, 13233 -- *http://www.ncbi.nlm.nih.gov/pubmed/27775035*

Hirokawa, N. (1982). Cross-linker system between neurofilaments, microtubules, and membranous organelles in frog axons revealed by the quick-freeze, deep-etching method. *J Cell Biol* 94, 129-42 -- *http://www.ncbi.nlm.nih.gov/pubmed/6181077*

Hirokawa, N. (1986). 270K microtubule-associated protein cross-reacting with anti-MAP2 IgG in the crayfish peripheral nerve axon. *J Cell Biol* 103, 33-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/3722268*

Hirokawa, N., Niwa, S., Tanaka, Y. (2010). Molecular motors in neurons: transport mechanisms and roles in brain function, development, and disease. *Neuron* 68, 610-638 -- *http://www.ncbi.nlm.nih.gov/pubmed/21092854*

Hoffman, P. N. (1995). Review : The Synthesis, Axonal Transport, and Phosphorylation of Neurofilaments Determine Axonal Caliber in Myelinated Nerve Fibers. *The Neuroscientist* 1, 76-83 -- *https://journals.sagepub.com/doi/abs/10.1177/107385849500100204*

Homma, N., Takei, Y., Tanaka, Y., Nakata, T., Terada, S., Kikkawa, M., Noda, Y., Hirokawa, N. (2003). Kinesin superfamily protein 2A (KIF2A) functions in suppression of collateral branch extension. *Cell* 114, 229-39 -- *http://www.ncbi.nlm.nih.gov/pubmed/12887924*

Howard, J. (2001). "Mechanics of motorproteins and the cytoskeleton." Sinauer Assoc., Sunderland --

Howes, S. C., Alushin, G. M., Shida, T., Nachury, M. V., Nogales, E. (2014). Effects of tubulin acetylation and tubulin acetyltransferase binding on microtubule structure. *Mol Biol Cell* 25, 257-66 -- *http://www.ncbi.nlm.nih.gov/pubmed/24227885*

Hummel, T., Krukkert, K., Roos, J., Davis, G., Klämbt, C. (2000). *Drosophila* Futsch/22C10 is a MAP1B-like protein required for dendritic and axonal development. *Neuron* 26, 357-370 -- *https://www.cell.com/neuron/fulltext/S0896-6273(00)81169-1*

Hur, E. M., Saijilafu, Lee, B. D., Kim, S. J., Xu, W. L., Zhou, F. Q. (2011). GSK3 controls axon growth via CLASP-mediated regulation of growth cone microtubules. *Genes Dev* 25, 1968-81 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=21937714*

Hyman, A. A., Chretien, D., Arnal, I., Wade, R. H. (1995). Structural changes accompanying GTP hydrolysis in microtubules: information from a slowly hydrolyzable analogue guanylyl-(alpha,beta)-methylene-diphosphonate. *J Cell Biol* 128, 117-25 -- *http://www.ncbi.nlm.nih.gov/pubmed/7822409*

Ichikawa, M., Bui, K. H. (2018). Microtubule inner proteins: a meshwork of luminal proteins stabilizing the doublet microtubule. *BioEssays* 40, 1700209 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/bies.201700209*

Ingber, D., Folkman, J. (1989). Tension and compression as basic determinants of cell form and function: Utilization of a cellular tensegrity mechanism. *In* "Cell Shape: Determinants, Regulation and Regulatory Role" (W. Stein, F. Bronner, Eds.), pp. 3-31. Academic Press, San Diego

Janke, C., Kneussel, M. (2010). Tubulin post-translational modifications: encoding functions on the neuronal microtubule cytoskeleton. *Trends Neurosci* 33, 362-72 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20541813*

Janning, D., Igaev, M., Sundermann, F., Bruhmann, J., Beutel, O., Heinisch, J. J., Bakota, L., Piehler, J., Junge, W., Brandt, R. (2014). Single-molecule tracking of tau reveals fast kiss-and-hop interaction with microtubules in living neurons. *Mol Biol Cell* 25, 3541-51 -- *http://www.ncbi.nlm.nih.gov/pubmed/25165145*

Jenkins, B. V., Saunders, H. A. J., Record, H. L., Johnson-Schlitz, D. M., Wildonger, J. (2017). Effects of mutating alpha-tubulin lysine 40 on sensory dendrite development. *J Cell Sci* 130, 4120-4131 -- *http://www.ncbi.nlm.nih.gov/pubmed/29122984*

Jiang, K., Faltova, L., Hua, S., Capitani, G., Prota, A. E., Landgraf, C., Volkmer, R., Kammerer, R. A., Steinmetz, M. O., Akhmanova, A. (2018). Structural basis of formation of the microtubule minus-end-regulating CAMSAP-katanin complex. *Structure* 26, 375-382 e4 -- *http://www.ncbi.nlm.nih.gov/pubmed/29395789*

Kabir, A. M., Inoue, D., Hamano, Y., Mayama, H., Sada, K., Kakugo, A. (2014). Biomolecular motor modulates mechanical property of microtubule. *Biomacromolecules* 15, 1797-805 -- *http://www.ncbi.nlm.nih.gov/pubmed/24697688*

Kader, M. A., Satake, T., Yoshida, M., Hayashi, I., Suzuki, A. (2017). Molecular basis of the microtubule-regulating activity of microtubule crosslinking factor 1. *PLoS One* 12, e0182641 -- *https://doi.org/10.1371/journal.pone.0182641*

Kalil, K., Dent, E. W. (2014). Branch management: mechanisms of axon branching in the developing vertebrate CNS. *Nat Rev Neurosci* 15, 7-18 -- *http://www.ncbi.nlm.nih.gov/pubmed/24356070*

Kandel, E. R., Schwartz, J. H., Jessell, T. M., Siegelbaum, S., Hudspeth, A. J. (2012). "Principles of neural science (5th edition)." McGraw-Hill Publishing

Kapur, M., Wang, W., Maloney, M. T., Millan, I., Lundin, V. F., Tran, T. A., Yang, Y. (2012). Calcium tips the balance: a microtubule plus end to lattice binding switch operates in the carboxyl terminus of BPAG1n4. *EMBO Rep* 13, 1021-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/22995871*

Karabay, A., Yu, W., Solowska, J. M., Baird, D. H., Baas, P. W. (2004). Axonal growth is sensitive to the levels of katanin, a protein that severs microtubules. *J Neurosci* 24, 5778-88 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids= 15215300*

Kawamura, R., Kakugo, A., Shikinaka, K., Osada, Y., Gong, J. P. (2008). Ring-Shaped Assembly of Microtubules Shows Preferential Counterclockwise Motion. *Biomacromolecules* 9, 2277-2282 -- *http://dx.doi.org/10.1021/bm800639w*

Kawamura, R., Kakugo, A., Shikinaka, K., Osada, Y., Gong, J. P. (2011). Formation of motile assembly of microtubules driven by kinesins. *Smart Mater Struct* 20, 124007 -- *http://dx.doi.org/10.1088/0964-1726/20/12/124007*

Keating, T. J., Peloquin, J. G., Rodionov, V. I., Momcilovic, D., Borisy, G. G. (1997). Microtubule release from the centrosome. *Proc Natl Acad Sci U S A* 94, 5078-83 -- *http://www.ncbi.nlm.nih.gov/pubmed/9144193*

Kellogg, E. H., Hejab, N. M. A., Poepsel, S., Downing, K. H., DiMaio, F., Nogales, E. (2018). Near-atomic model of microtubule-tau interactions. *Science* 360, 1242-1246 -- *http://science.sciencemag.org/content/sci/360/6394/1242.full.pdf*

Koh, K., Ishiura, H., Tsuji, S., Takiyama, Y. (2018). JASPAC: Japan Spastic Paraplegia Research Consortium. *Brain Sci* 8 -- *http://www.ncbi.nlm.nih.gov/pubmed/30104498*

Kononova, O., Kholodov, Y., Theisen, K. E., Marx, K. A., Dima, R. I., Ataullakhanov, F. I., Grishchuk, E. L., Barsegov, V. (2014). Tubulin bond energies and microtubule biomechanics determined from nanoindentation in silico. *J Am Chem Soc* 136, 17036-17045 -- *https://doi.org/10.1021/ja506385p*

Koper, A., Schenck, A., Prokop, A. (2012). Analysis of adhesion molecules and basement membrane contributions to synaptic adhesion at the *Drosophila* embryonic NMJ. *PLoS One* 7, e36339 -- *http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0036339*

Krebs, A., Goldie, K. N., Hoenger, A. (2004). Complex formation with kinesin motor domains affects the structure of microtubules. *J Mol Biol* 335, 139-53 -- *http://www.ncbi.nlm.nih.gov/pubmed/14659746*

Krendel, M., Mooseker, M. S. (2005). Myosins: tails (and heads) of functional diversity. *Physiology (Bethesda)* 20, 239-51 -- *http://www.ncbi.nlm.nih.gov/pubmed/16024512*

Krieg, M., Stühmer, J., Cueva, J. G., Fetter, R., Spilker, K., Cremers, D., Shen, K., Dunn, A. R., Goodman, M. B. (2017). Genetic defects in β-spectrin and tau sensitize *C. elegans* axons to movement-induced damage via torque-tension coupling. *Elife* 6, e20172 -- *https://doi.org/10.7554/eLife.20172*

Kuijpers, M., van de Willige, D., Freal, A., Chazeau, A., Franker, Mariella A., Hofenk, J., Rodrigues, Ricardo J. C., Kapitein, Lukas C., Akhmanova, A., Jaarsma, D., Hoogenraad, Casper C. (2016). Dynein regulator NDEL1 controls polarized cargo transport at the axon initial segment. *Neuron* 89, 461-471 -- *http://www.sciencedirect.com/science/article/pii/S0896627316000477*

Lacroix, B., van Dijk, J., Gold, N. D., Guizetti, J., Aldrian-Herrada, G., Rogowski, K., Gerlich, D. W., Janke, C. (2010). Tubulin polyglutamylation stimulates spastin-mediated microtubule severing. *J Cell Biol* 189, 945-54 -- *http://www.ncbi.nlm.nih.gov/pubmed/20530212*

Lam, A. T., Curschellas, C., Krovvidi, D., Hess, H. (2014). Controlling self-assembly of microtubule spools via kinesin motor density. *Soft Matter* 10, 8731-8736 -- *http://dx.doi.org/10.1039/C4SM01518E*

Lam, A. T., VanDelinder, V., Kabir, A. M. R., Hess, H., Bachand, G. D., Kakugo, A. (2016). Cytoskeletal motor-driven active self-assembly in *in vitro* systems. *Soft Matter* 12, 988-997 -- *http://dx.doi.org/10.1039/C5SM02042E*

Lamoureux, P., Heidemann, S. R., Martzke, N. R., Miller, K. E. (2010). Growth and elongation within and along the axon. *Dev Neurobiol* 70, 135-49 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids= 19950193*

Langford, G. M. (1980). Arrangement of subunits in microtubules with 14 profilaments. *J Cell Biol* 87, 521-6 -- *http://www.ncbi.nlm.nih.gov/pubmed/7430256*

Lansky, Z., Braun, M., Lüdecke, A., Schlierf, M., ten Wolde, Pieter R., Janson, Marcel E., Diez, S. (2015). Diffusible crosslinkers generate directed forces in microtubule networks. *Cell* 160, 1159-68 -- *http://www.cell.com/cell/abstract/S0092-8674(15)00129-4*

Lazarus, C., Soheilypour, M., Mofrad, Mohammad R. K. (2015). Torsional behavior of axonal microtubule
bundles. *Biophys J* 109, 231-239 --
http://www.sciencedirect.com/science/article/pii/S0006349515006128

Lazarus, J. E., Moughamian, A. J., Tokito, M. K., Holzbaur, E. L. (2013). Dynactin subunit p150(Glued) is
a neuron-specific anti-catastrophe factor. *PLoS Biol* 11, e1001611 --
http://www.ncbi.nlm.nih.gov/pubmed/23874158

Lee, G., Brandt, R. (1992). Microtubule bundling studies revisited: is there a role for MAPs? *Trends in Cell*
Biology 2, 286-289 -- *http://www.sciencedirect.com/science/article/pii/096289249290106W*

Leo, L., Yu, W., D'Rozario, M., Waddell, E. A., Marenda, D. R., Baird, M. A., Davidson, M. W., Zhou, B.,
Wu, B., Baker, L., Sharp, D. J., Baas, P. W. (2015). Vertebrate fidgetin restrains axonal growth by
severing labile domains of microtubules. *Cell Rep* 12, 1723-30 --
http://dx.doi.org/10.1016/j.celrep.2015.08.017

Leterrier, C. (2018). The axon initial segment: an updated viewpoint. *J Neurosci* 38, 2135-2145 --
http://www.jneurosci.org/content/jneuro/38/9/2135.full.pdf

Leterrier, C., Dubey, P., Roy, S. (2017). The nano-architecture of the axonal cytoskeleton. *Nature*
Reviews Neuroscience 18, 713-726 -- *http://dx.doi.org/10.1038/nrn.2017.129*

Letourneau, P. C., Shattuck, T. A., Ressler, A. H. (1987). "Pull" and "push" in neurite elongation:
observations on the effects of different concentrations of cytochalasin B and taxol. *Cell Motil*
Cytoskeleton 8, 193-209 --
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=
2891448

Li, S., Wang, C., Nithiarasu, P. (2018). Effects of the cross-linkers on the buckling of microtubules in cells.
J Biomech Eng -- *http://www.sciencedirect.com/science/article/pii/S0021929018301544*

Lin, S., Liu, M., Mozgova, O. I., Yu, W., Baas, P. W. (2012). Mitotic motors coregulate microtubule
patterns in axons and dendrites. *J Neurosci* 32, 14033-49 --
http://www.ncbi.nlm.nih.gov/pubmed/23035110

Liu, H., Bachand, G. D. (2013). Effects of Confinement on Molecular Motor-Driven Self-Assembly of Ring
Structures. *Cellular and Molecular Bioengineering* 6, 98-108 -- *https://doi.org/10.1007/s12195-012-*
0256-5

Liu, H., Spoerke, E. D., Bachand, M., Koch, S. J., Bunker, B. C., Bachand, G. D. (2008). Biomolecular
Motor-Powered Self-Assembly of Dissipative Nanocomposite Rings. *Advanced Materials* 20, 4476-
4481 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/adma.200801291*

Liu, L., Tuzel, E., Ross, J. L. (2011). Loop formation of microtubules during gliding at high density. *J Phys*
Condens Matter 23, 374104 -- *http://www.ncbi.nlm.nih.gov/pubmed/21862840*

Liu, M., Nadar, V. C., Kozielski, F., Kozlowska, M., Yu, W., Baas, P. W. (2010). Kinesin-12, a mitotic
microtubule-associated motor protein, impacts axonal growth, navigation, and branching. *J Neurosci*
30, 14896-906 --
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=
21048148

Liu, Z., Zhou, T., Ziegler, A. C., Dimitrion, P., Zuo, L. (2017). Oxidative Stress in Neurodegenerative
Diseases: From Molecular Mechanisms to Clinical Applications. *Oxid Med Cell Longev* 2017,
2525967 -- *http://www.ncbi.nlm.nih.gov/pubmed/28785371*

Lu, W., Fox, P., Lakonishok, M., Davidson, Michael W., Gelfand, Vladimir I. (2013). Initial neurite
outgrowth in *Drosophila* neurons is driven by Kinesin-powered microtubule sliding. *Current Biology*
23, 1018-1023 -- *http://www.sciencedirect.com/science/article/pii/S0960982213004910*

Lu, W., Gelfand, V. I. (2017). Moonlighting Motors: Kinesin, Dynein, and Cell Polarity. *Trends Cell Biol* --
http://www.ncbi.nlm.nih.gov/pubmed/28284467

Luria, I., Crenshaw, J., Downs, M., Agarwal, A., Seshadri, S. B., Gonzales, J., Idan, O., Kamcev, J.,
Katira, P., Pandey, S., Nitta, T., Phillpot, S. R., Hess, H. (2011). Microtubule nanospool formation by
active self-assembly is not initiated by thermal activation. *Soft Matter* 7, 3108-3115 --
http://dx.doi.org/10.1039/C0SM00802H

Maas, T., Eidenmuller, J., Brandt, R. (2000). Interaction of tau with the neural membrane cortex is
regulated by phosphorylation at sites that are modified in paired helical filaments. *J. Biol. Chem.* 275,
15733-40 -- *http://www.jbc.org.manchester.idm.oclc.org/content/275/21/15733*

Machin, N. A., Lee, J. M., Barnes, G. (1995). Microtubule stability in budding yeast: characterization and dosage suppression of a benomyl-dependent tubulin mutant. *Mol Biol Cell* 6, 1241-1259 -- *https://www.molbiolcell.org/doi/abs/10.1091/mbc.6.9.1241*

Mao, C.-X., Xiong, Y., Xiong, Z., Wang, Q., Zhang, Y. Q., Jin, S. (2014). Microtubule-severing protein Katanin regulates neuromuscular junction development and dendritic elaboration in *Drosophila*. *Development* 141, 1064-1074 -- *http://dev.biologists.org/content/141/5/1064.abstract*

Marner, L., Nyengaard, J. R., Tang, Y., Pakkenberg, B. (2003). Marked loss of myelinated nerve fibers in the human brain with age. *J Comp Neurol* 462, 144-52 -- *http://www.ncbi.nlm.nih.gov/pubmed/12794739*

McBride, H. M., Neuspiel, M., Wasiak, S. (2006). Mitochondria: more than just a powerhouse. *Curr Biol* 16, R551-60 -- *http://www.ncbi.nlm.nih.gov/pubmed/16860735*

McNally, F. J., Roll-Mecak, A. (2018). Microtubule-severing enzymes: From cellular functions to molecular mechanism. *J Cell Biol* 217, 4057-69 -- *http://jcb.rupress.org/content/early/2018/10/31/jcb.201612104*

McVicker, D. P., Millette, M. M., Dent, E. W. (2015). Signaling to the microtubule cytoskeleton: an unconventional role for CaMKII. *Dev Neurobiol* 75, 423-34 -- *http://www.ncbi.nlm.nih.gov/pubmed/25156276*

Medana, I. M., Esiri, M. M. (2003). Axonal damage: a key predictor of outcome in human CNS diseases. *Brain* 126, 515-30 -- *http://www.ncbi.nlm.nih.gov/pubmed/12566274*

Memet, E., Hilitski, F., Morris, M. A., Schwenger, W. J., Dogic, Z., Mahadevan, L. (2018). Microtubules soften due to cross-sectional flattening. *Elife* 7 -- *http://www.ncbi.nlm.nih.gov/pubmed/29856317*

Méphon-Gaspard, A., Boca, M., Pioche-Durieu, C., Desforges, B., Burgo, A., Hamon, L., Piétrement, O., Pastré, D. (2016). Role of tau in the spatial organization of axonal microtubules: keeping parallel microtubules evenly distributed despite macromolecular crowding. *Cell Mol Life Sci* 73, 3745-3760 -- *https://www.ncbi.nlm.nih.gov/pubmed/27076215*

Miller, K. E., Sheetz, M. P. (2006). Direct evidence for coherent low velocity axonal transport of mitochondria. *J Cell Biol* 173, 373-81 -- *http://www.ncbi.nlm.nih.gov/pubmed/16682527*

Mohan, R., John, A. (2015). Microtubule-associated proteins as direct crosslinkers of actin filaments and microtubules. *IUBMB Life* 67, 395-403 -- *https://iubmb.onlinelibrary.wiley.com/doi/abs/10.1002/iub.1384*

Monroy, B. Y., Sawyer, D. L., Ackermann, B. E., Borden, M. M., Tan, T. C., Ori-McKenney, K. M. (2018). Competition between microtubule-associated proteins directs motor transport. *Nat Commun* 9, 1487 -- *https://doi.org/10.1038/s41467-018-03909-2*

Morikawa, M., Yajima, H., Nitta, R., Inoue, S., Ogura, T., Sato, C., Hirokawa, N. (2015). X-ray and Cryo-EM structures reveal mutual conformational changes of Kinesin and GTP-state microtubules upon binding. *EMBO J* 34, 1270-86 -- *http://www.ncbi.nlm.nih.gov/pubmed/25777528*

Muto, E., Sakai, H., Kaseda, K. (2005). Long-range cooperative binding of kinesin to a microtubule in the presence of ATP. *J Cell Biol* 168, 691-6 -- *http://www.ncbi.nlm.nih.gov/pubmed/15738263*

Myers, K. A., Baas, P. W. (2007). Kinesin-5 regulates the growth of the axon by acting as a brake on its microtubule array. *J Cell Biol* 178, 1081-91 -- *http://www.ncbi.nlm.nih.gov/pubmed/17846176*

Myers, K. A., He, Y., Hasaka, T. P., Baas, P. W. (2006). Microtubule transport in the axon: Re-thinking a potential role for the actin cytoskeleton. *Neuroscientist* 12, 107-18 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16514008*

Nadar, V. C., Lin, S., Baas, P. W. (2012). Microtubule redistribution in growth cones elicited by focal inactivation of kinesin-5. *J Neurosci* 32, 5783-94 -- *http://www.ncbi.nlm.nih.gov/pubmed/22539840*

Nashchekin, D., Fernandes, Artur R., St Johnston, D. (2016). Patronin/Shot cortical foci assemble the noncentrosomal microtubule array that specifies the *Drosophila* anterior-posterior axis. *Dev Cell* 38, 61-72 -- *http://dx.doi.org/10.1016/j.devcel.2016.06.010*

Nguyen, M. D., Lariviere, R. C., Julien, J. P. (2000). Reduction of axonal caliber does not alleviate motor neuron disease caused by mutant superoxide dismutase 1. *Proc Natl Acad Sci U S A* 97, 12306-11 -- *http://www.ncbi.nlm.nih.gov/pubmed/11050249*

Ning, W., Yu, Y., Xu, H., Liu, X., Wang, D., Wang, J., Wang, Y., Meng, W. (2016). The CAMSAP3-ACF7 complex couples noncentrosomal microtubules with actin filaments to coordinate their dynamics. *Dev Cell* 39, 61-74 -- *http://www.sciencedirect.com/science/article/pii/S1534580716305998*

Nogales, E., Wolf, S. G., Khan, I. A., Luduena, R. F., Downing, K. H. (1995). Structure of tubulin at 6.5 A and location of the taxol-binding site. *Nature* 375, 424-7 -- *https://www.nature.com/articles/375424a0*

Noordstra, I., Liu, Q., Nijenhuis, W., Hua, S., Jiang, K., Baars, M., Remmelzwaal, S., Martin, M., Kapitein, L. C., Akhmanova, A. (2016). Control of apico-basal epithelial polarity by the microtubule minus-end-binding protein CAMSAP3 and spectraplakin ACF7. *J Cell Sci* 129, 4278-4288 -- *http://www.ncbi.nlm.nih.gov/pubmed/27802168*

O'Brien, E. T., Salmon, E. D., Erickson, H. P. (1997). How calcium causes microtubule depolymerization. *Cell Motility* 36, 125-135 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/%28SICI%291097-0169%281997%2936%3A2%3C125%3A%3AAID-CM3%3E3.0.CO%3B2-8*

O'Toole, M., Lamoureux, P., Miller, K. E. (2008). A physical model of axonal elongation: force, viscosity, and adhesions govern the mode of outgrowth. *Biophys J* 94, 2610-20 -- *http://www.ncbi.nlm.nih.gov/pubmed/18178646*

O'Toole, M., Lamoureux, P., Miller, Kyle E. (2015). Measurement of subcellular force generation in neurons. *Biophys J* 108, 1027-1037 -- *http://www.sciencedirect.com/science/article/pii/S0006349515001150*

Odde, D. J., Ma, L., Briggs, A. H., DeMarco, A., Kirschner, M. W. (1999). Microtubule bending and breaking in living fibroblast cells. *J Cell Sci* 112, 3283-3288 -- *http://jcs.biologists.org/content/joces/112/19/3283.full.pdf*

Pan, S., Chan, J. R. (2017). Regulation and dysregulation of axon infrastructure by myelinating glia. *Journal Cell Biol* 216, 3903-3916 -- *http://jcb.rupress.org/content/jcb/216/12/3903.full.pdf*

Papadopoulos, C., Orso, G., Mancuso, G., Herholz, M., Gumeni, S., Tadepalle, N., Jungst, C., Tzschichholz, A., Schauss, A., Honing, S., Trifunovic, A., Daga, A., Rugarli, E. I. (2015). Spastin binds to lipid droplets and affects lipid metabolism. *PLoS Genet* 11, e1005149 -- *http://www.ncbi.nlm.nih.gov/pubmed/25875445*

Park, J. H., Roll-Mecak, A. (2018). The tubulin code in neuronal polarity. *Curr Opin Neurobiol* 51, 95-102 -- *http://www.ncbi.nlm.nih.gov/pubmed/29554585*

Park, S. H., Zhu, P. P., Parker, R. L., Blackstone, C. (2010). Hereditary spastic paraplegia proteins REEP1, spastin, and atlastin-1 coordinate microtubule interactions with the tubular ER network. *J Clin Invest* 120, 1097-110 -- *https://www.jci.org/articles/view/40979*

Pascual-Ahuir, A., Manzanares-Estreder, S., Proft, M. (2017). Pro- and antioxidant functions of the peroxisome-mitochondria connection and its impact on aging and disease. *Oxid Med Cell Longev* 2017, 9860841 -- *http://www.ncbi.nlm.nih.gov/pubmed/28811869*

Pearce, S. P., Heil, M., Jensen, O. E., Jones, G. W., Prokop, A. (2018). Curvature-sensitive kinesin binding can explain microtubule ring formation and reveals chaotic dynamics in a mathematical model. *Bull Math Biol* 80, 3002-22 -- *https://tinyurl.com/yd43ncb9*

Peet, D. R., Burroughs, N. J., Cross, R. A. (2018). Kinesin expands and stabilizes the GDP-microtubule lattice. *Nat Nanotechnol* 13, 386-391 -- *https://www-nature-com.manchester.idm.oclc.org/articles/s41565-018-0084-4*

Penazzi, L., Bakota, L., Brandt, R. (2016). Microtubule dynamics in neuronal development, plasticity, and neurodegeneration. *Int Rev Cell Mol Biol* 321, 89-169 -- http://www.ncbi.nlm.nih.gov/pubmed/26811287

Perrot, R., Berges, R., Bocquet, A., Eyer, J. (2008). Review of the multiple aspects of neurofilament functions, and their possible contribution to neurodegeneration. *Mol Neurobiol* 38, 27-65 -- *http://www.ncbi.nlm.nih.gov/pubmed/18649148*

Peter, S. J., Mofrad, M. R. (2012). Computational modeling of axonal microtubule bundles under tension. *Biophys J* 102, 749-57 -- *https://www.ncbi.nlm.nih.gov/pubmed/22385845*

Pfenninger, K. H. (2009). Plasma membrane expansion: a neuron's Herculean task. *Nat Rev Neurosci* 10, 251-61 -- *http://www.ncbi.nlm.nih.gov/pubmed/19259102*

Pines, M. K., Housden, B. E., Bernard, F., Bray, S. J., Roper, K. (2010). The cytolinker Pigs is a direct target and a negative regulator of Notch signalling. *Development* 137, 913-22 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20150280*

Popov, S., Brown, A., Poo, M. M. (1993). Forward plasma membrane flow in growing nerve processes. *Science* 259, 244-6 -- *http://www.ncbi.nlm.nih.gov/pubmed/7678471*

Preitner, N., Quan, J., Nowakowski, Dan W., Hancock, Melissa L., Shi, J., Tcherkezian, J., Young-Pearse, Tracy L., Flanagan, John G. (2014). APC is an RNA-binding protein, and its interactome provides a link to neural development and microtubule assembly. *Cell* 158, 368-382 -- *http://www.sciencedirect.com/science/article/pii/S0092867414007478*

Prezel, E., Elie, A., Delaroche, J., Stoppin-Mellet, V., Bosc, C., Serre, L., Fourest-Lieuvin, A., Andrieux, A., Vantard, M., Arnal, I., Zhu, X. (2018). Tau can switch microtubule network organizations: from random networks to dynamic and stable bundles. *Molecular Biology of the Cell* 29, 154-165 -- *https://www.molbiolcell.org/doi/abs/10.1091/mbc.E17-06-0429*

Prior, R., Van Helleputte, L., Benoy, V., Van Den Bosch, L. (2017). Defective axonal transport: A common pathological mechanism in inherited and acquired peripheral neuropathies. *Neurobiol Dis* 105, 300-320 -- *http://www.ncbi.nlm.nih.gov/pubmed/28238949*

Prokop, A. (2013). The intricate relationship between microtubules and their associated motor proteins during axon growth and maintenance. *Neur Dev* 8, 17 -- *http://www.neuraldevelopment.com/content/8/1/17*

Prokop, A. (2016). Fruit flies in biological research. *Biological Sciences Review* 28, 10-14 -- *https://tinyurl.com/ybvpoqmw*

Prokop, A. (2018). Why funding fruit fly research is important for the biomedical sciences. *Open Access Govern* 20, 198-201 -- *https://tinyurl.com/y7b25jpm*

Prokop, A., Beaven, R., Qu, Y., Sánchez-Soriano, N. (2013). Using fly genetics to dissect the cytoskeletal machinery of neurons during axonal growth and maintenance. *J. Cell Sci.* 126, 2331-41 -- *http://dx.doi.org/10.1242/jcs.126912*

Prokop, A., Küppers-Munther, B., Sánchez-Soriano, N. (2012). Using primary neuron cultures of *Drosophila* to analyse neuronal circuit formation and function. *In* "The making and un-making of neuronal circuits in *Drosophila*" (B. A. Hassan, Ed.), Vol. 69, pp. 225-47. Humana Press, New York -- *http://dx.doi.org/10.1007/978-1-61779-830-6_10*

Prokop, A., Uhler, J., Roote, J., Bate, M. C. (1998). The *kakapo* mutation affects terminal arborisation and central dendritic sprouting of *Drosophila* motorneurons. *J. Cell Biol.* 143, 1283-1294 -- *http://jcb.rupress.org/content/143/5/1283.long*

Pronker, M. F., Lemstra, S., Snijder, J., Heck, A. J. R., Thies-Weesie, D. M. E., Pasterkamp, R. J., Janssen, B. J. C. (2016). Structural basis of myelin-associated glycoprotein adhesion and signalling. *Nature Communications* 7, 13584 -- *https://doi.org/10.1038/ncomms13584*

Qiang, L., Sun, X., Austin, T. O., Muralidharan, H., Jean, D. C., Liu, M., Yu, W., Baas, P. W. (2018). Tau does not stabilize axonal microtubules but rather enables them to have long labile domains. *Curr Biol* 28, 2181-2189 e4 -- *http://www.ncbi.nlm.nih.gov/pubmed/30008334*

Qiang, L., Yu, W., Andreadis, A., Luo, M., Baas, P. W. (2006). Tau protects microtubules in the axon from severing by katanin. *J Neurosci* 26, 3120-9 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16554463*

Qu, Y., Hahn, I., Lees, M., Parkin, J., Voelzmann, A., Dorey, K., Rathbone, A., Friel, C., Allan, V., Okenve Ramos, P., Sánchez-Soriano, N., Prokop, A. (2018). Efa6 regulates axon growth, branching and maintenance by eliminating off-track microtubules at the cortex. *bioRxiv* 10.1101/385658 -- *https://www.biorxiv.org/content/early/2018/09/03/385658*

Qu, Y., Hahn, I., Webb, S. E. D., Pearce, S. P., Prokop, A. (2017). Periodic actin structures in neuronal axons are required to maintain microtubules. *Mol Biol Cell* 28 296-308 -- *http://www.molbiolcell.org/content/early/2016/11/21/mbc.E16-10-0727*

Rajendran, L., Bali, J., Barr, M. M., Court, F. A., Kramer-Albers, E. M., Picou, F., Raposo, G., van der Vos, K. E., van Niel, G., Wang, J., Breakefield, X. O. (2014). Emerging roles of extracellular vesicles in the nervous system. *J Neurosci* 34, 15482-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/25392515*

Rao, M. V., Campbell, J., Yuan, A., Kumar, A., Gotow, T., Uchiyama, Y., Nixon, R. A. (2003). The neurofilament middle molecular mass subunit carboxyl-terminal tail domains is essential for the radial growth and cytoskeletal architecture of axons but not for regulating neurofilament transport rate. *J Cell Biol* 163, 1021-31 -- *http://www.ncbi.nlm.nih.gov/pubmed/14662746*

Rashedul Kabir, A. M., Wada, S., Inoue, D., Tamura, Y., Kajihara, T., Mayama, H., Sada, K., Kakugo, A., Gong, J. P. (2012). Formation of ring-shaped assembly of microtubules with a narrow size distribution at an air-buffer interface. *Soft Matter* 8, 10863-10867 -- *http://dx.doi.org/10.1039/C2SM26441B*

Ray, S., Meyhofer, E., Milligan, R. A., Howard, J. (1993). Kinesin follows the microtubule's protofilament axis. *J Cell Biol* 121, 1083-93 -- *http://www.ncbi.nlm.nih.gov/pubmed/8099076*

Reinsch, S. S., Mitchison, T. J., Kirschner, M. (1991). Microtubule polymer assembly and transport during axonal elongation. *J Cell Biol* 115, 365-79 -- *http://www.ncbi.nlm.nih.gov/pubmed/1717484*

Riancho, J., Gonzalo, I., Ruiz-Soto, M., Berciano, J. (2019). Why do motor neurons degenerate? Actualization in the pathogenesis of amyotrophic lateral sclerosis. *Neurología* 34, 27-37 -- *http://www.elsevier.es/en-revista-neurologia-english-edition--495-articulo-why-do-motor-neurons-degenerate-S2173580817301633*

Riano, E., Martignoni, M., Mancuso, G., Cartelli, D., Crippa, F., Toldo, I., Siciliano, G., Di Bella, D., Taroni, F., Bassi, M. T., Cappelletti, G., Rugarli, E. I. (2009). Pleiotropic effects of spastin on neurite growth depending on expression levels. *J Neurochem* 108, 1277-88 -- *https://doi-org.manchester.idm.oclc.org/10.1111/j.1471-4159.2009.05875.x*

Rieusset, J. (2017). Endoplasmic reticulum-mitochondria calcium signaling in hepatic metabolic diseases. *Biochim Biophys Acta Mol Cell Res* 1864, 865-876 -- *http://www.ncbi.nlm.nih.gov/pubmed/28064001*

Roos, J., Hummel, T., Ng, N., Klämbt, C., Davis, G. W. (2000). *Drosophila* Futsch regulates synaptic microtubule organisation and is necessary for synaptic growth. *Neuron* 26, 371-382 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10839356*

Roossien, D. H., Lamoureux, P., Miller, K. E. (2014). Cytoplasmic dynein pushes the cytoskeletal meshwork forward during axonal elongation. *J Cell Sci* 127, 3593-602 -- *http://www.ncbi.nlm.nih.gov/pubmed/24951117*

Roossien, D. H., Lamoureux, P., Van Vactor, D., Miller, K. E. (2013). *Drosophila* growth cones advance by forward translocation of the neuronal cytoskeletal meshwork *in vivo*. *PLoS One* 8, e80136 -- *http://www.ncbi.nlm.nih.gov/pubmed/24244629*

Roote, J., Prokop, A. (2013). How to design a genetic mating scheme: a basic training package for *Drosophila* genetics. *G3 (Bethesda)* 3, 353-8 -- *http://www.g3journal.org/content/3/2/353.full*

Rosenberg, K. J., Ross, J. L., Feinstein, H. E., Feinstein, S. C., Israelachvili, J. (2008). Complementary dimerization of microtubule-associated tau protein: Implications for microtubule bundling and tau-mediated pathogenesis. *Proc Natl Acad Sci U S A* 105, 7445-50 -- *http://www.ncbi.nlm.nih.gov/pubmed/18495933*

Sakaguchi, T., Okada, M., Kitamura, T., Kawasaki, K. (1993). Reduced diameter and conduction velocity of myelinated fibers in the sciatic nerve of a neurofilament-deficient mutant quail. *Neurosci Lett* 153, 65-8 -- *http://www.ncbi.nlm.nih.gov/pubmed/8510825*

Sakakibara, A., Ando, R., Sapir, T., Tanaka, T. (2013). Microtubule dynamics in neuronal morphogenesis. *Open Biology* 3, 130061 -- *https://royalsocietypublishing.org/doi/abs/10.1098/rsob.130061*

Salvadores, N., Sanhueza, M., Manque, P., Court, F. A. (2017). Axonal degeneration during aging and its functional role in neurodegenerative disorders. *Front Neurosci* 11 -- *https://www.frontiersin.org/article/10.3389/fnins.2017.00451*

Samsonov, A., Yu, J. Z., Rasenick, M., Popov, S. V. (2004). Tau interaction with microtubules *in vivo*. *J Cell Sci* 117, 6129-41 -- *http://www.ncbi.nlm.nih.gov/pubmed/15564376*

Sánchez-Soriano, N., Gonçalves-Pimentel, C., Beaven, R., Haessler, U., Ofner, L., Ballestrem, C., Prokop, A. (2010). *Drosophila* growth cones: a genetically tractable platform for the analysis of axonal growth dynamics. *Dev. Neurobiol.* 70, 58-71 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/dneu.20762*

Sánchez-Soriano, N., Tear, G., Whitington, P., Prokop, A. (2007). *Drosophila* as a genetic and cellular model for studies on axonal growth. *Neural Develop* 2, 9 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17475018*

Sánchez-Soriano, N., Travis, M., Dajas-Bailador, F., Goncalves-Pimentel, C., Whitmarsh, A. J., Prokop, A. (2009). Mouse ACF7 and *Drosophila* Short stop modulate filopodia formation and microtubule organisation during neuronal growth. *J Cell Sci* 122, 2534-42 -- *http://jcs.biologists.org/content/122/14/2534.long*

Satake, T., Yamashita, K., Hayashi, K., Miyatake, S., Tamura-Nakano, M., Doi, H., Furuta, Y., Shioi, G., Miura, E., Takeo, Y. H., Yoshida, K., Yahikozawa, H., Matsumoto, N., Yuzaki, M., Suzuki, A. (2017). MTCL1 plays an essential role in maintaining Purkinje neuron axon initial segment. *EMBO J* 36, 1227-1242 -- *http://emboj.embopress.org/content/embojnl/36/9/1227.full.pdf*

Savage, C., Hamelin, M., Culotti, J. G., Coulson, A., Albertson, D. G., Chalfie, M. (1989). mec-7 is a beta-tubulin gene required for the production of 15-protofilament microtubules in *Caenorhabditis elegans*. *Genes Dev* 3, 870-81 -- *http://www.ncbi.nlm.nih.gov/pubmed/2744465*

Saxton, W. M., Hollenbeck, P. J. (2012). The axonal transport of mitochondria. *J Cell Sci* 125, 2095-104 -- *http://www.ncbi.nlm.nih.gov/pubmed/22619228*

Sayas, C. L., Tortosa, E., Bollati, F., Ramirez-Rios, S., Arnal, I., Avila, J. (2015). Tau regulates the localization and function of End-binding proteins 1 and 3 in developing neuronal cells. *J Neurochem* 133, 653-67 -- *http://www.ncbi.nlm.nih.gov/pubmed/25761518*

Scaife, R., Margolis, R. L. (1990). Biochemical and immunochemical analysis of rat brain dynamin interaction with microtubules and organelles *in vivo* and *in vitro*. *J Cell Biol* 111, 3023-33 -- *http://www.ncbi.nlm.nih.gov/pubmed/2148566*

Schelski, M., Bradke, F. (2017). Neuronal polarization: From spatiotemporal signaling to cytoskeletal dynamics. *Mol Cell Neurosci* 84, 11-28 -- *http://www.ncbi.nlm.nih.gov/pubmed/28363876*

Schüle, R., Wiethoff, S., Martus, P., Karle, K. N., Otto, S., Klebe, S., Klimpe, S., Gallenmüller, C., Kurzwelly, D., Henkel, D., Rimmele, F., Stolze, H., Kohl, Z., Kassubek, J., Klockgether, T., Vielhaber, S., Kamm, C., Klopstock, T., Bauer, P., Züchner, S., Liepelt-Scarfone, I., Schöls, L. (2016). Hereditary spastic paraplegia: Clinicogenetic lessons from 608 patients. *Annals of Neurology* 79, 646-658 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/ana.24611*

Sharp, D. J., Ross, J. L. (2012). Microtubule-severing enzymes at the cutting edge. *J Cell Sci* 125, 2561-9 -- *http://www.ncbi.nlm.nih.gov/pubmed/22595526*

Shaw, G., Bray, D. (1977). Movement and extension of isolated growth cones. *Exp Cell Res* 104, 55-62 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=556695*

Sheng, Z. H. (2017). The interplay of axonal energy homeostasis and mitochondrial trafficking and anchoring. *Trends Cell Biol* 27, 403-416 -- *http://www.ncbi.nlm.nih.gov/pubmed/28228333*

Shin, S., Lim, S., Jeong, H., Kwan, L., Kim, Y. (2018). Visualization of tau–tubulin interaction in a living cell using bifluorescence complementation technique. *International Journal of Molecular Sciences* 19, 2978 -- *http://www.mdpi.com/1422-0067/19/10/2978*

Shin, S. C., Im, S.-K., Jang, E.-H., Jin, K. S., Hur, E.-M., Kim, E. E. (2019). Structural and molecular basis for katanin-mediated severing of glutamylated microtubules. *Cell Reports* 26, 1357-1367.e5 -- *https://doi.org/10.1016/j.celrep.2019.01.020*

Shoukier, M., Neesen, J., Sauter, S. M., Argyriou, L., Doerwald, N., Pantakani, D. V., Mannan, A. U. (2009). Expansion of mutation spectrum, determination of mutation cluster regions and predictive structural classification of SPAST mutations in hereditary spastic paraplegia. *Eur J Hum Genet* 17, 187-94 -- *http://www.ncbi.nlm.nih.gov/pubmed/18701882*

Shpetner, H. S., Vallee, R. B. (1989). Identification of dynamin, a novel mechanochemical enzyme that mediates interactions between microtubules. *Cell* 59, 421-32 -- *http://www.ncbi.nlm.nih.gov/pubmed/2529977*

Sirajuddin, M., Rice, L. M., Vale, R. D. (2014). Regulation of microtubule motors by tubulin isotypes and post-translational modifications. *Nat Cell Biol* 16, 335-344 -- *http://dx.doi.org/10.1038/ncb2920*

Skruber, K., Read, T.-A., Vitriol, E. A. (2018). Reconsidering an active role for G-actin in cytoskeletal regulation. *J Cell Sci* 131 -- *http://jcs.biologists.org/content/joces/131/1/jcs203760.full.pdf*

Smith, D. H., Nonaka, M., Miller, R., Leoni, M., Chen, X. H., Alsop, D., Meaney, D. F. (2000). Immediate coma following inertial brain injury dependent on axonal damage in the brainstem. *J Neurosurg* 93, 315-22 -- *http://www.ncbi.nlm.nih.gov/pubmed/10930019*

Solowska, J. M., Baas, P. W. (2015). Hereditary spastic paraplegia SPG4: what is known and not known about the disease. *Brain* 138, 2471-84 -- *http://www.ncbi.nlm.nih.gov/pubmed/26094131*

Song, Y., Kirkpatrick, L. L., Schilling, A. B., Helseth, D. L., Chabot, N., Keillor, J. W., Johnson, G. V., Brady, S. T. (2013). Transglutaminase and polyamination of tubulin: posttranslational modification for stabilizing axonal microtubules. *Neuron* 78, 109-23 -- *http://www.ncbi.nlm.nih.gov/pubmed/23583110*

Soppina, V., Herbstman, J. F., Skiniotis, G., Verhey, K. J. (2012). Luminal localization of alpha-tubulin K40 acetylation by cryo-EM analysis of fab-labeled microtubules. *PLoS One* 7, e48204 -- *http://www.ncbi.nlm.nih.gov/pubmed/23110214*

Sorbara, C. D., Wagner, N. E., Ladwig, A., Nikic, I., Merkler, D., Kleele, T., Marinkovic, P., Naumann, R., Godinho, L., Bareyre, F. M., Bishop, D., Misgeld, T., Kerschensteiner, M. (2014). Pervasive axonal

transport deficits in multiple sclerosis models. *Neuron* 84, 1183-90 --
http://www.ncbi.nlm.nih.gov/pubmed/25433639

Staff, N. P., Podratz, J. L., Grassner, L., Bader, M., Paz, J., Knight, A. M., Loprinzi, C. L., Trushina, E.,
Windebank, A. J. (2013). Bortezomib alters microtubule polymerization and axonal transport in rat
dorsal root ganglion neurons. *Neurotoxicology* 39, 124-31 --
http://www.ncbi.nlm.nih.gov/pubmed/24035926

Steinmetz, M. O., Prota, A. E. (2018). Microtubule-targeting agents: strategies to hijack the cytoskeleton.
Trends Cell Biol 28, 776-792 -- *https://doi.org/10.1016/j.tcb.2018.05.001*

Stewart, A., Tsubouchi, A., Rolls, M. M., Tracey, W. D., Sherwood, N. T. (2012). Katanin p60-like1
promotes microtubule growth and terminal dendrite stability in the larval class IV sensory neurons of
Drosophila. *The Journal of Neuroscience* 32, 11631-11642 --
http://www.jneurosci.org/content/jneuro/32/34/11631.full.pdf

Stokin, G. B., Lillo, C., Falzone, T. L., Brusch, R. G., Rockenstein, E., Mount, S. L., Raman, R., Davies,
P., Masliah, E., Williams, D. S., Goldstein, L. S. (2005). Axonopathy and transport deficits early in the
pathogenesis of Alzheimer's disease. *Science* 307, 1282-8 --
http://www.ncbi.nlm.nih.gov/pubmed/15731448

Stone, M. C., Rao, K., Gheres, K. W., Kim, S., Tao, J., La Rochelle, C., Folker, C. T., Sherwood, N. T.,
Rolls, M. M. (2012). Normal spastin gene dosage is specifically required for axon regeneration. *Cell*
Rep 2, 1340-50 -- *http://www.ncbi.nlm.nih.gov/pubmed/23122959*

Stroud, M. J., Nazgiewicz, A., McKenzie, E. A., Wang, Y., Kammerer, R. A., Ballestrem, C. (2014). GAS2-
like proteins mediate communication between microtubules and actin through interactions with end-
binding proteins. *J Cell Sci* 127, 2672-82 -- *http://www.ncbi.nlm.nih.gov/pubmed/24706950*

Sturgill, E. G., Ohi, R. (2013). Microtubule-regulating kinesins. *Current Biology* 23, R946-R948 --
http://www.sciencedirect.com/science/article/pii/S0960982213009883

Sturrock, R. R. (1987). Age-related changes in the number of myelinated axons and glial cells in the
anterior and posterior limbs of the mouse anterior commissure. *J Anat* 150, 111-127 --
https://www.ncbi.nlm.nih.gov/pubmed/3654327

Subramaniyan Parimalam, S., Tarhan, M. C., Karsten, S. L., Fujita, H., Shintaku, H., Kotera, H.,
Yokokawa, R. (2016). On-chip microtubule gliding assay for parallel measurement of tau protein
species. *Lab on a Chip* 16, 1691-1697 -- *http://dx.doi.org/10.1039/C5LC01486G*

Sudo, H., Baas, P. W. (2010). Acetylation of microtubules influences their sensitivity to severing by
katanin in neurons and fibroblasts. *J Neurosci* 30, 7215-26 --
http://www.ncbi.nlm.nih.gov/pubmed/20505088

Sumino, Y., Nagai, K. H., Shitaka, Y., Tanaka, D., Yoshikawa, K., Chate, H., Oiwa, K. (2012). Large-scale
vortex lattice emerging from collectively moving microtubules. *Nature* 483, 448-52 --
http://www.ncbi.nlm.nih.gov/pubmed/22437613

Takei, Y., Teng, J., Harada, A., Hirokawa, N. (2000). Defects in axonal elongation and neuronal migration
in mice with disrupted *tau* and *map1b* genes. *J. Cell Biol.* 150, 989-1000 --
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=
10973990

Tang-Schomer, M. D., Johnson, V. E., Baas, P. W., Stewart, W., Smith, D. H. (2012). Partial interruption
of axonal transport due to microtubule breakage accounts for the formation of periodic varicosities
after traumatic axonal injury. *Exp Neurol* 233, 364-72 --
http://www.ncbi.nlm.nih.gov/pubmed/22079153

Tao, J., Feng, C., Rolls, M. M. (2016). The microtubule-severing protein fidgetin acts after dendrite injury
to promote their degeneration. *J Cell Sci* 129, 3274-81 --
http://www.ncbi.nlm.nih.gov/pubmed/27411367

Tarrade, A., Fassier, C., Courageot, S., Charvin, D., Vitte, J., Peris, L., Thorel, A., Mouisel, E.,
Fonknechten, N., Roblot, N., Seilhean, D., Dierich, A., Hauw, J. J., Melki, J. (2006). A mutation of
spastin is responsible for swellings and impairment of transport in a region of axon characterized by
changes in microtubule composition. *Hum Mol Genet* 15, 3544-58 --
http://www.ncbi.nlm.nih.gov/pubmed/17101632

Tas, R. P., Chazeau, A., Cloin, B. M. C., Lambers, M. L. A., Hoogenraad, C. C., Kapitein, L. C. (2017).
Differentiation between oppositely oriented microtubules controls polarized neuronal transport.
Neuron -- *https://www.sciencedirect.com/science/article/pii/S0896627317310711*

Tedeschi, A., Bradke, F. (2016). Spatial and temporal arrangement of neuronal intrinsic and extrinsic mechanisms controlling axon regeneration. *Curr Opin Neurobiol* 42, 118-127 -- *http://www.ncbi.nlm.nih.gov/pubmed/28039763*

Ti, S.-C., Alushin, G. M., Kapoor, T. M. (2018). Human β-tubulin isotypes can regulate microtubule protofilament number and stability. *Developmental Cell* 47, 175-190.e5 -- *http://www.sciencedirect.com/science/article/pii/S1534580718306841*

Tint, I., Jean, D., Baas, P. W., Black, M. M. (2009). Doublecortin associates with microtubules preferentially in regions of the axon displaying actin-rich protrusive structures. *J Neurosci* 29, 10995-1010 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19726658*

Triclin, S., Inoue, D., Gaillard, J., Htet, Z. M., De Santis, M., Portran, D., Derivery, E., Aumeier, C., Schaedel, L., John, K., Leterrier, C., Reck-Peterson, S., Blanchoin, L., Thery, M. (2018). Self-repair protects microtubules from their destruction by molecular motors. *bioRxiv*, 499020 -- *https://www.biorxiv.org/content/biorxiv/early/2018/12/17/499020.full.pdf*

Trinczek, B., Ebneth, A., Mandelkow, E. M., Mandelkow, E. (1999). Tau regulates the attachment/detachment but not the speed of motors in microtubule-dependent transport of single vesicles and organelles. *J Cell Sci* 112, 2355-2367 -- *http://jcs.biologists.org/content/joces/112/14/2355.full.pdf*

Tymanskyj, S. R., Yang, B., Falnikar, A., Lepore, A. C., Ma, L. (2017). MAP7 regulates axon collateral branch development in dorsal root ganglion neurons. *J Neurosci* 37, 1648-1661 -- *http://www.ncbi.nlm.nih.gov/pubmed/28069923*

Valenstein, M. L., Roll-Mecak, A. (2016). Graded control of microtubule severing by tubulin glutamylation. *Cell* 164, 911-21 -- *https://www.sciencedirect.com/science/article/pii/S0092867416000593?via%3Dihub*

van der Vaart, B., van Riel, Wilhelmina E., Doodhi, H., Kevenaar, Josta T., Katrukha, Eugene A., Gumy, L., Bouchet, Benjamin P., Grigoriev, I., Spangler, Samantha A., Yu, Ka L., Wulf, Phebe S., Wu, J., Lansbergen, G., van Battum, Eljo Y., Pasterkamp, R. J., Mimori-Kiyosue, Y., Demmers, J., Olieric, N., Maly, Ivan V., Hoogenraad, Casper C., Akhmanova, A. (2013). CFEOM1-associated kinesin KIF21A is a cortical microtubule growth inhibitor. *Developmental Cell* 27, 145-160 -- *http://dx.doi.org/10.1016/j.devcel.2013.09.010*

van Haren, J., Wittmann, T. (2019). Microtubule plus end dynamics − do we know how microtubules grow? *BioEssays* 41, 1800194 -- *https://onlinelibrary.wiley.com/doi/abs/10.1002/bies.201800194*

VanDelinder, V., Adams, P. G., Bachand, G. D. (2016a). Mechanical splitting of microtubules into protofilament bundles by surface-bound kinesin-1. *Sci Rep* 6, 39408 -- *http://www.ncbi.nlm.nih.gov/pubmed/28000714*

VanDelinder, V., Brener, S., Bachand, G. D. (2016b). Mechanisms Underlying the Active Self-Assembly of Microtubule Rings and Spools. *Biomacromolecules* 17, 1048-56 -- *http://www.ncbi.nlm.nih.gov/pubmed/26842978*

Vemu, A., Atherton, J., Spector, J. O., Moores, C. A., Roll-Mecak, A. (2017). Tubulin isoform composition tunes microtubule dynamics. *Mol Biol Cell* 28, 3564-3572 -- *http://www.ncbi.nlm.nih.gov/pubmed/29021343*

Vemu, A., Szczesna, E., Zehr, E. A., Spector, J. O., Grigorieff, N., Deaconescu, A. M., Roll-Mecak, A. (2018). Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation. *Science* 361 -- *http://www.ncbi.nlm.nih.gov/pubmed/30139843*

Villarroel-Campos, D., Gonzalez-Billault, C. (2014). The MAP1B case: an old MAP that is new again. *Dev Neurobiol* 74, 953-71 -- *http://www.ncbi.nlm.nih.gov/pubmed/24700609*

Voelzmann, A., Hahn, I., Pearce, S., Sánchez-Soriano, N. P., Prokop, A. (2016a). A conceptual view at microtubule plus end dynamics in neuronal axons. *Brain Res Bulletin* 126, 226-37 -- *http://www.sciencedirect.com/science/article/pii/S0361923016301885*

Voelzmann, A., Liew, Y.-T., Qu, Y., Hahn, I., Melero, C., Sánchez-Soriano, N., Prokop, A. (2017). *Drosophila* Short stop as a paradigm for the role and regulation of spectraplakins. *Sem Cell Dev Biol* 69, 40-57 -- *http://www.sciencedirect.com/science/article/pii/S1084952117302124*

Voelzmann, A., Okenve-Ramos, P., Qu, Y., Chojnowska-Monga, M., del Caño-Espinel, M., Prokop, A., Sánchez-Soriano, N. (2016b). Tau and spectraplakins promote synapse formation and maintenance

through Jun kinase and neuronal trafficking. *eLife* 5, e14694 -- *https://elifesciences.org/content/5/e14694*

Wada, S., Rashedul Kabir, A. M., Ito, M., Inoue, D., Sada, K., Kakugo, A. (2015). Effect of length and rigidity of microtubules on the size of ring-shaped assemblies obtained through active self-organization. *Soft Matter* 11, 1151-1157 -- *http://dx.doi.org/10.1039/C4SM02292K*

Walczak, C. E., Gayek, S., Ohi, R. (2013). Microtubule-depolymerizing kinesins. *Annu Rev Cell Dev Biol* 29, 417-41 -- *http://www.ncbi.nlm.nih.gov/pubmed/23875646*

Wali, G., Sue, C. M., Mackay-Sim, A. (2018). Patient-Derived Stem Cell Models in SPAST HSP: Disease Modelling and Drug Discovery. *Brain Sci* 8 -- *http://www.ncbi.nlm.nih.gov/pubmed/30065201*

Wali, G., Sutharsan, R., Fan, Y., Stewart, R., Tello Velasquez, J., Sue, C. M., Crane, D. I., Mackay-Sim, A. (2016). Mechanism of impaired microtubule-dependent peroxisome trafficking and oxidative stress in SPAST-mutated cells from patients with Hereditary Spastic Paraplegia. *Sci Rep* 6, 27004 -- *https://doi.org/10.1038/srep27004*

Wang, J. T., Medress, Z. A., Barres, B. A. (2012). Axon degeneration: molecular mechanisms of a self-destruction pathway. *J Cell Biol* 196, 7-18 -- *http://www.ncbi.nlm.nih.gov/pubmed/22232700*

Waterman-Storer, C. M., Salmon, E. D. (1997). Actomyosin-based retrograde flow of microtubules in the lamella of migrating epithelial cells influences microtubule dynamic instability and turnover and is associated with microtubule breakage and treadmilling. *J Cell Biol* 139, 417-34 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9334345*

Weiss, D., Langford, G., Seitz-Tutter, D., Maile, W. (1991). Analysis of the gliding, fishtailing and circling motions of native microtubules. *Acta Histochem Suppl.* 41, 81-105

Wilson, C., Gonzalez-Billault, C. (2015). Regulation of cytoskeletal dynamics by redox signaling and oxidative stress: implications for neuronal development and trafficking. *Front Cell Neurosci* 9, 381 -- *http://www.ncbi.nlm.nih.gov/pubmed/26483635*

Winckler, B., Faundez, V., Maday, S., Cai, Q., Guimas Almeida, C., Zhang, H. (2018). The endolysosomal system and proteostasis: from development to degeneration. *J Neurosci* 38, 9364-9374 -- *http://www.ncbi.nlm.nih.gov/pubmed/30381428*

Wood, J. D., Landers, J. A., Bingley, M., McDermott, C. J., Thomas-McArthur, V., Gleadall, L. J., Shaw, P. J., Cunliffe, V. T. (2006). The microtubule-severing protein Spastin is essential for axon outgrowth in the zebrafish embryo. *Hum Mol Genet* 15, 2763-71 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16893913*

Wortman, J. C., Shrestha, U. M., Barry, D. M., Garcia, M. L., Gross, S. P., Yu, C. C. (2014). Axonal transport: how high microtubule density can compensate for boundary effects in small-caliber axons. *Biophys J* 106, 813-23 -- *https://www.cell.com/biophysj/fulltext/S0006-3495(14)00075-7*

Wozniak, K. M., Vornov, J. J., Wu, Y., Liu, Y., Carozzi, V. A., Rodriguez-Menendez, V., Ballarini, E., Alberti, P., Pozzi, E., Semperboni, S., Cook, B. M., Littlefield, B. A., Nomoto, K., Condon, K., Eckley, S., DesJardins, C., Wilson, L., Jordan, M. A., Feinstein, S. C., Cavaletti, G., Polydefkis, M., Slusher, B. S. (2018). Peripheral Neuropathy Induced by Microtubule-Targeted Chemotherapies: Insights into Acute Injury and Long-term Recovery. *Cancer Res* 78, 817-829 -- *http://www.ncbi.nlm.nih.gov/pubmed/29191802*

Wu, Y., Li, J., Zhou, J., Feng, Y. (2014). Dynamic long-term microstructural and ultrastructural alterations in sensory nerves of rats of paclitaxel-induced neuropathic pain. *Chin Med J (Engl)* 127, 2945-52 -- *http://www.ncbi.nlm.nih.gov/pubmed/25131233*

Wu, Y., Whiteus, C., Xu, C. S., Hayworth, K. J., Weinberg, R. J., Hess, H. F., De Camilli, P. (2017). Contacts between the endoplasmic reticulum and other membranes in neurons. *Proc Natl Acad Sci U S A* 114, E4859-e4867 -- *https://www.pnas.org/content/114/24/E4859.short*

Xu, K., Zhong, G., Zhuang, X. (2013). Actin, Spectrin, and associated proteins form a periodic cytoskeletal structure in axons. *Science* 339, 452-6 -- *http://www.ncbi.nlm.nih.gov/pubmed/23239625*

Xu, Z., Schaedel, L., Portran, D., Aguilar, A., Gaillard, J., Marinkovich, M. P., Théry, M., Nachury, M. V. (2017). Microtubules acquire resistance from mechanical breakage through intralumenal acetylation. *Science* 356, 328-332 -- *http://science.sciencemag.org/content/sci/356/6335/328.full.pdf*

Yamasaki, H., Itakura, C., Mizutani, M. (1991). Hereditary hypotrophic axonopathy with neurofilament deficiency in a mutant strain of the Japanese quail. *Acta Neuropathol* 82, 427-34 -- *https://link-springer-com.manchester.idm.oclc.org/article/10.1007/BF00293376*

Yan, C., Wang, F., Peng, Y., Williams, C. R., Jenkins, B., Wildonger, J., Kim, H. J., Perr, J. B., Vaughan, J. C., Kern, M. E., Falvo, M. R., O'Brien, E. T., 3rd, Superfine, R., Tuthill, J. C., Xiang, Y., Rogers, S. L., Parrish, J. Z. (2018). Microtubule acetylation is required for mechanosensation in *Drosophila*. *Cell Rep* 25, 1051-1065 e6 -- *http://www.ncbi.nlm.nih.gov/pubmed/30355484*

Yap, A. S., Duszyc, K., Viasnoff, V. (2018). Mechanosensing and Mechanotransduction at Cell-Cell Junctions. *Cold Spring Harb Perspect Biol* 10 -- *http://www.ncbi.nlm.nih.gov/pubmed/28778874*

Yaron, A., Schuldiner, O. (2016). Common and divergent mechanisms in developmental neuronal remodeling and dying back neurodegeneration. *Curr Biol* 26, R628-39 -- *http://www.ncbi.nlm.nih.gov/pubmed/27404258*

Yau, K. W., van Beuningen, S. F., Cunha-Ferreira, I., Cloin, B. M., van Battum, E. Y., Will, L., Schatzle, P., Tas, R. P., van Krugten, J., Katrukha, E. A., Jiang, K., Wulf, P. S., Mikhaylova, M., Harterink, M., Pasterkamp, R. J., Akhmanova, A., Kapitein, L. C., Hoogenraad, C. C. (2014). Microtubule minus-end binding protein CAMSAP2 controls axon specification and dendrite development. *Neuron* 82, 1058-73 -- *http://www.ncbi.nlm.nih.gov/pubmed/24908486*

Yin, X., Kidd, G. J., Ohno, N., Perkins, G. A., Ellisman, M. H., Bastian, C., Brunet, S., Baltan, S., Trapp, B. D. (2016). Proteolipid protein-deficient myelin promotes axonal mitochondrial dysfunction via altered metabolic coupling. *J Cell Biol* 215, 531-542 -- *http://www.ncbi.nlm.nih.gov/pubmed/27872255*

Yu, W., Qiang, L., Solowska, J. M., Karabay, A., Korulu, S., Baas, P. W. (2008). The microtubule-severing proteins spastin and katanin participate differently in the formation of axonal branches. *Mol Biol Cell* 19, 1485-98 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18234839*

Zala, D., Hinckelmann, M.-V., Yu, H., da Cunha, L., Menezes, M., Liot, G., Cordelières, Fabrice P., Marco, S., Saudou, F. (2013). Vesicular glycolysis provides on-board energy for fast axonal transport. *Cell* 152, 479-491 -- *http://www.sciencedirect.com/science/article/pii/S0092867412015516*

Zanic, M., Widlund, P. O., Hyman, A. A., Howard, J. (2013). Synergy between XMAP215 and EB1 increases microtubule growth rates to physiological levels. *Nat Cell Biol* 15, 688–693 -- *http://dx.doi.org/10.1038/ncb2744*

Zempel, H., Mandelkow, E. M. (2015). Tau missorting and spastin-induced microtubule disruption in neurodegeneration: Alzheimer Disease and Hereditary Spastic Paraplegia. *Mol Neurodegener* 10, 68 -- *http://www.ncbi.nlm.nih.gov/pubmed/26691836*

Zhang, D., Rogers, G. C., Buster, D. W., Sharp, D. J. (2007). Three microtubule severing enzymes contribute to the "Pacman-flux" machinery that moves chromosomes. *J Cell Biol* 177, 231-42 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17452528*

Zheng, J., Lamoureux, P., Santiago, V., Dennerll, T., Buxbaum, R. E., Heidemann, S. R. (1991). Tensile regulation of axonal elongation and initiation. *J Neurosci* 11, 1117-25 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=2010807*

Zheng, Y., Wildonger, J., Ye, B., Zhang, Y., Kita, A., Younger, S. H., Zimmerman, S., Jan, L. Y., Jan, Y. N. (2008). Dynein is required for polarized dendritic transport and uniform microtubule orientation in axons. *Nat Cell Biol* 10, 1172-80 -- *http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18758451*

Ziebert, F., Mohrbach, H., Kulić, I. M. (2015). Why Microtubules Run in Circles: Mechanical Hysteresis of the Tubulin Lattice. *Physical Review Letters* 114, 148101 -- *http://link.aps.org/doi/10.1103/PhysRevLett.114.148101*

Züchner, S., Noureddine, M., Kennerson, M., Verhoeven, K., Claeys, K., De Jonghe, P., Merory, J., Oliveira, S. A., Speer, M. C., Stenger, J. E., Walizada, G., Zhu, D., Pericak-Vance, M. A., Nicholson, G., Timmerman, V., Vance, J. M. (2005). Mutations in the pleckstrin homology domain of dynamin 2 cause dominant intermediate Charcot-Marie-Tooth disease. *Nat Genet* 37, 289-94 -- *http://www.ncbi.nlm.nih.gov/pubmed/15731758*

1728   **Figures**

1729

1730

Fig. 1 Specific properties of axons. Axons during the growth cone stage are shown in (**A**) and after
synaptic maturation in (**B**), differing primarily in certain stage-specific specialisations including growth
cones, synapses, electrical properties and glial interactions (here myelination; Pan and Chan, 2017). The
core machinery in the axon shaft can be expected to be similar at both stages: parallel continuous
bundles of extended but discontinuous MTs run all along axons serving as a structural backbone (see
Fig.2), a transport highway for axonal trafficking (driven by motor proteins), and a source for 'off-track'
MTs contributing to morphogenetic processes including branch formation, directed axon growth and
synapse formation/plasticity (green, orange, blue curved arrows); MT bundles are interspersed with
longitudinal actin trails (Leterrier et al., 2017), continuous networks of (smooth) ER (Gonzalez and Couve,
2014), and other membranous organelles including mitochondria (Saxton and Hollenbeck, 2012); axonal
membranes display regularly spaced periodic rings of cortical actin (Qu et al., 2017; Xu et al., 2013), an
unusually high number of ion-specific channel proteins and transporters to conduct nerve impulses
(Kandel et al., 2012), as well as adhesions with external structures including parallel axons (not shown),
glial processes (Pronker et al., 2016) and synaptic partner cells (Koper et al., 2012); a degree of
independence from cell-body derived proteins is provided by local translation machinery (Cioni et al.,
2018; Giuditta et al., 2002b) or supply from surrounding glia cells (not shown; Court et al., 2011; Frühbeis
et al., 2013; Giuditta et al., 2002a; Rajendran et al., 2014) . Note that the axon diameter in the region
between glia cells in B (referred to as Node of Ranvier) usually has a much smaller diameter than the rest
of the axon (Hoffman, 1995).

1750

Fig. 2 Axonal response to mechanical challenges. Continuous bundles of discontinuous MTs which are flexibly cross-linked (likely involving slip-bonds) are thought to provide a structural element that can respond to different forms of mechanical impact (as indicated in blue).

1754

Fig. 3 An interactome of MT-regulating and -associated mechanisms expected to contribute within the model of local axon homeostasis. Developing and mature neurons are shown at the bottom indicating that the close-up (magenta frame) might apply in both contexts. **1**-**16**) Potential mechanisms that can 'tame' MTs into bundled conformation: MT polymerisation (blue stippled arrows) is driven by molecular machinery centred on Eb1 (blue balls), further influenced by the tubulin-supply machinery (not shown) and shaft-binding proteins (**7**); polymerisation generates new MTs required for bundle formation (**8**) and turn-over (**14**); to integrate into bundles, extending MTs require guidance via actin-Eb1 cross-linkage along the axonal surface (**5**; Shot) or along pre-existing MTs through MT-MT cross-linkers (**9**; brown L). The same or other cross-linkers provide the structural glue that holds MT bundles together (**12**; brown L); some of them can also bind to actin (**2**), they protect from (or recruit) MT severing activity (**15**), and influence motor protein dynamics (**a**). MTs which have escaped any cross-linkage are eliminated by cortical collapse factors when approaching the axonal surface (**4**; Efa6) or by MT severing factors at MT-MT cross-points (**6**). The bundled MTs are discontinuous; their free minus ends are stabilised by CAMSAP/Patronin (Ptrn) together with katanin (black scissors; **13**), whereas non-polymerising MT plus ends are stabilised by other factors (not shown; e.g. CLASP or the Dynactin subunit p150/Glued; Hur et al., 2011; Lazarus et al., 2013). The dynein/Dynactin complex is believed to link cortical actin to MT bundles and drive them anterogradely (**10**), whereas Ptrn at minus ends may anchor MTs via spectraplakins to the axon cortex (**1**); spectraplakins may also link MTs directly to cortical actin (**2**) or to transmembrane receptors (**3**), and they are expected to perform further, still unexplored actin-independent bundle-promoting roles through their PRR domains (**11**). Tear-and-wear damages MTs (dashed green line), potentially affecting interaction with MT-binding proteins (**16**; red X); MT severing proteins might selectively eliminate such MTs (**16**; scissors) or MTs undergo repair (not shown). **A**-**E**) Mechanisms closely 'associated' with MT bundles: MT-associated motor proteins ('motor', solid orange arrows) drive axonal transport of (protein-loaded) vesicles (**A**), cytoplasmic factors including proteins, translational machinery (ribosomes) or RNAs (**B**), move other MTs (**B**, sliding), and position/rearrange organelles including mitochondria (**C**, mitos), ER, peroxisomes and endosome (**D**) - and this likely

includes mitochondrial fission and fusion (**E**). **a-e**) The motor-associated functions all act downstream of MT bundles because they require them to walk on; but they also act upstream: for example, the forces they generate (stippled orange arrows) are the potential cause for MT disorganisation (buckling shown in **d**); transport delivers required regulators and building blocks for bundle-maintaining processes (**b**); the proper regulation of organelles/endocytic compartments provides systemic factors that can orchestrate taming mechanisms, including intracellular free calcium or reactive oxygen species (Ca^{2+}, ROS; yellow cloud) as well as ATP required for many processes including actin dynamics, MT severing and MT motor activity (red stippled arrows; note that vesicular transport uses glycolysis to generate its own ATP; yellow star); *vice versa*, the MT severer spastin also regulates the ER through ATP-independent mechanisms (**e**), and MT-associated proteins (APC) regulate local translation events (**c**).

Fig. 4 MT bundle defects as cause or consequence of axon decay. **1**) Disease-inducing mutations/conditions can affect an MT-bundle regulator (e.g. dystonin; Voelzmann et al., 2017), thus causing MT bundle defects first which can, in turn, trigger axon decay. **2**) Disease-inducing mutations/conditions can affect systemic factors which, in turn cause MT bundle defects as an intermediate causative step in the cascade leading to axon decay (e.g. axonal transport fails, leading to MT bundle defects which then contribute to axon decay (e.g. Alzheimer's disease or ALS; Brandt and Bakota, 2017; Farah et al., 2003; Zempel and Mandelkow, 2015); this may occur even if MT regulators are affected, but these regulators mainly act in the cell body (e.g. dysregulation of the Golgi; Ferrier et al., 2013). **3**) MT bundle deterioration may be a mere consequence of axon decay, although this case will be difficult to disentangle from option 2, since MT bundle disintegration and axonal disassembly may occur in parallel, as observed in developmental or injury-induced axon degeneration; Bradke et al., 2012; Wang et al., 2012; Yaron and Schuldiner, 2016).

Fig. 5 Disorganisation of axonal MTs upon loss of different MT regulators in *Drosophila* primary neurons.
A) Normal neuron (wild-type, wt) with soma (asterisk), axon shaft (curved arrow) and growth cone (tip of most distal MT indicated by arrow head). **B)** *Eb1⁵* mutant neuron where the area of MT disorganisation is framed by a red stippled box and shown as close-up on the right. **C-E)** Similar close-ups shown for *Efa6^GX6[w-]*, *Khc^27* and *shot³* mutant neurons. All neurons were derived from wild-type or homozygous mutant embryos, mechanically and chemically dissociated, kept for 7days in pre-culture in a centrifuge tube to deplete any maternal gene product, mechanically and chemically dissociated again, plated on concanavalin A-coated glass coverslips and grown for 1day at 21°C, fixed and stained with anti-α-tubulin (DM1A, Sigma; procedures detailed elsewhere: Prokop et al., 2012). Scale bar in A represents 10 µm for the two neurons and 4 µm in close-ups.

Fig. 6 A molecular perspective of microtubule properties. **A)** Cross-section of a MT with 14 protofilaments (PF) and lateral view of a 13 PF MT, both in B-lattice configuration, where α-tubulins make lateral bonds with α-tubulins and ß with ß, except at the seam (magenta line: seam; dashed red line: PF). **B)** Close-up of an α/ß-tubulin heterodimer showing the various post-translational modification sites as indicated; note that the GTP of ß-tubulin in lattices is usually hydrolysed (GDP). **C)** A 13 PF MT (top), cut open at the seam and rolled out (bottom); the yellow line shows the diameter, the blue line follows the helical rise of laterally bonded tubulins; in 13 PF MTs, tubulins are precisely aligned at the seam (yellow arrow head) but shifted by three positions (3-start helix). **D)** When deviating from the 13 PF prototype, tubulins are misaligned at the seam (orange arrow head); when forced into alignment, the PFs skew, causing a super-twist of the MT as described by the 'lattice accommodation model' (Chrétien and Fuller, 2000; Langford, 1980); for certain PF numbers, MTs can form two alternative alignments, of which usually the version with

the lower helix start value (left) has a left-handed super-twist, the higher value is right-handed (Chrétien
and Fuller, 2000). **E)** MTs behave like rigid rods with a persistence length of up to 10 mm, but can be bent
down to diameters of ~1µm before they break; it has been reported that their cross-sectional profile may
flatten above a certain threshold (black arrow head), thus softening the tube. **F)** Lattices of GDP-tubulin
are 1-3% shorter than MTs that were polymerised with the non-hydrolysable GTP analogue GMPCPP, or
stabilised with taxol (orange structure binding α-tubulin 1:1, according to Nogales et al., 1995); binding of
kinesin-1 causes similar lengthening of tubulin (and additional compactions in the tubulin structure: yellow
stars) which may cause cooperative binding of further kinesins and induce curvature if occurring only on
one side of the MT; in extended taxol-bound MTs, bending forces were suggested to transfer tubulins on
the concave side into their short conformation as an energetically favoured condition. For further
references see main text.

1843
1844
1845

condition	diameters of curvature [µm] [a]	comments	ref.
standard tub, 10-20 µm taxol (after?) [b] polym.	1-1.4 [c]	waves and curls upon pinning	[1]
standard tub, 50 µM taxol during & after polym.; high MT density (2.5 MTs/µm²)	1-5	loops form through collision; loop duration up frequently >5 min; strong increase in loops at high MT conc.; decreasing loop radius with increasing contour	[2]
rhodamine-tub, 10 µm taxol after polym.; exposing to air bubble or n-heptane	1.1 (heptane), 1.8 (air)	MTs become reversibly unstable in non-polar conditions: 50% of MTs form loops as long as close to air bubble; effect absolutely requires kinesins	[3]
rhodamine-tub, 10 µM taxol after polym.	2.5-3.75 [c]	left-handed supertwist favours CCW rotation of loops; CCW rotation is preserved in spools	[4]
biotin-tub, 10 µm taxol after polym.; SA-linked	1-12.6, mean 3.9		
biotin-tub, 10 µm taxol after polym.; SA-linked	1-5, mean 2.3	up to 25 µm long straight bundles; pinning of tip induces spools or fishtailing; rare "unspooling" events	[5]
biotin-tub, 10 µm taxol after polym.; SA-linked; 1600, 870, 270 and 90 kinesins/µm²	ca. 2.4-4	highest spool density & lowest spool diameter @ highest kinesin density; pinning as main cause for spool formation	[6]
biotin-GTP-tub, 10 µm taxol after polym.; SA-linked	5.7 (@ 10.8 µm length), 3 (@ 3,7 µm)	spool diameters increase with MT length per condition; spool diameters: GMP-MTs (taxol) < GMPCPP-MTs (no taxol) < GMPCPP-MTs (taxol)	[7]
biotin-GMPCPP-tub, 10 µm taxol after polym.; SA-linked	18.8 (@ 10.3 µm length), 5.8 (@ 3.4 µm)		
biotin-GMPCPP-tub, no taxol; SA-linked	8.2 (@ 10 µm length), 4.3 (@ 3.4 µm)		
biotin-GTP-tub, 10 µm taxol (after?)[b] polym.; SA-linked	3.2 µm (@ 6µm length)	live imaging: pinning & collisions (simultaneous sticking) cause spool formation; spool formation is not activated by a Brownian ratchet type process	[8]

biotin-tub, 10 μm taxol after polym.; SA-linked; microfluidic device	2.7 (pinning), 6.2 (collisions)	live imaging: pinning & collisions (simultaneous sticking) cause spools of different diameters; pinning more frequent in flow cells than microfluidic device	[9]
biotin-tub, (taxol?)[b] polym.; SA-quantum dot-linked	1.2, mean 3.4	left/right-handed super-twist: CCW/CW rotation; rings form intertwined wreath-like structures; tendency to disassemble involving MT breakage, kinesins pulling (blocked by AMP-PNP), counteracted by SA (enhanced by biotin)	[10]
biotin-tub, 10 μm taxol after polym.; SA-quantum dot-linked; patterned kinesin carpets	1-5.3 and 3.1	smallest spool diameters on constrained carpets: 1-5.3μm on 5 μm stripes, 3.1 μm on 2 μm wide squares	[11]

Tab. 2 MT loop or spool formation in gliding assays under different conditions. a) primarily the lower range of mentioned diameters is listed; b) not clear from experimental section; c) measured from images. References [1] (Amos and Amos, 1991), [2] (Liu et al., 2011), [3] (Rashedul Kabir et al., 2012), [4] (Kawamura et al., 2008), [5] (Hess et al., 2005), [6] (Lam et al., 2014), [7] (Wada et al., 2015), [8] (Luria et al., 2011), [9] (VanDelinder et al., 2016b), [10] (Liu et al., 2008), [11] (Liu and Bachand, 2013).