Mapping the steroid response to major trauma from injury to recovery: a prospective cohort study

Mark A. Foster[1,2], Angela E. Taylor[3,4], Neil E. Hill[5], Conor Bentley[1,2], Jon Bishop[1,2], Julian F. Bion[6], Joanne L.

Fallowfield[7], David R. Woods[2,8], Irina Bancos[9], Mark M Midwinter[10], Janet M Lord[1,11,12*], Wiebke Arlt[3,4,12*]

* equal senior authors

[1]NIHR-Surgical Reconstruction and Microbiology Research Centre, Queen Elizabeth Hospital Birmingham,

University Hospitals Birmingham NHS Foundation Trust, Birmingham, B15 2WB, UK; [2]Royal Centre for

Defence Medicine, Queen Elizabeth Hospital Birmingham, B15 2TH, UK; [3]Institute of Metabolism and Systems

Research, University of Birmingham, Birmingham, B15 2TT, UK; [4]Centre for Endocrinology, Diabetes and

Metabolism, Birmingham Health Partners, University of Birmingham, Birmingham, B15 2TH, UK [5]Section of

Investigative Medicine, Imperial College London, W12 0NN, UK; [6]Intensive Care Medicine, Queen Elizabeth

Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK; [7]Institute of Naval

Medicine, Alverstoke, Hants., UK; [8]Leeds Beckett University, Leeds, UK; [9]Division of Endocrinology,

Metabolism and Nutrition, Department of Internal Medicine, Mayo Clinic, Rochester, Minnesota, USA;

[10]School of Biomedical Sciences, University of Queensland, Brisbane, Australia; [11]MRC-ARUK Centre for

Musculoskeletal Ageing Research, Institute of Inflammation and Ageing, University of Birmingham,

Birmingham, B15 2TT, UK; [12]NIHR Birmingham Biomedical Research Centre, University Hospitals

Birmingham NHS Foundation Trust and University of Birmingham, Birmingham, B15 2GW, UK

Corresponding author:

Lt Col Mark Anthony Foster, Academic Department of Military Surgery and Trauma, Royal Centre for Defence

Medicine, ICT Centre, Birmingham, B15 2SQ

m.foster@bham.ac.uk Tel: +47970 444328

Word Count: 2998

Short title: The steroid response to major trauma

Keywords: major trauma; systemic inflammatory response syndrome; stress response; steroids; DHEA;

testosterone**.**

Funding: The SIR Study was part of the Surgeon General's Casualty Nutrition Study (SGCNS), supported by

University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham. Additional funding

was provided by the Drummond Trust Foundation. WA and JML receive support from the National Institute for

Health Research (NIHR) Birmingham Biomedical Research Centre (Grant Reference Number BRC-1215-

20009).

Disclosure statement: The authors have nothing to disclose. The views expressed are those of the authors and

not necessarily those of the NIHR or the Department of Health and Social Care UK.

Abstract

Context: Survival rates after severe injury are improving, but complication rates and outcomes are variable.

Objective: This cohort study addressed the lack of longitudinal data on the steroid response to major trauma and during recovery.

Design: We undertook a prospective, observational cohort study from time of injury to six months post-injury at a major UK trauma center and a military rehabilitation unit, studying patients within 24 hours of major trauma (estimated New Injury Severity Score (NISS) >15).

Main outcome measures: We measured adrenal and gonadal steroids by tandem mass spectrometry, assessed muscle loss by ultrasound and nitrogen excretion, and recorded clinical outcomes (ventilator days, length of hospital stay, incidence of organ dysfunction and sepsis); results were analyzed by generalized mixed-effect linear models.

Findings: We screened 996 multiply injured adults, approached 106, and recruited 95 eligible patients; 87 survived. We analyzed all male survivors <50 years not treated with steroids (N=60; median age 27 [interquartile range 24-31] years; median NISS 34 [29-44]). Urinary nitrogen excretion and muscle loss peaked after one and six weeks, respectively. Glucocorticoid secretion remained within the reference range; the serum cortisol over cortisone ratio increased, normalizing eight weeks post-injury. Serum testosterone, DHEA and DHEAS decreased immediately after trauma and took two, four and more than six months, respectively, to recover; all three correlated with SOFA score and probability of sepsis.

Conclusion: The catabolic response to severe injury is accompanied by acute and sustained androgen suppression. Whether androgen supplementation improves health outcomes after major trauma should be investigated.

Précis

A cohort study in male survivors of major trauma revealed acute and sustained androgen suppression and protein catabolism including muscle loss. Serum androgens correlated with probability of sepsis.

INTRODUCTION

Over 5 million people worldwide die each year from serious injury(1), with almost 25% caused by road traffic collisions (RTC)(2). In England alone, there are 5400 trauma deaths and 20,000 severe injuries treated by the National Health Service annually(3). Since 2012, the establishment of 22 trauma centres in England has been accompanied by a 19% improvement in survival odds following injury(4). During this time, the UK also received severely injured military trauma patients from the conflict in Afghanistan(5,6).

Improvements in short-term outcomes have been achieved through early resuscitation and acute care(7), often informed by approaches pioneered on the battlefield. However, the improvement in survival is often offset during the weeks following acute major trauma, by the systemic inflammatory response syndrome (SIRS), associated with increased risks of infection, multi-organ dysfunction or failure (MOD/MOF), and death(8,9). Simultaneously, the hypothalamic-pituitary-adrenal axis (HPA) is thought to drive a hypermetabolic and overtly catabolic response. Pro-inflammatory cytokines activate the enzyme 11β-hydroxysteroid dehydrogenase type (11β-HSD1) responsible for tissue-specific activation of glucocorticoids through conversion of inactive cortisone to active cortisol(10), providing a link between SIRS and catabolism. Importantly, in this profound catabolic state, patients lose valuable lean muscle and suffer from increased rates of infection and poor wound healing. Nutritional strategies have proven disappointing (11,12). Moreover, the dynamic nature of this response, in particular beyond the first few days after injury and during recovery remains poorly described and understood, limiting the evidence base for novel therapeutic interventions.

To address this issue, we have undertaken a detailed prospective study of the endocrine and metabolic response to severe injury in military and civilian populations, recruiting patients within 24 hours of major trauma and following up on them for the six subsequent months. This was undertaken to identify predictive biomarkers and therapeutic targets as well as to explore the optimal timing for therapeutic interventions that could promote better recovery after severe traumatic injury.

METHODS

Study Design and Protocol

This prospective cohort study was conducted in the Royal Centre for Defense Medicine and the Queen Elizabeth Hospital Birmingham, a major UK trauma center and the primary receiving facility for UK military personnel injured abroad. Military and civilian trauma patients with an estimated New Injury Severity Score (NISS) >15 were recruited[13]. NISS was used to ensure those with significant extremity trauma but lower ISS were included[14]. Patients with significant head injury or pre-injury neoplastic conditions were not eligible. Informed consent was obtained from personal consultees until recovering capacity. The protocol was approved by the NRES Committee South West – Frenchay 11/SW/0177 and MOD REC 116/Gen/10.

A daily patient review allowed the injured to be clinically phenotyped. Bespoke study management software (Clinical RESearch Tool – CREST) tracked the patient and their clinical data were entered prospectively and used to calculate Acute Physiology and Chronic Health Evaluation II (APACHE II), Sequential Organ Failure Assessment (SOFA), and Simplified Acute Physiology Score (SAPS)[15,16]. Sepsis was defined using Bone's criteria of an infection associated with SIRS[17], current at the time of the study.

Blood and urine samples were collected within 24 hours of acute injury and at 3, 5, 10, 14, 21, 28 days and 2, 3, 4 and 6 months during recovery from major trauma. This challenging sampling schedule was supported by junior clinical staff to ensure blood sampling between 0730 and 0900hrs; serum was then separated and frozen at -80^{o}C for batched analysis. We separately collected morning blood samples from 37 healthy age- and sex-matched controls, to provide a comparator for the steroid data.

Assessment of protein catabolism

As a surrogate marker for muscle mass, we undertook longitudinal measurements of muscle thickness by a well validated method using portable ultrasound as described by Campbell *et al*[18,19]. Ultrasound measurements were taken from four different muscle sites; biceps brachii, radial forearm, rectus femoris and rectus abdominis at weekly intervals while in hospital and at 3, 4, 5 and 6 months following discharge. All ultrasound assessments were performed by two designated operators. Measurements were performed three times at each muscle site and the mean of the three measurements was recorded. The dominant arm was favored for

ultrasound assessment of muscle mass unless it was missing or unable to be measured where wounds were extensive. This non-invasive method was chosen over others such as creatine (methyl-d3) dilution (D3-creatine) (20) due to the strict dietary requirements for methy-d3 estimation, which was not practical in the context of major trauma; similarly, we did not undertake MRI measurement of muscle mass due to the risk associated with repeated transport of critically ill patients to scanning facilities (21).

Urinary urea excretion was measured and used to estimate Total Urinary Nitrogen (TUN) excretion as described by Milner et al(22) [Estimated Nitrogen Excretion: Urinary Urea Excretion (mmol/l) x 0.028 x 1.25 = Total Urinary Nitrogen excretion (g/l)].

Serum steroid analysis

Serum concentrations of adrenal and gonadal steroids were measured using liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis, employing a validated multi-steroid profiling method(23). In brief, serum steroids were extracted via liquid/liquid tert-butyl-methyl-ether (MTBE), evaporated, reconstituted, and analyzed by LC-MS/MS for cortisol and cortisone. Serum androgens and androgen precursors (DHEA, androstenedione, testosterone) were measured following oxime derivatisation(24,25). DHEA sulfate (DHEAS) was measured following protein precipitation(26,27).

Serum concentrations of sex hormone-binding globulin (SHBG) and luteinizing hormone (LH) were analyzed on the Roche Modular System (Roche Diagnostics, Lewes, UK) by two-site sandwich immunoassay using electrochemiluminesence technology.

Statistical Analysis

The raw data were evaluated by analysis of variance (ANOVA); in addition, paired and unpaired Student's t-test, Chi-Square Analysis and Mann-Whitney tests were used where appropriate.

Generalized linear mixed-effects models(28) were used to examine the change in variables over time. Patients were included in models as random effects to account for repeat measures over time on the same individuals. Time was modelled using restricted cubic splines(29) to allow for flexible relationships(30). Severity scores were modelled as Poisson distributions due to their skewness and non-negative ranges. Plots of

predicted average fixed effects with 95% confidence intervals were produced for the first four weeks and first six

months after injury as required. Analyses were conducted in R using libraries lme4, effects, rms and ggplot2.

Power calculation based on previous work[36] indicated a minimum sample size of 25 patients accommodate

variability in steroid analysis.

RESULTS

Patient recruitment and clinical characteristics of the final study cohort

We screened 996 multiply injured adults. The majority of the 889 excluded patients had a NISS≤15;

others had a significant head injury as their major injury component, and two were excluded due to a pre-injury

diagnosis of cancer. Of the 102 patients recruited into the study, two withdrew and re-assessment in five

revealed an actual NISS ≤15, leaving a study cohort of 95 patients (**Fig. 1A**).

Excluding eight fatalities and seven patients that had received steroid therapy, 80 survivors completed

sample collection over 6 months. To minimize confounders, we excluded the small groups of women (n=9) and

age-advanced men (n=11), leaving our final study cohort of 60 men <50 years of age (**Fig. 1B**).

A summary of the cohort characteristics is shown in **Fig. 2**. Median age was 27 (interquartile range

(IQR) 24-31) years, median NISS was 34 (IQR 29-44), and patient day 1 (=day of major trauma) APACHE II

score was 21 (IQR 14-25). Patients remained ventilated on the intensive care unit (ICU) for a median of 9 (IQR

5-16) days. Median length of hospital stay was 36 (IQR 19-56) days. Improvised Explosive Device (IED)

(n=33; 55%) and Road Traffic Collision (RTC) (n=11; 18%) were the most common causes of injury. Twenty-

five (42%) patients had at least one septic episode and most occurred in the second week (**Fig 2C**).

Glucocorticoid biosynthesis and metabolism after major trauma

Serum cortisol concentrations increased slightly after injury, peaking at 408 (IQR 249-511) nmol/L at

two weeks (**Fig. 3A; Suppl. Fig. 2**). However, concentrations remained within the wide range observed in

healthy controls. Serum concentrations of the inactive glucocorticoid metabolite cortisone were lower than

normal after injury, and increased slowly over time, but this trend was not significant (p=0·08) (**Fig. 3B**). The

serum cortisol-to-cortisone ratio, a marker of 11β-HSD activity (**Fig. 3C**), peaked at two weeks post-injury and returned to normal at around eight weeks.

Androgen biosynthesis and activation after major trauma

Serum concentrations of adrenal androgen precursor dehydroepiandrosterone (DHEA) were very low after injury (p<0·0001, as compared to healthy controls), but recovered to the normal range by three months post-injury (**Fig. 3D, Suppl. Fig. 1**). In contrast, its sulfate ester, DHEAS, demonstrated sustained suppression and median concentrations did not recover to values within the healthy reference range even at the end of the 6-month study period (**Fig. 3E**). Consequently, the serum DHEA-to-DHEAS ratio (**Fig. 3F**) increased by week 2 compared to controls and failed to return to normal during the entire study period. The serum cortisol-to-DHEAS ratio (**Fig. 3G**) increased after injury, peaking at 2 weeks, followed by a gradual decrease, but without returning to normal by the end of the 6-month study period.

Serum concentrations of the androgen precursor androstenedione (**Fig. 3H**) were below the reference range immediately after injury, recovering to the mid reference range at 2 weeks after injury, therefore, recovering much faster than serum DHEA, suggestive of rapid downstream activation of DHEA to androstenedione.

Serum testosterone (**Fig. 3I**) was very low following injury, starting to increase after two weeks, and recovering to the healthy sex- and age-matched reference range approximately eight weeks after injury. This was mirrored by acute suppression of serum LH immediately after injury, followed by recovery to the normal range approximately 2 weeks after injury (**Suppl. Fig. 2C+D**). Serum sex hormone-binding globulin (SHBG) (**Suppl. Fig. 2E+F**) concentrations were subnormal immediately post-injury but quickly returned to the healthy reference range between injury and day 7.

Protein catabolism after major trauma

The 24-hour total urinary nitrogen (TUN) excretion increased immediately after trauma, peaking at 25·0±16·1 g/day at the end of the first week, returning to below 15·0 g/day by week 4. The mean maximum rate of nitrogen excretion was 33·0±21·3 g/day (**Fig. 4A**). The normalization of TUN excretion coincided with the gradual recovery of adrenal and gonadal androgen production (**Figs. 4B+C**).

The biceps brachii muscle was the most reliable site for ultrasound measurement of muscle thickness; dressings, amputations and other wounds hampered the measurements in the other muscle areas. Changes in biceps brachii muscle thickness followed a U-shaped curve after injury, reaching a nadir at 6 weeks (day 1 after trauma compared with week 6, p=0·001). The mean muscle loss was 22·7±12·5% (**Fig. 4D**). Similar to TUN, muscle thickness recovered alongside gradually increasing adrenal and gonadal androgen production (**Figs. 4E+F**).

Clinical course of post-traumatic recovery and serum androgen dynamics

The relationship between adrenal and gonadal androgens and the Sequential Organ Failure Score (SOFA) and probability of sepsis are illustrated in **Fig. 5**. During the first four weeks, serum DHEA, DHEAS, and testosterone all correlated with the clinical SOFA score (autocorrelation factor (ACF) = 0·85, 0·90 and -0·79, respectively). The serum concentrations of all three steroids also showed strong associations with the probability of sepsis (R=-0·85, 0·85 and -0·97 for serum DHEA, DHEAS and testosterone, respectively). SOFA score and probability of sepsis also correlated strongly with the DHEA:DHEAS ratio (autocorrelation factor (ACF) = -0·94 and -0·96 respectively) and with the serum cortisol-to-DHEAS ratio, negatively for the SOFA score but positively for probability of sepsis (autocorrelation factor (ACF) = -0·81 and 0·89, respectively) (**Suppl. Fig 3**).

DISCUSSION

In this study, we have characterized the response of adrenal and gonadal steroids and catabolic metabolism to severe injury, describing the related dynamic changes for six months post-injury. Modelling the resulting data has allowed us to provide a detailed description of the transition from catabolism to anabolism during recovery from severe injury. Our data are the first to provide detailed adrenal and gonadal steroid data beyond the first days after trauma in a large cohort of patients, with all patients recruited prospectively and steroid analysis carried out by tandem mass spectrometry.

Our study revealed an initial phase of minor glucocorticoid activation with a transient increase in the serum cortisol-to-cortisone ratio. As summarized in a recent meta-analysis, previous data on serum cortisol after injury is limited to small cohorts derived from elective surgery, rarely followed up for more than two days(31).

In our study, serum cortisol quickly returned to normal following slight initial increases after acute trauma. In contrast, serum cortisone remained low for three months post-injury, resulting in a significant increase in the cortisol to cortisone ratio, an *in vivo* measure of the activity of 11β-hydroxysteroid dehydrogenase (11β-HSD) activity. The cortisol-activating enzyme 11β-HSD1 is the major enzyme converting inactive cortisone to cortisol and has been shown to be upregulated systemically and locally in response to inflammation, thereby dampening the inflammatory response(32,33). Skeletal muscle expresses 11β-HSD1(34) and previous studies reported increased 11β-HSD1 activity in an animal model of trauma haemorrhage(35) and recently improved wound healing in mice treated with 11β-HSD1 inhibitors(36); human data after trauma are lacking. Mirroring this, there is substantial evidence indicating a reduced cortisol clearance in critical illness, due to decreased cortisol inactivation in liver and kidney (37); this mechanism could also be responsible for the slight changes in cortisol and cortisone we observed.

In our study, we observed a pronounced and sustained loss of adrenal and gonadal androgen synthesis immediately following acute major trauma. The recovery of circulating DHEA and testosterone concentrations took two and four months post-injury, respectively, and DHEAS remained pathologically decreased at the end of the six-month follow-up period. In a mouse model of acute inflammation, sustained suppression of the expression of the DHEA sulfotransferase SULT2A1 and its sulfate donor enzyme, PAPSS2, have been described(60). We reviewed 23 previous studies that measured serum DHEA and DHEA(**Suppl. Table 1**) patients (**Suppl. Table 1**), but most followed patients only for a few days and relatively few patients suffered from acute trauma. One previous study measured both serum DHEA and DHEAS in 181 patients with septic shock and 31 patients with acute hip fracture(48); serum DHEAS was decreased in both groups, while DHEA was increased in the sepsis patients but decreased in the trauma patients, suggesting an inflammation-mediated downregulation of DHEA sulfation in the latter group, resulting in a dissociation between serum DHEA and DHEAS concentrations.

A number of previous studies have described an association of infection and mortality with low circulating DHEAS concentrations and a raised serum cortisol-to-DHEAS ratio in patients with trauma(42,43,61,62). In vitro studies have demonstrated that cortisol decreases neutrophil superoxide

production, which is counteracted by coincubation with DHEAS(42). Furthermore, we have previously shown

that DHEAS, but not DHEA, directly enhances neutrophil superoxide generation, a key mechanism of human

bactericidal function, via activation of protein kinase C-β, independent of androgen receptor signalling(63). In

this study in severely injured men younger than 50 years of age, we observed suppression of both serum DHEA

and DHEAS post-injury, indicating that the loss of adrenal androgen synthesis is a trauma-related event.

Importantly, we show for the first time that this decrease in circulating adrenal androgen precursors is sustained

for several months and that DHEAS remains low even six months post-injury.

Alongside the decrease in adrenal androgen synthesis, we observed a near complete loss of gonadal

testosterone production and pituitary LH secretion immediately after trauma. Both the gradual recovery of

adrenal and gonadal androgen production paralleled the decrease in catabolism, as assessed by urinary nitrogen

excretion and biceps muscle thickness. The suppression of the hypothalamus-pituitary-gonadal (HPG) axis after

severe injury shown in our study is supported by the literature(56,57,59,64). Our longitudinal data demonstrate

that suppression of the HPG axis is of shorter duration than that of the HPA axis. In traumatic brain injury

studies, a significant proportion of patients go on to develop anterior pituitary dysfunction including secondary

hypogonadiasm(65), (50,51), however, in our study traumatic brain injury was an exclusion criterion. While

limited data from patients with burns and critical illness have suggested a central, hypothalamic-pituitary cause

of trauma-related hypogonadism(66,67), the evidence prior to our study has been limited. Out data indicate a

central cause of suppression to the gonadotrophic axis, with a decrease in both pituitary LH and gonadal

testosterone. However, it cannot be excluded that the administration of opiates could causally contribute to the

induction of central hypogonadism(68).

Our cohort study reveals a loss of both adrenal and gonadal androgen production in young and middle-

aged men after major trauma. Androgens are important in wound healing, erythrocytosis, bone density and

muscle mass(69). The catabolic state that occurs following trauma thus presents a significant challenge. The

use of androgens to ameliorate catabolism has some precedent, as evidenced by the use of the synthetic

androgen, oxandrolone, that has some proven benefit in treating burn injury(70). A meta-analysis of 15 RCTs

including 806 burns patients by Li et al, showed significant benefits (P < 0.05) for using oxandrolone including

less net weight loss, lean body mass loss, nitrogen loss, donor-site healing time, and length of stay in the

catabolic and rehabilitative phases(71). The use of oxandrolone in major trauma was investigated in two

intensive care studies but no benefit was demonstrated(72,73).

The strengths of our study include its prospective nature, narrow age range of the patients, single gender,

single site for recruitment and analysis and detailed follow-up over six months as well as the measurement of

circulating steroid hormones by state-of-the-art tandem mass spectrometry. Analyzing a young cohort has also

reduced the confounding effects of age-related co-morbidities.

Our study is limited by the diverse nature of major trauma patients in relation to injury pattern and the

involvement of military casualties. The timing and number of observations during our study was pragmatic and

some statistical comparisons were made using modelled data. While we measured total cortisol and cortisone by

tandem mass spectrometry, we did not measure free cortisol or cortisol-binding globulin. The estimation of

nitrogen excretion was pragmatic due to the diverse nature of the patients and we were not able to record

nitrogen intake. Ultrasound estimation of muscle thickness was performed at four different body sites but many

individuals had limbs missing or extensive wounds that prevented measurements. While imperfect, the

longitudinal nature of these measurements allowed us to model these changes over time.

In conclusion, in this most detailed and first prospective study of the steroid response to major trauma,

we followed the patients from severe injury to six months of recovery, revealing pronounced and sustained

decreases in adrenal and gonadal androgen biosynthesis. Recovery of androgen production in the severely

injured patients was mirrored by a switch from catabolism to anabolism as reflected by recovery of muscle mass

and a decrease in nitrogen loss. Adrenal and gonadal androgens correlated with risk of sepsis. It is tempting to

suggest that an anabolic intervention with androgens or androgen precursors could have a beneficial effect on

health outcomes during recovery from major trauma. However, this will need to be investigated by future

intervention studies.

Acknowledgements

The SIR Study was part of the Surgeon General's Casualty Nutrition Study (SGCNS), supported by University Hospitals Birmingham NHS Foundation Trust (UHB) and the University of Birmingham. Additional funding was provided by the Drummond Trust Foundation. WA and JM receive support from the National Institute for Health Research (NIHR) Birmingham Biomedical Research Centre (Grant Reference Number BRC-1215-20009). The views expressed are those of the authors not necessarily those of the NIHR or the Department of Health and Social Care UK.

We thank staff from the NIHR Surgical Reconstruction and Microbiology Centre, Leah Duffy, Lauren Cooper, Peter Ip and Aisling Crombie; military nurses Michelle Taylor and Matt Daley; junior doctors from UHB, Abigail Routledge, Ben Booth, Rob Staruch and Researchers and staff from University of Birmingham, Hema Chahal, Donna M. O'Neil, Jon Hazeldine and Pete Hampson; Institute of Naval Medicine, Dr Adrian Allsopp, Sophie Britland, Dr Pieter Brown, Roz Cobley, Simon Delves, Anneliese Shaw, Dr Fran Gunner, Jenny Hayward-Karlsson; Defence Rehabilitation Centre Headley Court, Jakob Kristensen, Wg Cdr Alex Bennett; University of Surrey, Prof Susan Lanham-New; Imperial University, Prof Stephen Brett; Dr Kevin Murphy, Prof Gary Frost; members of SGCNS not already mentioned, Surg Cdr Jane Risdall, Andy Roberts, Lt Col Sandra Williams, and Col Duncan Wilson.

References

1.　**World Health Organization.** *Injuries and violence: the facts 2014*. 2014.

2.　**OECD I.** Road Safety Annual Report. *International Transport Forum* 2014.

3.　**Office GBNA.** *Major Trauma Care in England*. The Stationery Office; 2010.

4.　**Moran CG, Lecky F, Bouamra O, Lawrence T, Edwards A, Woodford M, Willett K, Coats TJ.**
Changing the System - Major Trauma Patients and Their Outcomes in the NHS (England) 2008-17.
EClinicalMedicine 2018;2:13–21. doi:10.1016/j.eclinm.2018.07.001.

5.　**Ministry of Defence.** Operations in Afghanistan. *https://www.gov.uk/guidance/uk-forces-operations-in-*
afghanistan/casualty-figures 2015:1–69.

6.　**Ministry of Defence.** UK forces: operations in Afghanistan. *https://www.gov.uk/guidance/uk-forces-*
operations-in-afghanistan/casualty-figures 2014:1–27 (Accessed 5 Jan 16.

7.　**Hodgetts TJ, Mahoney PF.** Military pre-hospital care: why is it different? *J R Army Med Corps*
2009;155(1):4–8.

8.　**Durham RM, Moran JJ, Mazuski JE, Shapiro MJ, Baue AE, Flint LM.** Multiple organ failure in
trauma patients. *J Trauma* 2003;55(4):608–616. doi:10.1097/01.TA.0000092378.10660.D1.

9.　**Lenz A, Franklin GA, Cheadle WG.** Systemic inflammation after trauma. *Injury* 2007;38(12):1336–
1345. doi:10.1016/j.injury.2007.10.003.

10.　**Esteves CL, Verma M, Róg-Zielińska E, Kelly V, Sai S, Breton A, Donadeu FX, Seckl JR, Chapman**
KE. Pro-inflammatory cytokine induction of 11β-hydroxysteroid dehydrogenase type 1 in A549 cells
requires phosphorylation of C/EBPβ at Thr235. *PLoS ONE* 2013;8(9):e75874.
doi:10.1371/journal.pone.0075874.

11.　**Arabi YM, Casaer MP, Chapman M, Heyland DK, Ichai C, Marik PE, Martindale RG, McClave**
SA, Preiser J-C, Reignier J, Rice TW, Van den Berghe G, van Zanten ARH, Weijs PJM. The
intensive care medicine research agenda in nutrition and metabolism. *Intensive Care Med*
2017;43(9):1239–1256. doi:10.1007/s00134-017-4711-6.

12.　**Reignier J, Boisramé-Helms J, Brisard L, Lascarrou J-B, Ait Hssain A, Anguel N, Argaud L,**
Asehnoune K, Asfar P, Bellec F, Botoc V, Bretagnol A, Bui H-N, Canet E, Da Silva D, Darmon M,
Das V, Devaquet J, Djibre M, Ganster F, Garrouste-Orgeas M, Gaudry S, Gontier O, Guérin C,
Guidet B, Guitton C, Herbrecht J-E, Lacherade J-C, Letocart P, Martino F, Maxime V, Mercier E,
Mira J-P, Nseir S, Piton G, Quenot J-P, Richecoeur J, Rigaud J-P, Robert R, Rolin N, Schwebel C,
Sirodot M, Tinturier F, Thévenin D, Giraudeau B, Le Gouge A, NUTRIREA-2 Trial Investigators,
Clinical Research in Intensive Care and Sepsis (CRICS) group. Enteral versus parenteral early
nutrition in ventilated adults with shock: a randomised, controlled, multicentre, open-label, parallel-group
study (NUTRIREA-2). *Lancet* 2018;391(10116):133–143. doi:10.1016/S0140-6736(17)32146-3.

13.　**Palmer C.** Major trauma and the injury severity score--where should we set the bar? *Annu Proc Assoc*
Adv Automot Med 2007;51:13–29.

14.　**Balogh ZJ, Varga E, Tomka J, Süveges G, Tóth L, Simonka JA.** The new injury severity score is a
better predictor of extended hospitalization and intensive care unit admission than the injury severity score
in patients with multiple orthopaedic injuries. *J Orthop Trauma* 2003;17(7):508–512.

15. **Le Gall J-R, Lemeshow S, Saulnier F.** A New Simplified Acute Physiology Score (SAPS II) Based on a
European/North American Multicenter Study. *JAMA* 1993;270(24):2957–2963.
doi:10.1001/jama.1993.03510240069035.

16. **Knaus WA, Draper EA, Wagner DP, Zimmerman JE.** APACHE II: a severity of disease classification
system. *Crit Care Med* 1985;13(10):818–829.

17. **Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, Schein RM, Sibbald WJ.**
Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The
ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of
Critical Care Medicine. In: Vol 101. 1992:1644–1655.

18. **Campbell IT, Watt T, Withers D, England R, Sukumar S, Keegan MA, Faragher B, Martin DF.**
Muscle thickness, measured with ultrasound, may be an indicator of lean tissue wasting in multiple organ
failure in the presence of edema. *Am J Clin Nutr* 1995;62(3):533–539.

19. **Reid CL, Campbell IT, Little RA.** Muscle wasting and energy balance in critical illness. *Clin Nutr*
2004;23(2):273–280. doi:10.1016/S0261-5614(03)00129-8.

20. **Clark RV, Walker AC, Miller RR, O'Connor-Semmes RL, Ravussin E, Cefalu WT.** Creatine (
methyl-d3) dilution in urine for estimation of total body skeletal muscle mass: accuracy and variability vs.
MRI and DXA. *Journal of Applied Physiology* 2018;124(1):1–9. doi:10.1152/japplphysiol.00455.2016.

21. **Buckinx F, Landi F, Cesari M, Fielding RA, Visser M, Engelke K, Maggi S, Dennison E, Al-Daghri
NM, Allepaerts S, Bauer J, Bautmans I, Brandi ML, Bruyère O, Cederholm T, Cerreta F,
Cherubini A, Cooper C, Cruz-Jentoft A, McCloskey E, Dawson-Hughes B, Kaufman J-M, Laslop A,
Petermans J, Reginster J-Y, Rizzoli R, Robinson S, Rolland Y, Rueda R, Vellas B, Kanis JA.** Pitfalls
in the measurement of muscle mass: a need for a reference standard. *J Cachexia Sarcopenia Muscle*
2018;9(2):269–278. doi:10.1002/jcsm.12268.

22. **Milner EA, Cioffi WG, Mason AD, McManus WF, Pruitt BA.** Accuracy of urinary urea nitrogen for
predicting total urinary nitrogen in thermally injured patients. *JPEN J Parenter Enteral Nutr*
1993;17(5):414–416. doi:10.1177/0148607193017005414.

23. **Ionita IA, Fast DM, Akhlaghi F.** Development of a sensitive and selective method for the quantitative
analysis of cortisol, cortisone, prednisolone and prednisone in human plasma. *J. Chromatogr. B Analyt.*
Technol. Biomed. Life Sci. 2009;877(8-9):765–772. doi:10.1016/j.jchromb.2009.02.019.

24. **Kushnir MM, Blamires T, Rockwood AL, Roberts WL, Yue B, Erdogan E, Bunker AM, Meikle
AW.** Liquid chromatography-tandem mass spectrometry assay for androstenedione,
dehydroepiandrosterone, and testosterone with pediatric and adult reference intervals. *Clinical Chemistry*
2010;56(7):1138–1147. doi:10.1373/clinchem.2010.143222.

25. **O'Reilly MW, Taylor AE, Crabtree NJ, Hughes BA, Capper F, Crowley RK, Stewart PM,
Tomlinson JW, Arlt W.** Hyperandrogenemia predicts metabolic phenotype in polycystic ovary
syndrome: the utility of serum androstenedione. *J Clin Endocrinol Metab* 2014;99(3):1027–1036.
doi:10.1210/jc.2013-3399.

26. **Chadwick CA, Owen LJ, Keevil BG.** Development of a method for the measurement of
dehydroepiandrosterone sulphate by liquid chromatography-tandem mass spectrometry. *Ann. Clin.*
Biochem. 2005;42(Pt 6):468–474. doi:10.1258/000456305774538175.

27. **Haring R, Wallaschofski H, Teumer A, Kroemer H, Taylor AE, Shackleton CHL, Nauck M, Volker**
U, Homuth G, Arlt W. A SULT2A1 genetic variant identified by GWAS as associated with low serum
DHEAS does not impact on the actual DHEA/DHEAS ratio. *Journal of Molecular Endocrinology*
2012;50(1):73–77. doi:10.1530/JME-12-0185.

28. **Breslow NE, Clayton DG.** Approximate Inference in Generalized Linear Mixed Models. *Journal of the*
American Statistical Association 2012;88(421):9–25. doi:10.1080/01621459.1993.10594284.

29. **Meyer MC.** Inference using shape-restricted regression splines. *Ann. Appl. Stat.* 2008;2(3):1013–1033.
doi:10.1214/08-AOAS167.

30. **Harrell FE, Lee KL, Mark DB.** Multivariable prognostic models: issues in developing models,
evaluating assumptions and adequacy, and measuring and reducing errors. *Stat Med* 1996;15(4):361–387.
doi:10.1002/(SICI)1097-0258(19960229)15:4<361::AID-SIM168>3.0.CO;2-4.

31. **Prete A, Yan Q, Al-Tarrah K, Akturk HK, Prokop LJ, Alahdab F, Foster MA, Lord JM,**
Karavitaki N, Wass JA, Murad MH, Arlt W, Bancos I. The cortisol stress response induced by
surgery: A systematic review and meta-analysis. *Clin Endocrinol* 2018;89(5):554–567.
doi:10.1111/cen.13820.

32. **Chapman KE, Coutinho A, Gray M, Gilmour JS, Savill JS, Seckl JR.** Local amplification of
glucocorticoids by 11beta-hydroxysteroid dehydrogenase type 1 and its role in the inflammatory response.
Ann NY Acad Sci 2006;1088(1):265–273. doi:10.1196/annals.1366.030.

33. **Ray JA, Meikle AW.** Cortisol-Cortisone Shuttle: A Functional Indicator of 11beta-HSD Activity. In:
Cortisol Physiology, Regulation and Health Implications. 2012:91–106.

34. **Hassan-Smith ZK, Morgan SA, Sherlock M, Hughes B, Taylor AE, Lavery GG, Tomlinson JW,**
Stewart PM. Gender-Specific Differences in Skeletal Muscle 11β-HSD1 Expression Across Healthy
Aging. *J Clin Endocrinol Metab* 2015;100(7):2673–2681. doi:10.1210/jc.2015-1516.

35. **Wang P, Ba ZF, Jarrar D, Cioffi WG, Bland KI, Chaudry IH.** Mechanism of adrenal insufficiency
following trauma and severe hemorrhage: role of hepatic 11beta-hydroxysteroid dehydrogenase. *Arch*
Surg 1999;134(4):394–401.

36. **Tiganescu A, Tahrani AA, Morgan SA, Otranto M, Desmoulière A, Abrahams L, Hassan-Smith Z,**
Walker EA, Rabbitt EH, Cooper MS, Amrein K, Lavery GG, Stewart PM. 11β-Hydroxysteroid
dehydrogenase blockade prevents age-induced skin structure and function defects. *The Journal of Clinical*
Investigation 2013;123(7):3051–3060. doi:10.1172/JCI64162.

37. **Boonen E, Vervenne H, Meersseman P, Andrew R, Mortier L, Declercq PE, Vanwijngaerden Y-M,**
Spriet I, Wouters PJ, Vander Perre S, Langouche L, Vanhorebeek I, Walker BR, Van den Berghe
G. Reduced cortisol metabolism during critical illness. *New England Journal of Medicine*
2013;368(16):1477–1488. doi:10.1056/NEJMoa1214969.

38. **Kolditz M, Halank M, Schulte-Hubbert B, Hoffken G.** Adrenal function is related to prognosis in
moderate community-acquired pneumonia. *Eur. Respir. J.* 2010;36(3):615–621.
doi:10.1183/09031936.00191709.

39. **Sharshar T, Bastuji-Garin S, De Jonghe B, Stevens RD, Polito A, Maxime V, Rodriguez P, Cerf C,**
Outin H, Touraine P, Laborde K. Hormonal status and ICU-acquired paresis in critically ill patients.
Intensive Care Med 2010;36(8):1318–1326. doi:10.1007/s00134-010-1840-6.

40. **Klouche K, Da Mota EF, Durant R, Amigues L, Corne P, Jonquet O, Beraud JJ.** Hypothalamic-
pituitary-adrenal axis reactivity and dehydroepiandrosterone sulfate plasma concentrations in the critically
ill elderly. *Age Ageing* 2007;36(6):686–689. doi:10.1093/ageing/afm113.

41. **Chinga-Alayo E, Villena J, Evans AT, Zimic M.** Thyroid hormone levels improve the prediction of
mortality among patients admitted to the intensive care unit. *Intensive Care Med* 2005;31(10):1356–1361.
doi:10.1007/s00134-005-2719-9.

42. **Butcher SK, Killampalli V, Lascelles D, Wang K, Alpar EK, Lord JM.** Raised cortisol:DHEAS ratios
in the elderly after injury: potential impact upon neutrophil function and immunity. *Aging Cell*
2005;4(6):319–324. doi:10.1111/j.1474-9726.2005.00178.x.

43. **Beishuizen A, Thijs LG, Vermes I.** Decreased levels of dehydroepiandrosterone sulphate in severe
critical illness: a sign of exhausted adrenal reserve? *Crit Care* 2002;6(5):434–438.

44. **Sharshar T, Bastuji-Garin S, Polito A, De Jonghe B, Stevens RD, Maxime V, Rodriguez P, Cerf C,
Outin H, Touraine P, Laborde K.** Hormonal status in protracted critical illness and in-hospital mortality.
Crit Care 2011;15(1):R47. doi:10.1186/cc10010.

45. **Van den Berghe G, Baxter RC, Weekers F, Wouters P, Bowers CY, Iranmanesh A, Veldhuis JD,
Bouillon R.** The combined administration of GH-releasing peptide-2 (GHRP-2), TRH and GnRH to men
with prolonged critical illness evokes superior endocrine and metabolic effects compared to treatment with
GHRP-2 alone. *Clin Endocrinol* 2002;56(5):655–669.

46. **Spratt DI, Longcope C, Cox PM, Bigos ST, Wilbur-Welling C.** Differential changes in serum
concentrations of androgens and estrogens (in relation with cortisol) in postmenopausal women with acute
illness. *J Clin Endocrinol Metab* 1993;76(6):1542–1547. doi:10.1210/jcem.76.6.8501162.

47. **Gottschlich MM, Khoury J, Warden GD, Kagan RJ.** An evaluation of the neuroendocrine response to
sleep in pediatric burn patients. *JPEN J Parenter Enteral Nutr* 2009;33(3):317–326.
doi:10.1177/0148607108325180.

48. **Arlt W, Hammer F, Sanning P, Butcher SK, Lord JM, Allolio B, Annane D, Stewart PM.**
Dissociation of serum dehydroepiandrosterone and dehydroepiandrosterone sulfate in septic shock.
Journal of Clinical Endocrinology & Metabolism 2006;91(7):2548–2554. doi:10.1210/jc.2005-2258.

49. **Mueller C, Blum CA, Trummler M, Stolz D, Bingisser R, Mueller C, Tamm M, Mueller B, Schuetz
P, Christ-Crain M.** Association of adrenal function and disease severity in community-acquired
pneumonia. Yende S, ed. *PLoS ONE* 2014;9(6):e99518–6. doi:10.1371/journal.pone.0099518.

50. **Van den Berghe G, de Zegher F, Wouters P, Schetz M, Verwaest C, Ferdinande P, Lauwers P.**
Dehydroepiandrosterone sulphate in critical illness: effect of dopamine. *Clin Endocrinol* 1995;43(4):457–
463. doi:10.1111/j.1365-2265.1995.tb02618.x.

51. **Dimopoulou I, Stamoulis K, Ilias I, Tzanela M, Lyberopoulos P, Orfanos S, Armaganidis A,
Theodorakopoulou M, Tsagarakis S.** A prospective study on adrenal cortex responses and outcome
prediction in acute critical illness: results from a large cohort of 203 mixed ICU patients. *Intensive Care
Med* 2007;33(12):2116–2121. doi:10.1007/s00134-007-0790-0.

52. **Osorio A, Vara-Thorbeck R, Rosell J, Osorio C, Ortega E, Ruiz-Requena E.** Dehydroepiandrosterone
sulfate and growth axis hormones in patients after surgery. *World J Surg* 2002;26(9):1079–1082.
doi:10.1007/s00268-002-6368-7.

53.   **Almoosa KF, Gupta A, Pedroza C, Watts NB.** Low Testosterone Levels are Frequent in Patients with
458   Acute Respiratory Failure and are Associated with Poor Outcomes. *Endocrine Practice*
459   2014;20(10):1057–1063. doi:10.4158/EP14003.OR.

460   54.   **Dolecek R.** Markers of bone remodelling, bone density and endocrine response after polytraumas and
461   burns. *OSTEOLOGICK BULLETIN* 2008;11(1):5–22.

462   55.   **Dossett LA, Swenson BR, Heffernan D, Bonatti H, Metzger R, Sawyer RG, May AK.** High Levels of
463   Endogenous Estrogens are Associated With Death in the Critically Injured Adult. *J Trauma*
464   2008;64(3):580–585. doi:10.1097/TA.0b013e31816543dd.

465   56.   **Folan MM, Stone RA, Pittenger AL, Stoffel JA, Hess MM, Kroboth PD.** Dehydroepiandrosterone,
466   dehydroepiandrosterone-sulfate, and cortisol concentrations in intensive care unit patients. *Crit Care Med*
467   2001;29(5):965–970.

468   57.   **Ilias I, Stamoulis K, Armaganidis A, Lyberopoulos P, Tzanela M, Orfanos S, Theodorakopoulou M,
469   Tsagarakis S, Dimopoulou I.** Contribution of endocrine parameters in predicting outcome of multiple
470   trauma patients in an intensive care unit. *Hormones (Athens)* 2007;6(3):218–226.

471   58.   **Brorsson C, Dahlqvist P, Nilsson L, Thunberg J, Sylvan A, Naredi S.** Adrenal response after trauma is
472   affected by time after trauma and sedative/analgesic drugs. *Injury* 2014;45(8):1149–1155.
473   doi:10.1016/j.injury.2014.02.001.

474   59.   **Bergquist M, Huss F, Fredén F, Mass GHC, 2016.** Altered adrenal and gonadal steroids biosynthesis in
475   patients with burn injury. *Clinical Mass Spectrometry* 2016;1:19–26. doi:10.1016/j.clinms.2016.10.002.

476   60.   **Kim MS, Shigenaga J, Moser A, Grunfeld C, Feingold KR.** Suppression of DHEA sulfotransferase
477   (Sult2A1) during the acute-phase response. *Am J Physiol Endocrinol Metab* 2004;287(4):E731–8.
478   doi:10.1152/ajpendo.00130.2004.

479   61.   **Lindh A, Carlström K, Eklund J, Wilking N.** Serum steroids and prolactin during and after major
480   surgical trauma. *Acta Anaesthesiol Scand* 1992;36(2):119–124. doi:10.1111/j.1399-6576.1992.tb03436.x.

481   62.   **Parker CR Jr, Baxter CR.** Divergence in adrenal steroid secretory pattern after thermal injury in adult
482   patients. *J Trauma* 1985.

483   63.   **Radford DJ, Wang K, McNelis JC, Taylor AE, Hechenberger G, Hofmann J, Chahal H, Arlt W,
484   Lord JM.** Dehydroepiandrosterone sulfate directly activates protein kinase C-beta to increase human
485   neutrophil superoxide generation. *Mol Endocrinol* 2010;24(4):813–821. doi:10.1210/me.2009-0390.

486   64.   **Brorsson C, Dahlqvist P, Nilsson L, Thunberg J, Sylvan A, Naredi S.** Adrenal response after trauma is
487   affected by time after trauma and sedative/analgesic drugs. *Injury* 2014:1–30.
488   doi:10.1016/j.injury.2014.02.001.

489   65.   **Berg C, Oeffner A, Schumm-Draeger PM, Badorrek F, Brabant G, Gerbert B, Bornstein S,
490   Zimmermann A, Weber M, Broecker-Preuss M, Mann K, Herrmann BL.** Prevalence of anterior
491   pituitary dysfunction in patients following traumatic brain injury in a German multi-centre screening
492   program. *Exp Clin Endocrinol Diabetes* 2010;118(2):139–144. doi:10.1055/s-0029-1225611.

493   66.   **Vogel AV, Peake GT, Rada RT.** Pituitary-testicular axis dysfunction in burned men. *J Clin Endocrinol*
494   *Metab* 1985;60(4):658–665. doi:10.1210/jcem-60-4-658.

495   67.   **Spratt DI, Bigos ST, Beitins I, Cox P, Longcope C, Orav J.** Both hyper- and hypogonadotropic
496   hypogonadism occur transiently in acute illness: bio- and immunoactive gonadotropins. *J Clin Endocrinol*
497   *Metab* 1992;75(6):1562–1570. doi:10.1210/jcem.75.6.1464665.

498   68.   **DANIELL H.** DHEAS Deficiency During Consumption of Sustained-Action Prescribed Opioids:
499   Evidence for Opioid-Induced Inhibition of Adrenal Androgen Production. *The Journal of Pain*
500   2006;7(12):901–907. doi:10.1016/j.jpain.2006.04.011.

501   69.   **Demling RH.** Comparison of the anabolic effects and complications of human growth hormone and the
502   testosterone analog, oxandrolone, after severe burn injury. *Burns* 1999;25:215–221.

503   70.   **Wolf SE, Edelman LS, Kemalyan N, Donison L, Cross J, Underwood M, Spence RJ, Noppenberger**
504   **D, Palmieri TL, Greenhalgh DG, Lawless M, Voigt D, Edwards P, Warner P, Kagan R, Hatfield S,**
505   **Jeng J, Crean D, Hunt J, Purdue G, Burris A, Cairns B, Kessler M, Klein RL, Baker R, Yowler C,**
506   **Tutulo W, Foster K, Caruso D, Hildebrand B, Benjamin W, Villarreal C, Sanford AP, Saffle J.**
507   Effects of oxandrolone on outcome measures in the severely burned: a multicenter prospective
508   randomized double-blind trial. *J Burn Care Res* 2006;27(2):131–9; discussion 140–1.
509   doi:10.1097/01.BCR.0000202620.55751.4F.

510   71.   **Li H, Guo Y, Yang Z, Roy M, Guo Q.** The efficacy and safety of oxandrolone treatment for patients
511   with severe burns: A systematic review and meta-analysis. *Burns* 2016;42(4):717–727.
512   doi:10.1016/j.burns.2015.08.023.

513   72.   **Bulger EM, Jurkovich GJ, Farver CL, Klotz P, Maier RV.** Oxandrolone does not improve outcome of
514   ventilator dependent surgical patients. *Ann Surg* 2004;240(3):472–8– discussion 478–80.

515   73.   **Gervasio JM, Dickerson RN, Swearingen J, Yates ME, Yuen C, Fabian TC, Croce MA, Brown RO.**
516   Oxandrolone in trauma patients. *Pharmacotherapy* 2000;20(11):1328–1334.

Legends

Figure 1.   Consort diagram. (A) recruitment process and (B) subgroup selection for analysis for sixty male

survivors of severe injury (NISS>15) under 50 years of age who had not been given exogenous steroids were

analysed**.**

Figure 2. Patient Characteristics of the Analysis Cohort. (A) Demographics, (B) Mechanism of Injury and

(C) the distribution of septic episodes for 60 male survivors from severe injury (NISS>15) under 50 years of age.

Figure 3. Serum steroids in 60 male survivors of severe injury (NISS>15) under 50 years of age. Serum

concentrations shown include cortisol (A), cortisone (B), the cortisol/cortisone ratio (C), DHEA (D), DHEAS

(E), the DHEA/ DHEAS ratio (F), the cortisol/DHEAS ratio (G), androstenedione (H), and testosterone (I). Data

are represented after modelling of the raw data (Suppl. Fig. 1) using a non-linear mixed effects model that

accounts for unbalanced repeated measures using a 4-knot cubic spline. Modelled data are shown as means and

95% confidence intervals.

Figure 4. The relationship between   (A) Urinary Nitrogen Excretion or   (B) biceps muscle thickness with

(B and D) DHEA and (C and F) testosterone, over time for young (<50), severely injured (NISS>15) males

who had survived and not been given anabolic steroids. Muscle thickness data was modelled using a mixed

effects technique; modelling time as a 6 and 7-knot restricted cubic spline respectively provided the best fit. Data

are means and 95% confidence intervals for model-based predicted fixed effects of time are shown.

Figure 5. Sequential Organ Failure Assessment (SOFA) score and probability of sepsis in relation to

endocrine response. SOFA and sepsis are related serum concentrations of DHEA (Panels A+B), DHEAS

(Panels C+D), and testosterone (E+F). Data were modelled using a non-linear mixed effects model that accounts

for unbalanced repeated measures using a 4-knot cubic spline. Modelled data are reported as means and 95%

confidence intervals.

Suppl. Table 1. Longitudinal Studies in critically ill patients analysing DHEA and DHEAS.

Suppl. Fig. 1. Serum steroid concentrations in 60 male survivors from severe injury (NISS>15) under 50

years of age. Serum concentrations of cortisol (A), cortisone (B), DHEA (C), DHEAS (D), androstenedione

(E), and testosterone (F) are shown as medians and interquartile ranges; the normal reference range is indicated

by the grey shaded box.

Suppl. Fig. 2: Male gonadal axis parameters during the first 4 weeks following injury. Luteinising hormone

(LH), sex hormone binding globulin (SHBG) and serum testosterone concentrations in male survivors <50 years

after severe injury (NISS>15). LH and SHBG were only measured in the military cohort, therefore represent

results from 43 individuals. Data are represented after modelling based on the raw data, using a non-linear mixed

effects model that accounts for unbalanced repeated measures using a 4-knot cubic spline.

Suppl. Fig 3. Sequential Organ Failure Assessment (SOFA) score and probability sepsis in relation to

endocrine response. SOFA and sepsis are related to the ratios of serum DHEA over serum DHEAS (Panels

A+B), and the ratio serum cortisol over serum DHEAS (Panels C+D) compared to serum (A), DHEA/DHEAS

ratio (C) Cortisol/DHEAS. Data were modelled using a non-linear mixed effects model that accounts for

unbalanced repeated measures using a 4-knot cubic spline. Modelled data are reported as means and 95%

confidence intervals.

Figure 1. Consort diagram. (A) recruitment process and (B) subgroup selection for analysis for sixty male survivors of severe injury (NISS>15) under 50 years of age who had not been given exogenous steroids were analysed**.**

A

Demographics	
Patients n=60	Median (IQR)
Age	27 (24-31)
GCS	14 (3-15)
ISS	25 (17-32)
NISS	34 (29-44)
TRISS	94 (56-98)
APACHE 2 (Dayl)	21 (14-25)
SAPS 2 (Dayl)	48 (26-54)
SOFA (Dayl)	9 (6-10)
Septic Episodes	1 (0-2)
RBCs (Day1), Units	10 (2-17)
FFP (Day 1), Units	9 (1-15)
Crystalloid/colloid (Day1), L	0 (0-1)
Ventilator Days	9 (5-12)
ICU Length of Stay, Days	9 (5-16)
Operative procedures	5 (3-7.5)
Hospital Length of Stay, Days	36 (19-56)

Figure 2. Patient Characteristics of the Analysis Cohort. (A) Demographics, (B) Mechanism of Injury and (C) the distribution of septic episodes for 60 male survivors from severe injury (NISS>15) under 50 years of age.

Figure 3. Serum steroids in 60 male survivors of severe injury (NISS>15) under 50 years of age. Serum concentrations shown include cortisol (A), cortisone (B), the cortisol/cortisone ratio (C), DHEA (D), DHEAS (E), the DHEA/ DHEAS ratio (F), the cortisol/DHEAS ratio (G), androstenedione (H), and testosterone (I). Data are represented after modelling of the raw data (Suppl. Fig. 1) using a non-linear mixed effects model that accounts for unbalanced repeated measures using a 4-knot cubic spline. Modelled data are shown as means and 95% confidence intervals.

Figure 4. The relationship between (A) Urinary Nitrogen Excretion or (B) biceps muscle thickness with (B and D) DHEA and (C and F) testosterone, over time for young (<50), severely injured (NISS>15) males who had survived and not been given anabolic steroids. Muscle thickness data was modelled using a mixed effects technique; modelling time as a 6 and 7-knot restricted cubic spline respectively provided the best fit. Data are means and 95% confidence intervals for model-based predicted fixed effects of time are shown.

Figure 5. Sequential Organ Failure Assessment (SOFA) score and probability of sepsis in relation to endocrine response. SOFA and sepsis are related serum concentrations of DHEA (Panels A+B), DHEAS (Panels C+D), and testosterone (E+F). Data were modelled using a non-linear mixed effects model that accounts for unbalanced repeated measures using a 4-knot cubic spline. Modelled data are reported as means and 95% confidence intervals.