Depiction of secondary metabolites and antifungal activity of *Bacillus velezensis* DTU001

Sagarika Devi[a], Heiko T. Kiesewalter[a], Renátó Kovács[b], Jens Christian Frisvad[a], Tilmann Weber[c], Thomas Ostenfeld Larsen[a], Ákos T. Kovács[a,*], Ling Ding[a,*]

[a] Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kgs Lyngby, Denmark

[b] Department of Medical Microbiology, Faculty of Medicine, University of Debrecen, 4032 Debrecen, Hungary

[c] Center for Biosustainability, Technical University of Denmark, 2800 Kgs Lyngby, Denmark

Corresponding authors: Ling Ding (lidi@dtu.dk) and Ákos T. Kovács (atkovacs@dtu.dk)

ABSTRACT

For a safe and sustainable environment, effective microbes as biocontrol agents are in high demand. Spore-forming *Bacilli* play an important role as effective and safe antifungal bacteria. To satisfy the need for novel and effective biocontrol strains, an airborne *Bacillus velezensis* isolate DTU001 was investigated due to its broad antifungal spectrum, and genomic and metabolomics investigations were performed. Genomic analysis of *B. velezensis* DTU001 revealed that it harbored the genetic potential to produce ribonuclease barnase, antibacterial bacillaene and macrolactin, and antifungal lipopeptides including iturins, fengycins and surfactins. The production of lipopeptides was uncovered in HPLC-HRMS metabolomics analysis. To test the antifungal efficacy, extracts of *B. velezensis* DTU001, together with standards of iturins, fengycins were tested against a range of twenty human or plant pathogenic fungi covering genera of *Penicillum*, *Aspergillus*, *Talaromyces*, *Candida*, and *Cryptococcus*. *B. velezensis* DTU001 extract exhibited strong and broad antifungal effect against the tested target species while some test fungi were insensitive to single exposure to iturin or fengycin. The isolate showed analogous biofilm formation to other closely related *Bacilli*. To further support the biocontrol properties of the isolate, coculture with *Candida albicans* demonstrated that *B. velezensis* DTU001 exhibited excellent antiproliferation effect against *C. albicans*. In summary, the described isolate is a potential antifungal agent with a broad antifungal spectrum that will help to avoid hazardous pathogenic fungi and provide alternative to toxicity caused by chemicals.

Keywords: *Bacillus velezensis*, *Candida albicans*, secondary metabolites, biocontrol, biofilm

1. Introduction

The prerequisite for benign and effective antifungal agents increases in parallel with the existing and escalating emergence of fungal pathogens resistant to current therapies (De Lucca and Walsh 1999), being relevant to both therapeutics and biocontrol. While infectious diseases impose serious public health burden and often have devastating consequences, plant diseases constitute an emerging threat to global food security. Additionally, extensive use of chemicals to control plant diseases has distressed the ecological balance of microbes inhabiting soil. This in turn has led to the appearance of resistant pathogenic strains, groundwater contamination, and obvious health risks to human beings. Thus, the development of environmental friendly alternatives to the currently used chemical pesticides for combating a variety of crop diseases is nowadays one of the biggest ecological challenges (Ongena and Jacques 2008). It is therefore vital to find antifungal agents that are non-hazardous to human health as well as to the environment.

Several microorganisms have so far been discovered for the development of biopesticides at commercial level. Among the various microorganisms, *Bacillus* spp., *Pseudomonas* spp., and some *mycorrhizal fungi.* are widely used groups for biocontrol (Rahman, 2018). Especially, the *Bacillus* genus was proven to be source an excellent source of antifungal agents and thus *Bacilli* are widely used as biocontrol agents (Chae et al. 1990; Shafi et al. 2017). The US Food and Drug Administration (US FDA) declared *Bacillus subtilis* as GRAS (generally recognized as safe) organisms for its use in food processing industries (Denner W 1996). Members of the *B. subtilis* groups were found to produce numerous highly active antifungal compounds (Grubbs, et al. 2017). The *B. subtilis* group contains the "Operational Group *Bacillus amyloliquefaciens*" that includes *B. amyloliquefaciens*, *Bacillus velezensis*, and *Bacillus siamensis* (Fan et al. 2017). Among these, *B. velezensis* FZB42 (formerly *B. amyloliquefaciens*

subsp. *plantarum* FZB42) is the most researched commercially used *Bacillus*-based biofertilizer and biocontrol agent in agriculture.

Antifungal cyclopeptides appeared to play a major role in biological control of plant pathogens (Chae et al. 1990; Verschuere et al. 2000). Composed of a peptide and a fatty acyl moiety, these amphiphilic metabolites have consolidated their position as potent therapeutic compounds, such as daptomycin, a lipopeptide antibiotic used in human medicine (Kanafani and Corey 2007). Antibacterial, -viral, -fungal, -tumor activity and immune-modulatory effects are some of their proven and emerging properties (Meena 2015). In addition, secondary metabolites seem to play an important role in bacterial-fungal interaction of *Bacilli* (Benoit et al., 2015) and other bacterial life styles, including surface spreading (Grau et al. 2015; Kearns 2010; Hölscher and Kovács 2017).

Here, we describe a new *Bacillus* isolate DTU001, which exhibited a broad and strong antifungal spectrum against different pathogenic fungi ranging from plant pathogens to human pathogens. Using *in silico* DNA hybridization, it was identified as *B. velenzensis*. To dissect the genetic potential as an antifungal agent, we genome-mined BGCs in 31 *Bacillus* species and compared their biosynthetic potential with a special focus on commercially applied strains. To detect the synthesized secondary metabolites of *B. velezensis* DTU001, the metabolic profile of the strain's crude extract was investigated.

Lipopeptides produced by *Bacilli*, such as iturin, fengycin and surfactin are believed to contribute to their antifungal potential. When novel isolates from the *B. subtilis* species complex are evaluated, the role of single lipopeptide in antifungal activity is scrutinized lacking systematic *in vitro* evaluation of the antifungal efficacy of the combination of these secondary metabolites. To fill in this gap, we also assayed the efficacy of purified iturin and fengycin in comparison to the crude extract of *B. velezensis* DTU001 using twenty critical human and plant

pathogens. In this report, we outline the phylogeny, genomics, metabolomics, biofilm formation, and antifungal effects of *B. velezensis* DTU001.

2. Materials and methods

2.1. Bacterial and fungal strains

B. velezensis DTU001 is an airborne bacterium, isolated indoor at the DTU campus, Kongens Lyngby Denmark. Twenty selected plant/human pathogens (Table 1) covering five genera *Aspergillus*, *Penicillium*, *Talaromyces*, *Cryptococcus* and *Candida* were obtained from the DTU Bioengineering IBT fungal collection. All microbes are maintained at the department.

2.2. Genome sequencing

Genomic DNA was isolated from *B. velezensis* DTU001 using Bacterial and Yeast Genomic DNA kit (EURx). Genome sequencing has been performed on PacBio RSII platform, *de novo* assembly and annotation were done using the RS HGAO Assembly (v3.0) and Prokka (v1.12b) software, respectively. The complete sequence has been deposited in NCBI (accession number CP035533) followed by automatic annotation via NCBI PGAP (Haft, et al. 2018).

2.3. Chemicals

Solvents used in this study were LC–MS grade, and all other chemicals were analytical grade. All were from Sigma-Aldrich (Steinheim, Germany) unless otherwise stated. Water was purified using a Milli-Q system (Millipore, Bedford, MA). ESI–TOF tune mix was purchased from Agilent Technologies (Torrance, CA, USA).

2.4. Digital DNA–DNA hybridization (dDDH)

The genome-to-genome-distance calculator (GGDC) version 2.1 provided by DSMZ (http://ggdc.dsmz.de/) was used for genome-based species delineation (Meier-Kolthoff et al. 2013) and genome-based subspecies delineation (Meier-Kolthoff et al. 2014). Distances were

calculated by (i) comparing two genomes using the chosen program to obtain HSPs/MUMs and (ii) inferring distances from the set of HSPs/MUMs using three distinct formulas. Next, the distances were transformed to values analogous to DDH. The DDH estimates were based on an empirical reference dataset comprising real DDH values and genome sequences. The DDH estimate resulted from a generalized linear model (GLM) which also provided the estimate's confidence interval (after the ± sign). Three formulas are available for the calculation: (1) HSP length/total length; (2) identities/HSP length; and (3) identities/total length. Formula 2, which is especially appropriate to analyze draft genomes, was used in this study.

2.5. Construction of the evolutionary tree

An initial whole genome phylogeny was inferred using the automlst tool (http:/automlst.ziemertlab.com; Ziemert et al., personal communication). This automatically identifies related strains based on a seed collection of sequences obtained from NCBI Genbank (*B. velezensis* DTU001, *B. thuringiensis* serovar Berliner ATCC 10792, *B. cereus* 03BB102, *B. megaterium* DSM319, *B. velezensis* FZB42), using a MASH-based algorithm, that extracts 80 phylogenetically relevant genes and builds a MLST phylogeny out of these data. The sequence alignment obtained from automlst was then manually refined (removal of very similar species). To infer the best phylogenetic model, jModeltest (v.2.1.10) was used with default parameters (Darriba et al. 2012). In all three evaluation criteria (Akaike Information Criterion AIC, Bayesian Information Criterion BIC and the Decision Therory Performance-based Selection), the GTR+I+G model was deemed most suitable. Thus, a maximum-likelihood tree was constructed in PhyML (v3.3.20180621) (Guindon et al. 2010) using the parameters suggested by jModeltest. The tree then was checked with Dendroscope (v3.5.10) (Huson and Scornavacca 2012) and plotted using the ETE3 toolkit (Huerta et al. 2016).

2.6. Genome mining of secondary metabolites

B. velezensis DTU001 and other 32 Bacillus strains were mined for the biosynthetic gene clusters using the antiSMASH platform (Weber et al. 2015) and BiG-SCAPE ((Navarro-Muñoz et al. 2018).

2.7. Agar-diffusion assay

Agar-diffusion assay was carried to test the anti-fungal activities. Ethyl acetate extract of *B. velezensis* DTU001 was dissolved in methanol (1 mg/mL). DMSO was used as a negative control. Initial screening of the bacterial extract was done by standard agar disc diffusion method (Bauer et al., 1966) and the zone of inhibition was noted. The 16 pathogenic filamentous fungi were grown on sterile MEAOX plates and the 4 yeasts were cultivated on PDA at 25 °C. 5-days old fungal plates and 2-days old yeast plates were taken for further investigation. Spore suspensions with a final cell concentration of $5x10^5$ cfu/mL of each microbial culture (fungi and yeast) were maintained for the disc diffusion assay. Plugs were harvested from 48 hr grown *B. velezensis* DTU001 from MEAOX plates and extracted in isopropanol-ethyl acetate-1% formic acid followed by 1 hr ultra-sonication. The supernatant was collected and dried utilizing nitrogen evaporator. The test sample (*B. velezensis* extract) and the standard compounds (iturin and fengycin) were dissolved in 0.5% DMSO to a concentration of 1 mg/mL each. Sterile Muller Hinton Agar plates were inoculated with the test organisms using lawn culture method. Discs of 6mm were cut at the centre of the sterile plates using a sterile cork borer. The discs were loaded with 50 µl of each test sample. The plates were maintained at 25 ± 2 °C for 5 days and observed for clear zone of inhibition. Iturin and fengycin standards were maintained as positive control. Blank assays in DMSO were

applied to avoid any possible effect of the solvent. Each experiment was conducted in triplicates for further confirmation and reproducibility of results.

2.8. Anti-Candida effects under planktonic and biofilm conditions

The impact of *B. velezensis* DTU001 on *C. albicans* SC5314 reference strain was assayed. *C. albicans* and *B. velezensis* were subcultured on Sabouraud dextrose agar and lysogeny broth (LB) agar, respectively. The growth of the individual strains in both monoculture and coculture was evaluated in order to examine the effect of *B. subtilis* on *C. albicans* planktonic growth. The final cell concentrations were adjusted to $2x10^5$ cells/ml in RPMI-1640 broth + LB broth (50:50 v/v %) both for *C. albicans* and *B. subtilis*. Test tubes were incubated with agitation in darkness at 35 °C. 100 µl culture was removed at 0, 2, 4, 6, 8, 12 and 24 hours, serially diluted tenfold in sterile physiological saline then plated onto either Sabouraud dextrose agar or LB agar. Plates were incubated for 24 and 48 hours at 35 °C for *B. subtilis* and *C. albicans*, respectively. Results are representative of three independent experiments and are expressed as mean ± SD (error bars), which was presented using GraphPad Prism 6.05.

For the biofilm assays, polystyrene flat-bottom 96-well microtiter plates (TPP, Trasadingen, Switzerland) were used for the formation of single and mixed species biofilms of *C. albicans* and *B. subtilis*. A volume of 200 µl of cell suspension ($1x10^6$ cells/mL in RPMI-1640 broth + LB broth (50:50 v/v%) for *C. albicans* and $1x10^6$ cells/mL in RPMI-1640 broth + LB broth (50:50 v/v%) for *B. subtilis*) was pipetted to each well for single biofilms, while 100 µL from each suspension ($2x10^6$ cells/mL for *C. albicans* with $2x10^6$ cells/ml for *B. subtilis*) was added for mixed species biofilms to wells assigned to endpoints 2, 4, 6, 8, 12 and 24 hours. The plates were incubated in darkness with static conditions. After 2, 4, 6, 8, 12 and 24 hours of incubation, the corresponding wells were washed three times with sterile physiological saline then 200 µL sterile saline was added into wells, scraped off the adhered cells from surfaces;

afterwards, aliquots of 100 μL were removed. Cells were released by mild sonication and vortexing then diluted serially ten-fold and plated onto either Sabouraud dextrose agar or LB agar. Plates were incubated for 24 and 48 hours at 35 °C for *B. subtilis* and *C. albicans*, respectively. Results are representative of three independent experiments and are expressed as mean ± SD (error bars), which was presented using GraphPad Prism 6.05.

2.9. Bacterial biofilm formation

Bacterial strains (*B. velezensis* DTU001, *B. subtilis* NCBI 3610 (Branda et al., 2001), *B. subtilis* ATCC6633 (ATCC collection), *B. subtilis* PS216 (Stefanic and Mandic-Mulec 2009), *B. velezensis* FZB42 (obtained from Bacillus Genetic Stock Centre), and *B. methylotrophicus* NCCB100236 (obtained from Bacillus Genetic Stock Centre) were inoculated in lysogeny broth (Lenox LB, Carl Roth, Germany) and grown overnight for 16 hours at 37°C at 225 rpm. For pellicle development 20 μl was inoculated in 2 mL MSgg medium and placed in a 24-well plate, while 5 μL spotted on MSgg medium solified with 1,5 % agar that was prepared as described previously (Holscher et al. 2016) (Branda, González-Pastor et al. 2001, Holscher, Dragos et al. 2016). Both pellicle and colony biofilms were incubated at incubated at 30°C static conditions and imaged after 3 days.

3.10. Secondary metabolites identification

A UHPLC–DAD–QTOF method was set up for screening, with typical injection volumes of 0.1–2 μL extract. Separation was performed on a Dionex Ultimate 3000 UHPLC system (Thermo Scientific, Dionex, Sunnyvale, California, USA) equipped with a 100×2.1 mm, 2.6 μm, Kinetex C_{18} column, held at a temperature of 40 °C, and using a linear gradient system composed of A: 20 mmol L^{-1} formic acid in water, and B: 20 mmol L^{-1} formic acid in

acetonitrile. The flow was 0.4 mL min^{-1}, 90 % A graduating to 100 % B in 10 min, 100 % B 10–13 min, and 90 % A 13.1–15 min.

Time-of-flight detection was performed using a maXis 3G QTOF orthogonal mass spectrometer (Bruker Daltonics, Bremen, Germany) operated at a resolving power of ~50000 full width at half maximum (FWHM). The instrument was equipped with an orthogonal electrospray ionization source, and mass spectra were recorded in the range m/z 100–1000 as centroid spectra, with five scans per second. For calibration, 1 µL 10 mmol L^{-1} sodium formate was injected at the beginning of each chromatographic run, using the divert valve (0.3–0.4 min). Data files were calibrated post-run on the average spectrum from this time segment, using the Bruker HPC (high-precision calibration) algorithm.

For ESI$^+$ the capillary voltage was maintained at 4200 V, the gas flow to the nebulizer was set to 2.4 bar, the drying temperature was 220 °C, and the drying gas flow was 12.0 L min^{-1}. Transfer optics (ion-funnel energies, quadrupole energy) were tuned on HT-2 toxin to minimize fragmentation. For ESI$^-$ the settings were the same, except that the capillary voltage was maintained at −2500 V. Unless otherwise stated, ion-cooler settings were: transfer time 50 µs, radio frequency (RF) 55 V peak-to-peak (Vpp), and pre-pulse storage time 5 µs.

3. Results and discussion

3.1. The genome of the indoor airborne isolate B. velezensis DTU001 encodes numerous secondary metabolites

B. velezensis DTU001 was isolated indoor at the DTU campus in Denmark and selected for detailed analysis based on its potential antimicrobial activity. The complete genome sequence of *B. velezensis* DTU001 consisted of a circular 3.927.210 bp chromosome with a GC content of 46.6%. Its 16S rRNA gene sequence exhibited 99% similarity to *B. velezensis* and *B. amyloliquefaciens* in genome database.

The phylogenetic affiliation of DTU001 isolate was further investigated using digital DNA:DNA-hybridization (dDDH) similarities and differences in genomic G+C content. (https://ggdc.dsmz.de/). These distances are transformed to values analogous to DDH using a generalized linear model (GLM) inferred from an empirical reference dataset comprising real DDH values and genome sequences. Compared to *B. velezensis* FZB42 (NC-009725.1), DDH value of 80.3%, difference in % G+C: 0.10. Meanwhile, there is a DDH analogous value of *B. amyloliquefaciens* (FN597644.1) 55.4%, difference in % G+C: 0.51. This result clearly supports that DTU001 is closer related to *B. velezensis*, where the DDH value exceeded the threshold of 70%.

A phylogenetic analysis with automlst (http://automlst.ziemertlab.com) was carried out, which automatically extracts 80 conserved genes, concatenates them, computes a multiple sequence alignment and builds a phylogenetic tree. The sequence alignment was manually checked, and a maximum-likelihood tree refined with jModelTest2 (Darriba et al. 2012) and PhyML (Guidon et al. 2010). According to the MLST analysis, *B. velezensis* DTU1 forms a clade with other *Bacilli* that were recently proposed to be assigned the new species *Bacillus lentimorphus* (Parks et al. 2018).

To determine the genetic potential of *B. velezensis* DTU001 as biocontrol strain, the complete genome was surveyed for biosynthetic gene clusters that identified nine gene clusters involved in nonribosomal synthesis of lipopeptides, polyketides, and siderophore bacillibactin. About 15.4 % of the whole DTU001 genome was devoted to secondary metabolites synthesis of nonribosomal synthesis of antimicrobial compounds and siderophores. In addition, genes for synthesis of the highly bioactive extracellular alkaline guanyl-preferring ribonuclease, previously detected in *B. amyloliquefaciens* (Hartley et al. 1972) was also identified in the genome of DTU001.

In order to compare the biosynthetic potentials between different *Bacilli*, *B. velezensis* together with other 17 species were mined for the genetic potential of secondary metabolite production using antiSMASH and BiG-SCAPE. All *Bacilli* related to *B. velezensis* DTU001 have the genetic potential to utilize bacillibactin as a siderophore to acquire irons from the environments (**Fig. 1**). All selected *B. amyloliquefaciens*, *B. velezensis*, and *B. subtilis* strains showed high similarities in BGC potentials and are considered as talented strains producing various lipopetides or polyketides. More distantly related *Bacillus* strains, such as members of the *Bacillus cereus* group showed a different secondary metabolite production genotype. All of these strains (*B. cereus*, *B. anthracis, B. thuringiensis*) seem to be unable of producing lipopeptides of the surfactin, iturin or fengycin families.

Fig. 1. Evolutionary relationships of taxa between *B. velezensis* DTU001 and other *Bacilli*. A maximum-likelihood phylogenetic tree based on 80 conserved genes of selected strains of *Bacilli* is shown; *B. megaterium* DSM 319 was selected as outgroup. The presence/absence of typical *Bacillus* BGCs is depicted on the right.

3.2. Antifungal characterization of B. velezensis

Iturins are well-known secondary metabolites important for biocontrol against plant pathogens such as *Xanthomonas campestris* subsp. *Cucurbitae, Pectobacterium carotovorum* subsp. *Carotovorum, Rhizoctonia solani, Fusarium graminearum* etc. (Gong et al. 2015; Zeriouh et al. 2011). However, the impact of iturins against medically-relevant fungal pathogens is less explored. Therefore, we tested the efficacy and target specificity of DTU001 against selected important plant and human pathogenic fungi. In addition, we compared the effectiveness of the DTU001 crude extract and the purified lipopeptides, iturin and fengycin. We have selected 17 important plant and human pathogenic filamentous fungi (Table 1) ranging from the genera *Penicillium*, *Aspergillus*, *Talaromyces*, *Candida* to *Cryptococcus*. Besides those notorious plant pathogens such as *Aspergillus uvarum* infecting grapes, citrus rotting *Penicillium ulaiense* and post-harvest pathogen *Penicillium expansum* infecting apples, several important human pathogens were tested, such as *Aspergillus niger* and *Candida*.

The organic extracts of *B. velezensis* DTU001 displayed strong antifungal effects against *A. uvarum*, *P. ulaiense* and *P. expansum* (Table1 and Fig. 2). Due to the susceptibility to infection of mature and overripe fruit, post-harvest treatment of fruit with fungicides has traditionally been the most common method of combating *P. expansum*. Also, bio-fungicides using active ingredients such as bacteria and yeast have been successful in preventing infection but are ineffective against existing infections (Konstantinou et al. 2011). Considering the size of the apple product industry and the large number of people that may come into contact with infected fruits, control of the *P. expansum* is vitally important. Interestingly both *Talaromyces atroroseus* and *Talaromyces amestolkiae* were inhibited by both fengycin (20 cm) and iturin (29, 26 cm) as well as the crude extract (Table 1 and Fig. 2). All the tested organisms showed less or no susceptibility towards fengycin.

Fig. 2. Extracts of *B. velezensis* DTU001 against fungi showing clear inhibition zones. Images left to right: *P. frequentans*, *A. niger* and *T. amestolkiae*. DMSO control loaded to the left hole, while the crude extract was applied to the right hole demonstrating the specific inhibition zone against the fungi.

In summary, the agar diffusion test showed that iturin and fengycin displayed selective anti-fungal activities against certain fungal pathogens, which means that some strains are insensitive to the addition of a single purified compound. For example, neither iturin nor fengycin displayed activities against *Penicillium sclerotigenum* and *Aspergillus calidoustus*. Overall, iturin displayed stronger and broader activities compared to fengycin. Compared to iturin and fengycin standards, extracts of *B. velezensis* DTU001 exhibited stronger inhibition activities and displayed a broader spectrum against test fungi including *P. sclerotigenum* and *A. calidoustus*. One explanation might be that *B. velezensis* produced a cocktail mixture which functions in a synergistic and more effective manner or additional secondary metabolite plays a role in the inhibition. However, the combined impact of the lipopeptides could avoid fast resistance evolution and help the bacterium in the environmental survival. In addition, these results also indicate that the organic extracts of *B. velezensis* DTU001 could be a better biofungicide than commercially used purified iturin, even though they were actually not that pure (see supplementary fig. XX). It seemed that *Aspergillus felis* harbored certain mechanisms to avoid anti-fungal effects from *B. velezensis* DTU001 as neither lipopeptide standards nor the bacterial crude extract displayed inhibition activities against this fungus.

3.3. B. velezensis DTU001 is able to lessen biofilms of C. albicans

As *B. velezensis* DTU001 showed broad anti-fungal activity spectrum, we further evaluated if the bacterial isolate can directly reduce the growth of the human pathogen fungus, *C. albicans* under planktonic conditions and during biofilm development. First, the growth of the fungus and bacterium was monitored in planktonic co-cultures and compared to the mono-cultures (Fig. 3). Importantly, the *B. velezensis* cell count in the coculture with *C. albicans* did not differ significantly compared monoculture. In contrast, *C. albicans* planktonic cells between 6 and 24 hours diminished in the presence of *B. velezensis* DTU001, therefore the bacterial isolate was able to inhibit the proliferation of the fungus in the co-cultures. Similarly, *B. velezensis* DTU001 was able to lessen the biofilms of *C. albicans* when co-cultured. The inhibitory effect was even higher that under planktonic conditions after 24 hours (2-log decrease in fungal cell count). These experiments further support the applicability of *B. velezensis* DTU001 as biocontrol agent even in the presence of the target organism.

Fig. 3. Planktonic (A) and biofilm (B) cultivation of *C. albicans* SC5314 and *B. velezensis* DTU001. Colony forming units were recorded at the indicated time points for *C. albicans* (open rhombus indicates *C. albicans* alone, open square denotes *C. albicans* in the presence of *B.*

subtilis) and *B. velezensis* (open circle indicates *B. subtilis* alone, filled circle denotes *B. velezensis* in the presence of *C. albicans*).

3.4. B. velezensis DTU001 is a potent biofilm former

The probiotic effects of *Bacilli* depend on the ability to form biofilm and production of a matrix, that acts as an immune system stimulator (Jones et al. 2014), protects from environmental stress (Kovacs et al. 2012), and is required for plant colonization (Beauregard et al, 2013; Gallegos-Monterrosa et al., 2016). Therefore, we have compared the biofilm formation of *B. velezensis* DTU001 to other *Bacillus* species known for biofilm formation. Both architecturally complex colonies on agar medium and air-liquid localized pellicles were tested. The biofilm development can be qualitatively monitored using these models: the more complex structures, the increased the biofilm development of *Bacilli* (Branda et al. 2001; Gallegos-Monterrosa et al. 2016). In addition, pellicle formation strongly depends on the matrix components, without which cell are unable to float on the air-medium interface (Dragos et al. 2018). Both the complex biofilm colonies and the pellicles showed robust biofilm formation by *B. velezensis* DTU001 that was comparable to other closely related isolates from the *B. subtilis* species complex (Fig. 4).

Fig. 4. Pellicle (above) and colony biofilms (below) of selected strains from the *B. subtilis* species complex. Scale bar indicates 5 mm.

3.5. Secondary metabolites characterization of B. velezensis DTU001

Based on the genome sequences and antifungal activity, *B. velezensis* DTU001 was predicted to produce various lipopeptides. To specifically demonstrate these secondary metabolites, LC-HRMS analysis was performed on extracts harvested from agar grown *B. velezensis* DTU001. Indeed, chemical analysis along with purified standards detected three groups of bioactive lipopeptides: iturins, fengycins, and surfactins (Fig. 5, Fig. S1, S2 and S3 in the supplemental data).

Fig. 5. Base peak chromatogram showing the presence of iturin, fengycin and surfactin produced by *B. velezensis* DTU001.

4. Conclusions

Through genomics and metabolomics investigation, we demonstrated that *B. velezensis* DTU001 harbored the capacity to produce bioactive lipopeptides, comparable to the commercially utilized *B. amyloliquefaciens* and *B. velezensis* strains. Production of bioactive iturin, fengycin and surfactin by this strain was confirmed by HPLC-HRMS. Phylogenetic distance between DTU001 and those commercially utilized *B. amyloliquefaciens* together with their shared chemistry makes DTU001 as an attractive biocontrol option. Additionally, our targeted comparative genomics study revealed the versatility of secondary metabolite production by the selected *Bacilli*.

Either iturin or fengycin were previously proposed to be solely responsible for the antifungal activities of isolates within the *B. subtilis* species complex, however, we propose here that a cocktail anti-fungal agent derived from the crude extract of *B. velezensis* DTU 001 functions more efficiently than single components. In summary, this newly described isolate can be utilized as a biocontrol agent with a broad antifungal spectrum and provides a sustainable alternative for replacement of hazardous chemicals.

Declaration of interests

None.

Acknowledgements

This work was supported by the Danish National Research Foundation (DNRF137) for the Center for Microbial Secondary Metabolites (CeMiSt). SD was supported by a Novozymes and Henning Holck Larsen fellowship during her stay at DTU.

Table 1

Bioactivity of iturin (1mg/mL), fengycin (1mg/ mL), and crude *B. velezensis* extract (1mg/ mL) against 16 human- and plant-pathogenic filamentous fungi and 4 yeasts

Name of the fungi	Strain information	Zone of inhibition (mm)		
		Iturin	Fengycin	*B. velezensis* supernatant
A. niger IBT27878	harvest decay of fresh and dried fruits and certain vegetables, nuts, beans, and cereals [28-29]	27	-	25
A. terreus IBT 22089	opportunistic human pathogen; crops pathogen, e.g. wheat, ryegrass, and potatoes [35]	20	-	19
A. felis IBT 34215	causative agent of Aspergillosis in humans, dogs and cats	-	-	-
A. flavus IBT32105		30	-	26
T. amestolkiae IBT23485	common cosmopolitan species that has been cultured from indoor house dust, sputum and lungs from cystic fibrosis patients, indoor air, wheat, soil, pineapple, sculptures and manure	26	20	24
T. atroroseus IBT11181		29	20	30
P. expansum IBT 34672		24	-	20
P. frequentans IBT 34879		25	-	23
P. italicum IBT 34168		30	-	33
P. digitatum IBT 34217		+	-	+
P. ulaiense 23038		30	-	30
P. arizonense IBT12857		+	-	+
P. solitum IBT34928		33	-	33
P. sclerotigenum IBT 30034	opportunistic human infection, associated with blue mold of yam in Korea	-	-	32
A. uvarum IBT32575	contamination of several agricultural products,	25	-	35

	including grape-derived products, coffee and cocoa [36-38]			
A. calidoustus IBT 28322	opportunistic human infection	-	-	32
C. albicans IBT 654	superficial and serious systemic diseases	21	-	19
C. albicans IBT 657		24	-	20
C. neoformans IBT 10488		30	-	31
C. neoformans IBT 10496		26	-	29

References

Bauer, A.W., Kirby, W.M., Sherris, J.C., Turck M., 1966. Antibiotic susceptibility testing by a standardized single disk method. Am. J. Clin. Pathol. 45, 493-496.

Beauregard, P. B., Chai, Y., Vlamakis, H., Losick R., Kolter, R., 2013. *Bacillus subtilis* biofilm induction by plant polysaccharides. Proc. Natl. Acad. Sci.110, E1621-1630.

Benoit, I., van den Esker, M.H., Patyshakuliyeva, A., Mattern, D.J., Blei, F., Zhou, M., Dijksterhuis, J., Brakhage, A.A., Kuipers, O.P., de Vries R.P., Kovacs, A.T., 2015. *Bacillus subtilis* attachment to *Aspergillus niger* hyphae results in mutually altered metabolism. Environ. Microbiol. 17, 2099-2113.

Branda, S.S., Gonzalez-Pastor, J.E., Ben-Yehuda, S., Losick R., Kolter R., 2001. Fruiting body formation by *Bacillus subtilis*. Proc. Natl. Acad. Sci. 98, 11621-11626.

Chae G.P., Shoda M., Kubota, H., 1990. Suppressive effect of *Bacillus subtilis* and its products on phytopathogenic microorganisms. J. Ferment. Bioeng. 69, 1-7.

Darriba, D., Taboada, G.L., Doallo R., Posada, D., 2012. jModelTest 2: more models, new heuristics and parallel computing. Nat Methods 9, 772.

De Lucca, A.J., Walsh T.J., 1999. Antifungal peptides: novel therapeutic compounds against emerging pathogens. Antimicrob Agents Chemother 43, 1-11.

Denner, W.G.T., 1996. The legislative aspects of the use of industrial enzymes in the manufacture of food and food ingredients. Industrial enzymology. New York: Stockton Press: 6.

Dragos, A., Kiesewalter, H., Martin, M., Hsu, C.Y., Hartmann, R., Wechsler, T., Eriksen, C., Brix, S., Drescher, K., Stanley-Wall, N., Kummerli, R., Kovacs, A.T., 2018. Division of labor during biofilm matrix production. Curr. Biol. 28, 1903-1913.

Fan, B., Blom, J., Klenk H.P., Borriss, R., 2017. *Bacillus amyloliquefaciens*, *Bacillus velezensis*, and *Bacillus siamensis* form an "operational group *B. amyloliquefaciens*" within the *B. subtilis* species complex. Front Microbiol 8, 22.

Gallegos-Monterrosa, R., Mhatre, E., Kovacs, A.T., 2016. Specific *Bacillus subtilis* 168 variants form biofilms on nutrient-rich medium. Microbiol 162, 1922-1932.

Gong, A.D., Li, H.P., Yuan, Q.S., Song, X.S., Yao, W., He, W.J., Zhang, J.B., Liao Y.C., 2015. "Antagonistic mechanism of iturin A and plipastatin A from *Bacillus amyloliquefaciens* S76-3 from wheat spikes against *Fusarium graminearum*. PLoS One 10, e0116871.

Grau, R. R., de Ona, P., Kunert, M., Lenini, C., Gallegos-Monterrosa, R., Mhatre, E., Vileta, D., Donato,V., Holscher, T., Boland, W., Kuipers, O.P., Kovacs, A.T. 2015. A duo of potassium-responsive histidine kinases govern the multicellular destiny of *Bacillus subtilis*. MBio 6, e00581.

Grubbs, K. J., Bleich, R.M., Santa Maria, K.C., Allen, S.E., Farag, S., AgBiome, T., Shank, E.A., Bowers, A.A., 2017. Large-scale bioinformatics analysis of Bacillus genomes uncovers conserved roles of natural products in bacterial physiology. mSystems 2, e00040-17

Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59, 307-321.

Haft, D. H., DiCuccio, M., Badretdin, A., Brover, V., Chetvernin, V., O'Neill, K., Li, W., Chitsaz, F., Derbyshire, M.K., Gonzales, N.R., Gwadz, M., Lu, F., Marchler, G.H., Song, J.S., Thanki, N., Yamashita, R.A., Zheng, C., Thibaud-Nissen, F., Geer, L.Y., Marchler-Bauer, A., Pruitt, K.D., 2018. RefSeq: an update on prokaryotic genome annotation and curation. Nucleic Acids Res 46, D851-D860.

Hartley, R.W., Rogerson, Rogersen Jr, D.L., Smeaton, J.R., 1972. Production and purification of the extracellular ribonuclease of *Bacillus amyloliquefaciens* (barnase) and its intracellular inhibitor (barstar). II. Barstar. Prep Biochem 2, 243-250.

Holscher, T., Dragos, A., Gallegos-Monterrosa, R., Martin, M., Mhatre, E., Richter, A., Kovacs, A.T., 2016. Monitoring spatial segregation in surface colonizing microbial populations. *J. Vis. Exp.* 116, e54752

Hölscher, T., Kovács, A.T., 2017. Sliding on the surface: bacterial spreading without an active motor. Environ. Microbiol. 19, 2537-2545.

Huerta-Cepas, J., Serra, F., Bork, P., 2016. ETE 3: Reconstruction, analysis, and visualization of phylogenomic data. Mol. Biol. Evol. 33,1635-1638.

Huson, D.H., Scornavacca, C., 2012. Dendroscope 3: an interactive tool for rooted phylogenetic trees and networks. Syst Biol 61, 1061-1067.

Shafi, J., Tian, H., Ji, M.S., 2017. Bacillus species as versatile weapons for plant pathogens: a review. Biotechnology & Biotechnological Equipment 31: 14.

Jones, S.E., Paynich, M.L., Kearns, D.B., Knight, K.L. 2014. Protection from intestinal inflammation by bacterial exopolysaccharides. J Immunol 192, 4813-4820.

Kanafani, Z.A., Corey, G.R. 2007. Daptomycin: a rapidly bactericidal lipopeptide for the treatment of Gram-positive infections. Expert Rev Anti Infect Ther 5, 177-184.

Kearns, D.B. 2010. A field guide to bacterial swarming motility. Nat. Rev. Microbiol. 8, 634-644.

Kovacs, A.T., van Gestel J., Kuipers, O.P., 2012. The protective layer of biofilm: a repellent function for a new class of amphiphilic proteins." Mol. Microbiol. 85, 8-11.

Meena KR, Kanwar, K.S. 2015. Lipopeptides as the antifungal and antibacterial agents: applications in food safety and therapeutics. Biomed Res Int 473050.

Meier-Kolthoff, J.P., Auch, A.F., Klenk, H.P., Goker, M., 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. BMC Bioinformatics 14, 60.

Meier-Kolthoff, J.P., Hahnke, R.L., Petersen, J., Scheuner, C., Michael, V., Fiebig, A., Rohde, C., Rohde, M., Fartmann, B., Goodwin, L.A., Chertkov, O., Reddy, T., Pati, A., Ivanova, N.N., Markowitz, V., Kyrpides, N.C., Woyke, T., Goker, M., Klenk, H.P., 2014. Complete genome sequence of DSM 30083(T), the type strain (U5/41(T)) of *Escherichia coli*, and a proposal for delineating subspecies in microbial taxonomy. Stand. Genomic Sci. 9, 2.

Navarro-Muñoz, J., Selem-Mojica, N., Mullowney, M., Kautsar, S., Tryon, J., Parkinson, E., De Los Santos, E., Yeong, M., Cruz-Morales, P., Abubucker, S., Roeters, A., Lokhorst, W., Fernandez-Guerra, A., Teresa Dias Cappelini, L., Thomson, R., Metcalf, W., Kelleher, N., Barona-Gomez, F., Medema, M.H., 2018. A computational framework for systematic exploration of biosynthetic diversity from large-scale genomic data. bioRxiv: 445270.

Ongena, M., Jacques, P., 2008. Bacillus lipopeptides: versatile weapons for plant disease biocontrol. Trends Microbiol 16, 115-125.

Parks, D. H., Chuvochina, M., Waite, D.W., Rinke, C., Skarshewski, A., Chaumeil, P.A., Hugenholtz, P., 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat. Biotechnol. 36, 996-1004.

Konstantinou, S., Karaoglanidis G. S. K., Bardas, G.A., 2011. Postharvest fruit rots of apple in Greece: pathogen incidence and relationships between fruit quality parameters, cultivar susceptibility, and patulin production. Plant Disease 95, 7.

Rahman,S.F.S.A , Eugenie Singh, E., Pieterse, C.M.J., Schenk, P. M. Emerging microbial biocontrol strategies for plant pathogens Plant Sci. 2018, 2, 102–111.

Stefanic, P., Mandic-Mulec, I., 2009. Social interactions and distribution of *Bacillus subtilis* pherotypes at microscale. J. Bacteriol. 191, 1756-1764.

Verschuere, L., Rombaut, G., Sorgeloos, P., Verstraete, W., 2000. Probiotic bacteria as biological control agents in aquaculture. Microbiol. Mol. Biol. Rev. 64, 655-671.

Weber, T., Blin, K., Duddela, S., Krug, D., Kim, H.U., Bruccoleri, R., Lee, S.Y., Fischbach, M.A., Muller, R., Wohlleben, W., Breitling, R., Takano, E., Medema, M.H., 2015. antiSMASH 3.0-a comprehensive resource for the genome mining of biosynthetic gene clusters. Nucleic Acids Res 43, W237-243.

Zeriouh, H., Romero, D., Garcia-Gutierrez, L., Cazorla, F.M., de Vicente, A., Perez-Garcia, A., 2011. The iturin-like lipopeptides are essential components in the biological control arsenal of Bacillus subtilis against bacterial diseases of cucurbits. Mol Plant Microbe Interact 24,1540-1552.