Interplay of primary bovine lymphocytes and *Mycobacterium avium* subsp. *paratuberculosis* shows distinctly different proteome changes and immune pathways in host-pathogen interaction

Kristina J.H. Kleinwort [1], Stefanie M. Hauck [2], Roxane L. Degroote [1], Karina Lutterberg [1], Armin M. Scholz [3], Christina Hölzel [4,5], Erwin P. Märtlbauer [4], Cornelia A. Deeg [1]*

1 Chair of Animal Physiology, Department of Veterinary Sciences, LMU Munich, Veterinärstraße 13, D-80539 Munich, Germany; E-Mail: K.Kleinwort@tiph.vetmed.uni-muenchen.de, r.degroote@tiph.vetmed.uni-muenchen.de, k.lutterberg@tiph.vetmed.uni-muenchen.de, Cornelia.Deeg@lmu.de

2 Research Unit for Protein Science, Helmholtz Zentrum Munich, German Research Center for Environmental Health GmbH, Munich, Germany; E-Mail: hauck@helmholtz-muenchen.de

3 Livestock Center of the Faculty of Veterinary Medicine, LMU Munich, Oberschleißheim, Germany; E-Mail: Armin.scholz@lvg.vet.med.uni-muenchen.de

4 Chair of Hygiene and Technology of Milk, Department of Veterinary Sciences, LMU Munich, Schönleutnerstr 8, D-85764 Oberschleißheim, Germany; E-Mail: E.Maertlbauer@mh.vetmed.uni-muenchen.de

5 Institute of Animal Breeding and Husbandry, Faculty of Agricultural and Nutritional Sciences, CAU Kiel, Hermann-Rodewald-Str. 6, 24098 Kiel, Germany; Email: choelzel@tierzucht.uni-kiel.de

* Author to whom correspondence should be addressed; Cornelia A. Deeg, Chair for Animal Physiology, Department of Veterinary Sciences, LMU Munich, Veterinärstr. 13, D-80539 Munich. E-mail: Cornelia.Deeg@lmu.de, Phone: + 49-89–2180-2551; Fax: +49-89-2180-2554.

Abstract: Mycobacterium avium subsp. paratuberculosis (MAP) is a pathogen causing paratuberculosis in cattle and small ruminants. During the long asymptomatic subclinical stage, high numbers of MAP are excreted and can be transmitted to food, where they survive many of the standard techniques of food decontamination. If these MAP are harmful to the consumers is currently under debate. In general, there is a lack of information regarding interaction of the hosts immune system with MAP.

In this study, we tested the interaction of peripheral blood lymphocytes (PBL) from cattle
with MAP in their exoproteomes/secretomes. Because in other mycobacterial infections,
the immune phenotype correlates with susceptibility, we additionally tested the
interaction of MAP with recently detected immune deviant cows.

In PBL, different biological pathways were enhanced in response to MAP dependent on
the immune phenotype of the host. PBL of control cows activated members of cell
activation and chemotaxis of leukocytes pathway as well as IL-12 mediated signaling. In
contrast, in ID cows CNOT1 was detected as highly abundant protein, pointing to a
different immune response, which could be favorable for MAP. Additionally, MAP
reacted different to the hosts. Their exoproteomes differed in either GroEL1 or DnaK
abundance, depending on the interacting immune response.

These findings point to an interdependent, tightly regulated response of MAP and the
immune system.

Keywords: Mycobacterium avium subsp. paratuberculosis, exoproteome, immune
capacity, IL12, CNOT, GroEL1, DnaK, ShinyGO
48

1. Introduction

Mycobacterium avium subsp. *paratuberculosis* (MAP) is a critical pathogen for cattle and
small ruminants, causing paratuberculosis with decreased milk production and, in some
animals, excessive loss of weight [1]. Paratuberculosis, also referred to as Johne's disease, is
world-wide endemic; no country or region has been found to be free of this disease [2].
Affected ruminants go through a long asymptomatic subclinical phase in which infection
cannot reliably be detected by standard diagnostic tests [3,4]. These subclinical infected
animals can already shed MAP, thereby contaminating dairy products or meat [5].
Diseased animals were shown to shed high numbers of MAP [6,7]. Since viable MAP were
found in pasteurized milk [8] and in dried dairy products like powdered infant formula [9]
and in raw fermented sausages [10], MAP are discussed as foodborne pathogens.
A similar pathology in the intestinal tissue of patients with intestinal tuberculosis and
paratuberculosis was described more than a century ago. Recently, an association between
MAP and Crohn's disease was shown, initiating a discussion about a possible relationship
of MAP in Crohn's pathogenesis [11]. Johne's and Crohn's disease share clinical and
histopathological similarities, MAP can survive standard pasteurization procedures and
MAP antibodies can be detected in Crohn's patients, where macrolide antibiotics
ameliorate disease [12]. In contrast, genotypes of MAP isolated from cattle and man are
different, there is a lack of evidence for uptake of contaminated food in respective patients
and MAP cannot consistently be isolated from Crohn's disease patients [13]. Neither in
cattle, nor in humans MAP is transmitted from mother to child during pregnancy. But

although MAP was detected widespread in many farms and different countries, the
incidence of Johne`s disease in ruminants is marginal [14]. Bacteria can survive for 2–10
years without causing obvious symptoms of infection in cows [15]. As seen in cattle farms,
susceptibility to MAP infection differs in human populations [16]. This points to a complex
disease in which several pathogens, environmental factors and an inappropriate immune
response in genetically susceptible hosts participate in the cause of disease [16]. Since an
enhanced susceptibility of the host contributes to pathogenesis in other mycobacteria
associated diseases (e.g. tuberculosis) [17], we wanted to gain further information about the
interplay of MAP with the immune system in hosts with different immune capacities. This
is also of interest because in cattle, the MAP eradication programmes that are solely based
on hygiene management are not very successful [18]. This could indicate certain reservoir
cows that host and spread MAP without developing any clinical signs.

Only little is known about the host-pathogen interaction of MAP and the immune
system of its hosts [19]. Functional differences in these responses could lead to aberrant
reactions in susceptible hosts [19].

Recently, we detected a functionally different immune capacity in 22% of cows from
different herds in Germany using differential proteome analyses [20]. These immune
deviant (ID) cows differ in their constitutive immune proteome and they regulate different
master immune regulators upon polyclonal immune cell stimulation. The phenotype is
functionally correlated with an increased prevalence of mastitis, indicating an impact on
the ability to fight infections [20].

All living microorganisms are exposed to changing environmental parameters that define
their habitats. Bacteria sense environmental changes and react to it with various stress
response mechanisms [21]. To gain information about the pathogenic mechanisms of MAP
and how they respond to different immune response signals of their hosts, we analysed the
changes in the proteomes of MAP after co-incubation with primary peripheral blood
derived leukocytes (PBL) of control and ID cows. Since the aim of this study was a better
understanding of the host-pathogen response, we analysed the exoproteomes of MAP and
the bovine peripheral blood derived lymphocytes.

The term exoproteome describes the protein content that can be found in the extracellular
proximity of a given biological system [22]. These proteins arise from cellular secretion,
other protein export mechanisms or cell lysis, but only the most stable proteins in this
environment will remain abundant [22]. These proteins play roles in the organism's
survival in extreme habitats such as saline environments [23]. Investigating the
exoproteome of the pathogen and the secretome of PBL provides expanded coverage of the
repertoire of proteins secreted in the stress response of MAP to different immune responses
and this is essential for understanding respective mechanisms. Accordingly, this study
aimed at providing a better understanding of the interplay of *Mycobacterium avium* subsp.

paratuberculosis (MAP) with the immune system on proteome level to identify the complex
network of proteins involved in host-foodborne bacteria communication.
110

2. Results

2.1. Differentially abundant proteins in secretomes and exoproteomes of PBL from control and
immune deviant hosts as response to in vitro infection with MAP

We investigated the immune response of primary PBL isolated from control and immune
deviant animals. After 48 hours of *in vitro* infection with live MAP, we harvested the mixed
secretomes and exoproteomes of PBL and MAP and analysed them with mass
spectrometry. Overall, we identified 826 proteins (811 bovine and 15 MAP proteins).
Cluster analysis confirmed significant differences in protein abundances between PBL of
control and ID cow in host-pathogen response (Figure 1, Venn diagramm). In secretome of
control cow, 90 proteins were differentially upregulated (Figure 1, ≥ 5-fold change of
expression, blue circle) in contrast to 38 proteins that were higher abundant in ID cow
(Figure 1, light red circle).

In the MAP exoproteome, although small in overall numbers, different protein abundances
were also detectable. In MAP co-incubated with control PBL, one protein (GroEl1) was
found with protein abundance change (Figure 2, blue circle; six-fold higher abundance) and
one protein (DnaK) was upregulated in exoproteome after co-incubation with ID PBL
(Figure 2, light red circle; 24-fold higher abundance).

2.2. Different biological pathways were regulated in the immune response in host PBL

Interpretation of large protein sets can be performed through enrichment analyses, using
published information for examination of overrepresentation of a known set of genes
within the input gene list [24,25]. Since many gene ontology (GO) terms are related or
redundant, we used a hierarchical clustering tree and network (ShinyGO v.050 [25]).

The top regulated biological process pathways in the secretome/excretome of the control
were all related to RNA splicing (Figure 3A). The top regulated immune pathways were
cellular immune responses of various leukocyte subsets, chemotaxis and the reaction to
IL-12 (Figure 3A).

In contrast, in secretome/excretome of ID PBL, the top regulated pathways were response
to stress and immune system process (Figure 3B). The enriched immune pathways
comprised distinct routes of immune response in ID PBL, focussing on closely related

positive regulations of cellular functions, complement activation and humoral immune
response (Figure 3B).

Visualization of network clearly indicates two major regulated networks in control PBL`s
secretome (Figure 4A) and one distinct, major enriched network in ID proteins (Figure 4B).

2.3. GroEl1 and DnAK were differentially abundant in MAPs interacting with different hosts

Interestingly, there was also a clear difference in the regulated proteins of the pathogen
itself, dependent on the co-cultivated host PBL (Table 1, supplementary table 2), although
the low number of proteins was not sufficient for in depth enrichment analyses of the MAP
exoproteomes. These differential proteome analyses clearly indicate substantial differences
on protein level of the foodborne pathogen MAP in the interaction with different host
immune capacities.

152 3. Discussion

Mycobacterium avium subsp. *paratuberculosis* (MAP) could be a foodborne pathogen, because
it is discussed in association with several diseases [26-28]. MAP were found in patients with
inflammatory bowel diseases like Crohn's disease and ulcerative colitis, as well as
autoimmune diseases like type 1 diabetes, multiple sclerosis, rheumatoid arthritis and
Hashimoto's thyroiditis (Garvey, 2018), but so far, a causal association has not yet been
proven in any of these cases. Since MAP can survive many of the standard techniques of
food decontamination (e.g. pasteurization), they are regularly found alive in pasteurized
milk (Gerrard et al., 2018) and in dried dairy products such as powdered infant formula
(Botsaris et al., 2016). If MAP play a role in pathogenesis of respective diseases, we think
that this must be associated with a certain type of susceptibility from these hosts. MAP is
ubiquitously found and only a minor proportion of consumers (if at all) is affected by
MAP-associated diseases. The same is true for cows: while they are often in contact with
MAP, resulting in high frequencies of seropositive animals – approximately 20% and at
least 3–5% in several countries [29,30] – Johne`s disease incidence is very rare. For example,
a total of 232 clinical cases of Johne`s disease were reported in Ireland from 1995 to 2002
[31], yielding an average annual rate of approximately 0.0005%, given a cattle population of
six million [32]. A study examining environmental samples from 362 dairy farms located in
all 10 provinces of Canada for detection of MAP by culture revealed true prevalence
estimates of 66% for farms in Western Canada, 54% in Ontario, 24% in Québec, and 47% in
Atlantic Canada [33].
From tuberculosis it is known that after infection with *Mycobacterium tuberculosis* (MTB),
there is a 10% probability that the host will develop active tuberculosis and the bacterium
may invade multiple organs [34]. Although 9 million new cases of active tuberculosis are

still reported annually, an estimated one-third of the world is infected with MTB while
remaining asymptomatic, defined as latent TB [35]. Among the individuals with latent TB,
only 5–10% will develop active tuberculosis disease in course of their lifetime, because they
effectively control the infection through their immune response [35]. This immune
response after MTB infection is highly complex as the bacteria have intricate immune
escape mechanisms [34].
For MAP, only little is known about host-pathogen reactions in general, and whether
different immune responses exist, but a thorough examination of the respective
mechanisms is of major importance to get functional data that allow a better understanding
and a substantiated risk assessment. In the post-genomic era, proteomics represents a key
discipline to perform in depth studies and identify the complex network of proteins
involved in such host-bacteria communication. In this study, we analysed the
exoproteomes/secretomes of MAP and host cow PBL with known, functionally different
immune capacities [20] using differential proteome analyses after *in vitro* infection.
Interestingly, there were significant differences in protein abundances secreted from
control and ID PBL. Ninety proteins were ≥ 5 fold higher abundant in control PBL after
interaction with MAP. In the control, the top regulated immune pathways described cell
activation and chemotaxis of leukocytes as well as IL-12 mediated signalling pathways. For
M. paratuberculosis it was shown, that IL-12 transcription is increased in infected bovine
macrophages within 6 hours [36], probably to enhance the developing T cell response [37].
To our knowledge, the role of IL-12 in MAP infections of cows has not been analysed so far,
but the IL-12 associated immune response should also be protective. In tuberculosis it was
shown that the induction of protective IFN-γ T cell responses against primary *M.*
tuberculosis infection clearly depends on IL-12 [38]. Mice lacking IL-12p40 cannot control
the growth of the MTB bacterial infection [39]. These findings in mouse models were
confirmed in man, where IL-12 was shown to be critical for preventing tuberculosis [40].
Therefore, we hypothesize that secretion of from IL-12indicates a protective immune
response against MAP infection in control PBL. In ID PBL on the other hand, CCR4-NOT
transcription complex, subunit 1 (CNOT1) was detected as highly abundant protein
(supplemental table 1). This novel finding of CNOT1 regulation in the bovine immune
system is very interesting, because CCR4-NOT complex members have recently been
shown to function as regulators ensuring repression of the MHCII locus in human cell lines
[41]. Poor MHC II expression can cause autoimmune or infectious diseases, since MHCII is
indispensable for adequate immune responses [41]. Additionally, CNOT proteins also
contribute to the downregulation of MHC class I gene expression by influencing
transcription and mRNA degradation [42]. Enriched network analyses of secreted proteins
from ID PBL revealed further major different functions of the 38 differentially abundant
proteins (Figure 4B). Members of complement activation pathway were enriched in ID PBL.

We think the regulated candidates in ID PBL merit further investigation in future studies to
clarify whether they indicate an immune response in favour of MAP infection or just
another way to successfully fight mycobacterial infections.

Interestingly, in the MAP exoproteome (although small in overall numbers) different
protein abundances were also detectable. Exoproteomes arise from cellular secretion, other
protein export mechanisms or cell lysis, but only the most stable proteins in this
environment will remain abundant [22]. In MAP co-incubated with control PBL, GroEL1
showed 6-fold higher abundance. GroEL1 belongs to the family of 60 kDa heat shock
proteins, also known as Cpn60s (GroELs) which are components of the essential protein
folding machinery of the cell, but are also dominant antigens in many infectious diseases
[43]. GroEL1 from MAP is highly immunogenic for cows [42]. The exact function of GroEL1
in MAP is not clarified so far, but it is highly similar to the respective protein in MTB
(Rv3417c), where GroEL1 is important for bacterial survival under low aeration by
affecting the expression of genes known for hypoxia response [43]. From co-cultivation of
MAP with ID PBL, on the other hand, DnaK emerged as differentially abundant protein.
DnaK is a HSP70 family chaperone protein with essential function in stress induced protein
refolding and DnaK loss is accompanied by disruption of membrane structure and
increased cell permeability [44]. DnaK is essentially required for cell growth in
mycobacteria due to a lack of redundancy with other chaperone systems [44]. This finding
from MAP – ID PBL co-cultivation points to regulation of important survival mechanisms
in MAP. However, these results must be interpreted with care because our analysed MAP
exoproteome only comprised 15 proteins.

4. Materials and Methods

4.1. Mycobacterium avium subsp. paratuberculosis (MAP)
The bacterial strain used in these experiments was *Mycobacterium avium* subsp.
paratuberculosis (DSM 44133), purchased from German Collection of Microorganisms and
Cell Cultures (DSMZ, Braunschweig, Germany). MAP were grown on Herrold`s egg yolk
agar (HEYM) (BD Biosciences, Heidelberg, Germany) for four weeks prior to harvesting for
the co-incubation experiment. MAP were yielded through rinsing the agar of the
cultivation tubes with phosphate buffered saline (PBS) and gentle scratching.

4.2. PBL isolation of control and immune deviant cows
Blood samples of cows were collected in tubes supplemented with 25.000 I.U. heparin.
Blood was then diluted 1:2 with PBS pH 7.2 and subsequently layered on density gradient
separating solution (Pancoll; PanBiotech, Aidenbach, Germany). After density gradient
centrifugation (room temperature, 290 × g, 25 min, brake off), PBL were obtained from
intermediate phase. Cells were then washed 3x in PBS (4°C). Withdrawal of blood was
permitted by the local authority Regierung von Oberbayern, Munich, permit no.

55.2-1-54-2532.3-22-12. For determination of control or ID status, PBL were tested in *in vitro*
proliferation assays as described [20]. Respective animals were tested at least 11 times,
before being assigned to control or ID status. The animals were from a MAP-free farm and
were tested negative for MAP antibodies and no MAP were detected in culture from feces.

4.3. Co-cultivation of MAP with primary bovine PBL

Primary bovine PBL ($2x10^7$) of control and ID cows were cultivated in 6 well plates in 3 ml
RPMI each. 15 μg of live MAP were added for 48 h to one well, control wells were not
infected. After 48 h, three technical replicates per experiment were centrifuged for 10 min at
350 x g and the supernatants were filtered through 0,5 μm filters (= exoproteome) and
stored at -20°C until filter aided sample preparation (FASP). Cell pellets were lysed and
total protein content measured with Bradford assay.

4.4. Proteolysis and LC-MS/MS mass spectrometry

10μg total protein was digested with LysC and trypsin by filter-aided sample preparation
(FASP) as described [45]. Acidified eluted peptides were analysed in the data-dependent
mode on a Q Exactive HF mass spectrometer (Thermo Fisher Scientific, Bremen, Germany)
online coupled to a UItimate 3000 RSLC nano-HPLC (Dionex). Samples were automatically
injected and loaded onto the C18 trap column and after 5 min eluted and separated on the
C18 analytical column (75μm IDx15cm, Acclaim PepMAP 100 C18. 100Å/size, LC Packings,
Thermo Fisher Scientific, Bremen, Germany) by a 90min non-linear acetonitrile gradient at
a flow rate of 250 nl/min. MS spectra were recorded at a resolution of 60000 and after each
MS1 cycle, the 10 most abundant peptide ions were selected for fragmentation.

4.5. Protein identification and label-free quantification

Acquired MS spectra were imported into Progenesis software (version 2.5 Nonlinear
Dynamics, Waters) and analyzed as previously described [6,46,47]. After alignment, peak
picking, exclusion of features with charge state of 1 and > 7 and normalization, spectra were
exported as Mascot Generic files and searched against a database containing all entries of
Mycobacterium avium subspecies *paratuberculosis* from NCBI Protein database combined
with the Ensembl bovine database (version 80) with Mascot (Matrix Science, Version 2.5.1).
Search parameters used were 10 ppm peptide mass tolerance, 20 mmu fragment mass
tolerance, one missed cleavage allowed, carbamidomethylation was set as fixed
modification, and methionine oxidation and deamidation of asparagine and glutamine as
variable modifications. Mascot integrated decoy database search was set to a false
discovery rate (FDR) of 1% when searching was performed on the concatenated mgf files
with a percolator ion score cut-off of 13 and an appropriate significance threshold p.
Peptide assignment was reimported to Progenesis Software. All unique peptides allocated
to a protein were considered for quantification. Proteins with a ratio of at least five-fold in
normalized abundance between control and ID or samples were defined as differentially
expressed.

4.6. Enriched pathway analyses

Abundances of the identified proteins were defined as differentially expressed based on the

threshold of protein abundance ratio of ≥1.5 and their assignement to MAP or bovine

exoproteome. Venn diagram was made with open source tool:

http://bioinformatics.psb.ugent.be/webtools/Venn/. The protein–protein interaction

network of differentially-accumulated proteins was analysed with GeneRanker of

Genomatix Pathway System (GePS) software (version 3.10; Intrexon Bioinformatics GmbH,

Munich, Germany; settings: Orthologous genes from H. sapiens were used for this ranking,

species bovine, analysed were proteins with ≥ 5-fold change). Further bioinformatic

analyses were conducted with open source software ShinyGO v0.50: http://ge-lab.org/go/

[25], P-value cutoff (FDR) was set to 0.05.

5. Conclusions

Our data provide novel information about MAP-leukocyte interaction, adding to a

more comprehensive picture of host-pathogen interactions. Co-incubation of MAP with

cells from animals with different immune capacities led to significant differences in PBL

secretomes and different immunological pathways enhanced in the hosts. In exoproteomes

of respective MAPs, GroEL1 and DnaK were differentially abundant. These analyses gave a

deeper insight into the different responses of host PBL and MAP bacteria. In this study,

several novel proteins were identified with changed abundance in host-pathogen

interaction. These candidates merit further investigations in the future to clarify their

functional role in infection control.

Author Contributions: C.D. conceived and designed the experiments; K.K., S.H., K.L., A.S.

and C.H. performed the experiments; K.K., S.H., R.D., E.M. and C.D. analyzed the data;

C.D. wrote the manuscript. All authors critically read the manuscript and approved the

final version to be published.

Funding: The IGF Project 18388 N of the FEI was supported via AiF within the program for

promoting the Industrial Collective Research (IGF) of the German Ministry of Economic

Affairs and Energy (BMWi), based on a resolution of the German Parliament.

Acknowledgments: The authors would like to thank Barbara Amann for excellent technical

support.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

CCR4   CCR4-NOT transcription complex, subunit 1

FASP   Filter-aided sample preparation

ID   Immune deviant

IL-12   Interleukin 12

HEYM Herrold`s egg yolk agar

MAP   *Mycobacterium avium* subsp. *paratuberculosis*

MTB   *Mycobacterium tuberculosis*

PBL   Peripheral blood derived lymphocytes

PBS   Phosphate buffered saline

References

1.   Yamamoto, T.; Murai, K.; Hayama, Y.; Kobayashi, S.; Nagata, R.; Kawaji, S.; Osaki,
M.; Sakakibara, S.-i.; Tsutsui, T. Evaluation of fecal shedding and antibody response
in dairy cattle infected with paratuberculosis using national surveillance data in
japan. *Preventive Veterinary Medicine* **2018**, *149*, 38-46.
2.   Li, L.; Katani, R.; Schilling, M.; Kapur, V. Molecular epidemiology of
mycobacterium avium subsp. Paratuberculosis on dairy farms. *Annual Review of*
Animal Biosciences **2016**, *4*, 155-176.
3.   Li, L.; Bannantine, J.P.; Campo, J.J.; Randall, A.; Grohn, Y.T.; Katani, R.; Schilling,
M.; Radzio-Basu, J.; Kapur, V. Identification of sero-reactive antigens for the early
diagnosis of johne's disease in cattle. *PLoS One* **2017**, *12*, e0184373.
4.   Hobmaier, B.F.; Lutterberg, K.; Kleinwort, K.J.H.; Mayer, R.; Hirmer, S.; Amann, B.;
Hölzel, C.; Märtlbauer, E.P.; Deeg, C.A. Characterization of plant lectins for their
ability to isolate mycobacterium avium subsp. Paratuberculosis from milk. *Food*
Microbiology **2019**, *82*, 231-239.
5.   Sweeney, R.W. Pathogenesis of paratuberculosis. *Veterinary Clinics of North America:*
Food Animal Practice **2011**, *27*, 537-546.
6.   Kuenstner jt, naser s, chamberlin w, borody t, graham dy, mcnees a, hermon-taylor
j, hermon-taylor a, dow ct, thayer w, et al: The consensus from the mycobacterium
avium ssp. Paratuberculosis (map) conference 2017. Front public health 2017;5:208.
7.   Machado, G.; Kanankege, K.; Schumann, V.; Wells, S.; Perez, A.; Alvarez, J.
Identifying individual animal factors associated with mycobacterium avium subsp.
Paratuberculosis (map) milk elisa positivity in dairy cattle in the midwest region of
the united states. *BMC Veterinary Research* **2018**, *14*, 28.
8.   Gerrard, Z.E.; Swift, B.M.C.; Botsaris, G.; Davidson, R.S.; Hutchings, M.R.; Huxley,
J.N.; Rees, C.E.D. Survival of mycobacterium avium subspecies paratuberculosis in
retail pasteurised milk. *Food Microbiology* **2018**, *74*, 57-63.
9.   Botsaris, G.; Swift, B.M.C.; Slana, I.; Liapi, M.; Christodoulou, M.; Hatzitofi, M.;
Christodoulou, V.; Rees, C.E.D. Detection of viable mycobacterium avium
subspecies paratuberculosis in powdered infant formula by phage-pcr and
confirmed by culture. *International Journal of Food Microbiology* **2016**, *216*, 91-94.
10.   Lorencova, A.; Babak, V.; Kralova, A.; Borilova, G. Survival of mycobacterium
avium subsp. Paratuberculosis in raw fermented sausages during production and
storage. *Meat Science* **2019**, *155*, 20-26.

11. Alcedo, K.P.; Thanigachalam, S.; Naser, S.A. Rhb-104 triple antibiotics combination
in culture is bactericidal and should be effective for treatment of crohn's disease
associated with mycobacterium paratuberculosis. *Gut pathogens* **2016**, *8*, 32-32.
12. Kuenstner, J.T.; Naser, S.; Chamberlin, W.; Borody, T.; Graham, D.Y.; McNees, A.;
Hermon-Taylor, J.; Hermon-Taylor, A.; Dow, C.T.; Thayer, W., *et al.* The consensus
from the mycobacterium avium ssp. Paratuberculosis (map) conference 2017.
Frontiers in public health **2017**, *5*, 208-208.
13. Mendoza, J.L.; Lana, R.; Díaz-Rubio, M. Mycobacterium avium subspecies
paratuberculosis and its relationship with crohn's disease. *World journal of*
gastroenterology **2009**, *15*, 417-422.
14. Sergeant, E.S.G.; McAloon, C.G.; Tratalos, J.A.; Citer, L.R.; Graham, D.A.; More, S.J.
Evaluation of national surveillance methods for detection of irish dairy herds
infected with mycobacterium avium ssp. paratuberculosis.
Journal of Dairy Science **2019**, *102*, 2525-2538.
15. Hermon-Taylor, J. Mycobacterium avium subspecies paratuberculosis in the
causation of crohn's disease. *World journal of gastroenterology* **2000**, *6*, 630-632.
16. Eslami, M.; Shafiei, M.; Ghasemian, A.; Valizadeh, S.; Al-Marzoqi, A.H.; Shokouhi
Mostafavi, S.K.; Nojoomi, F.; Mirforughi, S.A. Mycobacterium avium
paratuberculosis and mycobacterium avium complex and related subspecies as
causative agents of zoonotic and occupational diseases. *Journal of Cellular Physiology*
2019, *0*.
17. Scriba, T.J.; Coussens, A.K.; Fletcher, H.A. Human immunology of tuberculosis.
Microbiology Spectrum **2017**, *5*.
18. McAloon, C.G.; Roche, S.; Ritter, C.; Barkema, H.W.; Whyte, P.; More, S.J.; O'Grady,
L.; Green, M.J.; Doherty, M.L. A review of paratuberculosis in dairy herds — part 2:
On-farm control. *The Veterinary Journal* **2019**, *246*, 54-58.
19. Davis, W.C. On deaf ears, mycobacterium avium paratuberculosis in pathogenesis
crohn's and other diseases. *World journal of gastroenterology* **2015**, *21*, 13411-13417.
20. Lutterberg, K.; Kleinwort, K.J.H.; Hobmaier, B.F.; Hauck, S.M.; Nüske, S.; Scholz,
A.M.; Deeg, C.A. A functionally different immune phenotype in cattle is associated
with higher mastitis incidence. *Frontiers in immunology* **2018**, *9*, 2884-2884.
21. Guo, M.S.; Gross, C.A. Stress-induced remodeling of the bacterial proteome. *Current*
biology : CB **2014**, *24*, R424-R434.
22. Armengaud, J.; Christie-Oleza, J.A.; Clair, G.; Malard, V.; Duport, C. Exoproteomics:
Exploring the world around biological systems. *Expert Review of Proteomics* **2012**, *9*,
561-575.
23. Rubiano-Labrador, C.; Bland, C.; Miotello, G.; Armengaud, J.; Baena, S. Salt stress
induced changes in the exoproteome of the halotolerant bacterium tistlia
consotensis deciphered by proteogenomics. *PloS one* **2015**, *10*, e0135065-e0135065.
24. Merico, D.; Isserlin, R.; Stueker, O.; Emili, A.; Bader, G.D. Enrichment map: A
network-based method for gene-set enrichment visualization and interpretation.
PLOS ONE **2010**, *5*, e13984.
25. Ge, S.; Jung, D. Shinygo: A graphical enrichment tool for animals and plants. *bioRxiv*
2018, 315150.

26. Cossu, D.; Yokoyama, K.; Sakanishi, T.; Momotani, E.; Hattori, N. Adjuvant and
411   antigenic properties of mycobacterium avium subsp.
412   paratuberculosis on experimental autoimmune encephalomyelitis.
413   *Journal of Neuroimmunology* **2019**.

414   27. Pierce, E.S. Could mycobacterium avium subspecies paratuberculosis cause crohn's
415   disease, ulcerative colitis…and colorectal cancer? *Infectious Agents and Cancer* **2018**,
416   *13*, 1.

417   28. Sechi, L.A.; Dow, C.T. Mycobacterium avium ss. Paratuberculosis zoonosis – the
418   hundred year war – beyond crohn's disease. *Frontiers in Immunology* **2015**, *6*.

419   29. Boelaert, F.; Walravens, K.; Biront, P.; Vermeersch, J.P.; Berkvens, D.; Godfroid, J.
420   Prevalence of paratuberculosis (johne's disease) in the belgian cattle population.
421   *Veterinary Microbiology* **2000**, *77*, 269-281.

422   30. Nielsen, S.S.; Toft, N. A review of prevalences of paratuberculosis in farmed
423   animals in europe. *Preventive Veterinary Medicine* **2009**, *88*, 1-14.

424   31. Kennedy, A.E.; Da Silva, A.T.; Byrne, N.; Govender, R.; MacSharry, J.; O'Mahony, J.;
425   Sayers, R.G. The single intradermal cervical comparative test interferes with johne's
426   disease elisa diagnostics. *Frontiers in immunology* **2014**, *5*, 564-564.

427   32. Maher, P.; Good, M.; More, S. Trends in cow numbers and culling rate in the irish
428   cattle population, 2003 to 2006. *Irish veterinary journal* **2008**, *61*, 455-463.

429   33. Corbett, C.S.; Naqvi, S.A.; Bauman, C.A.; De Buck, J.; Orsel, K.; Uehlinger, F.;
430   Kelton, D.F.; Barkema, H.W. Prevalence of mycobacterium avium ssp.
431   paratuberculosis infections in canadian dairy herds. *Journal of Dairy*
432   *Science* **2018**, *101*, 11218-11228.

433   34. Zhai, W.; Wu, F.; Zhang, Y.; Fu, Y.; Liu, Z. The immune escape mechanisms of
434   mycobacterium tuberculosis. *International journal of molecular sciences* **2019**, *20*, 340.

435   35. Dye, C.; Scheele, S.; Dolin, P.; Pathania, V.; Raviglione, M.C.; for the, W.H.O.G.S.;
436   Monitoring, P. Global burden of tuberculosis: Estimated incidence, prevalence, and
437   mortality by country. *JAMA* **1999**, *282*, 677-686.

438   36. Weiss, D.J.; Evanson, O.A.; Moritz, A.; Deng, M.Q.; Abrahamsen, M.S. Differential
439   responses of bovine macrophages to mycobacterium avium subsp. Paratuberculosis
440   and mycobacterium avium subsp. Avium. *Infection and immunity* **2002**, *70*,
441   5556-5561.

442   37. Bannantine, J.P.; Stabel, J.R.; Laws, E.; D Cardieri, M.C.; Souza, C.D. Mycobacterium
443   avium subspecies paratuberculosis recombinant proteins modulate
444   antimycobacterial functions of bovine macrophages. *PloS one* **2015**, *10*,
445   e0128966-e0128966.

446   38. Khader, S.A.; Partida-Sanchez, S.; Bell, G.; Jelley-Gibbs, D.M.; Swain, S.; Pearl, J.E.;
447   Ghilardi, N.; Desauvage, F.J.; Lund, F.E.; Cooper, A.M. Interleukin 12p40 is required
448   for dendritic cell migration and t cell priming after mycobacterium tuberculosis
449   infection. *The Journal of experimental medicine* **2006**, *203*, 1805-1815.

450   39. Cooper, A.M.; Magram, J.; Ferrante, J.; Orme, I.M. Interleukin 12 (il-12) is crucial to
451   the development of protective immunity in mice intravenously infected with
452   mycobacterium tuberculosis. *The Journal of experimental medicine* **1997**, *186*, 39-45.

40.   Alcaïs, A.; Fieschi, C.; Abel, L.; Casanova, J.-L. Tuberculosis in children and adults:
Two distinct genetic diseases. *The Journal of experimental medicine* **2005**, *202*,
1617-1621.
41.   Rodríguez-Gil, A.; Ritter, O.; Saul, V.V.; Wilhelm, J.; Yang, C.-Y.; Grosschedl, R.;
Imai, Y.; Kuba, K.; Kracht, M.; Schmitz, M.L. The ccr4-not complex contributes to
repression of major histocompatibility complex class ii transcription. *Scientific*
reports **2017**, *7*, 3547-3547.
42.   Yang, C.-Y.; Ramamoorthy, S.; Boller, S.; Rosenbaum, M.; Rodriguez Gil, A.; Mittler,
G.; Imai, Y.; Kuba, K.; Grosschedl, R. Interaction of ccr4–not with ebf1 regulates
gene-specific transcription and mrna stability in b lymphopoiesis. *Genes &*
Development **2016**, *30*, 2310-2324.
43.   Sharma, A.; Rustad, T.; Mahajan, G.; Kumar, A.; Rao, K.V.S.; Banerjee, S.; Sherman,
D.R.; Mande, S.C. Towards understanding the biological function of the unusual
chaperonin cpn60.1 (groel1) of mycobacterium tuberculosis. *Tuberculosis* **2016**, *97*,
137-146.
44.   Fay, A.; Glickman, M.S. An essential nonredundant role for mycobacterial dnak in
native protein folding. *PLoS genetics* **2014**, *10*, e1004516-e1004516.
45.   Grosche, A.; Hauser, A.; Lepper, M.F.; Mayo, R.; von Toerne, C.; Merl-Pham, J.;
Hauck, S.M. The proteome of native adult müller glial cells from murine retina.
Molecular & cellular proteomics : MCP **2016**, *15*, 462-480.
46.   Hauck, S.M.; Hofmaier, F.; Dietter, J.; Swadzba, M.E.; Blindert, M.; Amann, B.;
Behler, J.; Kremmer, E.; Ueffing, M.; Deeg, C.A. Label-free lc-msms analysis of
vitreous from autoimmune uveitis reveals a significant decrease in secreted wnt
signalling inhibitors dkk3 and sfrp2. *J Proteomics* **2012**, *75*, 4545-4554.
47.   Hauck, S.M.; Lepper, M.F.; Hertl, M.; Sekundo, W.; Deeg, C.A. Proteome dynamics
in biobanked horse peripheral blood derived lymphocyte (pbl) with induced
autoimmune uveitis. *PROTEOMICS* **2017**, *17*, 1700013.
480
481

Captions for Figures

Figure 1: Overlap (Venn diagram) of differentially (≥ 5 fold) expressed proteins between
secretomes/exoproteomes of control cow (blue) and ID cow (red). From a total of 811
identified proteins, 90 were higher abundant in control and 38 in ID.

Figure 2: Overlap (Venn diagram) of differentially (≥ 5 fold) expressed proteins between
exoproteomes of MAP co-incubated with PBL of control cow (blue) and MAP co-incubated
with PBL of ID cow (red). From a total of 15 identified proteins, one different protein from
each specimen showed higher abundance.

Figure 3: Hierarchical clustering tree and network of related GO terms of differentially
expressed proteins in secretomes/exoproteomes of control PBL (A) and ID PBL (B)
illustrates marked differences. GO terms are grouped together based on how many genes
they share. The size of the solid circle corresponds to the enrichment false discovery rate.

Figure 4: Visualization of overlapping relationships among enriched gene-sets revealed A)
two major networks as shown by network view for enriched GO molecular component
terms in control PBL after MAP infection and B) in ID PBL after MAP infection. Related
GO terms are connected by a line whose thickness reflects percent of overlapping genes.
Size of the nodes corresponds to number of genes.

Tables

Table 1: Regulation of MAP proteins identified in secretomes/exoproteomes after
co-incubation with control and ID PBL.

	Accession	Gene symbol	Description	Ratio MAP Ctr/ID
1	AAS06815	GroEL1	pep:novel chromosome:GCA_000007865.1:Chromosome:4733338:4734954:1 gene:MAP_4265 transcript:AAS06815 description:""GroEL1""	6,07
2	ETB04387	F0F1 ATP synthase subunit alpha	pep:novel supercontig:GCA_000504785.1:contig000131:9486:11150:1 gene:O979_07265 transcript:ETB04387 description:""F0F1 ATP synthase subunit alpha""	3,26
3	ETB05569	DNA-binding protein	pep:novel supercontig:GCA_000504785.1:contig000072:8572:9225:1 gene:O979_04085 transcript:ETB05569 description:""DNA-binding protein""	2,47
4	AAS06727	RplN	pep:novel chromosome:GCA_000007865.1:Chromosome:4652017:4652385:1 gene:MAP_4177 transcript:AAS06727 description:""RplN""	2,34
5	AAS06486; ELP44387; ETB08817	GroEL2	pep:novel chromosome:GCA_000007865.1:Chromosome:4395922:4397547:1 gene:MAP_3936 transcript:AAS06486 description:""GroEL2""	1,12
6	AAS06693	Tuf	pep:novel chromosome:GCA_000007865.1:Chromosome:4620946:4622136:1 gene:MAP_4143 transcript:AAS06693 description:""Tuf""	0,88
7	ETB02420	enoyl-CoA hydratase	pep:novel supercontig:GCA_000504785.1:contig000238:14619:15410:1 gene:O979_11400 transcript:ETB02420 description:""enoyl-CoA hydratase""	0,86
8	ETB00930	2-isopropylmalate synthase	pep:novel supercontig:GCA_000504845.1:contig000411:16560:18359:-1 gene:O978_19245 transcript:ETB00930 description:""2-isopropylmalate synthase""	0,75

9	AAS06690	RpsL	pep:novel chromosome:GCA_000007865.1:Chromosome:4617754:4 618128:1 gene:MAP_4140 transcript:AAS06690 description:""RpsL""	0,67
10	AAS06681; ETB45918; ETB45924	RpoC	pep:novel chromosome:GCA_000007865.1:Chromosome:4606729:4 610679:1 gene:MAP_4131 transcript:AAS06681 description:""RpoC""	0,62
11	AAS06680	RpoB	pep:novel chromosome:GCA_000007865.1:Chromosome:4603087:4 606683:1 gene:MAP_4130 transcript:AAS06680 description:""RpoB""	0,48
12	AAS02778	ClpC	pep:novel chromosome:GCA_000007865.1:Chromosome:488298:49 0829:1 gene:MAP_0461 transcript:AAS02778 description:""ClpC""	0,46
13	ETB04840	GTP-binding protein YchF	pep:novel supercontig:GCA_000504785.1:contig000104:1736:2809:1 gene:O979_06015 transcript:ETB04840 description:""GTP-binding protein YchF""	0,43
14	ETB04389 AAS04768	ATP synthase subunit beta	pep:novel supercontig:GCA_000504785.1:contig000131:12100:1355 7:1 gene:O979_07275 transcript:ETB04389 description:""ATP synthase subunit beta""	0,26
15	AAS06390	DnaK	pep:novel chromosome:GCA_000007865.1:Chromosome:4295544:4 297415:1 gene:MAP_3840 transcript:AAS06390 description:""DnaK""	0,04

Captions for Supplemental Material

Supplementary table 1: All proteins identified in PBL secretome analyses of co-cultivation of MAP

with PBL from donors with different immune capacities.

Supplementary table 2: All proteins identified in MAP exoproteome analyses of co-cultivation of

MAP with PBL from donors with different immune capacities.

Data Availability Statement

The datasets generated and analyzed for this study are available in supplemental tables 1

and 2.