Adaptation to seasonal reproduction and thermal minima-related factors

drives fine-scale divergence despite gene flow in Atlantic herring populations

Angela P. Fuentes-Pardo[1], Christina Bourne[2], Rabindra Singh[3], Kim Emond[4], Lisa Pinkham[5],

Jenni L. McDermid[6], Leif Andersson[7,8,9], Daniel E. Ruzzante[1]

Affiliations:

[1] Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada.

[2] Fisheries and Oceans Canada, Northwest Atlantic Fisheries Centre, St John's, NL, Canada.

[3] Fisheries and Oceans Canada, St. Andrews Biological Station, St. Andrews, NB, Canada.

[4] Fisheries and Oceans Canada, Maurice Lamontagne Institute, Mont-Joli, QC, Canada.

[5] Department of Marine Resources, West Boothbay Harbor, Maine, USA.

[6] Fisheries and Oceans Canada, Gulf Fisheries Centre, Moncton, NB, Canada.

[7] Science for Life Laboratory, Department of Medical Biochemistry and Microbiology, Uppsala

University, Uppsala, Sweden.

[8] Department of Veterinary Integrative Biosciences, Texas A&M University, College Station,

USA.

[9] Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences,

Uppsala, Sweden

Corresponding authors: Angela P. Fuentes-Pardo (apfuentesp@gmail.com) and Daniel E.

Ruzzante (daniel.ruzzante@dal.ca), Dalhousie University, 1355 Oxford Street, B3H 4R2,

Halifax, Canada, +1 (902) 494 1688.

Abstract:

High connectivity and low potential for local adaptation have been common assumptions for most marine species, given their usual high fecundity and dispersal capabilities. Recent genomic studies however, have disclosed unprecedented levels of population subdivision in what were previously presumed to be panmictic or nearly panmictic species. Here we analyzed neutral and adaptive genetic variation at the whole-genome level in Atlantic herring (*Clupea harengus* L.) spawning aggregations distributed across the reproductive range of the species in North America. We uncovered fine-scale population structure at putatively adaptive loci, despite low genetic differentiation at neutral loci. Our results revealed an intricate pattern of population subdivision involving two overlapping axes of divergence: a temporal axis determined by seasonal reproduction, and a spatial axis defined by a latitudinal cline establishing a steep north-south genetic break. Genetic-environment association analyses indicated that winter sea-surface temperature is the best predictor of the spatial structure observed. Thousands of outlier SNPs distributed along specific parts of the genome spanning numerous candidate genes underlined each pattern of differentiation, forming so-called "genomic regions or islands of divergence". Our results indicate that timing of reproduction and latitudinal spawning location are features under disruptive selection leading to local adaptation in the herring. Our study highlights the importance of preserving functional and neutral intraspecific diversity, and the utility of an integrative seascape genomics approach for disentangling intricate patterns of intraspecific diversity in highly dispersive and abundant marine species.

Keywords: Population genomics, local adaptation, fisheries, management, conservation, pool-seq, whole genome re-sequencing.

Introduction:

Population subdivision and connectivity are important topics in evolutionary and

conservation biology, because they can help elucidate how local adaptation arises (Barrett &

Hoekstra, 2011; Lewontin, 2002) and can guide management plans aiming to protect intraspecific

genetic diversity, a determinant factor for population persistence in changing environments

(Allendorf, Hohenlohe, & Luikart, 2010). Yet, the scarcity of genomic resources for most

species, and the difficulty in determining the relative importance of genetic drift, gene flow, and

selection in shaping contemporary patterns of intraspecific genetic diversity, remain major

challenges (Ravinet et al., 2017). The increased power for assessing neutral and putatively

adaptive genetic variation with next-generation sequencing (NGS) technologies (Nosil & Feder,

2012) is helping to uncover unprecedented levels of genetic structure in what were previously

presumed to be panmictic or nearly panmictic species.

Marine species are outstanding examples of such paradigm shifts, as they have often been

expected and observed to exhibit low levels of population structure and low divergence potential

(Palumbi, 1994), given their high fecundity and dispersal capabilities (Hauser & Carvalho, 2008).

Recent genomic studies revealing fine-scale structuring are challenging this view [e.g., Atlantic

cod (*Gadus morhua*) (Bradbury et al., 2013); Atlantic herring (*Clupea harengus*) (Martinez

Barrio et al., 2016); American lobster (*Homarus americanus*) (Benestan et al., 2015)]. Various

mechanisms by which population structure could arise have been proposed, including:

oceanographic barriers, isolation-by-distance, larval and adult behavior, recent evolutionary

history (e.g. historical vicariance and secondary contact), and natural selection (Palumbi, 1994).

There is great interest in understanding how natural selection can lead to population divergence

and local adaptation, especially under the homogenizing effect of gene flow (Tigano & Friesen,

2016) because of its direct relationship with fitness, population persistence, and evolution.

However, the genetic basis of adaptive traits remains largely unknown (Barrett & Hoekstra,

2011). Genome scans performed with NGS methods are helping to identifying loci associated

with adaptive phenotypes (Jones et al., 2012; Tavares et al., 2018). Such loci typically show

elevated genetic divergence that is interpreted as a signature of selection. Nevertheless,

disentangling genomic signatures of selection from signatures of demographic history has been

limiting (Hoban et al., 2016). Species that are widely distributed are often exposed to diverse

ecological habitats where selection can result in local adaptation (Yeaman & Whitlock, 2011).

Therefore, highly fecund marine species inhabiting heterogeneous environments offer ideal

candidates for the study of ecological adaptation, since in these the effect of genetic drift is

minuscule and the effectiveness of natural selection is greater.

Atlantic herring is an abundant marine schooling pelagic fish that has colonized diverse

environments throughout the North Atlantic, including open ocean and the brackish waters of the

Baltic Sea. These characteristics, together with the increasing availability of genomic resources,

make this species ideal for investigating the genetic basis and mechanisms involved in ecological

adaptation. Juveniles and adults undertake annual migrations between feeding, overwintering,

and spawning areas. Herring matures at 3-4 years of age and can live to 20+ years (Benoît et al.,

2018). Spawning occurs mostly in spring and fall seasons at predictable times and locations near

shore, which suggests strong spawning site fidelity (McQuinn, 1997; Stephenson, Melvin, &

Power, 2009; Wheeler & Winters, 1984). Atlantic herring plays an important role in the marine

ecosystem, feeding on plankton and being preyed upon by numerous marine fish, birds and

mammals. It also sustains large fisheries throughout the North Atlantic (FAO, 2019), some of

which have experienced severe periods of decline and signs of recovery in the last century

(Britten, Dowd, & Worm, 2016; Engelhard & Heino, 2004; Overholtz, 2002; Simmonds, 2007).

The ecological, economic, and cultural importance of herring has therefore motivated research on

this species for more than a century (Stephenson et al., 2009); however, its complex life history

has made the description of its population structure elusive (Iles & Sinclair, 1982).

Numerous studies have examined the population structure of herring using different

genetic tools and at various spatial scales, mostly in the northeast (NE) Atlantic. Such studies

have observed low levels of population differentiation at neutral loci (Andersson, Ryman,

Rosenberg, & Ståhl, 1981; André et al., 2011; Jorgensen, Hansen, Bekkevold, Ruzzante, &

Loeschcke, 2005). The expansion of these studies to the use of thousands of single nucleotide

polymorphisms (SNPs) derived from various genomic techniques have revealed significant

genetic differentiation at putatively adaptive loci in relation to environmental gradients (Guo, Li,

& Merilä, 2016; Lamichhaney et al., 2012; Limborg et al., 2012). Moreover, the recent

development of a high-quality genome assembly for the Atlantic herring allowed the

identification of many millions of SNPs and a breakthrough in the possibility to study the genetic

basis of ecological adaptation in this species (Martinez Barrio et al., 2016). A few studies have

addressed this question in the northwest (NW) Atlantic (Kerr, Fuentes☐Pardo, Kho, McDermid,

& Ruzzante, 2018; Lamichhaney et al., 2017; McPherson, O'Reilly, & Taggart, 2004); while

they provided important insight on population structuring with seasonal reproduction and within

the southern region, and reported temporal stability of genomic divergence between spring and

fall spawners, they were limited by scarce sampling.

In the NW Atlantic, herring spawn from Cape Cod to southern Labrador (Bourne,

Mowbray, Squires, & Koen-Alonso, 2018; Sinclair & Iles, 1989) between April and November,

but spawning peaks in spring and fall. Spring- and fall-spawners are therefore the main spawning

types recognized in the region. The relative abundance of each reproductive strategy varies

geographically: in the north (northern Newfoundland) spring-spawners were historically more

abundant, at mid-range (Gulf of St. Lawrence) both strategies were common, and in the southern

extreme (Bay of Fundy, Scotian Shelf, Gulf of Maine) fall-spawners predominate (Melvin,

Stephenson, & Power, 2009). Changes in the prevalence of these components have been observed

in the last decade; in particular, a significant decline of spring-spawners and a moderate

abundance of fall-spawners in the Gulf of St. Lawrence (McDermid, Swain, Turcotte, Robichaud,

& Surette, 2018) and Newfoundland (Bourne et al., 2018). Such changes have been attributed to

varying elevated fishing mortality, declines in weight-at-age, and environmental conditions

(Melvin et al., 2009), suggesting that the effects of climate change on population persistence of

Atlantic herring are important. The concerning population declines (Britten et al., 2016)

emphasize the need to disentangle the population structure of NW Atlantic herring.

Here, we study neutral and adaptive variation of adult herring collected from 14 spawning

grounds distributed across the species' reproductive range in the NW Atlantic. The two

overarching questions were: *i*) What are the spatial scale and pattern of population structuring in

herring and what is the genetic basis of such structuring, and *ii*) What is the potential functional

effect of variant sites underlying population divergence and which mechanisms and

environmental variables are associated with population structure patterns? We used whole-

genome re-sequencing of pools of individuals [Pool-seq, (Schlötterer, Tobler, Kofler, & Nolte,

2014)] and individual genotyping along with multivariate statistical approaches, machine

learning algorithms, and oceanographic information, to address these questions. Considering the

particular attributes of the NW Atlantic Ocean (DFO, 1997; Townsend, Thomas, Mayer, Thomas,

& Quinlan, 2004) and the importance of environment for shaping population divergence in

herring, we predict that some of the divergent genomic regions exclusively found in Canada may

be strongly associated with local environmental conditions. Our results provide insight into how

population divergence arises in the presence of gene flow via temporal and spatial isolation and

will help inform management and conservation practices.

Materials and Methods:

Sample collection and DNA extraction

Adult herring (N=697) were collected from 14 inshore spawning aggregations distributed

across Atlantic Canada and the Gulf of Maine (Fig. 1A and Table 1). Collections took place

during the local spawning peak in the spring and fall seasons from 2012 to 2016. Sampling

locations correspond to areas with recurrent annual spawning and jointly represent most of the

reproductive range of the species in the NW Atlantic. Because of the presumed spawning site

fidelity and the mixing of populations during the non-spawning seasons, we targeted individuals

in reproductive condition to assess population definition. Individual muscle or fin tissue samples

were preserved in 95% ethanol at -20 ℃ until processing. DNA was isolated from the tissue

samples using a standard phenol chloroform protocol. DNA concentration (in ng/µl) was

measured in triplicates using the Quant-iT PicoGreen dsDNA assay (Thermo Fisher Scientific,

U.S.) and the Roche LightCycler 480 Instrument (Roche Molecular Systems, Inc., Germany).

DNA integrity was verified with 0.8% agarose gel electrophoresis using 0.5x TBE buffer and a

1Kbp molecular weight ladder.

Pool-sequencing and read quality filtering

Genome-wide patterns of genetic variation and population allele frequencies were

assessed for each spawning aggregation using the Pool-seq approach. This method consists of

performing whole-genome sequencing of pools of individuals using a single barcoded library,

which implies that only population level data is recovered (individual genotype information is

lost). In our case, each pool comprised equal amounts of DNA of ~50 individuals collected on the

same spawning ground (the terms spawning aggregation and sampling site will be

interchangeably used hereafter). Individual DNA were normalized to a common concentration

and pooled to a single tube using the liquid handling robot epmotion 5407 (Eppendorf,

Germany). Sequencing library preparation and shotgun sequencing were outsourced. In brief, a

single TruSeq Nano Illumina DNA library was built for each DNA pool (i.e. spawning

aggregation). AMPURE beads were used for fragment size selection, targeting an insert size of

~550 bp. The 14 pooled-DNA libraries were sequenced using paired-end 126-bp reads on an

Illumina Hiseq-2500 sequencer in two batches (5 libraries in 2015, 11 in 2016). Target read depth

of coverage per pool was 40-50x, for an estimated herring genome size of ~850 Mbp (Martinez

Barrio et al., 2016).

Quality of raw sequence reads of each pool was checked using FastQC v0.11.5 (Andrews,

2010), and jointly evaluated for the 14 pools with MultiQC v.1.3 (Ewels, Magnusson, Lundin, &

Käller, 2016). Low quality bases (Phred score <20) and Illumina adapters were trimmed-off the

reads, and reads shorter than 40 bp were removed from the dataset using Trimmomatic v.0.36

(Bolger, Lohse, & Usadel, 2014) [*parameters: ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10*

SLIDINGWINDOW:5:20 MINLEN:40]. High quality paired-reads remaining after filtering were

used for downstream analysis.

Read mapping, SNP calling and filtering

We adapted the Genome Analysis Toolkit (GATK) Best Practices workflow (Van der

Auwera et al., 2013) to variant discovery in Pool-seq data and to our computing infrastructure.

For this we first obtained a stitched version of the herring genome for optimal SNP caller

performance in the computer cluster available. Then, sequence reads of each pool were

independently aligned against the stitched herring genome using the Burrows-Wheeler Aligner

(BWA) v0.7.12-r1039 [*default parameters, MEM algorithm*] (Li, 2013). SNP calling was

performed using GATK v3.8 (McKenna et al., 2010) (see Fig. S1). Lastly, the raw variant calls

were filtered using GATK (Fig. S2), Popoolation2, and custom python scripts (See Supporting

Information for details). In Pool-seq applications, population allele frequencies are derived from

the total read counts supporting a variant site. Read coverage though, can be biased by

sequencing and read mapping artifacts (Dohm, Lottaz, Borodina, & Himmelbauer, 2008;

Kolaczkowski, Kern, Holloway, & Begun, 2011). To control for these factors and minimize their

potential effect on population allele frequency calculation, we applied the allele count correction

proposed by (Feder, Petrov, & Bergland, 2012; Kolaczkowski et al., 2011). Details on the

application of this correction method and population allele frequencies estimation can be found in

the Supporting Information.

Population structure

Based on the population allele frequencies, we examined genetic structure among

spawning aggregations with a Neighbor-Joining (NJ) phylogenetic tree and with pairwise F_{ST}

estimates. We computed pairwise Nei (1972) genetic distance with *Gendist* and built a NJ tree

with *Neighbor*, both programs implemented in the package PHYLIP v3.697 (Baum, 1989).

Bootstrapping was performed using the program *Seqboot* of PHYLIP, and the consensus tree was

visualized with FigTree (Rambaut, 2007). We estimated unbiased F_{ST} for pools ($\hat{F}_{ST}^{pool}$) between

all pairs of spawning aggregations using the R package *poolfstat* (Hivert, 2018). This algorithm

computes *F*-statistics equivalent to Weir & Cockerham (1984) estimates, while accounting for

random sampling of chromosomes that may occur during DNA pooling and sequencing in Pool-

seq applications.

Outlier loci detection and genome-wide patterns of differentiation

To identify loci potentially under selection, we performed genome scans for outlier loci

detection using Principal Component Analysis (PCA), as implemented in the R package *pcadapt*

v.4.0.2 (Luu, Bazin, & Blum, 2017). This algorithm assumes that divergent loci highly correlated

to population structure are likely under selection. Outlier loci are detected based on the

Mahalanobis distance calculated from the correlation coefficients between SNPs and a selected

number of *K* principal components (PCs) (i.e. PCA loadings).

We performed a genome scan for the first 13 PCs (default is *K*=number of pools-1, 14-1= 13)

using a minor allele frequency (MAF) of 0.05. Loci with Benjamini-Hochberg (BH) adjusted *P*-

values ≤0.01 were considered candidates for being under selection. To identify which PCs

explained the greatest proportion of genomic variance, we examined the scree plot generated by

pcadapt, as well as the allele frequency patterns revealed in heatmaps made with the R package

ComplexHeatmap (Gu, Eils, & Schlesner, 2016). The heatmaps depicted population allele

frequencies (standardized to the major allele) of the 200 outlier loci most correlated to each PC

(ranked by *P*-value in ascending order). We further explored the loci driving genomic

differentiation in the herring by performing, with *pcadapt*, component-wise genome scans for the

PCs exhibiting distinctive allele frequency patterns. To examine the distribution of outlier loci

across the herring genome, for each informative PC we obtained Manhattan plots depicting the

genomic position of outlier SNPs and their respective significance association value ($-\log_{10}P$-

value) using the R package *qqman* (S. D. Turner, 2014).

Identification of the most informative outlier loci

We ranked outlier loci based on their importance for classification to each of the

categories (or classes) of distinctive genomic patterns of differentiation in herring. For this we

used random forest (RF), a supervised learning algorithm implemented in the R package

randomForest (Liaw & Wiener, 2002). For the seasonal reproductive pattern, classes

corresponded to spring or fall. For the latitudinal pattern, classes were northern (SIL-S, SPH-S,

NTS-S, LAB-F, BLS-F, NDB-S, NDB-F, TRB-F, MIR-F, BDO-S, SCB-F), intermediate (MUS-

F, GEB-F), and southern (ME4-F) regions. The RF model was based on 50 individual genotypes

per spawning aggregation simulated from population allele frequencies using the R function

sample.geno implemented in *pcadapt* v3.0.4. For the RF runs, the parameter *mtry* was set to

default (equals to sqrt(*p*), where *p* is the number of loci); *ntree* was set to 1,000,000; and

sampsize was set to 2/3 of the class with the lower sample size. From a scatter-plot of importance

values generated by the random forest classifier (Mean Decrease in Accuracy, MDA), loci before

the point where the differences between importance values level-off ("elbow method") were

considered the most important (Goldstein, Hubbard, Cutler, & Barcellos, 2010).

Validation of a subset of outlier SNPs related to seasonal reproduction and to latitudinal

divergence

262   We validated some of the top candidate loci detected with Pool-seq data that showed

strong association with seasonal reproduction and latitudinal divergence with individual

genotypes. For this, we genotyped 240 individuals (30 individuals from 8 locations) in 40 SNPs

related to seasonal reproduction and 90 SNPs related to latitude using the Agena MassARRAY

SNP genotyping platform (Agena Bioscience, Inc.). These SNPs were chosen considering these

criteria: (*i*) top ranked based on importance values (Mean Decrease in Accuracy, MDA) obtained

from the random forest algorithm (as described in the previous section), (*ii*) had ≥150 bp of

flanking sequence for primer design, (*iii*) did not fall within or a few bases away from repetitive

regions and had fewer than 4 flanking SNPs, (iv) when two or more top ranked SNPs were

located within the same scaffold, the ones separated by ≥ 1Kbp were kept, in an attempt to

minimize redundancy in the panel). The application of these filters and the retrieval and

preparation of DNA sequences for primer design for the Agena platform were performed with

custom R scripts. A quality control of raw SNP genotypes was performed using PLINK (Purcell

et al., 2007), in which SNPs and individuals with more than 20% missing data, and SNPs with

minor allele frequency (MAF) lower than 0.01 were removed. We obtained a heatmap plot using

the R function *heatmap.2* of the R package *gplots* for the visual inspection of individual genotype

patterns. File format conversions required for missing data filtering and heatmap plotting were

conducted with PGDSpider (Lischer & Excoffier, 2012) and a custom python script (data was

transformed to PLINK format, then to VCF file format, and finally to 0,1,2 format).

Functional annotation of outlier loci

We investigated the potential effect on gene function of outlier SNPs associated with

seasonal reproduction and the latitudinal cline using SNPeff v4.1l (build 2015-10-03) (Cingolani

et al., 2012) [*default parameters*]. This program determines the position of a SNP with respect to

the constituents of a nearby gene within 5Kbp (i.e. exons, introns, 5'-UTR region, etc.), and

predicts its putative effect on gene and protein composition (i.e. synonymous and missense

mutations, premature stop codon, etc., a complete list of effects is described in the program

documentation). Variants located beyond 5Kbp of a gene were annotated as 'intergenic'. We

based this analysis on the current herring genome assembly and annotations (Martinez Barrio et

al., 2016). Further, we separately examined gene ontology (GO) terms of the genes annotated to

the outlier loci most strongly associated with seasonal reproduction and the latitudinal cline (-

$\log_{10}P$-value ≥ 7, equivalent to P-value $\leq 1\text{x}10^{-7}$, lower threshold commonly used for significant

association in human GWAS, (Fadista, Manning, Florez, & Groop, 2016; Panagiotou &

Ioannidis, 2012). Details of the analysis performed on the GO terms can be found in the

Supporting Information.

Genetic-Environment Association analysis

We performed redundancy analysis (RDA) and random forest (RF) regressions to identify

environmental variables significantly associated with spatial patterns of population divergence.

The environmental dataset used for these analyses consisted of sea surface temperature

(SST), sea bottom temperature (SBT), and sea surface salinity (SSS) for winter, spring, summer

and fall seasons, for a total of 12 oceanographic variables. These variables are relevant in

population structuring of numerous marine species in the NW Atlantic (Stanley et al., 2018).

To obtain environmental measures for each sampling location, we acquired monthly data

layers of SST, SBT, and SSS between 2008-2017 from NEMO 2.3 (Nucleus for European

Modelling of the Ocean), an oceanographic model developed by the Bedford Institute of

Oceanography, Canada. A detailed description of oceanic (Madec, Delecluse, Imbard, & Levy,

1998) and sea ice (Fichefet & Maqueda, 1997) model components can be found in Wang,

Brickman, Greenan, & Yashayaev (2016) and Brickman, Hebert, & Wang (2018). Data layers

were converted to an ASCII grid with a NAD83 projection (ellipse GRS80), they had a nominal

resolution of 1/12o (~5km^2), and a uniform land mask. Four seasonal bins, corresponding to

winter (January-February-March), spring (April-May-June), summer (July-August-September),

and fall (October-November-December), were averaged across 9 years in order to capture long-

term trends of oceanographic variation. Data extraction for the 14 geo-referenced locations was

conducted using custom R scripts (Stanley et al., 2018). Environmental data were standardized to

zero mean and unit variance in R for downstream analysis. Collinearity between environmental

variables was estimated with pairwise correlation coefficients computed with the function

pairs.panels of the R package *psych* (Revelle, 2018) (Fig. S11), and with variance inflation

factors (VIF) obtained from RDA models built with the R package *vegan* (Dixon, 2003). Prior to

RDA, the most collinear variables were removed based on biological/ecological criteria (Forester,

Lasky, Wagner, & Urban, 2018). Subsequently, remaining collinear variables were identified and

removed one by one in consecutive RDA runs based on their VIF. The variable with the highest

VIF was discarded in each run until all variables had a VIF < 5, following recommendations by

(Zuur, Ieno, & Elphick, 2010).

For RDA, we used the reduced environmental data as constraining variables for the

population allele frequencies of the top 500 outlier loci exhibiting the latitudinal pattern. RDA

runs were performed with the R package *vegan*, following Jeffery et al. (2018) and Lehnert et al.

(2018). Environmental variables that best explained genetic variance were identified using a bi-

directional stepwise permutational ordination method (1000 iterations) implemented in the R

function *ordistep*. Significance of the overall RDA model and of selected environmental variables

was assessed with analysis of variance (ANOVA) using 1000 permutations. In order to estimate

the proportion of the genetic variance independently explained by environment, geographic

distance, or both, we performed variance partitioning using partial redundancy analysis (pRDA),

either conditioned on geographic distance (Cartesian coordinates) or selected environmental

variables, respectively. Cartesian coordinates of each location, equivalent to the pairwise least-

cost geographic distance between locations accounting for land as barrier, were obtained with the

R function *CartDist* (Stanley & Jeffery, 2017). Concordance between Cartesian and geographic

coordinates was assessed with a linear regression (Fig. S3).

For RF regressions, we used the population allele frequencies of each outlier locus as

single response vectors and the 12 standardized environmental variables as predictors. A RF

regression was performed for each outlier locus with the R package *randomForest*, as described

in Lehnert et al. (2018) and Sylvester et al. (2018). Default parameters for regression were

applied to the RF runs (*mtry* = *p*/3, where *p* is the total number of predictors, or environmental

variables in this case), except that *ntree* was set to 10,000. The selected number of trees to grow

per run (*ntree*) assured Mean Decrease in Accuracy (MDA) convergence, as demonstrated in a

pilot test that compared MDA of predictors of 3 independent RF runs (correlation coefficient *r* =

0.9999, Fig. S4). Environmental variables were then ranked based on their relative importance to

explain genetic variance from the averaged MDA values across loci, and the mean residual

square error (MSE) of each location averaged across loci.

Isolation-by-distance pattern test

To evaluate whether global (all loci) and latitude-related population structure (subset of

loci) corresponded to an isolation-by-distance (IBD) pattern, we determined the significance of

the association between geographic and genetic distances for all possible pairs of sampled

spawning sites using Mantel tests (Mantel 1967) with 9999 permutations, implemented in the R

package *ade4* (Dray & Dufour, 2007). Genetic distances were linearized ($\hat{F}_{ST} = \frac{\hat{F}_{ST}}{1-\hat{F}_{ST}}$) (Rousset,

1997) with $\hat{F}_{ST}$ computed using all SNPs identified across the genome, in the first case, or solely

outlier SNPs strongly associated with latitudinal divergence, for the latter. Geographic distances

were estimated with the R package *CartDist* (Stanley & Jeffery, 2017) as the least-coast oceanic

distance in Km considering land as barrier.

Results:

Sampling distribution and pool-sequencing

A total of 697 adult herring from 14 spawning aggregations distributed in and around

Newfoundland and Labrador, the Gulf of St. Lawrence, Scotian Shelf, Bay of Fundy, and Gulf of

Maine in the NW Atlantic were included in this study (Fig. 1A, Table 1). We aimed to include in

the same pool only DNA from "ready-to-spawn" and "actively spawning" individuals collected

in the same area [gonadal maturity stage 5 and 6, respectively, (Bucholtz, Tomkiewicz, &

Dalskov, 2008)]. Yet, in some spawning aggregations (BDO-S, NDB-S, NDB-F, TRB-F, and

ME4-F, see pie charts in Fig. 1A) 25-50% of individuals were in "maturing" (stage 4) or

"resting" (stage 8) condition at the time of sampling. The designation of "S" or "F" in the

location name thus only reflects the season of collection and not necessarily the actual spawning

season of all fish included in the pool.

A total of ~800 GB of raw sequence data were obtained. After quality filtering and

adapter trimming, 6,119,940,640 reads of optimal quality (Phred score > 20) were available for

the genomic analysis. Read mapping statistics indicated that $> 98.8\%$ of read-pairs were correctly

aligned to the stitched version of the herring reference genome (mapping quality MQ > 48,

median insert size of 527 bp) (Table S1), confirming that misalignment errors, if present, were

negligible. Average read depth of coverage per pool ranged between 25x to 44x and varied

between sequencing batches [2015 batch mean 28.7 ± 4.0, 2016 batch mean 36.9 ± 2.6 (Table

S1). We monitored the potential effect of coverage variation in downstream analysis, in particular

for collections with lower coverage (TRB-S, NTS-S, and GEB-F). Variant calling resulted in

11,154,328 raw SNPs of which 2,189,380 passed quality filters and were retained for further

analysis.

Population structure

As observed in our previous study (Lamichhaney et al., 2017), spawning aggregations in

the NW Atlantic clustered according to reproductive season in a Neighbor-Joining tree, with

spring and fall spawning collections forming separate groups (Fig. 1B), although a few

exceptions were observed. BDO-S sample was in an intermediate position with respect to these

two main clusters, and a spring-collected sample in Newfoundland (NDB-S) clustered with the

fall group, suggesting it may be composed of a large proportion of fall spawners. A closer

examination of the fall group revealed clustering according to latitude. Southern collections in the

Scotian Shelf (MUS-F, GEB-F), Bay of Fundy (SCB-F), and Gulf of Maine (ME4-F) were

separated from northern collections in the Gulf of St. Lawrence (MIR-F, BLS-F), Newfoundland

(TRB-F, NDB-S, NDB-F) and Labrador (LAB-F). Such separation suggests genetic differences

may exist between herring inhabiting these two geographic regions.

The pairwise fixation index F_{ST} for pools ($\hat{F}_{ST}^{pool}$) ranged between 0.012 and 0.043,

indicating low levels of genetic structure among the 14 spawning aggregations studied (Fig. 1C,

pairwise F_{ST} values in Table S2). Nevertheless, three clear patterns of subtle genetic

differentiation were noticeable: i) between spring and fall spawners (SIL-S, SPH-S, NTS-S, vs.

others, $\hat{F}_{ST}^{pool}$ 0.022-0.043), ii) within spring spawners, the sample from the NW of the Gulf of St.

Lawrence (SIL-S) was the most genetically distinguishable ($\hat{F}_{ST}^{pool}$ ~0.030), and iii) within fall

spawners, the two southernmost collections (GEB-F and ME4-F) were the most divergent

($\hat{F}_{ST}^{pool}$ 0.020-0.031). In general, the largest genetic differentiation was observed between spring

spawners and the most southern collections ($\hat{F}_{ST}^{pool}$ ~0.040). Interestingly, the two spring-

collected samples BDO-S and NDB-S (two samples presumably containing both spring and fall

spawning individuals, see below) exhibited similar levels of differentiation ($\hat{F}_{ST}^{pool}$ 0.022-0.033)

with samples comprising solely spring spawners (SIL-S, SPH-S, NTS-S) as with samples

comprising solely fall spawners.

Outlier loci detection and genome-wide patterns of differentiation

A PCA-based whole-genome scan for the identification of SNPs putatively under

selection revealed two main axes of genomic differentiation in NW Atlantic herring: spawning

season, and geographic origin according to latitude. In a PCA plot based on 2,189,380 SNPs (Fig.

1D), spring and fall spawning herring were distinguishable along the first principal component

(PC1) (36% of variance explained). PC2 distinguished two collections, German Bank (GEB-F)

and Northumberland Strait (NTS-S) from the rest (Fig. S5). These two collections exhibited the

shallowest average sequencing coverage, suggesting this axis (PC2) is largely reflecting an

artefact of sequencing. PC2 was therefore ignored (Fig. S5). On PC3, the southernmost

collections, distributed on the Scotian Shelf, Bay of Fundy and Maine (MUS-F, SCB-F, GEB-F,

ME4-F), were differentiated from the aggregations in the Gulf of St. Lawrence, Newfoundland,

and Labrador (30% of the variance explained) that formed a tight cluster. The sample from Maine

(ME4-F) was the most differentiated of all, followed by German Banks (GEB-F), the

southernmost location sampled on the Scotian Shelf. Along PC1, BDO-S and SIL-S were

positioned in between the spring and fall spawners, BDO-S being closer to the fall samples and

SIL-S to the spring samples. NBD-S clustered tightly with the fall spawners. In general, with the

exception of the two southernmost samples (GEB-F and ME4-F), fall spawning aggregations

grouped more closely together than the spring spawning ones, suggesting that more genetic

differences may exist among the spring spawners than among fall spawners included in this

study.

440   In PC1, a total 14,724 outlier SNPs were detected (with Benjamini-Hochberg-adjusted P-

values and FDR $\leq$ 0.01). A Manhattan plot depicting significance values ($-\log_{10}P$-value) of

outlier loci for this PC disclosed numerous "peaks" or regions of divergence across the genome,

spanning about 18 scaffolds and numerous genes (Fig. 2A). The top SNPs of these scaffolds were

in the proximity of genes with known function in reproduction, such as *TSHR*, *ESRA*, *HERPUD2*,

CALM (Martinez Barrio *et al.* 2016). Moreover, a new set of candidate genes linked to seasonal

reproduction were *ISO3*, *SERTM1*, *SIPA1L1*, *CAMKK1, TMEM150C, CBLB, ENTPD5, KCNJ6,*

LPAR6 and *GPR119*, as they were near top outlier loci in the unique islands of differentiation

only observed in the NW Atlantic (Lamichhaney et al., 2017). A heatmap depicting standardized

population allele frequencies of the top 200 outlier loci from the scaffolds identified with RF

(ranked in descending order by -$\log_{10}P$-value) distinguished aggregations by spawning season

(Fig. 2B), with fall spawners fixed for one allele and almost all spring spawners fixed for the

alternative allele. The exceptions to this observation were three aggregations sampled in spring,

BDO-S, SIL-S, and NDB-S. The first two collections exhibited allele frequencies around 0.5,

while NDB-S showed population allele frequencies consistent with fall spawners. These results

indicate that BDO-S and SIL-S either correspond to a mixture of spring and fall spawning

individuals or to hybrids or both, and that NDB-S should be considered as a sample of fall

spawners, suggesting possible mislabeling.

In PC3, a total of 6,595 outlier loci were detected (with BH-adjusted *P*-values and FDR ≤

0.01). A Manhattan plot for this PC disclosed four main regions of divergence across the genome,

corresponding to scaffolds 44, 122, 869 and 958, and a small number of outlier loci from other

scaffolds (Fig. 2C). The top SNPs in the four main scaffolds were located within 5Kbp of the

genes *FAM129B, FNBP1, SH3GLB2*, and *GPR107*. A heatmap representing standardized

population allele frequencies of the top 200 outlier loci from the scaffolds identified with RF

(ranked in descending order by -$\log_{10}P$-value) revealed contrasting genetic patterns according to

latitude (Fig. 2D). In northern collections, including Labrador (LAB-F), Newfoundland (NDB-S,

NDB-F, TRB-F, SPH-S), Gulf of St. Lawrence (BLS-F, SIL-S, MIR-F, NTS-S), Bras D'Or lake

(BDO-S), and inner Bay of Fundy (SCB-F), one allele was close to fixation; in the southernmost

collection, in Maine (ME4-F), the alternative allele was in high frequency; and in intermediate

southern collections along the Scotian Shelf (MUS-F, GEB-F) allele frequencies were around

0.5. An extended examination of population allele frequencies of the 14,724 outlier SNPs

detected in PC1 (Fig. S8), revealed that additional SNPs from the four scaffolds showing the

latitudinal pattern were present in PC1 and showed the same pattern as the ones found in PC3

(3,378). Thus, these SNPs were removed from the PC1 set and added to the ones detected in PC3,

for a total of 11,346 SNPs associated with seasonal reproduction and 9,973 SNPs associated with

latitude.

A closer examination of the genomic distribution of outlier SNPs revealed that seasonal

reproduction-related outliers exhibited varying levels of significance (–$\log_{10}P$-value up to 30)

(Fig. 2A), were confined to a particular region within a scaffold (around 50-500 Kbp) and

spanned a given set of genes (Fig. S6). In contrast, latitude-related outliers showed similar

significance values ($-\log_{10}P$-value ~15) (Fig. 2C), were widely spread along scaffolds (covering

between 480 Kbp to 4.75 Mbp) and spanned numerous genes (Fig. S7).

Validation of a subset of outlier SNPs related to seasonal reproduction and to latitudinal

divergence

A total of 230 individuals (NDB-F: 30, NDB-S: 29, SIL-S: 27, NTS-S: 30, BDO-S: 28,

MUS-F: 29, GEB-F: 27, ME4-F: 30) and 52 and 74 SNPs related to seasonal reproduction and

latitudinal divergence, respectively, passed the missing rate and MAF quality filters. Heatmaps

depicting individual SNP genotypes for each of the two panels (Fig. S9) confirmed the overall

patterns of population allele frequencies of the two axes of divergence detected with Pool-seq

data (Fig. 2B,D), seasonal reproduction and latitude.

The SNP panel discriminating spawning season revealed that the spring-collected samples

SIL-S and BDO-S corresponded to a mixture of spring and fall spawners and putative hybrids,

the latter defined as heterozygous individuals at many of the loci showing a high degree of

fixation between groups. SIL-S comprised an even proportion of pure fall spawners and putative

hybrids with a few pure spring spawners, whereas BDO-S comprised mostly pure fall spawners

and a few hybrids and spring spawners. The other spring-collected samples, NTS-S, consisted of

mostly pure spring spawners and a few putative hybrids, while NDB-S corresponded to pure fall

spawners. In contrast, all the fall-collected samples genotyped (NDB-F. MUS-F, GEB-F and

ME4-F) corresponded to pure fall spawners, with a few heterozygous loci.

The SNP panel discriminating by latitude confirmed northern samples were characterized

by high frequency of one allele, while the alternative allele had greater frequency in the

southernmost sample (in Maine), although putative hybrids were present in both cases in varying

proportions. Intermediate locations (BDO-S, MUS-F, GEB-F) exhibited a genotypic cline of

increasing proportion of putatively hybrids towards the south.

Functional annotation of outlier loci

A total of 2,977 and 1,257 outlier SNPs associated with seasonal reproduction and

latitudinal divergence, respectively, were annotated with respect to a neighboring gene (within

5Kbp). For both cases, the majority of outlier SNPs were located within introns and intergenic

regions, or 5Kbp upstream or downstream of genes (Fig. 3A). A small number of outlier SNPs

were predicted as synonymous (~2%) or missense variants (1,6% and 0.9%, for spawning- and

latitude-related outliers, respectively).

Excluding intergenic variants and genes that did not correspond to an orthologous gene in

zebrafish, a list of 298 and 182 genes associated with seasonal reproduction and latitudinal

divergence in herring, respectively, resulted from the annotated outlier loci. For seasonal

reproduction-related genes, 126 had a GO term in the biological process category, 109 in the

cellular component category, and 120 in the molecular function category (Fig. S10A). For

latitude-related genes, 90 had a GO term in the biological process category, 72 in the cellular

component category, and 80 in the molecular function category; considered together, close to half

of the genes lacked GO classification. A comprehensive description of particular functions within

the three GO categories and the number of genes in each of them is presented in Fig. S10B).

The overrepresentation enrichment analysis (ORA) of both sets of candidate genes did not

reach statistical significance (FDR of 5%) (Table S3 and S4), likely due to the large number of

genes lacking GO annotation (Fig. S10). However, a closer examination of the top GO terms with

P-value < 0.05 (ranked in ascending P-values from ORA, Table S3 and S4, GO terms indicated

with an asterisk), suggested that seasonal reproduction-related candidate genes may participate in

biological processes such as metabolism of lipids, cell adhesion, biosynthesis of cellular

products, peptidyl-aminoacid modification, protein complex biogenesis, inositol lipid-mediated

signaling, developmental maturation, regulation of developmental process, and cellular

component organization (Fig. 3B-top, Table S3). These genes might primarily act in cellular

components such the endoplasmic reticulum and the whole membrane (Fig. 3B-middle) and play

a molecular function related to cell adhesion molecule and protein binding and lipid transporter

and transferase activities (Fig. 3B-bottom). The top GO terms of candidate genes associated with

latitudinal divergence were all involved in embryological and organ development processes (Fig.

3C-top, Table S4). These genes might act in cellular components such phosphatase complex,

collagen trimer, and in the extracellular region (Fig. 3C-middle), and participate in sulfur

compound binding and hydrolase and isomerase activities (Fig. 3C-bottom).

Genome-Environment Association analysis

Collinearity among several of the environmental variables examined and redundancy

analyses (See Supporting Information) allowed us to reduce the environmental data set to just

three variables: summer SBT, winter SST, and spring SSS. RDA indicated that winter SST

(Win_SST) (Fig. 4A) was the environmental variable that best explained the genetic variance of

outlier loci exhibiting the latitudinal cline ($F = 16.7$, $p = 0.001$, from *ordistep* function) (Fig. 4B).

No other temperature or salinity variable in the reduced environmental dataset was significant

(from ANOVA with 1000 permutations, significance value = 0.05). Spawning aggregations were

separated according to Win_SST on RDA axis 1, which explained 58.1% of the total genetic

variance ($R^2 = 0.58$, adjusted $R^2 = 0.55$). pRDA however, showed that the Win_SST-based RDA

model was no longer significant when the effect of geographic distance between sites was

removed from the model. A variance partitioning analysis revealed that the interaction between

environment and geographic distance explained the greatest proportion of clinal genetic variation

(44.9%).

In agreement with RDA results, RF regressions also indicated that Win_SST was the most

important environmental variable (MDA = 23.5), followed by Fall_SST (MDA = 21.8) (Fig. 4C).

The other temperature variables had lower importance (MDA < 10), and salinity measures were

the least important of all (MDA < 5). ME4-F, the southernmost spawning aggregation sampled,

exhibited the highest mean square error (MSE = 0.21), followed by SCB-F and MUS-F (MSE ~

0.05), whereas the other 10 collections had lower MSE, below 0.03 (Fig. 4D).

A closer examination of the map of the NW Atlantic depicting average Win_SST over the

last 9 years and the predominant population allele frequency of the 14 sites studied (Fig. 4A),

revealed that herring in "northern" collections in the Bay of Fundy, the Gulf of St. Lawrence, and

Newfoundland and Labrador were characterized by being exposed to temperatures below zero (-2

℃), whereas in "southern" collections they were mainly exposed to temperatures above zero (>2

℃).

Isolation-by-distance test

The Mantel test showed there is not a significant linear relationship between geographic

and genetic distances for all loci across the genome ($R^2 = 0.04$), whereas there is a significant

linear relationship ($R^2 = 0.30$) between geographic distance and genetic differentiation when only

looking at outlier SNPs exhibiting the latitudinal break in population allele frequencies between

northern and southern collections (Fig. 5).

Discussion:

Here we described patterns of genetic variation at the whole-genome level in Atlantic

herring populations distributed across the reproductive range of the species in North America.

This study represents the most comprehensive assessment of this kind in the region to date. We

uncovered fine-scale population structure at outlier loci putatively under selection, despite low

differentiation at selectively neutral loci. This observation is consistent with previous genetic

work on herring in both, the NE (Guo et al., 2016; Lamichhaney et al., 2012; Limborg et al.,

2012; Martinez Barrio et al., 2016; Teacher, André, Jonsson, & Merilä, 2013) and the NW

Atlantic (Lamichhaney et al., 2017; McPherson et al., 2004; McPherson, Stephenson, O'Reilly,

Jones, & Taggart, 2001). The large population sizes, high potential for gene flow, and minute

effect of genetic drift explain the low genetic differentiation observed at neutral loci (Palumbi,

1994). These conditions also favor the more efficient action of natural selection, which seems to

be behind the genetic differences observed at outlier loci.

While prior genomic studies disclosed genetic structure with seasonal reproduction and

salinity (Lamichhaney et al., 2012; Martinez Barrio et al., 2016), and others suggested structuring

along the salinity/temperature gradient in the Baltic Sea from dozens of markers While prior

genomic studies disclosed genetic structure with seasonal reproduction and salinity

(Lamichhaney et al., 2012; Martinez Barrio et al., 2016), and others suggested structuring along

the salinity/temperature gradient in the Baltic Sea from a dozens of markers (Gaggiotti et al.,

2009; Guo et al., 2016; Limborg et al., 2012), here we successfully disentangled two main

overlapping axes of divergence supported by thousands of outlier SNPs: seasonal reproduction

and a latitudinal cline defining a north-south genetic break. Our genetic-environment association

analyses indicated that winter sea-surface temperature is the best predictor of the spatial structure

observed. These results: demonstrate for the first time that herring from the north (Labrador,

Newfoundland, Gulf of St. Lawrence and Bay of Fundy) are genetically distinguishable from the

ones in the south (Scotian Shelf and Maine) regardless of their spawning season; indicating that

thermal-minima related factors are likely driving latitudinal genetic differentiation; and provide

additional evidence supporting the recently described multispecies biogeographic break in eastern

Nova Scotia (Stanley et al. 2018).

Outlier SNPs exhibited remarkable clustering, forming so-called "genomic regions of

divergence" (Nosil, Funk, & Ortiz-Barrientos, 2009; T. L. Turner, Hahn, & Nuzhdin, 2005), and

extreme allele frequency differences (i.e. alternative alleles were close to fixation in either spring-

or fall-spawning, or in northern- or the southernmost populations). Theory predicts that formation

of genomic regions of divergence (Schluter, 2009; Wu, 2001) and fixation of different alleles

conducive to opposing phenotypes often result from natural selection acting in contrasting

directions between environments (Vitti, Grossman, & Sabeti, 2013). Considering the

heterogeneous environmental properties of the Northwest Atlantic (Melvin et al., 2009;

Townsend et al., 2004) and having discarded an effect of genetic drift and an isolation-by-

distance pattern, we conclude that disruptive selection may be the main evolutionary force

involved in population structuring in the region.

A few exceptions to the allele fixation pattern were observed in both axes of divergence.

In seasonal reproduction outliers, two aggregations sampled in spring, BDO-S and SIL-S,

exhibited allele frequencies around 0.5 at SNPs being closed to fixation for opposite alleles in

other populations of spring- and fall-spawning herring. This observation suggests these

collections either correspond to a mixture of spring- and fall-spawners, or to a unique population

where allele diversity is favored. Individual genotypes of a subset of diagnostic SNPs of

spawning time confirmed BDO-S and SIL-S comprised a mixture of spring and fall spawners and

putative hybrids (i.e. heterozygous individuals at many of the loci showing a high degree of

fixation between groups). In latitude-related outliers, intermediate allele frequencies were

observed in MUS-F and GEB-F, two locations in southwestern Nova Scotia, mid-range in the

latitudinal cline. Interestingly, these locations are few kilometers south of the biographic barrier

described in the NW Atlantic (Stanley et al., 2018). Environmental conditions in the NW Atlantic

vary between years in relation to oceanographic global trends (Townsend et al., 2004). It is

possible then that populations in southwestern Nova Scotia experience significant inter-annual

environmental fluctuations during winter months, depending on the strengthening either of the

warm Gulf Stream flowing north or of the cold Labrador Current flowing south. Under these

dynamic circumstances, it is possible that balancing selection may be maintaining polymorphism

at these loci. Additional studies including an extended sampling in the southern region could be

used to test this hypothesis.

A closer examination of the genomic regions of divergence revealed they vary in size and

genomic location between the two axes of divergence. Seasonal reproduction-related outliers

were distributed across 18 scaffolds in which they spanned about 50-500 Kbp and a given set of

genes. In contrast, latitude-related outliers were mostly spread in four scaffolds, covering a larger

extension, from 480 Kbp to 4.75 Mbp, and larger number of genes. The observation that latitude-

related outliers were widely distributed and consistently divergent across four large scaffolds

suggests that they could be located within a chromosomal rearrangement. If this were the case,

the expectation would be that populations from the north were homozygous for one state of the

variant, the ones in southwest Nova Scotia were polymorphic, and in the Gulf of Maine were

homozygous for the alternative state of the variant. Further research supported by a linkage map,

not described yet for herring, is required for the evaluation of this hypothesis.

A bioinformatic evaluation of the functional effect of outlier SNPs disclosed that, for both

axes of divergence, the majority of SNPs were located within introns, intergenic regions, and

5Kbp upstream or downstream of genes, and a smaller proportion corresponded to missense

mutations (1,6% and 0.9%, for spawning- and latitude-related outliers, respectively). Mutations

in introns can modify regulatory domains, intron-exon boundaries and RNA splicing (Pagani &

Baralle, 2004); missense mutations result in a different amino acid; and mutations in regulatory

elements can modify gene expression (Epstein, 2009; Metzger et al., 2016; M. Nei, 2007). While

at this point is not possible to trace a direct link between single SNPs and gene function or

identify causal mutations, our observations suggest that single base changes in introns, protein-

coding, and regulatory regions may be involved in adaptive divergence in NW Atlantic herring,

in agreement with previous observations in the NE Atlantic (Martinez Barrio et al., 2016).

Gene annotation of top outlier SNPs confirmed that *TSHR*, *HERPUD2*, *SOX1*, *SOX11A*,

SYNE1, *SYNE2*, and *ESR2A* are candidate genes related to seasonal reproduction. These genes

have a known function in reproduction and were previously linked to spawning time in NE

Atlantic herring (Lamichhaney et al., 2017; Martinez Barrio et al., 2016). We discovered an

additional set of candidate genes, *ISO3*, *SERTM1*, *SIPA1L1*, *CAMKK1, TMEM150C, CBLB,*

ENTPD5, KCNJ6, LPAR6 and *GPR119*, corresponding to the genomic regions of differentiation

uniquely observed in the NW Atlantic (Lamichhaney et al., 2017), hence, they can potentially be

involved in local adaptation. Candidate genes related with the latitudinal cline are *FAM129B,*

FNBP1, SH3GLB2, and *GPR107*.

A qualitative examination of the top ranked GO terms indicated that candidate genes

related to seasonal reproduction may be involved in biological processes such as metabolism of

lipids, biosynthesis of cellular products, developmental maturation, regulation of developmental

process, and cellular component organization. Similarly, latitude-related candidate genes may

participate in embryological and organ development processes. These observations suggest that

outlier SNPs underlying the two axes of divergence may be involved in different physiological

pathways, and that natural selection along the latitudinal cline likely acts on early life stages, in

agreement with the proposed hypothesis for the multispecies climatic cline (Stanley et al., 2018).

It is likely that early life stages experience selection along the latitudinal cline given that larval

retention areas are in the proximity of spawning grounds (Stephenson et al., 2009). If selection

would act on juveniles or adults, which are highly migratory, then the pattern should not coincide

with spawning locations.

683   We provide genetic evidence that suggests timing of reproduction and latitudinal

684 spawning location are features under disruptive selection leading to local adaptation. Several

685 characteristics of herring biology and ecology seem to support this. For instance, (*i*) spawning

686 occurs at predictable times and locations, the timing differs among geographic regions

687 (Stephenson et al., 2009), and there is no evidence indicating that individual fish can switch

688 spawning season (Melvin et al., 2009); (*ii*) herring spawns once a year and exhibits spawning site

689 fidelity (Wheeler & Winters, 1984); (*iii*) spring- and fall-spawners differ in morphometric

690 characters, in life-history traits (fecundity, egg size and growth), and in phenotypic traits (number

691 of vertebrae and otolith shape) (Baxter, 1959; Cushing, 1967; Messieh, Anthony, & Sinclair,

692 1985); growth rate, otolith shape, and vertebral counts seem to be largely influenced by genetic

693 factors (Berg et al., 2018); (*iv*) early life stages spawned in different seasons and locations

694 experience contrasting environmental conditions (e.g. in the Gulf of St. Lawrence, eggs released

695 by spring spawners hatch after 30 days at 5°C, while eggs of fall spawners hatch after 10 days at

696 15°C; in Nova Scotia, eggs of fall spawners hatch in 11 days at 10°C) (Scott & Scott, 1988); (*v*)

697 larval retention areas occur near spawning grounds and are stable over time, in predictable

698 patterns related to oceanographic conditions (Stephenson et al., 2009); and (*vi*) genetic

699 differences between spring- and fall-spawners are temporally stable (Kerr et al., 2018). From this,

700 we then infer that timing of reproduction and latitudinal spawning location can be adaptive

701 strategies to increase offspring survival, particularly at vulnerable early life stages, in

702 environments that vary seasonally and geographically. When timing of reproduction is largely

703 heritable, the resulting temporal assortative mating may reduce gene flow between individuals

704 breeding at different times (Hendry & Day, 2005). In herring, gene flow may be limited between

705 early spring-spawners and late fall-spawners even if they are in sympatry (as their gonads are not

ripe at the same time, as we observed in our samples). How do hybrids occur? We hypothesize

hybridization could happen between late spring-spawners and early fall-spawners at geographic

areas where both reproductive strategies coexist (e.g. in the Gulf of St. Lawrence), and when the

onset of gonadal maturation coincides (likely temperature driven). Hybrids would survive then, if

they can cope with the local environmental conditions.

Although disruptive selection is a strong candidate for explaining latitudinal divergence in

herring, other mechanisms are possible. For example, additional biotic or abiotic factors that

covariate with temperature may be the actual drivers of adaptation. Pre- or post-zygotic

reproductive incompatibilities that coincide with latitude (but are not dependent on) can result in

the observed spatial genetic discontinuity (Bierne, Welch, Loire, Bonhomme, & David, 2011).

The current latitudinal break may actually reflect historical vicariance (Bradbury et al., 2010), not

contemporary population dynamics (Palumbi, 1994). Further studies are required to evaluate

these alternative hypotheses.

Even though valuable information was obtained through this study, there were some

limitations. In Pool-seq individual information is missed, thus it is not possible to correct

accidental mixing of individuals with different origin/spawning season. To avoid this, we

selected maturing and ripe fish collected in known spawning grounds during the local peak of

reproduction. Despite these precautions, we found evidence of some mixed aggregations (SIL-S

and BDO-S). Moreover, in the over-representation enrichment analysis statistical significance

was not reached. This outcome may have been influenced by the restriction that only herring

candidate genes with an zebrafish ortholog could be included, and that half of the total genes

mapped to zebrafish lacked a GO term. We expect with a more complete reference genome and

annotations, along with functional experiments, a better functional characterization of outlier loci

will be achieved.

Our findings have several implications and potential applications in fisheries. Firstly, our

results support the maintenance of separate management of spring- and fall-spawning

components currently in place across most of the region. Secondly, management units should be

revised in order to protect the functional intraspecific biodiversity revealed in this study,

specifically considering a climate change scenario as spring-spawners seem to be less resilient to

a warming ocean (Melvin et al., 2009). Thirdly, as we now have the molecular tools to

distinguish herring spawning in spring or autumn and in northern and southern regions, a subset

of outlier SNPs reported here can be used for genetic monitoring of stock composition already at

the larval stage and out of breeding seasons to minimize the risk of overexploitation of vulnerable

components within mixed stocks. And lastly, the current herring population models could be

revised as none of them are in complete agreement with our genetic data, as similarly noted by

McPherson et al. (2004). For instance, the discrete population concept proposes that gene flow is

limited, hybrids have reduced fitness, and local populations are reproductively isolated by fixed

spawning time, natal homing, spawning site fidelity, and larval retention areas with particular

hydrographic features (Sinclair, 1988; Sinclair & Iles, 1989). While our data agrees with most of

this, the presence of numerous putative hybrids suggests that gene flow may be more extensive

than expected under this model and they are viable. In the dynamic balance population concept

there is significant gene flow, no stable population structure, no fixed spawning time, no

philopatry, no larval retention areas, as populations respond to changing environmental

conditions (Smith & Jamieson, 1986). The temporal and spatial structuring we observed is

opposite to this model. And in the metapopulation concept (adopted migrant) there is repeated

homing to traditional spawning grounds defined by hydrographic features, migration and homing

patterns are socially transmitted, and significant gene flow can occur as vagrants are adopted by

non-natal local populations (McQuinn, 1997). This model implies an isolation-by-distance

pattern and that spawning time is not genetically determined (it is learned), contrary to our

observations.

In summary, our results confirm that Atlantic herring is a model system for the study of

ecological adaptation with gene flow in the wild (Lamichhaney et al., 2017; Martinez Barrio et

al., 2016), and provide insight into patterns and mechanisms of genomic divergence and local

adaptation despite gene flow in an abundant and highly dispersive marine fish.

Acknowledgements: We thank staff at Fisheries and Oceans Canada, the Maine Department of

Resources, Comeau's Seafoods Ltd, Cape Breeze Seafoods Ltd, and fishers Crystal and Donald

Kent, Gordon McKay, and many others for their valuable contribution in sample collection.

Thanks to Gregory McCracken for assistance in sampling and DNA extraction, to Gavin

Douglas, Emma Sylvester, Sarah Lehnert, Ryan Stanley, Simone Fior, Alan Bergland, Miguel

Carneiro, Michael Blum, and Mathieu Gautier for data analysis assistance. Many thanks to Ryan

Stanley for obtaining the oceanographic data layers used in this study. Library preparation and

shotgun sequencing was performed in The Centre for Applied Genomics of the Hospital for Sick

Children, Canada. Computations were conducted on the supercomputer Mp2 of the University of

Sherbrooke, managed by Calcul Québec and Compute Canada and funded by the Canada

Foundation for Innovation (CFI), the ministère de l'Économie, de la science et de l'innovation du

Québec (MESI) and the Fonds de recherche du Québec - Nature et technologies (FRQ-NT).

A.P.F.P. and D.E.R. thank the Killam Trust. A.P.F.P. thanks the Vanier Canada Graduate

Scholarship, the President's Award of Dalhousie University, the Nova Scotia Graduate

Scholarship, the Lett Fund and a Strategic grant to D.E.R for graduate studies funding. This study

was funded by NSERC Discovery and Strategic grants to D.E.R.

Data accessibility: Oceanographic data, and population-allele frequencies and individual SNP

genotype data for this study will be available in Dryad upon publication acceptance. Bash, python

and R scripts are available upon request to the authors.

Author contributions: D.E.R. and A.P.F.P. designed and conceived the study; C.B., R.S., K.E.,

L.P., and J.L.M provided herring samples; A.P.F.P contributed to tissue collection and

processing, and performed lab work and bioinformatics data analysis; D.E.R and L.A.

contributed to the interpretation of results; A.P.F.P. wrote the manuscript with input from D.E.R.

and L.A. All authors vetted and approved the manuscript before submission.

Competing interests: The authors declare no competing financial interests.

References:

Allendorf, F. W., Hohenlohe, P. A., & Luikart, G. (2010). Genomics and the future of conservation genetics. *Nature Reviews Genetics*, *11*(10), 697–709. https://doi.org/10.1038/nrg2844

Andersson, L., Ryman, N., Rosenberg, R., & Ståhl, G. (1981). Genetic variability in Atlantic herring (Clupea harengus harengus): description of protein loci and population data. *Hereditas*, *95*(1), 69–78. https://doi.org/10.1111/j.1601-5223.1981.tb01330.x

André, C., Larsson, L. C., Laikre, L., Bekkevold, D., Brigham, J., Carvalho, G. R., … Ryman, N. (2011). Detecting population structure in a high gene-flow species, Atlantic herring (Clupea harengus): direct, simultaneous evaluation of neutral vs putatively selected loci. *Heredity*, *106*(2), 270–280. https://doi.org/10.1038/hdy.2010.71

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. Retrieved May 1, 2018, from http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Barrett, R. D. H., & Hoekstra, H. E. (2011). Molecular spandrels: tests of adaptation at the genetic level. *Nature Review Genetics*, *12*(11), 767–780. https://doi.org/10.1038/nrg3015

Baum, B. R. (1989). PHYLIP: Phylogeny Inference Package. Version 3.2 . Joel Felsenstein. *The Quarterly Review of Biology*, *64*(4), 539–541. https://doi.org/10.1086/416571

Baxter, I. G. (1959). Fecundities of winte-spring and summer-autumn herring spawners. *J. Cons. Int. Explor. Mer.*, *25*, 73–80.

Benestan, L., Gosselin, T., Perrier, C., Sainte-Marie, B., Rochette, R., & Bernatchez, L. (2015). RAD genotyping reveals fine-scale genetic structuring and provides powerful population assignment in a widely distributed marine species, the American lobster (Homarus americanus). *Molecular Ecology*, *24*(13), 3299–3315. https://doi.org/10.1111/mec.13245

Benoît, H., Swain, D., Hutchings, J., Knox, D., Doniol-Valcroze, T., & Bourne, C. (2018). Evidence for reproductive senescence in a broadly distributed harvested marine fish. *Marine Ecology Progress Series*, *592*, 207–224. https://doi.org/10.3354/meps12532

Berg, F., Almeland, O. W., Skadal, J., Slotte, A., Andersson, L., & Folkvord, A. (2018). Genetic factors have a major effect on growth, number of vertebrae and otolith shape in Atlantic herring (Clupea harengus). *PLOS ONE*, *13*(1), e0190995. https://doi.org/10.1371/journal.pone.0190995

Bierne, N., Welch, J., Loire, E., Bonhomme, F., & David, P. (2011). The coupling hypothesis: Why genome scans may fail to map local adaptation genes. *Molecular Ecology*, *20*(10), 2044–2072. https://doi.org/10.1111/j.1365-294X.2011.05080.x

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bourne, C., Mowbray, F., Squires, B., & Koen-Alonso, M. (2018). 2017 Assessment of Newfoundland east and south coast Atlantic herring (Clupea harengus) stock complexes. *DFO Can. Sci. Advis. Sec. Res. Doc. 2018/026*, v + 45 p.

Bradbury, I. R., Hubert, S., Higgins, B., Borza, T., Bowman, S., Paterson, I. G., … Bentzen, P. (2010). Parallel adaptive evolution of Atlantic cod on both sides of the Atlantic Ocean in response to temperature. *Proceedings of the Royal Society of London B: Biological Sciences*, *277*(1701), 3725–3734. https://doi.org/10.1098/rspb.2010.0985

Bradbury, I. R., Hubert, S., Higgins, B., Bowman, S., Borza, T., Paterson, I. G., … Bentzen, P. (2013). Genomic islands of divergence and their consequences for the resolution of spatial structure in an exploited marine fish. *Evolutionary Applications*, *6*(3), 450–461. https://doi.org/10.1111/eva.12026

Brickman, D., Hebert, D., & Wang, Z. (2018). Mechanism for the recent ocean warming events on the Scotian Shelf of eastern Canada. *Continental Shelf Research*, *156*, 11–22. https://doi.org/10.1016/j.csr.2018.01.001

Britten, G. L., Dowd, M., & Worm, B. (2016). Changing recruitment capacity in global fish stocks. *Proceedings of the National Academy of Sciences*, *113*(1), 134–139.

843   https://doi.org/10.1073/pnas.1504709112

844 Bucholtz, R. H., Tomkiewicz, J., & Dalskov, J. (2008). *Manual to determine gonadal maturity of herring*
845   *(Clupea harengus L.)*. *DTU Aqua-report*. Retrieved from
846   http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Manual+to+determine+gonadal+
847   maturity+of+herring+(Clupea+harengus+L.)#0

848 Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., … Ruden, D. M. (2012). A
849   program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly*,
850   *6*(2), 80–92. https://doi.org/10.4161/fly.19695

851 Cushing, D. H. (1967). The grouping of herring populations. *J. Mar. Biol. Ass., U.K*, *47*, 193–208.

852 DFO. (1997). *State of the Ocean: Northwest Atlantic*. *DFO Science Stock Status Report G0-01*. Retrieved
853   from
854   https://login.proxy.lib.duke.edu/login?url=https://search.proquest.com/docview/1668269133?account
855   id=10598%0Ahttp://pm6mt7vg3j.search.serialssolutions.com?ctx_ver=Z39.88-
856   2004&ctx_enc=info:ofi/enc:UTF-8&rfr_id=info:sid/ProQ%3Aasfabiological&rft_val_fmt=info

857 Dixon, P. (2003). VEGAN, a package of R functions for community ecology. *Journal of Vegetation*
858   *Science*, *14*(6), 927–930. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

859 Dohm, J. C., Lottaz, C., Borodina, T., & Himmelbauer, H. (2008). Substantial biases in ultra-short read
860   data sets from high-throughput DNA sequencing. *Nucleic Acids Research*, *36*(16), e105–e105.
861   https://doi.org/10.1093/nar/gkn425

862 Dray, S., & Dufour, A.-B. (2007). The ade4 Package: Implementing the Duality Diagram for Ecologists.
863   *Journal of Statistical Software*, *22*(4). https://doi.org/10.18637/jss.v022.i04

864 Engelhard, G. H., & Heino, M. (2004). Maturity changes in Norwegian spring-spawning herring before,
865   during, and after a major population collapse. *Fisheries Research*, *66*(2–3), 299–310.
866   https://doi.org/10.1016/S0165-7836(03)00195-4

867 Epstein, D. J. (2009). Cis-regulatory mutations in human disease. *Briefings in Functional Genomics and*
868   *Proteomics*, *8*(4), 310–316. https://doi.org/10.1093/bfgp/elp021

869 Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: summarize analysis results for
870   multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048.
871   https://doi.org/10.1093/bioinformatics/btw354

872 Fadista, J., Manning, A. K., Florez, J. C., & Groop, L. (2016). The (in)famous GWAS P-value threshold
873   revisited and updated for low-frequency variants. *European Journal of Human Genetics*, *24*(8),
874   1202–1205. https://doi.org/10.1038/ejhg.2015.269

875 FAO. (2019). Species Fact Sheets: Clupea harengus (Linnaeus, 1758). Retrieved from
876   http://www.fao.org/fishery/species/2886/en

877 Feder, A. F., Petrov, D. A., & Bergland, A. O. (2012). LDx: Estimation of Linkage Disequilibrium from
878   High-Throughput Pooled Resequencing Data. *PLoS ONE*, *7*(11), e48588.
879   https://doi.org/10.1371/journal.pone.0048588

880 Fichefet, T., & Maqueda, M. A. M. (1997). Sensitivity of a global sea ice model to the treatment of ice
881   thermodynamics and dynamics. *Journal of Geophysical Research: Oceans*, *102*(C6), 12609–12646.
882   https://doi.org/10.1029/97JC00480

883 Forester, B. R., Lasky, J. R., Wagner, H. H., & Urban, D. L. (2018). Comparing methods for detecting
884   multilocus adaptation with multivariate genotype–environment associations. *Molecular Ecology*,
885   *27*(9), 2215–2233. https://doi.org/10.1111/mec.14584

886 Gaggiotti, O. E., Bekkevold, D., Jørgensen, H. B. H., Foll, M., Carvalho, G. R., Andre, C., & Ruzzante, D.
887   E. (2009). Disentangling the effects of evolutionary, demographic, and environmental factors
888   influencing genetic structure of natural populations: Atlantic herring as a case study. *Evolution*,
889   *63*(11), 2939–2951. https://doi.org/10.1111/j.1558-5646.2009.00779.x

890 Goldstein, B. A., Hubbard, A. E., Cutler, A., & Barcellos, L. F. (2010). An application of Random Forests
891   to a genome-wide association dataset: Methodological considerations & new findings. *BMC*

Genetics, *11*(1), 49. https://doi.org/10.1186/1471-2156-11-49

Gu, Z., Eils, R., & Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics*, *32*(18), 2847–2849.

https://doi.org/10.1093/bioinformatics/btw313

Guo, B., Li, Z., & Merilä, J. (2016). Population genomic evidence for adaptive differentiation in the Baltic

Sea herring. *Molecular Ecology*, *25*(12), 2833–2852. https://doi.org/10.1111/mec.13657

Hauser, L., & Carvalho, G. R. (2008). Paradigm shifts in marine fisheries genetics: ugly hypotheses slain

by beautiful facts. *Fish and Fisheries*, *9*(4), 333–362. https://doi.org/10.1111/j.1467-

2979.2008.00299.x

Hendry, A. P., & Day, T. (2005). Population structure attributable to reproductive time: isolation by time

and adaptation by time. *Molecular Ecology*, *14*(4), 901–916. https://doi.org/10.1111/j.1365-

294X.2005.02480.x

Hivert, V. (2018). Measuring genetic differentiation from Pool-seq data.

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., … Whitlock, M. C.

(2016). Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future

Directions. *The American Naturalist*, *188*(4), 379–397. https://doi.org/10.1086/688018

Iles, T. D., & Sinclair, M. (1982). Atlantic Herring: Stock Discreteness and Abundance. *Science*,

215(4533), 627–633. https://doi.org/10.1126/science.215.4533.627

Jeffery, N. W., Bradbury, I. R., Stanley, R. R. E., Wringe, B. F., Van Wyngaarden, M., Lowen, J. Ben, …

DiBacco, C. (2018). Genomewide evidence of environmentally mediated secondary contact of

European green crab (Carcinus maenas) lineages in eastern North America. *Evolutionary*

Applications, *11*(6), 869–882. https://doi.org/10.1111/eva.12601

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., … Kingsley, D. M.

(2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, *484*(7392), 55–

61. https://doi.org/10.1038/nature10944

Jorgensen, H. B. H., Hansen, M. M., Bekkevold, D., Ruzzante, D. E., & Loeschcke, V. (2005). Marine

landscapes and population genetic structure of herring (Clupea harengus L.) in the Baltic Sea.

Molecular Ecology, *14*(10), 3219–3234. https://doi.org/10.1111/j.1365-294X.2005.02658.x

Kerr, Q., Fuentes⬜Pardo, A. P., Kho, J., McDermid, J. L., & Ruzzante, D. E. (2018). Temporal stability

and assignment power of adaptively divergent genomic regions between herring (Clupea harengus)

seasonal spawning aggregations. *Ecology and Evolution*, (October), ece3.4768.

https://doi.org/10.1002/ece3.4768

Kolaczkowski, B., Kern, A. D., Holloway, A. K., & Begun, D. J. (2011). Genomic Differentiation

Between Temperate and Tropical Australian Populations of Drosophila melanogaster. *Genetics*,

187(1), 245–260. https://doi.org/10.1534/genetics.110.123059

Lamichhaney, S., Barrio, A. M., Rafati, N., Sundstrom, G., Rubin, C.-J., Gilbert, E. R., … Andersson, L.

(2012). Population-scale sequencing reveals genetic differentiation due to local adaptation in

Atlantic herring. *Proceedings of the National Academy of Sciences*, *109*(47), 19345–19350.

https://doi.org/10.1073/pnas.1216128109

Lamichhaney, S., Fuentes-Pardo, A. P., Rafati, N., Ryman, N., McCracken, G. R., Bourne, C., …

Andersson, L. (2017). Parallel adaptive evolution of geographically distant herring populations on

both sides of the North Atlantic Ocean. *Proceedings of the National Academy of Sciences*, *114*(17),

E3452–E3461. https://doi.org/10.1073/pnas.1617728114

Lehnert, S. J., DiBacco, C., Van Wyngaarden, M., Jeffery, N. W., Ben Lowen, J., Sylvester, E. V. A., …

Bradbury, I. R. (2018). Fine-scale temperature-associated genetic structure between inshore and

offshore populations of sea scallop (Placopecten magellanicus). *Heredity*, 1–12.

https://doi.org/10.1038/s41437-018-0087-9

Lewontin, R. C. (2002). Directions in Evolutionary Biology. *Annual Review of Genetics*, *36*(1), 1–18.

https://doi.org/10.1146/annurev.genet.36.052902.102704

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *ArXiv*
Preprint ArXiv, *00*(00), 3. https://doi.org/arXiv:1303.3997 [q-bio.GN]
Liaw, A., & Wiener, M. (2002). Classification and Regression by randomForest. *R News*, *2*(3), 18–22.
Limborg, M. T., Helyar, S., De Bruyn, M., Taylor, M. I., Nielsen, E. E., Ogden, R., … Bekkevold, D.
(2012). Environmental selection on transcriptome-derived SNPs in a high gene flow marine fish, the
Atlantic herring (Clupea harengus). *Molecular Ecology*, *21*(15), 3686–3703.
https://doi.org/10.1111/j.1365-294X.2012.05639.x
Lischer, H. E. L., & Excoffier, L. (2012). PGDSpider: an automated data conversion tool for connecting
population genetics and genomics programs. *Bioinformatics*, *28*(2), 298–299.
https://doi.org/10.1093/bioinformatics/btr642
Luu, K., Bazin, E., & Blum, M. G. B. (2017). pcadapt: an R package to perform genome scans for
selection based on principal component analysis. *Molecular Ecology Resources*, *17*(1), 67–77.
https://doi.org/10.1111/1755-0998.12592
Madec, G., Delecluse, P., Imbard, M., & Levy, C. (1998). *OPA8.1 Ocean general Circulation Model*
reference manual. France.
Martinez Barrio, A., Lamichhaney, S., Fan, G., Rafati, N., Pettersson, M., Zhang, H., … Andersson, L.
(2016). The genetic basis for ecological adaptation of the Atlantic herring revealed by genome
sequencing. *ELife*, *5*(MAY2016), 1–32. https://doi.org/10.7554/eLife.12081
McDermid, J. L., Swain, D. P., Turcotte, F., Robichaud, S. A., & Surette, T. (2018). Assessment of the
NAFO Division 4T southern Gulf of St. Lawrence Atlantic herring (Clupea harengus) in 2016 and
2017. *DFO Can. Sci. Advis. Sec. Res. Doc. 2018/052*, xiv + 122 p.
McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo, M. A.
(2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation
DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110
McPherson, A., O'Reilly, P. T., & Taggart, C. T. (2004). Genetic Differentiation, Temporal Stability, and
the Absence of Isolation by Distance among Atlantic Herring Populations. *Transactions of the*
American Fisheries Society, *133*(2), 434–446. https://doi.org/10.1577/02-106
McPherson, A., Stephenson, R. L., O'Reilly, P. T., Jones, M. W., & Taggart, C. T. (2001). Genetic
diversity of coastal Northwest Atlantic herring populations: implications for management. *Journal of*
Fish Biology, *59*(SUPPL. A), 356–370. https://doi.org/10.1006/jfbi.2001.1769
McQuinn, I. H. (1997). Metapopulations and the Atlantic herring. *Reviews in Fish Biology and Fisheries*,
7(3), 297–329. https://doi.org/10.1023/A:1018491828875
Melvin, G. D., Stephenson, R. L., & Power, M. J. (2009). Oscillating reproductive strategies of herring in
the western Atlantic in response to changing environmental conditions. *ICES Journal of Marine*
Science, *66*(8), 1784–1792. https://doi.org/10.1093/icesjms/fsp173
Messieh, S. N., Anthony, V., & Sinclair, M. (1985). Fecundities of Atlantic herring Clupea harengus L.
populations in the Northwest Atlantic. *ICES C.M. 1985/H:8.*, 22 pp.
Metzger, B. P. H., Duveau, F., Yuan, D. C., Tryban, S., Yang, B., & Wittkopp, P. J. (2016). Contrasting
Frequencies and Effects of cis - and trans -Regulatory Mutations Affecting Gene Expression.
Molecular Biology and Evolution, *33*(5), 1131–1146. https://doi.org/10.1093/molbev/msw011
Nei, M. (1972). Genetic distance between populations. *The American Naturalist*, *1062*(949), 283–292.
https://doi.org/10.1086/285153
Nei, M. (2007). The new mutation theory of phenotypic evolution. *Proceedings of the National Academy*
of Sciences, *104*(30), 12235–12242. https://doi.org/10.1073/pnas.0703349104
Nosil, P., & Feder, J. L. (2012). Genomic divergence during speciation: causes and consequences.
Philosophical Transactions of the Royal Society B: Biological Sciences, *367*(1587), 332–342.
https://doi.org/10.1098/rstb.2011.0263
Nosil, P., Funk, D. J., & Ortiz-Barrientos, D. (2009). Divergent selection and heterogeneous genomic
divergence. *Molecular Ecology*, *18*(3), 375–402. https://doi.org/10.1111/j.1365-294X.2008.03946.x

Overholtz, W. . (2002). The Gulf of Maine–Georges Bank Atlantic herring (Clupea harengus): spatial pattern analysis of the collapse and recovery of a large marine fish complex. *Fisheries Research*, *57*(3), 237–254. https://doi.org/10.1016/S0165-7836(01)00359-9

Pagani, F., & Baralle, F. E. (2004). Genomic variants in exons and introns: identifying the splicing spoilers. *Nature Reviews Genetics*, *5*(May), 389–396.

Palumbi, S. R. (1994). Genetic Divergence, Reproductive Isolation, and Marine Speciation. *Annual Review of Ecology and Systematics*, *25*(1), 547–572. https://doi.org/10.1146/annurev.es.25.110194.002555

Panagiotou, O. A., & Ioannidis, J. P. A. (2012). What should the genome-wide significance threshold be? Empirical replication of borderline genetic associations. *International Journal of Epidemiology*, *41*(1), 273–286. https://doi.org/10.1093/ije/dyr178

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., … Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics*, *81*(3), 559–575. https://doi.org/10.1086/519795

Rambaut, A. (2007). FigTree. Retrieved May 20, 2018, from http://tree.bio.ed.ac.uk/software/figtree/

Ravinet, M., Faria, R., Butlin, R. K., Galindo, J., Bierne, N., Rafajlović, M., … Westram, A. M. (2017). Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. *Journal of Evolutionary Biology*, *30*(8), 1450–1477. https://doi.org/10.1111/jeb.13047

Revelle, W. (2018). psych: Procedures for Personality and Psychological Research. Northwestern University, Evanston, Illinois, USA. Retrieved from https://cran.r-project.org/package=psych

Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics*, *145*(April), 1219–1228.

Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals — mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics*, *15*(11), 749–763. https://doi.org/10.1038/nrg3803

Schluter, D. (2009). Evidence for Ecological Speciation and Its Alternative. *Science*, *323*(5915), 737–741. https://doi.org/10.1126/science.1160006

Scott, W. B., & Scott, M. G. (1988). *Atlantic fishes of Canada. Canadian Bulletin of Fisheries and Aquatic Sciences, bulletin 219*. Toronto, CA: University of Toronto Press.

Simmonds, E. J. (2007). Comparison of two periods of North Sea herring stock management: success, failure, and monetary value. *ICES Journal of Marine Science*, *64*(4), 686–692. https://doi.org/10.1093/icesjms/fsm045

Sinclair, M. (1988). Marine Populations: an Essay on Population Regulation and Speciation (p. 252). Seattle: Washington Sea Grant/Univ. Wash. Press.

Sinclair, M., & Iles, T. D. (1989). Population regulation and speciation in the oceans. *J. Cons. Int. Explor. Mer*, *45*, 165–175.

Smith, P. J., & Jamieson, A. (1986). Stock discreteness in herrings: a conceptual revolution. *Fish. Res.*, 223–234.

Stanley, R. R. E., DiBacco, C., Lowen, B., Beiko, R. G., Jeffery, N. W., Van Wyngaarden, M., … Bradbury, I. R. (2018). A climate-associated multispecies cryptic cline in the northwest Atlantic. *Science Advances*, *4*(3), eaaq0929. https://doi.org/10.1126/sciadv.aaq0929

Stanley, R. R. E., & Jeffery, N. W. (2017). CartDist: Re-projection tool for complex marine systems. https://doi.org/10.5281/zenodo.802875

Stephenson, R. L., Melvin, G. D., & Power, M. J. (2009). Population integrity and connectivity in Northwest Atlantic herring: a review of assumptions and evidence. *ICES Journal of Marine Science*, *66*(8), 1733–1739. https://doi.org/10.1093/icesjms/fsp189

Sylvester, E. V. A., Beiko, R. G., Bentzen, P., Paterson, I., Horne, J. B., Watson, B., … Bradbury, I. R. (2018). Environmental extremes drive population structure at the northern range limit of Atlantic

salmon in North America. *Molecular Ecology*, *27*(20), 4026–4040.
https://doi.org/10.1111/mec.14849
Tavares, H., Whibley, A., Field, D. L., Bradley, D., Couchman, M., Copsey, L., … Coen, E. (2018).
Selection and gene flow shape genomic islands that control floral guides. *Proceedings of the*
National Academy of Sciences, *115*(43), 11006–11011. https://doi.org/10.1073/pnas.1801832115
Teacher, A. G., André, C., Jonsson, P. R., & Merilä, J. (2013). Oceanographic connectivity and
environmental correlates of genetic structuring in Atlantic herring in the Baltic Sea. *Evolutionary*
Applications, *6*(3), 549–567. https://doi.org/10.1111/eva.12042
Tigano, A., & Friesen, V. L. (2016). Genomics of local adaptation with gene flow. *Molecular Ecology*,
n/a-n/a. https://doi.org/10.1111/mec.13606
Townsend, D. W., Thomas, A. C., Mayer, L. M., Thomas, M. A., & Quinlan, J. A. (2004). Oceanography
of the Northwest Atlantic Shelf (1, W). In A. R. Robinson & K. H. Brink (Eds.), *The Sea: The*
Global Coastal Ocean: Interdisciplinary Regional Studies and Syntheses (pp. 1–57). Harvard
University Press.
Turner, S. D. (2014). qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots.
BiorXiv. https://doi.org/https://doi.org/10.1101/005165
Turner, T. L., Hahn, M. W., & Nuzhdin, S. V. (2005). Genomic Islands of Speciation in Anopheles
gambiae. *PLoS Biology*, *3*(9), e285. https://doi.org/10.1371/journal.pbio.0030285
Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A., …
DePristo, M. A. (2013). From FastQ Data to High-Confidence Variant Calls: The Genome Analysis
Toolkit Best Practices Pipeline. In *Current Protocols in Bioinformatics* (Vol. 11, p. 11.10.1-
11.10.33). Hoboken, NJ, USA: John Wiley & Sons, Inc.
https://doi.org/10.1002/0471250953.bi1110s43
Vitti, J. J., Grossman, S. R., & Sabeti, P. C. (2013). Detecting Natural Selection in Genomic Data. *Annual*
Review of Genetics, *47*(1), 97–120. https://doi.org/10.1146/annurev-genet-111212-133526
Wang, Z., Brickman, D., Greenan, B. J. W., & Yashayaev, I. (2016). An abrupt shift in the Labrador
Current System in relation to winter NAO events. *Journal of Geophysical Research: Oceans*, *121*(7),
5338–5349. https://doi.org/10.1002/2016JC011721
Weir, B. S., & Cockerham, C. C. (1984). Estimating F-Statistics for the Analysis of Population Structure.
Evolution, *38*(6), 1358. https://doi.org/10.2307/2408641
Wheeler, J. P., & Winters, G. . (1984). Homing of Atlantic herring (Clupea harengus harengus) in
Newfoundland waters as indicated by tagging data. *Can. J. Fish. Aquat. Sci.*, *41*, 108–117.
Wu, C.-I. (2001). The genic view of the process of speciation. *Journal of Evolutionary Biology*, *14*(6),
851–865. https://doi.org/10.1046/j.1420-9101.2001.00335.x
Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration-selection
balance. *Evolution*, *65*(7), 1897–1911. https://doi.org/10.1111/j.1558-5646.2011.01269.x
Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common
statistical problems. *Methods in Ecology and Evolution*, *1*(1), 3–14. https://doi.org/10.1111/j.2041-
210X.2009.00001.x
1078
1079

1080   **Figures:**

1081

1082

1083

Figure 1. Geographic location and population structure among 14 spawning aggregations in the NW
Atlantic. (A) Map depicting sampling locations in the Northwest Atlantic. Location names as described in
Table 1. Pie charts indicate the proportion of individuals in a given gonadal maturity stage for each
spawning aggregation. Dark and light blue: maturing individuals (stages 3 and 4, respectively), light and
dark orange: ready-to-spawn and actively spawning individuals (stages 5 and 6, respectively), pink: spent
individuals (recently spawned) (stage 7), and black: individuals with resting gonads (stage 8) (Bucholtz et
al., 2008). **(B)** NJ phylogenetic tree based on Nei's distance calculated from population allele frequencies
of 2,189,380 SNPs (percent bootstrap support is shown for all branches, based on 1000 bootstrapping).
The collections clustered according to reproductive season into two main groups, spring (blue oval) and
fall (red oval) spawners with a few exceptions. BDO-S was in an intermediate position between these two
groups. The spring-collected sample NDB-S was closer to fall spawners. Within the fall spawners,

collections clustered depending on the latitude, forming the southern and northern sub-groups. **(C)**
Heatmap depicting pairwise F_{ST} estimates based on population allele frequencies of 2,189,380 SNPs
(values presented in Table S2). Samples are ordered by collecting season with "S" indicating spring and
"F" fall. Within season of collection, samples are ordered by latitude. Shading represents the degree of
genomic divergence. Pairwise F_{ST} ranged between 0.012 and 0.043, indicating overall and varying low
levels of population genetic structure. The most significant genomic differentiation was observed between
spring and fall spawning aggregations. Within spring spawners, the location SIL-S appeared as the most
differentiated, and within fall spawners, the greatest genetic divergence was observed between the two
southernmost collections (GEB-F and ME4-F) and all others. Notably, two spring-collected samples
(BDO-S and NDB-S) showed similar patterns of differentiation with respect to spring spawners as other
fall spawners. Collection name abbreviations are as defined in Table 1. **(D)** Plot of principal components 1
and 3 explaining 36% and 30%, respectively, of the genetic variation among 14 herring spawning
aggregations in the NW Atlantic (based on 2,189,380 SNPs). Each dot represents a spawning aggregation.
Colors indicate spawning season, blue for spring, red for fall, and yellow for mix. Aggregations were
distinguishable by spawning season along PC1 and by geographic origin along PC3.
1110

1111
1112
Figure 2. Genome-wide patterns of differentiation associated with seasonal reproduction and
latitude for 14 Atlantic herring spawning aggregations in the Northwest Atlantic. (A-C) Manhattan
plots depicting the genomic position of outlier SNPs and their respective significance association value (–
log10*P*-value) obtained with *pcadapt*, **(A)** for PC1 and **(C)** for PC3. Each dot of the Manhattan plots
represents a single SNP locus. For the purpose of visualization, only outlier loci per PC are displayed
(14,726 for PC1 and 6,570 for PC3). The top-ranked 500 SNPs based on importance values from a RF
classifier are highlighted in blue. SNPs reported in (Lamichhaney et al., 2017) as highly associated with
seasonal reproduction are emphasized in red in (A). SNPs within the four scaffolds showing the latitudinal
pattern in PC3 but present in PC1 were denoted in yellow (A). When available, annotation of the closest
gene 5Kb upstream or downstream of both, the top SNP per scaffold and the SNPs reported in our
previous study, are shown. The SNP annotated as *TSHR** falls within the first exon of the *TSHR* gene
(unpublished Leif Andersson com. pers.). **(B and D)** Heatmaps depicting standardized population allele
frequencies of the top 200 outlier loci distinguishing collections by **(B)** seasonal reproduction (PC1), and
(D) latitude (PC3). Each row in the heatmaps corresponds to a collection site and each column to a SNP.
SNPs were ranked in descending order based on their significance association value, -log$_{10}$*P*-value (from
left to right). Cell colors represent the population allele frequency of the major allele; thus, purple
indicates fixation of the major allele (allele frequency of 1) whereas orange represents fixation of the
minor allele (major allele frequency of 0).
1131

Figure 3. Functional characterization of outlier SNPs. (**A**) Functional classification of outlier SNPs associated with seasonal reproduction (dark gray) and with latitude (light gray), counts. (**B-C**) Bar plots showing the relative proportion of genes in each of the top GO terms associated with (**B**) seasonal reproduction and (**C**) latitude, for each biological category (i.e. biological process, cellular component, and molecular function). Top GO terms corresponded to the ones with P-value < 0.05 (ranked in ascending order based on their P-values obtained from ORA, Table S3 and S4, GO terms with an asterisk). Gene counts are indicated within parenthesis.

Figure 4. Genetic-environment association analysis. (A) Map depicting winter sea surface temperature averaged between 2008-2017 and the predominant population allele frequencies at diagnostic SNPs in the 14 spawning aggregations included in this study. A purple circle represents the prevalent alleles fixed in northern collections, a light purple square corresponds to southern collections with intermediate allele frequencies, and an orange tringle shows fixation of the minor alleles in the southernmost collection in Maine. **(B)** Redundancy analysis plot based on population allele frequencies of the top 500 outlier loci, with respect to latitudinal cline, ranked with a random forest classifier. Each circle corresponds to a spawning aggregation and their color indicates the predominant population allele frequency; labelling as in Fig. 4A. The vector of the most significant environmental variable is shown in blue, in this case it is surface sea temperature in winter months (January-February-March) (F = 16.7, p = 0.001). The length of the vector indicates its level of correlation with genetic variance. RDA1 explained 58.1% of the genetic variance, RDA2 is shown just for plotting purposes. **(C-D)** Random forest regression results. **(C)** Mean decrease in accuracy (MDA) for 12 environmental variables averaged across all runs and loci. Bars in light gray correspond to temperature values and bars in dark gray to surface salinity. Within temperature measures, surface is denoted with "S" whereas bottom is indicated with "B". Average sea surface winter temperature is the most important variable explaining genetic variance, followed by Fall sea surface temperature. Salinity variables were the least important. **(D)** Mean squared error (MSE) for each location averaged across all runs and loci. The southernmost collection had the largest MSE, followed by the collection in the Bay of Fundy and in the Scotian Shelf.

1163
1164
Figure 5. Isolation-by-distance (IBD) test for 14 NW Atlantic herring populations based on all
neutral and outlier SNPs or on latitude-related SNPs only. Regression between linearized genetic
distance ($F_{ST}/1$-F_{ST}), calculated from either 2,189,371 SNPs (red "X"s) or from 6,595 latitude-related
outlier SNPs (open black circles), and geographic distance (in km) between pairs of populations. The
dashed red line and the continuous black line correspond to the best fit line in each case. R^2 values indicate
the correlation between geographic and genetic distance matrices used in the Mantel test (Mantel's test for
all SNPs: P < 0.001, $R^2 = 0.04$, 9999 replicates; for latitude-related SNPs only: P < 0.001, $R^2 = 0.30$, 9999
replicates). Note the IBD pattern is only observed in the latitude-related outlier SNPs, not in all SNPs.
1173
1174
1175

1176 **Tables:**

1177

1178 **Table 1.** Characteristics of the 14 herring spawning aggregations included in this study.

1179

Locality	Code	Sample size (N)	Geographic coordinates (longitude, latitude)		Sampling (dd/mm/yy)	Season	Salinity (PPM)	Sequencing year
Seven Islands	SIL-S	50	-66.33	50.09	06/06/2012	Spring	35	2016
Stephenville	SPH-S	48	-57.94	49.73	30/05/2012	Spring	35	2016
Northumberland Strait	NTS-S	50	-64.12	46.30	14/05/06	Spring	35	2015
Labrador	LAB-F	50	-55.50	52.25	24/08/2014, 22/08/2015	Fall	35	2016
Blanc Sablon	BLS-F	49	-57.31	51.38	13/08/2014	Fall	35	2016
Notre Dame Bay	NDB-S	50	-55.44	49.55	03/05/2015	Spring	35	2016
Notre Dame Bay	NDB-F	50	-55.47	49.55	26/10/2015	Fall	35	2016
Trinity Bay	TRB-F	50	-53.47	47.84	28/09/2014	Fall	35	2015
Miramichi	MIR-F	50	-63.96	47.04	25/08/2014	Fall	35	2016
Bras D'Or lake	BDO-S	50	-60.85	45.93	20/04/2016	Spring	25	2016
Scots Bays	SCB-F	50	-64.92	45.17	24/08/2015	Fall	35	2016
Musquodoboit	MUS-F	50	-63.10	44.63	28/10/2015	Fall	35	2016
German Banks	GEB-F	50	-66.33	43.45	28/08/2014	Fall	35	2015
Maine fishing area 514	ME4-F	50	-70.41	42.09	19/10/2015	Fall	35	2016

1180