1 **The interaction of compliance and activation on the force-length operating**

2 **range and force generating capacity of skeletal muscle**

3 S.M. Cox[1]*, K.L. Easton[2,6], H. Sanghvi[2,3], M. Cromie Lear[4], R.L. Marsh[5,6], S.L. Delp[4,7], J.

4 Rubenson[1]

[1]Biomechanics Laboratory, Kinesiology Department, The Pennsylvania State University,

University Park, PA 16802, USA;

[2]School of Sport Science, Exercise and Health, The University of Western Australia, Perth, WA,

6009, Australia

[3]Department of Biomedical Engineering, Linkoping University, 581 83 Linköping, Sweden

[4]Department of Mechanical Engineering, Stanford University, Stanford, CA, 94305, USA

[5]Department of Ecology and Evolutionary Biology, Brown University, Providence, RI, 02912,

USA

[6]Department of Biology, Northeastern University, Boston, MA, USA

[7]Departments of Bioengineering and Orthopedic Surgery, Stanford University, Stanford, CA,

94305, USA

*Corresponding Author E-mail: zanne@psu.edu Phone: 603-320-9955

Running Title: Activation dependent effects on muscle length

Abstract

Muscle and locomotor performance is influenced, in large part, by where muscles operate on their force-length curve. Here we explore how activation and tendon compliance interact to influence the operating lengths and force-generating capacity. To study this, we built a musculoskeletal model of the lower limb of the guinea fowl. To quantify the activation-compliance effects with this model, we simulated the force-length operating range during fixed-end fixed-posture contractions for 39 actuators under thousands of combinations of activation and posture and using three different muscle models: Muscles with non-compliant tendons, muscles with compliant tendons but no activation dependent shift in optimal fiber length (activation-dependent shifts in L0), and muscles with both compliant tendons and activation-dependent shifts in L0. We found that, at the extreme, activation dependent effects altered muscle fiber lengths up to 40% and increased or decreased force capacity up to 50% and 80%, respectively, during fixed-end contractions. Typically, activation-compliance effects reduce force generation and are dominated by the effects of tendon compliance at high activations. At low activation, however, activation-dependent shifts in L0 are equally important and can result in relative force changes for low compliance muscles of up to 60%. Because of the non-linear nature of these interactions, predicting their effects at the individual muscle or joint level are not straightforward. There are regions of the force-length curve in which muscles are most sensitive to compliance and there are troughs of influence where these factors have little effect on length and force capacity. These regions are hard to predict, though, because the magnitude and location of these areas of high and low sensitivity shift with compliance level. Here we provide a map for when these effects will meaningfully influence force capacity and an example of their contributions to force production during a static task, namely standing.

Introduction

Where muscles operate on their force-length relationship has important implications for muscle and locomotor performance. Most tangibly, muscle length affects muscle force and, therefore, joint torque capacity (Arnold et al., 2013; Blix, 1894; Gordon et al., 1966; Huxley and Simmons, 1971; Morgan et al., 2002; Rack and Westbury, 1969; Ramsey, R. W. & Street, 1941; Rassier et al., 1999; TerKeurs et al., 1978). One relatively unexplored, but potentially significant, factor influencing muscle operating lengths is muscle–tendon compliance. Zajac (Zajac, 1989) initially suggested that even in "isometric" (fixed-end) contractions, muscle fibers will shorten due to the stretch of the tendon, an effect exaggerated at higher activation levels due to greater forces. The result is that the operating ranges for muscles with compliant tendons shift to the left on the force-length curve with increasing activation. This theoretical prediction has been experimentally confirmed in several studies (Arnold and Delp, 2011; Azizi and Roberts, 2010; Fukunaga et al., 1997; Hawkins and Bey, 1997; Holt and Azizi, 2016; Lakatta and Jewell, 1977; Lemos et al., 2008; MacIntosh and MacNaughton, 2005; Mayfield et al., 2016; Rubenson et al., 2012; Sugisaki et al., 2011). More recently, it has been shown that tendon compliance is an important factor affecting muscle force-length operating ranges during human gait (Arnold and Delp, 2011). Yet, despite many studies demonstrating this effect for individual muscles or during specific tasks, we still do not have a broad understanding of when tendon compliance has a meaningful influence on muscle operating length and function. For instance: Is there a threshold of muscle-tendon compliance where the effect becomes functionally relevant? Does compliance have a consistent influence on the F-L operating range in all conditions (i.e. across different postures and muscle lengths) or are there conditions when it is more influential?

Complicating these questions is the

observation that, even in the absence of

muscle-tendon compliance, activation

can alter optimal muscle lengths.

While the exact mechanisms remain

unclear, a rightward shift of the plateau

region of the force-length curve occurs

with decreasing activation (Holt and

Azizi, 2014; Rack and Westbury,

1969; Roszek et al., 1994) such that

the optimal muscle length for force

Box 1: Glossary	
Muscle tendon unit:	MTU
Optimal muscle fiber lengths	L0
Sarcomere length	L_s
Fascicle length	L_f
Muscle length at 50% activation	L_{A50}
Passive muscle length	L_p
Physiological cross-sectional area	PCSA
Compliance: the ratio of optimal fiber length to tendon slack length	T/M ratio
Muscle Models	
-**N**o tendon **C**ompliance **N**o activation dependent **S**hift in optimal fiber length	NCNS
-Tendon **C**ompliance with **N**o activation Dependent **S**hift in optimal fiber length	CNS
-Tendon **C**ompliance and activation Dependent **S**hift in optimal fiber length	CS

production increases. Although an interaction between activation and compliance on muscle

operating lengths has been demonstrated (Arnold and Delp, 2011; Arnold et al., 2013; Lemos et

al., 2008; Lieber and Brown, 1992; Rubenson et al., 2013), no studies have systematically

explored the simultaneous influence of activation-dependent optimal fiber length and tendon

compliance, and its subsequent effect on force-length operating ranges. Nor have the relative

contributions of compliance and activation dependent shifts of the force-length curve been

quantified. Therefore, we do not know under what conditions we can or cannot safely ignore

these complicating factors in interpreting and predicting how muscle force varies with length

across any species.

The purpose of this study is to undertake the first

comprehensive assessment of how tendon

elasticity and activation-dependent shifts in

optimal fiber lengths combine to influence

force-length operating ranges. To this end, we

integrated a comparative and computational

approach, the combination of which is well

suited to illuminate these relationships. We

developed a computational musculoskeletal

model of the guinea fowl pelvic-limb (Fig. 1A),

a popular avian bipedal model for biomechanical

studies (Daley and Biewener, 2011; Daley et al.,

2006; Ellerby et al., 2005; Ellerby and Marsh,

2010; Gordon et al., 2015; Marsh et al., 2006;

Rubenson and Marsh, 2009). An animal

computational model has the advantage of

including highly detailed measurements of

musculoskeletal parameters, many of which are

Figure 1.Methodological approach. A) Musculoskeletal model developed in SIMM and implemented in OpenSim (see supplemental material. B) Since several muscles cross multiple joints, we varied starting MTU length by changing the posture and initial joint angle.

typically not included in current human models (see Supplemental Material). Additionally, in

this animal model the scope of muscle-tendon compliance (Table 1) is twice that of humans

(Arnold and Delp, 2011), amplifying the effects. Using a computational approach allowed us

iterate many more combinations of activation and posture than would be possible

experimentally. We simulated the force-length operating range for 39 actuators (Table 1) under

thousands of combinations of activation and posture to distill the influence of tendon elasticity

and activation on force-length operating range (Fig. 1). This computational approach enabled us

to extrapolate overall patterns and to systematically tease out the contributions of individual

factors; for example, the role of the initial passive (pre-activation) lengths of muscles, activation

dependent optimal fiber lengths, and tendon compliance on muscle operating length.

We addressed three questions: 1) Averaging across all muscles, how does activation level

interact with muscle-tendon unit (MTU) compliance, initial passive muscle length and/or

activation-dependent shifts in L0 to alter the length and force generating capacity of muscles? 2)

Under what combinations do these factors meaningfully influence force generating capacity of

an individual muscle? 3) How do these factors influence the force and/or torque generating

capacity during a low-activation isometric task like standing?

Methods

OpenSim Model development

Our approach to explore these questions was to build a realistic detailed musculoskeletal model.

This was done first in SIMM software (Delp and Loan, 1995) and subsequently converted to

OpenSim (Seth et al., 2018). This model incorporated several experimental measurements of

muscle and skeletal properties as well as steps to validate the model accuracy (Fig. 2). A

detailed description of each modelling step is provided in the Supplemental Material and

summarized below.

The modeling began by digitizing the 3D muscle-tendon paths of the model animal (Female,

1.45 kg). This was achieved by isolating the pelvic limb and systematically dissecting off each

muscle-tendon-unit (MTU) from the right limb (kept fresh/frozen). The muscle-tendon path

from origin to insertion was traced using an optical tracking system (Polaris, Northern Digital,

Waterloo, ON). A total of 39 MTUs were defined, with some large muscles being divided into

sub-muscles. Care was taken to capture the geometry of MTU paths across articulating surfaces,

which were used to define muscle wrapping surfaces in SIMM software. Muscle architecture was

measured on the model specimen as well as four additional animals (2 Females, 2 Males; 1.46 $\pm$

0.1 kg; mean $\pm$ SD) for comparison. Experimental muscle architecture measurements included

the muscle mass and free tendon mass and length. The left limb was fixed (10% neutral buffered

formalin) in mid-swing posture and used to measure muscle fiber lengths (L_f) and pennation

angles. Bundles of fascicles were isolated and used for sarcomere length measurements (L_s)

based on second harmonic generation using two-photon laser microscopy (Cromie et al., 2013).

Optimal muscle fiber lengths (L0) were defined as $L0 = L_f \cdot {2.36}/{L_s}$, where 2.36 is the length in

microns of the optimal sarcomere length in guinea fowl muscle (Carr et al., 2011; Llewellyn et

al., 2010). The Physiological Cross-sectional Area ($PCSA$) of the muscle was calculated as from

the L0, muscle mass (m_{mus}) and density (ρ_{mus}, 0.00106 g/mm^3) as: $PCSA = \frac{m_{mus}}{\rho_{mus} \cdot L_0}$. Maximal

isometric force (F_{100max}) for each muscle unit was calculated using a specific tension of 0.3

N/mm^2.

Anatomical and functionally relevant bone coordinate systems (BCS) were defined,

incorporating mathematically derived joint centers and axes. For the knee, ankle and

tarsometatarsus-phalangeal (TMP) joint, a helical axis and functional joint center were computed

from 3D motion capture data of adjacent segment motion (Rubenson et al., 2007). Ankle joint

translation and patella-complex motion was defined as a function of ankle and knee joint angle,

respectively (measured from 3D motion capture). The hip joint center was defined by directly

digitizing its location with the 3D pointer. Skeletal elements were 3D-scanned to generate bone

Figure 2. Model development framework and main steps. See Methods section and Supplementary Material for details.

mesh (.ply) files. The bone model files and muscle-tendon paths were transformed into the

relevant BCSs that were used to define the final model's joint coordinate systems (JCS; see

Supplementary Material for a full definition of BCSs and JCSs).

We measured tendon material properties from two separate animals (1F, 1.43 kg; 1M 1.49 kg) of

the free common tendon of the lateral, medial and intermedius gastrocnemius muscles (Achilles),

and the free tendon from the tibialis cranialis, digital flexor IV and extensor digitorum longus

(Bose EnduraTEC, ELectroForce 3200, Framingham, MA, USA). The tendons were

programmed to undergo a 5Hz sinusoidal cycle that approximated the duration of the stance

phase / swing phase of gait (Rubenson and Marsh, 2009). The clamp displacement was

programmed to produce force approximating F_{100max}. These data were used to generate muscle-

specific tendon load-elongation curves for the tested muscles and to generate a generic (average)

load-elongation curve for the remaining tendons. The tendon slack lengths were defined by

constraining the passive simulated muscle fiber length to match the experimental fiber length in

joint postures matching the experimental specimen. Segment inertial properties for the model

were obtained from previously collected data from six animals (See Rubenson and Marsh, 2009).

Finally, to assess the accuracy of our model we compared simulated data to experimental values

for both muscle moment arms and passive net joint moments.   Muscle moment arms were

computed from tendon travel experiments measured post mortem in separate specimens (4F, 2M;

1.47 ± 0.1 kg; mean $\pm$ SD). Passive net joint moments were measured for the hip joint (proximal

muscles) and ankle joint (distal muscles) in deeply anesthetized and nerve-blocked animals,

separate to that of the model specimen (3F, 2M; 1.49 ± 0.1 kg; mean ± SD). Details of these

experiments and results are presented in the Supplemental Material.

Computation of fiber lengths during isometric (fixed-posture) contractions

In this musculoskeletal model, we simulated fixed-end contractions across a range of activation

levels and limb configurations (Fig. 1B) intended to elicit the maximum *in vivo* range of muscle

lengths. MTU lengths were roughly prescribed by setting a posture that set an initial passive

length (short, medium and long) and then imposing smaller deviations in MTU lengths around

that initial position though variation of either the ankle (distal muscles) or hip angle (proximal

muscles) through 95% of the range for that joint during locomotion reported in the literature

across various experiments (Ellerby and Marsh, 2006, 2010; Henry et al., 2005; Rubenson and

Marsh, 2009). Each posture generated a range of passive MTU lengths across the limb

determined by fiber length, moment arm and tendon slack length of each muscle. At each of

these postures we implemented three different muscles models (1: No Compliance, No

activation-dependent shift in L0 (NCNS), 2: Compliance with No activation-dependent shift in

L0 (CNS) and, 3: Compliance with a 15% activation dependent Shift in L0 (CS) and simulated

fixed-end contractions at 5 activation levels (0%, 25%, 50%, 75%, 100%). Simulations for

muscle models without activation-dependent shift were performed on the unaltered model with

and without tendon compliance. Including activation dependent shifts of optimal fiber length

required altering the force-length curve for activation levels below 100%. For each activation

level, the force-length curve for all muscles was replaced in the OpenSim model with a shifted

curve (see supplemental material for details). A 15% activation-dependent shift in L0 (at zero

activation) was used because it represents a typical value reported in the literature for both

voluntary muscle activation and electrically stimulated isolated muscle under realistic

stimulation frequencies (de Brito Fontana and Herzog, 2016; Rack et al., 1969; Roszek et al.,

1994), and also because it is a value previously adopted in modeling of muscle force (Buchanan

et al., 2004; Lloyd and Besier, 2003). For each simulation, we extracted the normalized active

individual fiber force and normalized fiber length for each muscle. Passive muscle lengths were

defined as the muscle length at 0% activation. This resulted in 46,800 MTU lengths (Fig. 1B).

These results were analyzed in three ways.

Table 1. Tendon slack length (TSL), optimal fiber length (OFL) and their ratio listed for each muscle used for
analysis. Muscles are organized by group (proximal vs distal) and mean T/M Ratio (TSL/OFL) for each group
listed.

Muscle	TSL(m)	OFL(m)	T/M Ratio	Muscle	TSL(m)	OFL(m)	T/M Ratio
Hip flexion/Extension Muscles				Ankle Flexion/Extension Muscles			
CFP	0.027	0.037	0.71	DFii_deep_ant	0.182	0.011	16.55
FCLP_a	0	0.120	0.00	DFii_deep_post	0.185	0.015	12.33
FCLP_p	0.078	0.054	1.46	Deep_DFiii_major	0.196	0.016	12.33
IC_1	0	0.086	0.00	Deep_DFiii_minor	0.195	0.013	14.77
IC_2	0	0.085	0.00	EDL_dist2_DIGIII	0.193	0.012	15.69
ILPO_a	0.064	0.033	1.93	FHL_III	0.239	0.020	11.75
ILPO_b	0.017	0.083	0.21	FL_lat	0.084	0.049	1.73
ILPO_c	0	0.111	0.00	FL_mid	0.081	0.049	1.67
ILPR_a	0.044	0.041	1.09	IG	0.075	0.072	1.05
ILPR_b	0.057	0.026	2.19	LG	0.102	0.047	2.17
ISF_1	0.039	0.014	2.76	MG_lat	0.105	0.041	2.57
ISF_2	0.039	0.014	2.76	MG_mid	0.118	0.037	3.17
ITC_deep	0.019	0.023	0.83	MG_most_medial	0.120	0.028	4.23
ITC_r	0.006	0.023	0.28	TC_knee	0.119	0.029	4.07
ITC_super	0.012	0.028	0.42	TC_tibia	0.072	0.051	1.42
ITM	0.008	0.023	0.33	sup_DFII	0.198	0.013	15.19
PIFL_caud	0	0.075	0.00	supDFIII	0.183	0.037	4.97
PIFL_cran	0	0.049	0.00	FDL_III	0.214	0.037	5.78
PIFM_caud	0	0.075	0.00	**Mean TRF**			**7.82**
PIFM_cran	0	0.040	0.00				
FCM	0.003	0.74	0.01				
Mean TFR			**0.71**				

1. On average, how does activation level interact with MTU compliance, initial passive

muscle length and/or activation-dependent shift in L0 to influence the operating

range and force capacity of muscles?

To discern whether there are any useful broad patterns of interaction between MTU compliance,

(i.e. the ratio of tendon slack length to optimal fiber length, Table 1), initial passive muscle

length and activation dependent shifts in L0, we binned data into two levels of compliance

(Low< 2< High), and passive muscle length [ascending (A), plateau (P) or descending (D) limb

of the force-length curve] and evaluated how muscle length changed with activation level

between muscle models. For each group (see Table 1), we calculate average normalized muscle

length [$L/(L0_{100})$] and force (F/ F_{100max}) across all muscles in each group (Table 2) at each level

of activation (0%,25%,50%,75%,100%). Additionally, at each activation level we calculated

how muscle length changed with activation (ΔL) from its passive starting length. For instance,

ΔL at 50% activation would be given by

$$\Delta L \;=\; L_{A50} - L_{A0}$$

Where the muscle length at 50% and 0% activation (normalized to L0 at 100% activation) for

this passive length is given by L_{A50} and L_{A0} respectively. Main and interaction effects were

evaluated with a nested linear mixed effects model (Bates et al., 2015) with ΔL as the dependent

variable, muscle as a random factor and activation level (0,25,50,75,100), activation-dependent

shift in L0 (0,15 %), passive length of the muscle (A, P, D) as independent factors. Given that

activation dependent shifts in the optimal fiber length did not show a main or any interaction

effects, the statistical models were run both with and without this factor. Throughout this paper,

muscle lengths are reported as normalized by the optimum fiber length at 100% activation, $L0_{100}$, and forces by maximum force at 100% activation, F_{100max}.

Table 2. Triple nested experimental design

MTU compliance (T/M ratio)		Low			High		
Passive length (Ls)		Ascending	Plateau	Descending	Ascending	Plateau	Descending
Activation-dependent shift in L0 (S_{LO})	0%	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100
	15%	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100	1,25,50,75,100

2. Under what combinations of these factors is the force generating capacity of an individual muscle meaningfully altered? Specifically, we asked how compliance and activation-dependent shift in L0 alter muscle operating length and force capacity at high (100%) and low (25%) activation and which combinations of these factors alter the relative force capacity substantially.

Our second analysis aimed to quantify our results at a finer level of detail. While in the first analysis we binned our data into two compliance levels and three different passive length ranges, here we present the length and force data for each muscle at all simulated passive lengths (9216 iterations). Again, we isolated the influence of compliance and activation-dependent shifts in L0 by comparing resultant muscle lengths and forces between the three muscle-tendon models (NCNS, CNS, CS). Differences between models with and without compliance isolate the influence of compliance alone. Likewise, differences between models with compliance but with

and without activation-dependent shifts in L0 allow us to isolate the contribution of activation-

dependent shift to changes in operating length and force capacity. Specifically, the contribution

of activation-dependent shift in L0 to muscle operating length (ΔL) for a given muscle, passive

length and activation level was found by subtracting the muscle length of the CNS condition,

L_{CNS}, from that of the CS condition, L_{CS}.

$$\Delta L_{ADshift} = L_{CS} - L_{CNS}$$

The force values for each muscle with no compliance or shift (i.e. non-compliant tendon, NCNS)

were found by assuming no change in muscle length across activation levels and scaling the

force with activation. For example, for a muscle with a passive length of 0.809 at 100%

activation, F_{100}, is 0.726. At 50% activation, the calculated force with no compliance would be

0.363 or

$$F_{50} = 0.5 * F_{100}.$$

Since the significance of a change in muscle length or force will vary by context, we chose to

present these data in two ways; as absolute changes normalized by maximum force capacity at

100% activation (F_{100max}) and as relative changes between models. Absolute changes were

plotted for all data points as a function of MTU compliance and passive length normalized to

muscle length at 100% activation, $L0_{100}$ (Fig. 3). Absolute changes relative to maximal force

allowed us to quantify the contribution of each factor as well as painting a more detailed picture

of the interaction between these factors and the conditions in which each is most influential. Yet,

this approach can obscure the significance of changes at low activations. For instance, at 25%

activation, a change in force capacity of 5% of maximum force between models may represent a

relative change of anywhere from 20 to 80% of force capacity. To capture these relative effects,

we again binned our data, but this time into much smaller bin sizes (steps of 5% of Lo, T/M ratio

bins: 0.2 for low compliant muscles and 1 for high) and compared across models as described

above (NCNS, CNS, CS). For each bin we calculated the number of samples in each bin and the

percent of samples with a greater than 10% difference in force capacity between models. The

10% difference is an arbitrary cut-off below which we deemed the activation-compliance effect

to be of less importance in dictating muscle function. The percent difference between muscle-

tendon models was taken from the difference between the forces predicted by each model

divided by the force predicted by the less complex model. For example, the percent difference

between models with and without compliance (NCNS and CNS), was given by

$$\%\Delta F\ =\ 100\ *\ \frac{F_{CNS} - F_{NCNS}}{F_{NCNS}},$$

where F_{CNS} and F_{NCNS} are the forces predicted by the CNS model and NCNS models

respectively. With this analysis we aimed to identify a threshold of MTU compliance or range of

operating lengths where these factors can be safely ignored and the conditions in which simple

models that exclude these factors lead to errors.

3. What are the functional consequences of these factors? Specifically, how do these

activation dependent factors sum over many muscles acting across a joint to influence the

posture for maximal force and moment capacity at low activation (25%) and how does

this influence compare between a distal and proximal joint?

To explore these questions, we compared force and torque generated by the sum of muscles

acting across the ankle and the hip. We chose to evaluate the muscles acting across the hip and

ankle since on average they are composed of MTU's with very different levels of compliance

(T/M Ratio Hip:0.71, Ankle:7.82, Table 2) and did not overlap. We chose a consistent 25%

activation to match the lowest level of activation used to evaluate our second question above and

made it uniform to simplify the analysis. We simulated 25% activation of muscles acting across

these joints for each model (NCNS, CNS, CS) through a sweep of hip and ankle postures with

knee and TMP angle held constant at standing angles [Ankle and hip angle ranges as described

in Fig. 1. $Knee_{flex}$:-110^{o}, TMP_{flex}:-55^{o} (Henry et al., 2005)]. For each joint angle we extracted

the total force and torque acting across each joint as well as the mean moment arm weighted by

maximal muscle force, F_{100max} (Fig. 5). Maximum total muscle force and torque were found for

each joint for each model as well as the corresponding joint angle (Table 4). For reference, we

then calculated the percent difference between these joint angle-dependent values and the force

or torque in the standing posture. The goal of this analysis was to provide an estimate of the

combined influence of the activation-compliance dependent effects across many muscles at low

activation.

Results

1. On average, how does activation level interact with MTU compliance, passive

muscle length and/or activation dependent shifts in optimal fiber length to influence

the operating range and force capacity of muscles?

We found passive muscle length, activation and MTU compliance influenced how a muscle

changes length with increasing activation, while activation dependent shifts in optimal fiber

lengths did not (Fig. 3, Table 3). There were neither significant main nor interaction effects with

activation-dependent shifts in L0. When models were re-run without including activation-

dependent shifts in L0, we found 3rd order interaction terms between passive muscle length,

activation level and MTU compliance (Table 3). This implies that how muscle length changes

with activation level differs between muscle of low and high compliance and that this

relationship changes with the starting passive length of the muscle.

This complex interaction can be seen in Figure 3. While both high and low compliance muscles

become shorter with increasing activation, high compliance muscles show a larger effect. The

extent of the effect is also dependent on passive muscle length, with muscles with passive

lengths on the descending limb of the force-length curve showing larger length changes with

increasing activation for all levels of compliance (Fig. 3). This is explained on the basis that

force capacity on the descending limb, and thus the compliance effect, increases as the muscle

shortens towards L0.

Figure 3. The influence of activation, MTU compliance, starting muscle length and activation dependent shifts in L_0 on average muscle length. Circles represent the mean value at the five different activation levels and circles from the same group are connected by lines across activation. $L0_{100}$ is always a normalized fiber length of 1. Yellow circle marks the maximum force at 100% activation, F_{100max}. On average, muscle length decreases with increasing activation and is significantly influenced by interactions between starting length, compliance and activation level. Muscles with low MTU compliance (A) change length with activation less than muscles with high compliance (B). While AD-shifts in L_0 change operating length (difference between black and grey filled circles), they are not significant.

Table 3. Model parameters and results for linear mixed effects models with change in muscle length as dependent factor. Activation dependent shift in $L0$ (Shift) did not have significant main or interaction effect. Activation level, MTU compliance (TFR) and passive limb of the force-length curve all interact to influence operating muscle length.

Model		Shift*TMR*Limb*Act				TMR*Limb*Act			
	DF	Sum sq	Mean sq	F val	p-val	Sum sq	Mean sq	F val	p-val
Shift (1)	1	0.02	0.02	17.4	0.15				
T/M Ratio (2)	1	0.94	0.094	83.1	0.07	0.097	0.097	83.08	0.07
Limb (3)	**2**	**13.07**	**6.53**	**5768.2**	**1.7e-4**	**6.8**	**3.3**	**2808.9**	**3.6e-4**
Act (4)	**4**	**83.87**	**20.96**	**18502.8**	**1.75e-8**	**41.9**	**10.5**	**8964.4**	**7.4e-8**
1x2	1	0.017	0.167	14.70	0.01				
1x3	2	0.016	0.008	7.03	0.01				
2x3	**2**	**5.95**	**2.97**	**2625.2**	**3.08e-4**	**3.04**	**1.5**	**1300**	**7.7e-4**
1x4	4	0.018	0.005	4.07	0.1				
2x4	4	**39.83**	**9.96**	**8787.3**	**7.8e-8**	**19.98**	**4.99**	**4265.6**	**3.3e-7**
3x4	8	**14.49**	**1.81**	**1598.6**	**1.6e-10**	**7.5**	**0.94**	**804.5**	**2.6e-9**
1x2x3	2	0.005	0.002	2.27	0.2				
1x2x4	4	0.013	0.003	2.95	0.16				
1x3x4	8	0.022	0.003	2.37	0.16				
2x3x4	**8**	**5.78**	**0.722**	**637.4**	**6.4e-9**	**3.02**	**0.38**	**322.1**	**9.5e-8**
1x2x3x4	8	0.008	0.001	0.89	0.3				

2. **Under what combinations of these factors is the force generating capacity of an individual muscle meaningfully altered?** *How does compliance and activation dependent shifts in optimal fiber length alter muscle operating length and force capacity at high (100%) and low (25%) activation?*

<u>The influence of compliance</u>: Increasing compliance increases how much a muscle shortens with increasing activation. At 100% activation (Fig. 4A), the change in length can be as high as 40% fiber strain while at 25% activation that decreases to 25% muscle fiber strain (Fig. 4B). At high activation, the starting passive length which results in the greatest length changes increases with MTU compliance (Fig. 4A, shifting from ~ 1.0 to 1.25 L0) while at low activations the passive length that produces the greatest length changes is less influenced by compliance (Fig. 4B, shifting only from ~ 1.0 to 1.05 L0).The consequence of these length changes on the force

capacity depend on the passive muscle length, generally hampering force generation at short

passive lengths and amplifying force capacity at long passive lengths (Fig. 4D). Though, it is

interesting to note that change in force capacity does not linearly scale with passive muscle

length. At the tails of the force-length curve (short and long extremes), the force effects are

smaller due to the decreased force capacity. Thus, the passive length that results in the greatest

change in muscle length does not occur at the extremes nor does it align with the length that

results in the greatest change in force. The transition between decreasing and increasing force

capacity as a result of activation-compliance effects occurs at longer passive muscle lengths for

more compliant muscles. At 100% activation, the transition shifts from a normalized length of 1

at low compliance to 1.2 for high compliant muscles. The results at 25% activation, again, show

similar patterns, but with less dramatic variations (Fig. 4E).

<u>The influence of activation dependent shifts in optimal fiber length</u>: In comparison to the

Figure 4. The influence of compliance (A,B,D,E) and an activation dependent shift (C&F) in the optimal fiber length on muscle length (A-C) and force (D-E) as a function of the starting fiber length and MTU compliance (Designated by color).

influence of compliance, activation dependent shifts in optimal fiber length result in much

smaller changes in normalized muscle length. Muscle lengths change by at most 2% of $L0_{100}$

(Fig. 4C). Unlike the influence of compliance, though, shifts in optimal fiber length result in

longer as well as shorter muscles depending on passive muscle lengths. While intuition may

suggest that activation-dependent shifts alone would not alter a muscle's length, the variations

are likely the result of muscles finding an equilibrium with the compliant tendon. For a muscle

on the ascending limb of the force-length curve, for instance, a shift in the force-length curve to

the right decreases its force capacity. This results in the muscle being able to stretch the tendon

less. Thus, for any fixed muscle-tendon length, the equilibrium length of the muscle on the

ascending limb is longer in the presence of activation-dependent shifts. The opposite effect can

be seen on the descending limb (Fig. 4C), and again, these effects are amplified for more

compliant MTU's.

In contrast to the influence of compliance, which has little effect on the force generating capacity

of low compliance muscles, activation-dependent shift in $L0$ influence the force capacity of

muscles of all compliance levels, though the greatest effects are still seen in the most compliant

MTU's. Activation-dependent shifts in $L0$ can result in both decreased and increased force

capacity, depending on passive muscle length and compliance. In general, muscles with passive

lengths less than ~1.1 $L0$ decrease force capacity in the presence of activation-dependent shifts

while muscles above a normalized length of 1.1 $L0$ increase force capacity. The magnitude of

the change in force capacity with shift matches or exceeds the influence of compliance at 25%

activation.

In summary, at high activation, compliance results in muscle strain up to 40%. These changes

result in increases of force capacity for muscles on the descending limb of the force-length curve

(up to 60% of maximum force, F_{max}) and decreases on the ascending limb (of up to 80% of

maximum muscle force capacity). Muscle force is most sensitive to compliance effects on the

shallow ascending limb (Ls~0.7-0.9) or the middle descending limb (Ls~1.2-1.3) of the force-

length curve. At low activations, though activation-dependent shifts in L0 result in only small

changes in muscle length (at most 1-2% strain), they have a larger effect on force capacity than

compliance (Fig. 4F vs Fig. 4E) since the peak of the force-length curve is changing relative to

the muscle length. While the influence of activation-dependent shifts in L0 on force capacity

decreases with compliance, for muscles with no tendon, they can be as high as 5% of maximum

force capacity.

Which combinations of these factors alter the relative force capacity by more than 10%?

<u>Influence of compliance</u>: The range of passive lengths that result in a change of force capacity

increases with compliance and activation, as expected. For the most compliant MTU's, 88% of

starting passive muscle lengths result in a change in force capacity that is greater than 10% (Fig.

5A). While our data does not allow us to fully quantify the space, it does allow us to isolate

regions of high and low sensitivity to compliance. For instance, at both 100 and 25% activation,

all MTU's with a T/M ratio of as low at 0.8 show a meaningful (>10%) influence of compliance

on force capacity when the muscle's initial passive length is between 0.7 and 0.75 L0 (Figs.

5A&B). Thus, even low compliance MTU's have a region of operating lengths where

compliance meaningfully influences force capacity. There also exists a trough of influence where

compliance and AD shifts of L0 have little influence on force capacity. The location of this

trough varies with MTU compliance, though, shifting slightly to longer muscle lengths with

increasing compliance. For example, MTU's with a T/M ratio of 1.2 show no change in force

capacity with compliance when operating at 1.05 L0, while the passive length that has the

smallest influences on the force capacity of the most compliant muscles is closer to 1.15 at 25%

activation and 1.25 at 100% (Fig. 5C).

<u>Influence of activation-dependent shifts in L0</u>: Unlike the influence of compliance, activation-

dependent shifts alter the force capacity of muscles of all levels of compliance. While

compliance had very little influence on the force capacity for muscles with very little tendon,

activation-dependent shifts in L0 resulted in changes in force capacity of over 10% for muscles

with no tendon (T/M ratio = 0) across half of the operating range, primarily at lengths on the

ascending limb. Again, the influence increased with increasing compliance, such that for the

most compliant muscles activation-dependent shifts in L0 resulted in changes in force capacity

for all muscles across 95% of the possible initial passive lengths of the muscle. The trough of

influence for activation-dependent shifts in L0 followed similar patterns to that for compliance,

with minimal effect on force for stiff MTU's at lengths of 1.05 and for compliant muscles at

1.15.

To summarize, at high activation compliance considerably changes the force capacity for

muscles with a T/M ratio over 2 at nearly all passive lengths (95%). While change in force

capacity increases with compliance, all muscles with low compliance (i.e. T/M ratio as low as

0.7) are also affected when passive muscle lengths are in the middle of the ascending limb (Lp:

0.7-0.85). At 25% activation the results are similar but with decreased magnitude. Activation-

dependent shifts considerably influence the force capacity of all muscles on the ascending limb

of the force-length curve. For all of these effects, there is a trough of influence near or slightly

above L0 where activation does not alter force capacity appreciably.

Figure 5. The influence of compliance (at 100% (a) and 25% activation (b)) and AD shifts in L0 on force production [at 25% activation,(c)] as a function of MTU compliance and starting muscle length. Color designates the percent of results that saw a greater than 10% change in force production between models. Circle size represents the number of samples per bin. Y-axis is non-linear to ease visualization of results at low MTU compliance. Stars represent average compliance and operating length of the muscles acting at the hip (grey) and ankle (red) in CNS (b) and CS (c) conditions

3. How do these compliance and activation dependent factors influence the posture

which maximizes the force generating or moment capacity?

<u>Influence of compliance</u>: As expect, compliance has less of an effect on the force capacity for

muscles acting at the hip than for more distal muscles acting at the ankle. At the hip, the postures

that maximizes force and moment capacity change little with compliance (4^o and 6^o,

respectively). In contrast, at the ankle the postures that maximizes force and moment capacity

change by 26^o and 20^o, respectively ($^($Figs. 6a&b, Table 4). Likewise, we see very little difference

in force capacity at the hip between models that do and do not include compliance in a standing

posture (<-6%, blue vs. grey solid lines at 50^o Fig. 6a). When muscle moment arms are taken

into account we observe a reduction in moment capacity (-15%) compared to maximum values at

a more extended posture. At the ankle, muscles operating on the descending limb of the force-

length curve in a standing posture benefit from the change in length with compliance such that

Figure 6. Joint angle vs total force (a) and moment (b) acting across the hip (solid lines) and ankle (dashed lines) under three different muscle-tendon models. Average weighted moment arms for the hip (solid) and ankle (dashed) depicted in orange. While at the hip, the posture that produces maximum force and moment does not vary much between models (<9°), the maximum force and moment capacity decreases with both compliance and Ad-shift (~13%). In contrast, at the ankle the magnitude of maximum force and torque varies little between models (<3%), but the posture at which this maximum occurs varies by almost 40º.

they approach $L0_{100}$ and increase their force capacity by 27% and joint moment by +23% (CNS

model; blue vs grey dashed lines at 98°, Fig. 6a).

<u>Influence of activation-dependent shifts in L0</u>: The influence of activation-dependent shifts

shows a different pattern than that of compliance. Both the hip and ankle are equally influenced

by activation dependent shifts in L0.   At the hip, the postures that maximizes force capacity and

moment change by 16° and 8° when activation-dependent shifts in L0 are included in the model,

while the ankle the postures that maximizes force capacity and moment change by 8° and 10°

(Figs. 6a&b respectively, Table 4).

In a standing posture, muscles acting act the hip, since they operate on the ascending limb, are

shifted further away from optimal length when an activation-dependent shift in L0 is added to

the model and show a decrease in force and moment capacity (Δforce:-4%, Δmoment:-9%, Fig.

6A solid red line). The ankle, though, operates on the plateau of the force-length curve (~1.06)

and its force capacity increases slightly (Δforce:1.5%, Δmoment:+9%) when an activation-

dependent shift in L0 is added to the model (Fig. 6A, dashed red line).

In summary, at the hip, compliance decreases force capacity slightly more than AD shifts in L0

(-5% and -4%, respectively) in a standing posture, with both effects being small. At the ankle,

compliance increases the force generated at the ankle substantially (+26%), while activation-

dependent shifts have a much smaller effect (+2%). Similar effects are seen for moment

generating capacity.

Table 4. A comparison of functional influence of three different muscle-tendon models: No compliance No Shift
(NCNS), Compliance but No Shift (CNS) and Compliance and Shift (CS). For each model, the angle at maximum
force (Fmax°), average normalized fiber length (nFL), and total force (F) and moment (M) across the joint are listed
for both the hip (low compliance) and ankle joints (high compliance). Compliance and shifts in L0 *decrease the*
force and torque capacity at the hip while increasing them at the ankle.

	Hip T/M ratio: 0.71						**Ankle** T/M ratio 8.46						
	Max Values				Standing 50°			Max Values				Standing 98°		
	Force		Moment		nFL	F(N)	M (Nm)	Force		Moment		nFL	F(N)	M (Nm)
	Mag	°	Mag	°				Mag	°	Mag	°			
NCNS	65.8	50	0.67	24	0.97	65.8	.52	75.2	64	0.65	68	1.21	60.2	.53
CNS	62.1	56	0.62	20	0.95	62.0	.44	77.4	90	0.67	88	1.06	76.3	.65
CS	61.4	72	0.58	28	0.95	59.8	.48	78.2	100	0.66	96	1.05	78.1	.60

Discussion

The force-length relationship is one of the key components affecting a muscle's capacity to

produce force. And while there are multiple factors that are known to influence a muscle's

force-length operating range, their contribution and possible interactions have not been well

quantified. Here we tested the effect of, and interaction between, muscle tendon compliance,

activation and activation-dependent shifts of L0 on muscle operating length and force during

simulated fixed-end contractions in architecturally diverse muscles across a multitude of

conditions.

General effect of compliance and activation-dependent shifts of L0 on muscle length and force capacity

One of the overarching goals for this study was to identify general principles (if any) that describe how compliance and activation-dependent shifts of L0 influence muscle operating lengths. While the relationships were quite complex (see following section), we reveal several notable patterns. First, we found that in this species, on average, most muscles operate on the plateau or ascending limb of the force-length curve irrespective of the level of tendon compliance or activation-dependent shifts of L0. These averages incorporate data across the total possible *in vivo* length range of the muscles, including both the full range of passive lengths and activation levels. Because our analysis spans muscle force levels from near zero to maximum values, we do not expect that force-length ranges would deviate from this in dynamic contractions. The exception is possibly eccentric contractions under maximal activations where force can exceed F_{max} (Herzog et al., 2006). This comprehensive simulation of muscle operating lengths corroborates much of the literature showing that muscles typically do not operate on the descending limb of the force-length curve (Burkholder et al., 2001; Holt and Azizi, 2016; Rubenson et al., 2012). This provides further support for the idea that an animal's neuro-musculoskeletal structure is most often organized to operate on the ascending limb or plateau regions where muscles are inherently more stable and less susceptible to stretch-induced injury (Herzog et al., 1992, 1991; Lindstedt et al., 2001; Lutz and Rome, 1996; Morgan et al., 2002)

Our results also show that the most common effect of compliance is a leftward shift on the force-length curve leading to shorter muscle lengths and a decrease in force capacity. Since muscles, on average, operate on the ascending limb, increasing activation shortens muscles further below

L0 as the tendon stretches with greater force generation. The exception is for muscles that start at

passive lengths longer than $L0_{100}$. Under this condition, the result is an increase in force capacity

as the compliant muscle's fibers shorten towards L0, though some muscle lengths can shorten

beyond L0, first increasing and then decreasing force capacity. It is also notable that the average

effect of activation-dependent shifts of L0 generally mirror those described above for

compliance, although with much smaller effects. Thus, on average, the effects of compliance

dominates activation-dependent changes in operating lengths.

Finally, by comparing results that average across muscles acting at the hip (low compliance)

with those acting at the ankle (high compliance), we show that more compliant tendons will

generally amplify the effects described above. Given the proximal-distal gradient of compliance,

this analysis suggests that proximal muscles will show little activation-compliance effect

whereas distal muscles will be more significantly influenced. In summary, if specific knowledge

of a muscle's fiber, tendon and joint characteristics are lacking, the average data provide a

cursory prediction of the most likely influence of compliance and activation-dependent shifts of

L0 on muscle force-length operating ranges. As muscles increase activation, they will shorten

further away from L0 and generate less force than would be predicted by a linear force-activation

relationship. This effect increases with compliance and muscle passive length (up to L0).

Activation dependent shifts in L0 amplify this effect, but only minimally. However, these

generalizations should be used with caution because, as we will detail in the following sections,

they are error prone and can obscure the highly variable effect of compliance and/or activation-

dependent shifts of L0 on functionally-relevant changes in length and force capacity.

Under what conditions are the compliance and activation-dependent shifts in L0 meaningful?

While the analysis of average behavior paints results with a wide brush, it fails to capture the complex interactions between the muscle's starting passive length, compliance and activation-dependent shifts in L0. While there are trends in these interactions, it is important to emphasize that they are not broad. These relationships are very sensitive to a muscle's passive length and have large continuous gradients that shift with MTU compliance. Thus, small changes in the initial passive length can result in a large change in activation-compliance-dependent force capacity that generalizations fail to capture. For instance, while averages suggest that activation-dependent shifts in L0 have little influence on force capacity, this more detailed analysis shows that they can decrease or increase force capacity by as much as 15% of F_{100max}, resulting in a relative change of over 60% in force capacity at low activations. Likewise the average analysis underrepresented the extent of compliance on muscle length, which can be drastic, resulting in changes of up to 40% of L0. This is commensurate with the largest muscle strains seen in dynamic, high power movements (Askew and Marsh, 2002). Here we show this in a fixed-end 'isometric' contraction. Further, while the 'average' analysis concluded only a small effect of activation on muscle length in non-compliant muscles, this more detailed analysis highlighted that there is no abrupt threshold of MTU compliance below which the effects disappear. Rather there is a non-linear continuous gradient with compliance that varies across muscle operating lengths. There are regions of the force-length curve in which muscles are most sensitive to compliance and there are troughs of influence where these factors have little effect on force capacity. These regions are hard to predict, though, because the magnitude and location of these areas of high and low sensitivity shift with compliance level. Thus, rather than being able to provide an accurate rule of thumb for how and when compliance will alter a muscle's force

generating capacity, we identify conditions in which the effect is most drastic, the variables that

are most influential, and provide a guide for the possible errors in predicting muscle function if

muscle-tendon properties are not known.

While, on average, changes in muscle length were greatest for muscles starting at passive lengths

closer to optimal fiber length, L0, the influence of compliance on force capacity does not follow

suit. The length at which we see the greatest change in force shifts with MTU compliance. On

the ascending limb, stiff MTU's show the greatest drop in force at the shortest passive lengths

(dark blue regions of Fig. 4d), while compliant muscles see the greatest change when muscles

start at passive lengths just short of optimal fiber lengths (yellow regions of Fig. 4d). While it is

fair to say that compliance will always decrease the force generating capacity of muscles with

passive lengths on the ascending limb, the non-linear interactions between activation and

compliance make any approximations of the magnitude of the effect potentially inaccurate.

Depending on passive muscle length, the influence of compliance could decrease a muscle's

force capacity on the ascending limb anywhere from zero to 80%. While these numbers are

striking, this isn't even the region of the force-length curve that is most sensitive to the effects of

compliance. The region of most variability extends from just short of the plateau region of the

force-length curve and down a portion of the descending limb. For muscles within this region, at

100% activation compliance can result in either a decrease of or increase in force capacity of

well over 50% of F_{100max}, depending on the passive length. It is important to emphasize that

these conditions in which we see the greatest influence of activation on operating length are not

unusual. Within this region of highest variability are conditions we typically regard as 'optimal',

on or near the plateau. Thus, surprisingly, across the region of the force-length curve we

typically view force as relatively insensitive to length changes (i.e. the plateau), we find the area

of greatest sensitivity to activation-compliance effects. In summary, compliance results in

activation dependent changes in the operating length and force capacity of muscles, the

maximum values of which increase with increasing compliance. On the ascending limb, force

capacity decreases with increasing compliance and passive muscle length. On the plateau or

descending limb, the story is much more complex and can force capacity can either increase or

decrease depending on compliance and initial passive muscle length. Thus, at a muscle function

level we show that even in "isometric" conditions, the muscle-tendon unit is really 'dynamic'

with changes in muscle length, force capacity and activation requirements. One cannot

accurately assume force increases proportionally with activation; the activation-force

relationship is far from linear. Instead, it is likely a complex interaction of activation, muscle

passive length and MTU compliance. This interaction, although not well understood, suggests

great computational complexity for neural control of muscle force production.

The influence of activation-dependent shifts in L0 on force capacity generally follows a similar

pattern as described for compliance, increasing influence with increasing compliance. Again,

this can be explained by the dynamic nature of these interactions. The changes in force capacity

that come with a shift in L0 result in larger length changes in more compliant MTU's, which

result in further force changes. While these effects can be small in magnitude (less than 5% of

maximum force capacity for T/M ratio < 2), at low activation, this can represent a relatively large

change in force capacity (over 20%). In fact, at low activation, the influence of activation-

dependent shifts in L0 is as important, if not more important, than compliance effects in

determining a muscle's force capacity. Further, there is only a narrow region of muscle lengths

that result in 'no influence'; thus, the contribution of activation-dependent shifts in L0 are more

ubiquitous than our average analysis suggested, substantially altering the force capacity across a

wide range of MTU compliance and operating lengths at low activations. In summary, this

analysis allows us to provide information on what the most important variables are for accurately

predicting muscle force. Compliance is more important than activation-dependent shifts of L0 at

high activation, while both significantly alter the force capacity at low activation levels.

This has implications for musculoskeletal modeling and can provide a guide for what the

possible errors are in predicting muscle function if muscle-tendon properties are not known.

Models that assume non-compliant tendons could be over or underestimating muscle force

drastically. Specifically, this simplification would most often result in over estimation of force

generating capacity with muscles operating on the ascending limb and under estimation on the

descending limb. This is true, even within the less compliance range of human MTU's [lower

limb av. 3.06 (Arnold and Delp, 2011)]. At a T/M ratio of 3, for instance, our analysis shows

that for all but a narrow band of passive lengths, muscles show significant force deficits with

compliance. More striking is that even muscles with relatively short tendons (tendons shorter

than the muscle optimal fiber length, T/M rations lower than one) have regions of passive lengths

on the ascending limb where compliance result in significant changes in force capacity. This

implies that even low levels of compliance can significantly alter force capacity in some

conditions and that models that ignore or do not accurately define compliance should be

interpreted cautiously. More optimistically, muscle models that include compliance but ignore

activation-dependent shifts should capture the most dominant factors when simulating muscles at

high activations. But in cases where muscles are at submaximal activation (i.e. many activities),

ignoring activation-dependent shifts could lead to errors that mirror those of non-compliant

models; in general forces on the ascending limb are overestimated and those on the descending

limb underestimated.

What are the functional consequences of compliance / activation-dependent shifts of
L0?

While the major focus of this paper is to understand in detail the influence of compliance and

activation-dependent shifts of L0 on muscle operating lengths and force capacity, we also make a

first pass at assessing their possible functional influence. We do this in a task where our 'static'

analysis is most applicable, namely in a standing posture.

First, consistent with results from our other analyses, we find that force in the hip muscles, which

shorten down the ascending limb, decrease when both compliance and activation-dependent

shifts in L0 are included in the muscle models, whereas forces in the ankle muscles, which

shorten up the descending limb, increase with the addition of activation dependent effects. This

is also consistent with our finding that activation dependent effects generally have a larger

influence on more compliant MTU's. Models that include both activation dependent effects have

a greater influence of 'optimal force-generating posture' at the ankle than at the hip.

Surprisingly, though, the influence of activation-dependent shifts in L0 are greater for muscles

acting at the hip than those acting at the ankle, despite our results which show that activation-

dependent shifts generally have larger effects for more compliant muscles. This emphasizes two

take home points from this study. 1) Muscles with very little tendon are not immune to non-

linear activation dependent effects, and 2) as stated previously, the magnitude of activation-

dependent effects are very sensitive to the lengths at which the muscles operate. Here, the

muscles acting across the ankle fall into a trough of influence in the plateau region where

activation-dependent shifts in L0 have little impact (red star, Fig. 5c), while the muscles acting at

the hip operate on the ascending limb in a region of relatively high influence (grey star Fig. 5c).

Thus, again, we emphasize how highly sensitive activation-dependent effects are to muscle

passive lengths and how broad generalizations can lead to erroneous conclusions.

In a previous study, we also asked whether animals utilize postures that maximize torque

capacity during walking and/or running.   These analyses concluded that they did not, but were

based on simpler muscle models that did not incorporate activation dependent effects

[compliance or activation-dependent shifts of L0 (Hutchinson et al., 2015)]. This previous study

also did not separate the roles of length-dependent muscle force capacity and muscle moment

arms in dictating the effect of joint posture on moment capacity. Several experiments have

shown that peak isometric muscle force and moment arms do not necessarily coincide with the

joint angle that generates peak muscle or joint moments (Hoy et al., 1990; Lieber and Boakes,

1988a). Whether muscle force or moment arms have the largest contribution to peak moment is

variable. For example, for the same muscle (bi-articular frog semitendinosus), force capacity

dictates the muscle's peak knee joint moment (Lieber and Boakes, 1988b), but the muscle's

moment arm dictates the moment-angle profile at the hip (Lieber and Shoemaker, 1992). How

length-dependent force capacity and muscle moment arms influence observed postures remains

poorly understood.

If we assume that models that incorporate both compliance and activation-dependent shifts in L0

accurately capture muscle dynamics in this species and a conservative 25% muscle activation,

we do find some evidence for muscle-tendon mechanics and posture operating in concert.

Notably, we find that at the ankle, animals stand with a posture that maximizes both force and

joint moment, accommodating the strain in the muscle that occurs due to the high compliance of

its MTUs. These could offer support to the long-held theory that muscles operate at a length that

minimize force losses arising from their force-length relationship (Azizi, 2014; Azizi and

Roberts, 2010; Lieber and Ward, 2011; Rubenson et al., 2012; Smith et al., 2007).

However, the story is complex. At the hip, we see a difference of ~45° between postures that

maximize torque and force, with the animal choosing a posture halfway between. Standing

posture is ~22° more extended than optimal for force production and ~22° more flexed than the

posture that maximizes joint torque. This symmetry obscures the functional significance of these

differences, though. Since force capacity is less sensitive to changes in joint angle, a postural

difference of 22° results in a loss of only 20% in force capacity but decreases joint moment by

nearly 40%. It is interesting to note that adopting a posture that would increase moment capacity

at the hip (more erect) would also decrease the moment necessary to oppose gravity. Thus, a less

flexed hip angle could be more economically supported, yet is not adopted. While 25%

activation may overestimate muscle activations during standing (Rubenson and Marsh, 2009), a

lower activation level will only amplify these effects. Thus, this study, along with our previous

work and those of others, hint that there are likely many factors that influence preferred posture

and torque capacity is not universally prioritized. While our results are among the first to

systematically link length-dependent force capacity with posture, whether this specific

mechanical constraint strongly dictates posture is still far from clear.

Limitations and Future directions

While we have done these analyses assuming a particular level of activation dependent shift of the force-length curve, several studies have suggested that the level of shift my vary by species and may range from having no influence (de Brito Fontana and Herzog, 2016), to as large as 60% shift (Holt and Azizi, 2014). Our analysis of a 15% shift level was not intended to produce a definitive quantification of its influence, but as a first pass at evaluating the possible contribution and interaction effects. These analyses could be repeated for different levels in the AD shift in L0 and in dynamic conditions that include force-velocity effects to quantify how this influences the general conclusions. Likewise, care should be taken in extrapolating our results for species with different tendon properties for a similar muscle/tendon ratio could result in very different force and length changes. Regardless of these limitations, this analysis highlights the advantages of exploring these questions via musculoskeletal modeling, an approach which allows the generation of thousands of data points to elucidate patterns and trends that are not visible from studies of one or two muscles alone.

Author Contributions

S.M.C. and J.R. contributed to the conception and design of the study. S.M.C., K.E., H.S, S.L.D. and J.R. were the primary authors involved in the development of the musculoskeletal model and S.M.C developed the muscle simulations. M.C., R.L.M. and J.R. were the primary authors involved in the collection and analysis of experimental data. All authors contributed critically to data interpretation. S.M.C and J.R. drafted the initial manuscript and S.M.C., K.E., H.S, R.L.M, S.L.D. and J.R. contributed to editing the manuscript. S.M.C and J.R. contributed to figure preparation.

Funding

Research reported in this publication was supported in part through a Company of Biologists

Travelling Fellowship to J.R. and in part through The National Institute of Arthritis and

Musculoskeletal and Skin Diseases of the National Institutes of Health under award number

R21AR071588. . Sarcomere measurement work was supported by a Stanford Interdisciplinary

Graduate Fellowship to M.C.L. Work at Northeastern University was supported by a National

Institutes of Health (RO1 AR47337) and a National Science foundation grant (IOB-0542795) to

R.L.M. The content is solely the responsibility of the authors and does not necessarily represent

the official views of the National Institutes of Health or National Science Foundation.

Acknowledgements

706 The authors would like to thank Andrew Wong for his assistance with muscle dissection and

707 muscle-tendon path digitizing and spatial transformations. This study benefited from helpful

708 discussions with the Stanford Neuromuscular Biomechanics Lab fellows and students, in

709 particular Frank 'Clay' Anderson, Thor Besier, and Ajay Seth. Robert Siston helped develop the

711 optical tracking software and interface used to digitize muscle-tendon paths.

Data Availability

714 Upon acceptance, data will be made available on Data Dryad and model will be made public on

715 SimTK.

References

Arnold, EM, Delp, SL. 2011. Fibre operating lengths of human lower limb muscles during

walking. Philosophical Transactions of the Royal Society B: Biological Sciences,

366(1570): 1530–1539.

Arnold, EM, Hamner, SR, Seth, A, Millard, M, Delp, SL. 2013. How muscle fiber lengths and

velocities affect muscle force generation as humans walk and run at different speeds.

Journal of Experimental Biology, 216(11): 2150–2160.

Askew, G, Marsh, R. 2002. Muscle designed for maximum short-term power output: quail flight

muscle. Journal of Experimental Biology, 205: 2153–2160.

Azizi, E. 2014. Locomotor function shapes the passive mechanical properties and operating

lengths of muscle. Proceedings. Biological Sciences / The Royal Society, 281(1783):

20132914.

Azizi, E, Roberts, TJ. 2010. Muscle performance during frog jumping: influence of elasticity on

muscle operating lengths. Proceedings of the Royal Society B: Biological Sciences,

277(1687): 1523–1530.

Bates, D, Maechler, M, Bolker, B, Walker, S. 2015. Fitting Linear Mixed-Effects Models Using

{lme4}. Journal of Statistical Software, 67(1): 1–48.

Blix, M. 1894. Die Lange und die Spannung des Muskels. Skandinavisches Archiv Fur

Physiologie, 5(1): 173–206.

Buchanan, TS, Lloyd, DG, Manal, K, Besier, TF. 2004. Neuromusculoskeletal modeling:

Estimation of muscle forces and joint moments and movements from measurements of

neural command. Journal of Applied Biomechanics, 20(4): 367–395.

Burkholder, TJ, Lieber, RL, Burkholder, TJ. 2001. Sarcomere length operating range of

vertebrate muscles during movement. Journal of Experimental Biology, 204(Pt 9): 1529–

1536.

Carr, JA, Ellerby, DJ, Marsh, RL. 2011. Differential segmental strain during active lengthening

in a large biarticular thigh muscle during running. Journal of Experimental Biology, 214(Pt

20): 3386–95.

Cromie, MJ, Sanchez, GN, Schnitzer, MJ, Delp, SL. 2013. Sarcomere lengths in human extensor

carpi radialis brevis measured by microendoscopy. Muscle and Nerve, 48(2): 286–292.

Daley, MA, Biewener, AA. 2011. Leg muscles that mediate stability : mechanics and control of

two distal extensor muscles during obstacle negotiation in the guinea fowl. Philosophical

Transactions of the Royal Society B: Biological Sciences, 366: 1580–1591.

Daley, MA, Usherwood, JR, Felix, G, Biewener, AA. 2006. Running over rough terrain : guinea

fowl maintain dynamic stability despite a large unexpected change in substrate height.

Journal of Experimental Biology, 209: 171–187.

de Brito Fontana, H, Herzog, W. 2016. Vastus lateralis maximum force-generating potential

occurs at optimal fascicle length regardless of activation level. European Journal of Applied

Physiology, 116(6): 1267–1277.

Delp, SL, Loan, JP. 1995. A graphics-based software system to develop and analyze models of

musculoskeletal structures. Computational Biological and Medicine, 25(1): 21–34.

Ellerby, DJ, Henry, HT, Carr, JA, Buchanan, CI, Marsh, RL. 2005. Blood flow in guinea fowl

Numida meleagris as an indicator of energy expenditure by individual muscles during

walking and running. Journal of Physiology, 2: 631–648.

Ellerby, DJ, Marsh, RL. 2006. The energetic costs of trunk and distal-limb loading during

walking and running in guinea fowl Numida meleagris II . Muscle energy use as indicated

by blood flow. Journal of Experimental Biology, 209: 2064–2075.

Ellerby, DJ, Marsh, RL. 2010. The mechanical function of linked muscles in the guinea fowl

hind limb. Journal of Experimental Biology, 213(13): 2201–2208.

Fukunaga, T, Ichinose, Y, Ito, M, Kawakami, Y, Fukashiro, S. 1997. Determination of fascile

length and pennation in a contracting human muscle in vivo. Journal of Applied Physiology,

82(1): 354–8.

Gordon, AM, Huxley, AF, Julian, FJ. 1966. The variation in isometric tension with sarcomere

length in vertebrate muscle fibres. Journal of Physiology, 184(1): 170–192.

Gordon, JC, Rankin, JW, Daley, MA. 2015. How do treadmill speed and terrain visibility

influence neuromuscular control of guinea fowl locomotion? Journal of Experimental

Biology, 218: 3010–3022.

Hawkins, D, Bey, M. 1997. Muscle and tendon force-length properties and their interactions in

vitro. Journal of Biomechanics, 30(1): 63–70.

Henry, HT, Ellerby, DJ, Marsh, RL. 2005. Performance of guinea fowl Numida meleagris during

jumping requires storage and release of elastic energy. Journal of Experimental Biology,

208: 3293–3302.

Herzog, W, Lee, EJ, Rassier, DE, Schappacher, G, DuVall, M, Leonard, TR, Herzog, JA. 2006.

Residual Force Enhancement Following Eccentric Contractions: A New Mechanism

Involving Titin. Physiology, 31(3): 300–312.

Herzog, W, Leonard, TR, Renaud, JM, Wallace, J, Chaki, G, Bornemisza, S. 1992. Force-length

properties and functional demands of cat gastrocnemius, soleus and plantaris muscles.

Journal of Biomechanics, 25(11): 1329–1335.

Herzog, W, Read, LJ, ter Keurs, HEDJ. 1991. Experimental determination of force-length

relations of intact human gastrocnemius muscles. Clinical Biomechanics, 6(4): 230–238.

Holt, NC, Azizi, E. 2014. What drives activation-dependent shifts in the force-length curve?

Biology Letters, 10(9): 20140651-.

Holt, NC, Azizi, E. 2016. The effect of activation level on muscle function during locomotion:

are optimal lengths and velocities always used? Proceedings of the Royal Society B:

Biological Sciences, 283(1823): 20152832.

Hoy, M, Zajac, E, Gordon, E. 1990. A musculoskeletal model of the human lower extremity: The

effect of muscle, tendon, and moment arm on the moment-angle relationship of

musculotendon actuators at the hip, knee, and ankle. Journal of Biomechanics, 23(2): 157–

169.

Hutchinson, JR, Rankin, JW, Rubenson, J, Rosenbluth, KH, Siston, RA, Delp, SL. 2015.

Musculoskeletal modelling of an ostrich (Struthio camelus) pelvic limb: influence of limb

orientation on muscular capacity during locomotion. PeerJ, 3: e1001.

Huxley, AF, Simmons, RM. 1971. Proposed mechanism of force generation in striated muscle.

Nature, 233: 533–8.

Lakatta, EG, Jewell, BR. 1977. Length-dependent activation: its effect on the length-tension

relation in cat ventricular muscle. Circulation Research, 40(3): 251–7.

Lemos, RR, Epstein, M, Herzog, W. 2008. Modeling of skeletal muscle: The influence of tendon

and aponeuroses compliance on the force-length relationship. Medical and Biological

Engineering and Computing, 46(1): 23–32.

Lieber, R, Brown, CG. 1992. Sarcomere length Joint angle relationships of seven frogs hindlimb

muscles. Acta Anatomica, 145: 289–295.

Lieber, RL, Boakes, JL. 1988a. Muscle force and moment arm contributions to torque production

in frog hindlimb. American Journal of Physiology, 254(6 Pt 1): C769-72.

Lieber, RL, Boakes, JL. 1988b. Sarcomere length and joint kinematics during torque production

in frog hindlimb. American Journal of Physiology, 254: C759-68.

Lieber, RL, Shoemaker, SD. 1992. Muscle, joint, and tendon contributions to the torque profile

of frog hip joint. American Journal of Medicine, 263(3 Pt 2): R586–R590.

Lieber, RL, Ward, SR. 2011. Skeletal muscle design to meet functional demands. Philosophical

Transactions of the Royal Society B: Biological Sciences, 366(1570): 1466–1476.

Lindstedt, SL, LaStayo, PC, Reich, TE. 2001. When Active Muscles Lengthen: Properties and

Consequences of Eccentric Contractions. Physiology, 16(6): 256–261.

Llewellyn, ME, Barretto, RPJ, Delp, SL, Schnitzer, MJ, Program, B, Clark, JH, Engineering, B.

2010. Minimally invasive high-speed imaging of sarcomere contractile dynamics in mice

and humans. Nature, 454(7205): 784–788.

Lloyd, DG, Besier, TF. 2003. An EMG-driven musculoskeletal model to estimate muscle forces

and knee joint moments in vivo. Journal of Biomechanics, 36: 765–776.

Lutz, GJ, Rome, LC. 1996. Muscle function during jumping in frogs. I. Sarcomere length

change, EMG pattern, and jumping performance. The American Journal of Physiology, 271:

C563-70.

MacIntosh, BR, MacNaughton, MB. 2005. The length dependence of muscle active force:

considerations for parallel elastic properties. Journal of Applied Physiology (Bethesda,

Md. : 1985), 98(5): 1666–73.

Marsh, RL, Ellerby, DJ, Henry, HT, Rubenson, J. 2006. The energetic costs of trunk and distal-

limb loading during walking and running in guinea fowl Numida meleagris I . Organismal

metabolism and biomechanics. Journal of Experimental Biology, 209: 2050–2063.

Mayfield, DL, Launikonis, BS, Cresswell, AG, Lichtwark, GA. 2016. Additional in-series

compliance reduces muscle force summation and alters the time course of force relaxation

during fixed-end contractions. Journal of Experimental Biology, 219(September): jeb.143123.

Morgan, DL, Brockett, CL, Gregory, JE, Proske, U. 2002. The role of the length-tension curve in the control of movement. Sensorimotor Control of Movement and Posture, 508: 489–494.

Rack, P, Westbury, DR. 1969. The effects of length and stimulus rate on tension in the isometric cat soleus muscle. Journal of Physiology, 204: 443–460.

Rack, PM, Westbury, DR, Rack, PM. 1969. The effects of length and stimulus rate on tension in the isometric cat soleus muscle. Journal of Physiology, 204(2): 443–460.

Ramsey, R. W. & Street, SF. 1941. The isometric length-tension diagram of isolated skeletal muscle fibers of the frog. Protoplasma, 36(1): 157–190.

Rassier, DE, MacIntosh, BR, Herzog, W. 1999. Length dependence of active force production in skeletal muscle. Journal of Applied Physiology, 86(5): 1445–1457.

Roszek, B, Baan, GC, Huijing, PA. 1994. Decreasing stimulation frequency-dependent length-force characteristics of rat muscle. Journal of Applied Physiology, 77(4): 2115–2124.

Rubenson, J, Marsh, RL. 2009. Mechanical efficiency of limb swing during walking and running in guinea fowl (Numida meleagris). Journal of Applied Physiology, 106(1985): 1618–1630.

Rubenson, J, Pires, NJ, Loi, HO, Pinniger, GJ, Shannon, DG. 2012. On the ascent: the soleus operating length is conserved to the ascending limb of the force-length curve across gait mechanics in humans. Journal of Experimental Biology, 215(20): 3539–3551.

Rubenson, J, Sanghvi, H, Cromie, MJ, Easton, K, Marsh, RL, Delp, SL. 2013. Influence of tendon compliance and activation level on fibre operating lengths of skeletal muscle. *Annual Meeting of the Society for Integrative and Comparative Biology*, San Fancisco,

856   CA.

857   Seth, A, Hicks, JL, Uchida, TK, Habib, A, Dembia, L, et al. 2018. OpenSim : Simulating

858   musculoskeletal dynamics and neuromuscular control to study human and animal

859   movement. Computational Biology, 14(7): e1006223.

860   Smith, NC, Payne, RC, Jespers, KJ, Wilson, AM. 2007. Muscle moment arms of pelvic limb

861   muscles of the ostrich (Struthio camelus). Journal of Anatomy, 211(3): 313–324.

862   Sugisaki, N, Kawakami, Y, Kanehisa, H, Fukunaga, T. 2011. Effect of muscle contraction levels

863   on the force-length relationship of the human Achilles tendon during lengthening of the

864   triceps surae muscle-tendon unit. Journal of Biomechanics, 44(11): 2168–2171.

865   TerKeurs, H, Isazumi, K, Pollack, GH. 1978. The Sarcomere Length-Tension Relation in

866   Skeletal Muscle. Journal of General Physiology, 72(October): 565–592.

867   Zajac, FE. 1989. Muscle and tendon: properties, models, scaling, and application to

868   biomechanics and motor control. Critical Reviews in Biomedical Engineering, 17(4): 359–

869   411.