1

Vps34 PI 3-kinase controls thyroid hormone production by regulating thyroglobulin

iodination, lysosomal proteolysis and tissue homeostasis

Giuseppina Grieco[1*], Tongsong Wang[1*], Ophélie Delcorte[1*], Catherine Spourquet[1*], Virginie

Janssens[1], Aurélie Strickaert[2], Héloïse P. Gaide Chevronnay[1], Xiao-Hui Liao[3], Benoît Bilanges[4],

Samuel Refetoff[3], Bart Vanhaesebroeck[4], Carine Maenhaut[2], Pierre J. Courtoy[1*], and

Christophe E. Pierreux[1*#]

[1], Cell Biology Unit, de Duve Institute and Université Catholique de Louvain, 1200 Brussels,

Belgium;

[2], Institute of Interdisciplinary Research (IRIBHM); Université Libre de Bruxelles, 1070 Brussels,

Belgium;

[3], Departments of Medicine (X-H.L., S.R.) and Pediatrics and Genetics (S.R), The University of

Chicago, Chicago, Illinois 60637;

[4], UCL Cancer Institute; University College London, 72 Huntley Street, London WC1E 6DD, UK

[#], correspondence to christophe.pierreux@uclouvain.be

*, equal contributions

Running title: Vps34 controls TH synthesis

Key words: Vps34, thyroid, hormone, organification, autophagy

Dr. Giuseppina Grieco, g.pia@hotmail.it

Dr. Tongsong Wang, tswang0802@hotmail.com

Ms. Ophélie Delcorte, ophelie.delcorte@uclouvain.be

Ms. Catherine Spourquet, catherine.spourquet@uclouvain.be

Ms. Virginie Janssens, virginie.janssens@uclouvain.be

Dr. Aurélie Strickaert, aurelie.strickaert@ulb.ac.be

Dr. Héloïse Gaide-Chevronnay, heloisegaide@hotmail.com

Dr. Xia Hui Liao, xliao1@uchicago.edu

Dr. Benoit Bilanges, b.bilanges@ucl.ac.uk

Prof. Samuel Refetoff, srefetof@uchicago.edu

Prof. Bart Vanhaesebroeck, bart.vanh@ucl.ac.uk

Prof. Carine Maenhaut, cmaenhau@ulb.ac.be

Prof. Pierre J. Courtoy, pierre.courtoy@uclouvain.be

Prof. Christophe E. Pierreux, christophe.pierreux@uclouvain.be

Abstract (327 words, 350 allowed)

BACKGROUND: The production of thyroid hormones (T_3, T_4) depends on the organization of

the thyroid in follicles, which are lined by a monolayer of thyrocytes with strict apico-basal

polarization. This polarization supports vectorial transport of thyroglobulin for storage into, and

recapture from, the colloid. It also allows selective addressing of channels, transporters, ion

pumps and enzymes to their appropriate basolateral (NIS, SLC26A7 and Na^+/K^+-ATPase) or

apical membrane domain (Anoctamin, SLC26A4, DUOX2, DUOXA2 and TPO). How these actors

of T_3/T_4 synthesis reach their final destination remains poorly understood. The PI 3-kinase

(PI3K) isoform Vps34/PIK3C3 is now recognized as a main component in the general control

of vesicular trafficking and of cell homeostasis via the regulation of endosomal trafficking and

autophagy. We recently reported that conditional Vps34 inactivation in proximal tubular cells

in the kidney prevents normal addressing of apical membrane proteins and causes abortive

macroautophagy.

METHODS: Vps34 was inactivated using a Pax8-driven Cre recombinase system. The impact

of Vps34 inactivation in thyrocytes was analyzed by histological, immunolocalization and

mRNA expression profiling. Thyroid hormone synthesis was assayed by ^{125}I injection and

serum plasma analysis.

RESULTS: Vps34^{cKO} mice were born at the expected Mendelian ratio and showed normal

growth until postnatal day 14, then stopped growing and died at around 1 month of age. We

therefore analyzed thyroid Vps34^{cKO} at postnatal day 14. We found that loss of Vps34 in

thyrocytes causes: (i) disorganization of thyroid parenchyma, with abnormal thyrocyte and

follicular shape and reduced PAS$^+$ colloidal spaces; (ii) severe non-compensated

hypothyroidism with extremely low T_4 levels (0.75 ± 0.62 $\mu g/dL$) and huge TSH plasma levels

(19,300 ± 10,500 mU/L); (iii) impaired ^{125}I organification at comparable uptake and frequent

occurrence of follicles with luminal thyroglobulin but non-detectable T4-bearing

thyroglobulin; (iv) intense signal in thyrocytes for the lysosomal membrane marker, LAMP-1,

as well as thyroglobulin and the autophagy marker, p62, indicating defective lysosomal

proteolysis, and (v) presence of macrophages in the colloidal space.

CONCLUSIONS: We conclude that Vps34 is crucial for thyroid hormonogenesis, at least by

controlling epithelial organization, Tg iodination as well as proteolytic T_3/T_4 excision in

lysosomes.

 Introduction

The main function of the thyroid gland is to produce the hormones, T_3/T_4 and

calcitonin, which are essential for the regulation of metabolic processes (1,2). The production

of T_3 and T_4 depends on the correct tissue organization of thyroid epithelial cells, the

thyrocytes, into functional and independent units, the follicles. These are composed of a single

layer of polarized thyrocytes that form a spherical structure delineating an internal space or

lumen where the thyrocyte secretory product, thyroglobulin, is stored in a colloidal form, thus

called the colloid lumen*.* As thyrocytes communicate via gap junctions, each follicle functions

as an integrated unit. Thyroid hormone synthesis depends on apico-basal cell polarity that

allows the specific localization of channels, transporters, pumps and enzymes at the

appropriate membrane domains. Iodide from the bloodstream freely traverses the

fenestrated endothelium of the thyroid capillaries and is taken up into thyrocytes via the

basolaterally-localized Na^+/I^- symporter (NIS), thanks to a Na^+ gradient generated by the

Na^+/K^+-ATPase. An alternative basolateral transporter, SLC26A7, has been recently reported

to also control iodide uptake, although its role might be indirect (3). Iodide diffuses freely

within thyrocytes and is next transported across the apical membrane into the colloid space

via apically-localized transporters such as anoctamin or pendrin (SLC26A4). Iodide is then

rapidly oxidized into iodine by thyroperoxidase (TPO) located at the apical membrane, in the

presence of H_2O_2, generated by the DUOX2/DUOXA2 apical membrane complex. Iodine is

incorporated into accessible tyrosine residues close to the N- and C-termini of thyroglobulin

(Tg), a large protein secreted by thyrocytes into the colloid space. Iodotyrosine rearranges into

hormonogenic peptides, which are the direct T_3 and T_4 precursors. Thyroid hormone synthesis

thus requires basal localization of NIS, SLC26A7 and Na^+/K^+-ATPase, apical localization of

anoctamin, pendrin, TPO, DUOX2 and DUOXA2, as well as apical delivery of Tg into the colloid

lumen and endocytic uptake of iodoTg into thyrocyte lysosomes (1,2). Specific regulators of

thyroid follicular organization have recently been identified (4). Transcriptomic comparison of

thyroid FRT cells cultured in 2D monolayers and in 3D spherical follicles indeed revealed

involvement of structural and functional cell elements such as adherens and tight junctions

(cadherin-16), cytoskeleton proteins, ions channels, proteins involved in differentiation, and

components of the trafficking machinery (e.g. myosin-Vb, Rab17) (4).

There is strong evidence that vesicular trafficking is critical for apico-basal polarization

and epithelial function (5-7), but the role of vesicular trafficking in thyroid function is

incompletely understood. Vps34/*PIK3C3* (also referred to as type III PI3K) has long been

recognized as a main actor involved in the general control of endocytic vesicular trafficking (8-

13). Vps34/PIK3C3 also plays an important role in epithelial organization in *Drosophila* (14)

and in autophagy (12,15), both in the initiation of autophagosome formation and the

progression towards autophagosome-lysosome fusion (16,17).

We recently inactivated Vps34 in kidney proximal tubular cells (PTC) using Wnt4-Cre

and Pax8-Cre (18,19). Wnt4-Cre driven Vps34 excision caused major perinatal lethality, with

abnormalities in kidney cortex development and PTC apical differentiation in the surviving

mice. With the Pax8-Cre, which triggers recombination in nephrogenic tubules and in

nonvascular components of glomeruli two days later than Wnt4-Cre (20,21), we observed

normal mouse growth until postnatal (P) day 14 (19). Then, body weight levelled off and all

pups died at ~3-5 weeks of age, probably due to kidney failure. Structural and functional

studies of Vps34[cKO] kidneys revealed preserved PTC membrane polarity, but impaired apical

membrane protein trafficking, thus causing a general PTC dysfunction known as renal Fanconi-

like syndrome, manifested by polyuria and low-molecular weight proteinuria. Vps34[cKO] also

displayed impaired lysosome size/positioning and blocked autophagy, thereby causing cell

vacuolization. We concluded that Vps34 is a crucial component of the trafficking machinery

necessary for differentiated PTC function and is essential for overall PTC homeostasis (19).

Given that Pax8-Cre also triggers recombination in thyrocytes (21), we here investigated the

role of Vps34/PIK3C3 in thyroid function and homeostasis.

Materials and methods

Mice

Vps34$^{fl/fl}$ mice have been described (22). Pax8-Cre mice were obtained from Dr. M.

Busslinger (21). Vps34$^{fl/fl}$ mice were crossed with Pax8-Cre;Vps34$^{fl/+}$ mice to generate

conditional targeted excision of Vps34 exon 21 in the thyroid of 25% of offspring (Vps34^{cKO}).

Non-recombined Vps34$^{fl/fl}$ mice were used as control. Mice were treated according to the NIH

Guide for Care and Use of Laboratory Animals, and experiments were approved by the

University Animal Welfare Committee, Université Catholique de Louvain (2016/UCL/MD/006

and 2018/UCL/MD/026).

Plasma, tissue collection and histology

Blood was collected by eye sinus puncture at sacrifice (P14) under irreversible

anesthesia by xylazine 2% and ketamine 50 mg/ml (200 l/mice i.p.). Thyroid lobes were

excised, fixed by immersion in neutral-buffered formaldehyde (4% F) at 4°C under stirring

overnight. Samples were paraffin-embedded or equilibrated overnight in 20% sucrose and

embedded in Tissue-Tek Optimal Cutting Medium (Sakura Finetek) for cryostat sections.

TSH and T4 plasma concentrations

140   Plasma TSH concentrations were measured by a sensitive, heterologous,

141 disequilibrium double-antibody precipitation RIA as described (23). T4 concentration was

142 measured by coated-tube RIA (Siemens Medical Solution Diagnostics, Los Angeles, CA).

143 *Immunofluorescence*

144   Immunofluorescence was performed on 5 μm-thick frozen sections or on 6 μm-thick

145 paraffin sections (19). Antigen retrieval was promoted in citrate buffer, pH 6.0, at 98°C for 20

146 min using a Lab Vision Pretreatment Module™ (Thermo Scientific). After permeabilization with

147 PBS/0.3% Triton-X100 for 5 min, non-specific sites were blocked by 1-h incubation in PBS/0.3%

148 Triton-X100 with 10% bovine serum albumin (BSA) and 3% milk, followed by primary

149 antibodies (described in Supplementary Table I) in blocking buffer at 4°C overnight. After

150 extensive washing, sections were incubated with the appropriate AlexaFluor-secondary

151 antibodies in 10% BSA/0.3% Triton-X100 at room temperature for 1 h, extensively washed,

152 mounted with Faramount Aqueous Mounting Medium (Dako) and imaged on a spinning disk

153 confocal microscope using a Plan Apochromat 100x/1.4 Oil DIC objective (Cell Observer

154 Spinning Disk; Zeiss). For whole thyroid section recording, images were acquired using Zeiss

155 Pannoramic P250 slide scanner, stitched and analyzed using Case Viewer software.

156 *RT-qPCR*

157   Total RNA was extracted from thyroid lobes using TRIzol Reagent (Thermo Scientific),

158 as described (24). Aliquots of 500 ng RNA were reverse-transcribed by M-MLV reverse

159 transcriptase (Invitrogen) with random hexamers, as described (25). Primer sequences used

160 are described in Supplementary Table II. Real-time qPCR was performed as described (25), in

161 presence of 250 nM of specific primers with Kappa SYBR Fast qPCR Master Mix (Kapa

162 Biosystems) on a CFX96 touch real-time PCR Detection System (Bio-Rad). Data were analyzed

using the ΔΔCT method, using the geometric mean of -Actin and *Rpl27* as reference genes

(26).

[125]I uptake and organification

At postnatal day 8 (P8), mothers and litters were fed with an iodine-free diet for 24 h.

At P9, pups were injected intraperitoneally with 1 µCi [125]I (Perkin Elmer) and left for another

24 h before sacrifice and thyroid dissection. Both lobes were collected in 2 mM methimazole

(Sigma) and homogenized with a glass Potter. Total thyroid [125]I was measured with an

automatic gamma counter "Wizard[2]" (Perkin Elmer) before protein precipitation using 10%

trichloroacetic acid (TCA; Merck). After a single wash in TCA, radioactivity was measured in

the protein pellet. Percentage of protein-bound iodide was calculated using the ratio of

precipitated cpm/total thyroid cpm (27).

H_2O_2 level measurements

The hydrogen peroxide Assay kit from Abcam (ab102500) was used. At P14, thyroid

lobes were collected in ice-cold PBS and homogenized with a glass Potter. Proteins were

precipitated with perchloric acid 4 M, and supernatant was neutralized with KCl 3 M and KCl

1 M until pH was comprised between 6.5 and 8. Fifty l, corresponding to a ninth of a thyroid,

was incubated with OxiRed probe and horseradish peroxidase (HRP), according to the

instructions form the manufacturer. Fluorescence was measured with a GloMax fluorimeter

(Promega, Ex/Em = 535/587 nm).

Statistical analysis

All statistical analyses were determined by Prism software (GraphPad Software, La

Jolla, California, USA). RT-qPCR values were obtained by the ΔΔCT method and are expressed

as boxplots with median, 25th and 75th percentiles, and min-max whiskers. Each graph

represents the results form a minimum of 8 independent thyroid lobes from at least 3

different litters. Nonparametric statistical tests were used: Mann-Whitney for single

comparisons. Differences were considered statistically significant when $p<0.05$ (*); ** stands

for $p<0.01$; *** for $p<0.001$.

Results

Genetic construction and assessment of Vps34 inactivation

Ubiquitous Vps34 inactivation is embryonically lethal, with embryos dying around

embryonic day E8.5 (28). To study tissue-specific roles of Vps34 *in vivo,* we used a mouse line

which carries a floxed *Vps34* allele (Vps34fl) that allows to conditionally delete the loxP-flanked

exon 21 of the *Vps34* gene, which encodes a critical sequence of the lipid kinase domain of

Vps34 (22). This approach was designed to allow for the expression of a minimally-truncated,

catalytically-inactive Vps34 protein. However, upon conditional expression in

megakaryocytes, the level of this truncated Vps34 and its obligatory partner, Vps15, were

found to be decreased by 80-90% in the megakaryocyte lineage (thrombocytes), indicating

instability of truncated Vps34 protein (22). We will thus hereafter refer to the result of this

Vps34 truncation as Vps34 conditional knock-out (Vps34cKO).

We crossed homozygous Vps34fl/fl mice with Pax8-Cre;Vps34fl/+ mice, which is expected

to lead to a tissue-specific (thyroid and kidney) conditional excision of exon 21 from both

Vps34 alleles in 25% of the pups, which are further referred to as Vps34cKO mice. Vps34cKO

mice were born at the expected Mendelian ratio and showed normal development until

postnatal day 14 (P14), then stopped growing and died at around 1 month of age (19). We

therefore analyzed the thyroid of Vps34^{cKO} mice at P14. Of note, eye opening, a marker of

cerebral maturation was usually delayed till P10.

We first assessed the extent of Vps34 genetic excision in the thyroid by quantification

of *Vps34* mRNA using primer that are specific for the total *Vps34* or are exon 21-specific.

Compared to control littermate pups, we found a ~70 % reduction of exon 21-containing

Vps34 mRNA in cKO thyroid extracts. In comparison, we found no significant difference in the

abundance of *Vps34* mRNA containing exon 22-24 (Fig. 1A). Assuming that *Vps34* is equally

expressed in all thyroid cell types (expressing and non-expressing the Cre recombinase), this

~70 % decrease suggests that most thyrocytes in Vps34^{cKO} mice had undergone Cre-mediated

recombination.

Vps34^{cKO} thyroids show signs of hyperstimulation by thyroid stimulating hormone (TSH)

At P14, thyroid glands from Vps34^{cKO} were almost twice heavier as compared to

control glands (Fig. 1B). Histological staining using Periodic-Acid-Schiff (PAS, which in the

thyroid reflects production of the thyroglobulin glycoprotein) readily revealed striking

differences between thyroid glands of control and Vps34^{cKO} mice (Fig. 1C). Thyroid glands of

control mice appeared as assemblies of round follicular sections of variable diameter, filled

with colloid intensely and homogenously stained by PAS. Although histology of Vps34^{cKO} was

more variable, the three representative individual thyroids shown in Fig. 1C revealed irregular

follicle shape, especially at the periphery of the gland, and weaker or absent PAS staining (Fig.

1C and Suppl. Fig. 1). In addition, the PAS-negative follicular space were often filled with nuclei

(see Vps34^{cKO} #3). Quantification revealed that 90% of control follicles had a regular shape

and intense (+++) PAS staining (Suppl. Fig. 1). On the contrary, only 50% of Vps34^{cKO} follicles

displayed a regular shape and most of the follicles were weakly positive for PAS (++ or +)

(Suppl. Fig. 1). In addition, 10 to 50% of the Vps34[cKO] follicles showed no PAS staining (-) and

luminal cells were found in 40-70% of the follicles (Suppl. Fig. 1). Confocal

immunofluorescence microscopy for the epithelial marker E-cadherin confirmed

heterogeneity of Vps34[cKO] follicles, but also of thyrocytes. Hoechst labelling revealed the

presence of several nuclei in a significant fraction of lumina (Fig. 1D). These histological

features of follicle remodelling and colloid consumption up to exhaustion, suggested

perturbed thyroid function.

Vps34[cKO] causes severe non-compensated hypothyroidism

To directly test the hypothesis that Vps34[cKO] could impair thyroid hormone production,

plasma was collected at P14 and analyzed for the levels of T_4 and TSH. In P14 control pups,

values were 6.4 ± 1.5 μg/dL for T_4, and 75 ± 110 mU/L for TSH (Fig. 2). In Vps34[cKO] mice, we

found extremely low T_4 values (0.75 ± 0.62 μg/dL) and very high plasma levels of TSH (19,311

$\pm$ 10,482 mU/L) (Fig. 2). These data indicate severe, non-compensated hypothyroidism. This

observation, compatible with the growth retardation observed after P15 (19), could be

explained by mislocalization of one or several basolateral and/or apical actors involved in

thyroid hormone synthesis, as we reported in Vps34[cKO] kidney proximal tubular cells (19).

Vps34 cKO display reduced iodine organification

The reduced or absent PAS staining (Fig. 1C) and the low T_4 plasma levels in Vps34[cKO]

mice (Fig. 2) might be caused by defective basolateral iodine uptake, apical transport and/or

apical organification on thyroglobulin. Due to lack or very poor specificity of antibodies

directed against murine proteins involved in these processes, we first measured their mRNA

expression levels in total thyroids at P14 (Fig. 3). We observed no change in mRNA expression

of *Nis*, *Ano1*, *Tg*, *Tpo* and *Duox2,* a significant increase in *Tshr* and *Slc26a4* expression, and a

significant decrease of *Slc26a7* and *Duoxa2* mRNA levels. At this stage, we cannot conclude if

these changes contribute to the observed hypothyroidism of Vps34^{cKO}.

To further test whether basolateral NIS and SLC26A7 and apical ANO1, SLC26A4, TPO,

DUOX and DUOXA were all correctly localized, we functionally assayed their combined activity

by injecting iodine-deprived pups at P9 with ^{125}I. At P10, 24-h post-injection, thyroid lobes

were collected and radioactivity measured before and after protein precipitation by TCA (Fig.

4A). ^{125}I uptake was not statistically-different in Vps34^{cKO} and the three control genotypes (Fig.

4A), suggesting normal NIS and SLC26A7 function and thus basolateral localization. In marked

contrast, only 10% of thyroid ^{125}I was bound to proteins in Vps34^{cKO} compared to

approximately 55% in control genotypes at this stage (Fig. 4A). This indicates a major effect of

Vps34 on one or several apical proteins involved in iodine organification.

Normal *Ano1* and increased *Slc26a4/Pendrin* mRNA levels suggested correct, or even

improved, transfer of iodine in the follicular lumen. On the other hand, decreased *Duoxa2*

might impact on H_2O_2 production. We thus measured the production of H_2O_2 in control and

Vps34^{cKO} thyroid lobes. Surprisingly, the median level of H_2O_2 production in Vps34^{cKO} was

higher, even if we observed variability (Fig. 4B), indicating that the DUOX/DUOXA pair is

functional in Vps34^{cKO} thyroid. Altogether, these results suggest defective localization of one

or several apical actors involved in iodine organification.

Apical polarity is impaired in Vps34^{cKO}

Apico-basal polarity is essential for thyrocyte function and defects in polarity might

impact on the delivery, and thus localization, of actors involved in thyroid hormone synthesis.

As general markers to assess polarity, we used the basement membrane protein laminin, the

basolateral Na$^+$/K$^+$-ATPase, E-cadherin and -catenin, the apically-localized ERM family

member ezrin and the tight junction-associated protein ZO-1. Control thyrocytes displayed a

well-defined apico-basal polarity with laminin assembled on the basal side, with Na^+/K^+-

ATPase, E-cadherin and -catenin restricted to the basolateral membrane and separated from

apical ezrin by the tight junction, visualized by ZO-1 (Fig. 5A and 5B). In Vps34^{cKO} thyrocytes,

basal laminin was correctly assembled and Na^+/K^+-ATPase, E-cadherin and -catenin were

restricted to the basolateral membrane, indicating the presence of functional tight junctions.

However, on the apical side of Vps34^{cKO} thyrocytes, the ezrin signal was weaker or absent from

the lumen-facing pole of some thyrocytes (Fig. 5A), and most cells also lacked ZO-1 protein

labelling (Fig. 5B). These data suggest impaired apical polarity in Vps34^{cKO}, which might

contribute to defective localization of apical actors involved in iodine organification.

Vps34^{cKO} mice display defective thyroglobulin iodination

To further analyse the significance of the lack of PAS staining in a fraction of Vps34^{cKO}

follicles (Fig. 1C) and the defect in iodine organification (Fig. 4), we assessed thyroglobulin

protein expression and its associated T_4 hormonogenic peptide (I-Tg) by immunofluorescence.

Low magnification of control sections showed an identical distribution pattern of

thyroglobulin and I-Tg, homogeneously filling all round colloidal spaces (Fig. 6A). In Vps34^{cKO},

the thyroglobulin labelling was more heterogeneous, mainly due to the presence of cells in

the colloid. Remarkably, antibodies recognizing the T_4 hormonogenic peptide often failed to

label colloidal spaces even when containing Tg in the adjacent section (Fig. 6A), thereby

confirming the organification defect (Fig. 4A). Quantification revealed that only a quarter (25.2

± 11.9 %) of the follicles were labeled for T_4 hormonogenic peptide. Of specific interest,

whereas the Tg signal was restricted to the follicle lumen in control thyroid, high magnification

revealed that the Tg signal was frequently seen within Vps34^{cKO} thyrocytes (Fig. 6B,

arrowheads). These observations indicated that Vps34 deletion caused defective

thyroglobulin iodination, as well as reduced Tg exocytosis and/or excessive Tg endocytosis.

Impaired lysosomal function and I-Tg proteolysis in Vps34cKO thyroid

Vps34 is involved in endocytic trafficking and cKO of Vps34 in postnatal kidney

glomeruli podocytes causes a strong increase of the overall immunofluorescence signal for

the lysosomal membrane marker LAMP-1, indicating enhanced lysosome biogenesis (29,30).

We further reported that absence of Vps34 in kidney PTCs causes an increase in the actual

size of lysosomes, that were sometimes enlarged, and mislocalized toward the basal pole of

the cell (19). Given that lysosomal proteases in thyrocytes are important to excise

hormonogenic peptides to release free T_3 and T_4, we investigated LAMP-1-labeled structures

in Vps34cKO thyrocytes as a proxy for lysosomal function, and its co-localization with Tg and I-

Tg to demonstrate endocytosis. In our conditions, immunofluorescence on control thyroid

sections only produced a weak LAMP-1 signal in the E-cadherin-positive epithelial thyrocytes

(Fig. 7A and 7B). On the contrary, a strong and widespread LAMP-1 signal was observed in all

Vps34cKO thyrocytes. As shown in the enlarged boxes of Fig. 7A and 7B (separate emission

channels in white), LAMP-1-positive late endosomes/lysosomes were more abundant, and

sometimes enlarged in Vps34cKO thyrocytes as compared to control. In addition, the

intracellular Tg protein, found in most Vps34cKO thyrocytes (Fig. 7A, arrowheads), most often

colocalized with LAMP-1-positive structures. This indicates that colloidal Tg was endocytosed

by thyrocytes, but that trafficking to or Tg proteolysis within the lysosomes was slower or

impaired, as compared to controls. Of note, luminal cells also contained LAMP-1 structures

positive for Tg (Fig. 7A, right).

Additional supporting evidence for impaired lysosomal proteolysis came from the

analysis of I-Tg (Fig. 7B). Indeed, Vps34^{cKO} thyrocytes often contained intracellular signal for I-

Tg, colocalizing within LAMP-1 structures (Fig. 7B, enlarged boxes,). Thus, in addition to

reduced iodine organification, defective lysosomal proteolysis of T_3/T_4, if/when hormonogenic

peptides were still formed on Tg, was also accounting for the low T_3/T_4 plasma levels observed

in Vps34^{cKO} (Fig. 2).

Evaluation of autophagy in Vps34^{cKO} thyroid

The deletion of Vps34 in the kidney also causes a block of autophagy (19). We therefore

assessed the expression of p62 (also called sequestosome-1, or SQSTM1), a polyubiquitin-

binding protein that interacts with LC3b (microtubule associated protein 1 light chain) on the

autophagosome membrane and is normally continuously degraded by the autophagy process

(31). Since p62 accumulates when completion of autophagy is inhibited, p62 can be used as a

marker to study autophagic flux (32). Although we found a weak punctiform LC3 signal in

control thyrocytes, we observed no p62 signal, indicating normal autophagic flux in control

thyroid. As reported, few LAMP-1-labelled structures were found in control thyroid. In

contrast, much larger structures, mostly double-labeled for p62 and LC3, were easily detected

in the cytoplasm of Vps34^{cKO} thyrocytes. As the LAMP-1 signal was increased in Vps34^{cKO}, these

p62/LC3-positive punctae sometimes co-localized with LAMP-1. We concluded that LC3 could

be recruited on p62 aggregates but that progression to autophagosome maturation and fusion

with, and degradation by, lysosomes was arrested. This suggested that deletion of Vps34 in

thyrocytes abrogated the autophagic flux.

Recent evidences have demonstrated that p62 is also an activator of the antioxidant

KEAP1/NRF2 pathway, which impacts on thyrocyte physiology and Tg biology (33). We thus

examined the possibility that stabilized p62 would compete with NRF2 for KEAP1 binding in

Vps34^{cKO}, thus allowing NRF2 to reach the nucleus and activate gene expression. We

quantified the mRNA levels of *Nrf2* and its target genes *Nqo1*, *Gpx2* and *Txnrd1*. Although we

measured a slight decrease of *Nrf2* mRNA in Vps34^{cKO}, two of its target genes, namely the

quinone reductase, *Nqo1*, and the thioredoxin reductase 1, *Txnrd1*, were upregulated 1.9- and

3.5-fold, respectively (Fig. 8B). This observation supports the hypothesis that KEAP1

interaction with p62 is favored in Vps34^{cKO} and that NRF2 is thus displaced and able to migrate

to the nucleus to regulate gene expression.

We also evaluated macroautophagy/mitophagy by immunolocalizing TOM20 and

LAMP-1. In Vps34^{cKO} thyrocytes, TOM20-labeled structures did not colocalize with the more

abundant, enlarged LAMP-1+ structures (Suppl. Fig. 2A), as we also reported in Vps34^{cKO}

kidney proximal tubular cells (19). This suggested that sequestration of mitochondria into

autophagosomes is either not triggered, which is unlikely for altered cells, or abortive at P14

in Vps34^{cKO} thyrocytes, and/or that fusion of autophagosomes with lysosomes is prevented.

We then evaluated the chaperone-mediated autophagy by immunolocalizing LAMP-2A. As

compared to control thyroid, LAMP-2A-labeled structures were more abundant in Vps34^{cKO}

thyrocytes (Suppl. Fig. 2B), as we also reported in Vps34^{cKO} kidney proximal tubular cells (19).

Luminal cells present in Vps34^{cKO} follicles are macrophages

Finally, we investigated the origin of the cells present in the colloidal space. As

observed in Figure 7A, cells trapped in the follicular lumen showed a strong LAMP-1 signal

(Fig. 7A). However, they were negative for E-cadherin (Figs. 1D and 5A). In addition, we found

that luminal cells were also negative for the permanently-expressed thyrocyte-specific

transcription factor, TTF-1 (Fig. 9A), indicating that these cells were not derived from the

thyrocyte lineage. Lineage-tracing experiments on Pax8-Cre; Vps34$^{fl/fl}$;ROSA-STOP-YFP pups at

P14 further revealed YFP-positive signal only in follicle-delineating thyrocytes and not in the

luminal cells (Fig. 9B). Thus, luminal cells had never expressed Pax8 and were thus not derived

from thyrocyte progenitors. Although infiltration by C-cells was a possibilty, we failed to label

luminal cells for the Prox1 transcription factor. As luminal cells displayed LAMP-1 signal and

were also positive for Tg (Fig. 7A), we tested the possibility that luminal cells were infiltrating

macrophages. This was the case: luminal cells were labeled by the macrophage marker F4/80

(Fig. 10A), but not by E-cadherin. To confirm that colloid was consumed by infiltrated

macrophages, we performed a triple immunolabelling for F4/80, LAMP-1 and I-Tg. Luminal

F4/80-positive cells displayed huge LAMP-1-positive lysosomes filled with I-Tg (Fig. 10B). Thus,

colloid consumption (Tg in Fig. 7A and I-Tg in Fig. 9D) by infiltrated macrophages provides a

third explanation for the low T$_4$ plasma level observed in Vps34^{cKO}.

Discussion

In this study, we report that deleting Vps34 in thyrocytes by Pax8-Cre-driven

recombination causes several defects in the thyroid: (i) doubling of thyroid weight and

perturbed thyroid parenchyma organization, with reduced PAS$^+$ colloidal spaces; (ii) severe

hypothyroidism with collapsed plasma T$_4$ levels and very high TSH; (iii) strong decrease of ^{125}I

organification, at comparable ^{125}I uptake, and of T$_4$ formation on thyroglobulin (detected by

immunofluorescence as early as P3); (iv) defective apical polarization; (v) impaired lysosomal

proteolysis; (vi) infiltration of macrophages in the colloid. Some of these features, combined

with delayed "eye opening" and impaired postnatal growth after two weeks (18), phenocopy

the impact of *Duoxa2* KO in the thyroid (34).

DUOXA2 is a chaperone protein required for the correct localization of DUOX2 at the

apical pole of thyrocyte, where the complex (DUOX2/DUOXA2) produces H_2O_2. In the absence

of DUOXA2, H_2O_2 is not produced and subsequent thyroperoxidase-mediated oxidation of

iodide into reactive compounds and thyroglobulin iodination are abolished (34). The similarity

of the severity of hypothyroidism in Vps34[cKO] and *Duoxa2* KO mice prompted us pay a closer

look at DUOXA2 in Vps34[cKO]. We found that the expression of *Duoxa2* started decreasing after

P3, reaching two-fold lower values at P14. It is interesting to note that expression of *Duox2*

and *Duoxa2* mRNAs was quantitatively different, despite of the fact that they share the same

promoter (35). Whether decreased *Duoxa2* mRNA expression is reflected by equivalent two-

fold decrease of DUOX2/DUOXA2 complex at the apical pole is unknown. However, it is very

unlikely that a two-fold decrease would by itself so severely impact on thyroglobulin

iodination, since heterozygous mice for *Duoxa2* deletion have virtually no phenotype. In

addition, our measurements of H_2O_2 levels in total Vps34[cKO] thyroid extracts also support

normal, or increased, function of the DUOX2/DUOXA2 complex. Instead, based on the known

functions of Vps34 in kidney PTCs, where its inactivation causes lack of apical localization of

endocytic receptors (megalin, cubilin) and solute transporters (NaPi-IIa, SGLT-2) (19), it would

be extremely interesting to localize actors of thyroid hormonogenesis on thyroid sections, and

assess their basolateral or apical addressing in Vps34[cKO] thyrocytes. Unfortunately, reliable

antibodies to detect most of these proteins are not yet available for mice. Nevertheless, [125]I

uptake and processing experiments suggests normal localization and function of the sodium-

iodine symporter, NIS, and/or the alternative transporter SLC26A7 (3), and rather support the

hypothesis of defective apical localization of one or several actors involved in thyroid

hormonogenesis (Ano1, pendrin, TPO, DUOX, DUOXA).

Characteristic histopathological alterations combined with the very high plasma TSH

levels and two-fold increased *Tshr* expression in Vps34^{cKO} thyroid suggest that thyrocytes are

in a hyperstimulated state. For example, DuoxA2 KO, which display very high TSH levels,

present a 20- and 5-fold higher expression level of *Nis* and *Tpo*, as compared to controls (34).

However, the expression of these two sensitive target genes of the TSH signaling pathway, *Nis*

and *Tpo*, was surprisingly unchanged in Vps34^{cKO}, arguing against thyrocyte hyperstimulation.

We favor the hypothesis of TSHR mistrafficking into intracellular vesicles. Recent work in

Drosophila revealed that Vps34 inactivation or pharmacological inhibition using the small

molecule inhibitor, SAR405, caused alteration of cell polarity and disruption of epithelial

architecture by relieving LKB1 inhibition and triggering JNK activation (14). A role of Vps34 in

epithelial organization and polarity was also observed in 3D cultures of Caco-2 kidney cells

(14). It would be interesting to analyze the activation states of LKB1 and JNK in Vps34^{cKO}

thyroid, and, if modified, to cross Vps34^{cKO} with floxed LKB1 alleles.

The expression of the two SLC transporters, *Slc26a7* and *Slc26a4*, came out as a

surprise in Vps34^{cKO}. Decreased expression of Slc26a7 (4-fold) in Vps34^{cKO} may affect entry of

iodine in the thyroid. However, it should be mentioned that this alternative basolateral

transporter, SLC26A7, may only play an indirect role on iodine uptake (3). In addition,

expression of the main transporter, *Nis*, is at least 16-fold more important than that of

Slc26a7. This may explain why ^{125}I uptake was not affected. Expression of apical *Slc26a4*

(pendrin) in Vps34^{cKO} was even more dramatic with a 20-fold increase in expression. However,

we do not think that this would have affected apical transport of iodine. Indeed, in control

thyroids, expression level of *Slc26a4* were 60-fold less important than those of *anoctamin1* (6

Ct). This supports a more prominent role for *anoctamin1* in iodine apical transport, as

demonstrated by Twyffels (36). These changes in expression of *Slc26a7* and *Slc26a4* could

however impact on thyrocyte ion balance. Indeed, SLC26A7 and SLC26A4 transporters have

opposite action on chloride ions at the basolateral and apical membranes, respectively.

Decreased levels of SLC26A7 at the basolateral membrane may decrease the export of

chloride out of the thyrocyte (3). On the contrary the increase of SLC26A4 at the apical

membrane may increase the entry of chloride in the thyrocyte. Accumulation of chloride may

in turn decrease intracellular pH and causes cellular stress in Vps34^{cKO}.

The weak PAS staining in Vps34^{cKO} could be due to decreased exocytosis of Tg or

increased Tg endocytosis. We indeed very often observed intracellular Tg in Vps34^{cKO}. We

suggest that the defective process is the endocytic route rather than exocytosis because we

readily observed intracellular structures positive for iodinated Tg. This is rarely the case in

control thyrocyte where I-Tg is rapidly proteolytically processed. In addition, our work also

evidenced increased abundance and size of LAMP-1-positive late endosomes/lysosomes in

Vps34^{cKO} thyrocytes, with accumulation of Tg and the autophagic marker, p62, indicative of a

defective lysosomal trafficking and/or function. In addition to impaired Tg iodination, low T$_4$

plasma levels thus also probably result from impaired endocytic transport of iodo-Tg to

lysosomes and/or proteolytic excision of T$_3$/T$_4$ therein. Several studies have reported a role

for Vps34 in endocytic trafficking to lysosomes (13, 15, 30, 37) and activation of lysosomal

proteases (15). In our study on Vps34^{cKO} kidney PTCs, we also observed enlargement of

lysosomes and their filling by undigested material, labelled for a variety of antigens (19). In

Vps34^{cKO} thyrocytes, we also observed increased abundance and size of LAMP-1-positive

compartments filled with Tg and I-Tg. Based on these and previous findings showing that

Vps34 deletion leads to late endosome/lysosome enlargement and defective lysosomal

function (37), we assume a defective proteolytic excision of T$_3$/T$_4$ from iodo-Tg, analogous to

what we reported in cystinotic *Ctns$^{-/-}$* mice (38). Cystinosis is a lysosomal storage disease due

to deletion or inactivating mutations of the lysosomal cystine transporter, cystinosin (39).

Defective cystinosin impacts on lysosomal proteolysis, presumably due to endolysosomal

mislocalization, impaired pro-cathepsin maturation and altered luminal redox status;

autophagic flux is also altered (40). Like cystinotic patients, $Ctns^{-/-}$ mice show compensated

hypothyroidism with moderate increase of TSH, thyrocyte hyperplasia and proliferation,

combined with progressive colloid depletion and evidence of increased endocytosis into

colloid droplets. Iodo-thyroglobulin could be detected in $Ctns^{-/-}$ but not wild-type thyrocyte

lysosomes, under identical labelling conditions, further indicating defective proteolysis.

It has recently become clear that Vps34 may act on lysosome positioning, which is

crucial to autophagosome formation (41,42). Similar to the kidney PTC defect, we observed

an increased p62 signal in Vps34^{cKO} thyroids at P14. Accumulation of p62 and LC3b in Vps34^{cKO}

indicates a block in the autophagy process. In addition, p62 accumulation also suggests

activation of the antioxidant NRF2 pathway by competitive binding of p62 to KEAP1 (33). We

indeed observed increased expression of two NRF2 target genes, namely the quinone

reductase, *Nqo1*, and the thioredoxin reductase 1, *Txnrd1*. Considering the expected key role

of Vps34 in macroautophagy, our data on mitophagy do not prove, but are compatible with

the suggestion that this important homeostatic process is abrogated upon Vps34 deficiency.

A final observation deserving discussion is the occurrence of viable cells in the follicular

lumens of Vps34^{cKO} thyroids. Our first hypothesis was that defective endocytic and autophagic

routes to the lysosomes could impact thyrocyte homeostasis, thus causing cellular stress.

Cellular stress, induced by defective autophagosome and lysosomal function, or by activation

of stress kinase such as JNK, could explain shedding of thyrocytes cells into the colloidal space.

However, luminal cells of Vps34^{cKO} thyroids were negative for E-cadherin and the thyrocyte-

specific transcription factor, TTF-1. Furthermore, lineage tracing experiments demonstrated

that these luminal cells had never expressed Pax8 and thus did not derive from thyrocytes.

The second hypothesis was that the cellular stress and loss of tissue homeostasis would recruit

macrophages. Indeed, we found that luminal cells were labelled for the conventional

macrophage marker, F4/80, are proliferating and display high LAMP-1 signal. In addition, Tg

and I-Tg colocalized with the LAMP-1 structures. This observation contributes to explain the

colloid exhaustion observed in Vps34^{cKO} and could thus be responsible for the low levels of

circulating T$_4$. Although the signal(s) attracting macrophages to the colloid is(are) unknown,

the loss of tight junction integrity might facilitate their invasion.

Acknowledgements

This work was supported by grants from the Fonds pour la Recherche Scientifique (F.R.S-FNRS,

J.0076.18, Belgium), Université catholique de Louvain (Actions de Recherche concertées to

CEP, ARC 15/20-065), and Fondation Roi Baudouin. SR and X-HL were supported by a grant

DK15070 from the National Institutes of Health, USA. Work in the laboratory of BV was

supported by grants from the MRC (G0700755) and BBSRC (BB/I007806/1; BB/M013278/1).

The core facility for Imaging Cells and Tissues was also financed by National Lottery, Région

bruxelloise, Région wallonne, Université catholique de Louvain and de Duve Institute. The

authors thank Prof. C. Ris-Stalpers for iodo-thyroglobulin antibody, Abdelkadder El Kaddouri

and Claude Massart for technical help. GG held a fellowship from the Fonds pour la formation

à la Recherche dans l'Industrie et l'Agriculture (FRIA, Belgium), TW was postdoctoral

researcher supported by UCLouvain, OD is supported by Télévie, VJ by the Cystinosis Research

Foundation, CS is a UCLouvain teaching assistant, HGC was a postdoctoral researcher and CEP

is a Senior Research Associate at F.R.S-FNRS.

Name and address of corresponding author

Prof. Christophe E. Pierreux

Cell Biology Unit

de Duve Institute and Université Catholique de Louvain

75, Avenue Hippocrate

B-1200 Brussels

Belgium

Email: christophe.pierreux@uclouvain.be

References

1. Colin IM, Denef JF, Lengele B, Many MC, Gerard AC 2013 Recent insights into the cell biology

of thyroid angiofollicular units. Endocr Rev 34:209-238.

2. Carvalho DP, Dupuy C 2017 Thyroid hormone biosynthesis and release. Mol Cell Endocrinol

458:6-15.

3. Cangul H, Liao X-H, Schoenmakers E, Kero J, Barone S, Srichomkwun P, Iwayama H, Serra

EG, Saglam H, Eren E, Tarim O, Nicholas AK, Zvetkova I, Anderson CA, Karet Frankl FE,

Boelaert K, Ojaniemi M, Jääskeläinen J, Patyra K, Löf C, Williams ED, UK10K Consortium,

Soleimani M, Barrett T, Maher ER, Chatterjee VK, Refetoff S, Schoenmakers N 2018

Homozygous loss-of-function mutations in SLC26A7 cause goitrous congenital

hypothyroidism. J Clin Ivest Insight 3:e99631.

4. Koumarianou P, Gomez-Lopez G, Santisteban P 2017 Pax8 controls thyroid follicular polarity

through cadherin-16. J Cell Sci 130:219-231.

5. Shivas JM, Morrison HA, Bilder D, Skop AR 2010 Polarity and endocytosis: reciprocal

regulation. Trends Cell Biol 20:445-452.

6. Rodriguez-Boulan E, Macara IG 2014 Organization and execution of the epithelial polarity

programme. Nat Rev Mol Cell Biol 15:225-242.

7. Morrison HA, Dionne H, Rusten TE, Brech A, Fisher WW, Pfeiffer BD, Celniker SE, Stenmark

H, Bilder D 2008 Regulation of early endosomal entry by the Drosophila tumor

suppressors Rabenosyn and Vps45. Mol Biol Cell 19:4167-4176.

8. Bilanges B, Posor Y, Vanhaesebroeck B 2019 PI3K isoforms in cell signaling and vesicle

trafficking. Nat Rev Mol Cell Biol May 20. doi: 10.1038/s41580-019-0129-z. [Epub

ahead of print].

9. Lindmo K, Stenmark H 2006 Regulation of membrane traffic by phosphoinositide 3-kinases.

J Cell Sci 119:605-614.

10. Juhasz G, Hill JH, Yan Y,Sass M, Baehrecke EH, Backer JM, Neufeld TP 2008 The class III

PI(3)K Vps34 promotes autophagy and endocytosis but not TOR signaling in Drosophila.

J Cell Biol 181:655-666.

11. Backer JM 2008 The regulation and function of Class III PI3Ks: novel roles for Vps34.

Biochem J 410:1-17.

12. Jaber N, Zong WX 2013 Class III PI3K Vps34: essential roles in autophagy, endocytosis, and

heart and liver function. Ann NY Acad Sci 1280:48-51.

13. Bechtel W, Helmstädter M, Balica J, Hartleben B, Kiefer B, Hrnjic F, Schell C, Kretz O, Liu S,

Geist F, Kerjaschki D, Walz G, Huber TB 2013 Vps34 deficiency reveals the importance

of endocytosis for podocyte homeostasis. J Am Soc Nephrol 24:727-743.

14. O'Farrell F, Lobert VH, Sneeggen M, Jain A, Katheder NS, Wenzel EM, Schultz SW, Tan KW,

Brech A, Stenmark H, Rusten TE 2017 Class III phosphatidylinositol-3-OH kinase

controls epithelial integrity through endosomal LKB1 regulation. Nat Cell Biol 19:1412-

1423.

15. Ronan B, Flamand O, Vescovi L, Dureuil C, Durand L, Fassy F, Bachelot M-F, Lamberton A,

Mathieu M, Bertrand T, Marquette J-P, El-Ahmad Y, Filoche-Romme B, Schio L, Garcia-

Echeverria C, Goulaouic H, Pasquier B 2014 A highly potent and selective Vps34

inhibitor alters vesicle trafficking and autophagy. Nat Chem Biol 10:1013-1019.

16. Nascimbeni AC, Codogno P, Morel E 2017 Phosphatidylinositol-3-phosphate in the

regulation of autophagy membrane dynamics. FEBS J 284:1267-1278.

17. Dupont N, Nascimbeni AC, Morel E, Codogno P 2017 Molecular Mechanisms of

Noncanonical Autophagy. Intl Rev Cell Mol Biol 328:1-23.

18. Carpentier S, N'Kuli F, Grieco G, Van Der Smissen P, Janssens V, Emonard H, Bilanges B,

Vanhaesebroeck B, Gaide Chevronnay HP, Pierreux CE, Tyteca D, Courtoy PJ 2013

ClassIII phosphoinositide 3-kinase/VPS34 and dynamin are critical for apical endocytic

recycling. Traffic 14:933-948.

19. Grieco G, Janssens V, Gaide Chevronnay HP, N'Kuli F, Van Der Smissen P, Wang TS, Shan J,

Vainio S, Bilanges B, Jouret F, Vanhaesebroeck B, Pierreux CE, Courtoy PJ 2018

Suppression of Vps34 PI 3-kinase activity in kidney proximal tubules abrogates apical

trafficking and causes a Fanconi syndrome. Sci Rep 8:14133.

20. Shan J, Jokela T, Skovorodkin I, Vainio S 2010 Mapping of the fate of cell lineages generated

from cells that express the Wnt4 gene by time-lapse during kidney development.

Differentiation 79:57-64.

21. Bouchard M, Souabni A, Busslinger M 2004 Tissue-specific expression of cre recombinase

from the Pax8 locus. Genesis 38:105-109.

22. Valet C, Levade M, Chicanne G, Bilanges B, Cabou C, Viaud J, Gratacap MP, Gaits-Iacovoni

F, Vanhaesebroeck B, Payrastre B, Severin S 2017 A dual role for the class III PI3K,

Vps34, in platelet production and thrombus growth. Blood 130:2032-2042.

23. Pohlenz J, Maqueem A, Cua K, Weiss RE, Van Sande J, Refetoff S 1999 Improved

radioimmunoassay for measurement of mouse thyrotropin in serum: strain

differences in thyrotropin concentration and thyrotroph sensitivity to thyroid

hormone. Thyroid 9:1265-1271.

24. Delmarcelle AS, Villacorte M, Hick AC, Pierreux CE 2014 An ex vivo culture system to study

thyroid development. J Vis Exp 6:88.

25. Hick AC, Delmarcelle AS, Bouquet M, Klotz S, Copetti T, Forez C, Van Der Smissen P,

Sonveaux P, Collet JF, Feron O, Courtoy PJ, Pierreux CE 2013 Reciprocal

epithelial:endothelial paracrine interactions during thyroid development govern

follicular organization and C-cells differentiation. Dev Biol 381:227-240.

26. Dupasquier S, Delmarcelle AS, Marbaix E, Cosyns JP, Courtoy PJ, Pierreux CE 2014

Validation of housekeeping gene and impact on normalized gene expression in clear

cell renal cell carcinoma: critical reassessment of YBX3/ZONAB/CSDA expression. BMC

Mol Biol 15:9.

27. Massart C, Hoste C, Virion A, Ruf J, Dumont JE, Van Sande J 2011 Cell biology of H2O2

generation in the thyroid: investigation of the control of dual oxidases (DUOX) activity

in intact ex vivo thyroid tissue and cell lines. Mol Cell Endocrinol 343:32-44.

28. Zhou X, Takatoh J, Wang F 2011 The mammalian class 3 PI3K (PIK3C3) is required for early

embryogenesis and cell proliferation. PloS one 6:e16358.

29. Chen J, Chen MX, Fogo AB, Harris RC, Chen JK 2013 mVps34 deletion in podocytes causes

glomerulosclerosis by disrupting intracellular vesicle trafficking. J Am Soc Nephrol.

24:198-207.

30. Bechtel, W, Helmstädter M, Balica J, Hartleben B, Schell C, Huber TB 2013 The class III

phosphatidylinositol   3-kinase   PIK3C3/VPS34   regulates   endocytosis   and

autophagosome-autolysosome formation in podocytes. Autophagy 9:1097-1099.

31. Cohen-Kaplan V, Ciechanover A, Livneh I 2016 p62 at the crossroad of the ubiquitin-

proteasome system and autophagy. Oncotarget 7:83833-83834.

32. Bjorkoy G, Lamark T, Pankiv S, Overvatn A, Brech A, Johansen T 2009 Monitoring

autophagic degradation of p62/SQSTM1. Methods Enzymol 452:181-197.

33. Ziros PG, Habeso IG, Chartoumpekis DV, Ntalampyra E, Somm E, Renaud CO, Bongiovanni

M, Trougakos IP, Yamamoto M, Kensler TW, Santisteban P, Carrascao N, Ris-Stalpers

C, Amendola E, Liao XH, Rossich L, Thomasz L, Juvenal GJ, Refetoff S, Sykiotos GP 2018

NFE2-related transcription factor 2 coordinates antioxidant defense with thyroglobulin

production and iodination in the thyroid gland. Thyroid 28:780-798.

34. Grasberger H, De Deken X, Mayo OB, Raad H, Weiss M, Liao XH, Refetoff S 2012 Mice

deficient in dual oxidase maturation factors are severely hypothyroid. Mol Endocrinol

26:481-492.

35. Christophe-Hobertus C, Christophe D 2010 Delimitation and functional characterization of

the bidirectional THOX-DUOXA promoter regions in thyrocytes. Mol Cell Endocrinol

317:161-167.

36. Twyffels L, Strickaert A, Virreira M, Massart C, Van Sande J, Wauquier C, Beauwens R,

Dumont JE, Galietta LJ, Boom A, Kruys V 2014 Anoctamin-1/TMEME16A is the major

apical iodide channel of the thyrocyte. Am J Physiol Cell Physiol 307:1102-1112.

37. Jaber N, Mohd-Naim N, Wang Z, DeLeon JL, Kim S, Zhong H, Sheshadri N, Dou Z, Edinger

AL, Du G, Braga VM, Zong WX 2016 Vps34 regulates Rab7 and late endocytic trafficking

through recruitment of the GTPase-activating protein Armus. J Cell Sci 129:4424-4435.

38. Gaide Chevronnay HP, Janssens V, Van Der Smissen P, Liao XH, Abid Y, Nevo N, Antignac

C, Refetoff S, Cherqui S, Pierreux CE, Courtoy PJ 2015 A mouse model suggests two

mechanisms for thyroid alterations in infantile cystinosis: decreased thyroglobulin

synthesis due to endoplasmic reticulum stress/unfolded protein response and

impaired lysosomal processing. Endocrinology 156:2349-2364.

39. Gahl WA, Thoene JG, Schneider JA 2002 Cystinosis. N Engl J Med. 347:111–121.

40. Cherqui S, Courtoy PJ 2017 The renal Fanconi syndrome in cystinosis: pathogenic insights

and therapeutic perspectives. Nat Rev Nephr 13:115-131.

41. Hong Z, Pedersen NM, Wang L, Torgersen ML, Stenmark H, Raiborg C 2017 PtdIns3P

controls mTORC1 signaling through lysosomal positioning. J Cell Biol 216:4217-4233.

42. Korolchuk VI, Rubinsztein DC 2011 Regulation of autophagy by lysosomal positioning.

Autophagy 7:927-928.

	Figure legends

Figure 1. Genetic and histopathological characterization. A. Extensive genetic excision of

Vps34 exon 21. Compared with control (black boxes), Vps34^{cKO} thyroid (red boxes) shows ~70%

reduction in exon 21 mRNA level at P14. The unchanged mRNA level spanning exons 22-24

serves as control. Boxes with median and percentiles of 10 WT and 11 cKO samples; ***, p<

0.0001 by Mann-Whitney non-parametric test. *B. Increased thyroid weight.* Compared with

control (black boxes), Vps34^{cKO} thyroid (red boxes) shows a two-fold increase in thyroid weight.

Boxes with median and percentiles of 8 WT and 6 cKO samples; **, p< 0.01 by Mann-Whitney

non-parametric test. *C. Histopathological evidence for colloid exhaustion.* As compared to

control thyroid tissues (n=3) where all follicles show regular lumen filling with homogenous

and intense PAS (Periodic Acid Shiff) staining, Vps34^{cKO} thyroids (n=3) present fewer, mostly

centrally located, PAS-stained follicles and with weaker staining intensity, other follicles that

appear empty (for further quantification, see Supplementary Figure 1). *D. Vps34^{cKO} thyrocytes*

are altered and follicles contains abundant non-epithelial cells. Nuclei are labeled by Hoechst

(shown in white); thyrocyte basolateral contours are labeled for E-cadherin (red); two lumen

boundaries are delineated by broken lines. As compared to control thyroid follicles, Vps34^{cKO}

follicular structures are frequently irregular and containing additional cells inside the lumen

(arrowheads). These cells are not labeled for E-cadherin.

Figure 2. Vps34^{cKO} mice display severe hypothyroidism associated with high TSH levels. As

compared to control mice (black boxes), T$_4$ plasma level in Vps34^{cKO} is extremely low (red

boxes). Conversely, TSH levels are dramatically elevated. (10 WT and 9 cKO samples; ***, p=

0.0003 by Mann-Whitney non-parametric test).

Figure 3. Relative expression level of main actors of thyroid hormonogenesis at P14. Gene expression analysis by RT-qPCR, presented as boxes with median and percentiles. As compared to control thyroid (black boxes), expression in Vps34[cKO] (red boxes) is preserved for the thyroid-specific genes *Nis*, *Tpo*, *Tg* and *Duox2*. Expression of the TSH receptor (*Tshr*) and of *Slc26a4* is significantly increased in Vps34[cKO], while expression of *Slc26a7* and *Duoxa2* is significantly decreased. Boxes with median and percentiles of at least 9 control and 9 Vps34[cKO] samples; ***, p< 0.001 by Mann-Whitney non-parametric test.

Figure 4. Normal uptake of iodine but major defect of organification in Vps34[cKO] thyroid. A. Comparable ^{125}I-iodine uptake in the thyroid in control genotypes (black box) and Vps34[cKO] thyroid (red box), but much reduced ^{125}I bound to protein (in %) in Vps34[cKO] thyroid, suggesting defective organification. Boxes with median and percentiles of 39 (10 Flox/+, 19 Flox/Flox, 10 Cre;Flox/+) and 12 Vps34[cKO] (Cre;Flox/Flox) samples; ***, p< 0.001 by Mann-Whitney non-parametric test. **B.** Level of H_2O_2 in total thyroid extracts show a trend to an increase in Vps34[cKO] thyroid. Boxes with median and percentiles of 9 control and 9 Vps34[cKO] samples; p= 0.077 by Mann-Whitney non-parametric test.

Figure 5. Normal basolateral but impaired apical organization of tyrocytes. A. Thyroid sections from control (left) and Vps34[cKO] (right) labeled for laminin (red), ezrin (green) and E-cadherin (white). Nuclei are labeled by Hoechst (shown in blue). In control thyroid, laminin surrounds follicles composed of thyrocytes delineated by basolateral E-cadherin and apical ezrin. In Vps34[cKO], laminin normally surrounds follicles composed of thyrocytes with well-defined basolateral E-cadherin but weaker and less regular apical ezrin. Note here apical membrane bulging in the colloidal space. *B.* Thyroid sections from control (left) and Vps34[cKO]

(right) labeled for ZO-1 (red), Na$^+$/K$^+$-ATPase (green) or β-catenin (white). Nuclei are labeled

by Hoechst (shown in blue). In control and Vps34^{cKO} thyroid, Na$^+$/K$^+$-ATPase and β-catenin are

correctly localized and restricted to the basolateral pole of the thyrocytes, indicating the

presence of a tight junction. However, the tight junction-associated protein ZO-1 is only

detected at few apico/basolateral junctions.

Figure 6. Evidence for defective iodide organification in Vps34^{cKO} thyroid: follicular lumina

with Tg but devoid of iodinated-Tg. **A.** Thyroid sections from control (left) and Vps34^{cKO} (right)

labeled for E-cadherin (green) and thyroglobulin (red), recognized either for a core protein

epitope (Tg), or a hormonogenic peptide (I-Tg). Control follicular lumina are round and

uniformly labeled for both Tg and I-Tg. In Vps34^{cKO}, follicular structures are less regular, with

the majority containing Tg, but not the hormonogenic peptide. In Vps34^{cKO}, two serial (tilted)

sections are shown. ***B. Comparison of Tg labelling in control and Vps34^{cKO} thyrocytes.*** In control

follicles, Tg is essentially restricted to lumen, with little or no signal inside thyrocytes. In

Vps34^{cKO}, Tg is readily detected within thyrocytes as collections of submicrometric dots or

larger spheres (arrowheads). Of note, some luminal cells (broken line in middle panel) also

contain Tg-labeled spheres, indicating active endocytosis (arrow).

Figure 7. Vps34^{cKO} thyrocytes display increased LAMP-1 signal and impaired I-Tg proteolytic

processing. **A.** Thyroid sections from control (left) and Vps34^{cKO} (right) labeled for

thyroglobulin (red), E-cadherin (green) and lysosomal LAMP-1 (white). Nuclei are labeled by

Hoechst (shown in blue). In control, thyrocytes rarely contain intracellular Tg and LAMP-1

signal is weak. In Vps34^{cKO}, most thyrocytes have intracellular Tg (arrowheads) and LAMP-1

signal is much increased. Insets show magnification with separate emissions, then merged,

channels. Note colocalization of Tg with LAMP-1 in luminal cells (at right). **B.** Thyroid sections

from control (left) and Vps34^{cKO} (right) labeled for iodinated-thyroglobulin (I-Tg, red), E-

cadherin (green) and lysosomal LAMP-1 (white). Nuclei are labeled by Hoechst (shown in blue).

As compared to control follicles which show a weak signal for LAMP-1, and rare intracellular I-

Tg, LAMP-1 signal is intense in Vps34^{cKO} thyrocytes and intracellular I-Tg dots are frequently

observed (arrowheads). As shown in the enlargements below, LAMP-1-labeled structures in

Vps34^{cKO} thyrocytes are enlarged or vacuolated as compared to control.

Figure 8. Vps34^{cKO} thyrocytes show strong accumulation of p62 and LC3b dots. **A.** Thyroid

sections from control (left) and Vps34^{cKO} (right) labeled for p62 (red), LC3b (green) and

lysosomal LAMP-1 (white). Nuclei are labeled by Hoechst (shown in blue). Compared with

control follicles, which show low LAMP-1 and almost no signal for p62 and LC3b, all three

markers are very strong in Vps34^{cKO} thyroid sections. p62 perfectly co-localizes with LC3b and

labels large aggregates. These aggregates are adjacent to LAMP-1-positive structures. **B.**

Increased NRF2 signaling. Gene expression analysis by RT-qPCR, presented as boxes with

median and percentiles. As compared to control thyroid (black boxes), expression of *Nrf2* is

reduced in Vps34^{cKO} (red boxes), but level of two out of three NRF2 target genes, *Txnrd1* and

Nqo1 is increased. Boxes with median and percentiles of at least 9 control and 9 Vps34^{cKO}

samples; *, $p < 0.05$ and ***, $p < 0,001$ by Mann-Whitney non-parametric test.

Figure 9. Luminal cells in Vps34^{cKO} thyroid are not thyrocytes. ***A.*** Immunolabelling for

thyrocyte transcription factor-1 (TTF-1, green); nuclei are visualized with the Hoechst stain

(shown in white). In both control and Vps34^{cKO} thyroid, nuclei of thyrocytes circumscribing the

follicular lumina are all labeled by the thyrocyte-specific transcription-factor-1, TTF-1. As

indicated by the arrowheads, nuclei of luminal cells show no signal for TTF-1 (only white signal

representing Hoechst). *B.* Immunolabelling for E-cadherin (red) and YFP (green); nuclei are

visualized with the Hoechst stain (shown in blue). Representative images of Vps34cKO thyroid

sections reveal expression of YFP only in cells surrounding the follicular lumina, i.e. in cells

where Pax8-Cre has been active. Luminal cells do not derive from thyrocyte progenitors as

they are negative for YFP.

Figure 10. Luminal cells in Vps34^{cKO} thyroid are macrophages. *A.* Immunolabelling for E-

cadherin (red), and for the conventional macrophage marker (F4/80, green); nuclei are

visualized with the Hoechst stain (shown in white). In Vps34^{cKO} thyroid, F4/80 signal can be

found in the interstitium (arrows), but also in the colloidal space, thus identifying luminal cells

as infiltrating macrophages. **D. Luminal cells are taking up the colloid.** Immunolabelling for I-

Tg (red), F4/80 (green) and LAMP-1 (white). Nuclei are visualized with the Hoechst stain

(shown in blue). Luminal cells in Vps34^{cKO} have abundant LAMP-1 structures that surround I-Tg

droplets. Arrowheads show additional I-Tg in thyrocytes. Broken lines indicate luminal

contours.

GriecoFig1

GriecoFig2

GriecoFig3

GriecoFig4

GriecoFig5

GriecoFig6

GriecoFig7

GriecoFig8

A.

Control   Vps34^{cKO}

TTF-1   Hoechst

B.

Vps34^{cKO}

E-cadherin   YFP   Hoechst

GriecoFig9

GriecoFig10