1

Vps34/PIK3C3 deletion in thyroid impairs thyroid hormonogenesis and autophagic flux

Giuseppina Grieco[1], Tongsong Wang[1], Ophélie Delcorte[1,] Virginie Janssens[1], Catherine

Spourquet[1], Aurélie Strickaert[2], Héloïse P. Gaide Chevronnay[1], Xiao-Hui Liao[3], Benoît

Bilanges[4], Samuel Refetoff[3], Bart Vanhaesebroeck[4], Carine Maenhaut[2], Pierre J. Courtoy[1*],

and Christophe E. Pierreux[1*#]

[1], Cell Biology Unit, de Duve Institute and Université Catholique de Louvain, 1200 Brussels,

Belgium;

[2], Institute of Interdisciplinary Research (IRIBHM); Université Libre de Bruxelles, 1070 Brussels,

Belgium

[3], Departments of Medicine (X-H.L., S.R.) and Pediatrics and Genetics (S.R), The University of

Chicago, Chicago, Illinois 60637;

[4], UCL Cancer Institute; University College London, 72 Huntley Street, London WC1E 6DD, UK;

[#], correspondence to christophe.pierreux@uclouvain.be

[*], equal contributions

Running title : Vps34 controls TH synthesis

Key words : Vps34, thyroid, hormone, organification, autophagy

Dr. Giuseppina Grieco, g.pia@hotmail.it

Dr. Tongsong Wang, tswang0802@hotmail.com

Ms. Ophélie Delcorte, ophelie.delcorte@uclouvain.be

Ms. Virginie Janssens, virginie.janssens@uclouvain.be

Ms. Catherine Spourquet, catherine.spourquet@uclouvain.be

Dr. Aurélie Strickaeart, aurelie.strickaert@ulb.ac.be

Dr. Héloïse Gaide-Chevronnay, heloisegaide@hotmail.com

Dr. Xia Hui Liao, xliao1@uchicago.edu

Dr. Benoit Bilanges, b.bilanges@ucl.ac.uk

Prof. Samuel Refetoff, srefetof@uchicago.edu

Prof. Bart Vanhaesebroeck, bart.vanh@ucl.ac.uk

Prof. Carine Maenhaut, cmaenhau@ulb.ac.be

Prof. Pierre J. Courtoy, pierre.courtoy@uclouvain.be

Prof. Christophe E. Pierreux, christophe.pierreux@uclouvain.be

Abstract (327 words, 350 allowed)

BACKGROUND: The production of thyroid hormones (T_3, T_4) depends on thyroid organization

in follicles, lined by a monolayer of thyrocytes with strict apico-basal polarity. Polarization

supports vectorial transport of thyroglobulin for storage into, and recapture from, the colloid.

It also allows selective addressing of channels, transporters, pumps and enzymes to their

appropriate basolateral (NIS and Na^+/K^+-ATPase) or apical membrane domain (pendrin,

anoctamin, DUOX2, DUOXA2 and TPO). How these actors of T_3/T_4 synthesis reach their final

destination remains poorly understood. Vps34/PIK3C3 is now recognized as a main

component in the general control of vesicular trafficking and of cell homeostasis via

autophagy. We recently reported that conditional Vps34 inactivation in kidney proximal

tubular cells by Pax8-driven excision prevents normal addressing of apical membrane proteins

and causes abortive macroautophagy.

METHODS: Vps34 was inactivated using a Pax8-driven Cre recombinase system. The impact

of Vps34 inactivation in thyrocytes was analyzed by histological, immunolocalization and

mRNA expression profiling. Thyroid hormone synthesis was assayed by ^{125}I injection and by

serum plasma analysis.

RESULTS: Vps34^{cKO} mice were born at the expected Mendelian ratio and showed normal

growth until postnatal day 14, then stopped growing and died at around 1 month of age. We

therefore analyzed thyroid Vps34^{cKO} before postnatal day 14. We found that loss of Vps34 in

thyrocytes causes: (i) disorganization of thyroid parenchyma with abnormal thyrocyte and

follicular shape and reduced PAS$^+$ colloidal spaces; (ii) impaired ^{125}I organification at

comparable uptake and frequent occurrence of follicles with luminal thyroglobulin but non-

detectable T4-bearing thyroglobulin; (iii) severe non-compensated hypothyroidism with

extremely low T_4 levels (<0.25 ± 1.5 μg/dL) and huge TSH plasma levels (19,300 ± 10,500

mU/L); (iv) intense signal in thyrocytes for the lysosomal membrane marker, LAMP-1, as well

as thyroglobulin and the autophagy marker, p62, indicating defective proteolysis.

CONCLUSIONS: We conclude that Vps34 is crucial for thyroid hormonogenesis, at least by

controlling delivery of apical actors responsible for biogenesis of thyroid hormones on Tg as

well as defective proteolytic T_3/T_4 excision in lysosomes.

 Introduction

The main function of the thyroid gland is to produce the hormones, T_3/T_4 and

calcitonin, which are essential for the regulation of metabolic processes [1,2]. The production

of T_3 and T_4 depends on the organization of thyroid epithelial cells, the thyrocytes, into

functional and independent units, the follicles. These are composed of a single layer of

polarized thyrocytes that form a spherical structure delineating an internal space or lumen

where the thyrocyte secretory product, thyroglobulin, is stored in a colloidal form, thus called

the colloid lumen. As thyrocytes communicate via gap junctions, each follicle functions as an

integrated unit. Thyroid hormone synthesis depends on apico-basal cell polarity that allows

localization of channels, transporters, pumps and enzymes at the appropriate membrane

domains. Iodide from the bloodstream freely traverses the fenestrated endothelium of the

thyroid capillaries and is taken up into thyrocytes via the basolaterally localized Na^+/I^-

symporter (NIS), thanks to a Na^+ gradient generated by the Na^+/K^+-ATPase. An alternative

basolateral transporter, SLC26A7, has been recently reported to also control iodide uptake,

although its role might not be direct [3]. Iodide diffuses freely within thyrocytes and is next

transported across the apical membrane into the colloid space via apically localized

transporters such as pendrin or anoctamin. Iodide is then rapidly oxidized into iodine by

thyroperoxidase (TPO) located at the apical membrane, in the presence of H_2O_2, generated by

the DUOX2/DUOXA2 apical membrane complex. Iodine is incorporated into accessible

tyrosine residues close to N- and C-termini of thyroglobulin (Tg), a large protein secreted by

thyrocytes into the colloid space. Iodotyrosine rearranges into hormonogenic peptides, which

are the direct T_3 and T_4 precursors. Thyroid hormone synthesis thus requires basal localization

of NIS and Na^+/K^+-ATPase, apical localization of pendrin, anoctamin, TPO, DUOX2 and

DUOXA2, as well as apical delivery of Tg into the colloid lumen and endocytic uptake of iodoTg

into thyrocyte lysosomes (1,2). Specific regulators of thyroid follicular organization have

recently been identified (4). Transcriptomic comparison of thyroid FRT cells cultured in 2D

monolayers and in 3D spherical follicles indeed revealed involvement of structural and

functional cell elements such as adherens and tight junctions (cadherin-16), cytoskeleton

proteins, ions channels, proteins involved in differentiation, and components of the trafficking

machinery (e.g. myosin-Vb, Rab17)(4).

There is strong evidence that vesicular trafficking is critical for apico-basal polarization

and epithelial function (5-7), but the role of vesicular trafficking in thyroid function is

incompletely understood. Vps34/PIK3C3 (also referred to as type III PI3K) has long been

recognized as a main actor involved in the general control of endocytic vesicular trafficking (8-

12). Vps34/PIK3C3 also plays an important role in epithelial organization in *Drosophila* (12)

and in autophagy (11,14), both in the initiation of autophagosome formation and the

progression towards autophagosome-lysosome fusion (15,16). We recently inactivated Vps34

in kidney proximal tubular cells (PTC) using Wnt4-Cre and Pax8-Cre (17,18). Wnt4-Cre driven

Vps34 excision caused major perinatal lethality, with abnormalities in kidney cortex

development and PTC apical differentiation in the surviving mice. With the Pax8-Cre, which

triggers recombination in nephrogenic tubules and in nonvascular components of glomeruli

two days later than Wnt4-Cre (19,20), we observed normal mouse growth until postnatal (P)

day 14 (18). Then, body weight levelled off and all pups died at ~3-5 weeks of age, probably

due to kidney failure. Structural and functional studies of Vps34[cKO] kidneys revealed

preserved PTC membrane polarity, but impaired apical membrane protein trafficking, thus

causing a general PTC dysfunction known as renal Fanconi-like syndrome, manifested by

polyuria and low-molecular weight proteinuria. Vps34[cKO] also displayed impaired lysosome

size/positioning and blocked autophagy, thereby causing cell vacuolization. We concluded

that Vps34 is a crucial component of the trafficking machinery necessary for differentiated

PTC function and is essential for overall PTC homeostasis (18). As Pax8-Cre also triggers

recombination in thyrocytes, we here investigated the role of Vps34/PIK3C3 in thyroid

function and homeostasis.

Materials and methods

Mice

Vps34$^{fl/fl}$ mice have been described (20). Pax8-Cre mice were obtained from Dr. M.

Busslinger (20). Vps34$^{fl/fl}$ mice were crossed with Pax8-Cre;Vps34$^{fl/+}$ mice to generate

conditional targeted excision of Vps34 exon 21 in the thyroid of 25% of offspring (Vps34^{cKO}).

Vps34$^{fl/fl}$ mice were used as control. Mice were treated according to the NIH Guide for Care

and Use of Laboratory Animals, and experiments were approved by the University Animal

Welfare Committee, Université Catholique de Louvain (2016/UCL/MD/006 and

2018/UCL/MD/026).

Plasma, tissue collection and histology

Blood was collected by eye sinus puncture at sacrifice (P14) under irreversible

anesthesia by xylazine 2% and ketamine 50 mg/ml (200 μl/mice i.p.). Thyroid lobes were

excised, fixed by immersion in neutral-buffered formaldehyde (4% F) at 4°C under stirring

overnight. Samples were paraffin-embedded or equilibrated overnight in 20% sucrose and

embedded in Tissue-Tek Optimal Cutting Medium (Sakura Finetek) for 5 μm-thick cryostat

sections.

TSH and T4 plasma concentrations

Plasma TSH concentrations were measured by a sensitive, heterologous, disequilibrium double-antibody precipitation RIA as described (22). T4 concentration was measured by coated-tube RIA (Siemens Medical Solution Diagnostics, Los Angeles, CA).

Immunofluorescence

147   Immunofluorescence was performed on 5 µm-thick frozen sections or on 6 µm-thick paraffin sections (18). Antigen retrieval was promoted in citrate buffer, pH 6.0, at 98°C for 20 min using a Lab Vision Pretreatment Module™ (Thermo Scientific). After permeabilization with PBS/0.3% Triton-X100 for 5 min, non-specific sites were blocked by 1-h incubation in PBS/0.3% Triton-X100 with 10% bovine serum albumin (BSA) and 3% milk, followed by primary antibodies (described in Supplementary Table I) in blocking buffer at 4°C overnight. After extensive washing, sections were incubated with the appropriate AlexaFluor-secondary antibodies in 10% BSA/0.3% Triton-X100 at room temperature for 1 h, extensively washed, mounted with Faramount Aqueous Mounting Medium (Dako) and imaged on a spinning disk confocal microscope using a Plan Apochromat 100x/1.4 Oil DIC objective (Cell Observer Spinning Disk; Zeiss). For whole thyroid section recording, images were acquired using Zeiss Mirax Midi slide scanner, stitched and analyzed using Pannoramic Viewer software.

RT-qPCR

160   Total RNA was extracted from thyroid lobes using TRIzol Reagent (Thermo Scientific), as described (23). Aliquots of 500 ng RNA were reverse-transcribed by M-MLV reverse transcriptase (Invitrogen) with random hexamers, as described (21). Primer sequences used are described in Supplementary Table II. Real-time qPCR was performed as described (24), in presence of 250 nM of specific primers with Kappa SYBR Fast qPCR Master Mix (Kapa Biosystems) on a CFX96 touch real-time PCR Detection System (Bio-Rad). Data were analyzed

using the ΔΔCT method, using the geometric mean of *β-Actin* and *Rpl27* as reference genes

(25).

^{125}I uptake and organification

At postnatal day 8 (P8), mothers and litters were fed with an iodine-free diet for 24 h.

At P9, pups were injected intraperitoneally with 1 µCi ^{125}I (Perkin Elmer) and left for another

24 h before sacrifice and thyroid dissection. Both lobes were collected in 2 mM methimazole

(Sigma) and homogenized with a glass Potter. Total thyroid ^{125}I was measured with an

automatic gamma counter "Wizard^2" (Perkin Elmer) before protein precipitation using 10%

trichloroacetic acid (TCA; Merck). After a single wash in TCA, radioactivity was measured in

the protein pellet. Pourcentage of protein-bound iodide was calculated using the ratio of

precipitated cpm/total thyroid cpm (26).

Statistical analysis

All statistical analyses were determined by Prism software (GraphPad Software, La

Jolla, California, USA). RT-qPCR values were obtained by the ΔΔCT method and are expressed

as boxplots with median, 25th and 75th percentiles, and min-max whiskers. Each graph

represents the results form a minimum of 8 independent thyroid lobes from at least 3

different litters. Nonparametric statistical tests were used: Mann-Whitney for single

comparisons and Kruskal-Wallis followed by Dunn's post-test for more conditions. Differences

were considered statistically significant when $p < 0.05$ (*); ** stands for $p < 0.01$; *** for $p < 0.001$.

Results

Genetic construction and assessment of Vps34 inactivation

Global Vps34 inactivation is lethal at embryonic stage (E8.5) (27). To study the role of

Vps34 *in vivo,* we used a genetic approach to conditionally delete the loxP-flanked exon 21 of

the Vps34 gene, which encodes a critical sequence of the lipid kinase domain of Vps34 (21).

Tthis approach was designed to allow for the expression of a minimally truncated,

catalytically-inactive Vps34 protein. However, upon conditional expression in megakaryocytes,

the level of this truncated Vps34 and its obligatory partner, Vps15, were found to be

decreased by 80-90% in the megakaryocyte lineage (thrombocytes), indicating instability of

this truncated Vps34 protein (28). We will thus hereafter refer to the result of this type of

Vps34 deletion as Vps34 absence or Vps34[cKO].

We crossed homozygous Vps34[fl/fl] mice with Pax8-Cre;Vps34[fl/+] mice, which is expected

to lead to a tissue-specific (thyroid and kidney) conditional excision of exon 21 from both

Vps34 alleles in 25% of the pups (Vps34[cKO]). Vps34[cKO] mice were born at the expected

Mendelian ratio and showed normal development until postnatal day 14 (P14), then stopped

growing and died at around 1 month of age (18). We thus analyzed thyroid Vps34[cKO] at P14.

Of note, eye opening was usually delayed till P10.

We first assessed the extent of Vps34 genetic excision in the thyroid by quantification

of its mRNA using Vps34 exon 21-specific primers. Compared to control littermate pups, we

found a ~70 % reduction of exon 21-containing Vps34 mRNA in cKO thyroid extracts. In

comparison, we found no significant difference in the abundance of Vps34 mRNA containing

exon 22-24 (Fig. 1A). Assuming that Vps34 is equally expressed in all thyroid cell types

(expressing and non-expressing the Cre recombinase), this ~70 % decrease suggests that most

of the thyrocytes in cKO mice had undergone Cre-dependent recombination.

Vps34[cKO] thyroid show signs of hyperstimulation by thyroid stimulating hormone (TSH)

At P14, thyroid glands from Vps34^{cKO} were almost twice heavier as compared to

control glands (2.06 ± 0.13 mg (n=6) *versus* 1.03 ± 0.07 mg (n=8); p= 0.0024, Mann Whitney

test). Histological staining with Periodic-Acid-Schiff (PAS, which in the thyroid reflects

expression of the thyroglobulin glycoprotein) readily revealed striking differences between

thyroid glands of control and Vps34^{cKO} mice (Fig. 1B). Thyroid glands of controls appeared as

an assemblies of round follicular sections of variable diameter, filled with colloid intensely and

homogenously stained by PAS. In Vps34^{cKO} thyroids, follicle shape was often irregular,

especially at the periphery of the gland, and PAS staining was weaker or absent, indicating

empty lumina. Papillary projections were frequently seen in those empty lumina (Fig. 1B,

enlarged box). By confocal immunofluorescence microscopy, the epithelial marker, E-cadherin,

confirmed heterogeneity of Vps34^{cKO} follicles and revealed thicker thyrocytes. DAPI labelling

revealed the presence of several nuclei in a significant fraction of lumina (Fig. 1C). As these

histological features (follicle remodelling with thicker thyrocytes, papillary projections, and

colloid consumption up to exhaustion) are reminiscent to those caused by strong stimulation

of the thyroid gland by TSH, they suggested strong TSH stimulation of Vps34^{cKO} thyroid.

Vps34^{cKO} display defective thyroglobulin iodination and proteolysis

To further analyse the significance of the lack of PAS staining in a fraction of Vps34^{cKO}

follicles, we assessed thyroglobulin protein expression and its associated T$_4$ hormonogenic

peptide by immunofluorescence on adjacent sections. WT sections showed a perfectly

overlapping distribution pattern of thyroglobulin and T$_4$, homogeneously filling round colloidal

spaces (Fig. 2A). In Vps34^{cKO}, thyroglobulin labelling was much more heterogeneous and less

intense throughout the tissue. Remarkably, antibodies recognizing the T$_4$ hormonogenic

peptide often failed to label colloidal spaces even when containing Tg in the adjacent section

(Fig. 2A). Quantification revealed that only a quarter (25.2 ± 11.9 %) of the follicles were labeled for T_4. These observations indicated that Vps34 ablation caused altered epithelial organization, defective Tg production and/or excessive Tg consumption, as well as defective thyroglobulin iodination. This further suggested that Vps34^{cKO} mice suffered from impaired T_4 hormone biogenesis, thus would suffer from hypothyroidism. This observation, compatible with the growth retardation observed after P15, could be explained by mislocalization of one or several basolateral and/or apical actors involved in thyroid hormone synthesis, as reported in Vps34^{cKO} kidney proximal tubular cells (18). Of specific interest, whereas Tg signal was restricted to the follicle lumen in control thyroid, Tg dots were also seen within Vps34^{cKO} thyrocytes. This indicates that colloidal Tg was endocytosed by thyrocytes, but that trafficking to or Tg proteolysis within the lysosomes was impaired, thus suggesting defective lysosomal release of T_3/T_4, if/when hormonogenic peptides were still formed on Tg (Fig. 2B, b-d).

Vps34^{cKO} causes severe non-compensated hypothyroidism

To directly test the hypothesis that Vps34^{cKO} impairs thyroid hormone production, plasma was collected at P14 and analyzed for T_4 and TSH levels. In control pups at that age, values were 6.4 ± 1.5 μg/dL for T_4, and 75 ± 110 mU/L for TSH (Fig. 3). In Vps34^{cKO} mice, we found extremely low T_4 values, often near detection level (<0.25 ± 1.5 μg/dL) and very high plasma levels of TSH (19,300 ± 10,500 mU/L) (Fig. 3). These data indicate severe, non-compensated hypothyroidism.

Vps34 cKO display reduced iodine organification

The absence of T_4-bearing Tg in Tg-containing follicles (Fig. 2A) and the undetectable T_4 plasmatic levels in Vps34^{cKO} mice (Fig. 3) might be caused by defective basolateral iodine uptake, apical transport and/or apical organification. Due to lack or very poor specificity of

antibodies directed against the proteins involved in these processes, we first measured their mRNA expression levels in total thyroids at P14 (Fig. 4). We observed no change in expression of *Nis*, *Tpo* and *Duox2,* a significant increase in *Tshr expression*, and a significant decrease of *Duoxa2* mRNA level. At this stage, we cannot conclude if *Duoxa2* downregulation is a contributing cause for the observed hypothyroidism of Vps34[cKO].

To further test whether basolateral NIS and apical TPO, DUOX and DUOXA were all correctly localized, we functionally assayed their combined activity by injection of ^{125}I in pups at P9, followed by a 24-h iodine deprivation and collection of thyroid lobes at P10. Radioactivity was measured before and after protein precipitation by TCA (Fig. 5). ^{125}I uptake was not statistically different in Vps34[cKO] and the three control genotypes (Fig. 5A), suggesting normal NIS function and thus basolateral localization. In marked contrast, only 8% of thyroid ^{125}I was bound to proteins in Vps34[cKO] compared to approximately 55% in control genotypes at this stage (Fig. 5B). This indicates a major effect of Vps34 on one or several apical proteins involved in iodine organification.

Evaluation of lysosomal function and autophagy in Vps34[cKO] thyroid

Vps34 is involved in endocytic trafficking and autophagy. cKO of Vps34 in postnatal kidney glomeruli podocytes has been reported to cause a strong increase of the overall immunofluorescence signal for the lysosomal membrane marker LAMP-1, indicating promotion of lysosome biogenesis (29,30). We further reported that absence of Vps34 in kidney PTCs causes an increase in the actual size of lysosomes, that were sometimes enlarged, and mislocalized toward the basal pole of the cell (18). Given that lysosomal proteases in thyrocytes are important to excise hormonogenic peptides to release free T_3 and T_4, we investigated LAMP-1-labeled structures in Vps34[cKO] thyrocytes as a proxy for lysosomal

function. In our conditions, immunofluorescence on control thyroid sections only produced a

weak LAMP-1 signal in the E-cadherin-positive epithelial thyrocytes (Fig. 6). On the contrary,

a strong and widespread LAMP-1 signal was observed in all Vps34^{cKO} thyrocytes. As shown in

the enlarged box of Fig. 6, LAMP-1 positive late endosomes/lysosomes were enlarged and

sometimes vacuolated in Vps34^{cKO} thyrocytes as compared to control. This is reminiscent of

thyroid of cystinotic (*Ctns$^{-/-}$*) mice in which thyroid hormone production is also impaired (31).

The absence of Vps34 in kidney also causes a block of macroautophagy. We therefore

assessed the expression of p62 (also called sequestosome-1, or SQSTM1), a polyubiquitin-

binding protein that interacts with LC3 II (microtubule associated protein 1 light chain) on the

autophagosome membrane and is normally continuously degraded by the autophagy process

(32). Since p62 accumulates when completion of autophagy is inhibited, p62 can be used as a

marker to study autophagy flux (31). As shown in Fig. 7, no signal for p62 was detected in

control thyroid section, indicating normal autophagic flux. Conversely, Vps34^{cKO} thyrocytes

showed multiple p62 aggregates, indicating impaired autophagic flux, consistent with

defective trafficking to lysosomes. Further studies are necessary to determine if Vps34^{cKO}

causes abortive autophagy in thyrocytes as previously shown in kidney PTCs (18).

Vps34^{cKO} follicles display luminal cells of unknown nature

Surprisingly, while cells trapped in follicular lumen also showed a strong LAMP-1 signal

(Fig. 6), they were negative for for E-cadherin or TTF-1 expression, a permanently expressed

thyrocyte-specific transcription factor (Suppl Fig. 1A), indicating that these cells were not

derived from the thyrocyte lineage. In addition, lineage-tracing experiments on Pax8-Cre;

Vps34$^{fl/fl}$;ROSA-STOP-YFP pups at P14 revealed YFP positive signal only in follicle-delineating

thyrocytes and not in these luminal cells (Suppl Fig. 1B). Thus, luminal cells had never

expressed Pax8 and were thus not derived from thyrocyte progenitors. Although infiltration

by macrophages was an attractive alternative, we failed to label these cells for the

conventional macrophage marker, F4/80 (Suppl Fig. 1C). At this stage, we cannot conclude

about the origin of the cells trapped in follicular lumen of Vps34^{cKO} mice.

Further characterization of luminal cell turn-over, using immunofluorescence for

markers of proliferation (Ki67) and apoptosis (activated caspase 3), showed many proliferative

events and few apoptotic bodies (Suppl Fig. 1D). However, since proliferating cells are labeled

for Ki67 during most of the cell division time whereas apoptotic bodies are generally very

short-lived, it is not possible to assess the respective importance of proliferation and apoptosis

in this population. Of note, there was no appreciable change of proliferation and apoptotic

markers in the cKO thyrocytes themselves at P14, indicating that cell homeostasis was not

severely perturbed despite Vps34 absence at this stage.

Discussion

In this study we report that deleting Vps34 in thyrocytes by Pax8-Cre-driven

recombination causes several defects in the thyroid: (i) doubling of thyroid weight and

perturbed thyroid parenchyma organization, with reduced PAS$^+$ colloidal spaces; (ii) strong

decrease of ^{125}I organification, at comparable ^{125}I uptake, and of T$_4$ formation on thyroglobulin

(detected by immunofluorescence as early as P3); (iii) severe hypothyroidism with collapsed

plasma T$_4$ levels and very high TSH (documented at P14); (iv) delayed "eye opening" and

impaired postnatal growth after two weeks. All these features phenocopy the impact of

Duoxa2 KO (34).

DUOXA2 is a chaperone protein required for the correct localization of DUOX2 at the

apical pole of thyrocyte, where the complex (DUOX2/DUOXA2) produces hydrogen peroxide.

In the absence of DUOXA2, hydrogen peroxide is not produced and subsequent

thyroperoxidase-mediated oxidation of iodide into reactive compounds and thyroglobulin

iodination are abolished (34). The similarity of the severity of hypothyroidism in Vps34^{cKO} and

Duoxa2 KO mice prompted us pay a closer look at DUOXA2 in Vps34^{cKO}. We found that the

expression of *Duoxa2* started decreasing after P3, reaching two-fold lower values at P14. It is

interesting to note that expression of *Duox2* and *Duoxa2* mRNAs was quantitatively different,

despite of the fact that they share the same promoter (35). Whether decreased *Duoxa2* mRNA

expression is reflected by equivalent two-fold decrease of DUOX2/DUOXA2 complex at the

apical pole is unknown. However, it is very unlikely that a two-fold decrease would by itself so

severely impact on thyroglobulin iodination, since heterozygous mice for *Duoxa2* deletion

have virtually no phenotype. Instead, based on the known functions of Vps34 in kidney PTCs,

where its inactivation causes lack of apical localization of endocytic receptors (megalin,

cubilin) and solute transporters (NaPi-IIa, SGLT-2)(18), it would be extremely interesting to

localize DUOXA2 on thyroid sections, and assess apical addressing in Vps34^{cKO} thyrocytes, but

unfortunately no reliable antibodies are currently available. Nevertheless, ^{125}I uptake and

processing experiments support the hypothesis of defective apical localization of one or

several actors involved in thyroid hormonogenesis. While similar ^{125}I uptake in the thyroid of

control and Vps34^{cKO} at P10 suggests normal localization and function of the sodium-iodine

symporter, NIS, and/or the alternative transporter SLC26A7 (3), the major decrease of its

protein-bound fraction in Vps34^{cKO} are best explained by defective apical iodine transport or

organification.

Characteristic histopathological alterations combined with the very high plasma TSH

levels and two-fold increased *Tshr* expression in Vps34^{cKO} thyroid suggest that thyrocytes are

in a hyperstimulated state. However, the expression of the most sensitive target gene of the

TSH signaling pathway, *Nis*, was surprisingly not changed leading to the hypothesis of TSHR

mistrafficking into intracellular vesicles. Recent work in *Drosophila* revealed that Vps34

inactivation or pharmacological inhibition using the small molecule inhibitor SAR405, caused

alteration of cell polarity and disruption of epithelial architecture by relieving LKB1 inhibition

and triggering JNK activation (13). A role of Vps34 in epithelial organization and polarity was

also observed in 3D cultures of Caco-2 cells (13). It would be interesting to analyze the

activation states of LKB1 and JNK in Vps34^{cKO} thyroid, and, if modified, to cross Vps34^{cKO} with

LKB1 floxed alleles.

Our work also evidenced increased abundance and size of LAMP-1 positive late

endosomes/lysososmes in Vps34^{cKO} thyrocytes, with accumulation of Tg and the autophagic

marker, p62, indicative of a defective lysosomal trafficking and/or function. In addition to

impaired Tg iodination, defective thyroid hormone release might thus also result from

impaired endocytic transport of iodoTg to lysosomes or proteolytic excision of T_3/T_4 therein.

Several studies have reported a role for Vps34 for endocytic trafficking to lysosomes (12, 36,

37) and activation of lysosomal proteases (14). In our study on Vps34^{cKO} kidney PTCs, we also

observed enlargement of lysosomes and their filling by undigested material, labelled for a

variety of antigens (18)*.* In Vps34^{cKO} thyrocytes, we also observed increased abundance and

size of LAMP-1-positive compartments. Based on these and previous findings showing that

Vps34 deletion leads to late endosome/lysosome enlargement and defective lysosomal

function (38), we assume a defective proteolytic excision of T_3/T_4 from iodoTg, analogous to

what we reported in cystinotic *Ctns$^{-/-}$* mice (31). Lysosomal enlargement due to defective

degradation causes translocation of the transcription factor EB (TFEB), normally sequestered

in the cytosol, to the nucleus where it promotes the coordinate expression of genes involved

in lysosome biogenesis and autophagic pathways (39). TFEB translocation would readily

account for increased LAMP-1 signal.

It has recently become clear that Vps34 may act on lysosome positioning, which is

crucial to autophagosome formation (37,40). Similar to the kidney PTC defect, we observed

an increased p62 signal in Vps34^{cKO} thyroids at P14. Accumulation of p62 in Vps34^{cKO} indicates

a block in the autophagy process. It would be interesting to further analyse the routes to the

lysosomes, by analysing the activation state of Rab7, the lipidation of LC3II, and the expression

of LAMP2A, a marker for chaperone-mediated autophagy at lysosomes.

A final observation deserving discussion is the occurrence of luminal living cells.

Defective endocytic and autophagic routes to the lysosomes could impact thyrocyte

homeostasis, thus causing cellular stress. Cellular stress, induced by defective autophagosome

and lysosomal function, or by activation of stress kinase such as JNK, could explain shedding

of thyrocytes cells into the colloidal space. However, the cells that we found in the lumens of

Vps34^{cKO} thyroids were negative for E-cadherin and the thyrocyte-specific transcription factor,

TTF-1. Furthermore, using lineage tracing experiments, we demonstrated that these luminal

cells had never expressed Pax8 and thus did not derive from thyrocytes. Alternatively, we

examined whether luminal cells could derive from mesenchymal cells, in particular infiltrating

macrophages. We know that these cells have high LAMP-1 signal, are proliferating, and must

have invasive properties to cross the follicular monolayer and its surrounding basal lamina,

before reaching the colloid. However, luminal cells were not labelled for the conventional

macrophage marker, F4/80, admittedly not a universal macrophage identifier. The origin of

infiltrating cells and the cause of high LAMP1 are thus still unknown.

Acknowledgements

This work was supported by grants from the Fonds pour la Recherche Scientifique (F.R.S-FNRS,

J.0076.18, Belgium), Université catholique de Louvain (Actions de Recherche concertées to

CEP, ARC 15/20-065), and Fondation Roi Baudouin. SR and X-HL were supported by a grant

DK15070 from the National Institutes of Health, USA. Work in the laboratory of BV was

supported by grants from the MRC (G0700755) and BBSRC (BB/I007806/1; BB/M013278/1).

The core facility for Imaging Cells and Tissues was also financed by National Lottery, Région

bruxelloise, Région wallonne, Université catholique de Louvain and de Duve Institute. The

authors thank Abdelkadder El Kaddouri and Claude Massart for technical help. GG held a

fellowship from the Fonds pour la formation à la Recherche dans l'Industrie et l'Agriculture

(FRIA, Belgium), TW was postdoctoral researcher supported by UCLouvain, VJ is supported by

the Cystinosis Research Foundation, CS is a UCLouvain teaching assistant, HGC was a

postdoctoral researcher and CEP is a Senior Research Associate at F.R.S-FNRS.

Authors disclosures and contributions

BV is a consultant for Karus Therapeutics (Oxford, UK), iOnctura (Geneva, Switzerland) and

Venthera (Palo Alto, US) and has received speaker fees from Gilead. The other authors have

no competing financial interest. GG and TSW performed the morphological and molecular

studies and analyzed the data. OD, VJ, CS and AS participated in data collection and analysis.

HGC was involved in supervision of GG. BB and BV provided the Vps34 mice, XHL and SR

assayed T4 and TSH levels, and CM provided expertise with [125]I experiments. PJC and CEP

conceived, designed and supervised the project, and wrote the manuscript. All authors

have approved the final version of the manuscript.

Name and address of corresponding author

Prof. Christophe E. Pierreux

Cell Biology Unit

de Duve Institute and Université Catholique de Louvain

75, Avenue Hippocrate

B-1200 Brussels

Belgium

Email : christophe.pierreux@uclouvain.be

References

1. Colin IM, Denef JF, Lengele B, Many MC, Gerard AC 2013 Recent insights into the cell biology

of thyroid angiofollicular units. Endocr Rev 34:209-238.

2. Carvalho DP, Dupuy C 2017 Thyroid hormone biosynthesis and release. Mol Cell Endocrinol

458:6-15.

3. Cangul H, Liao X-H, Schoenmakers E, Kero J, Barone S, Srichomkwun P, Iwayama H, Serra

EG, Saglam H, Eren E, Tarim O, Nicholas AK, Zvetkova I, Anderson CA, Karet Frankl FE,

Boelaert K, Ojaniemi M, Jääskeläinen J, Patyra K, Löf C, Williams ED, UK10K Consortium,

Soleimani M, Barrett T, Maher ER, Chatterjee VK, Refetoff S, Schoenmakers N 2018

Homozygous loss-of-function mutations in SLC26A7 cause goitrous congenital

hypothyroidism. J Clin Ivest Insight 3:e99631.

4. Koumarianou P, Gomez-Lopez G, Santisteban P 2017 Pax8 controls thyroid follicular polarity

through cadherin-16. J Cell Sci 130:219-231.

5. Shivas JM, Morrison HA, Bilder D, Skop AR 2010 Polarity and endocytosis: reciprocal

regulation. Trends Cell Biol 20:445-452.

6. Rodriguez-Boulan E, Macara IG 2014 Organization and execution of the epithelial polarity

programme. Nat Rev Mol Cell Biol 15:225-242.

7. Morrison HA, Dionne H, Rusten TE, Brech A, Fisher WW, Pfeiffer BD, Celniker SE, Stenmark

H, Bilder D 2008 Regulation of early endosomal entry by the Drosophila tumor

suppressors Rabenosyn and Vps45. Mol Biol Cell 19:4167-4176.

8. Lindmo K, Stenmark H 2006 Regulation of membrane traffic by phosphoinositide 3-kinases.

J Cell Sci 119:605-614.

9. Juhasz G, Hill JH, Yan Y,Sass M, Baehrecke EH, Backer JM, Neufeld TP 2008 The class III PI(3)K

Vps34 promotes autophagy and endocytosis but not TOR signaling in Drosophila. J Cell

Biol 181:655-666.

10. Backer JM 2008 The regulation and function of Class III PI3Ks: novel roles for Vps34.

Biochem J 410:1-17.

11. Jaber N, Zong WX 2013 Class III PI3K Vps34: essential roles in autophagy, endocytosis, and

heart and liver function. Ann NY Acad Sci 1280:48-51.

12. Bechtel W, Helmstädter M, Balica J, Hartleben B, Kiefer B, Hrnjic F, Schell C, Kretz O, Liu S,

Geist F, Kerjaschki D, Walz G, Huber TB 2013 Vps34 deficiency reveals the importance

of endocytosis for podocyte homeostasis. J Am Soc Nephrol 24:727-743.

13. O'Farrell F, Lobert VH, Sneeggen M, Jain A, Katheder NS, Wenzel EM, Schultz SW, Tan KW,

Brech A, Stenmark H, Rusten TE 2017 Class III phosphatidylinositol-3-OH kinase

controls epithelial integrity through endosomal LKB1 regulation. Nat Cell Biol 19:1412-

1423.

14. Ronan B, Flamand O, Vescovi L, Dureuil C, Durand L, Fassy F, Bachelot M-F, Lamberton A,

Mathieu M, Bertrand T, Marquette J-P, El-Ahmad Y, Filoche-Romme B, Schio L, Garcia-

462   Echeverria C, Goulaouic H, Pasquier B 2014 A highly potent and selective Vps34

463   inhibitor alters vesicle trafficking and autophagy. Nat Chem Biol 10:1013-1019.

464 15. Nascimbeni AC, Codogno P, Morel E 2017 Phosphatidylinositol-3-phosphate in the

465   regulation of autophagy membrane dynamics. FEBS J 284:1267-1278.

466 16. Dupont N, Nascimbeni AC, Morel E, Codogno P 2017 Molecular Mechanisms of

467   Noncanonical Autophagy. Intl Rev Cell Mol Biol 328:1-23.

468 17. Carpentier S, N'Kuli F, Grieco G, Van Der Smissen P, Janssens V, Emonard H, Bilanges B,

469   Vanhaesebroeck B, Gaide Chevronnay HP, Pierreux CE, Tyteca D, Courtoy PJ 2013

470   ClassIII phosphoinositide 3-kinase/VPS34 and dynamin are critical for apical endocytic

471   recycling. Traffic 14:933-948.

472 18. Grieco G, Janssens V, Gaide Chevronnay HP, N'Kuli F, Van Der Smissen P, Wang TS, Shan J,

473   Vainio S, Bilanges B, Jouret F, Vanhaesebroeck B, Pierreux CE, Courtoy PJ 2018

474   Suppression of Vps34 PI 3-kinase activity in kidney proximal tubules abrogates apical

475   trafficking and causes a Fanconi syndrome. Sci Rep 8:14133.

476 19. Shan J, Jokela T, Skovorodkin I, Vainio S 2010 Mapping of the fate of cell lineages generated

477   from cells that express the Wnt4 gene by time-lapse during kidney development.

478   Differentiation 79:57-64.

479 20. Bouchard M, Souabni A, Busslinger M 2004 Tissue-specific expression of cre recombinase

480   from the Pax8 locus. Genesis 38:105-109.

481 21. Valet C, Levade M, Chicanne G, Bilanges B, Cabou C, Viaud J, Gratacap MP, Gaits-Iacovoni

482   F, Vanhaesebroeck B, Payrastre B, Severin S 2017 A dual role for the class III PI3K,

483   Vps34, in platelet production and thrombus growth. Blood 130:2032-2042.

484 22. Pohlenz J, Maqueem A, Cua K, Weiss RE, Van Sande J, Refetoff S 1999 Improved

485   radioimmunoassay for measurement of mouse thyrotropin in serum: strain

differences in thyrotropin concentration and thyrotroph sensitivity to thyroid

hormone. Thyroid 9:1265-1271.

23. Delmarcelle AS, Villacorte M, Hick AC, Pierreux CE 2014 An ex vivo culture system to study

thyroid development. J Vis Exp 6:88.

24. Hick AC, Delmarcelle AS, Bouquet M, Klotz S, Copetti T, Forez C, Van Der Smissen P,

Sonveaux P, Collet JF, Feron O, Courtoy PJ, Pierreux CE 2013 Reciprocal

epithelial:endothelial paracrine interactions during thyroid development govern

follicular organization and C-cells differentiation. Dev Biol 381:227-240.

25. Dupasquier S, Delmarcelle AS, Marbaix E, Cosyns JP, Courtoy PJ, Pierreux CE 2014

Validation of housekeeping gene and impact on normalized gene expression in clear

cell renal cell carcinoma: critical reassessment of YBX3/ZONAB/CSDA expression. BMC

Mol Biol 15:9.

26. Massart C, Hoste C, Virion A, Ruf J, Dumont JE, Van Sande J 2011 Cell biology of H2O2

generation in the thyroid: investigation of the control of dual oxidases (DUOX) activity

in intact ex vivo thyroid tissue and cell lines. Mol Cell Endocrinol 343:32-44.

27. Zhou X, Takatoh J, Wang F 2011 The mammalian class 3 PI3K (PIK3C3) is required for early

embryogenesis and cell proliferation. PloS one 6:e16358.

28. Bilanges B, Alliouchene S, Pearce W, Morelli D, Szabadkai G, Chung YL, Chicanne G, Valet

C, Hill JM, Voshol PJ, Colinson L, Peddie C, Ali K, Ghazaly E, Rajeeve V, Trichas G, Srinivas

S, Chaussade C, Salomon RS, Backer JM, Scudamore CL, Whitehead MA, Keaney EP,

Murphy LO, Semple RK, Payrastre B, Tooze SA, Vanhaesebroeck B 2017 Vps34 PI 3-

kinase inactivation enhances insulin sensitivity through reprogramming of

mitochondrial metabolism. Nat Commun 8:1804.

29. Chen J, Chen MX, Fogo AB, Harris RC, Chen JK 2013 mVps34 deletion in podocytes causes

glomerulosclerosis by disrupting intracellular vesicle trafficking. J Am Soc Nephrol.

24:198-207.

30. Bechtel, W, Helmstädter M, Balica J, Hartleben B, Schell C, Huber TB 2013 The class III

phosphatidylinositol 3-kinase PIK3C3/VPS34 regulates endocytosis and

autophagosome-autolysosome formation in podocytes. Autophagy 9:1097-1099.

31. Gaide Chevronnay HP, Janssens V, Van Der Smissen P, Liao XH, Abid Y, Nevo N, Antignac C,

Refetoff S, Cherqui S, Pierreux CE, Courtoy PJ 2015 A mouse model suggests two

mechanisms for thyroid alterations in infantile cystinosis: decreased thyroglobulin

synthesis due to endoplasmic reticulum stress/unfolded protein response and

impaired lysosomal processing. Endocrinology 156:2349-2364.

32. Cohen-Kaplan V, Ciechanover A, Livneh I 2016 p62 at the crossroad of the ubiquitin-

proteasome system and autophagy. Oncotarget 7:83833-83834.

33. Bjorkoy G, Lamark T, Pankiv S, Overvatn A, Brech A, Johansen T 2009 Monitoring

autophagic degradation of p62/SQSTM1. Methods Enzymol 452:181-197.

34. Grasberger H, De Deken X, Mayo OB, Raad H, Weiss M, Liao XH, Refetoff S 2012 Mice

deficient in dual oxidase maturation factors are severely hypothyroid. Mol Endocrinol

26:481-492.

35. Christophe-Hobertus C, Christophe D 2010 Delimitation and functional characterization of

the bidirectional THOX-DUOXA promoter regions in thyrocytes. Mol Cell Endocrinol

317:161-167.

36. Pasquier B 2015 SAR405, a PIK3C3/Vps34 inhibitor that prevents autophagy and synergizes

with MTOR inhibition in tumor cells. Autophagy 11:725-726.

532 37. Hong Z, Pedersen NM, Wang L, Torgersen ML, Stenmark H, Raiborg C 2017 PtdIns3P

533   controls mTORC1 signaling through lysosomal positioning. J Cell Biol 216:4217-4233.

534 38. Jaber N, Mohd-Naim N, Wang Z, DeLeon JL, Kim S, Zhong H, Sheshadri N, Dou Z, Edinger

535   AL, Du G, Braga VM, Zong WX 2016 Vps34 regulates Rab7 and late endocytic trafficking

536   through recruitment of the GTPase-activating protein Armus. J Cell Sci 129:4424-4435.

537 39. Settembre C, De Cegli R, Mansueto G, Saha PK, Vetrini F, Visvikis O, Huynh T, Carissilo A,

538   Palmer C, Klisch TJ, Wollenberg AC, Di Bernardo D, Chan L, Irazoqui JE, Ballabio A 2013

539   TFEB controls cellular lipid metabolism through a starvation-induced autoregulatory

540   loop. Nat Cell Biol 15:647-658.

541 40. Korolchuk VI, Rubinsztein DC 2011 Regulation of autophagy by lysosomal positioning.

542   Autophagy 7:927-928.

Supplementary Table I: Antibodies and use

Antibody	Supplier	Reference	Species	Dilution	Unmasking	Embedding
E-Cadherin	BD Biosciences	610182	mouse IgG2a	1/1000	- or +	paraffin or gelatin
Thyroglobulin	DAKO	M0781	mouse IgG1	1/500	+	paraffin
T4	MyBiosource	MBS211305	mouse IgG1	1/1000	+	paraffin
LAMP-1	Hybridoma Bank	1D4B	rat (Mab)	1/100	-	paraffin or gelatin
P62	ARP	03-GP62C	guinea pig	1/400	+	paraffin or gelatin
TTF1	Agilent	M357501	mouse IgG1	1/200	+	paraffin
F4/80	Abcam	Ab6640	rat	1/200	+	paraffin
YFP	Abcam	Ab6673	goat	1/250	+	Paraffin or gelatin
Ki67	BD Biosciences	556003	mouse IgG1	1/250	+	paraffin
Act. Casp. 3	Cell signalling	9661	rabbit	1/100	+	paraffin

 Supplementary Table II: Primers

gene	forward primer (5' → 3')	reverse primer (5' → 3')
Vps34 ex20-21	CACACAGTATCCAGCACAGC	TGAGACTGGACCCTATGGCA
Vps34 ex21-22	TTGGAGTTGGAGACCGGCA	TCCTTGTTCAGCTTCATCGGAG
Vps34 ex22-24	GATGGTGGAAGGGATGGGTG	CGCTCTCGTCGATCAGACTC
Cdh1	AGGGAGCTGTCTACCAAAGTG	CCAGTCTCGTTTCTGTCTTC
Nis	AGCAGGCTTAGCTGTATCCC	AGCCCCGTAGTAGAGATAGGAG
Tpo	TGCCAACAGAAGCATGGGCAAC	GCACAAAGTTCCCATTGTCCAC
Tshr	CTGCGGGGCAAAGAGTGTGC	AGGGGAGCTCTGTCAAGGCA
Tg	TGGGACGTGAAAGGGGAATGGTGC	GTGAGCTTTTGGAATGGCAGGCGA
Duox	TCCAGAAGGCGCTGAACAG	GCGACCAAAGTGGGTGATG
Duoxa2	CGTTAACATTACACTCCGAGGAACA	CAGAATGCCACCCACAGTGT
b-actin	TCCTGAGCGCAAGTACTCTGT	CTGATCCACATCTGCTGGAAG
Rpl27	GCCCTGGTGGCTGGAATTGACC	AAACTTGACCTTGGCCTCCCGC

Figure legends

Figure 1. Genetic and histopathological characterization. A. Extensive genetic excision of

Vps34 exon 21. Compared with control (black boxes), Vps34[cKO] thyroid (red boxes) shows ~70%

reduction in exon 21 mRNA level at P14. The unchanged mRNA level spanning exons 22-24

serves as control. Boxes with median and percentiles of 10 WT and 11 cKO samples; ***, p<

0.0001 by Mann-Whitney non-parametric test. *B. Histopathological evidence of follicle*

hyperactivity. As compared to control thyroid tissue where all follicles show lumen filling with

homogenous PAS (Periodic Acid Shiff) staining, Vps34[cKO] thyroid presents fewer, mostly

centrally located, PAS-stained follicles and with weaker staining intensity, other follicles that

appear empty (arrows), and still other with papillary projections into the luminal space

(arrowheads and enlarged box in the inset). All these alterations are characteristic of follicle

hyperactivity. *C. Vps34[cKO] thyrocytes are altered and follicles contains abundant non-*

epithelial cells. Nuclei are labeled by Hoechst (presented in white); thyrocyte basolateral

contours are labeled for E-cadherin (red); two lumen boundaries are delineated by broken

lines. As compared to control thyroid follicles, Vps34[cKO] follicular structures are frequently less

organized, circumscribed by thicker thyrocytes and containing additional cells inside the lumen

(yellow arrowheads). These cells are not labeled for E-cadherin (for further characterization,

see Supplementary Figure 1).

Figure 2. Evidence for defective iodide organification in Vps34[cKO] thyroid: follicular lumina

with Tg but devoid of iodinated-Tg. **A.** Semi-serial thyroid sections from control (left) and

Vps34[cKO] (right) labeled for E-cadherin (green) and thyroglobulin (red), recognized either for a

proteic epitope (Tg), or a hormonogenic peptide (T4). Control follicular lumina are round and

uniformly labeled for both Tg and T4. In Vps34^{cKO}, follicular structures are less regular, with the

majority that are filled of Tg (in this field), are without T4 signal. ***B. Comparison of Tg labelling***

in control (a) and Vps34^{cKO} thyrocytes (b-d). In control follicles (a), Tg is essentially restricted

to lumen, with little or no signal inside thyrocytes. In Vps34^{cKO} (b-d), Tg is often less abundant

in the lumen, but readily detected within thyrocytes as collections of submicrometric dots

(small arrowheads) or larger spheres (large arrowhead). Of note, some luminal cells (c, broken

line) also contain Tg-labeled spheres, indicating active endocytosis (arrow).

Figure 3. Vps34^{cKO} mice display severe hypothyroidism associated with strong TSH

production. As compared to control mice (black circles), T$_4$ blood level in Vps34^{cKO} is extremely

low, if not at the detection level (red circles). Conversely, TSH levels are dramatically elevated.

(10 WT and 9 cKO samples; ***, $p = 0.0003$ by Mann-Whitney non-parametric test).

Figure 4. Relative expression level of principal actors in thyroid hormonogenesis at P14. Gene

expression analysis by RT-qPCR, presented as boxes with median and percentiles. As

compared to control thyroid (black boxes), expression in Vps34^{cKO} (red boxes) is preserved for

the epithelial marker E-cadherin (*Cdh1)*, and the thyroid-specific genes *Nis*, *Tpo*, *Tg* and *Duox2.*

Expression of the TSH receptor (*Tshr*) is significantly increased in Vps34^{cKO}, consistent with a

TSH response, while expression of *Duoxa2* is significantly decreased. Boxes with median and

percentiles of 10 WT and 11 cKO samples; ***, $p= 0.0005$ by Mann-Whitney non-parametric

test.

Figure 5. Normal uptake of iodine but major defects of organification in Vps34^{cKO} thyroids.

A. Comparable ^{125}I-iodine uptake in the thyroid of the three control genotypes (black boxes)

and Vps34^{cKO} thyroid (red box). **B**. Compared to the three control genotypes, the percentage

of ^{125}I bound to protein is much reduced in Vps34^{cKO} thyroid, suggesting defective

organification. Boxes with median and percentiles of 10 Flox/+, 19 Flox/Flox, 10 Cre;Flox/+

and 10 Vps34^{cKO} (Cre;Flox/Flox) samples; ***, p< 0.001 by Kruskal-Wallis test with Dunn's

multiple comparison test.

Figure 6. Vps34^{cKO} thyrocytes and luminal cells display increased LAMP-1 signal (green).

Thyrocytes are labeled by E-cadherin (red). As compared to control follicles which uniformly

show a weak signal for the lysosomal marker LAMP-1, this signal is intense in Vps34^{cKO}

thyrocytes and in cells present in the colloidal space (arrows). Broken lines indicate luminal

contours. As shown in the enlargements below, LAMP-1-labeled structures in Vps34 cKO

thyrocytes are enlarged compared with WT (small arrowheads) or vacuolated (large

arrowheads).

Figure 7. Vps34^{cKO} thyrocytes show strong accumulation of p62 dots outside LAMP-1.

Immunolocalization of LAMP-1 (green), and p62 (red). Compared with control follicles, which

show a weak signal for the LAMP-1 and no signal for p62, both signals are very strong in

Vps34^{cKO} thyroid sections. p62 label aggregates up to 5 m in diameter (large arrowhead in

the enlargement). These aggregates show no signal for LAMP-1, indicating failure to reach

lysosomes.

Suppl Fig. 1. Further characterization of luminal cells in Vps34^{cKO} thyroid.

A. Luminal cells are not thyrocytes. Immunolabelling for thyrocyte transcription factor-1 (TTF-

1, green); nuclei are visualized with the Hoechst stain (represented in white for better

visualization). In both control and cKO thyroid, nuclei of thyrocytes circumscribing the

follicular lumina are all labeled by the thyrocyte-specific transcription-factor-1, TTF-1. As

indicated by the yellow arrowhead, nuclei of luminal cells show no signal for TTF-1 (only white

signal representing Hoechst). ***B. Luminal cells do not derive from thyrocyte progenitors.***

Immunolabelling for E-cadherin (red) and YFP (green); nuclei are visualized with the Hoechst

stain(blue). Representative images of Vps34cKO thyroid sections reveal expression of YFP only

in cells surrounding the follicular lumina, i.e. in cells where Pax8-Cre has been active. Luminal

cells do not derive from thyrocyte progenitors as they are negative for YFP. ***C. Luminal cells***

are not conventional macrophages. Immunolabelling for E-cadherin (red), and for the

conventional macrophage marker (F4/80, green) + Hoechst (white). In control thyroid, F4/80

signal is restricted to the interstitium. In Vps34^{cKO}, luminal cells identified by Hoechst show no

signal for F4/80 (yellow arrowheads). **D. Luminal cells are proliferating.** Immunolabelling for

E-cadherin (red), proliferation (Ki67, shown in white) and apoptosis (activated caspase 3,

green). In developing control thyroid at P14, Ki67 labels mostly round thyrocyte nuclei (arrow)

and elongated interstitial nuclei (arrowhead). In Vps34^{cKO} follicles, luminal cells show evidence

of proliferation (arrows) and apoptosis (arrowhead). Broken lines indicate luminal contours.

GriecoFig1

GriecoFig2

GriecoFig3

GriecoFig4

GriecoFig5

control   Vps34cKO

E-cadherin LAMP-1

GriecoFig6

GriecoFig7