Synthesizing Disparate LiDAR and Satellite Datasets through Deep Learning to Generate Wall-to-Wall Forest Inventories of New England

Elias Ayrey[a], Daniel J. Hayes[a], John B. Kilbride[b], Aaron R. Weiskittel[a]

[a] School of Forest Resources, University of Maine, 5755 Nutting Hall, Orono, Maine, 04469-5755, USA;
[b] College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, 114 Wilkinson Hall, Corvallis, OR 97331, USA

Abstract

Light detection and ranging (LiDAR) has become a common tool for generating remotely sensed forest inventories. However, regional modeling of forest attributes using LiDAR has remained challenging due to varying parameters between LiDAR datasets, such as pulse density. Here we develop a regional model using a three dimensional convolutional neural network (CNN). This is a form of deep learning capable of scanning a LiDAR point cloud as well as coincident satellite data, identifying features useful for predicting forest attributes, and then making a series of predictions. We compare this modeling approach to the standard modeling approach for making forest predictions from LiDAR data, and find that the CNN outperformed the standard approach by a large margin in many cases. We then apply our model to publicly available data over New England, generating maps of fourteen forest attributes at a 10 m resolution over 85 % of the region. Our estimates of attributes that quantified tree size were most successful. In assessing aboveground biomass for example, we achieved a root mean square error of 36 Mg/ha (44 %). Our county-level mapped estimates of biomass were in good agreement with federal estimates. Estimates of attributes quantifying stem density and percent conifer were moderately successful, with a tendency to underestimate of extreme values and banding in low density LiDAR acquisitions. Estimate of attributes quantifying detailed species groupings were less successful. Ultimately, we believe these maps will be useful to forest managers, wildlife ecologists, and climate modelers in the region.

1. Introduction

1.1 Overview

Over the past two decades light detection and ranging (LiDAR) has become a common tool for developing spatially explicit forest inventories (Naesset 1997). Measurements of point cloud datasets derived from LiDAR can be used to predict useful forest inventory attributes such as biomass, stem volume, tree count, and species (Means et al. 2000, Jensen et al. 2006, Lim and Treitz 2004). These inventories are useful for a wide range of applications, including assessing carbon stocks (Patenaude et al. 2004), assisting in precision forestry (Woods et al. 2011), and predicting wildlife habitat (Wulder et al. 2008, García-Feced et al. 2011).

Forest inventories are typically developed using the *area based* approach (White et al. 2013), where the forest is segmented into a series of grid cell-areas, ranging from 10 m to 1 ha in size. First, the LiDAR point cloud and the desired forest attribute (e.g. stem density) within each grid cell are each measured. Next, predictive models are then developed relating the field measurements to the LiDAR measurements. Finally, these models can then be applied to every grid cell across a landscape. The resulting maps are referred to as enhanced forest inventories (EFIs).

While LiDAR data is becoming increasingly available to the public, few studies have emphasized mapping whole regions, focusing instead on specific municipalities or individual parcels. One example of regional LiDAR modeling is Sweden, which recently completed a project to develop nation-wide forest inventory maps at a 12.5 m resolution (Nilsson et al. 2017). Similar maps have also been generated in Finland (Kangas et al. 2018). In Canada, large portions of Alberta's forests have had their vegetative functional groups mapped (Guo et al. 2017), and in

New Brunswick, a province-wide effort has resulted in near wall-to-wall LiDAR inventories (Dick

2019).

This study seeks to develop a regional EFI over the Acadian/New England forests, with a

total of 85 % coverage of the New England states. We overcome several challenges faced by

having disparate LiDAR datasets through the use of deep learning, and we incorporate other

remote sensing products such as spectral data, phenology, and disturbance history.

1.2 The current approach

One common difficulty in generating regional LiDAR inventories is that many regions are

comprised of a patchwork of LiDAR datasets acquired with various specifications, and with forest

analytics as a secondary objective. A standard measure for assessing LiDAR quality is pulse

density, which refers to the number of laser pulses landing within a given area (pls/m^2). Pulse

density can vary both between and within LiDAR acquisitions, and frequently regional LiDAR

collections consist of many acquisitions in which pulse density varies by up to an order of

magnitude.

This is particularly problematic because the standard approach for measuring a LiDAR

point cloud for the development of an EFI model is to take a series of summary statistics describing

point heights and vertical distributions. These include measures of the mean, variance, and vertical

quantiles, as well as proportions of points that fall above certain height thresholds (McGaughey

2009, Silva et al. 2017). Unfortunately, because these statistics are all dependent on the vertical

locations of points, these metrics are subject to change based on different acquisition parameters.

Many of these measurements are also highly correlated, and so there is a risk of model overfitting

without careful model selection (Junttila et al. 2015, Næsset et al. 2005).

Many studies have noted that varying pulse density can adversely affect EFI predictions. For example, in 2008 Gobakken and Næsset noted that the area based approach was strongly effected by pulse density. In 2015 Hansen et al. noted that EFI estimates could be subject to biases if predictions were made on point clouds of different densities than those used to train the model. Even differences in acquisition parameters related to LiDAR sensor type, pulse frequency, and flight altitude can result in different height metrics being generated over the same area of forest (Næsset 2009, Goodwin et al. 2006).

Seasonality also has a major impact on LiDAR height metrics (Næsset 2005, White et al. 2015, Villikka et al. 2012). In deciduous forests, the presence of leaves can result in LiDAR beams being intercepted higher in the canopy (Ørka et al. 2010). Taken together, the literature suggests that models developed using one LiDAR point cloud are often not applicable to another, prohibiting regional LiDAR modelling without unusually consistent LiDAR datasets (White et al. 2013).

1.3 Deep learning

Here we use deep learning to overcome these obstacles by developing a single model for predicting forest attributes that is applicable across many disparate LiDAR and satellite datasets. Deep learning is a form of machine learning and primarily refers to artificial neural networks of a sufficient complexity so as to interpret raw data, without a need for human-derived explanatory variables. These differ from simpler neural networks (such as perceptrons) which make estimates using a set of features. Recently, deep learning has proven itself successful at classifying imagery despite varying contextual information, such as light levels and background subject matter (Krizhevski et al. 2012, LeCun et al. 2015). Deep learning does so by identifying useful spatial features in the LiDAR without the need for human-derived explanatory variables such as height

metrics. These features often identify shapes and gradients in space that may be less subject to

change relative to one another with different acquisition parameters.

We implemented a type of spatial deep learning model called a convolutional neural

network (CNN). A CNN works by passing a series of moving kernels over spatial data. As the

model trains, the weights of those kernels are tuned to identify features useful for predicting the

dependent variables (such as the edges of objects). Deep CNNs stack many of these moving

windows atop one another, allowing the algorithm to quantify complex features. The three-

dimensional CNN proposed here uses a volumetric window to quantify a LiDAR point cloud that

has been binned into voxel-space. These 3D CNNs are thus able to quantify vertical as well as

horizontal features, and shapes such as tree crowns, providing a level of complexity not captured

by height metrics.

Early CNNs were first developed in the late 1990s and were used to classify hand written

digits (Lecun and Bengio 2005). The technique was largely underrepresented until advances in

computing power, technique, open tools popularized them in 2012 (Krizhevsky et al. 2012). Since

then, CNNs of increasing complexity have consistently outperformed models based on feature

extraction for computer vision tasks (Szegedy et al. 2015, Tiagman et al. 2014, He et al. 2016).

More recently, CNNs have been becoming a popular option for remote sensing problems. There

has been much success in using 2D CNNs to classify aerial imagery, hyperspectral, and LiDAR

data (Rizaldy et al. 2018, Ghamisi et al. 2017, Castelluccio et al. 2015). In relation to forestry,

some have used segmentation algorithms to isolate individual trees from LiDAR, then used 2D

CNNs to classify tree species (Guan et al. 2015, Ko et al. 2018). Work is also being done using

2D CNNs to identify individual tree crowns from high resolution imagery (Weinstein et al. 2019,

Li et al. 2016).

Progress has also been made in adapting CNNs to scan LiDAR point clouds and three

dimensional space. Similar CNNs that make use of voxels to quantify point clouds have been used

to identify household objects and malignancies in medical scans (Qi et al. 2016, Maturana and

Scherer 2015). In 2017 Qi et al. introduced PointNet, designed to interpret LiDAR data without

voxels, although this technique does not make full use of the spatial relationship between

neighboring points. In a remote sensing context, Maturana and Scherer (2015) used a 3D CNN to

identify helicopter landing zones in LiDAR. Most similar to this study, Ayrey and Hayes (2018)

tested a variety of CNN architectures to interpret LiDAR for the estimation of forest attributes.

1.4 Objectives

Our ultimate objective is to develop 10 m resolution EFI maps of fourteen forest attributes

over the New England region using LiDAR and satellite data. A secondary objective is to assess

the value of deep learning, and compare it to standard approaches for LiDAR modelling. A final

objective is to compare our mapped estimates of biomass, percent conifer, and tree count to

estimates derived via the design-based U.S. national forest inventory program. We will compare

estimates made at the pixel-level and at the county level using both inventory techniques, and

identify any obvious shortcomings. The end result will be a series of near wall-to-wall mapped

forest inventory estimates of the region, with an accurate assessment of error across space and

forest type. This will provide forest managers, ecologists, and climate modelers in the region with

an unprecedented amount of detailed information about the forest.

2. Materials and Methods

2.1 Forest attributes

Our goal was to develop a complete EFI over our study region, estimating many forest

attributes which may be useful to ecologists, forest managers, and modelers. The complete list is

as follows: aboveground biomass (BIOAG), total biomass (BIOTOT), tree count (TC), basal area

(BA), mean tree height (HT), quadratic mean diameter (QMD), percent conifer stems (PC), percent

spruce or fir stems (PSF), percent *pinus strobus* stems (PWP), inner bark volume (VOL),

merchantable volume (VOLM), inner bark conifer volume (VOLC), inner bark spruce or fir

volume (VOLSF), and inner bark *pinus strobus* volume (VOLWP). For brevity, at points

throughout this manuscript we will highlight only the results of BIOAG, PC, and TC estimation.

All other attributes can also be considered measurements of tree size, density, and species;

represented by these three.

142  *2.2 Training data*

143  For most applications deep learning require very large datasets (Russakovsky et al. 2015).

To meet this requirement we combined thirteen different forest inventories collected over thirty

two sites (Appendix Table A1). Within each inventory all trees greater than 10 cm in diameter

were measured spatially and species and diameter were measured. In several inventories heights

were only measured on only a subset of trees, so species-specific non-linear height-diameter

models were generated using a Chapman-Richards model form to impute tree height using the

existing height measurements at each site. Some inventories were also measured up to ten years

prior to LiDAR acquisition. In instances in which the temporal discrepancy between LiDAR and

field data exceeded two years, trees were projected forward in time using the Forest Vegetation

Simulator's Acadian Variant (Weiskittel et al. 2012).

153  Volume was estimated using regional taper equations (Li et al. 2012). A 10 cm cutoff was

used for merchantable volume. Biomass was estimated using the component ratio method

developed for the US Forest Inventory and Analysis (FIA) program (Woodall et al. 2011).

Each of the stem mapped inventories were aligned visually with the LiDAR to correct for GPS

error in plot location, and then segmented into 10x10 m grid cell plots. We selected this in order

to amplify the amount of unique plots available, while retaining plots large enough to contain

several whole tree crowns. We also strive to make the most spatially explicit models possible.

Nevertheless, these plots are small compared to plot sizes used in other studies, and could be prone

to edge effects.

We accounted for edge effects by using regional diameter to crown width equations to

project each tree's crown in space (Russell and Weiskittel 2011). Tree level basal area, biomass,

and volume allometry were then multiplied by the proportion that each tree's crown overlapped

the plot. Trees were therefore treated as areas containing biomass, rather than points which could

lie one or another side of a plot boundary. This mimics the method by which the remote sensing

instrument measures the trees, as LiDAR or satellite imagery has no means of measuring the

precise location of a tree's stem. Figure 1 demonstrates this correction. Preliminary testing using

a subset of the data indicated that this correction greatly improved model performance.

Figure 1. The percent of crown overlap of each tree in and around the plot was used as a modifier
on that tree's basal area, biomass, and volume. This allowed for the development of models which
more closely reflected what was visible to the remote sensing systems, while remaining unbiased
across multiple cells.

We also augmented the sample size of our training data by allowing out plots to overlap

one another by a maximum of 25 %, and by including multiple LiDAR acquisitions of the same

plot in the training dataset. The precise configuration of LiDAR returns will always vary between

acquisitions. Similar augmentation techniques, such as transforming or rotating input images

multiple times, and using adjacent still frames of videos have been successfully used in deep

learning for many years. Lastly, we included 500 plots with no trees on them to allow for better

predictions in low-vegetation environments. These were sampled randomly across Northern New

England using the National Land Cover Database. The associated LiDAR point clouds were then

manually checked for trees.

Ultimately we assembled 24,606 plots for model training and preliminary evaluation. Of

these, we randomly withheld 4,000 plots, 1,000 for model validation (to determine the optimal

stopping point during deep learning training), and 3,000 for model testing (used for model

comparison and selection). Augmented plots which were related to withheld plots were removed

from the training dataset, thus the final training set was comprised of 17,432 plots.

2.3 Remote Sensing Data

190   2.3.1. LiDAR

We aggregated many public LiDAR datasets from across the region, combining

acquisitions of varying pulse density and seasonality. Although much of the pubic LiDAR in the

US Northeast is flown leaf-off, we chose to develop models capable of functioning in either state

to allow for potential future integration with leaf-on Canadian Maritime data. The training data

ultimately consisted of a 53:47 split between leaf-off and leaf-on respectively.

The majority of the LiDAR used for this study was funded and hosted by the US Geological

Survey's national 3D Elevation Program (3DEP). This data was captured in leaf-off conditions

between 2006 and 2018 at resolutions ranging from 0.5 to 10 pls/m^2. We also incorporated LiDAR

data acquired by NASA Goddard's LiDAR Hyperspectral and Thermal Imager (G-LIHT) as well

as the National Ecological Observatory Network (NEON). These data were acquired in leaf-on

conditions over several of our training sites with pulse densities ranging from 5 to 16 pls/m^2.

Finally, we incorporated several private LiDAR datasets for training, including one over the

Penobscot Experimental Forest site, one over Baxter, and one over Noonan Research Forest in

New Brunswick. Each of these had an average pulse density of 5-6 pls/m^2, the first two were leaf-

off and the second leaf-on. Both pulse density and seasonality were included as model predictors.

206   2.3.2 Satellite variables

We also chose to include satellite derived spectral indices, disturbance metrics, and

phenology data in our models. Each of these are spatially contiguous across our study area, and

have proven useful for predicting forest attributes (Zheng et al. 2004, Pflugmacher et al. 2012,

Clerici et al. 2012). All satellite data processing was done in Google Earth Engine (Gorelick et al.

2017).

Using Sentinel-2b data, we generated maps of six spectral vegetation indices (Normalized

Burn Ratio: NBR, Normalized Difference Vegetation Index: NDVI, Normalized Difference

Moisture Index: NDMI, Red-Edge Chlorophyll Index; RECI, Greenness Index; GI, and Triangular

Chlorophyll Index; TCI). Using Landsat-8 three tassel-cap indices were developed (brightness,

greenness, and wetness). This imagery was acquired between 2015 and 2017, imagery from

between the 150[th] and 244[th] Julian days were used. All images were cloud-masked, then a single

median composite was developed for the study area.

We also incorporated disturbance history using Landsat 5-8 and the LandTrendr

disturbance detection algorithm (Kennedy et al. 2010). LandTrendr works by fitting a maximum

of seven linear segments to the yearly medians of a spectral band within each Landsat pixel.

Vertices identify dramatic changes in the spectral characteristics of that pixel in time, and often

correspond to disturbances. We ran LandTrendr over two tassel cap indices and NBR. Instances

in which LandTrendr identified a vertex in two or more of the three bands, within two years of one

another, were retained as disturbances. We then condensed this data into the year of last

disturbance and the magnitude of that disturbance (as a percent of vegetation index change).

Finally, we estimated growing season length across our study area using Moderate

Resolution Imaging Spectroradiometer (MODIS) data with a resolution of 500 m. This was derived

by subtracting the mean dates of greenness onset and senescence (Xhang et al 2006). This has been

demonstrated to correlate well with site quality, and so may aid models which primarily use height

to infer biomass or tree size.

2.4 Deep learning modelling:

2.4.1 Data preparation

We first prepared the LiDAR data to be scanned by the 3D-CNN by converting it from a

point cloud with each data point representing an X,Y, and Z value, to volumetric pixels (voxels).

A height-normalized point cloud was voxelized by segmenting each 10 x 10 x 35 m space

(representing a grid cell) into 40 x 40 x 105 bins, and then tallying the number of LiDAR points

within each bin. Thus, each voxel represented a space of 25 x 25 x 35 cm on our plot. We used

vertically rectangular voxels to reduce dimensionality and prevent horizontal features from being

lost. Exact voxel size was determined through qualitative testing of several size configurations

using a reduced model form. Ultimately the voxels took the form of a three-dimensional tensor,

over which the kernels of a CNN could be passed.

It has been noted that CNNs frequently perform better and train faster when applied to

standardized data. We tried several standardization techniques (z-score, prescience/absence,

ect…), however we noted no performance increase, possibly owing to the high number of empty

voxels in volumetric space.

247   2.4.2 Model architecture

Our deep learning model architecture was based loosely on Google's Inception-V3

(Szegedy et al. 2016), which was determined by Ayrey and Hayes (2018) to be better suited than

several other popular CNN architectures. The underlying model-form was converted to interpret

3D data, and care was taken to maintain similar proportional dimensionality to the original model

(designed to interpret images with a resolution of 224 x 224). The full model architecture is

visualized in Figure 2.

Figure 2. The full architecture of the Inception-VS convolutional neural network used to predict forest attributes from LiDAR and satellite data.

Inception-V3 consists of a series of preliminary convolution and pooling layers, followed by inception layers. Inception layers consist of a number of convolutions of varying sizes which are passed over the incoming data, each designed to pick up on different features, which are then

concatenated. Inception-V3 consists of nine inception layers back to back, with intermittent

pooling to reduce dimensionality. The final inception layer is fed into a fully connected layer for

a classification or regression prediction. Each convolution was followed by a ReLU activation

layer and batch normalization.

The model was first trained to estimate only BIOAG using LiDAR. We used a process

called transfer learning to initialize the weights of a more complex model using the weights from

the simpler one, which was designed to simultaneously predict all 14 of our forest attributes. Each

forest attribute was standardized using z-scores so that their values fell upon the same range. A

single loss function was then used to optimize the model to predict all forest variables (Equation

1).

[EQUATION 1 HERE]

We included the satellite data as side-channel information by first developing a multi-layer

perceptron to estimate BIOAG directly from the satellite variables. We took the weights from this

model and used them to initialize a subcomponent of the larger model, which produced a 40 x 40

tensor that was then concatenated onto the LiDAR voxels (Zhou et al. 2017).

276   2.4.3 Training

Deep learning models were developed in Python using Google's Tensorflow version 1.4

(Abadi et al. 2016). The model training process took place in three stages (1) A BIOAG model

using only LiDAR, (2) A model predicting all fourteen attributes using only LiDAR, and (3) a

model predicting all fourteen forest attributes using LiDAR and satellite metrics. Transfer learning

was used to preserve the learned weights with each additional level of complexity. Training time

for the first stage was approximately five days using an NVIDIA Tesla k80, the following stages

were trained more rapidly. This lengthy multi-step training process made cross-fold validation

highly impractical.

2.5 Traditional modelling

Deep learning models were compared to models trained using the standard suite of LiDAR

height metrics, derived using the Rlidar package (Silva et al. 2015). This produces a series of

summary statistics of the LiDAR return's heights and proportions above certain height thresholds.

We discarded metrics which made use of LiDAR intensity and return counts, as these could not be

normalized between the different LiDAR acquisitions. We filtered out points lower than 0.5 m

above ground, and used a 2 m threshold for many of the proportional metrics. Other studies in the

region have used similar cutoffs (Hayashi et al. 2014). We also included the aforementioned

satellite-derived metrics, as well as pulse density, and seasonality. In total 41 covariates were

derived from the LiDAR and satellites.

Random forest imputation in regression mode was used to model each of the forest

attributes (Breiman 2001). Other studies conducted on subsets of our dataset have demonstrated

that this modelling technique outperforms linear mixed-effects modelling (Ayrey and Hayes 2018,

Hayashi et al. 2015). We used a process called Variable Selection Using Random Forest to

eliminate unimportant predictors (Genuer et al. 2015). New models were then developed using

2000 decision trees and one-third variable selection at each node-split. These hyper-parameters

were fine-tuned using a subset of the data.

The random forest models were trained and validated using the aforementioned 17,432

training plots and 3,000 test plots. Although accuracy can be assessed using out-of-bag sampling,

we used the same validation scheme as the deep learning models due to data augmentation and

consistency.

2.6 Validation

The training, validation, and testing data derived from the thirteen individual forest inventories are likely not fully representative of the landscape, leading to problems with spatial-autocorrelation at the regional scale. We therefore performed two phases of validation. The first phase of validation made use of the 4,000 withheld plots. This was used for model comparisons between deep learning and traditional modelling, and to settle on the final model form.

The second phase of validation made use of an independent dataset, and was used to assess true model performance across the landscape. For this we used the United States Forest Service's FIA national inventory plots. These consisted of approximately 7,000 stratified-random plots with a nested sampling design within the states of Massachusetts, Vermont, New Hampshire, and Maine. We used these to determine the maps' error and bias, assess spatial-autocorrelation across the landscape, and compare our inventory estimates to FIA county-level estimates. We mapped Connecticut and Rhode Island as well, but chose not to assess error in these state as their forests increasingly represent a Mid-Atlantic forest type not fully represented by the training data. Nevertheless we thought maps of these areas might prove useful in future studies. We removed buildings from our maps using a building mask of the United States developed by Microsoft's Bing Maps Team using high resolution imagery (Bing Maps Team 2018).

The FIA plots consist of a nested plot design that consists of four 7.3 m radius subplots placed 36.6 m away from one another. The subplots have an area of 168 m^2, while the entire FIA plot taken as an aggregate has an area of 672 m^2. The individual subplot measurements were more affected by errors in plot location, as these were more subject to intra-canopy variability. The aggregate plot measurements were less prone to plot location errors, but did not represent the entire range of variability that one would expect in a 10 m grid cell. It is important for validation plots to

have roughly the same area as the grid cells being validated so that each have a similar range in

values. We therefore used the individual subplots to assess 10 m accuracy, and then used the entire

plot as an aggregate to also assess the map's accuracy were it resampled to 20 m. We decided not

to use FIA plots for model training for several reasons: (1) The FIA nested plot design was not

compatible with our edge correction technique. (2) FIA plot locations are imprecise and recorded

with a consumer-grade GPS, and frequently have an error greater than the size of our cells. (3) FIA

plot location data is not publically available and is subject to considerable restrictions, less so if

only used for extracting errors.

3. Results

3.1 Phase one validation and model comparison

The first validation phase made use of withheld plots to compare fourteen random forest

models using height and satellite metrics to two Inception-V3 CNNs. The first CNN made use of

only the LiDAR point cloud, while the second made use of the LiDAR point cloud and satellite

metrics. Results in terms of RMSE and bias are displayed in Table 1. In terms of RMSE, both

CNNs outperformed random forest in predicting 12/14 forest attributes. The two exceptions were

percent conifer and percent spruce fir. In terms of absolute bias, random forest outperformed both

CNNs in 7/14 metrics. It should be noted that in many comparisons of bias, the absolute difference

between models was negligible.

The comparison between CNNs with and without satellite metrics illustrated that the

satellite metrics are contributing to the final model's performance. In terms of RMSE, the CNN

with satellite metrics outperformed the one without 100 % of the time. The CNN without satellite

metrics had less absolute bias in predicting 3/14 metrics. Despite this, the proportional

improvement in terms of both RMSE and bias after the addition of satellite metrics was relatively

small. Additional testing of the random forest model with and without satellite metrics indicated a

greater relative improvement in predictive power.

Table 1. Results in terms of RMSE, RMSE as a percent of mean (%), and bias of three models. Random forest models trained using LiDAR height and satellite metrics, an Inception-V3 model trained using only the LiDAR point cloud, and an Inception-V3 model trained using the LiDAR point cloud and satellite metrics. Highlighted in green are the best values achieved.

	Random Forest with Satellite Metrics		Inception-V3 without Satellite Metrics		Inception-V3 with Satellite Metrics	
	RMSE (%)	Bias	RMSE (%)	Bias	RMSE (%)	Bias
BIOAG (Mg/ha)	48.5 (29.4)	-1.2	34.5 (20.9)	1.3	33.2 (20.1)	-1.5
PC (%)	13.3 --	-0.1	15.7 --	-2.3	14.2 --	-1.7
TC (#)	2.51 (37.2)	-0.01	2.10 (31.1)	-0.6	1.75 (26.1)	-0.03
BIOTOT (Mg/ha)	58.2 (29.1)	-1.4	41.5 (20.8)	1.2	40.0 (18.6)	-2.58
BA (m²)	0.083 (24.5)	-0.001	0.065 (19.3)	-0.004	0.063 (20.0)	-0.009
HT (m)	2.6 (15.2)	0.14	2.1 (12.0)	0.6	1.5 (9.1)	-0.07
QMD (cm)	5.9 (23.9)	0.2	4.3 (17.2)	0.2	3.8 (15.3)	0.1
PSF (%)	8.3 --	0.1	10.8 --	-3.6	9.5 --	-0.2
PWP (%)	8.6 --	-0.1	6.7 --	0.1	6.6 --	-0.3
VOL (m³)	0.81 (27.5)	-0.019	0.580 (19.7)	-0.013	0.558 (18.9)	-0.022
VOLM (m³)	0.751 (27.8)	-0.016	0.545 (20.1)	-0.012	0.524 (19.3)	-0.033
VOLD (m³)	0.756 (60.6)	-0.002	0.460 (36.9)	-0.023	0.446 (35.8)	0.001
VOLSF (m³)	0.31 (91.4)	0.007	0.227 (67.1)	0.016	0.238 (70.2)	-0.017
VOLWP (m³)	0.483 (94.2)	-0.016	0.388 (75.6)	-0.075	0.383 (74.6)	-0.039

* Forest attribute abbreviations are as follows: BIOAG = Aboveground biomass, PC = Percent conifer, TC = Tree Count, BIOTOT = Total biomass, BA = Basal area, HT = Mean tree height, QMD = Quadratic Mean Diameter, PSF = Percent spruce or fir, PWP = Percent *Pinus strobus*, VOL = Inner bark volume, VOLM = Merchantable volume, VOLD = Deciduous volume, VOLSF = Spruce or fir volume

 3.2 Phase two validation

     In the second phase of validation each of the mapped forest attributes were validated across

 Northern New England using FIA plots. Table 2 displays the results of this validation in terms of

 RMSE, RMSE as a percent of mean (nRMSE), and bias at both the subplot level and plot level.

 Model error at the subplot-level was considerably higher than error at a plot level with each forest

 attribute. This is to be expected given that smaller areas are more likely to contain extreme values,

 and the subplot values are more likely to be affected by GPS inaccuracy. The opposite trend could

 be seen in bias, with 10/14 attributes exhibiting greater bias at the plot-level.

Table 2. Results in terms of RMSE, RMSE as a percent of mean (%), and bias of the Inception-V3 model using FIA plot data for validation. Assessments were made using the FIA subplots (roughly corresponding to 10 m cell validation) and the FIA plots taken as an aggregate (roughly corresponding to 20 m accuracy).

		BIOAG (Mg/ha)	PC (%)	TC (#)	BIOTOT (Mg/ha)	BA (m²)	HT (m)	QMD (cm)	PSF (%)	PWP (%)	VOL (m³)	VOLM (m³)	VOLD (m³)	VOLSF (m³)	VOLWP (m³)
FIA 10 m	RMSE	54.3	26	4.77	65.3	0.120	3.9	7.1	22	15	0.944	0.870	0.613	0.400	0.587
	(%)	79	--	96	79	71	40	47	--	--	81	84	125	224	365
Assessment	Bias	0.2	-1.4	0.61	-0.6	0.003	-1.13	0.0	2.0	0.2	-0.049	-0.071	0.009	0.030	-0.004
FIA 20 m	RMSE	36.0	20.1	2.93	43.2	0.079	2.7	4.5	17.9	12.3	0.625	0.578	0.436	0.330	0.454
	(%)	44	--	56	43	41	24	27	--	--	46	48	61	155	235
Assessment	Bias	2.0	-2.8	0.57	1.57	-0.001	-1.06	0.5	2.4	-0.3	-0.059	-0.076	-0.041	0.044	-0.034

* Forest attribute abbreviations are as follows: BIOAG = Aboveground biomass, PC = Percent conifer, TC = Tree Count, BIOTOT = Total biomass, BA = Basal area, HT = Mean tree height, QMD = Quadratic Mean Diameter, PSF = Percent spruce or fir, PWP = Percent *Pinus strobus*, VOL = Inner bark volume, VOLM = Merchantable volume, VOLD = Deciduous volume, VOLSF = Spruce or fir volume, VOLWP = *Pinus strobus* volume

Results of the second phase of validation indicated that two phases were in fact necessary.

Plot level RMSE was worse than the RMSE obtained from the first phase of validation in 13/14

forest attributes, indicating that the withheld plots likely did not represent landscape heterogeneity.

In some forest attributes this difference was relatively minor. Performance of the tree count, mean

height, and species estimates were notably worse in the second phase of validation. The error of

each of these values increased between 41-90 % from that observed in the first phase of validation.

Overall, the error and biases of attributes representing tree size were considerably lower

than those representing species or stem density. Aboveground biomass, total biomass, basal area,

mean tree height, QMD, and volume all had a nRMSE less than 50 %. In contrast, tree count had

a nRMSE of 56 %. Model performance was worst in volume estimates of species groups. Estimates

of the deciduous volume had a relatively high nRMSE, 61 %. Estimates of spruce/fir, and *pinus*

strobus volume both had nRMSEs above 150 %, and could generally be considered not useful. We

did not assess nRMSE of the percent species group estimates. The RMSE and biases of percent

spruce/fir and percent *pinus strobus* were lower than that of percent conifer, but this is likely

because their average values are smaller. Qualitatively, the maps of percent conifer appeared

better, with the others suffering from more banding and local biases.

We also assessed model performance of BIOAG, PC, and TC spatially and by plotting their

predicted verses observed values. Figure 2 illustrates the residuals of each of the three forest

attributes. We note that the BIOAG residuals appear to be fairly evenly distributed across the

landscape, with consistent model biases not immediately apparent. Positive PC residuals appear to

be clustered mostly in Eastern Maine, where greater number of conifers are likely to be found,

indicating that the model is underestimating in areas of proportionally higher conifers. Likewise,

negative residuals in Vermont are an indication that the model is underestimating in areas with

proportionally fewer conifers. Tree count followed a similar trend, greater residuals were

encountered in more northern areas, which correspond to both greater stem density and more

industrial   forests.

Figure 2. Residuals are plotted for three of mapped forest attributes. Red areas denote model underestimation,
blue areas denote model overestimation.

These trends can also be observed in the predicted versus observed plots (Figure 3).

Biomass residuals fall relatively tightly along the 1:1 line. Little to no attenuation is observed at

higher biomass values. Percent conifer residuals seemed to indicate a tendency to overestimate in

low conifer environments, and underestimate in high conifer environments. Finally, tree count

residuals appeared to follow the 1:1 line in low-medium density conditions, but often severely

underestimated   tree   count   in   high-density   conditions.

Figure 3. Predicted versus observed plots using FIA plot-level validation.

3.3 County-level comparisons

With the region nearly completely mapped, we compared country level estimates derived

by summing the values of our map with FIA county-level design-based estimates. This data is

summarized in Table 3. We chose the 38 counties in Northern New England with complete LiDAR

coverage. Initially we used all FIA plots within a county measured within two years of the LiDAR

acquisition. We discovered however, that a large number of FIA plots without trees on them were

located in suburban environments with trees. This resulted in dramatic underestimates by the FIA

data of each forest attribute, so plots that were denoted as having no trees that fell within semi-

forested suburbs were removed.

Examining aboveground biomass, our predictions fell within the 95 % confidence interval

of the FIA's in 31/38 counties, and within 97.5 % confidence interval in 33/38 counties. Across

the landscape of these counties the FIA estimated 4 % more biomass then our maps. This is to be

expected given that our maps frequently had gaps between LiDAR acquisitions and occasionally

had missing LiDAR tiles. The FIA's lack of urban tree sampling likely also played a large role.

Eight of the counties were classified by the US Census Bureau as having an urban population

greater than 50 %. In these urbanized counties FIA estimates were an average of 13 % lower than

our own when including empty plots in suburban areas, and 11 % greater than our own after they

were removed.

In estimating percent conifer, 25/38 of our estimates fell within the 95 % confidence

interval of the FIA's estimate. In agreement with the map of residuals, the percentage of conifers

was significantly underestimated in 6/11 counties in Maine, and significantly overestimated in 4/8

counties in Vermont. In 16/19 counties in Massachusetts and New Hampshire, our estimate of

percent conifer was within the 95 % confidence interval of the FIA's. Across the entire landscape

the two estimates were within 0.2 % of one another.

County level tree count estimates fell within the FIA's 95 % confidence interval 30/38

times. Once again, counties in Maine with greater numbers of small trees were more likely to be

underestimated. In 5/12 Maine counties mapped estimates of tree count were significantly lower

than the FIA's estimates. Outside of Maine 24/27 counties had mapped estimates in agreement

with FIA estimates. This is likely owing to an overall greater stem density in Maine due to more

intensive forestry, and a general proclivity of boreal forests to be more tightly packed.

Table 3. County-level estimates of total aboveground biomass (BIOAG), percent conifer (PC), and tree count (TC) are compared using the FIA's design-based sampling and summations of our forest inventory maps. Blue denotes mapped estimates that fell within the 95 % confidence interval of the FIA's estimate, yellow denotes estimates that fell within 97.5 %, red denotes values that estimates that differed from the FIA.

State	County	FIA BIOAG (Petagrams)	EFI BIOAG (Petagrams)	FIA PC (%)	EFI PC (%)	FIA TC (Millions)	EFI TC (Millions)
Maine	Cumberland	21.47 ± 3.48	18.75	34.45 ± 5.70	26.71	106.57 ± 17.47	89.79
Maine	Hancock	29.25 ± 3.54	29.57	55.63 ± 5.09	43.25	277.55 ± 32.08	244.19
Maine	Kennebec	17.29 ± 2.95	16.72	30.04 ± 5.40	28.86	112.41 ± 19.54	92.92
Maine	Knox	7.52 ± 2.08	5.48	42.67 ± 9.24	30.68	52.47 ± 13.05	35.11
Maine	Lincoln	9.06 ± 2.07	8.31	35.02 ± 8.28	32.43	55.02 ± 14.33	48.59
Maine	Penobscot	56.87 ± 4.64	56.21	51.11 ± 3.58	43.26	585.06 ± 41.93	452.14
Maine	Piscataquis	64.96 ± 5.18	61.13	50.86 ± 3.49	46.27	654.97 ± 49.96	527.91
Maine	Sagadahoc	6.70 ± 1.51	5.18	39.53 ± 12.68	33.46	39.98 ± 9.97	29.71
Maine	Waldo	15.97 ± 2.32	11.29	38.70 ± 5.44	33.92	124.14 ± 18.50	71.14
Maine	Washington	42.80 ± 3.86	43.11	50.94 ± 3.74	46.53	422.44 ± 32.49	404.26
Maine	York	25.92 ± 3.32	22.34	30.78 ± 5.26	27.52	139.91 ± 16.95	105.06
Massachusetts	Barnstable	4.49 ± 1.54	4.43	36.53 ± 13.00	23.96	36.25 ± 12.78	41.28
Massachusetts	Berkshire	34.42 ± 4.43	32.59	18.73 ± 5.05	28.67	114.48 ± 15.41	126.32
Massachusetts	Bristol	11.92 ± 2.98	11.89	15.52 ± 5.34	23.90	52.97 ± 15.64	56.44
Massachusetts	Dukes	0.89 ± 0.54	0.85	20.49 ± 20.15	18.87	6.29 ± 3.02	8.73
Massachusetts	Essex	13.98 ± 2.10	11.15	23.61 ± 5.15	18.41	57.68 ± 12.13	39.25
Massachusetts	Franklin	23.98 ± 3.75	27.30	31.35 ± 5.86	29.99	86.87 ± 14.18	93.13
Massachusetts	Middlesex	20.76 ± 4.31	17.68	23.50 ± 8.09	21.57	60.96 ± 15.27	65.82
Massachusetts	Nantucket	0.34 ± 1.19	0.06	0.00	6.99	2.49 ± 8.75	0.75
Massachusetts	Norfolk	9.66 ± 3.07	7.91	14.92 ± 8.76	21.77	36.71 ± 11.71	33.72
Massachusetts	Plymouth	13.73 ± 3.66	14.43	33.96 ± 8.64	27.24	58.78 ± 14.02	67.32
Massachusetts	Suffolk	NO PLOTS	0.33	NO PLOTS	10.73	NO PLOTS	2.07
Massachusetts	Worcester	44.63 ± 5.77	41.17	22.91 ± 4.75	24.02	144.91 ± 18.86	154.38
New Hampshire	Belknap	10.47 ± 2.42	11.06	29.18 ± 8.56	25.45	50.78 ± 11.55	45.40
New Hampshire	Cheshire	26.48 ± 3.24	25.58	29.09 ± 5.57	31.83	105.08 ± 14.76	98.29
New Hampshire	Hillsborough	25.83 ± 4.00	23.54	28.78 ± 5.45	27.27	102.79 ± 15.68	92.32
New Hampshire	Merrimack	32.48 ± 4.44	24.62	31.80 ± 4.83	27.67	134.76 ± 17.09	99.81
New Hampshire	Rockingham	19.07 ± 3.92	17.89	32.19 ± 6.27	22.27	72.74 ± 14.78	72.68
New Hampshire	Strafford	9.39 ± 2.43	9.76	25.51 ± 8.13	27.51	38.43 ± 10.20	42.43
New Hampshire	Sullivan	16.11 ± 2.58	17.01	30.85 ± 7.20	33.55	74.51 ± 12.52	72.30
Vermont	Caledonia	15.18 ± 3.12	14.79	35.60 ± 8.20	33.14	85.38 ± 16.62	75.95
Vermont	Essex	10.58 ± 1.59	13.95	23.61 ± 5.15	39.51	76.14 ± 9.19	92.98
Vermont	Lamoille	13.73 ± 2.14	12.37	23.19 ± 8.16	35.57	60.14 ± 9.54	62.42
Vermont	Orange	18.18 ± 3.16	19.18	32.09 ± 8.22	30.81	78.32 ± 14.20	79.24
Vermont	Rutland	30.77 ± 3.00	25.65	19.01 ± 4.26	26.68	116.47 ± 11.76	110.83
Vermont	Washington	18.11 ± 2.92	19.55	29.39 ± 6.60	33.57	85.94 ± 13.20	90.94
Vermont	Windham	28.02 ± 3.38	27.60	26.10 ± 6.23	33.09	102.42 ± 12.16	103.68
Vermont	Windsor	30.81 ± 4.33	30.86	19.23 ± 5.52	29.96	112.53 ± 16.81	116.98

4. Discussion

Our results indicate that 3D convolutional neural networks can be used to estimate forest attributes from disparate LiDAR and satellite data. These models outperformed random forest models which use the standard approach for generating forest inventories from LiDAR. They could

also be effectively scaled to make regional high resolution maps/estimates which were often

statistically equivalent to traditional forest inventories.

4.1 Model comparison

Our first objective was to compare LiDAR derived inventory estimates made using CNNs

to estimates made using height metrics and random forest. We assessed this in our first phase of

validation, in which several models were developed from training data and assessed using withheld

plots. Random forest models trained using traditional height metrics and satellite data nearly

always had considerably greater error than the two CNNs we trained. This mirrors the results found

by Ayrey and Hayes (2018), in which 3D CNNs of varying architectures often outperformed linear

and random forest models. These results indicate that deep learning can be a more effective way

of modelling LiDAR than the standard approach of height metrics.

Random forest did do a better job in estimating percent conifer and percent spruce/fir.

These species breakdown estimates were more likely to rely on satellite spectral data than LiDAR

structural data, and so we speculate that random forest did a better job of making use of the satellite

covariates than our CNN. The CNN was trained to scan LiDAR voxels, the addition of satellite

covariates was made after this training, and in such a ways as to concatenate satellite data onto

voxel space. This may have been less than ideal. Zhou and Hauser (2017) outlined several methods

for including side-channel data into a CNN, and we tried each of them with mixed results before

settling on our concatenation method.

The CNNs did not outperform the random forest models in terms of bias. Half of the

random forest models had a lower absolute bias than the CNNs, indicating that both types of

models did approximately as well as one another. We did not observe any notable trends in biases

by forest attribute. Overall, we believe that the differences in absolute bias between models was

often low enough to have been owing to the random variations of the testing data.

The comparison between the CNNs with and without satellite metrics highlighted that the

CNN did benefit from spectral and disturbance information. Error decreased when estimating

every forest attribute, and bias nearly always decreased as well. In terms of magnitude this

improvement was often small, and our results indicate that even without the satellite metrics, a

CNN could be trained which outperforms random forest with height and satellite metrics. We

chose not to explore which satellite metrics were most useful, as deep learning models of this size

run no risk of overfitting with extraneous inputs. However, such an analysis would be possible

through a process similar to random forest's derived importance, and might be useful in narrowing

down necessary remote sensing datasets.

We noted that there were numerous advantages to working with a single deep learning

model aside from better performance. Our Inception-V3 model took a considerable amount of data

and time to train, however, once trained the model could be quickly be applied to large regional

LiDAR datasets. A single model predicting all 14 forest attributes presented less of a data-

management challenge than 14 separate models. We also hypothesize that our CNN would be less

likely to produce estimates in conflict with one another than 14 separate unconstrained models

(e.g. more merchantable volume than total volume). Finally, there is a considerable amount of

precedence in the field of deep learning for making use of pre-trained model weights to solve novel

problems (Pan and Yang 2010, Shin et al. 2016,). The rapid retraining of our CNNs indicates that

this model can easily be fine-tuned with local data, retuned with non-local data, or applied to

different problems to save modelers the effort of training a large CNN to interpret voxel space

with randomized weights. The weights from our CNN could be used to initialize CNNs with other

LiDAR-related objectives, such as individual tree crown segmentation or LiDAR classification.

4.2 Assessing performance

With the model form decided upon, we mapped all fourteen forest attributes across the

study area (Figure 4). We assessed the maps' performance with our second phase of validation,

which made use of independent FIA plots. We assessed accuracy at a 10 and 20 m level, as well

as at the county level.

Figure 4. Regional 10 m forest inventory maps developed over New England. (A) Aboveground biomass. (B) Percent conifer. (C) Tree Count.

We first used the individual FIA subplots to assess 10 m pixel-level error. These error

values were consistently quite high. Better performing estimates included BIOAG, BIOTOT, BA,

HT, QMD, VOL, and VOLM. Normalized RMSE values for many of these attributes ranged from

70 - 80 %. Worse performing attributes had nRMSE values that exceeded 100 %. The FIA plot

locations in this region are subject to considerable error, and the FIA notes that plot location error

can be up to 100 m (although in practice most plots are located within 12 m of their measured

location, Hoppus and Lister 2007). Examining pixel-level accuracy using these subplots was

problematic given the high degree of intra-canopy variation detected by 10 m pixels.

We next assessed plot-level error by taking the mean values of each of the subplots and

their extracted pixel values. The error obtained by this method is comparable to the error of a 20

m aggregate of our maps, as the aggregated subplots have roughly the same area. These error

values were much more favorable. Once again BIOAG, BIOTOT, BA, VOL, and VOLM had the

lowest error values, in the range of 40 - 50 % nRMSE. Mean tree height and QMD had lower

nRMSE, though one might expect this given their underlying distributions. The TC, and VOLD

estimates performed moderately well, with nRMSEs of 56 and 61 %. The PC estimate also

performed moderately well, with a RMSE of 20.1 %.

The forest attributes of more detailed species-groups were estimated poorly. These

included PSF and PWP, which had RMSEs of 17.9 and 12.3 % respectively. These error values

may seem low, but must be placed into context with the distribution and abundance of these species

on the landscape. Likewise, we considered VOLSF and VOLWP to also be poorly estimated, with

nRMSEs exceeding 100 %.

Our model's performance can be summarized as having done a good job at predicting

attributes closely related to tree size, having done a moderate job of predicting attributes related to

tree density and percent conifer, and having done a poor job of predicting attributes related to

species groupings. Other studies modeling forest attribute using LiDAR have likewise had more

difficulty in estimating stem density and species (Treitz et al. 2012, Hayashi et al. 2014, Shang et

al. 2017, Hudak et al. 2008). We can think of several reasons for why our models have also

underperformed here. (1) Although we incorporated satellite spectral indices useful for species

estimation, we believe that the model may have not made full use of them. (2) Stem density was

often underestimated in high density stands, but qualitatively the maps seemed to suffer from

banding in areas with low pulse density LiDAR (<3 pls/m^2). It is possible that the model was

making use of horizontal structural features in the canopy which could not be resolved in low

density LIDAR. Ayrey and Hayes (2018) determined that 3D CNNs do make use of horizontal

canopy structure (such as the edges of tree crowns). (3) Finally, in re-examining our loss function,

we find that 7/14 of our forest attributes were in some way related to tree size, while only one

attribute estimated stem density. It is possible that our unweighted loss function inadvertently

favored attributes estimating tree size, resulting in a model that identified features in the LiDAR

that were more predictive of sizes.

4.3 Performance in space and at the county-level

We assessed model performance in space as well as at the county level using aboveground

biomass, percent conifer, and tree count. Our map of biomass residuals across Northern New

England appeared to be relatively uniform. We noted no major trends in residuals in space,

indicating that we are representing biomass across the landscape well. This may be a good

indication that we have not reached the spatial extent of the model's tree-size estimation capability.

The predicted versus observed plot mirrored this result, with the residuals falling near the 1:1 line.

Notably, we did not experience any saturation of high biomass areas as is often the case with

regional remote sensing studies. Our biomass estimates fell within the 97.5 % confidence of FIA

biomass estimates in all but four counties. Those four counties appeared to have little in common

in terms of forest characteristics, nearness in space, or population density.

Overall, we considered our county-level mapped biomass estimates to be in good

agreement with FIA estimates. We also noted that in urban and urbanized counties, FIA estimates

of biomass could be made to underestimate or overestimate our own, depending on sampling

design. Removing supposedly empty FIA plots placed in suburban areas in which trees were

present in aerial imagery resulted in the FIA overestimating biomass relative to our maps.

Retaining these plots resulted in the FIA underestimating biomass relative to our own. This

suggests that our maps are doing a better job of quantifying urban and suburban biomass. By our

maps, this adds up to an additional 13 % biomass in urbanized counties. However, we should stress

that none of our training data made use of urban plots, and few of our training plots had trees

grown in the open. This improvement may only be the case of some estimate being better than

none at all.

The map of percent conifer residuals showed a systematic underestimation of percent

conifer in Eastern Maine, and overestimation in Vermont. These areas are inhabited by very high

and low proportions of conifers respectively. Examining the predicted versus observed plot, one

can see that percent conifer is often underestimated in highly coniferous plots, and overestimated

in highly deciduous plots. This trend can also be seen in the county-level comparisons of percent

conifer. Thirty four percent of counties in the region had percent conifer estimates outside the

FIA's 97.5 % confidence interval. Counties-level estimates in Maine frequently underestimated

percent conifer, while several counties in Vermont overestimated it. Counties in Massachussetts,

New Hamphire, and Southern Maine tended to have estimates similar to the FIA's. Expanding

these estimates beyond this geographic range would likely continue to result in a curtailing of the

extremes.

The map of tree count residuals was similar to the percent conifer in that there was a

consistent underestimation of trees in Northern Maine, where stem densities are naturally higher

due to the species assemblages and greater industrial disturbance results in younger forests. Unlike

percent conifer, no consistent underestimation of tree count was observed in the older, undisturbed

deciduous forests in Southern New England. The county-level estimates parallel this finding, with

the majority of the disagreement between the FIA tree count estimates being in Maine. The

predicted versus observed plot highlights a trend of saturation in very high density forests.

Intuitively, one might expect this result given that the structure of very dense forest stands resemble

one another despite dramatically different stem densities. At the extreme high end of stem density,

a point cloud representing a stand with 2000 trees per hectare looks very similar to a point cloud

representing 2500 trees per hectare. Satellite indices are also sometimes prone to the same type of

saturation at very high stem densities (Mohammadi et al. 2010). Outside the geographic range of

New England, tree count estimates may remain accurate in low density areas, but be

underestimated in high density areas (characterized by more boreal or industrial forests).

4.4 Mapping errors

Unfortunately, the regional maps of forest attributes suffered from several types of errors.

One source of these errors were the LiDAR acquisitions themselves, which often didn't entirely

overlap, or were collected improperly. Missing areas can often be observed between the gaps of

the nearly 40 different LiDAR acquisitions over the region. In central Connecticut and Eastern

Maine portions of the LiDAR were acquired with improper settings, resulting in vegetation being

severely under-represented.

Banding errors occurred with forest attributes that had moderate/worse performance (TC,

PC, VOLD, PSF, PWP, VOLSF, and VOLWP). These bands were more likely to occur in areas in

which pulse density fell below 3 pls/m^2, and followed scan angle trends. In environments of low

pulse density and high scan angle, these attributes were frequently underestimated, possibly owing

to less horizontal structure being captured by the LiDAR. Maps estimating tree size, such as

BIOAG, VOL, and BA, had few banding errors.

4.5 Our results in context

Several other studies have mapped aboveground biomass in this region and can be used to

place our model's performance in context. In one early example, Kellndorfer et al. used Landsat

and radar to map biomass across the Continental United States (CONUS), and achieved RMSE

values ranging from 42 - 48 Mg/ha over New England with 30 m pixels (Cartus et al. 2012). In

2008 Blackard et al. used MODIS to predict biomass across the CONUS at a 250 m resolution and

achieved an average absolute error in New England ranging between 49.7 – 60.4 Mg/ha. In a more

regional study, Cartus et al. (2012) mapped aboveground biomass in the Northeastern United

States using radar, and achieved RMSE estimates of between 46 – 47.3 Mg/ha with 150 m pixels,

but noted that increasing pixel size dramatically reduced error. A more recent study mapped

biomass in New England and Atlantic Canada using Landsat time-series data, this achieved a

RMSE of 49.6 Mg/ha using 30 m pixels (Kilbride 2018). In the context of these studies, our

aboveground biomass error of 36 Mg/ha at a 20 m resolution represents a considerable

improvement over existing remotely-sensed regional estimates.

Localized studies in experimental forests throughout the region can also be used for

comparison. In a 2016 study Hayashi et al. mapped stem volume using LiDAR and achieved

RMSEs of 46 m^3/ha and 82 m^3/ha in two experimental forests in Maine and New Brunswick (we

achieved an error of 62.5 m^3/ha). In a similar study, Hayashi et al. obtained RMSEs of 4993

trees/ha, 3.68 cm, and 13 m^2/ha for tree count, QMD, and basal area respectively using 20 m cells

at an experimental forest in Maine (Hayashi et al. 2014). Our regional models achieved errors of

293 trees/ha, 4.5 cm, and 7.9 m^2/ha for these attributes; outperforming the local models in

estimating tree count and basal area. Another study at an experimental forest in Massachusetts

used large footprint LiDAR and radar to estimate biomass, achieving a RMSE of 66.6 Mg/ha with

25 m cells (Ahmed et al. 2010). Taken collectively, our regional inventories seem to perform on

par or better than local inventories in the region.

4.6 Conclusions

In this study we mapped the forests of New England at a 10 m resolution, making estimates

of fourteen different forest inventory attributes. We did so through the use of disparate LiDAR

datasets as well as satellite spectral, phenological, and disturbance data. Our method of modeling

these attributes was somewhat novel, and made use of three dimensional convolutional neural

networks which are a form of deep learning. These CNNs outperformed standard modeling

techniques, and proved themselves useful for large-scale mapping, making use of disparate data,

and reducing complexity.

We validated our maps using an external dataset derived from the USFS's forest inventory

and analysis program. We concluded that the most successful estimates were those attributes that

quantified tree size, moderately successful estimates were those that quantified tree density or

percent coniferous, and less successful estimates were those that quantified more specific species

groupings. In particular, we found our biomass estimates to be in good agreement with the FIA's

across the region.

We believe that both the models and the maps generated by this study will be of use to

many other scientists. The weights from the CNN trained here could be used as a starting point to

train models making estimates over different forests, or to other LiDAR-related remote sensing

problems. Likewise, the maps developed here can assist with wildlife mapping, precision forestry,

and carbon stock estimation in the region.

Acknowledgments: We thank NASA Goddard's G-LiHT team and NEON for the use of their LiDAR. We thank Kathryn Miller and the National Park Service for the use of their Acadia National Park field data. We thank Eben Sypitowski and Baxter State Park for the use of their Scientific Forest Management Area CFI field plots. We thank Brian Roth and the University of Maine's Cooperative Forestry Research Unit for the use of their LiDAR and field data, as well as for considerable technical assistance. We thank Keith Kanoti and the University of Maine for the use of their CFI field plots. We thank the New Hampshire Division of Forests and Lands, Caroline A. Fox Research and Demonstration Forest for the use of their CFI field plot data. We thank David Orwig and Harvard Forest for the use of their Megaplot field inventory. We thank Noonan research forest for the use of their stem mapped plots. We thank Laura Kenefic and the U.S. Forest Service's Penobscot Experimental Forest for the use of their LiDAR and field inventory data.

Funding details: This work was supported by the Maine Agricultural and Forest Experiment Station, Publication Number XXXX. This project was supported by the USDA National Institute of Food and Agriculture, McIntire-Stennis project number ME0-XXXX through the Maine Agricultural & Forest Experiment Station.

Conflicts of Interest: The authors declare no conflict of interest.

Data availability statement: Computer code for this work can be found online at https://github.com/Eayrey/3D-Convolutional-Neural-Networks-with-LiDAR. Generated inventory maps can be found online at XXXXXXXXXXXXXXXXX

References

Abadi, M., Barham, P., Chen, J., Chen, Z., Davis, A., Dean, J., Devin, M., Ghemawat, S., Irving, G., & Isard, M. (2016). Tensorflow: a system for large-scale machine learning. In, OSDI (pp. 265-283)

Ahmed, R., Siqueira, P., Bergen, K., Chapman, B., & Hensley, S. (2010). A biomass estimate over the harvard forest using field measurements with radar and lidar data. In, Geoscience and Remote Sensing Symposium (IGARSS), 2010 IEEE International (pp. 4768-4771): IEEE

Ayrey, E., & Hayes, D.J. (2018). The Use of Three-Dimensional Convolutional Neural Networks to Interpret LiDAR for Forest Inventory. Remote Sensing, 10, 649

Blackard, J., Finco, M., Helmer, E., Holden, G., Hoppus, M., Jacobs, D., Lister, A., Moisen, G., Nelson, M., & Riemann, R. (2008). Mapping US forest biomass using nationwide forest inventory data and moderate resolution information. Remote Sensing of Environment, 112, 1658-1677

Breiman, L. (2001). Random forests. Machine Learning, 45, 5-32

Cartus, O., Santoro, M., & Kellndorfer, J. (2012). Mapping forest aboveground biomass in the Northeastern United States with ALOS PALSAR dual-polarization L-band. Remote Sensing of Environment, 124, 466-478

Castelluccio, M., Poggi, G., Sansone, C., & Verdoliva, L. (2015). Land use classification in remote sensing images by convolutional neural networks. arXiv preprint arXiv:1508.00092

Dick, A. (2019). Enhanced Forest Inventory (EFI) Adoption in New Brunswick: Progress to Date and Future Directions. In, CIF E-Lecture Series. Canadian Wood Fibre Centre: Natural Resources Canada

García-Feced, C., Tempel, D.J., & Kelly, M. (2011). LiDAR as a tool to characterize wildlife habitat: California spotted owl nesting habitat as an example. Journal of Forestry, 109, 436-443

Genuer, R., Poggi, J.-M., & Tuleau-Malot, C. (2015). VSURF: an R package for variable selection using random forests. The R Journal, 7, 19-33

Ghamisi, P., Höfle, B., & Zhu, X.X. (2017). Hyperspectral and LiDAR data fusion using extinction profiles and deep convolutional neural network. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 10, 3011-3024

Goodwin, N.R., Coops, N.C., & Culvenor, D.S. (2006). Assessment of forest structure with airborne LiDAR and the effects of platform altitude. Remote Sensing of Environment, 103, 140-152

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. Remote Sensing of Environment, 202, 18-27

Guan, H., Yu, Y., Ji, Z., Li, J., & Zhang, Q. (2015). Deep learning-based tree classification using mobile LiDAR data. Remote Sensing Letters, 6, 864-873

Guo, X., Coops, N.C., Tompalski, P., Nielsen, S.E., Bater, C.W., & Stadt, J.J. (2017). Regional mapping of vegetation structure for biodiversity monitoring using airborne lidar data. Ecological informatics, 38, 50-61

Hayashi, R., Kershaw, J.A., & Weiskittel, A. (2015). Evaluation of alternative methods for using LiDAR to predict aboveground biomass in mixed species and structurally complex forests in northeastern North America. MCFNS, 7, 49-65

Hayashi, R., Weiskittel, A., & Kershaw Jr, J.A. (2016). Influence of prediction cell size on LiDAR-derived area-based estimates of total volume in mixed-species and multicohort forests in northeastern North America. Canadian Journal of Remote Sensing, 42, 473-488

Hayashi, R., Weiskittel, A., & Sader, S. (2014). Assessing the feasibility of low-density LiDAR for stand inventory attribute predictions in complex and managed forests of northern Maine, USA. Forests, 5, 363-383

He, K., Zhang, X., Ren, S., & Sun, J. (2016). Deep residual learning for image recognition. In, Proceedings of the IEEE conference on computer vision and pattern recognition (pp. 770-778)

Hoppus, M., & Lister, A. (2007). The status of accurately locating forest inventory and analysis plots using the Global Positioning System. In, In: McRoberts, Ronald E.; Reams, Gregory A.; Van Deusen, Paul C.; McWilliams, William H., eds. Proceedings of the seventh annual forest inventory and analysis symposium; October 3-6, 2005; Portland, ME. Gen. Tech. Rep. WO-77. Washington, DC: US Department of Agriculture, Forest Service: 179-184.

Hudak, A.T., Crookston, N.L., Evans, J.S., Hall, D.E., & Falkowski, M.J. (2008). Nearest neighbor imputation of species-level, plot-scale forest structure attributes from LiDAR data. Remote Sensing of Environment, 112, 2232-2245

Jensen, J.L.R., Humes, K.S., Conner, T., Williams, C.J., & DeGroot, J. (2006). Estimation of biophysical characteristics for highly variable mixed-conifer stands using small-footprint lidar. Canadian Journal of Forest Research, 36, 1129-1138

Junttila, V., Kauranne, T., Finley, A.O., & Bradford, J.B. (2015). Linear Models for Airborne-Laser-Scanning-Based Operational Forest Inventory With Small Field Sample Size and Highly Correlated LiDAR Data. IEEE Transactions on Geoscience and Remote Sensing, 53, 5600-5612

Kangas, A., Astrup, R., Breidenbach, J., Fridman, J., Gobakken, T., Korhonen, K.T., Maltamo, M., Nilsson, M., Nord-Larsen, T., & Næsset, E. (2018). Remote sensing and forest inventories in Nordic countries–roadmap for the future. Scandinavian Journal of Forest Research, 33, 397-412

Kennedy, R.E., Yang, Z., & Cohen, W.B. (2010). Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr—Temporal segmentation algorithms. Remote Sensing of Environment, 114, 2897-2910

Kilbride, J.B. (2018). Forest Disturbance Detection and Aboveground Biomass Modeling Using Moderate-Resolution, Time-Series Satellite Imagery

Ko, C., Kang, J., & Sohn, G. (2018). Deep Multi-task Learning for Tree Genera Classification. ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci, 153-159

Krizhevsky, A., Sutskever, I., & Hinton, G.E. (2012). Imagenet classification with deep convolutional neural networks. In, Advances in neural information processing systems (pp. 1097-1105)

Krizhevsky, A., Sutskever, I., & Hinton, G.E. (2012). Imagenet classification with deep convolutional neural networks. In, Advances in neural information processing systems (pp. 1097-1105)

LeCun, Y., & Bengio, Y. (1995). Convolutional networks for images, speech, and time series. The handbook of brain theory and neural networks, 3361, 1995

LeCun, Y., Bengio, Y., & Hinton, G. (2015). Deep learning. nature, 521, 436

Li, R., Weiskittel, A., Dick, A.R., Kershaw Jr, J.A., & Seymour, R.S. (2012). Regional stem taper equations for eleven conifer species in the Acadian region of North America: development and assessment. Northern Journal of Applied Forestry, 29, 5-14

Li, W., Fu, H., Yu, L., & Cracknell, A. (2016). Deep learning based oil palm tree detection and counting for high-resolution remote sensing images. Remote Sensing, 9, 22

Lim, K.S., & Treitz, P.M. (2004). Estimation of above ground forest biomass from airborne discrete return laser scanner data using canopy-based quantile estimators. Scandinavian Journal of Forest Research, 19, 558-570

Maturana, D., & Scherer, S. (2015). 3d convolutional neural networks for landing zone detection from lidar. In, Robotics and Automation (ICRA), 2015 IEEE International Conference on (pp. 3471-3478): IEEE

McGaughey, R.J. (2009). FUSION/LDV: Software for LIDAR data analysis and visualization. US Department of Agriculture, Forest Service, Pacific Northwest Research Station: Seattle, WA, USA, 123

Means, J.E., Acker, S.A., Fitt, B.J., Renslow, M., Emerson, L., & Hendrix, C.J. (2000). Predicting forest stand characteristics with airborne scanning lidar. Photogrammetric Engineering and Remote Sensing, 66, 1367-1372

Mohammadi, J., Shataee Joibary, S., Yaghmaee, F., & Mahiny, A. (2010). Modelling forest stand volume and tree density using Landsat ETM+ data. International Journal of Remote Sensing, 31, 2959-2975

Næsset, E. (2005). Assessing sensor effects and effects of leaf-off and leaf-on canopy conditions on biophysical stand properties derived from small-footprint airborne laser data. Remote Sensing of Environment, 98, 356-370

Næsset, E. (2009). Effects of different sensors, flying altitudes, and pulse repetition frequencies on forest canopy metrics and biophysical stand properties derived from small-footprint airborne laser data. Remote Sensing of Environment, 113, 148-159

Naesset, E. (1997). Determination of mean tree height of forest stands using airborne laser scanner data. ISPRS Journal of Photogrammetry and Remote Sensing, 52, 49-56

Nilsson, M., Nordkvist, K., Jonzén, J., Lindgren, N., Axensten, P., Wallerman, J., Egberth, M., Larsson, S., Nilsson, L., & Eriksson, J. (2017). A nationwide forest attribute map of Sweden predicted using airborne laser scanning data and field data from the National Forest Inventory. Remote Sensing of Environment, 194, 447-454

Ørka, H.O., Næsset, E., & Bollandsås, O.M. (2010). Effects of different sensors and leaf-on and leaf-off canopy conditions on echo distributions and individual tree properties derived from airborne laser scanning. Remote Sensing of Environment, 114, 1445-1461

Pan, S.J., & Yang, Q. (2010). A survey on transfer learning. IEEE Transactions on knowledge and data engineering, 22, 1345-1359

Patenaude, G., Hill, R., Milne, R., Gaveau, D., Briggs, B., & Dawson, T. (2004). Quantifying forest above ground carbon content using LiDAR remote sensing. Remote Sensing of Environment, 93, 368-380

Pflugmacher, D., Cohen, W.B., & Kennedy, R.E. (2012). Using Landsat-derived disturbance history (1972–2010) to predict current forest structure. Remote Sensing of Environment, 122, 146-165

Qi, C.R., Su, H., Mo, K., & Guibas, L.J. (2017). Pointnet: Deep learning on point sets for 3d classification and segmentation. Proc. Computer Vision and Pattern Recognition (CVPR), IEEE, 1, 4

Rizaldy, A., Persello, C., Gevaert, C., Oude Elberink, S., & Vosselman, G. (2018). Ground and Multi-Class Classification of Airborne Laser Scanner Point Clouds Using Fully Convolutional Networks. Remote Sensing, 10, 1723

Russakovsky, O., Deng, J., Su, H., Krause, J., Satheesh, S., Ma, S., Huang, Z., Karpathy, A., Khosla, A., & Bernstein, M. (2015). Imagenet large scale visual recognition challenge. International Journal of Computer Vision, 115, 211-252

Russell, M.B., & Weiskittel, A.R. (2011). Maximum and largest crown width equations for 15 tree species in Maine. Northern Journal of Applied Forestry, 28, 84-91

Shang, C., Treitz, P., Caspersen, J., & Jones, T. (2017). Estimating stem diameter distributions in a management context for a tolerant hardwood forest using ALS height and intensity data. Canadian Journal of Remote Sensing, 43, 79-94

Shin, H.-C., Roth, H.R., Gao, M., Lu, L., Xu, Z., Nogues, I., Yao, J., Mollura, D., & Summers, R.M. (2016). Deep convolutional neural networks for computer-aided detection: CNN architectures, dataset characteristics and transfer learning. IEEE transactions on medical imaging, 35, 1285-1298

Silva, C.A., Crookston, N.L., Hudak, A.T., Vierling, L.A., Klauberg, C., & Silva, M.C.A. (2017). Package 'rLiDAR'. In

Szegedy, C., Liu, W., Jia, Y., Sermanet, P., Reed, S., Anguelov, D., Erhan, D., Vanhoucke, V., & Rabinovich, A. (2015). Going deeper with convolutions. In, Proceedings of the IEEE conference on computer vision and pattern recognition (pp. 1-9)

Szegedy, C., Vanhoucke, V., Ioffe, S., Shlens, J., & Wojna, Z. (2016). Rethinking the inception architecture for computer vision. In, Proceedings of the IEEE conference on computer vision and pattern recognition (pp. 2818-2826)

Taigman, Y., Yang, M., Ranzato, M.A., & Wolf, L. (2014). Deepface: Closing the gap to human-level performance in face verification. In, Proceedings of the IEEE conference on computer vision and pattern recognition (pp. 1701-1708)

Team, B.M. (2018). Microsoft Releases 125 million Building Footprints in the US as Open Data. In, Bing blogs: Microsoft

Treitz, P., Lim, K., Woods, M., Pitt, D., Nesbitt, D., & Etheridge, D. (2012). LiDAR sampling density for forest resource inventories in Ontario, Canada. Remote Sensing, 4, 830-848

Villikka, M., Packalén, P., & Maltamo, M. (2012). The suitability of leaf-off airborne laser scanning data in an area-based forest inventory of coniferous and deciduous trees. Silva Fennica, 46

Weinstein, B., Marconi, S., Bohlman, S., Zare, A., & White, E. (2019). Individual tree-crown detection in RGB imagery using self-supervised deep learning neural networks. bioRxiv, 532952

Weiskittel, A., Russell, M., Wagner, R., & Seymour, R. (2012). Refinement of the Forest Vegetation Simulator Northeast variant growth and yield model: Phase III. Cooperative Forest Research Unit. Orono, ME: University of Maine, School of Forest Resources, 96-104

White, J.C., Arnett, J.T., Wulder, M.A., Tompalski, P., & Coops, N.C. (2015). Evaluating the impact of leaf-on and leaf-off airborne laser scanning data on the estimation of forest inventory attributes with the area-based approach. Canadian Journal of Forest Research, 45, 1498-1513

White, J.C., Wulder, M.A., Varhola, A., Vastaranta, M., Coops, N.C., Cook, B.D., Pitt, D., & Woods, M. (2013). A best practices guide for generating forest inventory attributes from airborne laser scanning data using an area-based approach. The Forestry Chronicle, 89, 722-723

Woodall, C.W., Heath, L.S., Domke, G.M., & Nichols, M.C. (2011). Methods and equations for estimating aboveground volume, biomass, and carbon for trees in the US forest inventory, 2010

Woods, M., Pitt, D., Penner, M., Lim, K., Nesbitt, D., Etheridge, D., & Treitz, P. (2011). Operational implementation of a LiDAR inventory in Boreal Ontario. The Forestry Chronicle, 87, 512-528

Wulder, M.A., Bater, C.W., Coops, N.C., Hilker, T., & White, J.C. (2008). The role of LiDAR in sustainable forest management. The Forestry Chronicle, 84, 807-826

Yi, D., Zhou, M., Chen, Z., & Gevaert, O. (2016). 3-D convolutional neural networks for glioblastoma segmentation. arXiv preprint arXiv:1611.04534

Zhang, X., Friedl, M.A., & Schaaf, C.B. (2006). Global vegetation phenology from Moderate Resolution Imaging Spectroradiometer (MODIS): Evaluation of global patterns and comparison with in situ measurements. Journal of Geophysical Research: Biogeosciences, 111

Zheng, D., Rademacher, J., Chen, J., Crow, T., Bresee, M., Le Moine, J., & Ryu, S.-R. (2004). Estimating aboveground biomass using Landsat 7 ETM+ data across a managed landscape in northern Wisconsin, USA. Remote Sensing of Environment, 93, 402-411

Zhou, Y., & Hauser, K. (2017). Incorporating side-channel information into convolutional neural networks for robotic tasks. In, Robotics and Automation (ICRA), 2017 IEEE International Conference on (pp. 2177-2183): IEEE

Appendix Table A1. A list of field inventories used for model training and the first phase of validation. Also included, the area those inventories represented, the number of LiDAR field plots, and the number of LiDAR acquisitions. When inventories covered multiple sites, those sites are listed.

Inventory	Sites	Location(s)	Area (km²)	Number of Plots	LiDAR Acquisitions
Acadia National Park	Mount Desert Isle au Haut Schoodic Point	-68.294, 44.339 -68.627, 44.032 -68.065, 44.351	671	128	2
Baxter State Park	Scientific Forest Management Area	-69.000, 46.176	87	882	1
Bartlett Experimental Forest Echidna		-71.286, 44.064	0.1	46	1
Cooperative Forestry Research Unit	Austin Pond Alder Stream Dow Road Golden Road Harlow Road Katahdin Ironworks Lazy Tom Lake Macwahoc Penobscot Experimental Forest Ronco Cove Rump Road Sarah Road Schoolbus Road St. Aurelie Summit Week's Brook Weymouth Point	-69.705, 45.193 -69.798, 45.369 -69.609, 45.996 -68.675, 45.719 -67.842, 45.646 -69.367, 45.489 -69.456, 45.726 -68.286, 45.798 -68.608, 44.844 -69.634, 45.680 -71.018, 45.193 -70.911, 44.817 -70.778, 44.841 -70.161, 46.259 -68.480, 45.096 -68.522, 46.217 -69.308, 45.947	30000	935	3
Carbon Monitoring System		-69.764, 45.589	4645	414	3
University of Maine Forests	Demeritt Forest Penobscot Experimental Forest	-68.678, 44.933 -68.608, 44.844	55	912	3
Fox Forest		-71.911, 43.138	9	581	1
Harvard Forest Megaplot		-72.176, 42.538	0.4	6646	2
Harvard Forest Echidna		-72.182, 42.531	0.1	90	2
Holt Experimental Forest		-69.772, 43.871	0.1	1001	3
Howland Experimental Forest		-68.742, 45.206	2	556	2
Howland Echidna		-68.742, 45.206	0.1	80	2
Noonan Research Forest		-66.439, 45.977	0.1	25	1
Penobscot Experimental Forest		-68.608, 44.844	4	409	3
Null Plots		Regional		500	1

Aboveground Biomass	
	0 Kg
	1000
	2000+ Kg