Regulation of hippocampal memory by mTORC1 in somatostatin interneuron

J. Artinian[1#], A. Jordan[1], A. Khlaifia[1], E. Honoré[1], A. La Fontaine[1], A-S. Racine[1], I.

Laplante[1], J-C. Lacaille[1]*

[1]Department of Neuroscience and GRSNC, University of Montreal, C.P. 6128, Succ.

Centre-ville, Montreal, QC H3C 3J7, Canada.

*Corresponding author: jean-claude.lacaille@umontreal.ca

[#]Present address: University of Bordeaux and INRA, Nutrition and Integrative

Neurobiology, UMR 1286, Bordeaux, France.

Abstract

Translational control of long-term synaptic plasticity via Mechanistic Target Of

Rapamycin Complex 1 (mTORC1) is crucial for hippocampal learning and memory. The

role of mTORC1 is well-characterized in excitatory principal cells but remains largely

unaddressed in inhibitory interneurons. Here we used cell type-specific conditional

knockout strategies to alter mTORC1 function selectively in somatostatin (SOM)

inhibitory interneurons (SOM-INs). We found that up- and down-regulation of SOM-IN

mTORC1 activity bi-directionally regulates contextual fear and spatial memory

consolidation. Moreover, contextual fear learning induced a metabotropic glutamate

receptor type 1 (mGluR1) mediated long-term potentiation (LTP) of excitatory input

synapses onto hippocampal SOM-INs, that was dependent on mTORC1. Finally, the

induction protocol for mTORC1-mediated late-LTP in SOM-INs regulated Schaffer

collateral pathway LTP in pyramidal neurons. Thus, mTORC1 activity in somatostatin interneurons contributes to learning-induced persistent plasticity of their excitatory synaptic inputs and hippocampal memory consolidation, uncovering a role of mTORC1 in inhibitory circuits for memory.

Introduction

Long-term synaptic plasticity is a prime candidate cellular substrate for learning and memory (*1-3*). In the hippocampus, long-term potentiation (LTP) at excitatory synapses of pyramidal cells is crucial for contextual as well as spatial learning and memory (*4*). In analogy to long-term memory consolidation, LTP in pyramidal cells displays an early phase, lasting minutes that is mediated by post-translational changes, and a late phase, lasting hours that requires transcription and translation (*3, 5*). Mechanistic Target of Rapamycin (mTOR) is a serine/threonine kinase that associates with two distinct protein complexes, mTOR Complex 1 and 2 (mTORC1 and 2), to control cell growth, proliferation and migration (*6*). mTORC1, of which Raptor is an essential component, plays a central role in cell growth by regulating protein synthesis and turnover, as well as lipid, nucleotide and glucose metabolism (*6*). In mature neurons, mTORC1 regulates protein synthesis during long-lasting synaptic plasticity and memory (*7, 8*). This role of mTORC1 in hippocampal memory was clearly established in principal cells (*7*).

In the hippocampus, excitatory neurons are regulated by highly heterogeneous inhibitory interneurons that display various morphological and physiological properties, as well as diverse patterns of connectivity and protein expression (*9*). Plasticity of

hippocampal interneurons is also implicated in learning. Hippocampus-dependent learning is associated with structural plasticity of interneuron afferent connectivity (*10*) and modulation of their excitatory inputs (*11*). In addition, excitatory synapses onto hippocampal interneurons demonstrate multiple forms of LTP (*12, 13*). However, whether mTORC1 plays a role in these learning-related changes in inhibitory interneurons remains mostly unknown.

In the hippocampus, somatostatin (SOM) expressing interneurons (SOM-INs) are a subgroup of GABAergic interneurons that, in CA1, receive excitation from local principal pyramidal cells and provide feedback dendritic inhibition (*9*). They include the so-called Oriens-Lacunosum/Moleculare (O-LM) cells, bistratified cells and long-range projecting cells (*9*). SOM-INs regulate pyramidal cell synaptic integration (*14*), control their rate and burst firing (*15*), modulate their synaptic plasticity (*16*) and support contextual fear learning (*17*). Excitatory synapses onto SOM-INs express a form of LTP dependent on type 1a metabotropic glutamate receptors (mGluR1a) (*18, 19*). Additionally, a late form of mGluR1a-dependent LTP is present in O-LM interneurons in slice cultures (*8*) that lasts 24 hours and is dependent on mTORC1-mediated translation (*8, 20*). Thus SOM-INs represent an interesting subpopulation of interneurons to investigate mTORC1 function.

To determine if mTORC1 activity in SOM-IN plays a role in hippocampal memory, we used conditional knock-out mouse strategies to bi-directionally manipulate mTORC1 activity selectively in SOM-INs by targeting an essential component of mTORC1, the Regulatory-Associated Protein of mTOR (Raptor), or a repressor of mTORC1, the Tuberous Sclerosis Complex 1 (TSC1). We found that mTORC1 activity in somatostatin

interneurons contributes to hippocampus-dependent memory formation and to learning-

induced persistent plasticity at somatostatin interneuron input synapses, uncovering an

important function of mTORC1 activity in inhibitory local circuits for memory

consolidation.

Results

We first established that mGluR1a- and mTORC1-dependent late-LTP (*8, 20*)

occurs at excitatory synapses onto CA1 SOM-INs in mice expressing EYFP under the

control of the SOM promoter (*Sst*[ires-Cre];*Rosa26*[lsl-EYFP]). As previously reported for mature

hippocampus (*19*), the vast majority of CA1 EYFP-expressing interneurons in slice

cultures from *Sst*[ires-Cre];*Rosa26*[lsl-EYFP] mice were immuno-positive for SOM and located in

stratum oriens (Fig. S1A). Whole-cell recordings from CA1 EYFP-expressing SOM-INs

revealed that EPSCs were potentiated at 24h after repeated chemical mGluR1 stimulation

(Fig. S1B). EPSC amplitude and potency were increased, and paired-pulse ratio was

decreased, compared to sham treated slices (Fig. S1C, D and G). EPSC potentiation was

prevented by the mGluR1a antagonist LY367385 (Fig. S1E and G) or the mTOR

inhibitor PP242 (Fig S1F and G). Thus, persistent late-LTP occurs at synapses onto

SOM-INs and is dependent on mGluR1a and mTORC1 signaling.

Cell-specific Rptor deletion prevents mTORC1 signaling in SOM-INs

mTORC1 is a key regulator of translation in long-term synaptic plasticity (*7, 8*).

We investigated the role of mTORC1 in SOM-INs in transgenic mice with a cell-specific

homozygous knock-out of *Rptor* in SOM-INs (*Sst*[ires-Cre];*Rosa26*[lsl-EYFP];*Rptor*[fl/fl] mice,

termed Som-Raptor-KO) and wild-type control mice ($Sst^{\text{ires-Cre}}$;$Rosa26^{\text{lsl-EYFP}}$;$Rptor^{\text{WT/WT}}$,

termed Som-Raptor-WT). First we characterized the loss of mTORC1 function in SOM-

INs in these mutant mice. The number of CA1 EYFP-expressing SOM-INs immuno-

positive for Raptor was drastically reduced in Som-Raptor-KO mice compared to -WT,

confirming a Raptor expression deficit in SOM-INs (WT: n=7, KO: n=8; Mann-Whitney

test, P=0.0013; Fig. 1A). Next we confirmed the cell-type selectivity of $Rptor$ deletion.

Western blots assays of Raptor in hippocampal lysates (Fig. 1B) and cultured slices (Fig.

1C) showed no difference in total hippocampal Raptor and p-S6 expression in lysates

(Raptor: WT: n=6, KO: n=6; t_{10}=-0.8, P=0.44; p-S6S235/236: WT: n=3, KO: n=3; t_4=-

0.3, P=0.81) and cultured slices (Raptor: WT: n=2, KO: n=2; Mann-Whitney, P=1; p-

S6S235/236: WT: n=3, KO: n=3; t_2=-0.8, P=0.49) in Som-Raptor-KO and -WT mice,

indicating that hippocampal principal neurons are unaffected by conditional deletion of

$Rptor$ in SOM-INs.

We then verified impairment of mTORC1 signaling in SOM-INs using

immunofluorescence (*8*). Repeated mGluR1 stimulation in hippocampal slices of Som-

Raptor-WT mice increased phosphorylation of ribosomal S6 protein, a downstream

effector of mTORC1, in SOM-INs, and this effect was prevented by the mTORC1

inhibitor rapamycin (Fig. 1D and F). Repeated mGluR1 stimulation failed to increase S6

phosphorylation in SOM-INs in Som-Raptor-KO mice, confirming the loss of mTORC1

signaling (Fig. 1E and F; two-way ANOVA, $F_{2,17\text{(interaction)}}$=11.6, P=0.002; Bonferroni's

tests: WT-sham [n=3] vs. WT-DHPG [n=3], P=0.0002; WT-DHPG vs. WT-

DHPG+rapamycin [n=3], P=0.0006; KO-sham [n=3] vs. KO-DHPG [n=3], P=1; KO-

DHPG vs. KO-DHPG+rapamycin [n=3], P=1; WT-DHPG vs. KO-DHPG, P=0.008).

Next we assessed if mTORC1-mediated late-LTP was affected in SOM-INs. In

slice cultures of Som-Raptor-WT mice, repeated mGluR1 stimulation (Fig. 1G),

increased amplitude (sham: n=10, DHPG: n=10; t-test with Welch correction, $t_{11.2}$=5.8,

P=0.0001) and potency (Mann-Whitney, WT: P=0.0003), and reduced paired-pulse ratio

(t-test with Welch correction, $t_{13.2}$=3.2, P=0.007), of EPSCs in SOM-INs, relative to

sham-treatment (Fig. 1H and I). Repeated mGluR1 stimulation failed to produce

potentiation of EPSCs in Som-Raptor-KO mice (sham: n=12, DHPG: n=10; amplitude:

Mann-Whitney tests, P=0.39; potency: P=0.34; paired-pulse ratio: t-test with Welch

correction, t_{20}=-0.7, P=0.48), indicating a block of persistent plasticity at excitatory

synapses onto SOM-INs (Fig. 1H and I). SOM-INs basal excitatory synaptic transmission

was intact as EPSCs were similar in Som-Raptor-KO and -WT mice in sham-treated

conditions. Heterozygous Som-Raptor-KO mice failed to show a deficit in late-LTP (Fig.

S1H). Together, these results show that conditional *Rptor* knockout in SOM-INs results

in deficient mTORC1-mediated signaling and synaptic plasticity in SOM-INs.

As control, we verified that the numbers of CA1 SOM-INs were unaffected in

Som-Raptor-KO relative to -WT mice (Fig. S2A and B), indicating that *Rptor* deletion

driven by the SOM promoter did not alter SOM-INs proliferation and migration. In

addition, because mTORC1 is a key regulator of protein synthesis implicated in cell

growth (*6*), we used whole-cell current-clamp recordings and reconstruction of biocytin-

filled SOM-INs in acute slices to show that general somatic and dendritic morphology, as

well as basic membrane properties of CA1 SOM-INs were not deficient in Som-Raptor-

KO mice (Fig. S2C to F and Fig. S3). These results indicate deficits in mTORC1

signaling and synaptic plasticity, with relatively intact morphological and membrane

properties, in SOM-INs of conditional KO mice.

SOM-IN specific Rptor deletion impairs hippocampus-dependent long-term

memory

mTORC1 is a key regulator of translation in hippocampal long-term memory (*7*).

Since hippocampal SOM-INs have been proposed to support contextual fear memories

(*17, 21*), we examined the consequences of impairment of SOM-IN mTORC1 function in

hippocampus-dependent memory tasks. We first verified in the open-field test that Som-

Raptor-KO mice showed normal anxiety level and no impairment in locomotion, relative

to -WT mice (Fig. S4). We next tested the mice in contextual fear learning and context

discrimination (Fig. 2A). Som-Raptor-KO and -WT mice froze identically in response to

footshocks, indicating normal anxiety and sensorimotor gating (WT: n= 55, KO: n=51;

Mann-Whitney, *P*>0.05; Fig. 2B). In the short-term memory test (1h), Som-Raptor-KO

and -WT mice showed similar freezing (WT: n= 11, KO: n=10; Mann-Whitney, *P*=0.70;

Fig. 2C), indicating intact short-term contextual fear memory. However, Som-Raptor-KO

mice displayed reduced freezing compared to -WT mice in the long-term memory test

(24h, WT: n= 37, KO: n=34; t_{69}=4.12, *P*=0.0001; Fig. 2C), revealing a long-term

contextual fear memory deficit. Mice were also tested for context discrimination in one of

two novel contexts varying in similarity with the training context (Fig. 2A). When

exposed to a novel context, Som-Raptor-WT mice froze more in the similar (n=13) than

in the distinct (n=13) context (Mann-Whitney, *P*=0.0021; Fig. 2D), manifesting context

generalization and discrimination, respectively. Som-Raptor-KO mice showed reduced

level of freezing in the similar novel context (n=14) compared to Som-Raptor-WT mice

(Mann-Whitney, P=0.0009; Fig. 2D). Similar results were obtained using discrimination

ratios to assess context discrimination normalized to the freezing level in the training

context (two-way ANOVA, $F_{1,47(interaction)}$=5.4, P=0.024; Tukey's tests, WT-similar vs.

WT-distinct: P=0.03; KO-similar vs. KO-distinct: P=0.96; WT-similar vs. KO-similar:

P=0.003; WT-distinct vs. KO-distinct: P=0.50; Fig. 2D), indicating impairment in

context generalization.

Fear conditioning strongly relies on amygdala function where SOM-INs play

crucial roles (*22, 23*). We subjected mice to auditory-cued fear conditioning (Fig. 2E), a

task that does not engage dorsal hippocampus, to assess if conditional *Rptor* deletion in

SOM-INs affected amygdala function. Som-Raptor-KO and -WT mice showed similar

levels of freezing in response to tone-shock presentations (WT: n= 16, KO: n=15; Mann-

Whitney, P>0.05) and in the long-term memory probe test (Mann-Whitney, Pre-Tone and

Post-Tone: P=0.57; Fig. 2F), indicating that mTORC1 function in SOM-INs is not

required for dorsal hippocampus-independent cued fear memory.

To investigate further the behavioral role of mTORC1 in SOM-INs, we examined

spatial reference memory in the Barnes maze, a hippocampus-dependent spatial learning

and memory task (Fig. 2G). Som-Raptor-WT (n=16) and -KO (n=15) mice performed

similarly during acquisition (errors: Friedman ANOVA, WT: P=0.0008; KO: P=0.036

and Mann-Whitney tests: S1: P=0.75, S2: P=0.87, S3: P=0.71, S4: P=0.08; latency:

Friedman ANOVA, WT and KO: P<0.0001 and Mann-Whitney tests: S1: P=0.89, S2:

P=0.78, S3: P=0.86, S4: P=0.013; distance: repeated measures ANOVA,

$F_{3,29(session)}$=16.2, P<0.0001; $F_{1,29(genotype)}$=1.5, P=0.23; $F_{3,29(interaction)}$=0.7, P=0.53),

indicating intact spatial learning (Fig. 2H and fig. S5A). During the memory probe test,

Som-Raptor-KO mice performed poorly with increased number of errors (Mann-

Whitney, $P=0.017$), latency ($P=0.007$) and travelled distance ($P=0.005$) in the primary

search for the target, indicating a deficit in long-term spatial memory at the single trial

level, relative to -WT mice (Fig. 2I and fig. S5A). During the total search period, Som-

Raptor-KO mice spent less time in the target quadrant than -WT mice (two-way

ANOVA, $F_{3,116(interaction)}=5.4$, $P=0.002$ and Tuckey's tests, WT-target vs. KO-target:

$P=0.028$; Fig. 2J and fig. S5A), indicating again an impairment in long-term spatial

memory. In addition, Som-Raptor-KO mice visited less the target hole, relative to -WT

mice, demonstrating a deficit in spatial memory precision (visits: Wilcoxon tests, WT:

$P=0.0005$; KO: $P=0.002$ and Mann-Whitney test, WT-target vs. KO-target: $P=0.007$;

selective search ratio: t-test against 0.5, WT: $t_{15}=10.2$, $P<0.0001$; KO: $t_{14}=5.1$, $P=0.0002$;

WT vs. KO: $t_{29}=3.9$, $P=0.0006$; Fig. 2K and fig. S5A). Mice solve the Barnes maze using

thygmotactic/serial and/or hippocampus-dependent spatial strategies (*24, 25*). During the

training protocol, Som-Raptor-WT mice rapidly manifested thygmotactic behaviors to

solve the task, visiting holes in a serial manner but progressively adopted a spatial

strategy, navigating directly to the target hole (Cochran test, $P=0.015$; Fig. 2M and fig.

S5B). Som-Raptor-KO mice did not significantly use spatial strategy but solved the maze

by relying mostly on thygmotactic or mixed strategies (Cochran test, $P=0.67$, WT: n=256

total trials, KO: n=240; z-score WT vs. KO: $P=0.016$; Fig. 2M and Fig. S5B), indicating

an impairment in hippocampus-dependent spatial memory. Together, these results reveal

that mTORC1 activity in SOM-INs is necessary for intact long-term spatial and

contextual fear memory.

Conditional Tsc1 knock-down augments mTORC1 signaling in SOM-INs

Tuberous Sclerosis Complex (TSC) is a key negative regulator of mTORC1 under

basal conditions and upstream signaling pathways inhibit TSC to activate mTORC1 (*6*).

We thus examined if *Tsc1* knock-down in SOM-INs would be sufficient to upregulate

mTORC1 activity and promote hippocampal memory. Hence we used transgenic mice

with a cell-specific heterozygous knock-out of *Tsc1* in SOM-INs (*Sst*[ires-Cre];*Rosa26*[lsl-

EYFP];*Tsc1*[WT/fl] mice, termed Som-TSC1-KO; and *Sst*[ires-Cre];*Rosa26*[lsl-EYFP];*Tsc1*[WT/WT],

termed Som-TSC1-WT). We first confirmed the cell-type selectivity of *Tsc1* knock-out.

Western blots assays showed no difference in phosphorylated S6 protein in hippocampal

lysates in Som-TSC1-KO and -WT mice (WT, n=12, KO: n=13; Mann-Whitney, *P*=0.25;

Fig. 3B), indicating unaffected mTORC1 signaling in hippocampal principal neurons.

Using immunofluorescence, basal level of phosphorylation of ribosomal S6 protein was

increased in SOM-INs in slices of Som-TSC1-KO mice relative to -WT mice (WT, n=6,

KO: n=6; t_5=8.4, *P*=0.0004; Fig. 3C to E). Repeated mGluR1 (Fig. 3A) stimulation

increased S6 phosphorylation in SOM-INs of Som-TSC1-WT relative to sham-treated

(t_5=3.2, *P*=0.024), but did not in -KO mice (t-test with Welch correction, $t_{6.1}$=-0.8,

P=0.46; Fig. 3C to E). Thus conditional Tsc1 knock-down increased basal mTORC1

activity and occluded mGluR1-induced mTORC1 activation in SOM-INs.

We then determined the effect of *Tsc1* knock-down on mTORC1-mediated

synaptic plasticity in SOM-INs. First, current-clamp electrophysiological characterization

of SOM-INs in slices of Som-TSC1-KO mice showed no major difference in basic

membrane or firing properties, except a reduction in input resistance (Fig. S6). Next, we

examined the effects of Tsc1 knock-down on chemically-induced late-LTP in SOM-INs

(Fig. 3A). First we found that basal synaptic transmission was unaffected in the mutant

mice. In sham treated slices, EPSC potency was unchanged in Som-TSC1-KO relative to

-WT mice (Fig. 3F and G). In acute slices of WT mice, like in cultured slices, repeated

mGluR1 stimulation was necessary to induce persistent potentiation of EPSC potency,

relative to sham treatment (Fig. 3F and G). However in slices of Som-TSC1-KO mice, a

single mGluR1 stimulation was sufficient, but repeated treatment failed, to induce

persistent potentiation of EPSC potency (two-way ANOVA, $F_{2,71(\text{interaction})}=19$, $P<0.0001$,

Bonferroni tests, WT-Sham [n=14] vs. WT-1xD [n=9]: $P=0.99$, WT-Sham vs. WT-3xD

[n=16]: $P=0.0014$, TSC1KO-Sham [n=14] vs. TSC1KO-1xD [n=12]: $P=0.0033$,

TSC1KO-Sham vs. TSC1KO-3xD [n=12]: $P=0.98$; Fig. 3F and G), indicating a

facilitation of induction of late-LTP in SOM-INs with *Tsc1* knock-down. We verified that

SOM-IN chemical late-LTP in acute slices was mTORC1-mediated and found that

persistent potentiation of EPSC potency by repeated mGluR1 stimulation was blocked in

SOM-INs of Som-Raptor-KO mice (WT: n=9, KO: n=10; t-test with Welch correction:

$t_{12.7}=0.5$, $P=0.65$; Fig. 3F and G). Chemical late-LTP in acute slices was associated with

unchanged paired-pulse ratio (WT vs TSC1, two-way ANOVA, $F_{2,71(\text{interaction})}=0.3$,

$P=0.76$, RaptorKO-Sham vs RaptorKO-3xD: t-test with Welch correction: $t_{9.5}=0.05$,

$P=0.96$; Fig. 3F and G), indicating some difference in mechanism with cultured slices.

Together, these results suggest that conditional mTORC1 upregulation in SOM-INs

results in a lower threshold for induction of persistent synaptic plasticity in SOM-INs,

and in an impairment of plasticity elicited by a normal induction paradigm.

Conditional Tsc1 knock-down in SOM-INs augments hippocampus-dependent long-term memory

Next we determined at the behavioral level the effects of *Tsc1* knock-down in SOM-INs. Som-TSC1-KO mice showed normal anxiety and slightly reduced locomotion in the open-field test (Fig. S7). Then we examined contextual fear memory and context discrimination (Fig. 4A). Som-TSC1-KO and -WT mice responded similarly to footshocks (WT: n=28, KO: n=31; Mann-Whitney tests, *P*>0.05; Fig. 4B), indicating intact sensorimotor gating. One hour after conditioning, Som-TSC1-KO and -WT mice showed similar freezing (WT: n=12, KO: n=13; t_{23}=-0.76, *P*=0.46; Fig. 4C), indicating intact short-term contextual fear memory. However, in the 24h memory probe test, Som-TSC1-KO mice displayed increased freezing compared to -WT mice (WT: n=16, KO: n=18; t-test with Welch correction, $t_{22.2}$=-3.08, *P*=0.005; Fig. 4C), revealing an augmentation of long-term contextual fear memory. When exposed to a novel distinct context, Som-TSC1-KO mice showed higher level of freezing (WT: n=8, KO: n=8; t_{16}=3.57, *P*=0.0027) and lower discrimination ratio (t_{16}=3.55, *P*=0.003) relative to -WT mice, indicating impairment in context discrimination (Fig. 4D). These results suggest that increased mTORC1 activity in SOM-INs is sufficient to augment long-term contextual fear memory and generalization. Next, we examined whether amygdala function was affected. Som-TSC1-KO mice were subjected to auditory-cued fear conditioning (Fig. 4E) and showed no difference in freezing during the training (WT: n=11, KO: n=13; Mann-Whitney tests, *P*>0.05; Fig. 4F) or 24h memory probe test (Mann-Whitney, Pre-Tone: *P*=0.16; Post-Tone: *P*=0.56; Fig. 4G), indicating that

upregulated mTORC1 activity in SOM-INs does not alter this dorsal hippocampus-

independent memory task.

Finally, we investigated spatial memory performance of Som-TSC1-KO mice in

the Barnes maze (Fig. 4H). Som-TSC1-KO mice showed fewer errors, shorter latency

and travelled distance to target compared to WT during the acquisition (WT: n=21, KO:

n=23; errors: repeated two-way ANOVA, $F_{1,42(genotype)}$=9.47, P=0.004; latency: Friedman

ANOVA, WT and KO: P<0.0001 and Mann-Whitney tests, S1: P=0.26, S2: P=0.012, S3:

P=0.0098, S4: P=0.048; distance: repeated two-way ANOVA, $F_{1,42(genotype)}$=6.09,

P=0.018; Fig. 4I and Fig. S8) and the long-term memory test (Mann-Whitney tests,

errors: P=0.0003, latency: P=0.0009; distance: P=0.002; Fig. 4J and Fig. S8),

demonstrating better spatial learning and memory performance. During the total search

period, Som-TSC1-KO mice visited more the target hole (Mann-Whitney, P=0.047) and

showed a higher selective search ratio than the -WT mice (t-test against 0.5, WT:

t_{20}=4.61, P=0.0002; KO: t_{22}=9.76, P<0.0001; WT vs. KO: t_{42}=-2.91, P=0.006; Fig. 4K

and Fig. S8), indicating a better selective search of the target in Som-TSC1-KO mice.

These changes observed after *Tsc1* knock-down are the converse of changes

found after *Rptor* deletion. Knock-down of *Tsc1* in SOM-INs upregulated mTORC1

activity and promoted hippocampal learning and memory. Thus mTORC1 activity level

in SOM-INs appears to determine the level and contextual precision of hippocampal

memories.

Contextual fear learning induces mTORC1-mediated plasticity at SOM-IN

excitatory synapses

Next we examined whether the effects of conditional genetic manipulations of

mTORC1 on hippocampus-dependent memory and SOM-IN synaptic plasticity could be

related. We tested if contextual fear learning produces potentiation at SOM-INs

excitatory synapses via mTORC1 using ex vivo whole-cell recordings in acute slices 24h

after training (Fig. 5A). SOM-INs from conditioned Som-Raptor-WT mice showed

increased spontaneous EPSC frequency (naive: n=15; CFC: n=13; Mann-Whitney,

P=0.015) and amplitude (Fig. 5B and C), increased minimally-evoked EPSC potency

(naive: n=11; CFC: n=14; Mann-Whitney, P=0.0004) associated with decreased minimal

stimulation intensity and unchanged paired-pulse ratio (Fig. 5D and E), as well as

potentiation of evoked EPSC input-output function (Fig. 5F and G), compared to naive

mice. Thus, conditioning induces potentiation of excitatory synapses onto SOM-INs

lasting 24h. The learning-induced potentiation was mTORC1-dependent. At 24h post-

training, SOM-INs of Som-Raptor-KO mice failed to show any potentiation of EPSCs

relative to naive mice (spontaneous EPSC amplitude: naive: n=11; CFC: n=14; two-way

ANOVA, $F_{1,49(interaction)}$=5.1, P=0.004; Tukey's tests, WT-naive vs. WT-CFC, P=0.007;

KO-naive vs. KO-CFC, P=0.69; WT-naive vs. KO-naive: P=1; minimally-evoked EPSC

potency: KO-naive [n=10] vs. KO-CFC [n=12]: P=0.87; paired-pulse ratio: two-way

ANOVA, $F_{1,43(interaction)}$=1.6, P=0.22; minimal stimulation intensity: two-way ANOVA,

$F_{1,42(interaction)}$=5.3, P=0.026; Tukey's tests, WT-naive vs. WT-CFC: P=0.005; KO-naive

vs. KO-CFC, P=0.99; WT-naive vs. KO-naive: P=0.31; input-output gain: two-way

ANOVA, $F_{1,34(interaction)}$=8.8, P=0.006; Tukey's tests,   WT-naive [n=9] vs. WT-CFC

[n=10]: P=0.0001, KO-naive [n=9] vs. KO-CFC [n=10], P=0.89; WT-naive vs. KO-

naive: P=0.99; Fig. 5B to G). SOM-INs from naive Som-Raptor-WT and -KO mice

showed similar spontaneous and evoked EPSCs (Mann-Whitney, spontaneous EPSC frequency: P=0.32; minimally-evoked EPSC potency: P=0.25; Fig. 5B to G), indicating unchanged basal synaptic transmission. Furthermore, administration of the mGluR1 antagonist JNJ16259685 to Som-Raptor-WT mice 30 min before conditioning prevented the learning-induced potentiation (spontaneous EPSC frequency: vehicle [VEH]: n=15, JNJ: n=14; Mann-Whitney, P=0.014; amplitude: t_{27}=2.13, P=0.0424; minimally-evoked EPSC potency: VEH: n=14, JNJ: n=12; t_{24}=5.31, P<0.0001; minimal stimulation intensity: t_{24}=-4.32, P=0.0002; paired-pulse ratio: t_{24}=-0.61, P=0.55; input-output gain: VEH: n=12, JNJ: n=11; Mann-Whitney, P=0.0092; Fig. 5H). These results indicate that contextual fear learning induces mGluR1- and mTORC1-mediated persistent potentiation at SOM-INs excitatory synapses, suggesting that the effects of conditional genetic manipulations of mTORC1 on hippocampus-dependent memory and SOM-IN synaptic plasticity may be linked.

mTORC1-mediated late-LTP induction protocol in SOM-INs regulates SC-LTP in pyramidal cells

CA1 area is the output of the hippocampus and LTP at CA3-CA1 pyramidal cell synapses is considered a crucial phenomenon induced by learning and supporting memory (*1, 2*). O-LM cells are a major subtype of dendrite-projecting SOM-INs (*9*) that provide differential regulation of SC and temporo-ammonic pathways onto CA1 pyramidal cells (*16*): they inhibit distal dendrites and downregulate LTP in the temporo-ammonic pathway; whereas they contact inhibitory interneurons in stratum radiatum and upregulate LTP at SC inputs (SC-LTP) by disinhibition (*16, 19, 26, 27*). Consequently,

induction of mGluR1a-mediated early-LTP at excitatory synapses onto SOM-INs

upregulates SC-LTP of pyramidal cells, indicating that plasticity at SOM-INs input

synapses regulates metaplasticity of the CA1 network (*19*). Therefore, we next examined

if mTORC1-mediated plasticity at SOM-IN input synapses controlled in a longer-lasting

manner their output and regulated metaplasticity of the SC CA1 network. First, we

established that late-LTP was elicited at SOM-INs synapses using electrical stimulation

in acute slices (Fig. 6A). Repeated TBS stimulation at the oriens/alveus border in slices

of Som-Raptor-WT mice induced, at 2h post-induction, increases in EPSC amplitude

(control: n=12; TBS: n=12; Mann-Whitney tests, *P*=0.004) and potency (*P*=0.023), and

decreases in paired-pulse ratio (*P*=0.026), relative to unstimulated slices (Fig. 6B and C).

In contrast, SOM-INs from Som-Raptor-KO mice failed to show potentiation of EPSCs

(control: n=11; TBS: n=11; Mann-Whitney, amplitude: *P*=0.74; potency: *P*=0.65; paired-

pulse ratio: *P*=0.6; Fig. 6B and C), indicating that late-LTP elicited by repeated TBS was

mediated by mTORC1.

Having established a TBS protocol inducing mTORC1-mediated late-LTP in

SOM-INs, we investigated its ability to control their output and upregulate SC-LTP of

pyramidal cells in a persistent manner. We applied the repeated TBS induction protocol

at the oriens/alveus border and two hours later applied weak high-frequency stimulation

(wHFS) in the stratum radiatum to induce SC-LTP in pyramidal cells (Fig. 6D). SC-LTP

magnitude was increased in slices of Som-Raptor-WT that received 2h previously the

SOM-INs late-LTP induction protocol, relative to previously unstimulated slices (control:

n=13; TBS: n=15; Mann-Whitney, *P*=0.03; Fig. 6G and I). The upregulation of SC-LTP

by repeated TBS was prevented in slices from Som-Raptor-KO mice (control: n=9; TBS:

n=13; KO: Mann-Whitney, $P=0.89$; WT vs. KO: $t_{26}=2.3$, $P=0.029$; Fig. 6H and I),

indicating that the upregulation of SC pathway plasticity required mTORC1-signaling,

and thus likely mTORC1-mediated late-LTP, in SOM-INs. The block of SC-LTP

regulation was not due to alterations in SC basal synaptic properties since input-output

function (WT: n=9; KO: n=10; repeated two-way ANOVA, $F_{8,135(interaction)}=0.2$, $P=0.7$;

Fig. 6E) and paired-pulse facilitation (two-way ANOVA, $F_{3,76}=0.39$, $P=0.76$; Fig. 6F)

were similar in Som-Raptor-KO and -WT mice. These results show that repeated

episodes of TBS induces long-lasting mTORC1-mediated plasticity at SOM-INs

synapses that may result in long-lasting regulation of their output and CA1 network

metaplasticity.

Together, our findings suggest that learning induces SOM-IN mTORC1 activity,

resulting in persistent LTP at SOM-IN excitatory input synapses, and that this SOM-IN

plasticity may in turn regulate CA1 SC synaptic plasticity. These SOM-IN mTORC1

cellular and circuit mechanisms could contribute to mTORC1 inhibitory circuit changes

in contextual and spatial memory.

Discussion

We used cell-type specific transgenic mouse approaches to reveal several novel

insights about mTORC1 activity in SOM-INs for hippocampal function. At the

behavioral level, we uncovered that loss of mTORC1 function in SOM-INs impaired

contextual fear and spatial long-term memories, but spared sensory-motor gating,

hippocampus-dependent short-term contextual memory and hippocampus-independent

long-term auditory-cued fear memory. In contrast, upregulation of mTORC1 activity in

SOM-INs by cell-specific conditional knockdown of *Tsc1* augmented spatial and contextual fear memories, and impaired discrimination. At the cellular level, bi-directional regulation of mTORC1 activity in SOM-INs differentially regulated mGluR1-mediated late-LTP at SOM-IN excitatory inputs. Moreover, contextual fear learning persistently increased afferent excitatory synaptic strength in SOM-INs via mGluR1 and mTORC1. At the network level, the SOM-IN late-LTP induction protocol upregulated metaplasticity of the SC pathway in pyramidal cells, in mTORC1-dependent manner. Our findings uncover a role of SOM-IN mTORC1 in learning-induced plasticity of their synapses and in hippocampal long-term memory consolidation.

Raptor is an essential component of mTORC1 which plays a central role in cell growth (*6*) in addition to regulation of protein synthesis during synaptic plasticity in mature neurons (*7*). We verified in control experiments that conditional recombination of *Rptor* in cells expressing Cre recombinase under control of the somatostatin promoter, resulted in effective loss of mTORC1 function sufficiently late not to impair development of SOM-INs. Som-Raptor-KO mice showed normal density of EYFP expressing SOM-INs in CA1 hippocampus, intact soma and dendritic morphology of individually labelled SOM-INs, as well as intact dendritic spine size and density. SOM-INs showed no change in basic membrane properties, except for a moderate increase of firing rate during sustained depolarization, as well as intact basal transmission and short-term plasticity of excitatory synaptic inputs. Thus the late-occurring conditional deletion of *Rptor* in SOM-Raptor-KO mice allowed us to examine mTORC1 function in mature SOM-INs. Similar observations were made with *Tsc1* knockdown that did not affect SOM-IN basic membrane properties, except a modest reduction of input resistance, nor basal

transmission and short-term plasticity of excitatory synaptic inputs. *Rptor* deletion and

Tsc1 knockdown were not restricted in the hippocampus but occurred in all SOM-

expressing cells and could possibly affect hippocampus-independent functions at the

behavioral level. However, the intact anxiety, the moderate change in locomotion and the

absence of freezing difference between WT and KO mice during CFC and short-term

contextual fear memory indicate that hippocampus-independent functions such as

sensorimotor gating and attention were spared in both conditional KO mice lines. Fear

learning-induced synaptic potentiation in SOM-INs in the amygdala gates auditory-cued

fear expression (*22, 23*). In the present study, conditional deletion of *Rptor* and *Tsc1*

knockdown in SOM-INs did not affect dorsal hippocampus-independent long-term

auditory-cued fear memory, indicating that learning-induced LTP in amygdala SOM-INs

crucial for cued fear memory may not require mTORC1 mechanisms. Thus, learning-

related SOM-INs synaptic plasticity mechanisms are heterogeneous with mTORC1

mechanisms prominent in hippocampal SOM-INs synaptic plasticity and showing some

brain region-specificity.

The control of protein synthesis by mTORC1 in synaptic plasticity and

memory consolidation is well-characterized in CA1 pyramidal cells (*7*). Our findings

suggest that some aspects of mTORC1 function in synaptic plasticity are cell type-

specific. We show that activation of mGluR1 in SOM-INs results in mTORC1-mediated

late-LTP of excitatory synapses (*8*). However, activation of group I mGluRs in pyramidal

cells was shown to induce mTORC1-mediated long-term depression of excitatory

synapses (*28*). More recently, mTORC2 but not mTORC1 was associated with group I

mGluR-mediated long-term depression (LTD) of CA1 pyramidal cells and novel object

recognition (*29*). Thus, mGluR-mediated mTORC1, and perhaps even mTORC2,

translational control likely regulates different mRNAs to achieve depression versus

potentiation of synapses in pyramidal cells and SOM-INs, respectively. In addition, in

other hippocampal interneurons which provides perisomatic inhibition of pyramidal cells,

endocannabinoid-dependent long-term depression of their output synapses onto

pyramidal cells requires presynaptic mTORC1-mediated protein synthesis to

downregulate GABA release (*30*). The identification of specific mRNAs controlled by

mTORC1 in different neurons classes will be important to clarify mTORC1 function in

synaptic plasticity.

Cell type-specificity of mTORC1 function is consistent with findings that

mTORC1 regulation of translation via one of its major effector, the translational

repressor eukaryotic Initiation Factor 4E Binding Protein (eIF4E-BP), controls excitatory

synaptic transmission differently in pyramidal cells and interneurons. In pyramidal cells,

knock-out of eIF4E-BP results in enhanced basal excitatory synaptic transmission (*31*), as

well as in a lower threshold for late-LTP induction and in impairment of late-LTP elicited

by normal induction paradigms (*32*). In contrast, in hippocampal O-LM inhibitory

interneurons, eIF4E-BP knockout does not affect basal excitatory transmission, lowers

the threshold for late-LTP induction and does not affect late-LTP elicited by normal

induction paradigm (*8*). Our results in SOM-INs of Som-TSC1-KO mice are largely

similar to those in eIF4E-BP knockout. Conditional knockdown of *Tsc1* in SOM-INs

increased mTORC1 activity as indicated by increase S6 phosphorylation in basal

conditions. Moreover, the increased basal activity occluded mGluR1-induced mTORC1

activity in SOM-INs. Consistent with increased mTORC1 activity, the threshold for

mGluR1-dependent late-LTP induction was lowered in SOM-INs of Som-TSC1-KO

mice, with no change in basal excitatory transmission. Thus, upregulated mTORC1

activity, in combination with ongoing synaptic activity, is not sufficient to increase

synaptic strength and mGluR1 activation is also required. Interestingly, repeated mGluR1

activation failed to induce late-LTP in SOM-INs of Som-TSC1-KO mice, suggesting the

presence of a mechanism of homeostatic control downstream of mTORC1, possibly via

S6 kinase/4E-BP2/eIF4E pathways, to prevent excessive or unselective translation of

mRNAs (*32*).

467   We demonstrated that Som-TSC1-KO mice manifested increased contextual fear

and spatial memories as well as altered context discrimination, suggesting that

upregulating mTORC1 activity in SOM-INs is sufficient to alter hippocampal memory

consolidation. These effects of *Tsc1* knockdown are converse to those of *Rptor* deletion

and suggest that the level of mTORC1 activity in SOM-INs determines the level of

persistent synaptic plasticity in SOM-INs and memory consolidation. Tuberous Sclerosis

is a genetic disease due to mutation in TSC genes resulting in excessive mTORC1

activity and severe neurologic and psychiatric manifestations (*33*). Our findings in Som-

TSC1-KO mice raise the interesting possibility that dysregulated mTORC1 activity and

synaptic plasticity in SOM-INs contribute to cognitive impairments in Tuberous

Sclerosis.

478   O-LM interneurons represent a major group of SOM-INs in CA1

hippocampus. They inhibit the distal temporo-amonic inputs of pyramidal cells in stratum

lacunosum-moleculare (*9, 17*), and send axonal projections in stratum radiatum to inhibit

interneurons that, in turn, inhibit SC inputs of pyramidal cells (*16*). O-LM interneurons

differentially control plasticity of the two inputs to pyramidal cells: they inhibit and

restrict LTP of temporo-amonic synapses; and they disinhibit and promote LTP of SC

synapses (*16, 19, 26, 27*). Hebbian mGluR1a-mediated LTP at excitatory input synapses

onto O-LM interneurons is translated into increased synaptically-driven action potential

firing (*34*), providing a synaptic basis for increased inhibition of their downstream

targets. Furthermore, induction of Hebbian mGluR1a-mediated LTP in SOM-INs

upregulates for a period of minutes LTP in the SC pathway of pyramidal cells (*19*). Here

we found that mTORC1-mediated late-LTP in SOM-INs upregulated for a period of

hours LTP in the SC pathway. Interneurons expressing SOM in the CA1 hippocampus

include not only O-LM cells but also bistratified cells and projection cells (*9*). In the

present study, 90% of the biocytin labeled cells in our recordings were O-LM cells,

clearly indicating a role of this interneuron subtype in SOM-IN mTORC1 actions in CA1.

The development of new biological markers selective of other SOM interneuron subtypes

will be necessary to assess their specific functional contributions.

We identified different physiological conditions in which mTORC1 activity is

necessary for persistent enhancement of SOM-INs excitatory synapses: activation of

mGluR1 in cultured (*8, 20*) and acute slices, stimulation of afferent fibers in acute slices,

and contextual fear learning. Our results suggest that the persistent potentiation of

excitatory synapses onto SOM-INs elicited in these conditions share some common

mechanisms. Contextual fear conditioning may engage activity of CA1 pyramidal cell

excitatory synapses and activation of local SOM-INs to induce Hebbian mGluR1-

mediated late-LTP at these synapses (*8, 20*). Hence, our results are consistent with a

model (Fig. S9) that hippocampal learning i) induces persistent mGluR1 and mTORC1-

mediated late-LTP at CA1 SOM-INs synapses due to coincident activation of presynaptic

local pyramidal cells and postsynaptic SOM-INs, ii) resulting in a long-lasting

disinhibition of SC inputs to pyramidal cells due to increased inhibition of intercalated

inhibitory interneurons, effectively leading to an upregulation of CA1 network

metaplasticity by enhancing LTP at pyramidal cell SC synapses, and iii) causing

improved consolidation of accurate hippocampal memory. In association with long-term

depression of inhibition from cholecystokinin (CCK)-positive interneurons in stratum

radiatum (*35*), and with the action of long-range inhibitory projections from entorhinal

cortex (*36*), mTORC1-mediated LTP in SOM-INs may be a complementary mechanism

for disinhibition of SC inputs in pyramidal cells (*27*). Indeed during spatial navigation, it

was recently reported that, as the animal crosses a place field, CA1 place cells decrease

their synaptic coupling with parvalbumin-expressing interneurons and increase it with

SOM-INs, causing a switch of pyramidal cell inhibition from perisomatic/proximal

dendritic to distal dendritic compartments, and allowing CA3 excitatory inputs to gain

control over entorhinal excitatory inputs in driving pyramidal cell firing in a short-term

fashion (*37*). Thus a crucial feature of mTORC1-mediated SOM-INs plasticity in

spatial/contextual information encoding by CA1 pyramidal cells may be to promote

internal representations by the hippocampal CA3 pathway while dampening external

representations via the extra-hippocampal entorhinal inputs at longer timescales.

In conclusion, our findings suggest that mTORC1 activity in SOM-INs contributes to

efficient learning and accurate hippocampal memory consolidation and is necessary for

learning-induced persistent potentiation of excitatory inputs of CA1 SOM-INs. At the

network level, SOM-IN mTORC1 mechanisms contribute to regulation of metaplasticity

of the SC pathway in pyramidal cells. Thus, mTORC1-mediated plasticity in SOM-INs

might open a permissive temporal window for distributed learning such as spatial

learning, and memory retrieval.

Materials and Methods

Subjects. All animal procedures and experiments were performed in accordance with the

Université de Montréal animal care committee regulations. Mice were group housed (2-4

per cage), maintained with food and water ad libitum and on a 12 hr light/dark cycle with

all testing performed during the light phase. Knock-in mice with an internal ribosome

entry site (IRES)-linked Cre recombinase gene downstream of the *Sst* locus ($Sst^{ires-Cre}$

mice, Jackson Laboratory #013044, Bar Harbour, ME). $Sst^{ires-Cre}$ mice were crossed with

$Rosa26^{lsl-EYFP}$ reporter mice (Ai3, Jackson Laboratory #007903) for Cre-dependent

Enhanced Yellow Fluorescent Protein (EYFP) expression in SOM-INs. Sst^{ires-}

^{Cre};$Rosa26^{lsl-EYFP}$ mice were crossed with floxed *Rptor* mice (Jackson Labs #013188) for

cell-specific knock-out of *Rptor* in SOM cells. $Sst^{ires-Cre}$;$Rosa26^{lsl-EYFP}$;$Rptor^{WT/fl}$

heterozygous offsprings were backcrossed together to obtain $Sst^{ires-Cre}$;$Rosa26^{lsl-}$

^{EYFP};$Rptor^{fl/fl}$ homozygous knock-out mice (termed Som-Raptor-KO mice), and compared

to the *Rptor* wild-type mice $Sst^{ires-Cre}$;$Rosa26^{lsl-EYFP}$;$Rptor^{WT/WT}$ (termed Som-Raptor-WT

mice). All strains were maintained on a C57BL/6N background. $Sst^{ires-Cre}$;$Rosa26^{lsl-EYFP}$

mice were crossed with floxed *Tsc1* mice (Jackson Labs # 005680) for cell-specific

knock-down of *Tsc1* in SOM cells. $Sst^{ires-Cre}$;$Rosa26^{lsl-EYFP}$;$Tsc1^{WT/fl}$ heterozygous

offsprings (termed Som-TSC1-KO mice) were compared with $Sst^{ires-Cre}$;$Rosa26^{lsl-}$

^{EYFP};$Tsc1^{WT/WT}$ control mice (termed Som-TSC1-WT mice).

Molecular biology and immunohistochemistry experiments were performed on

mice of both sex, electrophysiology experiments in mature acute slices and behavioral

tests were performed on male mice only.

EYFP-labelled interneuron distribution. Distribution of EYFP-labelled interneurons

was determined by fluorescence microscopy from hippocampal sections of Som-Raptor-

WT and -KO mice (6-12 weeks-old). Animals were deeply anesthetized intra-peritoneally

with sodium pentobarbital (MTC Pharmaceuticals, Cambridge, ON), perfused

transcardially with ice-cold 0.1M phosphate buffer (PB) and 4% para-formaldehyde in

0.1M PB (PFA) and the brain isolated. Post-fixed brains were cryoprotected in 30%

sucrose and coronal brain sections (50 μm thick) were obtained using freezing microtome

(Leica SM200R; Leica, Wetzlar, Germany). Sections were mounted in ProLong® Gold

(Invitrogen, Carlsbad, CA) and examined using an epifluorescence microscope (Nikon

Eclipse E600; Nikon, Tokyo, Japan). EYFP-positive interneuron density was assessed in

the hippocampal CA1 oriens layer.

Hippocampal slice culture. Organotypic hippocampal slice cultures were obtained from

Som-Raptor-WT and -KO mice (4-5 day-old), as described previously (*8*). In brief, the

brain was removed and dissected in HBSS (Invitrogen)-based medium. Cortico-

hippocampal slices (400 μm thick) were then obtained using a McIlwain tissue chopper

(The Cavey Laboratory Engineering Co. Ltd, Surrey, UK). After dissection, slices were

placed on Millicell culture plate inserts (Millipore, Burlington, MA) in OptiMEM

(Invitrogen) supplemented with Glutamax I and horse serum (Invitrogen) kept at 37°C in

a humidified atmosphere (95% air, 5% CO_2) for 3–7 d.

Chemical induction protocol for late-LTP. Slices received three applications (10 min

duration each at 30 min intervals) of the mGluR1/5 agonist (S)-3,5-

dihydroxyphenylglycine (DHPG, 5 µM; Abcam, Cambridge, UK) in the presence of the

mGluR5 antagonist 2-methyl-6 (phenylethynyl)-pyridine (MPEP, 25 µM; Tocris

Bioscience, Bristol, UK). Inhibitors were applied concomitantly from 30 min before to 30

min after DHPG/MPEP treatment. LY367385, rapamycin and PP242 were purchased

from Tocris, Millipore, and LC Laboratories (Woburn, MA), respectively. After

treatments, cultured and acute slices were allowed to recover for 24h and 2h,

respectively, before recordings. Experimenters were blind to all treatment groups and

mice genotype.

Immunofluorescence. Co-localization of Raptor immunofluorescence was determined in

EYFP-positive interneurons. Coronal brain sections (50 µm thick) were permeabilized

with 0.5% Triton X-100 in PBS (15 min) and unspecific binding was blocked with 10%

normal goat serum in 0.1% Triton X-100/PBS (1h). Mouse monoclonal Raptor (1/500;

Millipore) antibody was incubated 48 hours at 4ºC. Sections were subsequently incubated

at room temperature with Rhodamine-conjugated goat anti-mouse IgG1 (1/200; 90 min;

Jackson Immunoresearch Laboratories, West Grove, PA). Sections were mounted in

ProLong® Gold and imaged as for EYFP-labelled interneuron distribution. The number

of EYFP-positive interneurons in CA1 stratum oriens with co-localization of Raptor

immunofluorescence were counted and expressed as percent of the total EYFP-positive

cells.

Co-localization of somatostatin immunofluorescence in EYFP-labelled

interneurons was performed on organotypic slices obtained as described before. Slices

were fixed during 24h with 4% PFA, cryoprotected in 30% sucrose and re-sectioned (50

µm thick) using freezing microtome. Sections were permeabilized 30 min with 0.5%

Triton X-100/PBS and unspecific binding blocked as above. Sections were incubated

overnight at 4ºC with rabbit polyclonal somatostatin 28 antibody (1/2000; Abcam) and

subsequently at room temperature with Alexa 594-conjugated goat anti-rabbit IgGs

(1/500, 90 min; Abcam). Sections were mounted, imaged and co-localization measured as

described above for Raptor immunofluorescence.

S6 immunophosphorylation assay. Mice (3-5 weeks-old) were deeply anesthetized with

sodium pentobarbital and perfused transcardially with ice-cold ACSF containing (in

mM): 110 choline-chloride, 2.5 KCl, 7 $MgCl_2$, 26 $NaHCO_3$, 7 dextrose, 1.3 ascorbic acid

and 0.5 $CaCl_2$, and saturated with 95% O_2-5% CO_2. Coronal hippocampal slices (300

µm) were obtained using a vibratome (Leica VT 1000S) and transferred at room

temperature into normal oxygenated ACSF containing (in mM): 124 NaCl, 2.5 KCl, 1.25

NaH_2PO_4, 2 $MgCl_2$, 2 $CaCl_2$, 26 $NaHCO_3$, 10 dextrose, 1.3 ascorbic acid. After 1 hour

recovery period, slices received the late-LTP chemical induction protocol at 31-33ºC as

described above. Slices were fixed after treatment and cryoprotected in 30% sucrose 24

hours later. Slices were re-sectioned in 50 µm thick sections using a freezing microtome

(Leica SM200R). Sections were permeabilized with 0.3% Triton X-100 in PBS (15 min)

and unspecific binding was blocked with 10% normal goat serum in 0.1% Triton X-100/PBS (1h). Rabbit polyclonal phospho-S6 ribosomal protein (1/200; anti-phospho-S6S235/236; Cell Signaling, Danvers, MA) antibodies were incubated 48 hours at 4ºC. Sections were subsequently incubated at room temperature with Alexa 594-conjugated goat anti-rabbit IgGs (1/500; 90 min; Jackson Immunoresearch). Images were acquired using confocal microscope (LSM510, Zeiss, Oberkochen, Germany) at excitation 488 and 543 nm. Images from different treatment/groups were acquired using the exact same parameters. Cell fluorescence was quantified using Fiji (ImageJ, NIH) by comparing integrated density in cells corrected for background.

Western Blotting. Total hippocampus (10 weeks-old mice) and slice cultures (3 slices cortex-free pooled per sample) were collected and protein extracted using ice-cold radioimmunoprecipitation assay buffer containing: 50 mM Tris pH 7.4, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.5% sodium desoxycholate, 0.1% sodium dodecyl sulfate (SDS), 200 μM NaF, 200 μM Na3VO4 and protease inhibitor (Cocktail inhibitor set I; Millipore) (20 min, 4ºC). Lysates were centrifuged at 19 000 g (20 min, 4ºC) and protein concentration from supernatant was determined according to bicinchoninic acid method using bovine serum albumin as standard. Fifteen to thirty micrograms of proteins (slice culture or total hippocampus extracts respectively) were separated by 7% (Raptor) or 12% (p-S6) SDS-polyacrylamide gel electrophoresis and transferred onto polyvinilidene fluoride membrane. The membranes were blocked with 5% non-fat skin milk dissolved in Tris-buffered saline-0.1% tween 20 pH 7.4 (1h30, room temperature) and incubated with rabbit polyclonal anti-phospho-S6S235/236 (1/1000; Cell Signaling) or rabbit

monoclonal anti-Raptor (1/500; Cell Signaling) overnight at 4ºC. Membranes were then

incubated with horseradish peroxidase-conjugated anti-rabbit IgGs (1/20000; Jackson

Immunoresearch) for 1h30 at room temperature. Immunoreactive bands were detected by

enhanced chemiluminescence plus (Perkin Elmer, Waltham, MA). Membranes were next

stripped with buffer containing 0.2 M glycine pH 2.2, 0.1% SDS and reprobed with

antibodies detecting level of total S6 (1/2000; Cell Signaling) and/or tubulin (1/1000;

Cell Signaling) overnight at 4ºC. All immunoreactive bands were scanned with a desktop

scanner and quantified using Quantity One software (BioRAD, Hercules, CA).

Acute hippocampal slice preparation. Acute slices were prepared from 7- to 10-week

old Som-Raptor-WT and -KO mice. Animals were anesthetized with isoflurane

inhalation and the brain was rapidly removed and placed in ice-cold sucrose-based

cutting solution containing the following (in mM): 75 sucrose, 87 NaCl, 2.5 KCl, 1.25

NaH_2PO_4, 7 $MgSO_4$, 0.5 $CaCl_2$, 25 $NaHCO_3$, 25 glucose, 11.6 ascorbic acid and 3.1

pyruvic acid, pH 7.4, and 295 mOsmol/L. A block of tissue containing the hippocampus

was prepared and 300 or 400 μm (for whole-cell and field recordings, respectively)

transverse hippocampal slices were cut with a Leica VT1000S vibratome. Slices were

transferred for recovery for 30 min to a holding chamber in artificial cerebral spinal fluid

(ACSF) containing the following (in mM): 124 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 1.3

$MgSO_4$, 2.5 $CaCl_2$, 26 $NaHCO_3$, and 10 glucose (pH 7.3–7.4, 295–305 mOsmol/L) at

30°C and subsequently maintained at room temperature (20–22°C) for at least 90 min

until use. Both cutting solution and ACSF were saturated with 95% O_2/5% CO_2.

Whole-cell recordings. For experiments in cultured slices, culture plate inserts were

transferred to ACSF containing the following (in mM): 124 NaCl, 2.5 KCl, 1.25

NaH2PO4, 4 MgSO4, 4 CaCl2, 26 NaHCO3, and 10 glucose (pH 7.3–7.4, 295–305

mOsmol/L) maintained at room temperature for at least 30 min until use. Acute and

cultured slices were transferred to a submersion chamber perfused (3–4 ml/min) with

ACSF at 31 ± 0.5°C, CA1 and CA3 regions were disconnected by a surgical cut and

slices kept for an additional 30 minutes submerged before recording. EYFP-expressing

CA1 interneurons were identified using an upright microscope (Nikon Eclipse, E600FN),

equipped with a water-immersion long-working distance objective (40x, Nomarski

Optics), epifluorescence and an infrared video camera. Whole-cell voltage-clamp

recordings were obtained using borosilicate glass pipettes (2-5 MΩ; WPI) filled with

intracellular solution containing the following (in mM): 120 CsMeSO3, 5 CsCl, 2

MgCl2, 10 HEPES, 0.5 EGTA, 10 Na2-phosphocreatine, 2 ATP-Tris, 0.4 GTP-Tris, 0.1

spermine, 2 QX314, and 0.1% biocytin, pH 7.2–7.3, and 280 ± 5 mOsmol. For whole-cell

current-clamp recordings, the intracellular solution contained the following (in mM): 120

KMeSO4, 10 KCl, 10 HEPES, 0.5 EGTA, 10 Na2-phosphocreatine, 2.5 MgATP, 0.3

NaGTP, and 0.1% biocytin (pH 7.4, 300 mOsmol/L). Data was acquired using a

Multiclamp 700B amplifier (Molecular Devices, San Jose, CA), digitized at 20 kHz using

Digidata 1440A and pClamp 10 (Molecular Devices). Recordings were low-pass filtered

at 2 kHz. Access resistance was regularly monitored during experiments and data were

included only if the holding current was stable and access resistance varied less than 20%

of initial value. EPSCs were recorded in the presence of (2R)-amino-5-phosphonovaleric

acid (AP5; 50 µM, Sigma, St-Louis, MO) and SR-95531 (gabazine; 5 µM, Sigma) to

block NMDA and GABAA receptors, respectively. Data were acquired and analysed by

an experimenter blinded for the genotype and the treatment of the animals.

Spontaneous EPSCs were recorded in voltage-clamp mode over a period of 1-5

min and 300 consecutive EPSCs analyzed for frequency and amplitude (MiniAnalysis,

Synaptosoft, Fort Lee, NJ). Evoked EPSCs were elicited using constant current pulses

(50 µs duration) via an ACSF-filled bipolar theta-glass electrode (Harvard Apparatus,

Holliston, MA) positioned approximately 100 µm lateral to the recorded cell soma at the

border between CA1 stratum oriens and the alveus. Putative single-fiber EPSCs were

evoked at 0.5 Hz using minimal stimulation (success rate = 40-50%). EPSC potency

(EPSC amplitude excluding failures) was averaged over a 10-15 min period (pClamp 10).

Because the failure rate was an adjusted parameter, we used EPSC potency to

characterize amplitude changes in evoked transmission. Input-output function was

studied by delivering current pulses of incremental intensity (0-600 µA, 50 µA steps); 3-

10 trials per pulse intensity were delivered and responses averaged (including failures) to

determine EPSC amplitude (initial EPSC peak). Linear regressions were applied on

individual averaged responses; the slope (synaptic gain) and x-intercept (minimal

stimulation intensity) of the linear regression of the input-output relationship were

measured.

Membrane and firing properties of EYFP-labeled SOM interneurons were

measured in current-clamp recordings (*38*). Resting membrane potential was measured

with the holding current I=0 pA immediately after break-in. Input resistance (Rm) was

measured using a linear regression of voltage deflections ($\pm$ 15 mV max) in response to

current steps (500 ms, 5 pA increment, holding membrane potential -60 mV), excluding

responses with voltage deflections with activation of voltage-dependent conductance (voltage sag or ramps). Membrane time constant was calculated from the mean responses to 20 successive hyperpolarizing current pulses (-5 pA; 500 ms), determined by fitting voltage responses with a single exponential function and used to calculate specific membrane capacitance. Action potential (AP) threshold was taken as the voltage at which the slope trajectory reached 10 mV/ms. AP amplitude was the difference in membrane potential between threshold and peak. AP half-width was calculated as AP duration at half-amplitude. Fast afterhyperpolarization (fAHP) amplitude was calculated as the difference between threshold and the negative peak after the AP. The time difference between the current pulse onset and the AP peak was defined as AP latency. The rheobase was measured as the minimal current amplitude necessary to evoke an action potential. AP properties were measured for the first AP elicited by a 500 ms depolarizing current pulse just sufficient to reach threshold. The sag index was determined from a series of negative current steps (500 ms duration, 10 pA steps). From the V–I plots, the peak negative voltage deflection (Vhyp) and the steady-state voltage deflection (Vsag, calculated for the last 50 ms of the current step) were used to calculate the index as the ratio (Vrest – Vsag) / (Vrest – Vhyp), for current injections corresponding to Vsag = -80 mV.

Electrical stimulation protocol for late-LTP induction. Late-LTP was induced by electrical stimulation in acute slices using a concentric bipolar Pt/Ir electrode (FHC, Bowdoin, ME) positioned in the stratum oriens close to the alveus. Late-LTP induction protocol consisted of 4 trains of theta-burst stimulations (TBS) at 5 min intervals. Each

TBS train consisted of 3 episodes (at 30 s intervals) of 5 bursts (at 250 ms inter-burst

intervals) of 4 pulses at 100 Hz (*18, 19*). Whole cells recordings were obtained from

EYFP-positive interneurons located approximately 100 μm lateral to the stimulating

electrode at 2h after late-LTP induction or after a similar period in unstimulated slices.

LTP was assessed by recording EPSCs evoked by minimal stimulation through an ACSF-

filled bipolar theta-glass electrode positioned at approximately the same site as the

stimulating electrode used for LTP induction.

Field recordings. Field EPSPs (fEPSPs) were recorded as previously (*19*) in CA1

stratum radiatum with glass electrodes (1–3 MΩ; WPI, Saratosa, FL) filled with ASCF.

Schaffer collaterals were stimulated (0.1 ms duration; 30s-1) using a concentric bipolar

Pt/Ir stimulating electrode (FHC) placed in stratum radiatum proximal to the CA3 region.

A second concentric bipolar Pt/Ir stimulating electrode was positioned in the

oriens/alveus border proximal to the subiculum for theta-burst conditioning trains (as

described above). Field potentials were recorded with a differential extracellular

amplifier (Microelectrode AC Amplifier Model 1800, A-M Systems, Sequim, WA),

filtered at 2 kHz, digitized at 10 kHz (Digidata 1440A), and analyzed with pClamp10

(Molecular Devices). Stimulus intensity was adjusted to elicit 35% of maximal fEPSP.

Basal synaptic transmission was assessed by recording fEPSP in response to incremental

stimulations (20 to 180 μA, 20 μA steps). Short term plasticity was assessed by fEPSP in

response to paired-pulse stimulations with incremental delays (20 to 110 ms, 30 ms

steps). LTP was induced at CA1 Schaffer collateral synapses through the stimulating

electrode in the stratum radiatum by a weak high-frequency stimulation train (wHFS; 750

ms, 100 Hz). Two hours before LTP induction in the Schaffer collateral pathway, the

conditioning TBS protocol to induce late-LTP in SOM interneurons (as described above)

was applied at the oriens/alveus border. fEPSP slope was measured at 10–90% of fEPSP

amplitude.

Confocal imaging and cell reconstruction. Neurons were labeled with 0.1 % biocytin

added to the whole cell patch pipette internal solution. At the end of experiments,

hippocampal slices were postfixed overnight at room temperature in 4%

paraformaldehyde in 0.1M phosphate buffer and 0.1 % Triton. The morphology of

stained neurons was revealed using 1/1000 cy3-conjugated streptavidin (Rockland

Immunochemical, Pottstown, PA) incubated overnight at room temperature. Cy3 was

excited at 543 nm wavelength and images were captured using an LSM 880 confocal

microscope (Zeiss). Stack images (1 μm steps) were acquired through a 40x water-

immersion objective to perform three-dimension reconstructions and Sholl analysis of

SOM interneurons using the Simple Neurite Tracer and Sholl Analysis Fiji plugins

(ImageJ, NIH). SOM interneuron reconstructions revealed that in each group (Som-

Raptor-WT and -KO mice), 90% displayed all features of OLM type of SOM

interneurons, the remaining 10% showing dense axonal arborisation in strata oriens and

radiatum, typical of bistratified cells or axonal collateral running out from the

hippocampus typical of projection cells (*9*).

Behavioral experiments. Before behavioral experiments, mice were gently handled

daily for three days (~1 min per mouse) to habituate mice to the experimenter and reduce

the stress related to the experimental handling. Mice were 8-13 weeks old. The

experimenter was blind to the genotype of the mice. In all experiments, mice were first

video-tracked at 25 frames per second and their movements subsequently analyzed using

a position tracking system (Smart 3.0, PanLab, Barcelona, Spain). Mice performed the

open-field test first; the Barnes maze test started the next day and the fear conditioning

occurred 5 days after the end of the Barnes maze session. Half of the mice were subjected

to fear conditioning only.

Open-field test. Mice were allowed to freely explore a home-made circular (Som-Raptor-

WT and -KO mice, 60 cm diameter, 25 cm height) or square (Som-TSC1-WT and -KO

mice, PanLab) open-field for 5 min. Locomotion was evaluated by measuring the number

of zone transitions using a pattern divided into 25 (circular OF) or 21 (squared OF) zones

of equal size. Anxiety was assessed by measuring the time spent in a 20 cm radius/side

central zone versus in a 20 cm broad peripheral annulus.

Fear conditioning. Mice were trained in conditioning chambers that were housed in

sound- and light-isolated cubicles (Coulbourn Instruments, Holliston, MA). The

chambers contained a stainless steel grid floor, overhead LED lighting, camera and were

supplied with background noise (60 dB) by an air extractor fan. The experimental

protocol was based on Ruediger and coworkers (*10*) with slight modifications. The

training context was rectangular with transparent walls and was cleaned with 1% acetic

acid before and after each trial. Two neutral contexts were designed to assess contextual

generalization and discrimination. One had a triangular shape, transparent walls and was

cleaned with 70% ethanol before and after each trial. This context was considered novel

but similar to the training context because they shared some features. Another neutral context had a circular shape, opaque black and reflective walls, Plexiglas floor and was cleaned with 70% ethanol before and after each trial. This context was considered novel and distinct to the training context. Freezing was assessed using FreezeFrame (Coulbourn Instruments). Once placed in the conditioning chamber, mice were allowed to freely explore for 2.5 min, and then received 5 presentations of unconditioned and conditioned stimuli (1 s foot shock, 0.8mA; where indicated, 10 kHz tone for 10 s, 70 dB sound pressure level, 30 s interleaved). The last 1 s of each tone was paired with the unconditioned stimulus. To test for cued fear conditioning, mice explored the novel distinct context for 2.5 min, followed by 5 tone presentations (tone-dependent freezing). Contextual fear conditioning involved the same protocol, but without the tone component. To test for contextual fear memory, mice were returned to the training context during a test period of 2.5 min, at 1 or 24 h after conditioning, to assess short- or long-term memory, respectively. To test for contextual discrimination after fear conditioning, a within-subjects design was used. On the test day, 5 h after the test in the training context, mice were pseudo-randomly distributed across the two novel neutral contexts and freezing was assessed during 2.5 min. Discrimination ratio was calculated as the amount of freezing in (training context) / (training context + novel context) (*39*). A ratio of 1 indicates that mice were able to discriminate the contexts perfectly, and a ratio of 0.5 means that they were unable to discriminate. For a subset of training-induced persistent plasticity experiments, mice were subjected to contextual fear conditioning 30 min after IP injection of 1 mg/kg JNJ16259685 (Tocris) or vehicle.

Barnes maze. This test was used to assess hippocampus-dependent spatial learning (*24*) using an elevated platform (PanLab). The experimental protocol was based on Sunyer and coworkers (*40*) with slight modifications. Mice were trained to use spatial cues to find a small dark escape chamber under the platform termed "target". The target is retained at the same position relative to the room, while the platform is rotated with each trial to discourage use of the intra-maze odor cues. In addition, the platform, the starting chamber and the escape box were thoroughly cleaned (Versaclean 10 %) between every single trial to reduce any possible scent trails. Assignment of target location was balanced among experimental groups. An aversive pulsed noise (85 dB) went on during each trial and switched off when the mouse entered in the target. One day after a familiarization period (day 1), mice were trained in 4 daily sessions with an inter-trial interval of 15 min for 4 days. During the training, mice were left 10 s in a dark cylinder (starting chamber) in the center of the maze, allowed to freely explore the maze for a maximum duration of 3 min and left 1 min in the escape box after each trial. If a mouse failed to find the t by the end of the 3 min period, it was gently guided to it. Video tracking and data analysis were performed using Smart 3.0 (PanLab). On day 6, the mice were exposed to a probe trial in which the escape box was closed. Mice were allowed to explore the maze for 3 min. The number of errors, the latency and the distance travelled before the first reaching of the target (primary search) were collected. During the total search (90 s), the time spent in the quadrants (target, left, right and opposite, excluding a 15 cm diameter central zone) and the number of visits for each hole were collected. Non-target data was defined as the average number of visits in the 19 non-target holes. The selective search ratio was defined as the number of visits in (target) / (target + non-target). A ratio of 1 indicates

that a mouse visited only the target hole, and a ratio of 0.5 means that it visited target and

non-target holes equally. Search strategies were also collected and defined into three

categories: 1) direct (spatial) – moving directly to the target hole or to an adjacent hole

before visiting the target; 2) serial (thygmotactism) – the first visit to the target hole

preceded by visiting at least two adjacent holes in serial manner, in clockwise or counter-

clockwise direction; and 3) random (or mixed) – hole searches separated by crossing

through the centre of the maze or unorganized search (*24, 25*). One search strategy was

attributed for every single trial of the training and for the primary search of the probe test.

Statistical analysis. No statistical methods were used to predetermine the sample size,

but our sample sizes are similar to (or larger than) those generally employed in the field.

Statistical analysis was performed using OriginPro 2016 (OriginLab, Northampton, MA).

Data were tested for normality and homoscedasticity using the Shapiro-Wilk and the

Brown–Forsythe tests, respectively. We used Student t tests (with Welch corrections for

heteroscedasticity), two-way ANOVA with Tukey's pairwise comparison tests (with

Bonferroni adjustments for multiple comparisons) and two-way repeated-measures

ANOVA when data passed normality and homoscedasticity assumptions. Mann-Whitney

tests and Friedman ANOVA were used when it was not the case. All the tests were two-

sided. In the figures, data are expressed as arithmetic mean and standard error of the

mean (mean ± SEM). Asterisks denote statistical significance as calculated by the

specified statistical tests (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$, ns, not significant).

References and Notes

1. J. R. Whitlock, A. J. Heynen, M. G. Shuler, M. F. Bear, Learning induces long-term potentiation in the hippocampus. Science 313, 1093-1097 (2006).

2. J. H. Choi, S. E. Sim, J. I. Kim, D. I. Choi, J. Oh, S. Ye, J. Lee, T. Kim, H. G. Ko, C. S. Lim, B. K. Kaang, Interregional synaptic maps among engram cells underlie memory formation. Science 360, 430-435 (2018).

3. E. R. Kandel, Y. Dudai, M. R. Mayford, The molecular and systems biology of memory. Cell 157, 163-186 (2014).

4. R. A. Nicoll, A Brief History of Long-Term Potentiation. Neuron 93, 281-290 (2017).

5. T. Abel, P. V. Nguyen, M. Barad, T. A. Deuel, E. R. Kandel, R. Bourtchouladze, Genetic demonstration of a role for PKA in the late phase of LTP and in hippocampus-based long-term memory. Cell 88, 615-626 (1997).

6. R. A. Saxton, D. M. Sabatini, mTOR Signaling in Growth, Metabolism, and Disease. Cell 169, 361-371 (2017).

7. M. Costa-Mattioli, W. S. Sossin, E. Klann, N. Sonenberg, Translational control of long-lasting synaptic plasticity and memory. Neuron 61, 10-26 (2009).

8. Ran, I. Laplante, C. Bourgeois, J. Pepin, P. Lacaille, M. Costa-Mattioli, J. Pelletier, N. Sonenberg, J. C. Lacaille, Persistent transcription- and translation-dependent long-term potentiation induced by mGluR1 in hippocampal interneurons. The Journal of neuroscience : the official journal of the Society for Neuroscience 29, 5605-5615 (2009).

9. K. A. Pelkey, R. Chittajallu, M. T. Craig, L. Tricoire, J. C. Wester, C. J. McBain,

Hippocampal GABAergic Inhibitory Interneurons. Physiol Rev 97, 1619-1747

(2017).

10. S. Ruediger, C. Vittori, E. Bednarek, C. Genoud, P. Strata, B. Sacchetti, P. Caroni,

Learning-related feedforward inhibitory connectivity growth required for memory

precision. Nature 473, 514-518 (2011).

11. J. S. Polepalli, H. Wu, D. Goswami, C. H. Halpern, T. C. Sudhof, R. C. Malenka,

Modulation of excitation on parvalbumin interneurons by neuroligin-3 regulates the

hippocampal network. Nature neuroscience 20, 219-229 (2017).

12. J. G. Pelletier, J. C. Lacaille, Long-term synaptic plasticity in hippocampal feedback

inhibitory networks. Progress in brain research 169, 241-250 (2008).

13. D. M. Kullmann, A. W. Moreau, Y. Bakiri, E. Nicholson, Plasticity of inhibition.

Neuron 75, 951-962 (2012).

14. M. Lovett-Barron, G. F. Turi, P. Kaifosh, P. H. Lee, F. Bolze, X. H. Sun, J. F.

Nicoud, B. V. Zemelman, S. M. Sternson, A. Losonczy, Regulation of neuronal input

transformations by tunable dendritic inhibition. Nature neuroscience 15, 423-430,

S421-423 (2012).

15. S. Royer, B. V. Zemelman, A. Losonczy, J. Kim, F. Chance, J. C. Magee, G. Buzsaki,

Control of timing, rate and bursts of hippocampal place cells by dendritic and somatic

inhibition. Nature neuroscience 15, 769-775 (2012).

16. R. N. Leao, S. Mikulovic, K. E. Leao, H. Munguba, H. Gezelius, A. Enjin, K. Patra,

A. Eriksson, L. M. Loew, A. B. Tort, K. Kullander, OLM interneurons differentially

modulate CA3 and entorhinal inputs to hippocampal CA1 neurons. Nature

neuroscience 15, 1524-1530 (2012).

17. M. Lovett-Barron, P. Kaifosh, M. A. Kheirbek, N. Danielson, J. D. Zaremba, T. R.

Reardon, G. F. Turi, R. Hen, B. V. Zemelman, A. Losonczy, Dendritic inhibition in

the hippocampus supports fear learning. Science 343, 857-863 (2014).

18. Y. Perez, F. Morin, J. C. Lacaille, A hebbian form of long-term potentiation

dependent on mGluR1a in hippocampal inhibitory interneurons. Proceedings of the

National Academy of Sciences of the United States of America 98, 9401-9406

(2001).

19. C. Vasuta, J. Artinian, I. Laplante, S. Hebert-Seropian, K. Elayoubi, J. C. Lacaille,

Metaplastic Regulation of CA1 Schaffer Collateral Pathway Plasticity by Hebbian

MGluR1a-Mediated Plasticity at Excitatory Synapses onto Somatostatin-Expressing

Interneurons(1,2,3). eNeuro 2, (2015).

20. Ran, I. Laplante, J. C. Lacaille, CREB-dependent transcriptional control and quantal

changes in persistent long-term potentiation in hippocampal interneurons. The Journal

of neuroscience : the official journal of the Society for Neuroscience 32, 6335-6350

(2012).

21. L. C. Schmid, M. Mittag, S. Poll, J. Steffen, J. Wagner, H. R. Geis, I. Schwarz, B.

Schmidt, M. K. Schwarz, S. Remy, M. Fuhrmann, Dysfunction of Somatostatin-

Positive Interneurons Associated with Memory Deficits in an Alzheimer's Disease

Model. Neuron 92, 114-125 (2016).

22. H. Li, M. A. Penzo, H. Taniguchi, C. D. Kopec, Z. J. Huang, B. Li, Experience-
dependent modification of a central amygdala fear circuit. Nature neuroscience 16,
332-339 (2013).

23. M. A. Penzo, V. Robert, J. Tucciarone, D. De Bundel, M. Wang, L. Van Aelst, M.
Darvas, L. F. Parada, R. D. Palmiter, M. He, Z. J. Huang, B. Li, The paraventricular
thalamus controls a central amygdala fear circuit. Nature 519, 455-459 (2015).

24. C. A. Barnes, Memory deficits associated with senescence: a neurophysiological and
behavioral study in the rat. J Comp Physiol Psychol 93, 74-104 (1979).

25. G. B. Fox, L. Fan, R. A. LeVasseur, A. I. Faden, Effect of traumatic brain injury on
mouse spatial and nonspatial learning in the Barnes circular maze. J Neurotrauma 15,
1037-1046 (1998).

26. J. I. Hurtado-Zavala, B. Ramachandran, S. Ahmed, R. Halder, C. Bolleyer, A.
Awasthi, M. A. Stahlberg, R. J. Wagener, K. Anderson, R. M. Drenan, H. A. Lester,
J. M. Miwa, J. F. Staiger, A. Fischer, C. Dean, TRPV1 regulates excitatory
innervation of OLM neurons in the hippocampus. Nat Commun 8, 15878 (2017).

27. J. Artinian, J. C. Lacaille, Disinhibition in learning and memory circuits: New vistas
for somatostatin interneurons and long-term synaptic plasticity. Brain Res Bull 141,
20-26 (2018).

28. J. L. Banko, L. Hou, F. Poulin, N. Sonenberg, E. Klann, Regulation of eukaryotic
initiation factor 4E by converging signaling pathways during metabotropic glutamate
receptor-dependent long-term depression. The Journal of neuroscience: the official
journal of the Society for Neuroscience 26, 2167-2173 (2006).

29. P. J. Zhu, C. J. Chen, J. Mays, L. Stoica, M. Costa-Mattioli, mTORC2, but not

mTORC1, is required for hippocampal mGluR-LTD and associated behaviors. Nature

neuroscience 21, 799-802 (2018).

30. T. J. Younts, H. R. Monday, B. Dudok, M. E. Klein, B. A. Jordan, I. Katona, P. E.

Castillo, Presynaptic Protein Synthesis Is Required for Long-Term Plasticity of

GABA Release. Neuron 92, 479-492 (2016).

31. Ran, C. G. Gkogkas, C. Vasuta, M. Tartas, A. Khoutorsky, I. Laplante, A. Parsyan, T.

Nevarko, N. Sonenberg, J. C. Lacaille, Selective regulation of GluA subunit synthesis

and AMPA receptor-mediated synaptic function and plasticity by the translation

repressor 4E-BP2 in hippocampal pyramidal cells. The Journal of neuroscience: the

official journal of the Society for Neuroscience 33, 1872-1886 (2013).

32. J. L. Banko, F. Poulin, L. Hou, C. T. DeMaria, N. Sonenberg, E. Klann, The

translation repressor 4E-BP2 is critical for eIF4F complex formation, synaptic

plasticity, and memory in the hippocampus. The Journal of neuroscience: the official

journal of the Society for Neuroscience 25, 9581-9590 (2005).

33. M. Costa-Mattioli, L. M. Monteggia, mTOR complexes in neurodevelopmental and

neuropsychiatric disorders. Nature neuroscience 16, 1537-1543 (2013).

34. Croce, J. G. Pelletier, M. Tartas, J. C. Lacaille, Afferent-specific properties of

interneuron synapses underlie selective long-term regulation of feedback inhibitory

circuits in CA1 hippocampus. The Journal of physiology 588, 2091-2107 (2010).

35. J. Basu, K. V. Srinivas, S. K. Cheung, H. Taniguchi, Z. J. Huang, S. A. Siegelbaum, A cortico-hippocampal learning rule shapes inhibitory microcircuit activity to enhance hippocampal information flow. Neuron 79, 1208-1221 (2013).

36. J. Basu, J. D. Zaremba, S. K. Cheung, F. L. Hitti, B. V. Zemelman, A. Losonczy, S. A. Siegelbaum, Gating of hippocampal activity, plasticity, and memory by entorhinal cortex long-range inhibition. Science 351, aaa5694 (2016).

37. Fernandez-Ruiz, A. Oliva, G. A. Nagy, A. P. Maurer, A. Berenyi, G. Buzsaki, Entorhinal-CA3 Dual-Input Control of Spike Timing in the Hippocampus by Theta-Gamma Coupling. Neuron 93, 1213-1226 e1215 (2017).

38. L. Tricoire, K. A. Pelkey, B. E. Erkkila, B. W. Jeffries, X. Yuan, C. J. McBain, A blueprint for the spatiotemporal origins of mouse hippocampal interneuron diversity. The Journal of neuroscience: the official journal of the Society for Neuroscience 31, 10948-10970 (2011).

39. B. J. Wiltgen, A. J. Silva, Memory for context becomes less specific with time. Learning & memory 14, 313-317 (2007).

40. B. Sunyer, S. Patil, H. Höger, G. Lubec, Barnes maze, a useful task to assess spatial reference memory in the mice. (2007).

Acknowledgments

General: We thank J. Pepin for help with slice cultures and chemical late-LTP experiments, and members of the Lacaille laboratory for helpful discussions and comments on the manuscript.

Funding: This work was supported by the Canadian Institutes of Health Research (operating grants to J.-C.L.; MOP-10848 and MOP-125985), the Fonds de la Recherche du Québec en Santé [FRQS Group grant to J.-C.L.; Groupe de Recherche sur le Système Nerveux Central (GRSNC)]. J.-C.L. is the recipient of the Canada Research Chair in Cellular and Molecular Neurophysiology (950-231066). J.A. was supported by GRSNC and FRQS Fellowships; A.L.F. by a FRQS Fellowship.

Author contributions: J.A., I.L. and J.C.L. designed and directed experiments; J.A., A.J., A.K., E.H., A.L.F., A.S.R. and I.L. performed the experiments and analysed data; I.L. generated and validated transgenic mice lines; J.A. and J.C.L. conceived the project and wrote the paper with contributions from I.L.

Competing interests: The authors declare no competing interests.

Data and materials availability: Requests for materials should be addressed to J.C.L.

 Figures

Fig. 1: Cell-specific deficit in Raptor expression impairs mTORC1 signaling and late-LTP in SOM-INs of Som-Raptor-KO mice. (**A**) Left, Representative images of Raptor immunopositive (red) EYFP$^+$ (green) CA1 SOM-INs (arrows, co-labeling) in Som-Raptor-KO relative to -WT mice. Right, Percentage of Raptor$^+$ cells relative to EYFP$^+$ cells. (**B**) Left, Representative Western blots of Raptor and phospho-S6$^{S235/236}$ from hippocampal lysates. Right, Raptor (normalized to tubulin) and p-S6 (normalized to S6 and tubulin) levels (relative to Som-Raptor-WT mice). (**C**) Same as in (**B**) but in hippocampal cultured slices. (**D-E**) Representative confocal images illustrating EYFP$^+$ CA1 SOM-INs (green), S6$^{S235/236}$ phosphorylation (red) and co-labeling (merged) in Som-Raptor-WT (**D**) and -KO (**E**) mice, after sham, DHPG/MPEP and DHPG/MPEP in the presence of 1 μM rapamycin treatments. Arrows point to EYFP$^+$ SOM-INs with p-S6 co-labeling. (**F**) Phosphorylated S6$^{S235/236}$ level relative to sham treatment in Som-Raptor-WT mice. (**G**) Diagram of chemical late-LTP experimental protocol in cultured slices. (**H**) Representative EPSCs evoked by minimal stimulation in SOM-INs. Traces are superimposed 20 consecutive events (EPSCs + failures, black), average EPSC (including failures, red) and average of EPSC pairs (20 events) evoked by paired-pulse stimulation. (**I**) EPSC amplitude (including failures), EPSC potency (excluding failures) and paired-pulse ratio of SOM-INs 24h after DHPG/MPEP relative to sham treatment in Som-Raptor-WT and -KO mice.

Fig. 2: Som-Raptor-KO mice show impaired long-term contextual fear and spatial memory. (**A**) Diagram of the contextual fear conditioning protocol. (**B**) Percentage of time freezing after each footshock during the training session for Som-Raptor-WT and -KO mice (0: before the first footshock). (**C**) Percentage of time freezing during the probe tests at 1h (left) and at 24h (right). (**D**) Percentage of time freezing during the contextual discrimination test in a novel context (left) and discrimination ratio (right). (**E**) Diagram of the auditory-cued fear conditioning protocol. (**F**) Percentage of time freezing for Som-Raptor-WT and -KO mice during the training session (left) and during the probe test at 24h (right). (**G**) Diagram of the spatial learning protocol in the Barnes maze. (**H**) Performance in the Barnes maze during the training of Som-Raptor-WT and -KO mice. (**I**) Spatial memory performance during the primary search of the probe test. Errors, latency and distance before the first visit of the target. (**J**) Percentage of time spent in each quadrant of the maze during the total search period of the probe test. L: left, T: target, R: right, O: opposite quadrants. (**K**) Number of visits to all escape holes (left), number of visits expressed as target vs. average of all non-target holes (middle) and selective search ratios (right). (**M**) Percentage of trials using a spatial strategy over the training protocol. The dashed line represents chance (10%).

Fig. 3: *Tsc1* knockdown increases mTORC1 signaling and facilitates mGluR1-mediated late-LTP in SOM-INs. (**A**) Diagram of chemical late-LTP experimental protocol in acute slices. (**B**) Left, Representative Western blots of phosphorylated $S6^{S235/236}$ from hippocampal lysates. Right, p-$S6^{S235/236}$ (normalized to S6 and tubulin) levels (relative to Som-TSC1-WT mice). (**C, D**) Representative confocal images illustrating $EYFP^+$ CA1 SOM-INs (green), $S6^{S235/236}$ phosphorylation (red) and co-labeling (merged) in Som-TSC1-WT (**C**) and -KO (**D**) mice, after sham and DHPG/MPEP treatments. Arrows point to $EYFP^+$ SOM-INs with p-S6 co-labeling. (**E**) p-$S6^{S235/236}$ level in the different groups relative to sham treatment in Som-TSC1-WT mice. (**F**) Representative EPSCs evoked by minimal stimulation in SOM-INs. Traces are superimposed 20 consecutive events (EPSCs + failures, black), average EPSC (including failures, orange) and average of EPSC pairs (20 events) evoked by paired-pulse stimulation. (**G**) EPSC potency (excluding failures) and paired-pulse ratio of SOM-INs 2h after sham, single and repeated DHPG/MPEP treatment in WT, Som-TSC1-KO and Som-Raptor-KO mice.

Fig. 4: *Tsc1* **deletion in SOM-INs increased long-term contextual fear and spatial memories but impaired context discrimination.** (**A**) Diagram of the contextual fear conditioning protocol. (**B**) Percentage of time freezing after each footshock during the training session for Som-TSC1-WT and -KO mice (0: before the first footshock). (**C**) Percentage of time freezing during the probe tests at 1h and at 24h after conditioning. (**D**) Percentage of time freezing (left) and discrimination ratio (right) during the contextual discrimination test. (**E**) Diagram of the auditory-cued fear conditioning protocol. (**F, G**) Percentage of time freezing during the training session (**F**) and during the probe test 24h after (**G**) for Som-TSC1-WT and -KO mice. (**H**) Diagram of the spatial learning protocol in the Barnes maze. (**I**) Learning curves for Som-TSC1-WT and -KO mice during the training session. (**J**) Memory performance during the primary search of the probe test. (**K**) Number of visits to all escape holes (left) and selective search ratio (right) during the total search of the probe test.

Fig. 5: Contextual fear learning induces mGluR1- and mTORC1-mediated persistent LTP at excitatory synapses onto SOM-INs. (**A**) Diagram of the experimental protocol in acute slices. a: alveus, o: oriens, p: pyramidale, r: radiatum, lm: lacunosum-moleculare, stim: stimulation pipette, rec: recording pipette. (**B**) Representative traces of spontaneous synaptic activity in SOM-INs from naive and trained Som-Raptor-WT and -KO mice. (**C**) Spontaneous EPSC frequency and amplitude. (**D**) Representative traces of EPSCs evoked by minimal stimulation in the different conditions. (**E**) Minimally-evoked EPSC potency, paired-pulse ratio and minimal stimulation intensity. (**F**) Representative traces of evoked input-output function in SOM-INs in the different conditions. (**G**) Input-output gain of SOM-INs as the slope of individual linear regressions. (**H**) Synaptic properties of SOM-INs from CFC-trained Som-Raptor-WT mice treated with mGluR1 antagonist JNJ16259685 relative to vehicle (VEH). Spontaneous EPSC frequency and amplitude, minimally-evoked EPSC potency, minimal stimulation intensity and paired-pulse ratio, and input-output gain.

1103

1104

Fig. 6: mTORC1-mediated late-LTP induction protocol in SOM-INs upregulates SC-LTP in principal cells. (**A**) Recording configuration for late-LTP induction and whole-cell recording 2h later. (**B**) Representative traces of EPSCs evoked by minimal stimulation in control condition and after repeated TBS stimulation, in Som-Raptor-WT and -KO mice. Traces are superimposed 20 consecutive events (EPSCs + failures, black), average EPSC (including failures, orange) and average of EPSC pairs (20 events) evoked by paired-pulse stimulation. (**C**) EPSC amplitude, potency (excluding failures) and paired-pulse ratio. (**D**) Recording configuration and simplified diagram of underlying CA1 network. SC: Schaffer collaterals, TA: temporo-ammonic pathway, PC: pyramidal cell, Som: SOM-INs, rad: radiatum interneuron. (**E**) Representative traces and amplitude of field EPSPs in response to incremental SC stimulation in Som-Raptor-WT and -KO mice. (**F**) Representative traces and ratio of field EPSPs amplitude in response to SC paired-pulse stimulations. (**G**) Top: Description of the stimulation and induction protocol. Left: representative traces of field EPSPs in response to SC stimulation 10 min before (1) and 30 min after weak HFS (wHFS, 100 Hz, 750 ms) stimulation (2) to induce SC-LTP, in control condition and 2h after the SOM-INs late-LTP induction protocol (repeated TBS stimulation) in Som-Raptor-WT mice. Right: Graph of LTP of field EPSP slope induced by wHFS. (**H**) Same as (**G**) in Som-Raptor-KO mice. (**I**) SC-LTP upregulation 2h after SOM-INs late-LTP induction protocol, normalized to the averaged control SC-LTP.

Supplementary Materials

Fig. S1. Chemically-induced persistent LTP at excitatory synapses onto CA1 SOM-INs

depends on mGluR1a and mTOR.

Fig. S2. Intact SOM interneuron numbers and morphology in Som-Raptor-KO mice.

Fig. S3. Membrane and firing properties of SOM-INs from Som-Raptor-KO mice.

Fig. S4. Anxiety and locomotor activity of Som-Raptor-KO mice in the open field test.

Fig. S5. Spatial memory deficits in Som-Raptor-KO mice.

Fig. S6. Membrane and firing properties of SOM-INs from Som-TSC1-KO mice.

Fig. S7. Anxiety and locomotor activity of Som-TSC1-KO mice in the open-field test.

Fig. S8. Spatial memory strengthening in Som-TSC1-KO mice.

Fig. S9. A model for regulation of hippocampal memory by mTORC1 in CA1 SOM

interneurons.