1 Host-pathogen transcriptomics of macrophages, Mucorales and their endosymbionts: a
2 polymicrobial pas de trois
3

4 Authors
5 Poppy Sephton-Clark[1], José F Muñoz[2], Herbert Itabangi[1], Kerstin Voelz[1], Christina A Cuomo[2],
6 Elizabeth R Ballou[1]*
7

[1]*Institute of Microbiology and Infection, School of Biosciences, University of Birmingham,*
Edgbaston, Birmingham, B15 2TT, UK

[2]*Infectious Disease and Microbiome Program, Broad Institute of MIT and Harvard,*
Cambridge, Massachusetts, USA

*To whom correspondence should be addressed
ERB: e.r.ballou@bham.ac.uk
14

Abstract

16
Mucorales spores, the causative agents of mucormycosis, interact with the innate immune
system to cause acute, chronic, or resolving infection. Understanding the factors that
influence disease initiation and progression is key to understanding mucormycosis and
developing new treatments. Complicating this, mucormycosis can be caused by a number of
species that span the Mucorales genus and may be host to bacterial endosymbionts. This
study sets out to examine the differences between two species in the Mucorales order by
characterising their differential interactions with the innate immune system, and their
interactions with environmental bacterial endosymbionts. Through a holistic approach, this
study examines the transcriptional responses of *Rhizopus delemar* and *Rhizopus*
microsporus, two of the most commonly diagnosed species, to innate immune cells. This
study also examines the immune cell response and assesses the variation in these
responses, given the presence or absence of bacterial endosymbionts within the fungi. We
see that the fungal response is driven by interaction with innate immune cells. The effect of
the bacterial endosymbiont on the fungus is species specific, with a minimal in the absence
of stress, but strongly influencing fungal transcripts during interaction with innate immune
cells. In contrast, we observe that the macrophage response varies depending on the
infecting fungal species, but also depending on endosymbiont status. The most successful
macrophages elicit a pro-inflammatory response, and we see that through germination
inhibition macrophage survival is enhanced. This work reveals species-specific host
responses to related Mucorales spores and shows that bacterial endosymbionts impact the
innate immune cell response.
38

Introduction

40
41 Mucormycosis is a devastating, environmentally acquired fungal infection caused by a
42 variety of Mucorales species. Understanding the interaction between Mucorales spores and
43 innate immune cells is key to understanding mucormycosis. Studies frequently focus upon
44 the interaction of a single species of the Mucorales order with innate immune cells (Warris

et al. 2005; Chamilos et al. 2008; Schmidt et al. 2013; Kraibooj et al. 2014; Inglesfield et al. 2018). However, between Mucorales species, and across infecting isolates, numerous phenotypic and genomic differences can be observed (Hoffmann et al. 2013; Mendoza et al. 2014; Schwartze et al. 2014). Whether each species within this family employs the same mechanism to cause infection remains unclear, however the range of virulence profiles displayed across the family indicates there may be diversified mechanisms at work (Petraitis et al. 2013). For example, it has been shown that pathogenicity of these species is linked to the variable copy number of the gene encoding CotH, a family of cell surface proteins important for the spores' interactions with and adherence to endothelial cells. Iron-scavenging and melanisation pathways have also been shown to play a role in pathogenicity (Chibucos et al. 2016, Andrianaki et al. 2018). In addition to fungal genome plasticity and transcriptional responses to host stimuli, fungal-host interactions can be further modulated by bacterial endosymbionts (Itabangi et al., 2019; Partida-Martinez & Hertweck 2005). The majority of Mucorales species have been shown to harbour bacterial endosymbionts whose species can vary between Mucorales isolates (Ibrahim et al. 2008; Kobayashi & Crouch 2009; Mondo et al. 2017; Itabangi et al. 2019). We have recently shown that a bacterial endosymbiont influences the outcome of *Rhizopus microsporus* infections in both zebrafish and murine models through modulation of both fungal and phagocyte phenotypes (Itabangi et al. 2019). However, the impact of endosymbiont on fungal and host transcriptional response remains unclear.

While disease can be caused by several species in the Mucorales order, *Rhizopus delemar* and *Rhizopus microsporus* are responsible for the majority of infections (Liu et al. 2018). Risk factors for developing the disease include, but are not limited to: innate immune cell deficiencies, uncontrolled diabetes, immune suppression, skin barrier breaches and iron overload (Baldin & Ibrahim 2017). With limited antifungal treatments effective against these invasive infections, there is often a need for surgical debridement or amputation (Spellberg et al. 2016). In disseminated cases, mucormycosis may present with a mortality rate of over 90% (Mendoza et al. 2014). Reports of chronic mucormycosis have also emerged, affecting both immunocompromised and immunocompetent patients (Alan et al. 2019). The innate immune response to Mucorales spores is key to infection control: macrophages are required to induce a differential immune response on contact with *Rhizopus* spp. and may inhibit germination or kill hyphal forms upon contact in healthy individuals (Ghuman & Voelz 2017; Andrianaki et al. 2018). Single and multi-species studies have shown that phagosome maturation is arrested by melanin within the cell walls of *Aspergillus* spp. and *Rhizopus* spp., however iron limitation allows macrophages to more effectively kill *Rhizopus* spp.(Liu et al. 2015; Ibrahim et al. 2010; Andrianaki et al. 2018). Several works comparing *Aspergillus* spp. to *Rhizopus* spp. have revealed similar immunostimulatory capacities, but differences in their responses to host stress (Warris et al. 2005; Chamilos et al. 2008; Schmidt et al. 2013; Kraibooj et al. 2014). Exploring and understanding fungal responses to the host is essential to improving our understanding of mucormycosis, yet it remains unclear how Mucorales species respond to, and interact with, the innate immune system, and to what extent this varies by species.

Our work explores the interplay between the innate immune system, *R. delemar,* and *R. microsporus* using isolates of both species found to harbour bacterial endosymbionts (Itabangi et al. 2019). We investigate the differences between these two fungal species, how

they respond transcriptionally to innate immune cells, and how their respective bacterial
endosymbionts affect this interaction. We also investigate the transcriptional response of
innate immune cells to these infectious spores and determine how this interaction is
influenced by the presence of an endosymbiont. We reveal a large difference between the
fungal transcriptional profiles of *R. delemar* and *R. microsporus* during *in vitro* monoculture.
There is a small conserved response to exposure to innate immune cells, including key
changes in cell wall genes, consistent with germination. Conversely, we see that the host
innate immune response differs significantly between fungal species and is also influenced
by the presence or lack of an endosymbiont. The innate immune response to *R. delemar* and
R. microsporus mirrors the relative aggressiveness of infection between these two species.
We also observe that through the activation of innate immune cells, or upon inhibition of
chitin synthase, we can improve the ability of innate immune cells to control the fungal
spores. Our work represents a broad analysis of the transcriptional interplay between innate
immune responders and infectious Mucorales spores, revealing species-species differences
which question the current model of 'one species represents all', when it comes to
mucormycosis.
108

Results

Experimental Design

We set out to investigate paired transcriptional responses of host and fungal cells, whilst
also exploring the influence of the endosymbiont on this interaction (Figure 1a). Fungal
spores from *Rhizopus delemar* and *Rhizopus microsporus* were either cured via ciprofloxacin
treatment to remove the bacterial endosymbiont (cured) or maintained in media permissive
to bacterial endosymbiosis (wt). Cured spores were passaged twice in the absence of
ciprofloxacin to limit the impact of the drug on transcriptional responses. The cured and wt
spores of both *R. delemar* and *R. microsporus* were allowed to swell in sabouraud broth
until 95% of the population had reached mid isotropic phase. Due to the differences in
germination rates between the species (Figure 1b), this occurred at 2 hours for *R. delemar*
and 4 hours for *R. microsporus*. Swollen spores were then used to infect the J774.1 murine
macrophage-like cell line. Fungal spores were co-cultured with macrophages for one hour,
after which unengulfed spores were removed, and phagocytosed spores were incubated
within the macrophages for a further two hours. The cells from the resulting infection were
processed to explore their transcriptional response to this infection scenario (Figure 1a).
Macrophages that had phagocytosed fungal spores were isolated and sequenced via the
10X Genomics Chromium Single Cell Sequencing platform. Macrophages left unexposed to
the fungi were used as a negative control. RNA was also isolated from fungal spores (cured
and wt) which had been engulfed by macrophages, and this was sequenced with a bulk
RNA-Seq approach. Unexposed fungi (cured and wt) were incubated in macrophage media
for a matched time and used as a negative control. The data shows the fungal response to
phagocytosis by macrophages, as well as the fungal response to the presence of its
endosymbiont. The macrophage response to the two species is also revealed, a response
which appears to differ when the endosymbiont is present for both fungal species.
136

Comparative genomics predicts alternative transcriptional responses

In order to better understand differential disease progression, we chose to compare *R. delemar* and *R. microsporus*, as they cause a large proportion of mucormycosis infections but appear morphologically dissimilar. *R. delemar* germinates more quickly than *R. microsporus*, with 50% of spores germinating by 3 hours, and a spore body size which reaches 12.8 μm over the course of germination (Figure 1b, c). *R. microsporus* germinates at a significantly slower rate, taking 6 hours for 50% of spores to germinate, with a final spore body size of 13.9 μm (Figure 1b, c). To better understand their relationship to one another, we compared the gene content to explore the similarity and differences between the two species. Previous work has established that the *R. delemar* genome (45.3 Mb, 17,513 genes) is larger, and contains an increased number of genes, compared to *R. microsporus* genome (26 Mb, 10,959 genes) (Ma et al. 2009; Mondo et al. 2017). Our results show that, compared to *R. microsporus*, the genome of *R. delemar* is enriched for genes with protein domains (PFAM) associated with ion binding, carbohydrate derivative binding, nucleic acid binding, cytoskeletal protein binding, poly(A) binding, NAD+ ADP-ribosyltransferase activity, protein kinase C activity, translation initiation factor binding and inorganic phosphate transmembrane transporter activity (Supplemental Figure S1, Figure 1d). *R. microsporus* is enriched for genes with protein domains corresponding to nucleoside phosphate binding, early endosome activity and DNA repair complex activity (Supplemental Figure S1, Figure 1d). The clear differences illustrated by genome size and gene content indicate the likelihood of alternative transcriptional responses.

Alternative transcriptional profiles are presented by *R. delemar* and *R. microsporus* in response to innate immune cells

First we examined overall trends in fungal responses to phagocytosis, obtained through our bulk RNA-Seq approach. We analysed the signal obtained from *R. delemar* and *R. microsporus* samples with principle component analysis (PCA) (Figure 2). We observed large differences between the transcriptomes of both fungal species, when exposed or unexposed to macrophages, while the presence or absence of their respective endosymbionts had a weak but differential effect on PCA. The presence or absence of the endosymbiont appears to have very little bearing on the transcriptional patterns displayed by *R. delemar*, as samples fell into two distinct clusters, most strongly influenced by macrophage status (Figure 2a). *R. microsporus* exhibits a similar trend upon exposure to macrophages, however the presence of the endosymbiont also influenced clustering (Figure 2b). We investigate the phenotypic consequences of this interaction in the companion paper by Itabangi et al., showing that the presence of the endosymbiont *Ralstonia pickettii* impacts fungal cell wall organization, resistance to host-relevant stress, spore germination efficiency, and pathogenesis (Itabangi et al., 2019). Therefore, we focus here on the transcriptional analysis of the host-pathogen-endosymbiont interaction across the two species.

There are 2,493 genes that are significantly differentially expressed (Log fold change > 2; false discovery rate < 0.05) in *R. delemar* across all conditions (Figure 3a), while *R. microsporus* only exhibits 40 genes significantly differentially expressed across all conditions (Log fold change > 2; false discovery rate < 0.05) (Figure 3b). The theme of a muted

transcriptional response from *R. microsporus* is also seen within pairwise comparisons of
conditions. Pairwise comparisons of differential expression across each experimental
condition show similar trends in responses between *R. delemar* and *R. microsporus*,
however *R. microsporus* responds with a reduced gene set (Figure 4). Pairwise comparisons
showed the biggest shift in transcriptional response when comparing phagocytosed fungal
spores to those unexposed to macrophages, regardless of endosymbiont status. When
phagocytosed (Supplemental Figure S2), *R. microsporus* upregulates genes enriched in GO
categories corresponding to thiamine metabolism, sulfur metabolism, glycerol metabolism,
alcohol dehydrogenase activity and transmembrane transporter activity (hypergeometric
test, corrected *P* value < 0.05). This is consistent with the fungal response seen to
macrophage stress (Parente-Rocha et al. 2015), and the micronutrient scavenging response
to nutritional immunity (Ballou and Wilson 2016; Shen et al., 2018; Andrianaki et al., 2018).
Phagocytosed *R. microsporus* downregulated genes enriched in GO categories
corresponding to rRNA processing, ribosome biogenesis and ribosome localization
(hypergeometric test, corrected *P* value < 0.05) (Supplemental Figure S2), consistent with
growth arrest within the phagolysosome (Inglesfield et al. 2018; Andrianaki et al. 2018).
200
Comparisons of *R. delemar* conditions reveal that, upon phagocytosis, spores upregulate
genes enriched in KEGG classifications corresponding to MAPK signalling, phenylalanine
metabolism, tyrosine metabolism, glutathione metabolism and fatty acid synthesis
(hypergeometric test, corrected *P* value < 0.05). Upregulation of these processes is
consistent with melanin biosynthesis (Eisenman et al. 2011; Andrianaki et al. 2018) and
intra-phagosomal survival (Yadav et al., 2011; Lorenz and Fink, 2005). Unexposed *R. delemar*
spores upregulate genes enriched in KEGG classifications corresponding to ketone body
synthesis, protein processing via the endoplasmic reticulum, amino sugar and nucleotide
sugar metabolism (hypergeometric test, corrected *P* value < 0.05). This is consistent with
metabolic activation and cell wall biogenesis (Figure 5).
211
R. delemar and *R. microsporus* harbour distinct bacterial endosymbionts (Itabangi et al.,
2019). When comparing transcriptional profiles of wt and cured spores incubated in serum-
free DMEM (sfDMEM) for 3 hours (time matched to phagocytosis assay), we observed very
few transcriptional changes in either species in response to curing (Supplemental Table 1).
In *R. delemar,* a single gene, predicted to be a putative protein phosphatase, was repressed.
In *R. microsporus,* three genes were induced: an autophagy-related protein, a C2H2 zinc
finger transcription factor, and ribosomal protein L2.
219
Despite these small changes, loss of the endosymbiont significantly impacted the
transcriptional responses of both fungal species to macrophages (Figure 4). Specifically, we
observed an overall increase in the number of fungal genes differentially regulated upon
phagocytosis for both species. Exposure of wt *R. microsporus* to macrophages induced the
expression of one gene with no known function and repressed 6 genes. The repressed genes
include an autophagy-related protein and a zinc finger transcription factor, as well as 4 un-
annotated genes (Figure 4, Supplemental Table S1). In contrast, when cured *R. microsporus*
spores were exposed to macrophages, 277 genes were significantly induced and 82
repressed, compared to unexposed cured spores (Figure 4). Induced genes were enriched
(hypergeometric test, corrected *P* value < 0.05) for the following GO categories: organelle
organisation, pre-ribosome and ribosome activity, ATPase activity, hydrolase activity,

pyrophosphatase activity, helicase activity, nucleic acid binding, RNA metabolism, nitrogen
metabolism, chromatin silencing. Repressed genes were enriched (hypergeometric test,
corrected *P* value < 0.05) for the following GO categories: oxidoreductase activity, hydrogen
sulphide metabolism, glycolysis, sulphur metabolism, hexose catabolism, siderophore activity,
iron assimilation, nitrogen metabolism, carboxylic acid metabolism. This suggests an overall
failure to properly respond to host stresses such as iron starvation in the absence of the
bacterial endosymbiont.
238
A similar impact of the endosymbiont was observed for *R. delemar*. While the overall fungal
response to phagocytosis is characterized by a robust transcriptional response, induced
genes in wt samples (1137 genes, Figure 4) were enriched for KEGG classifications
corresponding to: alanine metabolism, PPAR signalling, aromatic compound biosynthesis
and degradation, lysine metabolism, lipid metabolism, MAPK signalling, sugar metabolism,
tyrosine metabolism, secondary metabolite biosynthesis (Figure 5). Repressed genes in wt
samples (472 genes, Figure 4) were enriched for KEGG classifications corresponding to:
carbohydrate metabolism, secondary metabolite biosynthesis, ketone body processing,
protein processes, MAPK signalling (Figure 5). In contrast, induced genes in cured samples
(1285 genes, Figure 4) were enriched for KEGG classifications corresponding to: Fatty acid
metabolism, DNA replication, amino acid metabolism, glycan metabolism, pyruvate
metabolism, secondary metabolite processing (Figure 5). Repressed genes in cured samples
(878 genes, Figure 4) were enriched for KEGG classifications corresponding to: sugar
metabolism, amino acid metabolism, lipid metabolism, MAPK signalling, NOD-like receptor
signalling. Again, this suggests that the endosymbiont has an overall suppressive impact on
fungal transcription in response to macrophage challenge.
255
Next, we directly compared the transcription patterns of genes shared across the two fungal
species. When comparing the transcriptional responses of orthologous genes shared by *R.*
delemar and *R. microsporus*, we saw only a small proportion behave similarly (213 genes).
When phagocytosed, wt spores from both species upregulate orthologues genes involved in
fatty acid catabolism, transcription, regulation via polymerase II, and organelle organization.
Phagocytosed cured spores from both species upregulate orthologues genes involved in
RNA processing, chromosome organization and condensed chromosome pathways. When
unexposed, we see wt spores upregulate orthologous genes involved in translocation,
protein binding, siderophore activity, cobalmin processing, and post-translational protein
targeting. Cured unexposed spores upregulate orthologous genes with roles in siderophore
activity and transferase activity. Overall, *R. delemar* and *R. microsporus* both respond
transcriptionally to the presence of macrophages, however the size and composition of this
response differs between species.
269
Finally, we examined the regulation of genes predicted to be involved in ferrous iron
transport, as previous work has linked iron scavenging to survival within the phagolysosome
(Andrianaki et al., 2018). There are 12 genes in the *R. delemar* genome with predicted
ferrous iron roles. While 8 showed no significant change over the tested conditions, 3
(ROG3_006623, ROG3_007727, ROG3_011864) appeared highly expressed in wt and cured
phagocytosed spores, compared to unexposed spores. The last gene, ROG3_009943, is
highly expressed in wt spores unexposed to macrophages. Together, this suggests there

may be condition dependent specialization in the expression of ferrous iron transport in *R.*
delemar.
279

Innate immune cell transcription varies with fungal species and endosymbiont presence

282
To investigate the innate immune response when challenged with *R. delemar* and *R.*
microsporus, we carried out single cell RNA-Seq of J774.A1 murine macrophages, unexposed
and exposed for 3 hours to the four types of pre-swollen spores (Figure 1a). Transcription of
both challenged and unchallenged macrophages displayed underlying population
heterogeneity (Supplemental Figure S3). To identify the transcriptional patterns of genes
responding to the spores, we focused on the expression of a subset of genes previously
identified as immune response genes (Muñoz et al. 2018). Principle component analysis of
the aggregated transcriptional data shows there is a clear difference in transcription
between macrophages that have and have not been exposed to the fungi (Figure 6). Across
all exposed conditions, relative to unexposed macrophages, there was a profile consistent
with cytokine activation, response to stimulus, and activation of the NF-Kb pathway. This
was accompanied by repression of CCL5, which is involved in T-cell recruitment (Figure 7).
However, different macrophage profiles can be seen in response to the two fungal species,
and these are further influenced by the presence of the endosymbiont. While the response
to wt *R. delemar* shows the most deviation from the macrophage-only control, exposure to
cured *R. delemar* also elicited a strong and distinct macrophage response (Figures 6,7).
Exposure to wt *R. delemar* elicits increased expression of general markers of activation,
including GTPase activity and MHC class II protein binding (LAG3 repressor of T-cell
activation, H2-M2, IFN-gamma induced IIGP1, MX1, KCTD14, PNP2), growth factor binding,
IL1 receptor agonist activity and endocytosis (SERPINE1, ENG, FGFBP3, GM8898, GCNT2,
IL1F6). Specifically, we observe modest increases in the expression of IFN-γ responsive
CXCL10 (3.1 fold) and IRG1/IRG11 (5.9 fold), pro-inflammatory SAA3 (3.5 fold), and ENPP4
(2.8 fold), but also induction of the M2 polarizing PSTPIP2/21 (6.7 fold), the IL-4 responsive
signaling modulator CISH (5.4 fold), and the vascular damage responsive F3/F31 (5.9 fold)
(Martinez et al. 2013). These latter genes are not as strongly induced during exposure to *R.*
microsporus, which may reflect the aggressive nature of infection by *R. delemar* relative to
R. microsporus. In contrast, infection with cured *R. delemar* showed a decrease in the
induction of these M2-polarisation markers (PSTPIP2, 3.5 fold relative to uninduced). The
transcriptional profile is instead shifted to include increased transcription of genes involved
in G protein signaling and phosphoinositide binding (PDE7B, CCL1, SCARF1, RGS16,
PLEKHA4) (Figure 7).
314
A similar change in macrophage polarization was observed during exposure to wt and cured
R. microsporus. The phenotypic analysis of this is discussed more fully in Itabangi et al. 2019.
For both wt and cured *R. microsporus,* expression of IFN-γ responsive CXCL10, SAA3, and
ENPP4 was comparable to unexposed macrophages. Compared to *R. delemar,* exposure to
wt *R. microsporus* induced a more limited expression of genes with roles in cytokine
activation, ERK1 and ERK2 regulation, and regulation of NF-kB cascade (Figure 7). There was
also a weaker induction of the vascular damage responsive F3/F31 genes, and a relatively
stronger upregulation of the PLA2G16 phospholipase, TRIM30D, SLC1A2 and the M2
polarizing IL-6. However, other key polarizing genes, particularly PSTPIP2/21, SAA3, and

ENPP4 were only weakly induced (Supplemental Table S2). Overall, this profile suggests a
weak M1-like activation consistent with poor phagocytosis and reduced overall antifungal
activity that we observe in macrophages interacting with endosymbiont-harbouring spores
(Itabangi et al., 2019).
328
Finally, cured *R. microsporus* induced a strong pro-inflammatory response, which included
upregulation of CXCL3, the neutrophil chemoattractant, consistent with our observations of
differences in phagocyte recruitment in zebrafish upon infection with wt vs. cured spores
(Itabangi et al., 2019). Markers of NF-kB activation were also strongly induced in this
population. Cured *R. microsporus* also strongly induced the expression of TNFRSF8 (CD30), a
marker of lymphocyte activation occasionally associated with subcutaneous fungal
infections. Overall, this is suggestive of a shift to a more pro-inflammatory profile. In our
companion paper, we observed that cured *R. microsporus* is more sensitive to phagocyte-
mediated killing and phagocyte recruitment compared to wt (Itabangi et al., 2019). We
therefore went on to test whether a more successful response to the spores could be
mounted via the induction of a pro-inflammatory response.
340
341 ## Chitin synthase inhibition and pro-inflammatory priming regulate infection outcome
342
R. delemar exhibits rapid germination followed by hyphal extension (Sephton-Clark et al.,
2018). The transcriptional profile we observe here in macrophages exposed to *R. delemar* is
consistent with a strong damage response, likely prompted by the germination of *R.
delemar* spores. We therefore hypothesised that macrophages may be better able to
control the infection if the fungi were slowed in their developmental progress. The necessity
of genes involved in chitin synthesis and regulation appeared important for both *R. delemar*
and *R. microsporus* in response to phagocytosis (Supplemental Figure S4). In previous work
we also highlighted the importance of chitin synthase for germination (Sephton-Clark et al.
2018). When fungal spores were pre-treated with the chitin synthesis inhibitor Nikkomycin Z
(24 µg/ml), spores failed to germinate, displayed less chitin/chitosan in their outer cell wall
(Supplemental Figure S5) and macrophage survival was increased at 7.5 hours post infection
(Figure 8). At Nikkomycin Z concentrations lower than those used to pre-treat spores for
phagocytosis, we see the spores are able to swell, however development appears halted
after swelling (Supplemental Figure S5). As the macrophages are better able to control
these spores, this suggests that spores undergoing the initial stages of germination may
offer less of a challenge for the macrophages.
359
Our transcriptional data show a strong M2 alternative activation signal during *R. delemar*-
macrophage interaction, but a weaker M2 polarisation during *R. microsporus*-macrophage
interaction that was further shifted towards NF-kB-mediated M1 upon endosymbiont cure.
We therefore hypothesized that shifting the macrophage polarization towards M1 classical
activation might have a protective effect upon Mucorales infection. Consistent with this, the
pre-treatment of macrophages with NF-kB activating lipopolysaccharide (LPS) significantly
improved the ability of macrophages to control *R. microsporus*. At 7.5 hours post infection,
59.7% of macrophages survived when pre-treated with LPS, compared to 24.6% without
(Figure 8).
369

Discussion

In this work we show the fungal response to innate immune cells differs by species in the Rhizopodaceae family. Although *R. delemar* and *R. microsporus* share a small conserved response to exposure to macrophages, the majority of their response differs. We show that the ability to germinate prior to phagocyte control appears to be key to virulence, as blocking spore germination with the chitin synthase inhibitor Nikkomycin Z improves macrophage survival. A range of germination and virulence phenotypes can be seen throughout the Mucorales, and this highlights a need for further investigation into these differences, to better understand the infections they cause.

We have also shown that the fungal transcriptional response appears largely unperturbed by the presence, or lack, of an endosymbiont, in the absence of stress. However, the presence of an endosymbiont greatly effects the response of and to the host. It has been shown that endosymbionts influence asexual development and sporulation through the regulation of Ras2 (Mondo et al. 2017). While we observed limited changes in fungal transcription when comparing wt and cured spores, we did observe changes in the expression of fungal genes upon exposure to macrophages. In our companion paper, Itabangi et al show that the presence of an endosymbiont is important for virulence of *R. microsporus*. The endosymbiont enhances virulence through the secretion of an anti-phagocytic factor. The endosymbiont also impacts organisation of the fungal plasma membrane, as well as environmental stress resistance and resistance to macrophage-mediated killing (Itabangi et al., 2019). We show here that this is mirrored by the transcriptional response of the WT and cured spores of both *Rhizopus* species to macrophages, and in the macrophage response to infection.

As anticipated, activation of pro-inflammatory pathways increased macrophage survival in response to the spores. This consolidates several studies which demonstrate improved innate immune cell response to fungal pathogens when primed (Rogers et al. 2013; Municio et al. 2013; Blasi et al. 1995). It has been shown that the early immune response to a *Mucor circinelloides* infection is dependent on the formation of a pro-inflammatory TNF-α-expressing granuloma-like structure that controls but does not kill spores in the zebrafish model (Inglesfield et al. 2018). Ungerminated *Rhizopus* spores are highly resistant to ROS/RNS which may mediate survival within granulomata. Consistent with this, the induction of a strong proinflammatory response by cured *R. microsporus*, hypersensitive to ROS/RNS stress, allows macrophages to better control the spores.

Interestingly, we see the chitin synthase inhibitor, Nikkomycin Z, is able to inhibit germination in both *R. delemar* and *R. microsporus*. This demonstrates the requirement of chitin synthase for spore development in these species. Treated spores swell, but do not polarise. The ability to germinate also has consequences for virulence. The ability of macrophages to control swollen spores, but not subsequently polarised ones, highlights that the developmental stage of the spore is key to innate immune success and control. Whilst many fungal pathogens are virulent in an ungerminated form, germination appears a key virulence factor for these filamentous fungi. This is confirmed by work which shows that the developmental stage of both *R. microsporus* and *Rhizopus oryzae* spores impacts pathogenesis in zebrafish and murine models (Itabangi et al., 2019; Andrianaki et al., 2018).

Specifically, infection with pre-swollen spores leads to evasion of macrophage-mediated
immunity and increased pathogenesis (Itabangi et al., 2019; Andrianaki et al., 2018). In both
models, spore clearance is dependent on phagocyte recruitment, and we show that
phagocyte recruitment (in Itabangi et al.) and activation (Figure 7) is influenced by
endosymbiont status (Itabangi et al., 2019; Andrianaki et al., 2018). In addition, we
extended this analysis by revealing profound differences in the host response to two closely
related *Rhizopus* species. In particular, we observe a M2/damage-associated response
during infection with wt *R. delemar* spores that is shifted towards an M1 protective
response upon infection with cured *R. microsporus* spores. We reinforce this finding through
experimental modulation of macrophage polarization, showing that exposure to strongly
M1-polarizing LPS is sufficient to reduce macrophage killing by wt *R. delemar* spores.
Therefore, our data provide a framework for beginning to understand differences in the
relative virulence of pathogenic Mucorales species and underpin our finding of cross-
kingdom fungal-bacterial symbiosis influencing mammalian disease.
431

Methods

Fungal Culture

R. delemar and *R. microsporus* were cultured with Sabouraud dextrose agar (SDA) or broth
(10 g/liter mycological peptone, 20 g/liter dextrose), sourced from Sigma-Aldrich, at room
temperature. Spores were harvested, 10 days after plating, with phosphate-buffered saline
(PBS), centrifuged for 3 min at 3,000 rpm, and washed. Appropriate concentrations of
spores were used for further experiments as indicated. To cure spores of their respective
bacterial endosymbionts, spores were cultured with ciprofloxacin, as described in Itabangi
et al. 2019. Once cured, spores were subcultured at least twice in ciprofloxacin-free media
before use.

Macrophage Culture

444 Macrophages from the J774.A1 cell line were cultured in Dulbecco's Modified Eagle
445 Medium, (complemented with 10% foetal bovine serum, 1% penicillin, 1% streptomycin and
446 1% L-glutamine). Macrophages were grown at 37°C, in 5% CO_2.

Phagocytosis Assay

Macrophages were incubated for one hour in serum-free DMEM prior to infection. Spores
were pre-swollen in SAB (2 hr for *R. delemar,* 4 hr for *R. microsporus).* Washed spores were
incubated at 5:1 MOI with 1×10^5 macrophages as described in Itabangi et al, to ensure that
>95% of macrophages contained one spore or more. After a 1 hour of incubation, excess
spores were washed off the surface and the macrophages were incubated for a further 2
hours, before processing for RNA-Seq experiments. For live cell imaging experiments,
images were taken starting immediately after the excess spores were removed.

Live Cell Imaging

458 Time course images were taken to determine how LPS pre-treatment (100ng/ml) (Myers et
459 al. 2010) of macrophages, and Nikkomycin Z pre-treatment (120ug/ml over the course of
460 swelling in SAB) of spores effected phagocytic outcome. Images were taken at 20x on a Zeiss
461 Axio Observer, with images taken every 5 minutes. Bright-field and fluorescent images were

462 then analysed using ImageJ V1.

Comparative Genomics and Enrichment Analysis
464 Fishers exact test was used to detect enrichment of Pfam terms between *R. delemar* and *R.*
465 *microsporus*, terms with a corrected P value of < 0.01 were considered significant.
466 Orthologue genes of *R. delemar* and *R. microsporus* were identified using blast+. R (version
467 3.3.3) was used to carry out hypergeometric testing of KEGG and GO terms to determine
468 enrichment.

Bulk RNA-Seq
RNA was extracted from spores which had either been incubated with the macrophages, or
incubated in DMEM for the equivalent time period. To remove the macrophages, triton at
1% was used to lyse the macrophages, the resulting solution was then centrifuged for 3 min
at 3,000 rpm, and washed, leaving only spores. The DMEM control also received the same
treatment. To extract total RNA, the washed samples were immediately immersed in TRIzol
and lysed via bead beating at 6,500rpm for 60s. Samples were then either immediately
frozen at -20°C and stored for RNA extraction or placed on ice for RNA extraction. After lysis,
0.2 ml of chloroform was added for every 1 ml of TRIzol used in the sample preparation.
Samples were incubated for 3 min and then spun at 12,000 g at 4°C for 15 min. To the
aqueous phase, an equal volume of 100% ethanol (EtOH) was added, before the samples
were loaded onto RNeasy RNA extraction columns (Qiagen). The manufacturer's
instructions were followed from this point onwards. RNA quality was checked by Agilent,
with all RNA integrity number (RIN) scores above 7 (Schroeder et al. 2006). One microgram
of total RNA was used for cDNA library preparation. Library preparation was done in
accordance with the NEBNext pipeline, with library quality checked by Agilent. Samples
were sequenced using the Illumina HiSeq platform; 100-bp paired-end sequencing was
employed (2 x 100 bp) (>10 million reads per sample).

Single Cell RNA-Seq
For single cell sequencing experiments, macrophages were infected with fungal spores, as
outlined above. Uninfected macrophages, used as a negative control, were treated in the
same manner and underwent mock washes and media changes at identical time points to
infected macrophages. Macrophages were isolated and released from the bottom of their
wells with accutase, as per manufacturer's instructions (Technologies n.d.). Once in solution,
the macrophages were loaded onto the 10X genomics single cell RNA sequencing pipeline
for single cell isolation and library preparation. In total, 1082 single cells were sequenced.
The libraries were sequenced on the Illumina Sequencing Platform.

Data Analysis
For the bulk RNA-Seq data, FastQC (version 0.11.5) was employed to ensure the quality of
all samples. Hisat2 (version 2.0.5) was used to align reads to the indexed genome of
Rhizopus delemar RA 88-880 (PRJNA13066, Ma et al. 2009) and the indexed genome of
Rhizopus microsporus (Mondo et al. 2017). HTSeq (version 0.8.0) was used to quantify the
output (Anders et al. 2015). Trinity and edgeR (Robinson et al. 2009)(version 3.16.5) were
then used to analyse differential expression (Grabherr et al. 2013). For the single cell RNA-
Seq data, the 10X genomics analysis pipeline (Loupe Cell Browser V 2.0.0, Cell Ranger
Version V 2.0.0) was used to align reads to the *mus musculus* genome (version MM10), and

quantify the output. For single cell analysis, the samples were then aggregated using this
pipeline, to allow comparisons between samples.

Data Availability
Data will be available upon request.
511

Acknowledgments

We are grateful to the University of Birmingham's Genomics Services Facility and to
Deborah Croom-Carter for her technical support.

Author contributions
PSC conceived and designed the experiments, collected the data, performed the analysis
and interpretation, and wrote the manuscript. JFM, KV and CAC contributed to
interpretation of the data, and contributed to the manuscript. ERB conceived and designed
the experiments, contributed to interpretation of the data, and wrote the manuscript.
521

Funding

PSC was supported by a BBSRC MIBTP PhD Studentship (BB/M01116X/1). This work was
supported by a Wellcome Trust Seed award to KV (108387/Z/15/Z). HI was supported by the
Wellcome Trust Strategic Award in Medical Mycology and Fungal Immunology (097377).
CAC and JFM were funded by the National Institute of Allergy and Infectious Diseases,
National Institutes of Health, under award U19AI110818 to the Broad Institute. ERB was
supported by the UK Biotechnology and Biological Research Council (BB/M014525/1) and a
Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society
(211241/Z/18/Z).

531
532

533 ## References
534

Alan, E.C. et al., 2019. An Emergent Entity⬚: Indolent Mucormycosis of the Paranasal Sinuses
. A Multicenter Study. , (2193), pp.92–100.

Anders, S., Pyl, P.T. & Huber, W., 2015. HTSeq-A Python framework to work with high-
throughput sequencing data. *Bioinformatics*, 31(2), pp.166–169.

Andrianaki, A.M. et al., 2018. Iron restriction inside macrophages regulates pulmonary host
defense against Rhizopus species. *Nature Communications*, 9(1). Available at:
http://dx.doi.org/10.1038/s41467-018-05820-2.

Baldin, C. & Ibrahim, A.S., 2017. Molecular mechanisms of mucormycosis—The bitter and
the sweet. *PLOS Pathogens*, 13(8), p.e1006408. Available at:
http://dx.plos.org/10.1371/journal.ppat.1006408.

Berger, U. V. & Hediger, M.A., 2006. Distribution of the glutamate transporters GLT-1
(SLC1A2) and GLAST (SLC1A3) in peripheral organs. *Anatomy and Embryology*, 211(6),
pp.595–606.

Blasi, E. et al., 1995. Role of nitric oxide and melanogenesis in the accomplishment of
anticryptococcal activity by the BV-2 microglial cell line. *Journal of Neuroimmunology*,
58(1), pp.111–116.

Chamilos, G. et al., 2008. Zygomycetes Hyphae Trigger an Early , Robust Proinflammatory
Response in Human Polymorphonuclear Neutrophils through Toll-Like Receptor 2
Induction but Display Relative Resistance to Oxidative Damage ⬚. , 52(2), pp.722–724.

Chibucos, M.C. et al., 2016. An integrated genomic and transcriptomic survey of
mucormycosis-causing fungi. *Nature Publishing Group*, 7, pp.1–11. Available at:
http://dx.doi.org/10.1038/ncomms12218.

Gerrick, K.Y. et al., 2018. Transcriptional profiling identifies novel regulators of macrophage
polarization. *Plos One*, 13(12), p.e0208602. Available at:
http://dx.plos.org/10.1371/journal.pone.0208602.

Ghuman, H. & Voelz, K., 2017. Innate and Adaptive Immunity to Mucorales. *Journal of Fungi*,
3(4), p.48. Available at: http://www.mdpi.com/2309-608X/3/3/48.

Grabherr, M.G.., Brian J. Haas, Moran Yassour Joshua Z. Levin, Dawn A. Thompson, Ido Amit,
Xian Adiconis, Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua Chen, Evan
Mauceli, Nir Hacohen, Andreas Gnirke, Nicholas Rhind, Federica di Palma, Bruce W., N.
& Friedman, and A.R., 2013. Trinity: reconstructing a full-length transcriptome without
a genome from RNA-Seq data. *Nature Biotechnology*, 29(7), pp.644–652.

Hoffmann, K. et al., 2013. The family structure of the Mucorales⬚: a synoptic revision based
on comprehensive multigene-genealogies. , pp.57–76.

Hou, Y. et al., 2014. The inhibitory effect of IFN-gamma on protease HTRA1 expression in
rheumatoid arthritis. *Journal of immunology (Baltimore, Md.⬚: 1950)*, 193(1), pp.130–
138.

Ibrahim, A.S. et al., 2008. Bacterial endosymbiosis is widely present among zygomycetes but
does not contribute to the pathogenesis of mucormycosis. *The Journal of infectious
diseases*, 198(7), pp.1083–1090.

Inglesfield et al., 2018. Robust Phagocyte Recruitment Controls the Opportunistic Fungal
Pathogen Mucor circinelloides in Innate Granulomas In Vivo. , 9(2), pp.1–20.

Kobayashi, D.Y. & Crouch, J.A., 2009. Bacterial/Fungal interactions: from pathogens to
mutualistic endosymbionts. *Annual review of phytopathology*, 47, pp.63–82.

Kraibooj, K. et al., 2014. Virulent strain of Lichtheimia corymbifera shows increased
phagocytosis by macrophages as revealed by automated microscopy image analysis.
Mycoses, 57(s3), pp.56–66.

Liu, M. et al., 2018. Comparative genome-wide analysis of extracellular small RNAs from the
mucormycosis pathogen Rhizopus delemar. *Scientific Reports*, 8(1), p.5243. Available
at: http://www.nature.com/articles/s41598-018-23611-z.

Luckett-Chastain, L. et al., 2016. IL-6 influences the balance between M1 and M2
macrophages in a mouse model of irritant contact dermatitis. *The Journal of*
Immunology, 196(1 Supplement), p.196.17--196.17. Available at:
http://www.jimmunol.org/content/196/1_Supplement/196.17.

Ma, L.J. et al., 2009. Genomic analysis of the basal lineage fungus Rhizopus oryzae reveals a
whole-genome duplication. *PLoS Genetics*, 5(7).

Martinez, F.O. et al., 2013. Genetic programs expressed in resting and IL-4 alternatively
activated mouse and human macrophages: similarities and differences. *Blood*, 121(9),
pp.e57-69.

Mendoza, L. et al., 2014. Human Fungal Pathogens of Mucorales and Entomophthorales.
Cold Spring Harbor perspectives in medicine, 10(1101). Available at:
http://perspectivesinmedicine.cshlp.org/.

Mondo, S.J. et al., 2017. Bacterial endosymbionts influence host sexuality and reveal
reproductive genes of early divergent fungi. *Nature Communications*, 8(1), p.1843.
Available at: https://doi.org/10.1038/s41467-017-02052-8.

Municio, C. et al., 2013. The Response of Human Macrophages to β-Glucans Depends on the
Inflammatory Milieu. *PLoS ONE*, 8(4).

Muñoz, J.F. et al., 2018. Coordinated host-pathogen transcriptional dynamics revealed using
sorted subpopulations and single, Candida albicans infected macrophages. *bioRxiv*,
p.350322. Available at: http://biorxiv.org/content/early/2018/06/19/350322.abstract.

Myers, J.T., Tsang, A.W. & Swanson, J.A., 2010. Activation of Murine Macrophages. *Current*
Protocols Immunology, 171(10), pp.5447–5453.

Owen, J.L. & Mohamadzadeh, M., 2013. Macrophages and chemokines as mediators of
angiogenesis. *Frontiers in Physiology*, 4 JUL(July), pp.1–8.

Petraitis, V. et al., 2013. Increased virulence of Cunninghamella bertholletiae in
experimental pulmonary mucormycosis: correlation with circulating molecular
biomarkers, sporangiospore germination and hyphal metabolism. *Medical mycology*,
51(1), pp.72–82.

Raggi, F. et al., 2017. Regulation of human Macrophage M1-M2 Polarization Balance by
hypoxia and the Triggering receptor expressed on Myeloid cells-1. *Frontiers in*
Immunology, 8(SEP), pp.1–18.

Robinson, M.D., McCarthy, D.J. & Smyth, G.K., 2009. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1),
pp.139–140.

Rogers, H. et al., 2013. Role of bacterial lipopolysaccharide in enhancing host immune
response to candida albicans. *Clinical and Developmental Immunology*, 2013.

Schmidt, S. et al., 2013. Immunobiology Rhizopus oryzae hyphae are damaged by human

natural killer (NK) cells , but suppress NK cell mediated immunity. *Immunobiology*,
218(7), pp.939–944. Available at: http://dx.doi.org/10.1016/j.imbio.2012.10.013.

Schroeder, A. et al., 2006. The RIN: an RNA integrity number for assigning integrity values to
RNA   measurements.   *BMC   molecular   biology*,   7(1),   p.3.   Available   at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1413964&tool=pmcentrez
&rendertype=abstract.

Schwartze, V.U. et al., 2014. Gene Expansion Shapes Genome Architecture in the Human
Pathogen Lichtheimia corymbifera: An Evolutionary Genomics Analysis in the Ancient
Terrestrial Mucorales (Mucoromycotina). *PLoS Genetics*, 10(8).

Sephton-Clark, P.C.S. et al., 2018. Pathways of Pathogenicity: Transcriptional Stages of
Germination in the Fatal Fungal Pathogen Rhizopus delemar. *mSphere*, 3(5), pp.1–16.

Spellberg, B., Edwards, J. & Ibrahim, A., 2016. Novel Perspectives on Mucormycosis:
Pathophysiology , Presentation , and Management. , 18(3), pp.556–569.

Szulzewsky, F. et al., 2015. Glioma-associated microglia/macrophages display an expression
profile different from M1 and M2 polarization and highly express Gpnmb and Spp1.
PLoS ONE, 10(2), pp.1–27.

Technologies,   S.,   Accutase   Directions   For   Use.   Available   at:
https://cdn.stemcell.com/media/files/pis/29637-
PIS_1_0_2.pdf?_ga=2.10286005.1436903603.1548350382-38118485.1548350382.

van Tol, S. et al., 2017. The TRIMendous Role of TRIMs in Virus–Host Interactions. *Vaccines*,
5(3), p.23. Available at: http://www.mdpi.com/2076-393X/5/3/23.

Warris, A. et al., 2005. Cytokine responses and regulation of interferon-gamma release by
human mononuclear cells to Aspergillus fumigatus and other filamentous fungi. ,
(November), pp.613–621.

Ye, R.D. & Sun, L., 2015. Emerging functions of serum amyloid A in inflammation. *Journal of
Leukocyte   Biology*,   98(6),   pp.923–929.   Available   at:
http://www.jleukbio.org/cgi/doi/10.1189/jlb.3VMR0315-080R.

Figure legends

651
Figure 1 Phenotypic and genomic comparisons. A) Experimental setup: Pre-swollen (Sab)
WT or cured fungal spores from *Rhizopus delemar* (2hr) and *Rhizopus microsporus* (4hr) in
mid-isotropic phase were co-cultured with J774.1 murine macrophages (sfDMEM) at 5:1
MOI for 1hr, washed to remove unengulfed spores, then co-incubated for a further 2 hr and
RNA for fungi and host cells harvested. Pre-swollen spores (Sab) incubated in sfDMEM for
3hr served as the negative control. B) Germination percentage over time for *R. delemar* and
R. microsporus grown in Sab. C) Spore sizes for *R. delemar* and *R. microsporus* grown in SAB.
D) Pfam terms enriched (corrected P < 0.01) in *R. delemar* vs *R. microsporus* genomes, with

colour indicating the extent of enrichment (Log_{10} Count).

Figure 2 Principal component analysis of fungal genes differentially expressed across all
samples. A) *R. delemar* wt and cured, macrophage engulfed or sfDMEM control B) *R.*
microsporus wt and cured, macrophage engulfed or sfDMEM control. Replicates (n=3) are
shown for each sample.

Figure 3 Clustering of fungal transcriptional changes. A) Heatmap displaying differentially
expressed genes in *R. delemar*. Expression levels are plotted in Log2, space and mean-
centered (FDR < 0.001) B) Heatmap displaying differentially expressed genes in *R.*
microsporus. Expression levels are plotted in Log2, space and mean-centered (FDR < 0.001)

Figure 4 Differential expression of fungal genes in compared conditions. A) The number of
genes significantly differentially expressed (multiple corrected P value < 0.05) between
samples. Blue bars indicate genes with an increase in expression (LogFC > 2), whilst orange
bars indicate genes with a decrease in expression (LogFC < -2).

Figure 5 Gene functions of genes differentially expressed in *R. delemar*. Enriched KEGG
categories for the up/down regulated genes over sample comparisons. The enrichment of
the category is indicated by the colour bar. White corresponds to no enrichment, and yellow
to red corresponds to the given P value of the enrichment.

Figure 6 Principal component analysis of macrophage genes differentially expressed
across all samples. Single cell sequencing was performed on uninfected and infected
macrophages. Transcriptional data from the experiment was analysed with the 10X
genomics analysis pipeline, and aggregated prior to principle component analysis.

Figure 7 Clustering of macrophage transcription. Heatmap displaying immune response
genes significantly differentially expressed between macrophage populations. Expression
levels are plotted in Log2 (FDR < 0.001)

Figure 8 Macrophage survival following exposure to *R. delemar* and *R. microsporus* spores.
Macrophages +/- LPS pre-treatment were infected (MOI 5:1) with fungal spores, pre-swollen
in SAB consistent with single cell experiments. Macrophages were infected with fungal
spores that were pre-treated with +/- Nikkomycin Z (24 μg/ml; n=3 for each sample).
Macrophage survival was determined 7 hr post infection. Significant differences between
samples is indicated by (*= p < 0.01).

Supplementary Figure 1 Table of Pfam terms found to be enriched (corrected P < 5 x 10^{-8}) in
R. delemar and *R. microsporus* upon genome comparison.

Supplementary Figure 2 Clustering of fungal transcription with GO annotation. Heatmap
displaying all differentially expressed genes, across all conditions, in *R. microsporus*.
Expression levels are plotted in Log2, space and mean-centered (FDR < 0.001).

Supplementary Figure 3 Clustering and heterogeneity in single cell analysis of J774.1
macrophages. Single cell plot generated by loupe cell browser. Colors indicate experimental
conditions as indicated. Spacial distribution indicates relative similarity across all detected
transcripts. 1082 single cells represented.

Supplementary Figure 4 Expression of chitin synthase genes. A) Heatmap showing the
expression of chitin synthase genes in *R. delemar* B) Heatmap showing the expression of
chitin synthase genes in *R. microsporus*

Supplementary Figure 5 Chitin synthase inhibition of *R. delemar* and *R. microsporus*
germination. *R. delemar* and *R. microsporus* treated with Nikkomycin Z in SAB at indicated
concentrations for the indicated times. Fluorescence indicates calcofluor white staining, and
thus the availability of chitin/chitosan in the cell wall. Labels indicate time and
concentration of inhibitor.

Supplemental Table 1 Selected genes differentially regulated in pairwise comparisons, with
Pfam annotations provided where available.

Supplementary Table 2 Table displaying the LogFC values for a subset of genes displayed in
Figure 7.