Stanford Accelerated Intelligent Neuromodulation Therapy for Treatment-Resistant Depression (SAINT-TRD)

Eleanor J. Cole[#], Katy H. Stimpson[#], Brandon S. Bentzley, Merve Gulser, Kirsten Cherian, Claudia Tischler, Romina Nejad, Heather Pankow, Elizabeth Choi, Haley Aaron, Flint M. Espil[1], Jaspreet Pannu, Xiaoqian Xiao, Dalton Duvio, Hugh B. Solvason, Jessica Hawkins, Austin Guerra, Booil Jo, Kristin S. Raj, Charles Debattista, Jennifer Keller, Alan F. Schatzberg, Keith D. Sudheimer[&*], Nolan R. Williams[&*]

[#]Co-first authors
[*]Corresponding Authors
[&]Co-Senior Authors

Abstract

Background: Current treatments for depression are limited by suboptimal efficacy, delayed response, and frequent side effects. Intermittent theta-burst stimulation (iTBS) is a non-invasive brain stimulation treatment which is FDA-approved for treatment-resistant depression. Recent studies suggest several improvements could be made to iTBS by 1) precision targeting of the left dorsolateral prefrontal cortex (L-DLPFC) to subgenual anterior cingulate cortex (sgACC) circuit, 2) treating with multiple sessions per day at spaced intervals and 3) applying a higher overall pulse dose of stimulation.

Objective: Examine the feasibility, tolerability, and preliminary efficacy of an accelerated, high-dose, fcMRI-guided iTBS protocol for treatment-resistant depression (TRD) termed 'Stanford Accelerated Intelligent Neuromodulation Therapy (SAINT)'.

Methods: Thirty-one participants with TRD received open-label SAINT. Resting-state functional connectivity MRI (fcMRI) was used to individually target the region of L-DLPFC most anti-correlated with sgACC. Fifty iTBS sessions (1800 pulses per session, 50-minute inter-session interval) were delivered as 10 daily sessions over 5 consecutive days at 90% resting motor threshold (adjusted for cortical depth). Neuropsychological testing was conducted before and after SAINT.

Results: Twenty-eight of 31 participants (90.32%) met criteria for remission (≤10 on the MADRS) and all 31 were remitted on measures of suicidal ideation. Neuropsychological testing demonstrated no negative cognitive side-effects. There were no seizures or other severe adverse events.

Discussion: Our highly accelerated, high-dose, iTBS protocol with fcMRI-guided targeting (SAINT) was well tolerated and safe. Efficacy was strikingly high, especially for this treatment-resistant population. Double-blinded sham-controlled trials are required to confirm the high remission rate found in this initial study.

Introduction
Depression is the leading cause of disability worldwide, and approximately 800,000 suicides are completed each year [1,2]. Current FDA-approved antidepressant treatments do not achieve remission in the majority of patients with treatment-resistant depression (TRD) [3–5], are limited by tolerability [6] and have extended treatment durations, which do not match the imminent risk to suicidal patients [7–9]. New antidepressant treatments are needed that are safe, tolerable, rapid-acting, durable, and more effective than current interventions.

Repetitive transcranial magnetic stimulation (rTMS) delivered to the left dorsolateral prefrontal cortex (L-DLPFC) is an FDA-approved non-invasive brain stimulation technique for TRD [10]. rTMS involves passing an electrical current through a magnetic coil placed on the scalp, producing a high-intensity magnetic field that passes through the scalp, skull and meninges and excites neuronal tissue [11]. Repeated high-frequency excitation of the same brain region results in successive strengthening of synapses through a process known as long-term potentiation (LTP) [12,13] causing lasting changes in functional connectivity [12,14]. The antidepressant mechanism of rTMS is hypothesized to be the result of strengthened indirect inhibitory functional connectivity from L-DLPFC to sgACC [14,15].

A more efficient form of rTMS, known as intermittent theta-burst stimulation (iTBS), has been developed, which has significantly shortened the duration of treatment sessions from 37 minutes to 3 minutes [16] and produces equivalent antidepressant responses [17,18]. FDA-approved rTMS and iTBS courses involve daily stimulation sessions for six weeks, achieving remission in 32% of patients and response in 49%, with an open-label, non-inferiority design [17]. Studies suggest that the efficacy of iTBS could be improved by accelerated delivery [19–21], higher overall stimulation doses [9,22,23] and individualized targeting [14,24].

This study aimed to examine the feasibility, safety and preliminary efficacy of an accelerated, high-dose iTBS protocol using functional connectivity magnetic resonance imaging (fcMRI)-guided targeting. Five consecutive days of 10-daily iTBS sessions (1800 pulses per session) were delivered to the region of the L-DLPFC that was most anti-correlated to the sgACC in each individual [25]. Individualized functional connectivity-guided targeting, accelerated delivery and high-dose stimulation were predicted to collectively result in higher response and remission rates than FDA-approved TMS protocols. This protocol was termed 'Stanford Accelerated Intelligent Neuromodulation Therapy (SAINT)' to distinguish this protocol from other attempts at accelerated iTBS delivery without individualized targeting and/or high dose [26,27]. We recently published a smaller series demonstrating efficacy in a cohort with extremely severe and treatment refractory depression [25].

Methods

Participants
Thirty-three participants with TRD (aged 19-78, 20 female) were recruited for this study. Twenty-four participants had a diagnosis of Major Depressive Disorder (MDD), six participants had a diagnosis of bipolar disorder (two bipolar 1) and three participants had depression in Parkinson's Disease (DPD). Diagnoses of depression or bipolar disorder were confirmed by the study psychiatrist who performed the Mini International Neuropsychiatric Interview (MINI).

Participants were required to have a Hamilton Depression Rating Scale 17-item (HDRS-17) score higher than 20 and not have responded to at least one antidepressant medication

(minimum trial duration of six weeks) to be eligible for the study. Urine samples were collected to screen for drug use and pregnancy. Participants were recruited through the Depression Research Clinic at Stanford University, Facebook advertising and clinic referrals. All participants provided written informed consent and the study was approved by the Stanford University Institutional Review Board.

One participant was excluded for having a motor threshold that was too high (>90% machine output) and one participant with a history of high anxiety dropped out after the first day of stimulation due to high anxiety levels (Young Mania Rating Scale (YMRS) and HDRS-17 scores had not increased from baseline). This resulted in a final participant sample of 31 participants (aged 19-78, 19 female). See Table 1 for participant demographic information and treatment history. Participants were required to maintain the same antidepressant regimen throughout the study enrollment (see Supplementary Table 7 for information regarding the medication participants were taking during study enrollment).

Table 1: Demographic information and treatment history for all participants (n=31)

Participant info	Mean (SD)
Gender (male:female)	12:19
Age	48.10 (17.06)
Age of onset of depression	22.45 (9.88)
Duration of depression	25.65 (15.86)
Number of antidepressant failures (lifetime)[1]	8.84 (6.03)
Number of adjunctive medications (lifetime)[2]	1.48 (1.39)
Number of participants attempted rTMS	14*
Number of participants attempted ECT	7**
Maudsley Staging Method Score	11.16 (2.04)

[1]Adequate antidepressant trials defined as a minimum of six weeks
[2]Medications defined as adequate augmentation strategies according to the Anti-depressant Treatment History Form (ATHF).
*One remitter, all other participants did not respond.
**One remitted to bilateral but did not respond to unilateral, all other participants were non-responders.

Functional magnetic resonance imaging (fMRI)
Before the stimulation course, each participant had both structural magnetic resonance imaging (MRI) and resting-state functional MRI (fMRI) scans. All participants were screened for MRI safety prior to any scanning procedures. All MRI scans were acquired using a 3T GE Discovery MR750 scanner with a 32-channel imaging coil at the Center for Cognitive and Neurobiological Imaging at Stanford, using a 3x accelerated multiband imaging sequence (TR=2 seconds). The resting state scan consisted of a single 8-minute eyes-open resting scan.

Two participants did not take part in the MRI portion of the study as one participant had a BMI which was above the limit of the scanner (BMI >35) and one participant had a spinal stimulator. For these participants, their heads were registered to the MNI brain and anatomical coordinates

previously shown to result in superior clinical outcomes compared to the standard 5cm rule were used for coil placement (–46, 45, 38) [82]. This target has also shown to be more anti-correlated to the sgACC than the area identified by the 5cm rule [14].

<u>Stanford Accelerated Intelligent Neuromodulation Therapy</u>
A Magventure Magpro X100 (MagVenture A/S, Denmark) system was used to deliver sessions of iTBS; 60 cycles of 10 bursts of 3 pulses at 50Hz were delivered in 2 second trains (5Hz) with an 8 second inter-train interval. Stimulation sessions were delivered hourly [19–21]. Ten sessions were applied per day (18,000 pulses/day) for five consecutive days (90,000 pulses in total). See Figure 2. Stimulation was delivered at 90% resting motor threshold [83,84]. A depth correction [85] was applied to the resting motor threshold to adjust for difference in the cortical depth of the individual's functional target compared to the primary motor cortex in order to consistently achieve 90% rMT in the intended functional target but stimulation was never delivered above 120% rMT for safety. The Localite Neuronavigation System (Localite GmbH, Sankt Augustin, Germany) was used to position the TMS coil over the individualized stimulation target. Participants in full remission prior to the end of the five-day stimulation course, were given the option to either finish early or complete the full course. One participant with a diagnosis of Bipolar type 1 finished treatment after the first day and one participant with Parkinson's finished treatment after 3 days. All other participants completed the full five-day treatment course.

<u>Figure 1</u>

Figure 1: This figure illustrates the three factors which make SAINT different from current FDA-approved iTBS protocols: Precise targeting (A), accelerated delivery (B) and high pulse dose [both per session (B) and overall (C)]. A) Stimulation was delivered to the region of the L-DLPFC (1) which was most anti-correlated to the sgACC (2) due to previous neuroimaging findings which suggest the higher the anti-correlation between the stimulated region of the L-DLPFC and the sgACC, the better the clinical outcome [14,24,44]. B) Unlike, FDA-approved protocols which only deliver one iTBS session per day, our SAINT protocol delivered 10 sessions per day. iTBS sessions were delivered 50 minutes apart because previous research has shown that iTBS sessions delivered 50-90 minutes apart have a cumulative effect [19–21]. In comparison, iTBS sessions delivered only 40 minutes apart or less, have shown not have a cumulative effect [19–21,40,42]. 1800 pulses per session was chosen as this is the only pulse dose that has been explored in a blinded iTBS trial [49]. Additionally, 1800 pulses has been shown to produce long-lasting changes in cortical excitability [100] and optimally produce intended cellular changes [101]. C) Our study administered the highest overall pulse-dose of any published study to date. A dose-response curve has never been constructed for clinical TMS but a previous study showed 61% of non-responders to a standard 6-week TMS course responded with further treatment [23] suggesting that FDA-approved protocols are under-dosing (the purple dotted line provides an example of where the FDA-approved dose could be on the curve). Our SAINT protocol

delivered 5x the FDA-approved dose. See supplementary material for more information regarding protocol development.

Clinical assessments

Prior to receiving any stimulation, participants' depressive symptoms and suicidal ideation were assessed using the Hamilton Depression Rating Scale-17 item (HDRS-17), Montgomery-Asberg Depression Rating Scale (MADRS) and Columbia-Suicide Severity Rating Scale (C-SSRS, suicidal ideation subscale). A self-report measure was also used; the Beck Depression Inventory II (BDI-II). The degree of treatment-resistance was calculated using the Maudsley Staging Method [86], and symptoms of mania were screened for using the Young Mania Rating Scale (YMRS). At the end of each day of stimulation (ten sessions), depressive symptoms were assessed using the Hamilton Depression Rating Scale 6-item (HDRS-6); participants also completed the BDI-II. The YMRS was completed daily to ensure hypomania had not been induced in any participants [87–89]. The same clinical assessments used at baseline were conducted approximately 72 hours after the stimulation course (immediate follow-up) as well as two weeks and one month after the stimulation course in order to assess the durability of the response.

A neuropsychological test battery was administered before SAINT and approximately 72 hours after the SAINT course in order to capture any neurocognitive side effects. The Hopkins Verbal Learning Test – Revised (HVLT-R) [90], the Brief Visuospatial Memory Test – Revised (BVMT-R) [91], subtests from the Wechsler Adult Intelligence Scale (4th Ed.; WAIS-IV) [92] and various tests from the Delis Kaplan Executive Function System (D-KEFS) [93] were used. These neuropsychological tests have been shown to have good psychometric properties [90,91,93,94] and assessed verbal learning and memory, visuospatial learning and memory, working memory, attention, cognitive inhibition, cognitive switching, problem solving, and verbal fluency. See supplementary material for detailed information about the neuropsychological test battery.

fMRI analysis for target generation

Personalized L-DLPFC targets were generated for each participant using the resting-state scan. All analyses were conducted in a participant's own brain space (i.e., not warped to standardized-brain space). Resting-state scans were pre-processed according to typical methods using Statistical Parametric Mapping (SPM12) software. The resting-state scans were motion corrected and resliced. The T1 weighted structural scan was then co-registered with the resting-state scans. The resting-state scans were then spatially smoothed with a 3mm Gaussian kernel, detrended using a linear model of the global signal [95], and band-pass filtered to preserve the typical resting-state frequencies (0.1Hz-0.01Hz). Next, the estimation parameters to warp the T1 weighted structural image into Montreal Neurological Institute (MNI) space was calculated using SPM segmentations based on tissue probability maps. These normalization parameters were inverted and applied to MNI space regions of interests (ROIs) for the L-DLPFC (Brodmann area 46) and the sgACC (BA25). This inverse normalization served to map the MNI space ROIs onto the individual participant's brain. The participant-space ROIs were then resliced, smoothed, and binarized to match the dimensions of the resting state scans.

The participant-space ROI for the L-DLPFC formed the search area for the optimal TMS coil placement. Two separate algorithms were used to determine coil placement. The first algorithm sorted each of the DLPFC and sgACC voxels into functional sub-units using a hierarchical agglomerative clustering algorithm. Median time series were then created for each functional subunit and the correlation coefficients were calculated between all median time series

extracted from all functional subunits of the L-DLPFC and sgACC. The second algorithm determined the optimal L-DLPFC subunit to target based on three factors: the net correlation/anti-correlation of the L-DLPFC subunit with sgACC subunits, the size of the subunit and the spatial concentration of the subunit. See supplementary methods for more details on these algorithms. 3D maps of the whole brain correlation coefficient of the selected L-DLPFC subunit were then created and used to target the coil placement using the Localite TMS Navigation software (Localite GmbH, Sankt Augustin, Germany).

Clinical outcome analysis
All statistical analyses were conducted using IBM SPSS Version 22. The level of statistical significance was set at $p = 0.05$. Missing data were not imputed for any statistical analysis. Statistical analyses were planned independently by BSB and reviewed by EJC, NWR, BJ and AFS. These analyses were then carried out by BSB and EJC.

Our primary outcome measure was change in MADRS scores from baseline to immediately following SAINT, and MADRS scores were used to calculate response and remission rates. Reductions in HDRS-17, HDRS-6 and BDI-II scores were used as secondary outcome measures of depression severity. Response was defined as ≥50% reduction in these scales. Remission was defined as MADRS score ≤10 [96], ≤7 on HDRS-17 [97] ≤4 on the HDRS-6 [98], BDI≤12 [99]. A floor effect of SAINT treatment was observed across all scales, and initial linear mixed models produced residuals that were not normally distributed. Thus, changes in MADRS, HDRS-17, and BDI-II were assessed with generalized linear mixed models (GLMM) that used a compound symmetry covariance structure, Satterthwaite approximation of degrees of freedom, and robust estimation of coefficients to handle violations of model assumptions. Fixed effects of time, diagnosis (MDD vs. non-MDD), and prior non-response to conventional rTMS were assessed, as well as time x diagnosis and time x response interactions. All post-hoc pairwise comparisons were Bonferroni-corrected.

Eleven participants were re-treated once their symptoms returned to baseline and again met study entry criteria. A similar GLMM was used to assess whether SAINT re-treatment was able to produce equivalent antidepressant responses. Fixed effects of time, treatment course, and their interaction were assessed.

Daily HDRS-6 scores were used to calculate the number of days of stimulation required to reach responder criterion (<50% HDRS-6) and remission criterion (<5 on HDRS-6). Kaplan-Meier survival analysis using the Breslow test of equality of survival distributions was used to determine if there were significant differences in the number of days to reach response and remission criteria for participants who had a history of conventional rTMS non-response compared to those who did not.

Suicidality was assessed using the Columbia Suicide Severity Rating Scale (C-SSRS, suicidal ideation subscale), item 3 of the HDRS-17, and item 10 of the MADRS. Response was defined as ≥50% reduction in these scores from baseline and remission was defined as a score of 0. Response was only calculated if the baseline score was >0. Remission was defined as a score of 0 for each of these scales. Scores were ordinal and changes in scores were assessed with generalized linear models with a multinomial link, compound symmetry covariance structure, Satterthwaite approximation of degrees of freedom, and robust estimation of coefficients to handle violations of model assumptions.

Scores on the neuropsychological tests before and after SAINT were compared using paired t-tests. The data for the Hopkins Verbal Learning Test (HVLT) total score, HVLT Delayed Recall score, the Brief Visuospatial Memory Test (BVMT)-Delayed Recall and the number of rule violations on the D-KEFS Tower Test violated the assumption of normality so non-parametric (Wilcoxon signed-rank) tests were used to evaluate SAINT-induced changes in performance on these three tasks. Neuropsychological test data were available for 22 participants.

Results

Safety
No serious adverse events occurred. The only side-effects participants reported were fatigue and some discomfort at both the stimulation site and in the facial muscles during stimulation. The neuropsychological test battery showed no negative cognitive side-effects following SAINT. Performance significantly improved on measures of cognitive inhibition [D-KEFS color-word inhibition task; $t(20)=2.98$, $p=.007$, $d=.65$], cognitive switching [DKEFS trail making test condition 4, Alternating Letter-Number Switching; $t(19)=2.50$, $p=.02$, $d=.56$], problem solving [DKEFS Tower Test Total Achievement Score; $t(20)=-2.70$, $p=.01$, $d=.59$]; and rule violations on the D-KEFS Tower Test; $z=-2.68$, $p=.007$, $r=.4$]. Due to the large number of comparisons made, the significant improvements in performance observed did not survive the correction for multiple comparisons (Bonferroni-corrected significance level $p<.004$). There were no significant changes on any of the other neurocognitive tasks, see supplementary material.

Depression symptoms
Depressive symptoms decreased from baseline immediately following SAINT with an 87.66% reduction in the mean Montgomery-Asberg Depression Rating Scale (MADRS) score from 37.68 (SD=7.24) to 4.65 (SD=6.46). Generalized linear mixed-model (GLMM) analysis revealed a significant effect of time ($F_{3,27}=106.742$, $P<.001$) on mean MADRS scores with all follow-up time points being significantly lower than baseline (Bonferroni-corrected pairwise comparisons, $P<.001$). These results were recapitulated for the Hamilton Depression Rating Scale (HDRS-17, $F_{3,26}=124.128$, $P<.001$) and the Beck Depression Inventory (BDI, $F_{3,29}=33.692$, $P<.001$). The response rate (≥50% reduction in MADRS) was 90.32%, and all responders were in remission (MADRS score ≤10). Results were similar across all clinical assessments (see Table 2). One month following SAINT, 59% of participants continued to meet responder criteria (see Supplementary Table 1 for response rates at one month).

A major hypothesis of SAINT is that conventional rTMS delivers insufficient cumulative stimulation to induce response and remission from depression for some patients. We tested this by including participants who had a history of non-response to conventional rTMS (rTMS non-responders, n=13). We found that conventional rTMS non-responders had significantly greater MADRS scores (GLMM, responder, $F_{1,14}=15.875$, $P=.001$; responder x time, $F_{3,35}=3.837$, $P=.018$) at all time points except baseline (Bonferroni-corrected pairwise comparisons, $P<.001$) compared to the rest of the participant sample (see Figure 2 and Supplementary Tables 5 and 6).

Table 2: Clinical assessment scores for all participants immediately following SAINT; mean (SD, n) or % (n)

	Pre-SAINT	Post-SAINT	Responders (%)[1]	Remission (%)[2]
MADRS	37.68 (7.24, 31)	4.65 (6.46, 31)	90.32 (31)	90.32 (31)
HDRS-17	27.84 (5.19, 31)	4.48 (5.09, 31)	90.32 (31)	83.87 (31)
HDRS-6	15.10 (2.77, 31)	2.42 (3.30, 31)	87.10 (31)	83.87 (31)
BDI-II	31.96 (12.42, 25)	6.38 (7.08, 21)	89.47 (19)	85.71 (21)
C-SSRS[3]	1.38 (1.06, 26)	0.00 (0.00, 23)	100.00 (18)	100.00 (23)
HDRS-Q3	1.45 (0.77, 31)	0.03 (0.18, 31)	96.42 (28)	96.77 (31)
MADRS-Q10	2.42 (0.99, 31)	0.06 (0.36, 31)	96.67 (30)	96.77 (31)

[1]Response defined as ≥50% reduction in score.
[2]Remission defined ≤7 on HDRS-17 [97] ≤4 on the HDRS-6 [98], ≤10 on MADRS [96], BDI≤12 [99] and C-SSRS=0 [102].
[3]Suicidal ideation subscale
MADRS: Montgomery-Asberg Depression Rating Scale, HDRS-17: Hamilton Depression Rating Scale-17 item, HDRS-6: Hamilton Depression Rating Scale-6 item, BDI-II: Beck Depression Inventory II, C-SSRS: Columbia-Suicide Severity Rating Scale.
Analyses limited to participants with a diagnosis of Major Depressive Disorder without Parkinson's disease (MDD, n=22) were performed to make comparisons to pivotal iTBS and TMS trials [28–30]. See Supplementary Table 2 for demographic information for this subsample. GLMM analysis showed no significant effect of diagnosis ($F_{1,13}$=0.027, P=.873) on mean MADRS scores at any time point (diagnosis x time, $F_{3,28}$=2.509, P=.80). The overall response and remission rates were similar to this subsample with MDD, see Supplementary Tables 3 and 4 for response and remission rates for this subsample immediately and one month after SAINT respectively.

Days of treatment until response and remission
The mean number of days of SAINT until participants met the response criterion (≥50% reduction in HDRS-6 score) was 2.46 (SD=1.26, ~25 10-min treatments, n=28, daily HDRS-6 scores missing for 2 participants, 1 participant did not respond by the HDRS-6 criterion), and the mean number of days to achieve remission (HDRS-6 score <5) was 2.96 (SD=1.48, ~30 10-min treatments, *n*=26, 3 participants did not achieve remission by the HDRS-6 criterion). Similar time courses were seen for participants with MDD. See Figure 3 for percentage change in HDRS-6 score with each day of stimulation, for all participants (2A) and participants with MDD (2B).

Kaplan-Meier survival analysis revealed that participants who had previously not responded to a 6-week rTMS treatment course (n=13) required more days of treatment to achieve responder criterion [X^2=8.047, P=.005, mean 3.08 days (SD=0.67), ~31 treatments] and remission criterion [X^2=7.894, P=.005, mean 3.70 days, (SD=.95), ~37 treatments). See Figure 4. None of the rTMS non-responders met responder criterion after the first day of aiTBS; notably, the first day of SAINT provides an equivalent amount of stimulation as six weeks of FDA-approved iTBS.

Figure 2: Percentage change in MADRS score immediately following the five-day stimulation course, two weeks and four weeks afterwards for TMS non-responders (red, n=13) and all other participants (blue, n=18). rTMS non-responses showed significantly lower percentage reduction in MADRS score at all timepoints after SAINT than the rest of the participant sample: (Immediate, n=18/13, p<0.05; 2 weeks, n=18/13 p<0.01; 4 weeks, n=17/12 p<0.01)

Figure 3

Figure 3: Mean percentage change in Hamilton Depression Rating Scale score (6-item version) with each day of stimulation for A) All participants (n=29, daily HDRS-6 scores missing for two participants), B) MDD participants (n=21, daily HDRS-6 scores missing for one participant). Dotted lines indicate responder criteria and error bars represent standard error.

Figure 4

Figure 4: Mean percentage change in Hamilton Depression Rating Scale score (6-item version) with each day of stimulation for A) All TMS non-responders (n=13), B) All other participants (n=16), C) MDD TMS non-responders (n= 10), D) All other MDD participants (n=11). Dotted lines indicate responder criteria and error bars represent standard error.

Re-treatment efficacy
Eleven participants were re-treated after they no longer met responder criterion and again met study entry criteria (mean time between treatments was 20.27 weeks, SD=14.07). There were no significant differences in MADRS scores between treatment courses (GLMM, $F_{1,155}$=0.192, P=0.662), nor was there a treatment x time interaction (GLMM, $F_{3,155}$=0.607, P=0.611). Baseline and follow up MADRS scores were all similar between treatment courses (Bonferroni-corrected

pairwise comparisons, P>0.05). See Table 3 for percentage change in MADRS score for initial SAINT and re-treatment for each participant.

Table 3: Percentage change in MADRS score for initial and re-treatment

Participant	Diagnosis	Initial % change MADRS	Re-treatment % change MADRS
3	MDD	91.43	85.71
4	MDD	97.67	100.00
11	BPD	100.00	86.11
14	PD	97.50	88.89
19	MDD	100.00	100.00
21	MDD	83.64	84.44
25	BPD	100.00	100.00
28	MDD	44.12	76.47
31	BPD	100.00	100.00
34	MDD	100.00	97.44
38	MDD	90.91	95.45
Mean (SD)		91.39 (16.56)	92.23 (8.24)

Suicidality scales

Changes in suicidality scales were assessed with generalized linear modeling (GLM) with a multinomial link. Following SAINT there were significant reductions in the Columbia-Suicide Severity Rating Scale [(C-SSRS, suicidal ideation subscale) (GLM, $X^2(2)=17.086$, $P<0.001$)], item 3 of the HDRS-17 (GLM, $X^2(3)=48.976$, $P<0.001$), and item 10 of the MADRS (GLM, $X^2(3)=59.133$, $P<0.001$) at all follow up time points (X^2, $P<0.001$). All participants remitted on these measures of suicidal ideation (see Table 2). One month following SAINT, 95% of participants continued to be remitted (see Supplementary Tables 3 and 4).

Discussion

The aim of this study was to examine the safety, feasibility and preliminary efficacy of an accelerated high-dose iTBS, fcMRI-guided treatment protocol (SAINT) for treatment-resistant depression (TRD). We found that SAINT significantly reduced depressive symptoms and suicidal ideation in patients with severe TRD within 5 days without negative cognitive side-effects or other adverse events. The 90% remission rate we observed is substantially higher than remission rates for open-label studies using standard FDA-approved rTMS treatment protocols [17,31,32]. This is also higher than what has been reported for electroconvulsive therapy (ECT) treatment of TRD (~48%[4]) and for ketamine treatment of TRD (14.3-66.6%[33–35]). This remission rate was observed despite the inclusion of participants who had previously failed FDA-approved rTMS and/or ECT treatments. The potential high efficacy of SAINT may be due to the combination of the accelerated delivery of iTBS sessions, the high-dose of stimulation and the individualized targeting method used.

Although our initial observed response and remission rates are atypically high compared to other studies that have tested rTMS protocols for treatment of depression (rTMS/iTBS/cTBS), similar response rates have been found in deep brain stimulation trials for depression and Parkinson's Disease [36,37]. The high efficacy of our SAINT protocol complements evidence from

basic neuroscience research and human physiology data, which suggest that multiple spaced daily iTBS sessions have an enhanced effect compared to the same number of single daily sessions [19–21,38,39]. Studies that have applied theta-burst protocols (cTBS/iTBS) to the motor cortex in humans have shown that two spaced stimulation sessions produce greater [40] and longer lasting [39,41] changes in cortical excitability than single stimulation sessions. Rodent studies conducted using hippocampal slices have shown that multiple iTBS sessions delivered with inter-session intervals of 50-90 minutes have a cumulative effect on dendritic spine enlargement, a process involved in synaptic strengthening [19–21,38]. In comparison, iTBS sessions delivered with inter-session intervals of 40 minutes or less do not have a cumulative effect on dendritic spines[19–21]. Similarly, two iTBS sessions delivered to the prefrontal or motor cortex in humans only 15 minutes apart have shown not to increase cortical excitability more than a single iTBS session [40,42]. This could explain the limited efficacy of a previously reported accelerated iTBS protocol, which used an inter-session interval of only 15 minutes [26,27].

The individualized targeting method used in our study may have also contributed to the high response and remission rates. The L-DLPFC is a large brain area that consists of several subregions, some of which are correlated and some anti-correlated with sgACC activity [43]. Targeting rTMS to the subregions of L-DLPFC that have higher functional anti-correlation with sgACC have been shown to produce a superior antidepressant response [14,24,44]. Defining L-DLPFC using common techniques such as scalp-based measurements, structural MRI scans or fMRI activation patterns could result in stimulating a subregion of the L-DLPFC that is correlated rather than anti-correlated with the sgACC and therefore not drive the desired increase in anti-correlation between L-DLPFC and the sgACC [45]. The standard '5cm rule' scalp-based measurement has been shown to miss the L-DLPFC completely in >1/3 of cases [46]. A retrospective study conducted by Fox and colleagues found that individual differences in the degree of anti-correlation between the stimulated subregion of the L-DLPFC and the sgACC accounted for over 70% of the variability in clinical efficacy [14]. A recent interleaved TMS-fMRI study showed that stimulating the subregion of the L-DLPFC that displays the greatest degree of functional connectivity with the sgACC resulted in stimulation propagation to the sgACC in all participants [47]. In comparison, in a separate study when the L-DLPFC was defined anatomically (border of BA9/BA46), stimulation propagated to the sgACC in only 44% of participants [48]. In the aiTBS study by the Baeken group [27] and a blinded iTBS trial [49] the same anatomical target (border of BA9/BA46) was utilized, which may have contributed to the limited efficacy of these approaches. By stimulating the subregion of the L-DLPFC that is most anti-correlated with the sgACC in each individual in this study, we may have reduced this variability in signal propagation and maximized treatment efficacy.

The high efficacy of our SAINT protocol also suggests that FDA-approved protocols may be under-dosing. Our protocol administered five-times the pulse dose of the FDA-approved iTBS protocol (90,000 pulses in comparison to the standard 18,000 iTBS pulses[17]). Prior studies found that 61% of individuals who do not respond to an initial six weeks of stimulation responded with additional rTMS treatment sessions [23], and higher pulse doses are associated with higher response and remission rates [50,51]. Our SAINT protocol applied the equivalent amount of stimulation as a six-week treatment protocol each day of stimulation (18,000 pulses/day [17]). Thirty-one percent of participants in our study met responder criteria after the first day of stimulation (n=9/29, daily HDRS-6 missing for 2 participants), this response rate is equivalent to response rates found after six-weeks of daily iTBS/rTMS sessions in individuals of this treatment-resistance level [29,30,52,53]. None of the prior rTMS non-responders in our study responded after the first day of SAINT (see Figure 4). These data indicate that a non-response

to a standard six-week treatment protocol may reflect the need for a higher stimulation dose, for the majority of patients [23], which matches with recent trajectory data [54]. The apparent need for a higher number of stimulation pulses is consistent with other neuromodulation modalities, such as deep brain stimulation in Parkinson's where a mean of 500,000 pulses of stimulation are delivered to the subthalamic nucleus (STN) or the globus pallidus (GPi) each day [55]. The need for a higher overall dose of stimulation may have also contributed to the limited efficacy of a previous attempt of an accelerated iTBS protocol, which only delivered 32,400 pulses [26]. Our study administered the highest number of pulses per day and highest overall pulse-dose of any study we are currently aware of [25].

Prior rTMS non-responders in our study not only required more stimulation sessions to induce a clinically significant response but also showed less-durable responses than the rest of the participant sample. The high relapse rate at one month in this subsample is reflective of the ECT data which show general relapse rates of 62.7% at 3 months and 78% at 6 months [56] with 25% of treatment-resistant individuals relapsing after one week [57] and an average length of wellness of 8.6 weeks [58]. It is possible that depressed individuals with a higher degree of treatment-resistance display neuroplasticity impairments [59]. The higher number of stimulation sessions required to induce antidepressant responses could be due to deficits in processes involved in the early stages of LTP, such as AMPA receptor phosphorylation [59,60]. The shorter duration of antidepressant responses could be the result of dysfunction in later LTP processes that mediate long-lasting changes to synaptic strength [41]. These processes include the synthesis of new proteins and gene expression changes such as the activation of transcription factors (e.g. CREB) and the induction of plasticity-enhancing genes such as BDNF [59]. Highly treatment-resistant individuals may require maintenance iTBS therapy [61] or even implanted epidural cortical stimulators [62,63] to sustain antidepressant responses.

The short duration of our SAINT protocol, the apparent reproducibility of responses and the lack of cognitive side-effects, provide potential benefits over currently available treatments. The short duration of our SAINT protocol means SAINT could provide a rapid means of treating patients with active suicidality. Ketamine and ECT are currently available alternatives. Ketamine/s-ketamine has recently been FDA-approved for treatment-resistant depression; however, ketamine has several limitations. Approximately 11% of patients report the dissociative symptoms as very disturbing [64], remission rates for ketamine/s-ketamine are lower than the remission rate we observed for SAINT [33,35,65], very few studies have investigated the anti-depressant efficacy of ketamine beyond a single infusion [66,67] and the opioid mechanism of action poses a potential risk [68]. The other available rapid-acting treatment is ECT, for which less than 2% of eligible patients receive due to concerns regarding cognitive side-effects and stigma [69,70]. In the majority of patients, ECT also takes two weeks or longer to produce remission from suicidal ideation [71].

Our study has several limitations. It is limited by the small sample size and open-label design; a double-blind controlled trial is required to determine the efficacy of our SAINT protocol in comparison to an identical schedule of sham stimulation sessions. However, individuals with the same degree of treatment-refractoriness as the participants included in this study (severe treatment-resistance; >11 Maudsley Staging Method), have previously shown no placebo response to iTBS sessions of 1800 pulses [49]. Additionally, the most recent deep brain stimulation (DBS) sgACC trial for TRD included individuals with similar treatment-resistance levels to the patients in this report and had a response rate of only 20% [72]. A greater degree of placebo response would be expected for DBS as placebo response has generally been related

to the degree of invasiveness of the intervention [73]. Other limitations of our study include the single stimulation site [74–76], the use of fixed stimulation frequencies [54,77], fixed inter-session intervals [77,78] and the lack of state-dependent stimulation [79]. It is possible that different stimulation sites may be optimal for certain patients based on their symptom profile [74–76]. The use of individualized stimulation frequencies may result in quicker and more durable responses [77,80]. Some individuals may require slightly different inter-session intervals due to differences in cortical excitability profiles [78,81]. Finally, recent studies have shown that applying stimulation in particular brain states using real-time electroencephalography (EEG)-triggered transcranial magnetic stimulation (EEG-TMS) can increase cortical responses to stimulation [79]. Limitations regarding the clinical application of SAINT include the high cost and additional time required to use neuronavigation equipment, acquire MRI scans and process the resulting data.

In conclusion, the high-dose of stimulation, accelerated delivery and fcMRI-guided individualized targeting method are preliminarily safe, well tolerated, feasible, and associated with high rates of remission from depressive episodes. The high remission rate we observed with SAINT was found despite the inclusion of participants who had previously not responded to rTMS and/or ECT. These rates are almost double those observed in TRD individuals receiving open-label ECT [4], which is currently the 'gold standard' treatment for TRD. Our data suggest that FDA-approved TMS protocols may be under-dosing and could potentially benefit from individualized targeting methods and accelerated delivery. The efficacy of SAINT in treating suicidal ideation and the short duration of the protocol suggest SAINT could provide a means of rapidly ensuring the safety of suicidal patients. Larger, double-blinded, placebo-controlled trials are required to confirm the promisingly high response and remission rates found in this initial study.

References

1.   Reddy M. Depression: The disorder and the burden. *Indian J Psychol Med*. 2010;32(1):1-2. doi:10.4103/0253-7176.70510

2.   World Health Organisation. WHO | Suicide data. WHO Website. doi:10.1109/TSE.2011.123

3.   Berlim MT, Van Den Eynde F, Tovar-Perdomo S, Daskalakis ZJ. Response, remission and drop-out rates following high-frequency repetitive transcranial magnetic stimulation (rTMS) for treating major depression: A systematic review and meta-analysis of randomized, double-blind and sham-controlled trials. *Psychol Med*. 2014;44(2):225-239. doi:10.1017/S0033291713000512

4.   Heijnen WT, Birkenhäger TK, Wierdsma AI, Van Den Broek WW. Antidepressant pharmacotherapy failure and response to subsequent electroconvulsive therapy: A meta-analysis. *J Clin Psychopharmacol*. 2010;30(5):616-619. doi:10.1097/JCP.0b013e3181ee0f5f

5.   Sinyor M, Schaffer A, Levitt A. The Sequenced Treatment Alternatives to Relieve Depression (STAR*D) trial: A review. *Can J Psychiatry*. 2010;55(3):126-135. doi:10.1177/070674371005500303

6.   Penn E, Tracy DK. The drugs don't work? antidepressants and the current and future pharmacological management of depression. *Ther Adv Psychopharmacol*. 2012;2(5):179-188. doi:10.1177/2045125312445469

7.   Blier P. Optimal use of antidepressants: when to act? *J Psychiatry Neurosci*. 2009;34(1):80.

8.   Husain MM, Rush AJ, Fink M, et al. Speed of response and remission in major depressive disorder with acute electroconvulsive therapy (ECT): A consortium for research in ECT (CORE) report. *J Clin Psychiatry*. 2004;65(4):485-491.

9.   Perera T, George MS, Grammer G, Janicak PG, Pascual-Leone A, Wirecki TS. The Clinical TMS Society Consensus Review and Treatment Recommendations for TMS Therapy for Major Depressive Disorder. *Brain Stimul*. 2016;9(3):336-346. doi:10.1016/j.brs.2016.03.010

10.   George MS, Talyor JJ, Short EB. The Expanding Evidence Base for rTMS Treatment of Depression. *Curr Opin Psychiatry*. 2013;26(1):13-18. doi:10.1097/YCO.0b013e32835ab46d.The

11.   Taniguchi M, Cedzich C, Schramm J, et al. Transcranial magnetic stimulation and the human brain. *Neurosurgery*. 2003. doi:10.1038/35018000

12.   Huerta PT, Volpe BT. Transcranial magnetic stimulation, synaptic plasticity and network oscillations. *J Neuroeng Rehabil*. 2009;6(7). doi:10.1186/1743-0003-6-7

13.   Ogiue-Ikeda M, Kawato S, Ueno S. The effect of repetitive transcranial magnetic stimulation on long-term potentiation in rat hippocampus depends on stimulus intensity. *Brain Res*. 2003;993(1-2):222-226. doi:10.1016/j.brainres.2003.09.009

14.   Fox MD, Buckner RL, White MP, Greicius MD, Pascual-Leone A. Efficacy of transcranial magnetic stimulation targets for depression is related to intrinsic functional connectivity with the subgenual cingulate. *Biol Psychiatry*. 2012;72(7):595-603. doi:10.1016/j.biopsych.2012.04.028

15.   Kito S, Hasegawa T, Takamiya A, et al. Transcranial Magnetic Stimulation Modulates Resting EEG Functional Connectivity Between the Left Dorsolateral Prefrontal Cortex and Limbic Regions in Medicated Patients With Treatment-Resistant Depression. *J Neuropsychiatry Clin Neurosci*. 2016;29(2):155-159. doi:10.1176/appi.neuropsych.15120419

16.   Huang YZ, Rothwell JC. The effect of short-duration bursts of high-frequency, low-

intensity transcranial magnetic stimulation on the human motor cortex. *Clin Neurophysiol*. 2004;115(5):1069-1075. doi:10.1016/j.clinph.2003.12.026

17. Blumberger DM, Vila-Rodriguez F, Thorpe KE, et al. Effectiveness of theta burst versus high-frequency repetitive transcranial magnetic stimulation in patients with depression (THREE-D): a randomised non-inferiority trial. *Lancet*. 2018;391(10131):1683-1692. doi:10.1016/S0140-6736(18)30295-2

18. Blumberger DM, Vila-Rodriguez F, Dunlop K, et al. Intermittent theta-burst versus 10 Hz left dorsolateral prefrontal rTMS for treatment resistant depression: preliminary results from a two-site, randomized, single blind non-inferiority trial. *Brain Stimul*. 2015;8(2):329. doi:10.1016/j.brs.2015.01.067

19. Smolen P, Zhang Y, Byrne JH. The right time to learn: Mechanisms and optimization of spaced learning. *Nat Rev Neurosci*. 2016;17(2):77-88. doi:10.1038/nrn.2015.18

20. Kramar EA, Babayan AH, Gavin CF, et al. Synaptic evidence for the efficacy of spaced learning. *Proc Natl Acad Sci*. 2012;109(13):5121-5126. doi:10.1073/pnas.1120700109

21. Lynch G, Kramár EA, Babayan AH, Rumbaugh G, Gall CM. Differences between synaptic plasticity thresholds result in new timing rules for maximizing long-term potentiation. *Neuropharmacology*. 2013;64:27-36. doi:10.1016/j.neuropharm.2012.07.006

22. George MS, Wassermann EM, Kimbrell T a, et al. Mood improvement following daily left prefrontal repetitive transcranial magnetic stimulation in patients with depression: a placebo-controlled crossover trial. *Am J Psychiatry*. 1997;154(12):1752-1756. doi:10.1176/ajp.154.12.1752

23. Yip AG, George MS, Tendler A, Roth Y, Zangen A, Carpenter LL. 61% of unmedicated treatment resistant depression patients who did not respond to acute TMS treatment responded after four weeks of twice weekly deep TMS in the Brainsway pivotal trial. *Brain Stimul*. 2017;10(4):847-849. doi:10.1016/j.brs.2017.02.013

24. Weigand A, Horn A, Caballero R, et al. Prospective Validation That Subgenual Connectivity Predicts Antidepressant Efficacy of Transcranial Magnetic Stimulation Sites. *Biol Psychiatry*. 2018;84(1):28-37. doi:10.1016/j.biopsych.2017.10.028

25. Williams NR, Sudheimer KD, Bentzley BS, et al. High-dose spaced theta-burst TMS as a rapid-acting antidepressant in highly refractory depression. *Brain*. 2018;141(3):e18. doi:10.1093/brain/awx379

26. Desmyter S, Duprat R, Baeken C, Van Autreve S, Audenaert K, van Heeringen K. Accelerated Intermittent Theta Burst Stimulation for Suicide Risk in Therapy-Resistant Depressed Patients: A Randomized, Sham-Controlled Trial. *Front Hum Neurosci*. 2016;10:1-7. doi:10.3389/fnhum.2016.00480

27. Duprat R, Desmyter S, Rudi DR, et al. Accelerated intermittent theta burst stimulation treatment in medication-resistant major depression: A fast road to remission? *J Affect Disord*. 2016;200:6-14. doi:10.1016/j.jad.2016.04.015

28. George MS, Lisanby SH, Avery D, et al. Daily left prefrontal transcranial magnetic stimulation therapy for major depressive disorder: a sham-controlled randomized trial. *Arch Gen Psychiatry*. 2010;67(5):507-516. doi:10.1001/archgenpsychiatry.2010.46

29. Levkovitz Y, Isserles M, Padberg F, et al. Efficacy and safety of deep transcranial magnetic stimulation for major depression: A prospective multicenter randomized controlled trial. *World Psychiatry*. 2015;14(1):64-73. doi:10.1002/wps.20199

30. O'Reardon JP, Solvason HB, Janicak PG, et al. Efficacy and Safety of Transcranial Magnetic Stimulation in the Acute Treatment of Major Depression: A Multisite Randomized Controlled Trial. *Biol Psychiatry*. 2007;62(11):1208-1216. doi:10.1016/j.biopsych.2007.01.018

31. Carpenter LL, Janicak PG, Aaronson ST, et al. TRANSCRANIAL MAGNETIC

STIMULATION (TMS) FOR MAJOR DEPRESSION: A MULTISITE, NATURALISTIC, OBSERVATIONAL STUDY OF ACUTE TREATMENT OUTCOMES IN CLINICAL PRACTICE. *Depress Anxiety*. 2012;29(7):587-596. doi:10.1002/da.21969

32. Janicak PG, Nahas Z, Lisanby SH, et al. Durability of clinical benefit with transcranial magnetic stimulation (TMS) in the treatment of pharmacoresistant major depression: Assessment of relapse during a 6-month, multisite, open-label study. *Brain Stimul*. 2010;3(4):187-199. doi:10.1016/j.brs.2010.07.003

33. Diamond PR, Farmery AD, Atkinson S, et al. Ketamine infusions for treatment resistant depression: A series of 28 patients treated weekly or twice weekly in an ECT clinic. *J Psychopharmacol*. 2014;28(6):536-544. doi:10.1177/0269881114527361

34. Mathew SJ, Murrough JW, Aan Het Rot M, Collins KA, Reich DL, Charney DS. Riluzole for relapse prevention following intravenous ketamine in treatment-resistant depression: A pilot randomized, placebo-controlled continuation trial. *Int J Neuropsychopharmacol*. 2010;13(1):71-82. doi:10.1017/S1461145709000169

35. Shiroma PR, Johns B, Kuskowski M, et al. Augmentation of response and remission to serial intravenous subanesthetic ketamine in treatment resistant depression. *J Affect Disord*. 2014;155(1):123-129. doi:10.1016/j.jad.2013.10.036

36. Riva-Posse P, Choi KS, Holtzheimer PE, et al. A connectomic approach for subcallosal cingulate deep brain stimulation surgery: prospective targeting in treatment-resistant depression. *Mol Psychiatry*. 2017;23(4):843-849. doi:10.1038/mp.2017.59

37. Groiss SJ, Wojtecki L, Südmeyer M, Schnitzler A. Deep brain stimulation in Parkinson's disease. *Ther Adv Neurol Disord*. 2009;2(6):20-28. doi:10.1177/1756285609339382

38. Harris KM, Kater SB. Dendritic Spines: Cellular Specializations Imparting Both Stability and Flexibility to Synaptic Function. *Annu Rev Neurosci*. 1994;17(1):341-371. doi:10.1146/annurev.ne.17.030194.002013

39. Tse NY, Goldsworthy MR, Ridding MC, et al. The effect of stimulation interval on plasticity following repeated blocks of intermittent theta burst stimulation. *Sci Rep*. 2018;8(1):8526. doi:10.1038/s41598-018-26791-w

40. Nettekoven C, Volz LJ, Kutscha M, et al. Dose-Dependent Effects of Theta Burst rTMS on Cortical Excitability and Resting-State Connectivity of the Human Motor System. *J Neurosci*. 2014;34(20):6849-6859. doi:10.1523/JNEUROSCI.4993-13.2014

41. Goldsworthy MR, Pitcher JB, Ridding MC. The application of spaced theta burst protocols induces long-lasting neuroplastic changes in the human motor cortex. *Eur J Neurosci*. 2012;35(1):125-134. doi:10.1111/j.1460-9568.2011.07924.x

42. Chung SW, Rogasch NC, Hoy KE, Fitzgerald PB. The effect of single and repeated prefrontal intermittent theta burst stimulation on cortical reactivity and working memory. *Brain Stimul*. 2018;11(3):566-574. doi:10.1016/j.brs.2018.01.002

43. Hoshi E. Functional specialization within the dorsolateral prefrontal cortex: A review of anatomical and physiological studies of non-human primates. *Neurosci Res*. 2006;54(2):73-84. doi:10.1016/j.neures.2005.10.013

44. Cash RFH, Zalesky A, Thomson RH, Tian Y, Cocchi L, Fitzgerald PB. Subgenual Functional Connectivity Predicts Antidepressant Treatment Response to Transcranial Magnetic Stimulation: Independent Validation and Evaluation of Personalization. *Biol Psychiatry*. 2019:e1-e3. doi:10.1016/j.biopsych.2018.12.002

45. Fox MD, Halko MA, Eldaief MC, Pascual-Leone A. Measuring and manipulating brain connectivity with resting state functional connectivity magnetic resonance imaging (fcMRI) and transcranial magnetic stimulation (TMS). *Neuroimage*. 2012;62(2012):2232-2243. doi:10.1016/j.neuroimage.2012.03.035

46. Mir-Moghtadaei A, Caballero R, Fried P, et al. Concordance between BeamF3 and MRI-

neuronavigated target sites for repetitive transcranial magnetic stimulation of the left dorsolateral prefrontal cortex. *Brain Stimul.* 2015;8(5):965-973. doi:10.1016/j.brs.2015.05.008

47. Oathes DJ, Zimmerman J, Duprat R, et al. Individualized non-invasive brain stimulation engages the subgenual anterior cingulate and amygdala. 2018.

48. Vink JJT, Mandija S, Petrov PI, van den Berg CAT, Sommer IEC, Neggers SFW. A novel concurrent TMS-fMRI method to reveal propagation patterns of prefrontal magnetic brain stimulation. *Hum Brain Mapp.* 2018;39(11):4580-4592. doi:10.1002/hbm.24307

49. Li CT, Chen MH, Juan CH, et al. Efficacy of prefrontal theta-burst stimulation in refractory depression: A randomized sham-controlled study. *Brain.* 2014;137(7):2088-2098. doi:10.1093/brain/awu109

50. Kaster TS, Daskalakis ZJ, Noda Y, et al. Efficacy, tolerability, and cognitive effects of deep transcranial magnetic stimulation for late-life depression: a prospective randomized controlled trial. *Neuropsychopharmacology.* 2018;43(11):2231-2238. doi:10.1038/s41386-018-0121-x

51. Jorge RE, Robinson RG, O'Brien JT. Top cited papers in international psychogeriatrics: 5. a controlled study of repetitive transcranial magnetic stimulation as a treatment of depression in the elderly. *Int Psychogeriatrics.* 2009;21(5):855-857. doi:10.1017/S1041610209990603

52. Fregni F, Marcolin MA, Myczkowski M, et al. Predictors of antidepressant response in clinical trials of transcranial magnetic stimulation. *Int J Neuropsychopharmacol.* 2006;9(6):641-654. doi:10.1017/S1461145705006280

53. George MS, Lisanby SH, Avery D, et al. Daily left prefrontal transcranial magnetic stimulation therapy for major depressive disorder: A sham-controlled randomized trial. *Arch Gen Psychiatry.* 2010;67(5):507-516. doi:10.1001/archgenpsychiatry.2010.46

54. Kaster TS, Downar J, Vila-Rodriguez F, et al. Trajectories of Response to Dorsolateral Prefrontal rTMS in Major Depression: A THREE-D Study. *Am J Psychiatry.* 2019:1-9. doi:10.1176/appi.ajp.2018.18091096

55. Williams NR, Okun MS. Deep brain stimulation (DBS) at the interface of neurology and psychiatry. *J Clin Invest.* 2013;123(11):4546-4556. doi:10.1172/JCI68341

56. Jelovac A, Kolshus E, McLoughlin DM. Relapse following successful electroconvulsive therapy for major depression: A meta-analysis. *Neuropsychopharmacology.* 2013;38(12):2467-2474. doi:10.1038/npp.2013.149

57. Prudic J, Haskett RF, Mulsant B, et al. Resistance to antidepressant medications and short-term clinical response to ECT. *Am J Psychiatry.* 1996;153(8):985-992. doi:10.1176/ajp.153.8.985

58. Prudic J, Olfson M, Marcus SC, Fuller RB, Sackeim HA. Effectiveness of electroconvulsive therapy in community settings. *Biol Psychiatry.* 2004;55(3):301-312. doi:10.1016/j.biopsych.2003.09.015

59. Pittenger C, Duman RS. Stress, Depression and Neuroplasticity: A Convergence of Mechanisms. *Neuropsychopharmacology.* 2008;33(1):88-109. doi:10.1038/sj.npp.1301574

60. Caudal D, Godsil BP, Mailliet F, Bergerot D, Jay TM. Acute stress induces contrasting changes in AMPA receptor subunit phosphorylation within the prefrontal cortex, amygdala and hippocampus. *PLoS One.* 2010;5(12). doi:10.1371/journal.pone.0015282

61. Pridmore S, Erger S, Rybak M, Kelly E, May T. Early relapse (ER) transcranial magnetic stimulation (TMS) in treatment resistant major depression. *Brain Stimul.* 2018;11(5):1098-1102. doi:10.1016/j.brs.2018.05.013

62. Williams NR, Short EB, Hopkins T, et al. Five-Year Follow-Up of Bilateral Epidural

Prefrontal Cortical Stimulation for Treatment-Resistant Depression. *Brain Stimul*. 2016;9(6):897-904. doi:10.1016/j.brs.2016.06.054

63.   Williams NR, Bentzley BS, Hopkins T, et al. Optimization of epidural cortical stimulation for treatment-resistant depression. *Brain Stimul*. 2018;11(1):239-240. doi:10.1016/j.brs.2017.09.001

64.   Correia-Melo FS, Argolo FC, Araújo-de-Freitas L, et al. Rapid infusion of esketamine for unipolar and bipolar depression: A retrospective chart review. *Neuropsychiatr Dis Treat*. 2017;13:1627-1632. doi:https://dx.doi.org/10.2147/NDT.S135623

65.   Murrough JW, Perez AM, Pillemer S, et al. Rapid and longer-term antidepressant effects of repeated ketamine infusions in treatment-resistant major depression. *Biol Psychiatry*. 2013;74(4):250-256. doi:10.1016/j.biopsych.2012.06.022

66.   Grady SE, Marsh TA, Tenhouse A, Klein K. Ketamine for the treatment of major depressive disorder and bipolar depression: A review of the literature. *Ment Heal Clin*. 2017;7(1):16-23. doi:10.9740/mhc.2017.01.016

67.   Murrough JW, Iosifescu D V., Chang LC, et al. Antidepressant efficacy of ketamine in treatment-resistant major depression: A two-site randomized controlled trial. *Am J Psychiatry*. 2013;170(10):1134-1142. doi:10.1176/appi.ajp.2013.13030392

68.   Williams NR, Heifets BD, Blasey C, et al. Attenuation of antidepressant effects of ketamine by opioid receptor antagonism. *Am J Psychiatry*. 2018;175(12):1205-1215. doi:10.1176/appi.ajp.2018.18020138

69.   Slade EP, Jahn DR, Regenold WT, Case BG. Association of electroconvulsive therapy with psychiatric readmissions in US hospitals. *JAMA Psychiatry*. 2017;74(8):798-804. doi:10.1001/jamapsychiatry.2017.1378

70.   Sackeim HA. Modern electroconvulsive therapy: Vastly improved yet greatly underused. *JAMA Psychiatry*. 2017;74(8):779-780. doi:10.1001/jamapsychiatry.2017.1670

71.   Kellner CH, Fink M, Knapp R, et al. Relief of expressed suicidal intent by ECT: A consortium for research in ECT study. *Am J Psychiatry*. 2005;162(5):977-982. doi:10.1176/appi.ajp.162.5.977

72.   Holtzheimer PE, Husain MM, Lisanby SH, et al. Subcallosal cingulate deep brain stimulation for treatment-resistant depression: a multisite, randomised, sham-controlled trial. *The Lancet Psychiatry*. 2017;4(11):839-849. doi:10.1016/S2215-0366(17)30371-1

73.   Brim RL, Miller FG. The potential benefit of the placebo effect in sham-controlled trials: Implications for risk-benefit assessments and informed consent. *J Med Ethics*. 2013;39(11):703-707. doi:10.1136/medethics-2012-101045

74.   Bakker N, Shahab S, Giacobbe P, et al. RTMS of the dorsomedial prefrontal cortex for major depression: Safety, tolerability, effectiveness, and outcome predictors for 10 Hz versus intermittent theta-burst stimulation. *Brain Stimul*. 2015;8(2):208-215. doi:10.1016/j.brs.2014.11.002

75.   Feffer K, Fettes P, Giacobbe P, Daskalakis ZJ, Blumberger DM, Downar J. 1 Hz rTMS of the right orbitofrontal cortex for major depression: Safety, tolerability and clinical outcomes. *Eur Neuropsychopharmacol*. 2018;28(1):109-117. doi:10.1016/j.euroneuro.2017.11.011

76.   Downar J, Fettes P, Dunlop K, et al. 116. Distinctive Mechanisms of Action for DLPFC-, DMPFC-, and OFC-rTMS in Major Depression. *Biol Psychiatry*. 2018;83(9):S47-S48. doi:10.1016/j.biopsych.2018.02.134

77.   Maeda F, Keenan JP, Tormos JM, Topka H, Pascual-Leone A. Interindividual variability of the modulatory effects of repetitive transcranial magnetic stimulation on cortical excitability. *Exp Brain Res*. 2000;133(4):425-430. doi:10.1007/s002210000432

78.   Du X, Summerfelt A, Chiappelli J, Holcomb HH, Hong LE. Individualized brain inhibition

and excitation profile in response to paired-pulse TMS. *J Mot Behav*. 2014;46(1):39-48. doi:10.1080/00222895.2013.850401

79. Schaworonkow N, Triesch J, Ziemann U, Zrenner C. EEG-triggered TMS reveals stronger brain state-dependent modulation of motor evoked potentials at weaker stimulation intensities. *Brain Stimul*. 2019;12(1):110-118. doi:10.1016/j.brs.2018.09.009

80. Chung SW, Sullivan CM, Rogasch NC, et al. The effects of individualised intermittent theta burst stimulation in the prefrontal cortex: A TMS-EEG study. *Hum Brain Mapp*. 2019;40(2):608-627. doi:10.1002/hbm.24398

81. Oberman L, Eldaief M, Fecteau S, Ifert-Miller F, Tormos JM, Pascual-Leone A. Abnormal modulation of corticospinal excitability in adults with Asperger's syndrome. *Eur J Neurosci*. 2012;36(6):2782-2788. doi:10.1111/j.1460-9568.2012.08172.x

82. Fitzgerald PB, Hoy K, McQueen S, et al. A randomized trial of rTMS targeted with MRI based neuro-navigation in treatment-resistant depression. *Neuropsychopharmacology*. 2009;34(5):1255-1262. doi:10.1038/npp.2008.233

83. Chung SW, Rogasch NC, Hoy KE, Sullivan CM, Cash RFH, Fitzgerald PB. Impact of different intensities of intermittent theta burst stimulation on the cortical properties during TMS-EEG and working memory performance. *Hum Brain Mapp*. 2018;39(2):783-802. doi:10.1002/hbm.23882

84. Chung SW, Hoy KE, Fitzgerald PB. Theta-burst stimulation: A new form of tms treatment for depression? *Depress Anxiety*. 2015;32(3):182-192. doi:10.1002/da.22335

85. Stokes MG, Chambers CD, Gould IC, et al. Simple Metric For Scaling Motor Threshold Based on Scalp-Cortex Distance: Application to Studies Using Transcranial Magnetic Stimulation. *J Neurophysiol*. 2005;94(6):4520-4527. doi:10.1152/jn.00067.2005

86. Fekadu A, Wooderson SC, Markopoulou K, Cleare AJ. The maudsley staging method for treatment-resistant depression: Prediction of longer-term outcome and persistence of symptoms. *J Clin Psychiatry*. 2009;70(7):952-957. doi:10.4088/JCP.08m04728

87. Ozten E, Sayar GH, Karamustafalioglu O. Hypomanic shift observed during rTMS treatment of patients with unipolar depressive disorder: Four case reports. *Ann Gen Psychiatry*. 2013;12(1):1-5. doi:10.1186/1744-859X-12-12

88. Krstic J, Ilic T. Switch to hypomania induced by repetitive transcranial magnetic stimulation and partial sleep deprivation added to antidepressant: A case report. *Vojnosanit Pregl Med Pharm J Serbia*. 2014;71(2):207-210. doi:10.2298/vsp1402207k

89. Xia G, Gajwani P, Muzina DJ, et al. Treatment-emergent mania in unipolar and bipolar depression: Focus on repetitive transcranial magnetic stimulation. *Int J Neuropsychopharmacol*. 2008;11(1):119-130. doi:10.1017/S1461145707007699

90. O'neil-Pirozzi TM, Goldstein R, Strangman GE, Glenn MB. Testre-test reliability of the Hopkins Verbal Learning Test-Revised in individuals with traumatic brain injury. *Brain Inj*. 2012;26(12):1425-1430. doi:10.3109/02699052.2012.694561

91. Benedict RHB, Groninger L, Schretlen D, Dobraski M, Shpritz B. Revision of the brief visuospatial memory test: Studies of normal performance, reliability, and, validity. *Psychol Assess*. 1996. doi:10.1037/1040-3590.8.2.145

92. Wechsler D. WAIS-IV administration and scoring manual. 2008.

93. DELIS DC, KRAMER JH, KAPLAN E, HOLDNACK J. Reliability and validity of the Delis-Kaplan Executive Function System: An update. *J Int Neuropsychol Soc*. 2004;10(02):301-303. doi:10.1017/s1355617704102191

94. Hartman DE. Test review Wechsler Adult Intelligence Scale IV (WAIS IV): Return of the gold standard. *Appl Neuropsychol*. 2009;16(1):85-87. doi:10.1080/09084280802644466

95. Macey PM, Macey KE, Kumar R, Harper RM. A method for removal of global effects from fMRI time series. *Neuroimage*. 2004;22(1):360-366.

doi:10.1016/j.neuroimage.2003.12.042

96. Zimmerman M, Posternak MA, Chelminski I. Derivation of a definition of remission on the Montgomery-Asberg depression rating scale corresponding to the definition of remission on the Hamilton rating scale for depression. *J Psychiatr Res*. 2004;38(6):577-582. doi:10.1016/j.jpsychires.2004.03.007

97. Leucht S, Fennema H, Engel R, Kaspers-Janssen M, Lepping P, Szegedi A. What does the HAMD mean? *J Affect Disord*. 2013;148(2-3):243-248. doi:10.1016/j.jad.2012.12.001

98. Timmerby N, Andersen JH, Søndergaard S, Østergaard SD, Bech P. A Systematic Review of the Clinimetric Properties of the 6-Item Version of the Hamilton Depression Rating Scale (HAM-D6). *Psychother Psychosom*. 2017;86(3):141-149. doi:10.1159/000457131

99. Schulte-Van Maaren YWM, Carlier IVE, Zitman FG, et al. Reference values for major depression questionnaires: The Leiden Routine Outcome Monitoring Study. *J Affect Disord*. 2013;149(1-3):342-349. doi:10.1016/j.jad.2013.02.009

100. Peinemann A, Reimer B, Löer C, et al. Long-lasting increase in corticospinal excitability after 1800 pulses of subthreshold 5 Hz repetitive TMS to the primary motor cortex. *Clin Neurophysiol*. 2004;115(7):1519-1526. doi:10.1016/j.clinph.2004.02.005

101. Volz LJ, Benali A, Mix A, Neubacher U, Funke K. Dose-dependence of changes in cortical protein expression induced with repeated transcranial magnetic theta-burst stimulation in the rat. *Brain Stimul*. 2013;6(4):598-606. doi:10.1016/j.brs.2013.01.008

102. Posner K, Brown GK, Stanley B, et al. The Columbia-suicide severity rating scale: Initial validity and internal consistency findings from three multisite studies with adolescents and adults. *Am J Psychiatry*. 2011;168(12):1266-1277. doi:10.1176/appi.ajp.2011.10111704

103. Grimm S, Beck J, Schuepbach D, et al. Imbalance between Left and Right Dorsolateral Prefrontal Cortex in Major Depression Is Linked to Negative Emotional Judgment: An fMRI Study in Severe Major Depressive Disorder. *Biol Psychiatry*. 2008;63(4):369-376. doi:10.1016/j.biopsych.2007.05.033

104. Koenigs M, Grafman J. The functional neuroanatomy of depression: Distinct roles for ventromedial and dorsolateral prefrontal cortex. *Behav Brain Res*. 2009;201(2):239-243. doi:10.1016/j.bbr.2009.03.004

105. Fitzgerald PB, Laird AR, Maller J, Daskalakis ZJ. A meta-analytic study of changes in brain activation in depression. *Hum Brain Mapp*. 2008;29(6):683-695. doi:10.1002/hbm.20426

106. Hamani C, Mayberg H, Stone S, Laxton A, Haber S, Lozano AM. The subcallosal cingulate gyrus in the context of major depression. *Biol Psychiatry*. 2011;69(4):301-308. doi:10.1016/j.biopsych.2010.09.034

107. Waters AC, Mayberg HS. Brain-Based Biomarkers for the Treatment of Depression: Evolution of an Idea. *J Int Neuropsychol Soc*. 2017;23(9-10):870-880. doi:10.1017/s1355617717000881

108. Lozano AM, Mayberg HS, Giacobbe P, Hamani C, Craddock RC, Kennedy SH. Subcallosal Cingulate Gyrus Deep Brain Stimulation for Treatment-Resistant Depression. *Biol Psychiatry*. 2008;64(6):461-467. doi:10.1016/j.biopsych.2008.05.034

109. Philip NS, Barredo J, Aiken E, Carpenter LL. Neuroimaging Mechanisms of Therapeutic Transcranial Magnetic Stimulation for Major Depressive Disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2018;3(3):211-222. doi:10.1016/j.bpsc.2017.10.007

110. Nahas Z, Teneback CC, Kozel A, et al. Brain Effects of TMS Delivered Over Prefrontal Cortex in Depressed Adults. *J Neuropsychiatry Clin Neurosci*. 2014;13(4):459-470. doi:10.1176/jnp.13.4.459

111. Baeken C, Marinazzo D, Everaert H, et al. The impact of accelerated HF-rTMS on the

subgenual anterior cingulate cortex in refractory unipolar major depression: Insights from [18]FDG PET brain imaging. *Brain Stimul*. 2015;8(4):808-815. doi:10.1016/j.brs.2015.01.415

112.   HAINDL W, MALHI GS, CROKER VM, et al. High (15 Hz) and low (1 Hz) frequency transcranial magnetic stimulation have different acute effects on regional cerebral blood flow in depressed patients. *Psychol Med*. 2003;33(6):997-1006. doi:10.1017/s0033291703007955

113.   Nyffeler T, Cazzoli D, Hess CW, Müri RM. One session of repeated parietal theta burst stimulation trains induces long-lasting improvement of visual neglect. *Stroke*. 2009;40(8):2791-2796. doi:10.1161/STROKEAHA.109.552323

114.   Cazzoli D, Müri RM, Schumacher R, et al. Theta burst stimulation reduces disability during the activities of daily living in spatial neglect. *Brain*. 2012;135:3426-3439. doi:10.1093/brain/aws182

115.   Janicak PG, Professor of Psy-chiatry is, Carpenter L. The Efficacy of Transcranial Magnetic Stimulation for Major Depression: A Review of the Evidence. *Psychiatr Ann*. 2014;44(6):284-292. doi:10.3928/00485713-20140609-06

116.   Modirrousta M, Meek BP, Wikstrom SL. Efficacy of twice-daily vs once-daily sessions of repetitive transcranial magnetic stimulation in the treatment of major depressive disorder: A retrospective study. *Neuropsychiatr Dis Treat*. 2018;14:309-316. doi:10.2147/NDT.S151841