The parietal operculum preferentially encodes heat pain and not salience

Björn Horing, Christian Sprenger, Christian Büchel

Björn Horing, Ph.D. (corresponding author)

Affective Neuroscience Group

Department of Systems Neuroscience

University Medical Center Hamburg-Eppendorf

e-mail b.horing@uke.de

phone +49 040 7410 - 57890

Christian Sprenger, M.D.

Affective Neuroscience Group

Department of Systems Neuroscience

University Medical Center Hamburg-Eppendorf

Christian Büchel, M.D.

Affective Neuroscience Group

Department of Systems Neuroscience

University Medical Center Hamburg-Eppendorf

Abstract

Substantial controversy exists as to which part of brain activity is genuinely attributable to pain-

related percepts, and which activity is due to general aspects of sensory stimulation, such as its

salience. The challenge posed by this question rests largely in the fact that pain is per se highly

salient, a characteristic which therefore has to be matched by potential control conditions. Here, we

used a unique combination of functional magnetic resonance imaging, behavioral and autonomic

measures to address this longstanding debate in pain research.

Subjects rated perceived intensity in a sequence alternating between heat and sound stimuli.

Neuronal activity was monitored using fMRI. Either modality was presented in six different

intensities, three of which lay above the pain threshold (for heat) or the unpleasantness threshold

(for sound). We performed our analysis on 26 volunteers in which psychophysiological responses (as

per skin conductance responses) did not differ between the two stimulus modalities. Having thus

ascertained a comparable amount of stimulus salience, we analyzed pain-related stimulus response

functions, and contrasted them with those of the salience-matched acoustic control condition.

Furthermore, analysis of fMRI data was performed on the brain surface to circumvent blurring issues

stemming from the close proximity of several regions of interest located in heavily folded cortical

areas. We focused our analyses on insular and periinsular regions which are strongly involved in

processing of painful stimuli. We employed an axiomatic approach to determine areas showing

higher activation in painful compared to non-painful heat, and at the same time showing a steeper

stimulus response function for painful heat as compared to unpleasant acoustic stimuli. Intriguingly,

an area in the posterior parietal operculum emerged whose response showed a pain preference, and

where we can unequivocally exclude salience as explanation.

This result has important implications for the interpretation of functional imaging findings in pain

research, as it clearly demonstrates that there are areas whose pain-related activity is not due to

general stimulus characteristics such as salience. Conversely, several areas did not conform to the

formulated axioms to rule out general factors as explanations.

Introduction

Pain is a multidimensional experience, including sensory-discriminative, affective-motivational, cognitive-evaluative as well as motor components [1], and is defined as "an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage" [2]. Following the advent of brain imaging, recurring patterns of brain activity following painful stimuli were summarized as a "pain matrix", comprising primary and secondary somatosensory cortices, cingulate cortices, as well as the insular subregions, among other structures [3,4].

This activity has frequently been attributed to pain per se. However, it has been pointed out that precisely because pain is a composite sensation, some of the observed activation may or may not be exclusively pain-related [5,6]. These studies provided evidence that in many cortical regions, activation is observed for both painful and non-painful (such as tactile or auditory) stimuli. Hence, general processes such as stimulus salience were put forward as an alternative interpretation. These contributions have led to lively controversy [7–9]. Recently, the authors positing the initial challenge to the "pain matrix" concept revisited these issues [10], and reemphasized that great care should be taken experimentally to match non-painful control modalities , which has frequently been neglected in previous studies.

In addition to the question of stimulus salience, many experiments have relied on the use of single stimulus intensities to characterize neuronal responses, when using painful stimulation and compared these responses to a non-painful control condition. However, such approaches disregard the possibility of modality-specific baseline activation, further compounding the issue to properly account for nonspecific activation [11]. A possible solution is to employ multiple stimulus intensities, which allows for the characterization of modality-specific stimulus-response functions [12,13], and a comparison of these between modalities.

Here, we address these issues, and present a novel approach that allows to directly test whether

there are cortical regions that can be defined as salience detectors or show preferential pain

processing. We employed heat and sound as stimulus modalities. Stimuli were presented in

alternating modalities in a within-subjects design. Of each modality, we used six graded intensities –

three below and three above the pain and unpleasantness threshold, respectively. This allowed us to

determine stimulus-response functions of physical intensities or their percepts, and relate those to

neuronal activity [12–14]. Importantly, auditory and thermal intensities were calibrated as

equisalient using an objective autonomic measure (skin conductance responses; SCR) [15].

We paid particular attention to insular and periinsular regions, especially the posterior insula and the

parietal operculum (the secondary somatosensory cortex), all of which have been reported as early

components of pain-responsive cortical areas [4,9,16–18].

To define areas as preferentially pain-processing, our analyses followed an axiomatic approach which

posits several logical conditions to be met to make a valid inference (see [19], for a similar approach

in pain avoidance). Within this rigorous approach, we formulated the following set of conditions to

preclude the possibility that activity in an area could be explained by salience alone: The effect of

painful stimulation should be larger than that of non-painful heat (axiom 1); the effect of painful

stimulation should be larger than that of (salience-matched) unpleasant sound (axiom 2); the

relationship of ratings and BOLD should be stronger for painful heat than for non-painful heat (axiom

3); the positive relationship of pain ratings and BOLD responses should be stronger for painful heat

than for (salience-matched) unpleasant sound (axiom 4).

Results

Heat stimuli were presented using a CHEPS thermode, sounds were 1kHz beeps presented binaurally

via headphones. For a brief overview, see Figure 1. Details are provided in the Method section.

Figure 1. Experimental procedures. **A.** Visual analogue scales (VAS) for rating heat (left) and sound stimuli

(right). The midpoint signifies the pain threshold, corresponding to a score of 0 in a conventional VAS. **B.**

Thermode arrangement on a subject's forearm. Patch C was used for calibration, patches 1|2 were used in

counterbalanced fashion for experimental sessions 1 and 2. **C.** Protocol by time. A visual cue (white fixation

cross turning red) announced the upcoming stimulus (either heat or sound) and stayed visible throughout

stimulation, which was 8 seconds at plateau (roughly 9.5 seconds all in all, depending on calibration). Subjects

were then prompted to rate the stimulus. After rating, the white fixation cross reappeared, to turn red again

for the next cue. Stimulus modalities were always alternating.

Sample

A core prerequisite of our analysis strategy was that both modalities (heat and sound) were matched

with respect to salience. Although previous studies based salience estimates on ratings, this can be

problematic when comparing sound and heat stimuli due to differential scaling. We therefore

selected skin conductance responses (SCR) as an objective readout parameter linked to salience

[15,20,21]. Consequently, our approach is based on comparable SCR for sound and heat stimuli. This

necessitated the selection of suitable subjects and experimental sessions that fulfilled this criterion

(see Methods). Analysis included N=26 subjects (50% female, mean age±SD 25.8±3.6; see S1 Table

for more detailed sample characteristics).

Skin conductance results

As intended by stimulus matching, no significant difference between modalities prevailed (p=0.177)

(random intercept model; Figure 2). SCR increased by intensity (t(308)=7.797, p=1e-13). There was

no interaction between intensity and modality (p=0.514).

Figure 2. Skin conductance responses following heat (A) and sound stimuli (B). The pain and unpleasantness

thresholds were located between intensities 3 and 4, as per calibration.

Stimulus calibration results

Mean heat pain threshold was at 43.5±1.1°C (range 40.5 to 45.4) and corresponded to 50 points on a

0 to 100 point visual analogue scale (VAS). Temperatures for stimulus intensities below and above

pain threshold, corresponding to VAS targets of 25, 35, 45, 55, 65 and 75, were 41.8±1.5°C,

42.5±1.3°C, 43.2±1.1°C, 43.9±1.0°C, 44.6±1.0°C and 45.3±1.1°C, respectively. Mean unpleasantness

threshold was at 83.0±6.7dBA (range 69.0-99.8). For additional details on heat and sound calibration,

see Methods, and S2 Table.

Behavioral results

The analysis of subjective ratings of sound and heat stimuli revealed a significant effect of modality

(t(320)=7.820, p=8e-14; average sound rated estimate±SE 13.3±1.7 VAS points higher than average

heat) (random intercept model; Figure 3) and a main effect of intensity (t(320)=42.014, p=2e-16;

11.8±0.3 VAS points per intensity step). The interaction between intensity and modality was also

significant (t(320)=-4.1529, p=4e-5; 2.3±0.6 VAS points shallower slope in sound, per intensity step).

Figure 3. Behavioral ratings following heat (A) and sound stimuli (B). The pain and unpleasantness thresholds were located between intensities 3 and 4, as per calibration.

Imaging results

For either modality, a mask was used that was obtained from main effect activations a) larger than

the respective comparator modality and b) larger than baseline (S1A Figure; see Methods for details).

The same mask was applied to all contrasts reported in the following, with the exception of

conjunction analyses, which were performed without mask. Application of the masks constrains the

analyses to areas consistently activated during the respective modality.

Main effects of modality

To test for intermodal differences, we contrasted the main effects for heat and sound (Figure 4).

Figure 4. Differential effects of heat (orange) and sound (blue). Significant differences were found in the

parietal operculum (H1, H2) and dorsal posterior insula (H3) for heat; in the superior temporal gyrus (S1) and

Heschl's gyri for sound. **A.** Activations are thresholded at p(uncorrected)<0.001 and overlaid on an average

brain surface for display purposes. The black line delineates the region of interest used for correction for

multiple comparisons. See S2 Figure for peak locations in brain volume slices. **B.** Poststimulus plots of fMRI

activation over all stimulus intensities (mean±SE). Subplots H1 through H3 show that heat-related activation

(orange) dominates in the analyzed time frames (seconds 2.2 through 10.8, see Methods), while subplot S1

shows increased sound activation (blue).

The parietal operculum (secondary somatosensory cortex; peak MNI coordinates x=51, y=-30, z=28,

Z=5.62, p(corrected)=1e-05; second peak at x=59, y=-23, z=25, Z=5.221, p(corrected)=1e-04) and

dorsal posterior insula (x=40, y=-21, z=19, Z=4.175, p(corrected)=0.012) showed stronger activation

for heat as compared to sound. Conversely, Heschl's gyri (primary auditory cortex; x=64, y=-24, z=7,

Z=Inf, p(corrected)=4e-16) showed stronger activation for sound stimuli.

Of note, areas activated by either modality show no overlap, as determined via conjunction analyses,

even at a liberal threshold of p(uncorrected)<0.001, of contrasts of heat or sound larger than

baseline activation. The conjunction analysis did not use any masking; regardless, it did not yield

significant results.

Parametric modulation by stimulus intensity

Irrespective of modality, main effects can be confounded by unspecific effects associated with the

generic occurrence of an external stimulus, such as orientation and response preparation. Therefore,

we performed an analysis investigating stimulus response functions (SRFs), i.e., testing for stronger

BOLD responses for higher stimulus intensities.

We contrasted both modalities to identify areas with diverging SRFs within those areas showing a

main effect of either modality, as determined above. For heat, we identified activity in the parietal

operculum (x=57, y=-30, z=31, Z=3.999, p(corrected)=0.026) whose SRF diverges from that of the

sound modality (Figure 5A). For sound, no significant activity prevailed, that is, no relationship of

intensity and brain activity was found within the region of interest. Closer inspection of the time-

course of the SRF in the heat modality (Figure 5B) indicates that the SRF's maximum slope coincides

with the peak of the main effect, that is, the modulation of the main effect by intensity is strongest

when the main effect itself is strongest.

B Parietal operculum (SII)

Figure 5. Differential modulation by stimulus intensity for heat (orange) and sound (blue). Significant

differences were found in the parietal operculum (H1) for heat. **A.** Activations are thresholded at

$p(uncorrected)<0.001$ and overlaid on an average brain surface for display purposes. The black line delineates

the region of interest used for correction for multiple comparisons. See S3 Figure for peak positions in brain

volume slices, and S3B for sound activation in S1. **B.** Poststimulus plots of fMRI activation in vertex H1 during

heat (orange) and sound (blue). The colored patches at the right axes show the stimulus duration. The lower

left (y-)axes show the parametric modulation affecting the main effect (average size of the effect along the

lower right (x-)axes): A straight line parallel to the y-axis indicates no change of the BOLD response depending

on stimulus intensity, whereas the a sloped main effect along the y-axis indicates parametric modulation. In

this area, the main effect of heat is mostly positively modulated by stimulus intensity, that is, higher stimulus

intensities induce a higher extent of BOLD. The highest main effect activation occurs around second 10.8

(corresponding to scan 6), coinciding with the steepest slope of parametric modulation by intensity (y-axis).

Again, an unmasked conjunction analysis at a liberal threshold yielded no significant overlap of both

modalities, when comparing contrasts with an SRF slope larger than zero for either modality.

Parametric modulation by ratings

Although relevant, physical stimulus intensity might not be directly mapped to neuronal activity, as a

sensory signal undergoes multiple levels of processing before it reaches cortical areas. We therefore

performed an additional analysis, where we investigated whether areas show BOLD responses that

are correlated with subjects' behavioral ratings.

This analysis revealed that activity in the parietal operculum (x=55, y=-37, z=26, Z=5.091,

p(corrected)=2e-04; x=58, y=-14, z=18, Z=4.312, p(corrected)=0.008) and the dorsal anterior insula

(x=36, y=0, z=14, Z=4.276, p(corrected)=0.009; x=41, y=1, z=14, Z=3.849, p(corrected)=0.047) (Figure

6) showed a positive relationship to perceived intensity. This agrees with and extends results from

the previous analysis in which BOLD responses were correlated with stimulus intensity in the parietal

operculum. For sound, no significant activity prevailed, that is, no relationship of ratings and brain

activity was found.

Figure 6. Differential modulation by ratings for heat (orange). Significant differences were found in the parietal

operculum (H1, H2), central operculum (H3) and dorsal anterior insula (H4). **A.** Activations are thresholded at

p(uncorrected)<0.001 and overlaid on an average brain surface for display purposes. The black line delineates

the region of interest used for correction for multiple comparisons. See S4 Figure for peak positions in brain

volume slices. **B.** Poststimulus plots of fMRI activation in vertices H1 through H4 during heat (orange) and

sound (blue). The colored patches at the right axes show the stimulus duration. The lower left (y-)axes show

the parametric modulation by ratings that are affecting the main effect.

As before, in the unmasked conjunction analysis of contrasts where rating correlated with the BOLD

responses, no regions with significant overlap were found.

Imaging results distinguishing stimuli perceived below and above thresholds

So far, all analyses pooled over non-painful and painful heat percepts. To further investigate pain-

related responses, we separated those stimuli reported as non-painful from those reported as painful

(i.e., subthreshold versus suprathreshold), and similarly for unpleasant versus non-unpleasant

sounds. We followed an axiomatic approach to identify areas where activity under painful

stimulation could neither be explained by an overlap with activity under non-painful heat (as would

be the case, e.g., in thermosensitive areas), or by an overlap with activity following unpleasant sound

(e.g., in areas processing stimulus salience). In particular, we posited that a region can be

characterized as preferentially pain-processing if the following conditions hold:

▪ Axiom 1: The effect of suprathreshold – i.e., painful – stimulation should be larger than that

of subthreshold – i.e., heat – stimulation.

▪ Axiom 2: The effect of suprathreshold heat stimulation should be larger than that of

suprathreshold sound stimulation.

▪ Axiom 3: The relationship of ratings and BOLD – i.e., the slope of the stimulus response

function – should be stronger for suprathreshold heat than for subthreshold heat.

■   Axiom 4: The relationship of ratings and BOLD should be stronger for suprathreshold heat

than for suprathreshold sound.

Each of the axioms was evaluated at a significance threshold of p=0.05, corrected for family-wise

error. After joint application of each axiom, analysis revealed activation in the posterior parietal

operculum (x=56, y=-37, z=25, Z=5.519, p(corrected)=0.035) (Figure 7), adjacent to the supramarginal

gyrus.

Figure 7. Areas that fulfill the axiomatic requirements of differential activation during pain compared to heat

and sound. In detail, these axioms were 1) a larger effect of suprathreshold heat compared to subthreshold

heat, 2) a larger effect of suprathreshold heat compared to suprathreshold sound, 3) a stronger relationship of

BOLD with pain ratings than with heat ratings, 4) a stronger relationship of BOLD with pain ratings than with

unpleasantness ratings. Significant activation was found in the parietal operculum (H1). **A.** Activations are

thresholded at p(uncorrected)<0.001 and overlaid on an average brain surface. The black line delineates the

SVC mask. See S5 Figure for peak positions in brain volume slices. **B.** Poststimulus plots of fMRI activation in

vertex H1 during heat (orange) and sound (blue). The shaded patch in the center signifies the pain threshold

(for heat) and unpleasantness threshold (for sound). The colored patches at the right axes show the stimulus

duration.

Discussion

This study aimed to identify regions relevant for heat pain processing, and to determine whether

their activation can be explained by salience. We used individually calibrated, parametrically graded

heat stimuli, and an auditory control condition. Heat and sound stimuli were matched for arousal as

indicated by similar skin conductance responses. Furthermore, we employed surface-based analyses

to mitigate spatial inaccuracies of 3D smoothing.

Main effects for heat were identified in the parietal operculum and the posterior insula, main effects

of acoustic stimuli were observed in the superior temporal gyrus. More importantly, in the parietal

operculum, we observed a differential correlation of brain activity with ratings above versus below

the heat pain threshold, concurrent with a differential correlation with ratings under painful heat

versus unpleasant sound. As we have matched both modalities for salience, these results

unequivocally rule out that activity in this area is simply related to stimulus salience, and suggests a

more dedicated role in heat pain processing.

Using SCR as an autonomic readout of arousal [15,20,21] allowed us to establish comparable salience

of the stimulus material, independent of any behavioral assessments. Although salience can be

assessed psychometrically [5] and research exists to establish concurrent validity of salience ratings

within individual modalities [22], to our knowledge, such ratings have not been validated cross-

modally. It is likely that salience ratings are scaled differently according to some modality-specific

perceptual range. Our results support this notion, as we have observed a prevailing difference in

behavioral ratings between the two modalities (sound was, on average, rated as more aversive, but

had a shallower slope with increasing intensities). This means that a reliance on behavioral ratings

alone could compound SCR dissimilarities between comparator modalities.

With six graded stimulus intensities per modality, our design allowed for the assessment of stimulus

response functions as opposed to simple mean comparisons between a single intensity and a low-

level baseline, or between single sub- and suprathreshold stimuli. Apart from physical intensities, this

also allowed us to use a large range of individual ratings as predictors. Using these perceived

intensities, we were able to directly investigate competing modes of encoding. For example, a brain

area may encode heat intensity, regardless of pain, or it may be inactive below threshold but encode

pain intensity above threshold [12,13,23]. In the analysis distinguishing between sub- and

suprathreshold stimuli, we see a clear pain-intensity-related response in the parietal operculum

(Figure 7). While not a main focus of this paper, we do see a shift in SRFs even within small cortical

distances: For example, an area rostral (x=55, y=-26, z=26) to the more heat pain-dedicated posterior

parietal operculum (x=56, y=-37, z=25) fails to register differences in the parametric modulation by

sub- and suprathreshold heat (Figure 8).

Figure 8. Distinction of areas with stimulus response functions corresponding to the axioms, or not (heat

modality only). A. The slopes of subthreshold (heat) versus suprathreshold (pain) activation as described in

Figure 7. Heat slopes are shallower than pain slopes. B. Slopes of heat and pain activation in an opercular

vertex slightly anterior to A, as determined per conjunction of heat and pain parametric modulation. Slopes are

more aligned, preventing the contrasted activation (pain>heat) of reaching significance. Note that this is not a

formal comparison to A. **C.** Location of the vertices described in A and B. B is in an adjacent area about 1 cm

surface distance rostral from A.

Areas in the insula and surrounding cortical areas are characterized by extreme cortical folding. We

therefore implemented a subject specific surface-based fMRI analysis. This prevents contamination

of gray matter voxels by signal from white matter and cerebrospinal fluid. Furthermore, surface-

based analyses circumvent potential issues arising from three-dimensional smoothing which

accidently mixes signals from structures adjacent in three-dimensional space which are actually

distant from each other. For example, the parietal operculum is directly adjacent to the superior

temporal gyrus in three-dimensional space, but their neurons are separated by the entire insular

fold. Smoothing with a three-dimensional kernel therefore includes activity across the lateral sulcus

(alongside noise from white matter and corticospinal fluid), thereby increasing error terms and

decreasing sensitivity of the respective comparisons. Importantly, in this case, three-dimensional

smoothing could also generate erroneous overlaps between conditions. Surface-based analyses have

been found to increase sensitivity and reduce deviations when normalizing from native to standard

space [24].

In contrast to previous multimodal studies, we explicitly chose modalities where the aversiveness

would be generated by virtue of physical intensity. This is naturally the case with painful stimulation,

but several studies did not use aversive stimulation in non-painful control modalities (for example

[9], who used low-intensity tactile stimulation). Consequently, the acoustic modality was chosen

because stimuli can be generated in close analogy to heat, by altering the physical intensity of the

stimuli.

For our analyses of neuronal activity, we have focused on the posterior insula and adjacent areas.

The insula is of particular interest, because its involvement in pain has been well-documented

[4,9,25]. Furthermore, it has been hypothesized to perform polymodal magnitude estimation [23,26],

and is also involved in salience processes [27,28]. Unambiguous data concerning the involvement of

the insular cortex in pain processing also comes from direct cortical stimulation studies [29–31].

Consequently, it is a prime candidate to assess overlaps and differences in activation patterns.

With its reliable activation following painful stimulation, we can unequivocally establish the parietal

operculum as an important area of heat pain processing, whose activity cannot be explained by

stimulus salience. The area not only shows increased activation when comparing pain and other

modalities (heat, sound), but also exhibits a monotonic increase with perceived pain. The peak of the

BOLD response following pain clearly coincides with the largest modulation by behavioral ratings of

pain, roughly 8 seconds after stimulus onset (Figure 7B). Importantly, this area has close functional

connections with the posterior insula [32], another area of interest [9,16].

Interestingly, we were not able to replicate earlier findings [5,33] of substantial overlap of activation

regardless of modality, even at a lower threshold. This might be related to the differences in stimulus

parameters between the studies: Previous multisensory studies have used rapid onset stimuli of very

short duration, whereas ours were considerably longer (8 seconds plateau, circa 9.5 seconds with

upward/downward slopes). It is possible that with increasing brevity and suddenness of the stimuli,

the extent of unspecific orientation responses and other attention related processes is

disproportionally larger, and therefore a larger overlap of neuronal activation can be observed [34]. If

true, this overlap would naturally be determined, to a large extent, by unspecific and not pain-

related activations such as salience.

Brain responses evoked by stimuli in different sensory modalities might follow different time courses;

systematic differences may, for example, arise from different conduction speeds of fibers relaying

auditory (mostly very fast Aα fibers), thermoceptive (mostly slow C fibers) and nociceptive input (Aδ

and C fibers), compounded by the fact that thermal stimulation (in this setup) occurs at a distal site

compared to auditory stimulation. Therefore, in analogy to [5], we opted for analyzing the time-

course of all imaging data by using finite impulse responses as basis functions. This largely avoids the

constraints and biases implicit in comparing mean activations obtained by pre-defined hemodynamic

response functions.

Some limitations apply to the present research.

The study used stimuli of mild to moderate aversiveness (calibrated to a maximum of 50 if rescaled

to a conventional, 0-100 suprathreshold visual analogue scale). This aspect, too, could be amended

to cover a broader range, albeit increasing the risk of carry-over effects such as sensitization,

particularly with longer stimulus duration. Additionally, the use of only a single trial-based, post-

stimulus rating of stimulus intensity could be criticized. In fact, one common recommendation for

pain measurement is to distinguish multiple pain dimensions [1], most frequently intensity and

unpleasantness [35], although these aspects tend to be highly correlated in non-interventional

designs [36,37]. Given the SCR-based approach to equalize salience and to include more stimulus

repetitions, we opted against multiple VAS for protocol reasons, namely ease of measurement and to

avoid confusion.

While we have identified areas preferentially active in painful heat as compared to unpleasant

sound, we cannot claim that these areas are specific for pain. In fact, it is important to note that

specificity cannot be ascertained with a limited number of control conditions [10,38]. We concur that

the notion of specificity is more academic in nature than might benefit the field [3]. The preferences

of certain areas to process various inputs – whether visual, acoustic, nociceptive – is best construed

as a matter of degree, that is, a question of specialization rather than specificity, as has been

suggested for functions unrelated to pain [39]. Nevertheless, the rigorous axiomatic approach allows

for a strong hypothesis ascribing the parietal operculum a dedicated role in pain processing.

Materials and Methods

The protocol was approved by the local Ethics Committee (Ethikkommission der Ärztekammer Hamburg, vote PV4745) and conformed to the standards laid out by the World Medical Association in the Declaration of Helsinki. Participants gave written informed consent prior to participation.

Exclusion criteria

A list of exclusion criteria is provided in Table 1.

Table 1. Exclusion criteria.

- Age younger than 18, older than 40
- Sufficient visual acuity, correction with contact lenses only
- Conditions disqualifying for MR-scanners (e.g. claustrophobia, wearing a pacemaker)
- Ongoing participation in pharmacological studies, or regular medication intake (e.g. analgesics)
- Analgesics use 24h prior to the experiment
- Pregnancy or breastfeeding
- Chronic pain condition
- Manifest depression (as per Beck Depression Inventory II, BDI-II, cutoff 14 as per [40])
- Somatic symptom disorder (as per Patient Health Questionnaire, PHQ15, cutoff 10 as per [41])
- Other neurological, psychiatric or dermatological conditions
- Inner ear conditions
- Head circumference >60 cm (for second cohort, due to device constraints)

Psychophysiological recordings

Electrodermal activity was measured with MRI-compatible electrodes on the thenar and hypothenar

of the left hand. Electrodes were connected to Lead108 carbon leads (BIOPAC Systems, Goleta, CA,

USA). The signal was amplified with an MP150 analog amplifier (also BIOPAC Systems). It was

sampled at 1000 Hz using a CED 1401 analog-digital converter (Cambridge Electronic Design,

Cambridge, UK) and downsampled to 100 Hz for analysis.

Analysis was performed using the Ledalab toolbox for MATLAB [42]. Single subject data were

screened for artifacts which were removed if possible by using built-in artifact correction algorithms.

Using a deconvolution procedure, we computed phasic skin conductance (SCR). SCR occurring after

stimulus onset and within stimulus duration was used for measuring autonomic arousal as a proxy

for stimulus salience. Response windows were defined by visual inspection, per modality: between

2.0 s and 4.5 s for heat, and between 1.5 s and 4.0 s for sound. Results were log- and z-transformed

to reduce the impact of intra- and interindividual outliers [15]. Subsequently, SCR was averaged

within subjects for two modalities (heat/sound) and six stimulus intensities each, yielding twelve

values per person.

SCR was used because it is an objective measure of general sympathetic activity, and therefore a

measure of arousal and stimulus salience [15,20,21] which is routinely used in assessing painful

[14,43,44] as well as acoustic stimulation [45].

fMRI acquisition and preprocessing

Functional and anatomical imaging was performed using a TRIO 3T MR Scanner (Siemens, Erlangen,

Germany) with a 12-channel head coil. An fMRI sequence of 36 transversal slices of 2 mm thickness

was acquired using T2*-weighted gradient echo-planar imaging (EPI; 2150 ms TR, 25 ms TE, 80° flip

angle, 2x2x2 mm voxel size, 1 mm gap, 216x216x107 mm field of view, acceleration factor of 2 with

generalized autocalibrating partially parallel acquisitions reconstruction, GRAPPA). Coverage did not

include the apical parts of the frontal/parietal lobes. Additionally, a T1-weighted MPRAGE anatomical

image was obtained for the entire head (voxel size 1x1x1 mm, 240 slices).

For each subject, fMRI volumes were realigned to the mean image in a two-pass procedure, and co-

registered to the anatomical image using affine transformations. Anatomical images were segmented

into tissue types, and individual brain surfaces generated, using the CAT12 toolbox for SPM (Christian

Gaser & Robert Dahnke, http://www.neuro.uni-jena.de/cat/).

Analysis of imaging data

Subject-level analyses were performed on the 3D (volume) data in native space without smoothing,

using an implicit mask at 0.6 to facilitate subsequent (surface) processing. We computed general

linear models to identify brain structures involved in the processing of each stimulus modality, as

well as the encoding of intensities within those modalities (volume data not shown). All analyses

were performed with seventh order FIR basis functions, of which bins 2 to 6 are considered when

comparing conditions. This amounts to seconds 4.3 through 12.9 post stimulus onset. Realignment

(motion) parameters as well as regressors obtained from ventricular motion were included as

nuisance variables, to mitigate motion-related artifacts.

We first set up a model including one regressor for stimulus main effects in each modality. Another

two regressors – one linear, one quadratic – encoding stimulus intensities 1 through 6 were added

per modality, as parametric modulators. The second model likewise included main effects, and

behavioral ratings as linear and quadratic parametric modulators. Finally, the third model further

distinguished the two modalities in stimuli perceived as below and above the respective thresholds

(pain for heat stimuli, unpleasantness for sound stimuli), yielding four main effect regressors

(subthreshold heat, suprathreshold heat – i.e., pain –, subthreshold sound, suprathreshold sound).

Behavioral ratings were again included as linear parametric modulators; quadratic modulation was

not considered to preclude overfitting.

Results from subject-level analyses were mapped to brain surfaces obtained via the CAT12

segmentation procedure. The mapped subject-level results were then resampled to correspond to

surface cortical templates, and smoothed with a 6 mm full width-half maximum 2D kernel. Group-

level analyses were performed including the mapped contrasts, which are described in the Results

section.

Masking was used to distinguish either modality, as the ANOVAs employed are unsigned and in

principle detect differences in activation regardless of direction. Therefore, we obtained signed (that

is, unconstrained by p values) masks from calculating a conjunction from significant voxels of a) a t-

test contrasting the average main effects of either modality (i.e., where activation following heat was

larger than that following sound, and vice versa), and b) a t-test contrasting either modality to low-

level baseline (i.e., where activation following heat – or sound, respectively – was larger than zero.

This yielded a single mask for both modalities, which was applied to all analyses (unless otherwise

noted) (S1B Figure).

For the purpose of this study, we focused on the hemisphere contralateral to the stimulation, in our

case the right hemisphere. In general, the larger part of activity following pain is contralateral to the

stimulation site, but is known to be bilateral in several key areas such as the secondary

somatosensory cortex and the insula [46].

Furthermore, we focused on the insula and directly adjacent areas for small volume correction of

significance level. In particular, we included the granular insular cortex (Ig1, Ig2) as well as the

parietal operculum (OP1, OP2) and primary auditory cortex (Te1.1), using the SPM Anatomy Toolbox

(version 2.2b [47]). This mask was mapped to a template brain surface, then smoothed with a 4 mm

2D kernel to close gaps. The resulting binary mask (S1A Figure) was roughly centered around

previously reported coordinates (x=[-]34, y=-20, z=18) involving areas putatively dedicated to pain

processing [9]. It was used for small volume correction of second level analyses, where results were

considered after correction for family-wise error rate of $p<0.05$.

Psychometry

Owing to the study's aim to compare two stimulus modalities (heat and sound), they had to be

presented and rated in an analogous fashion. Therefore, while retaining the intuitive descriptor

"painfulness" for rating noxious heat (as composite measure of intensity and unpleasantness), we

settled on "unpleasantness" as descriptor for sounds. This also seemed warranted considering the

high correlation of intensity and unpleasantness measures in heat pain [37], while unpleasantness is

one of the definitional criteria of pain [2].

Furthermore, since we wanted to use graded stimuli both below and above the respective thresholds

(pain threshold for heat, unpleasantness threshold for sound), we deviated from the more common

simple visual analogue scales (VAS) and devised two partitioned 0 to 100 VAS for both modalities

(Figure 1A).

For heat, it captured both painful and non-painful sensations. Subjects were instructed to indicate

heat intensity in absence of pain in the 0 through 49 range, and heat pain intensity in the 50 through

100 range. Hence, anchors were displayed for "no sensation" (0), "minimal pain" (50), and

"unbearable pain" (100). Pain was operationalized as the presence of sensations other than pure

heat intensity, such as stinging or burning, as per the guidelines of the German Research Network on

Neuropathic Pain [48].

Likewise, for sound, both unpleasant and non-unpleasant sensations were captured by the VAS.

Subjects were instructed to indicate loudness in absence of unpleasantness in the 0 through 49

range, and loudness unpleasantness in the 50 through 100 range. Anchors were displayed for

"inaudible" (0), "minimally unpleasant" (50), and "extremely unpleasant" (100). Unpleasantness was

operationalized as a bothersome quality of the sound emerging at a certain loudness.

Heat stimuli and calibration

Heat stimuli were delivered using a CHEPS thermode (Medoc, Ramat-Yishai, Israel). Stimulation sites

were located on the radial surface of the forearm. Three separate sites were used for calibration and

either experimental session, to avoid changes in heat/pain perception due to repeated stimulation.

Around the middle of the forearm (half distance between crook of the arm and distal wrist crease;

see Figure 1B), three stimulation sites were marked prior to the experimental sessions. For

calibration, a medial site on the distal part of the forearm was used; for experimental sessions 1 and

2, two adjacent proximal sites were used, in counterbalanced order. During both calibration

procedure and experimental sessions, baseline temperature was set to 35°C, and rise and fall rate

were set to 15°C per second. The duration of heat stimuli was set to eight seconds at target

temperature (plateau), except for preexposure stimuli whose plateau duration was zero (and thus

only consisted of temperature up- and downramping).

A two-step stimulus calibration was performed for each subject, to determine three temperatures

below the individual pain threshold, and three above. Calibration was performed with the MR-

scanner running the same sequence as during the actual experimental sessions, to mimic ambient

conditions [49]. fMRI data from calibration was later discarded.

In a first calibration step, the pain threshold was determined. Subjects were preexposed to four brief

heat stimuli. Preexposure started at 42°C and each consecutive stimulus was increased by 0.5°C, up

to 43.5°C. If a subject indicated the last stimulus as painful, starting temperature for the following

procedure was set to 43°C, else to 44°C. We then used a probabilistic tracking procedure for

threshold determination, assuming a normal distribution of pain perception around the actual

threshold [50]. Eight full-length stimuli were presented and received a binary rating (painful or not

painful). Depending on the rating of the previous stimulus, each consecutive stimulus was set to a

higher or lower temperature according to the probability informed by previous pivot points. The final

temperature was defined as threshold intensity.

In a second calibration step, eight stimuli unevenly spaced around threshold intensity (from -2°C to

+1.6°C, with smaller intervals towards ±0°C) were rated on the partitioned VAS described above.

After the procedure, linear regression was used to calculate target temperatures H1 through H6, to

obtain subthreshold VAS ratings of 25, 35 and 45 (H1–H3), and suprathreshold VAS ratings of 55, 65

and 75 (H4–H6).

These six intensities were used throughout the experimental sessions.

Sound stimuli and calibration

Sound stimuli were delivered using MR-compatible headphones (NordicNeuroLabs, Bergen, Norway).

A pure sound (frequency 1000 Hz, sampling rate 22050 Hz) was generated using MATLAB. A log

function was used to translate increases in (physical) amplitude to smooth gradual increases in

(psychoacoustic) loudness, to mimic the heat stimuli's temperature ramps. Like the heat stimuli,

sound stimuli were presented for eight seconds at target loudness (plateau), and the scanner was

running a dummy EPI sequence throughout to mimic actual conditions [51].

A two-step stimulus calibration was performed for each subject, to determine three sounds below

the individual unpleasantness threshold, and three above. The general procedure was analogous to

the one used for heat.

In a first calibration step, the individual loudness unpleasantness threshold (in percent of maximum

amplitude of ~100 dB, allowing for safe exposure even at maximum intensities [52]) was determined

by an ascending methods of limits-procedure. Six sounds of gradually increasing loudness were

played. The calibration sounds differed in the steepness of the loudness ramps, taking between 9 and

15 seconds to reach peak amplitude. Subjects were asked to indicate the point where the loudness

became unpleasant. The mean of the last four of the six stimuli was defined as threshold loudness.

In a second calibration step, 16 stimuli unevenly spaced around threshold loudness were presented

(from -15% to +15%, smaller intervals towards ±0%), with stimulus characteristics set to mimic those

of heat stimuli (roughly 0.75 s ramps up and down, plus 8 s plateau loudness). As with heat ratings,

linear regression was used to calculate target amplitudes S1 through S6, namely to obtain

subthreshold VAS ratings of 25, 35 and 45 (S1–S3), and suprathreshold VAS ratings of 55, 65 and 75

(S4–S6). For the second cohort, VAS targets were informed by the corresponding mean SCR

amplitude of the first cohort (see "Differences between first and second cohort").

Finally, ramping characteristics of sound stimuli (the seconds it took to plateau) were set to

correspond to those of the respective intensity's heat stimuli, such that corresponding intensities of

both modalities had an identical overall length (ramps plus plateau).

Stimulus presentation during experimental sessions

After calibration, the thermode stimulation site was changed, and the first experimental session

commenced. Heat and sound stimuli were presented in alternation, so that trials of the same

modality were spaced with an intertrial interval of approximately 30 seconds. Each trial followed the

same basic structure (Figure 1C).

Within each modality, the six intensities were pseudorandomized in microblocks. Randomization was

performed such that each sequence of six stimuli contained one instance of each intensity. It was

further constrained such that the very first stimulus was never chosen from the highest two

intensities, and two consecutive intensities were never more than 3 intensity steps different (e.g.,

the intensity following heat intensity 1 could not exceed heat intensity 4).

After changing thermode stimulation site again, session 2 commenced with identical protocol (albeit

different randomization).

Visual cues and VAS rating scales were displayed in the scanner using back-projection via a 45° mirror

placed atop the head coil.

Selection of subsample for analysis with comparable SCR between modalities

In total we assessed two cohorts of 32 subjects and 26 subjects. To obtain an "SCR-equalized"

subsample from all subjects (N=58 with 2 sessions, that is a total of 116 experimental sessions), in a

first step, we excluded all sessions where the correlation between ratings (that is, perceived stimulus

intensity/unpleasantness) and SCR was lower than or equal to zero, so that only subjects with a

positive correlation in both modalities were eligible for the next step.

In a second step, we used Bayes factors [53,54] to determine the flipping point where modality

became obsolete as explanatory variable. Bayes factors express the ratio of the marginal likelihood of

the data under the compared models; since they consider the number of free parameters, they allow

for the selection of the "better" model (best fit to the data and most parsimonious). For every

session, we obtained the mean SCR (log-transformed and normalized values) for both modalities;

sessions with the largest predominance of heat-SCR were then consecutively removed. After each

removal, we obtained the Bayes factors for the remaining sample, comparing the model with

intensity only as predictor to that with modality added as predictor. Once the Bayes factor dropped

below 1 (meaning that the addition of modality as predictor did not serve to improve the model), we

stopped the pruning procedure. This relatively permissive criterion for session inclusion was chosen

in order to preserve as many sessions as possible.

This procedure yielded a sample where modality did not contribute to explaining the SCR data (as

indicated by recalculating the random intercept model described under "Skin conductance results"),

with 26 unique subjects contributing 33 sessions. From the first cohort, 15 subjects contributed 19

sessions, from the second cohort, 11 subjects contributed 13 sessions to the SCR-equalized analysis.

Differences between first and second cohort

Since we had determined that not every person's skin conductance responded to both modalities to

a comparable extent, we set out to select a subsample of persons who had comparable SCR. To reach

a sufficient number of such "responders", we had to perform an additional data collection.

Because of logistical reasons (scanner upgrade in January 2018), some parameters of fMRI

acquisition had to be modified for the new PRISMA 3T MR Scanner (Siemens, Erlangen, Germany).

Instead of a 12-channel head coil, we had to employ a 20-channel head coil. Delivery of the auditory

stimulus was performed with a CONFON headphone (Cambridge Research Systems Ltd, Rochester,

United Kingdom). These measures necessitated the exclusion of subjects with head circumference

above 60cm.

Furthermore, to facilitate increased SCR responding to sound, we increased the amplitude of the

sound stimuli. Using calibration data from the first data collection and linear extrapolation, we

calculated sound VAS targets required to induce SCRs of an amplitude comparable to those of heat

VAS targets of the same intended intensity 1 through 6. We determined that corresponding to our

heat VAS targets of 25, 35, 45, 55, 65, 75 (see "Heat stimuli and calibration"), we would need to apply

sound amplitudes inducing sound VAS targets of 48, 59, 70, 82, 93, 105. Furthermore, during subject

instruction, we emphasized the fact that the amplitude of sound stimuli was not within pathological

range. This was done to prevent overly cautious subject behavior, following anecdotal evidence from

the first cohort that sound stimuli were associated with higher safety concerns than heat stimuli.

Statistical analyses

All analyses were performed using MATLAB (version R2017b) and SPM12 (version 6906).

Significance level was set to $p=0.05$ for psychophysiological and behavioral data, whereas imaging

results were corrected using family-wise error rate adjustment at $p<0.05$. For visualization,

activations are thresholded at p(uncorrected)<0.001 and overlaid on an average brain surface. All

coordinates are reported in Montreal Neurological Institute (MNI) space.

Skin conductance data and behavioral ratings were analyzed using linear mixed models with random

intercepts [55], with centering of predictors following recommendations [56].

Group-level analyses of imaging data were performed as within-subjects ANOVA (cf. "Analysis of

imaging data" above, and the respective Results sections).

Acknowledgements

This work was supported by European Research Council Advanced Grant ERC-2010-AdG_20100407

and Deutsche Forschungsgemeinschaft Grant SFB 936 Project A06. Björn Horing was supported by

the Alexander von Humboldt-Foundation (Feodor Lynen Return Fellowship). We thank Jürgen

Finsterbusch, Katrin Bergholz, Waldemar Schwarz and Kathrin Wendt for technical assistance during

MR data collection, and Lara Austermann, Tim Dretzler, Katharina Ebel and Matthias Kerkemeyer for

their assistance with data collection, and finally Christian Gaser for helpful comments concerning the

CAT12 toolbox.

Conflicts of interest

The authors declare no conflict of interest.

Corresponding author

Correspondence should be addressed to Dr. Björn Horing, Department of Systems Neuroscience,

University Medical Center Hamburg-Eppendorf, Martinistrasse 52, 20246 Hamburg, Germany.

b.horing@uke.de

References

1.   Moayedi M, Davis KD. Theories of pain: from specificity to gate control. J Neurophysiol.

2013;109(1):5–12. doi: 10.1152/jn.00457.2012

2.   International Association for the Study of Pain (IASP). Pain terms: a list with definitions and

notes on usage. Recommended by the IASP Subcommittee on Taxonomy. Pain. 1979;6(3):249.

3.   Tracey I, Mantyh PW. The cerebral signature for pain perception and its modulation. Neuron.

2007;55(3):377–91. doi: 10.1016/j.neuron.2007.07.012

4.   Garcia-Larrea L, Peyron R. Pain matrices and neuropathic pain matrices: A review. Pain.

2013;154:S29–43. doi: 10.1016/j.pain.2013.09.001

5.   Mouraux A, Diukova A, Lee MC, Wise RG, Iannetti GD. A multisensory investigation of the

functional significance of the "pain matrix." Neuroimage. 2011;54(3):2237–49. doi:

10.1016/j.neuroimage.2010.09.084

6.   Legrain V, Iannetti GD, Plaghki L, Mouraux A. The pain matrix reloaded: A salience detection

system for the body. Prog Neurobiol. 2011;93(1):111–24. doi:

10.1016/j.pneurobio.2010.10.005

7.   Davis KD, Bushnell MC, Iannetti GD, St Lawrence K, Coghill R. Evidence against pain specificity

in the dorsal posterior insula. F1000Research. 2015;4(2):362. doi:

10.12688/f1000research.6833.1

8.   Salomons T V., Iannetti GD, Liang M, Wood JN. The "pain matrix" in pain-free individuals.

JAMA Neurol. 2016;73(6):4–5. doi: 10.1001/jamaneurol.2016.0653

9.   Segerdahl AR, Mezue M, Okell TW, Farrar JT, Tracey I. The dorsal posterior insula subserves a

fundamental role in human pain. Nat Neurosci. 2015;18(4):499–500. doi: 10.1038/nn.3969

10.   Mouraux A, Iannetti GD. The search for pain biomarkers in the human brain. Brain.

2018;141(12):3290–307. doi: 10.1093/brain/awy281

11.   Büchel C, Geuter S, Sprenger C. Comparing Painful Stimulation vs Rest in Studies of Pain.

JAMA Neurol. 2016;73(10):1258. doi: 10.1001/jamaneurol.2016.2989

12.   Bornhövd K, Quante M, Glauche V, Bromm B, Weiller C, Büchel C. Painful stimuli evoke

different stimulus-response functions in the amygdala, prefrontal, insula and somatosensory

cortex: a single-trial fMRI study. Brain. 2002;125(Pt 6):1326–36.

13.   Loggia ML, Edwards RR, Kim J, Vangel MG, Wasan AD, Gollub RL, et al. Disentangling linear

and nonlinear brain responses to evoked deep tissue pain. Pain. 2012;153(10):2140–51. doi:

10.1016/j.pain.2012.07.014

14.   Tiemann L, Hohn VD, Ta Dinh S, May ES, Nickel MM, Gross J, et al. Distinct patterns of brain

activity mediate perceptual and motor and autonomic responses to noxious stimuli. Nat

Commun. 2018;9(1):4487. doi: 10.1038/s41467-018-06875-x

15.   Boucsein W. Electrodermal activity. 2nd ed. New York: Springer; 2012.

16.   Bastuji H, Frot M, Perchet C, Hagiwara K, Garcia-Larrea L. Convergence of sensory and limbic

noxious input into the anterior insula and the emergence of pain from nociception. Sci Rep.

2018;8(1):13360. doi: 10.1038/s41598-018-31781-z

17.   Bastuji H, Frot M, Perchet C, Magnin M, Garcia-Larrea L. Pain networks from the inside:

Spatiotemporal analysis of brain responses leading from nociception to conscious perception.

Hum Brain Mapp. 2016;37(12):4301–15. doi: 10.1002/hbm.23310

18.   Pomares FB, Faillenot I, Barral FG, Peyron R. The "where" and the "when" of the BOLD

response to pain in the insular cortex. Discussion on amplitudes and latencies. Neuroimage.

2013;64:466–75. doi: 10.1016/j.neuroimage.2012.09.038

19.   Roy M, Shohamy D, Daw N, Jepma M, Wimmer GE, Wager TD. Representation of aversive

prediction errors in the human periaqueductal gray. Nat Neurosci. 2014;17(11):1607–12. doi:

10.1038/nn.3832

20. D'Hondt F, Lassonde M, Collignon O, Dubarry A-S, Robert M, Rigoulot S, et al. Early brain-body

impact of emotional arousal. Front Hum Neurosci. 2010;4:33. doi:

10.3389/fnhum.2010.00033

21. Bradley MM, Codispoti M, Cuthbert BN, Lang PJ. Emotion and motivation I: Defensive and

appetitive reactions in picture processing. Emotion. 2001;1(3):276–98. doi: 10.1037/1528-

3542.1.3.276

22. Kayser C, Petkov CI, Lippert M, Logothetis NK. Mechanisms for Allocating Auditory Attention:

An Auditory Saliency Map. Curr Biol. 2005;15(21):1943–7. doi: 10.1016/j.cub.2005.09.040

23. Baliki MN, Geha PY, Apkarian A V. Parsing pain perception between nociceptive

representation and magnitude estimation. J Neurophysiol. 2009;101(2):875–87. doi:

10.1152/jn.91100.2008

24. Hutchison JL, Hubbard NA, Brigante RM, Turner M, Sandoval TI, Hillis GAJ, et al. The efficiency

of fMRI region of interest analysis methods for detecting group differences. J Neurosci

Methods. 2014;226:57–65. doi: 10.1016/j.jneumeth.2014.01.012

25. Geuter S, Boll S, Eippert F, Büchel C. Functional dissociation of stimulus intensity encoding and

predictive coding of pain in the insula. Elife. 2017;6. doi: 10.7554/eLife.24770

26. Moayedi M, Weissman-Fogel I. Is the insula the "how much" intensity coder? J Neurophysiol.

2009;102(3):1345–7. doi: 10.1152/jn.00356.2009

27. Taylor KS, Seminowicz DA, Davis KD. Two systems of resting state connectivity between the

insula and cingulate cortex. Hum Brain Mapp. 2009;30(9):2731–45. doi: 10.1002/hbm.20705

28. Eckert MA, Menon V, Walczak A, Ahlstrom J, Denslow S, Horwitz A, et al. At the heart of the

ventral attention system: the right anterior insula. Hum Brain Mapp. 2009;30(8):2530–41. doi:

10.1002/hbm.20688

29.   Mazzola L, Isnard J, Peyron R, Mauguière F. Stimulation of the human cortex and the

707   experience of pain: Wilder Penfield's observations revisited. Brain. 2012;135(2):631–40. doi:

708   10.1093/brain/awr265

709   30.   Mazzola L, Faillenot I, Barral F-G, Mauguière F, Peyron R. Spatial segregation of somato-

710   sensory and pain activations in the human operculo-insular cortex. Neuroimage.

711   2012;60(1):409–18. doi: 10.1016/j.neuroimage.2011.12.072

712   31.   Ostrowsky K, Magnin M, Ryvlin P, Isnard J, Guenot M, Mauguière F. Representation of pain

713   and somatic sensation in the human insula: a study of responses to direct electrical cortical

714   stimulation. Cereb Cortex. 2002;12(4):376–85.

715   32.   Garcia-Larrea L, Mauguière F. Pain syndromes and the parietal lobe. In: Handbook of Clinical

716   Neurology. 2018 [cited 2019 Jan 18]. p. 207–23. doi: 10.1016/B978-0-444-63622-5.00010-3

717   33.   Liberati G, Klöcker A, Safronova MM, Ferrão Santos S, Ribeiro Vaz J-G, Raftopoulos C, et al.

718   Nociceptive Local Field Potentials Recorded from the Human Insula Are Not Specific for

719   Nociception. Apkarian AV, editor. PLOS Biol. 2016;14(1):e1002345. doi:

720   10.1371/journal.pbio.1002345

721   34.   Baliki MN, Geha PY, Fields HL, Apkarian AV. Predicting value of pain and analgesia: Nucleus

722   accumbens response to noxious stimuli changes in the presence of chronic pain. Neuron.

723   2010;66(1):149–60. doi: 10.1016/j.neuron.2010.03.002

724   35.   Price DD, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio

725   scale measures for chronic and experimental pain. Pain. 1983;17(1):45–56. doi:

726   10.1016/0304-3959(83)90126-4

727   36.   Duncan GH, Bushnell CM, Lavigne GJ. Comparison of verbal and visual analogue scales for

728   measuring the intensity and unpleasantness of experimental pain. Pain. 1989;37(3):295–303.

729   doi: 10.1016/0304-3959(89)90194-2

37.   Quiton RL, Greenspan JD. Across- and within-session variability of ratings of painful contact

heat stimuli. Pain. 2008;137(2):245–56. doi: 10.1016/j.pain.2007.08.034

38.   Lashkari D, Vul E, Kanwisher N, Golland P. Discovering structure in the space of fMRI

selectivity profiles. Neuroimage. 2010;50(3):1085–98. doi: 10.1016/j.neuroimage.2009.12.106

39.   Kanwisher N. Functional specificity in the human brain: A window into the functional

architecture of the mind. Proc Natl Acad Sci. 2010;107(25):11163–70. doi:

10.1073/pnas.1005062107

40.   Beck AT, Steer RA, Brown GK. Manual for the Beck Depression Inventory-II. San Antonio, TX:

Psychological Corporation Press; 1996.

41.   Kroenke K, Spitzer RL, Williams JBW. The PHQ-15: validity of a new measure for evaluating the

severity of somatic symptoms. Psychosom Med. 2002;64(2):258–66.

42.   Benedek M, Kaernbach C. A continuous measure of phasic electrodermal activity. J Neurosci

Methods. 2010;190(1):80–91. doi: 10.1016/J.JNEUMETH.2010.04.028

43.   Geuter S, Gamer M, Onat S, Büchel C. Parametric trial-by-trial prediction of pain by easily

available physiological measures. Pain. 2014;155(5):994–1001. doi:

10.1016/j.pain.2014.02.005

44.   Loggia ML, Juneau M, Bushnell MC. Autonomic responses to heat pain: Heart rate, skin

conductance, and their relation to verbal ratings and stimulus intensity. Pain.

2011;152(3):592–8. doi: 10.1016/j.pain.2010.11.032

45.   Bach DR, Flandin G, Friston KJ, Dolan RJ. Modelling event-related skin conductance responses.

Int J Psychophysiol. 2010;75(3):349–56. doi: 10.1016/J.IJPSYCHO.2010.01.005

46.   Coghill RC, Sang CN, Maisog JM, Iadarola MJ. Pain Intensity Processing Within the Human

Brain: A Bilateral, Distributed Mechanism. J Neurophysiol. 1999;82(4):1934–43. doi:

10.1152/jn.1999.82.4.1934

47.   Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, et al. A new SPM toolbox

755   for combining probabilistic cytoarchitectonic maps and functional imaging data. Neuroimage.

756   2005;25(4):1325–35. doi: 10.1016/J.NEUROIMAGE.2004.12.034

757   48.   Rolke R, Magerl W, Campbell KA, Schalber C, Caspari S, Birklein F, et al. Quantitative sensory

758   testing: A comprehensive protocol for clinical trials. Eur J Pain. 2006;10(1):77–88. doi:

759   10.1016/j.ejpain.2005.02.003

760   49.   Ellerbrock I, May A. MRI scanner environment increases pain perception in a standardized

761   nociceptive paradigm. Brain Imaging Behav. 2015;9(4):848–53. doi: 10.1007/s11682-014-

762   9345-5

763   50.   Awiszus F. TMS and threshold hunting. Suppl Clin Neurophysiol. 2003;56:13–23. doi:

764   10.1016/S1567-424X(09)70205-3

765   51.   Hall DA, Summerfield AQ, Gonçalves MS, Foster JR, Palmer AR, Bowtell RW. Time-course of

766   the auditory BOLD response to scanner noise. Magn Reson Med. 2000;43(4):601–6.

767   52.   National Institute for Occupational Safety and Health (NIOSH). Criteria for a Recommended

768   Standard: Occupational Noise Exposure. Revised Criteria [Internet]. 1998.

769   53.   Kass RE, Raftery AE. Bayes Factors. J Am Stat Assoc. 1995;90(430):773–95. doi:

770   10.1080/01621459.1995.10476572

771   54.   Rouder JN, Morey RD. Default Bayes Factors for model selection in regression. Multivariate

772   Behav Res. 2012;47(6):877–903. doi: 10.1080/00273171.2012.734737

773   55.   Raudenbush SW, Bryk AS. Hierarchical linear models: Applications and data analysis.

774   Thousand Oaks, CA: Sage; 2002.

775   56.   Hofmann DA, Gavin MB. Centering decisions in Hierarchical Linear Models: Implications for

776   research in organizations. J Manage. 1998;24(5):623–41. doi: 10.1177/014920639802400504

Supporting information

 S1 Table. Sample descriptive statistics.

Questionnaire	Construct	Mean±SD	Sample range	Possible range
BDI-II [SR 1,2]	Depression	2.0±2.2	0-8	0-63
PHQ15 [SR 3]	Somatization	3.9±2.3	0-9	0-30
FPQ [SR 4]				
severe	Fear of pain	28.9±8.3	11-42	10-50
minor	Fear of pain	14.0±4.5	10-27	10-50
PVAQ [SR 5]	Pain vigilance and awareness	34.6±8.7	21-54	0-80
PSQ [SR 6]	Pain sensitivity	45.1±14.8	17-73	0-140
PRSS [SR 7]				
Catastrophizing	Pain catastrophizing	8.6±5.4	2-21	0-45, higher more catastrophizing
Coping	Pain coping	29.8±5.2	19-38	0-45, higher more active coping
STAI [SR 8,9]				
Trait	Trait anxiety	32.0±6.9	21-49	20-80
State	State anxiety (pre experiment)	31.5±6.2	22-51	20-80
MDMQ [SR 10]				
GoodBad A	Mood: Good vs bad (pre exp.)	17.7±1.9	12-20	4-24, the higher the better
AwakeTired A	Mood: Awake vs tired (pre exp.)	15.2±2.9	8-19	4-24, the higher the more awake

CalmNervous A	Mood: Calm vs nervous (post exp.)	16.5±2.6	10-20	4-24, the higher the calmer
GoodBad B	Mood: Good vs bad (pre exp.)	17.2±1.7	12-19	4-24, the higher the better
AwakeTired B	Mood: Awake vs tired (pre exp.)	10.1±2.4	6-16	4-24, the higher the more awake
CalmNervous B	Mood: Calm vs nervous (post exp.)	17.2±2.1	12-20	4-24, the higher the calmer

782   **S2 Table**. Average calibrated sound intensities 1 through 6, which were used as stimuli during the experiment.

783   Cohort 2 received higher intensities, see Methods for rationale.

	Cohort 1 (n=15)			Cohort 2 (n=11)		
Intensity	Target VAS	dBA mean±SD	Range	Target VAS	dBA mean±SD	Range
1	25	70.5±7.4	57.7-82.9	48	86.1±5.9	77.8-99.3
2	35	74.2±6.6	62.4-85.9	59	92.2±5.9	81.2-101.5
3	45	77.9±6.0	67.2-88.7	70	95.6±6.0	84.4-102.5
4	55	81.4±5.5	70.8-91.2	82	97.7±5.0	87.7-103.0
5	65	84.8±5.3	74.3-93.4	93	99.1±3.9	90.5-103.0
6	75	87.8±5.2	77.8-95.3	105	100.3±3.0	93.3-103.0

787　**S1 Figure**. Binary and signed masks used for analyses. **A.** Binary mask used for small volume correction used for

788　all analyses (unless otherwise noted), delineated by the black line. **B.** Signed mask used for covering heat

789　(orange) or sound (blue) contrasts.

793　**S2 Figure**. Location of peak voxels for modality main effects (H1 through H3 for heat, S1 for sound). Also see

794　Figure 4.

Parietal operculum (SII)

57 -30 31

798   **S3 Figure**. Location of peak voxel for parametric modulation by heat intensity. Also see Figure 5.

A

[H1]   55 -37 26

[H2]   58 -14 18

[H3]   36 0 14

[H4]   41 1 14

801   **S4 Figure**. Location of peak voxels for parametric modulation by ratings (H1 through H4). Also see Figure 6.

Parietal operculum (SII)

56 -37 25

S5 Figure. Location of peak voxel for activation corresponding to the three axioms. Also see Figure 7.

Supporting information – References

SR 1. Beck AT, Steer RA, Brown GK. Manual for the Beck Depression Inventory-II. San Antonio, TX:

Psychological Corporation Press; 1996.

SR 2. Hautzinger M, Keller F, Kühner C. BDI-II. Beck-Depressions-Inventar. Revision. 2nd ed.

Frankfurt: Pearson Assessment; 2009.

SR 3. Kroenke K, Spitzer RL, Williams JBW. The PHQ-15: validity of a new measure for evaluating the

severity of somatic symptoms. Psychosom Med. 2002;64(2):258–66.

SR 4. McNeil DW, Rainwater AJ. Development of the Fear of Pain Questionnaire-III. J Behav Med.

1998;21(4):389–410.

SR 5. McCracken LM. "Attention" to pain in persons with chronic pain: A behavioral approach.

Behav Ther. 1997;28(2):271–84. doi: 10.1016/S0005-7894(97)80047-0

SR 6. Ruscheweyh R, Marziniak M, Stumpenhorst F, Reinholz J, Knecht S. Pain sensitivity can be

assessed by self-rating: Development and validation of the Pain Sensitivity Questionnaire.

Pain. 2009;146(1):65–74. doi: 10.1016/j.pain.2009.06.020

SR 7. Flor H, Behle DJ, Birbaumer N. Assessment of pain-related cognitions in chronic pain patients.

Behav Res Ther. 1993;31(1):63–73.

SR 8. Spielberger CD, Gorsuch RL, Lushene RE. Manual for the State-Trait Anxiety Inventory. Palo

Alto, CA: Consulting Psychologists Press; 1970.

SR 9. Laux L, Glanzmann P, Schaffner P, Spielberger CD. Das State-Trait-Angstinventar. Weinheim:

Beltz; 1981.

SR 10. Steyer R, Schwenkmezger P, Notz P, Eid M. Der Mehrdimensionale Befindlichkeitsfragebogen

(MDBF). Handanweisung. Göttingen: Hogrefe; 1997.