A stable pollination environment limits current but not potential evolution of

floral traits

Maria Clara Castellanos[1,2], Javier Montero-Pau[3,4], Peio Ziarsolo[3], Jose Miguel Blanca[3], Joaquin

Cañizares[3], Juli G. Pausas[2]

[1] School of Life Sciences, University of Sussex, Brighton, BN1 9QG, UK

[2] CIDE-CSIC, Montcada, Valencia, Spain

[3] COMAV, Universitat Politècnica de València, Valencia, Spain

[4] Department of Biochemistry and Molecular Biology, Universitat de València, Valencia, Spain

Corresponding author: *Maria Clara Castellanos* *m.c.castellanos@sussex.ac.uk*

Running title: Floral evolutionary potential

Abstract

Plants' vast variation in floral traits at a macroevolutionary level is often interpreted as the result of

adaptation to pollinators. However, field studies often find no evidence of pollinator-mediated

selection on flowers. This could be explained by periods of stasis, when selection is relaxed under

stable conditions, followed by pollinator changes that provide innovative selection. We asked if

periods of stasis are caused by stabilizing or absence of other forms of selection on floral traits, or by

low trait heritability even if selection is present. We studied *Ulex parviflorus*, a plant predominantly

pollinated by one bee species across its range. We measured heritability and evolvability of floral

traits, using genome-wide molecular relatedness in a wild population, and combined this with

estimates of selection. We found evidence for both stabilizing selection and low trait heritability as

explanations for stasis in flowers. The area of the standard petal is under stabilizing selection, but

the variability observed in the wild is not heritable. A separate trait, floral size, in turn presents high

heritability, but is not currently under selection. We show how a stable pollination environment can

lead to a lack of evolutionary change, yet maintain heritable variation to respond to future selection

pressures.

Keywords: evolvability, floral evolution, heritability, keel flower, quantitative genetics in the wild,

Ulex parviflorus

Introduction

Flowering plants exhibit a striking diversity in floral form and function, and because flowers are reproductive organs, the causes and dynamics of their evolution are crucial for understanding plant biodiversity. Much of the variation in floral traits at a macroevolutionary level is often interpreted as the result of adaptations to pollinators (Fenster et al. 2004). Experimental studies also confirm that many floral traits can be subject to selection by pollinators (reviewed by Parachnowitsch and Kessler 2010, Caruso et al. 2018). However, field studies measuring pollinator-mediated evolution of floral traits often find sporadic and erratic evidence for strong selection taking place in wild populations (Harder and Johnson 2009). In their review, Harder and Johnson found that only about 1/3 of the studies reported significant selection on floral traits. A possible reason for this 'paradox' is the likely prevalence of periods of stasis, where pollinator-mediated selection on flowers is relaxed under stable conditions, interrupted by more unstable periods where pollinator changes can provide innovative selection (e.g. Galen 1989, Harder and Johnson 2009).

For pollinator-mediated evolution to take place in the wild, floral phenotypic traits must not only be under selection but also harbour enough heritable variation. Periods of stasis can thus be the consequence of stabilizing or a lack of directional/disruptive selection on traits, or alternatively, they can also be the result of low levels of heritable variation even if selection is present. An appropriate model to study the role of these two non-exclusive scenarios would be a plant with a stable single dominant pollinator. Under these stable conditions, floral traits can be expected to experience low levels of pollinator-driven innovative selection, but still be heritable. Heritable variation in floral traits has been shown for numerous species in the greenhouse (reviewed in Ashman and Majetic 2006, Opedal 2018), and in a few field studies (Schwaegerle and Levin 1990, Mazer and Schick 1991, Campbell 1996, Galen 1996). Thus a relaxation of selection could be the most likely explanation for stasis in floral traits in populations with stable pollination environments.

However, it also possible that trait heritability is lower in wild conditions than indicated by estimates under artificially reduced environmental variation. Traditional greenhouse and common garden studies of heritability allow for control of local environments and genetic background, but heritability values measured under controlled conditions can be systematically overestimated compared with wild conditions (Conner et al. 2003, Winn 2004). This can be caused by higher environmental variability in the field, as well as decreased expression of additive variance, or potential differences in survival, all leading to smaller heritability estimates. The alternative of

measuring heritability directly in the field, although being more realistic, was until recently

constrained by difficulties in designing complex crossing and planting experiments (see Campbell

1996), or in establishing relatedness among individual plants growing in the wild. This has now

changed thanks to access to large and highly informative molecular markers (Castellanos et al. 2011,

Stanton-Geddes et al. 2013). Using genome-wide markers to measure genetic similarity of plants

growing in the wild (in the form of a relatedness matrix, G), it is possible to estimate the proportion

of the phenotypic covariance that is explained by relatedness (i.e. heritability) in the focal floral trait

(Ritland 1996). This approach can incorporate environmental factors in the statistical estimation of

heritability, to provide us with an ecologically realistic view of what plant populations are

experiencing in natural conditions and help us understand the role of genetic variation in evolution

(Campbell 1996, Kruuk et al. 2014).

We study the consequences of a stable pollination environment on floral traits by focusing on a plant

with a dominant pollinator, the Mediterranean gorse (*Ulex parviflorus*). Observations across its

current distribution show that honey bees (*Apis mellifera*) are currently the prevailing pollinator,

with consistently low visitation rates, including in areas with low human influence. Dominance of

honey bee visitation was observed by Herrera (1988; no % of visits reported) and Reverté et al.

(2016; 63% of visits were by honey bees) in coastal populations in southern and eastern Spain

respectively, and has also been observed in inland populations in Cazorla, Spain (93% of visits; C.M.

Herrera pers com.). Pollinator-mediated selection on flowers is expected in this plant because *Ulex*

and relatives (the large legume subfamily Faboidae) often have complex irregular butterfly-type

flowers ("papilionoid" or "keel" flowers, Fig. 1) believed to be specialized on bee pollination, with

traits that both enhance pollinator attraction and mechanical interactions that improve pollination

(Westerkamp 1997). In such system, we predict 1) a relaxation of directional or disruptive (i.e.

innovative) selection of floral traits, and/or 2) low trait heritability as a consequence of genetic

erosion over time.

To test these predictions, we measured trait heritability and natural selection on the same plant

individuals in a wild population; this allowed us to assess the potential for evolution in response to

current and future selection. To our knowledge, this is the first time this approach is used

successfully to study floral traits. We measured floral morphology and pollinator visits, along with

natural selection, genetic correlation, evolvability, and heritability of the floral traits to 1) determine

if floral traits in a stable pollination environment are currently under selection and show heritable

102 variation, thus evolving in response to selection, and 2) if not, to establish if the causes for a lack of

103 evolutionary response are related to low selection, low heritability or both.

105 **Materials and Methods**

107 **Study species and sampling locations**

108 *Ulex parviflorus* Pourr. (Mediterranean gorse; Fabaceae) is a thorny perennial shrub that lacks true

109 leaves in the adult stage and grows up to 2 m. Species in the genus *Ulex* have yellow hermaphroditic

110 flowers visited and pollinated by large-bodied bees, in a similar way to other species in the tribe

111 Genistae (Herrera 2001). Flowers do not produce nectar and the bees visit to collect pollen, but to

112 be able to do so, they need to be heavy enough to actively trigger the explosive mechanism for

113 pollen release. Reproductive organs in these flowers are enclosed by specialized petals, the keel and

114 the wings (Fig. 1). The insect presses the keel petals with the hind legs and this pressure powerfully

115 releases the concealed stamens and stigma upwards, placing a cloud of pollen grains on the ventral

116 side of the bee. After a visit flowers do not recover their original shape, with stigmas and style now

117 protruding from the keel, and are rarely visited by large bees again, but can receive visits by smaller

118 insects like hoverflies and solitary bees. *Ulex parviflorus* is self-compatible but depends on

119 pollinators to set fruit (Herrera 1987). Flowering starts in the winter and can last for a few months

120 into the spring.

122 The species is widespread along the western Mediterranean coast from southern France to southern

123 Portugal. It is a successful colonizer of oldfields resulting from abandoned human activities, as well

124 as recently burnt areas, thanks to numerous adaptations to recruitment after fire (Pausas et al.

125 2012, Pausas and Moreira 2012, Pausas et al. 2017). The seeds form a persistent bank in the soil,

126 where they remain dormant until the heat produced during a fire breaks dormancy and stimulates

127 germination in post-fire conditions (Moreira et al. 2010). Current landscapes in eastern Spain are a

128 mosaic of oldfields and postfire shrubland (Pausas & Millán 2019), where *Ulex parviflorus* is very

129 abundant and distributed continuously from the lowlands up to 900m of altitude (Fig S1 in

130 supplementary materials). As a consequence there is very low genetic differentiation in the study

131 area (Moreira et al. 2014; see also Supplementary methods and Fig. S3), and different stands cannot

132 be considered distinct populations. For our sampling, we selected six sites within this continuous

133 population, aiming to capture the variability of mature *U. parviflorus* stands in the area (Table S1,

134 Fig. S1). By sampling at different altitudes, for example, we include variability in floral traits along the

135 elevational gradient (Fig. S2 in supplementary materials). At each site, we tagged 40 individual plants

(240 plants in total) for phenotypic and genotypic characterization as described below. Individuals

were at least 5 m apart from each other and blooming at the time of sampling.

Pollinator censuses

To quantify the diversity of floral visitors and visitation rates, we ran multiple three-minute long

pollinator censuses at different times of the day, for up to five hours of observations per locality, on

two separate days during peak blooming in 2014. We also ran censuses in two localities in 2013,

again during peak blooming (Montserrat and Cheste). Each census recorded the number and identity

of visitors to patches of flowers on haphazardly chosen individuals. We counted the number of

flowers included in each census to estimate the per-flower visitation rate.

Floral phenotypes

We collected five haphazardly selected flowers from each individual plant for phenotypic

characterization of two floral traits that function as proxies for flower showiness and flower size.

The area of the upwards-facing petal, or standard, plays a key role in flower showiness, as it is the

largest and more visible petal in these typical papilionoid flowers (Fig.1; standard petals are also

often called flag or banner petals). We removed standards from all flowers when fresh, pressed

them flat individually in a plant press until dry. We then used scanned images of the standards to

measure their surface area with the Image-J analysis freeware (Schneider et al. 2012).

Flower size is important in the Genistae as it can determine the size of the insects that can visit the

flowers (Herrera 2001, Córdoba and Cocucci 2011). Size was estimated as the dry weight of flowers

(calyx and corolla) after removing the standard petal and the pedicel, and carefully brushing off all

pollen grains. Flowers were pressed and oven-dried at 40ºC for 48 hours and weighed to the nearest

0.01 mg.

These traits were chosen because they can be expected to play an important role in the interaction

with pollinators and thus be under natural selection driven by pollinators (see Study Species above).

As is the case in many complex flowers, the two traits studied can be expected to co-vary (Herrera

2001), and analyses below are designed to take this into consideration. We have no reason to

suspect that there is variability in these traits with flower age (see also Herrera 2001). We have

never observed florivory in this species and thus doubt that herbivores will directly select for the two

focal traits in this study.

Plant genotyping

Fresh terminal twigs were collected from each tagged individual plant and dried in silica gel previous

to DNA extraction. The extraction was performed using the Speedtools plant DNA extraction kit

(Biotools, Madrid, Spain), with modifications to the manufacturer's protocol to optimize DNA

quantity and quality extracted for this highly lignified species. We used the Genotyping-by-

Sequencing (GBS) protocol to identify single nucleotide polymorphisms (SNPs) across the genome

(Elshire et al. 2011). Illumina libraries for our 240 individuals were constructed by digesting genomic

DNA with a restriction enzyme. The GBS protocol was followed twice for each plant after separate

digestions with *PstI* and *EcoT22I*, in order to increase the number of high quality SNPs. Library

construction and sequencing was performed by the Genomic Diversity facility at Cornell University

(USA). SNP calling was implemented using the UNEAK pipeline (Lu et al. 2013) in the TASSEL v.3

software package (Bradbury et al. 2007), designed for data sets without a reference genome.

The final SNP dataset used for the analysis of relatedness below excluded loci that were not

genotyped in at least 90% of individual plants. The minimum allele frequency allowed to retain loci

was set to MAF > 0.01. We also excluded individuals with low genotyping rates (under 85% of loci).

After applying these filters, we also manually removed remaining loci with extreme values of

observed heterozygosity (under 2% and higher than 98%), after estimating oHET with PLINK

command –Hardy (Purcell et al. 2007).

Fitness estimates and phenotypic selection

We estimated fruit set in the same 40 individual plants in each locality as a proxy for female

reproductive success. For this, we labelled a representative flowering twig in each plant during

flowering peak. When fruits were already developing (browning capsules) a few weeks later, we

collected the labelled twig in a paper envelope. Back in the laboratory we measured 10 cm of twig to

calculate a) the number of fruits developing normally, and b) scars left by all flowers produced by

the twig, clearly visible under a dissecting microscope. From this we calculated fruit set as the

proportion of flowers that develop into a fruit. The majority of fruits had one (71% of 3200 fruits

examined) or two seeds (25%), with a mean number of 1.22 seeds/fruit across all individuals.

We estimated selection parameters to test for both linear and non-linear selection on the two floral

traits, using fruit set as the response fitness variable in the models. Because floral weight and

standard area show a significant phenotypic correlation (even though floral weight did not include

the standard, Pearson r = 0.43, P < 0.001), we estimated selection gradients in addition to selection

differentials. Selection differentials provide univariate estimates of selection without considering
other traits, while gradients provide estimates on correlated traits. By estimating the four selection
parameters - standardized linear (S), and quadratic (c) selection differentials, and standardized linear
(β) and quadratic (γ) selection gradients - we can explore direct and indirect selection on the floral
traits. Linear parameters test for directional selection, while quadratic parameters measure
potential stabilizing or disruptive selection.
210
We used generalised additive models (GAM) to measure selection parameters on absolute fitness
values, following the approach developed by Morrisey and Sakreda (2013). This approach provides
quantitative estimates of selection differentials and gradients for non-normal fitness components,
testing for both linear and quadratic selection. We fitted GAMs for binomial fruit set data (fruits
developed in relation total flowers), using a logit link function and assuming a binomial error
distribution with the *mgcv* package in R. We used univariate GAMs to estimate selection
differentials, and included both floral traits into a bivariate model to estimate selection gradients. To
control for local effects, we included locality as a random factor in all models. Models included
additive spline effects on all factors. Differential and gradient parameters were estimated based on
numerical approximations of first and second partial derivatives of relative fitness, averaged over
the distribution of observed phenotype. To calculate the significance of selection differentials and
gradients, we used the bootstrap approach (n= 1000 samples) implemented in the *gsg* package in R
(Morrissey and Sakrejda 2013).
224
SNP-based relatedness and quantitative genetic parameters
Pairwise relatedness between all pairs of individuals was estimated from the similarity of their SNP
genotypes. To estimate G, the genome-wide relatedness matrix among all pairs of individuals, we
used the realized relatedness method of VanRaden (2008) and Astle and Balding (2009) as
implemented in the *kin* function of package *synbreed* in *R* (Wimmer et al. 2012; see details in
Supplementary methods). Relatedness values under this approach are a measure of excess allele
sharing compared to unrelated individuals. As a consequence, negative values can be common and
correspond to individuals sharing fewer alleles than expected given the sample.
To estimate additive genetic variance (and then heritability and evolvability) we used a linear
mixed 'animal model' approach to model the phenotypic variance in floral traits while including the
variance explained by relatedness (Wilson et al. 2010). We included the elevation above sea level as
a fixed effect to account for environmental variability among plants, because elevation is the main
factor that varies among localities (Fig. S2) and this could affect floral traits as in other species

(Herrera 2005). In addition to the additive genetic effects (see model below), models included two

more random effects: the site of origin of each plant, to account for unmeasured local

environmental effects that could co-vary with genetic variation, and the individual identity to

account for intra-individual effects (a "permanent environment" effect in Wilson et al. 2010),

because we had five flower replicates per plant. We ran a univariate model for each of the two

floral traits studied, specified as:

245                              $y = X\beta + Z_1a + Z_2s + Z_3i + e$

where y is the vector of floral trait values, β is the vector of fixed effects (with X as the incidence

matrix), Z_1, Z_2 and Z_3 are incidence matrices for the random effects a (individual identity to partition

additive genetic effects), s (the locality), i (individual identity to model intra-individual effects caused

by differences among replicate flowers from the same individual), and e is the residual error. The

variance-covariance structure of random factor a in the model is defined by $G \cdot V_a$, where G is the

genome-wide relatedness matrix between plant pairs, and V_a is the additive variance to be

estimated. To test for the effect of not including the spatial and environmental predictors in the

models, we also ran a 'naïve' version of each model that included only the relatedness and individual

effects (Castellanos et al. 2015). We ran Bayesian animal models using package *MCMCglmm* for *R*

(Hadfield 2010) with both floral weight and standard petal area modeled as continuous traits. For

modelling the standard area, we used parameter expanded priors for the distribution of variance

components following the χ^2 distribution with one degree of freedom. Each analysis was iterated

long enough to obtain 5000 independent chains (see supplementary methods and Table S2 for

model details, scripts and prior selection).

Narrow sense heritability (h^2) was then estimated as the proportion of the total phenotypic

variance assigned to the individual (i.e. to the additive genetic variance, V_a):

$$h^2 = \frac{V_a}{V_a + V_s + V_i + V_e}$$

where V_s is the variance explained by the site of origin, V_i is the intra-individual variance in the trait,

and V_e is the residual variance. We also estimated the narrow sense evolvability (*e*), i.e. the mean-

standardized additive genetic variance, $e = V_a / x^2$, where *x* is the trait mean. *e* reflects the expected

percentage of change of a trait under a unit strength of selection per generation (Houle 1992,

Hansen et al. 2003) and provides an estimate of evolvability that is independent of trait variation

and comparable across traits.

In addition, we estimated the genetic correlation (r_G) between floral weight and standard area by
running a bivariate animal model in *MCMCglmm*. In this case we used the same fixed and random
factors as in the univariate models above (see supplementary methods for prior information).
274

Results

Pollinators

We recorded 364 visits to 22522 censused flowers in 28 hours of observations across the six *U.*
parviflorus localities. Of those, 331 (92%) were visits by the honeybee *Apis mellifera*. Further 25 visits
were by *Bombus sp.* individuals (7%). The remaining 3 visits were to already open flowers by small
coleoptera and a hoverfly, both unlikely to contact stigmas and carry out pollination. Across sites, we
found an average visitation rate of 0.015 (±0.057) visits per 3-minute census to an individual flower,
which translates into a visit every 3.3 hours, on average. Visitation rates were similar when
comparing localities, except for one where visits were significantly more frequent (Simat average
visitation rate= 0.03 visits per census).
285

Floral phenotypes and selection

287 Flowers show considerable variation in the two traits measured, flower weight and standard petal
288 area, both within and across localities. A variance partition analysis showed that the variance in both
289 traits across the five flowers sampled per plant was negligible, so the selection analysis below was
290 run using mean floral values for each individual plant (see also Herrera 2001).
291

We found no evidence of linear directional selection on floral traits, either in univariate models (s
coefficients) or models of correlated selection incorporating both floral variables (β coefficients,
Table 1). However, we found evidence for univariate quadratic effects in both traits (c coefficients)
but only standard area shows significant quadratic gradients (γ coefficients). This suggests that floral
weight is not under direct selection, while there is strong evidence for stabilising selection on
standard petal area (Fig. 2).
298

Genomic markers and population genetic structure

The GBS sequencing approach yielded a large number of polymorphic SNPs across individuals
(261,775 SNPs before quality filtering). After MAF and heterozygosity filtering, we retained 10,421
high-quality SNPs that were present in at least 90% of individuals across all localities. The analyses
below use this dataset to estimate genomic relatedness; however, we also tested for the effect of
retaining a larger number of SNPs (with presence in at least 50% of the individuals, which leads to a

higher number of genotypes imputed by *synbreed*, see Supplementary methods). Analysis with this

larger dataset produced the same qualitative results, suggesting that retaining more (but highly

imputed) markers did not add valuable information on the relatedness among our study plants.

Therefore, all analyses below use the smaller dataset with 10,421 SNPs.

Heritability, evolvability and genetic correlation

Pairwise relatedness among sampled individuals varied markedly and was overall relatively low

(average values ranging from -0.09 to 0.79, but with most values <0.2), even within locality (Fig. S4),

supporting the prevalence of outcrossing in this species. The low population genetic structure and

the presence of variance in relatedness provide the conditions for a reliable estimation of heritability

in the field in this species (Ritland 1996).

We found significant estimates of heritability and evolvability in flower weight (h^2 = 0.14, e = 0.42%;

Table 2). For standard area, our models instead detected very low additive variance, yielding very

low h^2 and e in this case (h^2 = 0.001, e < 0.001%; Table 2). For both traits, Deviance Information

Criterion (DIC) values for the heritability naïve models were larger than for the complete model

(Table S2), indicating a better fit for the latter. The naïve models included only the relatedness

among individuals and neither environmental nor spatial predictors, and showed estimated h^2 values

substantially higher than our final estimates (Table 2).

Our bivariate analysis found a low genetic correlation between the two floral traits that is

indistinguishable from zero (r_G = 0.06); in addition, credible intervals were large (-0.139 to 0.381), so

we cannot confidently support the presence of a genetic correlation.

Discussion

We provide an example of a stable pollination environment that has led to a lack of innovative

selection, yet maintaining enough heritable variation for responding to possible novel selection

pressures, at least in some traits. In *Ulex parviflorus*, we found evidence for both stabilizing selection

and low trait heritability as alternative explanations for lack of evolution in flowers. Specifically, the

area of the standard petal is currently under stabilizing selection, but the variability we observe in

the field is not heritable. Floral weight, in turn, presents high heritability, but is not currently under

selection.

Stable pollinator communities are potentially a common feature for many plant species under even

environmental conditions. For the particular case of *Ulex parviflorus*, current evidence shows that

honey bees are the most frequent pollinators in all surveyed populations, including the one studied

here and other localities (Herrera 1988, Reverté et al. 2016, C.M. Herrera pers com). Other species in

the genus, including *U. europaeus, U. minor* and *U. galli*, present a higher diversity of large bees

among their visitors (several species of *Bombus* and *Andrena*; Kirchner and Bullock 1999, Bowman et

al. 2008, Falk 2011). The dominance of honey bees in *Ulex parviflorus* populations could be seen as a

consequence of the large anthropogenic influence across its range; however, *U. parviflorus*

populations in an area with low human influence and high pollinator diversity (Sierra de Cazorla, see

Herrera 2018) corroborates the predominance of honey bees as pollinators of this species.

Regardless of the reasons for the low pollinator diversity, our study provides evidence on how stable

conditions can lead to lack of current evolution in floral traits.

On the opposite side of the spectrum, field studies that do detect pollinator-mediated directional

selection on unmanipulated floral traits often focus on plants that are exposed to changing

pollinators, either in different parts of the species range (Herrera et al. 2006, Anderson et al. 2010)

or in hybrid contact zones where there is selection against hybridization (Campbell et al. 2018).

Taken together, current evidence supports the idea that pollination-driven floral evolution takes

place mostly during evolutionarily innovative periods driven by to changing pollinators.

Stabilizing selection is expected in floral traits that influence the accuracy of the flower-pollinator

interaction (Cresswell 2000, Armbruster et al. 2009). It is difficult to establish how common

stabilizing selection is on floral traits in wild plants, because studies do not measure non-linear

selection as often as directional selection (Harder and Johnson 2009, Caruso et al. 2018). For the

standard petal in *Ulex*, we detected stabilizing selection for intermediate surface area. The size of

this "flag" petal is expected to play an important role on pollinator attraction by increasing the floral

colourful display (Fig. 1), so that selection against smaller sizes is expected. Too large standard petals

could be selected against if they incur a higher cost for the plant. This cost could be even higher if

large standard petals are developmentally restricted to overall larger flowers; however, our genetic

correlation estimates suggest that the association of standard petal area with floral size is weak. This

is consistent with a previous study that carefully dissected the role of the different petals in another

keel flower; in *Collaea argentina,* Córdoba et al. (2015) found that the standard petal is not

functionally integrated with another set of floral traits that collectively regulate the enclosing

mechanism of stamens and pistil. That is, the mechanics of protecting the enclosed rewards in these

flowers can be independent of pollinator attraction as we expected, and selection can vary across

floral parts.

Floral morphological traits are often found to present heritable variation (reviewed by Ashman and

Majetic 2006, Opedal 2018); however, most of the studies in these reviews were performed in

controlled environments. Our field estimates of heritability fall within the lower range of those

summarized in Fig. 1 of Altman and Majetic (2006), as expected from field values compared to

greenhouse estimates. We found that flower weight shows significant heritability, but no detectable

heritability in the standard petal area. Comparing petals in papilionoid flowers, Herrera (2001) found

that the standard had higher phenotypic variance than other petals across Genisteae, and argued

that its role in pollination was smaller than for the keel petals, in a similar way as Córdoba et al.

(2015). This and our results suggest that this petal might be prone to high environmentally-induced

variation, which increases the exposure to stabilising selection, but does not lead to evolutionary

change.

Heritability estimates have been criticised as poor standardized measures of evolutionary potential

in realistic ecological settings, in part because of the covariance between environmental and genetic

effects (Houle 1992, Hansen et al. 2011). In this study, we estimate heritability directly in the field,

statistically controlling for environmental variation, and in the same individuals used to estimate

natural selection. In this context, field heritability estimates provide a very useful approach to

understand the current evolutionary potential at the population level, precisely because we are

interested in the role of environmental effects on the phenotypic variance, as exposed to natural

selection. An alternative measure of evolutionary potential, evolvability, uses the mean of trait

values to standardize the additive genetic variance and provides a comparable estimate of

proportional change in a trait value after selection (Hansen et al. 2003). Our estimates of evolvability

here confirm our findings in heritability, also showing near-zero evolutionary potential for the

standard petal area, but higher values for flower weight. In the latter case, evolvability is estimated

to be significant but small (under 1% of the trait mean value), suggesting that change in this trait

would not be fast unless submitted to strong selection. This value of evolvability is within the range

of evolvability values estimated for floral size specifically across plant species, as summarised in a

recent review (Opedal 2018).

Our estimate of genetic correlation between the two focal traits suffers from a low sample size to

run a bivariate animal model and needs to be interpreted with caution. However, the lack of a

genetic correlation is not surprising given that we cannot detect significant additive genetic variation
in one of the trait (the area of the standard petal). This does contrast with the fact that there is a
significant phenotypic correlation between the two traits, but as suggested by previous studies,
phenotypic correlations are not always good predictors of genetic correlations, even in highly
integrated organs as flowers (Gómez et al. 2009). Again, this is consistent with the decoupling of
petals found in a related species with keel flowers (Córdoba et al. 2015). It is thus possible that the
phenotypic correlation is caused by shared environmental factors that affect both traits in *Ulex*
flowers, further confirming the importance of studying evolutionary potential in field realistic
conditions.
415
Even though we could not detect a genetic correlation between the two floral traits studied here, a
caveat in our analysis is that we do not include selection on other (unmeasured) potentially
correlated traits. Another potential source of problems is that *Ulex* flowers are hermaphroditic and
thus likely subject to selection via both male and female reproductive success. Our estimates of
selection here are based on fruit set alone, and we cannot rule out that the two focal traits might be
under selection through male function (van Kleunen and Burczyk 2008). However, the two traits
studied here can be expected to affect pollen dispersal in similar ways as pollen deposition (and thus
seeds sired), because the trigger mechanism forces both male and female reproductive organs to
make contact with the bees at the same time. This means that factors affecting seed set and seed
sire are probably highly related in keel flowers.
426
This study adds to a series of recent works using large sets of molecular markers to study
quantitative genetics in wild populations, mostly focused on animals (Perrier et al. 2018), but also on
plants (Castellanos et al. 2015). Studies comparing the accuracy of SNP-based relatedness matrices
compared to pedigrees are consistently showing that they can be very good approximations, as long
as a large number of markers and a good sample of individuals is available (Bérénos et al. 2014,
Perrier et al. 2018). This is therefore an exciting time for studying the evolution of traits directly in
the wild, because field-based estimates of evolutionary potential provide new avenues to
understand basic evolutionary questions (such as stasis and the role of plasticity in trait variation),
but also the potential for wild organisms to respond to new selection pressures including those
imposed by anthropogenic environmental change. In the specific case of flowers, our findings
suggest that low-diversity pollination environments as those caused by anthropogenic pollination
declines can lead to reduced selection pressures, reduced opportunity for selection, and stasis
(Caruso et al. 2018), while exposure to new pollinators can lead to novel evolutionary change.

Conclusion

Relative stasis can be prevalent in contemporary populations, yet heritable phenotypic variance can be present; in combination with potential genetic correlations, this provides the potential to respond to novel selection. Selection on floral traits is not restricted to pollinators, as herbivores and abiotic factors can also be agents of selection (reviewed by Caruso et al. 2018). Regardless of the source of selection, our findings contribute to explain the macroevolutionary patterns of floral evolution where novel phenotypes are ubiquitous (exceptions are often related to very generalised pollination that is stable over evolutionary time, see Vasconcelos et al. 2019). Populations can experience stable conditions with undetectable innovative selection, but at the same time harbour genetically based variability to evolve under new conditions.

Acknowledgements

We are grateful to Abel Rubira, Santiago Donat-Caerols, Yedra García, Eva Sánchez, Paula Cassá and Lucia Tortajada for valuable help in the field. S. Donat-Caerols also provided help with genomic data analysis. Jorge Sellés helped with DNA extractions. We thank C.M. Herrera for sharing unpublished pollination data. Dr J. Brines kindly allowed access to his land in Simat. Financial support came from projects TREVOL and FILAS (CGL2012-39938, CGL2015-64086) from the Spanish Government. MCC was partly supported by the European Union's Horizon 2020 programme under the Marie Sklodowska-Curie grant agreement No 706365.

References

Anderson, B., R. Alexandersson, and S. D. Johnson. 2010. Evolution and coexistence of pollination ecotypes in an African *Gladiolus* (Iridaceae). Evolution **64**:960-972.

Armbruster, W. S., T. F. Hansen, C. Pélabon, R. Pérez-Barrales, and J. Maad. 2009. The adaptive accuracy of flowers: measurement and microevolutionary patterns. Annals of Botany **103**:1529-1545.

Ashman, T., and C. Majetic. 2006. Genetic constraints on floral evolution: a review and evaluation of patterns. Heredity **96**:343-352.

Astle, W., and D. J. Balding. 2009. Population structure and cryptic relatedness in genetic association studies. Statistical Science **24**:451-471.

Bérénos, C., P. A. Ellis, J. G. Pilkington, and J. M. Pemberton. 2014. Estimating quantitative genetic parameters in wild populations: a comparison of pedigree and genomic approaches. Molecular Ecology **23**:3434-3451.

Bowman, G., M. Tarayre, and A. Atlan. 2008. How is the invasive gorse Ulex europaeus pollinated during winter? A lesson from its native range. Plant Ecology **197**:197-206.

Bradbury, P. J., Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss, and E. S. Buckler. 2007. TASSEL: software for association mapping of complex traits in diverse samples. Bioinformatics **23**:2633-2635.

Campbell, D. R. 1996. Evolution of floral traits in a hermaphroditic plant: field measurements of heritabilities and genetic correlations. Evolution **50**:1442-1453.

Campbell, D. R., A. Faidiga, and G. Trujillo. 2018. Clines in traits compared over two decades in a
plant hybrid zone. Annals of Botany **122**:315–324.
Caruso, C. M., K. E. Eisen, R. A. Martin, and N. Sletvold. 2018. A meta-analysis of the agents of
selection on floral traits. Evolution.
Castellanos, M. C., J. M. Alcántara, P. J. Rey, and J. M. Bastida. 2011. Intra-population comparison of
vegetative and floral trait heritabilities estimated from molecular markers in wild *Aquilegia*
populations. Molecular Ecology **20**:3513-3524.
Castellanos, M. C., S. C. González-Martínez, and J. Pausas. 2015. Field heritability of a plant
adaptation to fire in heterogeneous landscapes. Molecular ecology **24**:5633-5642.
Conner, J. K., R. Franks, and C. Stewart. 2003. Expression of additive genetic variances and
covariances for wild radish floral traits: comparison between field and greenhouse
environments. Evolution **57**:487-495.
Córdoba, S. A., S. Benitez-Vieyra, and A. A. Cocucci. 2015. Functional modularity in a forcible flower
mechanism: relationships among morphology, biomechanical features and fitness.
Evolutionary ecology **29**:719-732.
Córdoba, S. A., and A. A. Cocucci. 2011. Flower power: its association with bee power and floral
functional morphology in papilionate legumes. Annals of Botany **108**:919-931.
Cresswell, J. E. 2000. Manipulation of female architecture in flowers reveals a narrow optimum for
pollen depostion. Ecology **81**:3244-3249.
Elshire, R. J., J. C. Glaubitz, Q. Sun, J. A. Poland, K. Kawamoto, E. S. Buckler, and S. E. Mitchell. 2011. A
robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PloS one
6:e19379.
Falk, S. J. 2011. A survey of the bees and wasps of fifteen chalk grassland and chalk heath sites within
the East Sussex South Downs. Self-published, downloaded from www.bwars.com (January
2019).
Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004. Pollination
syndromes and floral specialization. Annu. Rev. Ecol. Evol. Syst. **35**:375-403.
Galen, C. 1989. Measuring pollinator-mediated selection on morphometric floral traits: bumblebees
and the alpine sky pilot, Polemonium viscosum. Evolution **43**:882-890.
Galen, C. 1996. Rates of floral evolution: adaptation to bumblebee pollination in an alpine
wildflower, Polemonium viscosum. Evolution **50**:120-125.
Gómez, J., M. Abdelaziz, J. Muñoz-Pajares, and F. Perfectti. 2009. Heritability and genetic correlation
of corolla shape and size in *Erysimum mediohispanicum*. Evolution **63**:1820-1831.
Hadfield, J. D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The
MCMCglmm R Package. Journal of Statistical Software **33**:1-22.
Hansen, T. F., W. S. Armbruster, M. L. Carlson, and C. Pélabon. 2003. Evolvability and genetic
constraint in *Dalechampia* blossoms: genetic correlations and conditional evolvability.
Journal of Experimental Zoology **296B**:23-39.
Hansen, T. F., C. Pélabon, and D. Houle. 2011. Heritability is not evolvability. Evolutionary Biology
38:258.
Harder, L. D., and S. D. Johnson. 2009. Darwin's beautiful contrivances: evolutionary and functional
evidence for floral adaptation. New Phytologist **183**:530-545.
Herrera, C. M. 2018. Complex long-term dynamics of pollinator abundance in undisturbed
Mediterranean montane habitats over two decades. Ecological Monographs.
Herrera, C. M., M. C. Castellanos, and M. Medrano. 2006. Geographical context of floral evolution:
towards an improved research programme in floral diversification. Pages 278-294 *in* L. D.
Harder and S. C. H. Barrett, editors. Ecology and evolution of flowers. Oxford University
Press.
Herrera, J. 1987. Flower and fruit biology in southern Spanish Mediterranean shrublands. Annals of
the Missouri Botanical Garden **74**:69-78.

Herrera, J. 1988. Pollination relationships in southern Spanish Mediterranean shrublands. Journal of
Ecology:274-287.
Herrera, J. 2001. The variability of organs differentially involved in pollination, and correlations of
traits in Genisteae (Leguminosae: Papilionoideae). Annals of Botany **88**:1027-1037.
Houle, D. 1992. Comparing evolvability and variability of quantitative traits. Genetics **130**:195-204.
Kirchner, F., and J. Bullock. 1999. Taxonomic separation of *Ulex minor* Roth. and *U. gallii* Planch.:
morphometrics and chromosome counts. Watsonia **22**:365-376.
Kruuk, L. E., A. Charmantier, and D. Garant. 2014. The study of quantitative genetics in wild
populations. Quantitative genetics in the wild:1-15.
Mazer, S. J., and C. T. Schick. 1991. Constancy of population parameters for life-history and floral
traits in *Raphanus sativus* L. II. Effects of planting density on phenotype and heritability
estimates. Evolution **45**:1888-1907.
Moreira, B., M. C. Castellanos, and J. Pausas. 2014. Genetic component of flammability variation in a
M editerranean shrub. Molecular ecology **23**:1213-1223.
Moreira, B., J. Tormo, E. Estrelles, and J. Pausas. 2010. Disentangling the role of heat and smoke as
germination cues in Mediterranean Basin flora. Annals of Botany **105**:627-635.
Morrissey, M. B., and K. Sakrejda. 2013. Unification of regression-based methods for the analysis of
natural selection. Evolution **67**:2094-2100.
Opedal, Ø. H. 2018. The evolvability of animal-pollinated flowers: towards predicting adaptation to
novel pollinator communities. New Phytologist.
Parachnowitsch, A. L., and A. Kessler. 2010. Pollinators exert natural selection on flower size and
floral display in *Penstemon digitalis*. New Phytologist **188**:393-402.
Pausas, J. G., G. A. Alessio, B. Moreira, and G. Corcobado. 2012. Fires enhance flammability in *Ulex*
parviflorus. New Phytologist **193**:18-23.
Pausas, J. G., J. E. Keeley, and D. W. Schwilk. 2017. Flammability as an ecological and evolutionary
driver. Journal of Ecology **105**:289-297.
Pausas J.G. & Millán M.M. 2019. Greening and browning in a climate change hotspot: the
Mediterranean Basin. *BioScience* 96:143-151.
Pausas, J. G., and B. Moreira. 2012. Flammability as a biological concept. New Phytologist **194**:610-
613.
Perrier, C., B. Delahaie, and A. Charmantier. 2018. Heritability estimates from genomewide
relatedness matrices in wild populations: Application to a passerine, using a small sample
size. Molecular ecology resources **18**:838-853.
Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. De
Bakker, and M. J. Daly. 2007. PLINK: a tool set for whole-genome association and
population-based linkage analyses. The American Journal of Human Genetics **81**:559-575.
Reverté, S., J. Retana, J. M. Gómez, and J. Bosch. 2016. Pollinators show flower colour preferences
but flowers with similar colours do not attract similar pollinators. Annals of Botany **118**:249-
257.
Ritland, K. 1996. Marker-based method for inferences about quantitative inheritance in natural
populations. Evolution **50**:1062-1073.
Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of image
analysis. Nature methods **9**:671.
Schwaegerle, K. E., and D. A. Levin. 1990. Quantitative genetics of seed size variation in *Phlox*.
Evolutionary ecology **4**:143-148.
Stanton-Geddes, J., J. B. Yoder, R. Briskine, N. D. Young, and P. Tiffin. 2013. Estimating heritability
using genomic data. Methods in Ecology and Evolution **4**:1151-1158.
van Kleunen, M., and J. Burczyk. 2008. Selection on floral traits through male fertility in a natural
plant population. Evolutionary ecology **22**:39-54.
VanRaden, P. M. 2008. Efficient Methods to Compute Genomic Predictions. Journal of Dairy Science
91:4414-4423.

Vasconcelos, T. N. C., M. Chartier, G. Prenner, A. C. Martins, J. Schönenberger, A. Wingler, and E.
Lucas. 2019. Floral uniformity through evolutionary time in a species-rich tree lineage. New
Phytologist **221**:1597-1608.
Westerkamp, C. 1997. Keel blossoms: bee flowers with adaptations against bees. Flora **192**:125-132.
Wilson, A. J., D. Réale, M. N. Clements, M. M. Morrissey, E. Postma, C. A. Walling, L. E. B. Kruuk, and
D. H. Nussey. 2010. An ecologist's guide to the animal model. Journal of Animal Ecology
79:13-26.
Wimmer, V., T. Albrecht, H. Auinger, and C. Schoen. 2012. *synbreed*: a framework for the analysis of
genomic prediction data using R. Bioinformatics **28**:2086-2087.
Winn, A. A. 2004. Natural selection, evolvability and bias due to environmental covariance in the
field in an annual plant. Journal of Evolutionary Biology **17**:1073-1083.

Table 1 Directional and quadratic selection differentials and coefficients (± standard errors) for the
two floral traits studied.

Trait	Differential		Gradient	
	Directional, S	Quadratic, c	Directional, β	Quadratic, γ
Standard petal area	-0.003 ± 0.003 ns	**-0.002 ± 0.000 *****	-0.041 ± 0.039 ns	**-0.102 ± 0.029 *****
Flower weight	-0.018 ± 0.033 ns	**-0.070 ± 0.023 ****	-0.022 ± 0.047 ns	-0.038 ± 0.021 ns
Interaction				0.000 ± 0.002 ns

595 **Table 2** Estimates of heritability h^2 and evolvability e (with 95% credibility intervals, CI) for floral
596 traits in wild *Ulex parviflorus*. 'Naïve' heritability models did not include spatial or environmental
597 predictors.
598

	Naïve h^2 model		Final h^2 model		Evolvability	
	h^2	CI	h^2	CI	e	CI
Standard petal area	0.76	0.60 - 0.81	0.001	0.00 – 0.27	<0.001%	0.00 - 1.91
Flower weight	0.71	0.60 - 0.80	0.14	0.03 - 0.34	0.42%	0.11 - 1.21

Figure legends

Figure 1. Flowers of *Ulex parviflorus*. (a) Flowers previous to a visit with standard petal extended and

reproductive organs enclosed by the keel petals and calix. (b) Pressed standard petal. (c) Flower

after being "triggered" by a bee visit, showing all petals and exposed reproductive organs. (Photo

credits: (a) MC Castellanos, (c) J. Quiles).

Figure 2. Fruit set as a function of the two floral traits measured, (a) standard petal area and (b)

flower weight (a proxy for floral size). Lines are generalised additive model fits with shaded areas

showing 95% confidence intervals.

Figure 1

Fig. 2