Sex Solves Haldane's Dilemma.

Donal A. Hickey

Department of Biology, Concordia University

7141 Sherbrooke West,

Montreal, Quebec H4B 1R6

Canada

and

G. Brian Golding

Department of Biology, McMaster University

1280 Main St. West,

Hamilton, Ontario L8S 4K1

Canada

Corresponding author: Donal A. Hickey, Department of Biology,

Concordia University, 7141 Sherbrooke West,

Montreal, Quebec H4B 1R6, Canada.

E-mail: donal.hickey@concordia.ca

Abstract

The cumulative reproductive cost of multi-locus selection has been seen as a
potentially limiting factor on the rate of adaptive evolution. In this paper, we show that
Haldane's arguments for the accumulation of reproductive costs over multiple loci are
valid only for a clonally reproducing population of asexual genotypes. We show that a
sexually reproducing population avoids this accumulation of costs. Thus, sex removes a
perceived reproductive constraint on the rate of adaptive evolution. The significance of
our results is twofold. First, the results demonstrate that adaptation based on multiple
genes – such as selection acting on the standing genetic variation - does not entail a huge
reproductive cost as suggested by Haldane, provided of course that the population is
reproducing sexually. Secondly, this reduction in the cost of natural selection provides a
simple biological explanation for the advantage of sex. Specifically, Haldane's
calculations illustrate the evolutionary disadvantage of asexuality; sexual reproduction
frees the population from this disadvantage.

Keywords: cost of selection; recombination; standing genetic
variation;

46   evolution of sex.

Introduction

Adaptive evolution depends on reproductive excess because disfavored alleles must be culled in order for favored alleles to increase in frequency within the selected population. Such culling, in the absence of a sufficient reproductive excess, would result in a decrease in population size and, eventually, it could lead to the extinction of the population. Haldane (1957; 1960) referred to the culling of disfavored types as the "cost of natural selection". In addition to defining this cost, Haldane estimated how great this cost might be during the substitution of one allele for another at a single locus. He then argued that, given independent selection at different genetic loci, this cost would be approximately additive over loci (Haldane 1957; 1960). In other words, if the cost of allelic substitution at one locus equals x, then the cost for L loci would equal Lx. This would mean that evolutionary adaptations that involved allelic substitutions at many genetic loci could prove to be "too costly", given the available reproductive excess. Thus, simultaneous selection at many loci might lead to extinction of the population or, at best, it would constrain the rate of adaptive change. This problem has been referred to as Haldane's dilemma (VanValen 1963).

Haldane's calculations of the cost of natural selection had a major impact on the field of evolutionary biology. For example, Dodson (1962) wondered if the cost were too high to allow the adaptive divergence between humans and chimpanzees, while Kimura used the concept of selective cost to support his theory of neutral evolution (Kimura 1968; 1995; Kimura and Maruyama 1969; Jensen et al 2019). Other studies have pointed out various possible ways in which the cost of allele substitution at many loci might be reduced. But all of these solutions involve changing some of the starting assumptions. For example, Maynard Smith (1968) and Sved (1968) pointed out that the cost could be reduced if we assume some form of truncation selection rather than independent selection, while Grant and Flake (1974) showed that the cost could be reduced in a sub-divided population. These solutions are not very satisfying, however, since they involve modifying Haldane's original assumptions of independent positive selection in a single undivided population. Moreover, while these modifying assumptions can reduce the accumulation of costs over loci, they do not eliminate it.

In this study, we wished to revisit the problem using Haldane's original

assumptions about population structure and the form of the selection function. We point

out, however, that Haldane (1957) did not take sexual reproduction into account. In this

paper, we use numerical simulations to demonstrate that Haldane's argument about the

summation of the costs over multiple loci does not apply to a sexually reproducing

population – and this is true regardless of the selection scheme or the population

structure. In the Discussion below, we explain why Haldane's concept of cumulative

costs applies only to a quasi-infinite asexual population. A randomly outbreeding sexual

population, on the other hand, avoids the problem envisaged by Haldane.

Haldane's reasoning (Haldane 1957) for the cost of natural selection contains two

parts. First, there is the calculation of the reproductive cost of substituting one allele for

another at a single locus during the process of adaptive evolution. As an example,

Haldane cited the rapid spread of industrial melanism among populations of the peppered

moth, *Biston betularia*, in the coal-mining areas of northern England (Kettlewell 1956).

In order for the melanic types to become common, the non-melanic genotypes must be

removed from the population. It is this removal of the existing type that constitutes the

cost of natural selection. In general, the time to fixation of the new type is a function of

the reproductive capacity and the initial frequency (see Felsenstein 1971).

The second part of Haldane's argument is about extending the idea of the selective

cost of adaptive evolution to many loci. For example, he argues that industrialization

might have simultaneously selected for other traits in *B. betularia*, in addition to

melanism. Given that the cost is a function of the initial frequency of the favored

genotype, and based on the fact that the expected frequency of genotypes carrying

multiple favorable alleles (at different loci, given linkage equilibrium between loci) is

much lower than the frequency of genotypes carrying a single favorable allele, Haldane

reasoned that the cost of natural selection at multiple loci would be larger than at a single

locus. In fact, he estimated that there is an approximate summation of these costs over

many loci. Thus, it would take approximately twice as long for a genotype carrying two

favorable alleles to reach fixation as it would for a single mutant (Grant and Flake 1974).

His conclusion was that *B. betularia* might not have sufficient reproductive capacity to

allow adaptive evolution at ten loci simultaneously.

As we demonstrate in this study, it is this accumulation of costs over several loci
that is avoided in a sexually outbreeding population. Specifically, in a sexual population
we do not need to assume an initial population that is large enough to contain at least one
individual with the optimum genotype. And it is precisely this assumption of a very large
initial population that led Haldane to his conclusion that the costs of selection would
accumulate over loci.

116
117
118

Materials and Methods

In our simulations, we wished to recreate as closely as possible the situation
envisaged by Haldane. Therefore, we considered a single undivided population of
100,000 diploid individuals in which rare favorable alleles were already present at several
loci. As in Haldane's model, the initial population was in linkage equilibrium. Also
following Haldane's example, selection at different loci was independent, i.e., fitness
effects were multiplicative with no epistasis and no dominance (heterozygotes had a
fitness intermediate between the homozygotes). The number of offspring produced by
each individual was determined by sampling a Poisson distribution with a mean of 2.
Thus, the population conformed to Haldane's idea of 'modest reproductive capacity". All
parents were diploid and had the same average fertility; selection was implemented on
offspring viability based on the relative genotypic fitness values; i.e., the fittest
individuals had the highest probability of survival from zygote to adult.

In addition to replicating Haldane's model for an asexual population, we did a
parallel set of simulations for a diploid, outbreeding sexual population. For the sexual
populations, loci are evenly spaced along a linear chromosome. Recombination was
implemented as follows: a single recombination event occurred between the two
chromosomes in a diploid parent. This is equivalent to one chiasma per meiotic
chromosome pair. The position of this recombination event along the chromosome was
chosen at random; in the case where we consider 100 loci, this means that the probability
of recombination between adjacent loci was 1%. The resulting recombinant represents a
sexual gamete. These gametes were then paired at random to produce diploid zygotes for

the next generation. All sexual parents produced the same average number of diploid

zygotes, i.e., an average of two zygotes per parent. This means that there was no selection

at the haploid stage of the life cycle. Viability selection was implemented on the sexual

offspring, in the same way as for the asexual population.

We performed numerical simulations of populations containing 100,000 diploid

individuals. We did ten replicates of each simulation and the results are shown in the

Figures below. The initial frequency of the favored allele is 0.05 at each locus. The

fitness advantage of a favored allele is 0.02 and fitness interactions are multiplicative

between loci. Individuals produced an average of two offspring each and selection

occurred through differential viability that was dependent on the relative genotypic

fitness value.

We did a series of simulations in which we varied the number of loci under

selection. Specifically, we did simulations for two, four and one hundred selected loci;

this illustrates how the system behaves as we increase the number of loci under

independent selection. In each case, we compared the outcome of selection in asexual and

sexual populations. Following Haldane's example, we assumed that the favorable alleles

were already segregating at a low frequency in the initial population, and that the

population was in linkage equilibrium. The source code for the simulations is available

on GitHub (https://github.com/gbgolding/evolutionSex).

Results

The results described below illustrate two main points. First, we show why it was

necessary for Haldane (1957) to implicitly assume a progressively larger initial

population as the number of loci under selection increased. The reason was that

Haldane's model did not include recombination between the selected loci; therefore, the

initial population had to be large enough to contain at least one individual that contained

all of the favorable mutations in its genotype. Secondly, we show that there is no need to

increase the initial population size for multi-locus selection in a sexually outbreeding

population. This is because recombination will automatically produce the genotype with

the maximum number of favored alleles later on during the selection process (see

Discussion below).

First, we performed simulations of the situation where there is only one locus

undergoing selection. This allowed us to check the accuracy of our simulation method

against the theoretical predictions of classical population genetics and it also provided a

reference point for the subsequent simulations involving more than one locus. The results

are presented in Figure 1. The results of ten replicate simulations are shown in Figure 1a,

and the expected rise in allele frequency based on classical population genetics theory (Li

1955 page 258) is shown in Figure 1b. From the Figure, we see that the simulation results

match the theoretical prediction very well and that it takes approximately 400 generations

for the favored allele to rise from its initial frequency of 0.05 to near fixation.

Having established that the simulation provides the expected results for selection

at a single locus, we then performed simulations of selection acting at multiple loci in

both sexual and asexual populations. The results for two loci are shown in Figure 2,

Panels A and B. We see that the asexual and sexual populations yield different results

even when we are dealing with only two loci. This can be explained as follows. Given

that the initial frequencies of the favored alleles at the two loci are 0.05, then the expected

frequency of chromosomes carrying both favored alleles is one in four 400. But the

expected frequency of diploids carrying two such chromosomes is only one in 160,000;

therefore, we may not see even a single such double homozygote in the initial population.

In most replicates of the simulation, the expected single double homozygote is indeed

missing, leaving the fittest genotype with three rather than four favorable alleles (see

Figure 2, Panel A). In one "lucky" replicate, the initial population does contain a double

homozgote. Of course, if we had assumed a higher initial frequency for the favored

alleles – say 0.1 rather than 0.05 – then the initial population would have contained

several double homozygotes of the favored combination in all replicates. On the other

hand, the problem would have become even more acute if we had used lower initial allele

frequencies, say 0.01. In contrast to the asexual population, however, the sexual

population can generate diploid genotypes carrying four favorable alleles in all ten

replicates as the allele frequencies rise at the individual loci (see Figure 2, Panel B). In

other words, recombination compensates for the absence of the optimal genotype in the

initial population.

The results for selection acting simultaneously at four loci are shown in Figure 2,

Panels C and D. In this case, we begin to see an even greater difference between the

asexual and sexual populations. The reason for this difference is that the initial asexual

population lacks many high fitness genotypes. And the reason for this absence is that the

expected frequency of such individuals – given the initial allele frequencies of 0.05 and

linkage equilibrium – is exceedingly small. For example, the expected frequency of

diploid genotypes which are homozygous for the favorable allele at all four loci is less

than 10^{-10}. Consequently, it would require a population size of billions of diploid

individuals in order to see even a single individual of the optimal genotype within the

initial population. From Figure 2c we can see that the fittest genotypes in the asexual

population contained only four or five out of the eight possible favorable alleles – giving

average allele frequencies after selection of 0.5 and 0.625, respectively at the end of the

simulation. Of course, the sexual population is also lacking these high fitness genotypes

initially, but this is not a problem because random recombination can assemble such

genotypes as the individual allele frequencies rise in response to selection. For example,

once the allele frequencies have reached a frequency of 0.5 in an outbreeding sexual

population, we expect to see some individuals with the optimal genotype begin to appear

within the population of 100,000 diploid individuals. This explains why the average

allele frequency approaches a value of 1.0 in all replicates of the sexual population (see

Figure 2d).

Finally, the results for selection acting on 100 loci are shown in Figure 2, Panels E

and F. In this case, there is a striking difference between the asexual and sexual

populations. The initial population contains only low fitness genotypes where we expect

an average of only five favorable alleles per chromosome – given that there are 100 loci

and a frequency of 0.05 for the favorable allele at each locus (see Hickey and Golding

2018). The asexual population is again limited to selecting among such low-fitness

genotypes. In contrast to this, the sexual population can use recombination to gradually

assemble chromosomes with increasing numbers of favorable alleles per chromosome as

the frequencies of the favorable alleles rise in response to selection. At the end of the

selection period, the sexual population consists of individuals that are homozygous for

the favored allele at all one hundred loci. Table 1 shows a summary of the response to

selection in both the asexual and the sexual populations. As can be seen in Table 1, the

asexual population is constrained by the maximum fitness in the initial population; in

contrast to this recombination continues to produce increasing maximum fitnesses as the

selection proceeds. This allows the sexual population to advance beyond the highest

fitness value in the initial population. The complete numerical results for the simulations

of 100 loci are provided in Supplementary Tables S1 and S2.

These results show that, in a sexually outbreeding population, the frequency of

favorable alleles at many loci can respond simultaneously to independent selection at

rates that are very similar to the predicted rate at a single locus (compare Figure 2, Panel

F and Figure 1, Panel B). This implies that the rate of response to selection at any given

locus is not greatly impeded by selection at other loci. In other words, many loci can

respond simultaneously to independent selection – but only in a sexually reproducing

population. In summary, our results show that an asexual population cannot respond to

multi-locus selection as efficiently as a sexual population because it cannot generate

higher-fitness genotypic combinations as the allelic frequencies increase in response to

selection. Moreover, the relative disadvantage of the asexual population increases as the

number of loci under selection increases (compare Panels A, C and E in Figure 2).

Discussion

As stated in the introduction, Haldane (1957) argued that the costs of natural

selection accumulate over loci, i.e., the reproductive cost of allele substitution at two loci

would be twice the cost at a single locus. Consequently, the time required for substitution

should be twice as long. In general, if it takes t generations for a gene substitution at a

single locus, it should take Lt generations for substitutions at L loci. Applying this logic

to our model, given that it takes 400 generations for allelic substitution at one locus, it

should take 40,000 generations for substitutions at 100 loci. This prediction is in stark

contrast to our finding that it takes only 400 generations for allelic substitution at all 100

loci (see Figure 2, Panel F). How can we explain this major difference between Haldane's

prediction and our findings?

Clearly, Haldane's scenario does not explain our simulation results as shown in

Figure 2. The reason for the discrepancy is, we believe, that Haldane's argument assumed

that the optimal genotypic combination had to be already present in the initial population

and that genotypes could be considered as fixed entities that reproduce themselves from

one generation to the next. But these assumptions do not hold for a sexually outbreeding

population. First, the population is usually not large enough to contain even a single

individual with the vanishingly rare optimal genotypic combination; this fact has already

been pointed out by Ewens (1972) and Maynard Smith (1976). Secondly, specific

genotypes are broken down by recombination each generation and replaced by other

genotypic combinations (Hickey and Golding 2018). The process of recombination

involves the continual disassembly of existing genotypes and reassembly of new

genotypes from one generation to the next. Because of this process, however, those

genotypes which are expected to be vanishingly rare when the allele frequencies are low

will be automatically generated through recombination once the allele frequencies begin

to rise in response to selection (Muller 1964; Hickey and Golding 2018). Consequently,

there is no need for the massive culling that would be required in a very large initial

population that contained an extremely rare optimal genotype. Thus, in an outbreeding

sexual population, recombination solves the perceived problem of costs that are

cumulative over different genetic loci. In Haldane's scenario, it is necessary to "grow"

the optimal genotypic combination from an initial, vanishingly rare frequency to fixation.

In practice, however, this optimal combination is produced by recombination only near

the end of the process. This avoids the huge cumulative cost of natural selection.

Haldane's theory is mathematically very elegant; what happens in nature is

mathematically more muddled, but it is much more efficient. In nature, selection favors a

wide variety of suboptimal combinations which are then recombined. In this way,

recombination can harvest all favorable alleles from any genetic background. It is often

not until quite late in the selection process that the optimal genotypic combination first

appears. For example, even after the allele frequencies at each of 100 loci have risen to a

frequency of 0.9, the probability of randomly generating a chromosome with a favorable

allele at all 100 loci is still less than 0.0001. By the time the individual alleles have reached a frequency of 0.99, however, more than one third of all chromosomes in the population are expected to contain the optimal combination - i.e., a favorable allele at all 100 loci. In other words, rather than increasing slowly from an infinitesimally low initial frequency as suggested by Haldane (1957), the optimal combination is generated through recombination near the end of the selection process.

The strategy used by nature is somewhat analogous to the process of parallel computing. Since recombination decouples the allelic changes at a given locus from changes at other loci, it effectively decomposes multi-locus selection into several simultaneous instances of individual selection. Haldane (1957) envisaged that natural selection would have to do an exhaustive search through all possible genotypic combinations; instead, what happens in nature is a massively parallel heuristic search. The process of alternating selection and recombination enables the sexual population to trace a relatively narrow path through the myriad of possible genotypic combinations until, eventually, it arrives at the optimum genotypic combination. A comparison between natural selection and parallel computing has also been made by Wilf and Ewens (2010) although those authors were referring to the sequential fixation of favorable alleles during long-term evolution, rather than to the effects of recombination within a population.

The evolution of sex is a problem that has puzzled biologists for many decades (see Williams 1975; Maynard Smith 1978; Kondrashov 2018 for general reviews). Maynard Smith (1971) asked "what use is sex?" Our answer to that question is that recombination avoids the potentially increasing costs of natural selection as the number of genes under selection increases. This answer was already hinted at in the literature, ever since Fisher (1930) stated that the advantage of sexual reproduction is proportional to the number of genetic loci under selection. Muller (1932; 1964) developed Fisher's idea and concluded that recombination among many loci could allow a sexual population to evolve "hundreds to millions of times faster" than a comparable asexual population. This same idea of an increasing advantage for recombination as the number of loci under selection increases, has been echoed in later papers (e.g., Crow and Kimura 1965; Otto and Barton 2001; Colegrave 2002; Iles et al 2003; Park and Krug 2013; Edhan et al 2017). Park and Krug (2013) have shown that the advantage of recombination is

proportional to L, where L is the number of loci under selection. But the majority of these

previous studies deal with the effect of recombination on newly-arising favorable

mutations (e.g., Christiansen et al 1998) whereas the focus of this study was on the

standing genetic variation that is already present in the population. Also, none of these

previous studies offers a biological explanation of why the advantage of sex should

increase as we consider more loci under selection. Our explanation for this correlation is

that an asexual population would be subject to an increasing reproductive cost as the

number of loci increases – as shown by Haldane (1957). But since sexual reproduction

eliminates the accumulation of costs over loci (as shown in this study), this leads to an

increasing relative advantage of sex as the number of loci under selection increases.

Rather than saying that the advantage of sexual reproduction increases as the number of

loci under selection increases, it might be more accurate to say that there is an increasing

disadvantage to asexual reproduction as the number of loci under selection increases (see

Figure 2, Panels A, C and E).

It is interesting to note that Haldane (1957; 1960) did not see the relevance of

Muller's ideas about recombination to the problem of reproductive costs that accumulated

over loci. It is perhaps even more interesting that Muller (1964) also did not see the

connection between his own ideas and the problem posed by Haldane (1957; 1960). But

neither did their colleagues at the time see the connection (e.g. VanValen 1963). Indeed,

we believe that we are the first authors to have pointed out the direct link between the

ideas of Muller (1932; 1964) and those of Haldane (1957; 1960). And we have to admit

that it was only because of the counterintuitive nature of our own simulation results that

we were forced to re-examine Haldane's argument in the light of recombination. One

could say that the solution to Haldane's Dilemma has been "hiding in plain sight" for six

decades.

Our results support models of adaptation based on the standing genetic variation

(see Barrett and Schluter 2007; Orr 2005; Reznick 2016). As noted by Messer et al

(2016) rapid phenotypic evolution suggests that many genomic loci may contribute to

strongly selected traits. Molecular studies have confirmed that indeed several loci are

generally involved during rapid evolution (Burke et al 2010; Teotonio et al 2009).

Moreover, there is direct experimental evidence that recombination facilitates the rapid

response to multi-locus selection (McDonald et al 2016; Kosheleva and Desai 2018). The

relevance of our work is that it removes any concerns about the high cumulative costs of

multi-locus selection as argued by Haldane (1957), provided of course that there is

recombination occurring between the selected genes.

In a genetically polymorphic sexual population multi-locus genotypes are highly

ephemeral. This is because existing genotypes are merely a small random sample of a

much larger set of possible genotypes (Ewens 1972; Edhan 2017). In this study, we

assumed – for computational convenience – that there were only 100 biallelic loci. The

early, conservative estimates of genetic variation in natural populations concluded that

approximately twenty percent of the loci within the eukaryotic genome are polymorphic

(Lewontin and Hubby 1966). If we assume that there are 4,000 polymorphic loci – with

two alleles at each locus – there are $2^{4,000}$ possible genotypic combinations, a number

which is greater than $10^{1,000}$. This number is far greater than the total number of atoms in

the observable universe – which has been estimated to be approximately 10^{80} (see Hogan

2000). Darwin stated that "endless forms" have evolved as a result of natural selection.

We could say that this process has been facilitated through the production of effectively

endless genotypic combinations as a result of sexual reproduction and recombination.

In conclusion, our results show that outbreeding and recombination allow many

genes to respond rapidly to independent selection at the same time. In a sexual

population, selection on one gene does not slow the rate of adaptive response at another

genetic locus. This is because recombination allows allelic change at one locus to become

independent of the changes at other loci. Consequently, even those organisms with

limited population sizes and modest reproductive capacity can evolve efficiently. Haldane

(1957) predicted that the cost of multi-locus selection would pose a particularly difficult

challenge for such species. Perhaps this explains why sex tends to be most common

among species with limited population sizes and modest reproductive capacity.

Acknowledgements

This work was supported by a Discovery Grant from NSERC Canada to GBG

(grant no. 140221-10). We are grateful to Richard Lewontin for stimulating discussions

during the early development of this work. We would like to thank Chi Kuen Cheung for

his support and encouragement.

References

Barrett, R.D.H. and Schluter, D. 2007. Adaptation from standing genetic variation.

Trends Ecol. Evol. 23: 38-44.

Burke, M.K., Dunham, J.P., Shahrestani, P., Thornton, K.R., Rose, M.R. and Long, A.D.

2010. Genome-wide analysis of a long-term evolution experiment in Drosophila.

Nature 467: 587-590.

Christiansen, F.B., Otto, S.P., Bergman, A. and Feldman, M.W. 1998. Waiting with and

without recombination: the time to production of a double mutant. Theoret. Pop.

Biol. 53: 199-215.

Colegrave, N. 2002. Sex releases the speed limit on evolution. Nature 420: 664-666.

Crow, J.F. and Kimura, M. 1965. Evolution in sexual and asexual populations.

Amer. Natur. 99: 439-450.

Dodson, E.O. 1962. Note on the cost of natural selection. Amer. Natur. 96: 123-126.

Edhan, O., Hellman, Z. and Sherill-Rofe, D. 2017. Sex with no regrets: How sexual

Reproduction uses a no regret learning algorithm for evolutionary advantage.

J Theor Biol. 426, 67-81.

Ewens, W. J. 1972. The substitutional load in a finite population. Amer. Natur. 106:

273-282.

Felsenstein, J. 1971. On the biological significance of the cost of gene substitution. Amer.

Natur. 105: 1-12.

Fisher, R.A. 1930. *The Genetical Theory of Natural Selection*. Oxford: Clarendon.

Grant, V. and Flake, R.H. 1974. Population structure in relation to the cost of selection.

Proc. Nat. Acad. Sci USA 71: 1670 -1671.

Haldane, J.B.S. 1957. The cost of natural selection. J. Genet. 55: 511-524.

Haldane, J.B.S., 1960. More precise expressions for the cost of natural selection.

J. Genet. 57: 351-360.

Hickey, D.A. and Golding, G.B. 2018. The advantage of recombination when selection is

acting at many loci. J. Theoret. Biol. 442: 123-128.

Hogan, C.J. 2000. Why the universe is just so. Rev. Mod. Phys. 72: 1149-1161.

Iles, M.M., Walters, K. and Cannings, C. 2003. Recombination can evolve in large finite

populations given selection on sufficient loci. Genetics 165: 2249-2258.

Jensen, J.D., Payseur, B.A., Stephan, W., Aquadro, C.F., Lynch, M., Charlesworth, D.

and Charlesworth, B. 2019. The importance of the neutral theory in 1968 and 50

years on: a response to Kern and Hahn 2018. Evolution 73: 111-114.

Kettlewell, H.B.D. 1956. Further selection experiments on industrial melanism in the

Lepidoptera. Heredity 10: 287-303.

Kimura, M. 1968. Evolutionary rate at the molecular level. Nature 217: 624-626.

Kimura, M. and Maruyama, T. 1969. The substitutional load in a finite population.

Heredity 24: 101-114.

Kimura, M. 1995. Limitations of Darwinian selection in a finite population.

Proc. Nat. Acad. Sci. USA 92: 2343-2344.

Kondrashov, A.S. 2018. Through sex, nature is telling us something important.

Trends Genet. 34: 352-361.

Kosheleva, K. and Desai, M.M. 2018. Recombination alters the dynamics of adaptation

on standing variation in laboratory yeast populations. Mol. Biol. Evol. 35: 180-

201.

Lewontin, R.C. and Hubby, J.L. 1966. A molecular approach to the study of genic

Heterozygosity in natural populations. II. Amount of variation and degree of

heterozygosity in natural populations of Drosophila pseudoobscura . Genetics 54,

595-609.

Li, C.C. 1955. *Population Genetics*, p 258. (University of Chicago Press).

Maynard Smith, J. 1968. "Haldane's dilemma" and the rate of evolution. Nature 219: 1114-1116.

Maynard Smith, J. 1971. What use is sex? J. Theoret. Biol. 30: 319-335.

Maynard Smith, J. 1976. What determines the rate of evolution? Amer. Nat. 110: 331-338.

Maynard Smith, J. 1978. *The Evolution of Sex*. (Cambridge University Press).

McDonald, M. J., Rice, D. P. and Desai, M. M. 2016. Sex speeds adaptation by altering the dynamics of molecular evolution. Nature 531, 233-236.

Messer, P.W., Ellner, S.P. and Hairston, N.G. 2016. Can population genetics adapt to rapid evolution? Trends Genet. 32: 408-418.

Muller, H.J. 1932. Some genetic aspects of sex. Amer. Natur. 66: 118-138.

Muller, H.J. 1964. The relation of recombination to mutational advance. Mutat. Res. 1: 2 - 9.

Orr, H.A. 2005. The genetic theory of adaptation: a brief history. Nat. Rev. Genet. 6: 119-127.

Otto, S.P. and Barton, N. 2001. Selection for recombination in small populations. Evolution 55, 1921-1931.

Park, S.-C. and Krug, J. 2013. Rate of adaptation in sexuals and asexuals: a solvable model of the Fisher-Muller effect. Genetics 195: 941-955.

Reznick, D. 2016. Hard and soft selection revisited: how evolution by natural selection works in the real world. J. Hered. 107: 3-14.

Sved, J.A. 1968. Possible rates of gene substitution in evolution. Amer. Natur. 102: 283-293.

Teotónio, H., Chelo, I.M., Bradic, M., Rose, M.R. and Long, A.D. 2009. Experimental evolution reveals natural selection on standing genetic variation. Nat Genet. 41, 251-257.

VanValen, L. 1963. Haldane's dilemma, evolutionary rates, and heterosis. Amer. Natur. 97: 185-190.

Wilf, H.S. and Ewens, W.J. 2010. There's plenty of time for evolution. Proc. Nat. Acad.

479   Sci. USA 107: 22454-22456.

480   Williams, G. C. 1975. *Sex and evolution*. Princeton University Press, Princeton,

481   New Jersey.

485 Table 1. Response to multi-locus selection in asexual and sexual populations.

486 ____

	Asexual		Sexual	
	Av. fitness	Max. fitness	Av. fitness	Max. fitness
Generation				
0	1.22	1.64	1.22	1.70
10	1.27	1.64	1.27	1.71
20	1.33	1.64	1.33	1.88
30	1.40	1.64	1.41	2.00
40	1.47	1.64	1.50	2.43
50	1.53	1.64	1.62	2.43
60	1.57	1.64	1.77	2.85
70	1.59	1.64	1.96	3.26
80	1.61	1.64	2.21	3.68
90	1.62	1.64	2.52	4.39
100	1.63	1.64	2.91	4.94

____

The average fitness and maximum fitness are shown for each tenth generation, up
to generation 100; the complete data for all generations are shown in
Supplemental Tables T1 and T2. The population size was 100,000 diploid
individuals. The initial frequency of the favorable allele at each of 100 the loci
was 0.05. The selective advantage of each favorable allele was 0.02. The initial
population was in linkage equilibrium. Fitness effects were multiplicative between
loci.

Figure 1. Selection at a single locus, given an initial frequency of 0.05 and a selective

advantage of 0.02.

The population size was 100,000 diploid individuals. Panel A Shows simulation

results for 400 generations (10 replicates, results for individual replicates are shown).

Panel B shows the theoretical, predicted rise in frequency of a favored allele at a single

locus given an initial frequency of 0.05 and a selective advantage of 0.02.

Figure 2. Selection at multiple loci, given an initial frequency of 0.05 for the favorable

alleles and a selective advantage of 0.02.

The graph shows the rise in the average frequency of favorable alleles during the

course of selection. The reason that this frequency does not reach a value of 1.0 in the

asexual populations is because the highest fitness genotype in the initial populations did

not contain the maximum number of favorable alleles (see text). Panels A and B show the

results for asexual and sexual populations, respectively, when selection is acting at two

loci. Results for each of ten replicate simulations are shown separately. Panels C and D

show comparable results for four loci. Panels E and F show the results for 100 loci.