Brief Communication

Air mass source determines airborne microbial diversity at the ocean-

atmosphere interface of the Great Barrier Reef marine ecosystem

Stephen D.J. Archer[1,2#], Kevin C Lee[2#], Tancredi Caruso[3], Katie King-Miaow[2], Mike

Harvey[4], Danwei Huang[5], Benjamin J Wainwright[5], Stephen B Pointing[1,5,6]

[1] Yale-NUS College, National University of Singapore, Singapore 138527

[2] Institute for Applied Ecology New Zealand, Auckland University of Technology, Auckland

1142, New Zealand

[3] School of Biological Sciences and Institute for Global Food Security, Queen's University

Belfast, Belfast BT9 7BL, Northern Ireland, UK

[4] National Institute of Water and Atmospheric Research (NIWA), Wellington 6021, New

Zealand

[5] Department of Biological Sciences, National University of Singapore, Singapore 117558

[6] Institute of Nature and Environmental Technology, Kanazawa University, Kanazawa

920-1151, Japan

# These authors contributed equally.

The authors declare no competing interests.

Running title: Airborne microbial diversity above Great Barrier Reef

Keywords: aerobiology, atmospheric microbiology, bio-aerosol, coral symbionts, Great

Barrier Reef, marine microbiology, microbial biogeography.

1 **Abstract**

2 The atmosphere is the least understood biome on Earth despite its critical role as a

3 microbial transport medium. The influence of surface cover on composition of airborne

4 microbial communities above marine systems is unclear. Here we report evidence for a

5 dynamic microbial presence at the ocean-atmosphere interface of a major marine

6 ecosystem, the Great Barrier Reef, and identify that recent air mass trajectory over an

7 oceanic or continental surface associated with observed shifts in airborne bacterial and

8 fungal diversity. Relative abundance of shared taxa between air and coral microbiomes

9 varied between 2.2-8.8% and included those identified as part of the core coral

10 microbiome. We propose that this variable source of atmospheric inputs may in part

11 contribute to the diverse and transient nature of the coral microbiome.

12

13 **Main text**

14 Airborne microbial transport is central to dispersal outcomes [1] and several studies

15 have demonstrated diverse microbial biosignatures are recoverable from the

16 atmosphere. Microbial transport has been shown to occur across inter-continental

17 distances above terrestrial habitats [2–4]. Variation has been recorded seasonally [5, 6],

18 with underlying land use [7], and due to stochastic weather events such as dust storms

19 [8]. Above marine systems the abundance of microorganisms decreases exponentially

20 with distance from land [9], but relatively little is known about potential patterns in

21 biodiversity for airborne microorganisms above the oceans. Here we test the hypothesis

22 that persistent microbial inputs to the ocean-atmosphere interface of the Great Barrier

23 Reef ecosystem vary according to surface cover (i.e. land vs. ocean) during transit of the

24 air-mass.

The Great Barrier Reef is an ideal model system for research on bio-aerosols

because incoming air mass during the average residence time for microorganisms in air

[10] arises from two distinct sources: a terrestrial continental source in Australia

transported across east and northeast dust paths and an oceanic source in the Coral Sea

(Fig. 1a). Our study took advantage of a persistent flat-calm sea state during September-

October 2016 (www.marineweather.net.au). This minimised interference from

microorganisms that are aerosolised by marine spray in heavier sea states at the Great

Barrier Reef [11]. We recovered massive bulk-phase air samples at 25m above sea level

using a high-volume liquid impinger apparatus (Coriolis μ, Bertin Technologies, France)

[12] during a voyage of the *RV Investigator* to circumnavigate the reef (Supplementary

Information, Supplementary Methods). We used the National Oceanic and Atmospheric

Administration (NOAA) HYSPLIT-WEB model

(https://ready.arl.noaa.gov/HYSPLIT.php) to identify back trajectories of air mass

during the average residence time for microbial cells in air [10]. Back trajectories for air

mass could be delineated clearly into those with recent transit over either continental

Australia (continental path) or the Coral Sea (oceanic path) (Fig. 1a; Supplementary

Information, Fig. S1). The concurrent concentration of atmospheric radon gas measured

in real time was consistently higher from back trajectories originating over continental

Australia compared with those that originated from the ocean (Mann-Whitney U Test, P

= 0.003). This measurement further validated the binning of air mass into a continental

or oceanic origin.

Consistent with predictions that microbial biomass is extremely low and

unpredictable over the oceans, we achieved recoverable DNA from 27 of 53 bulk-phase

air samples (mean recovery 0.23 ng/m^3, STDEV 0.21 ng/m^3; Supplementary

Information, Table S1) with no significant difference in DNA yield between sample

groups (Mann-Whitney U Test, P = 0.37). High-throughput sequencing of the bacterial

16S rRNA gene and fungal Internal Transcribed Spacer (ITS) region were performed as

previously described [12] and phylogenetic analysis of amplicon sequence variants

(ASVs) were used to estimate diversity [13] (Supplementary Information,

Supplementary Methods). Sequencing of control samples revealed very low recovery of

putative contaminant microbial signatures. A total of only 17 out of 1403 bacterial and 5

out of 3775 fungal sequences were statistically classified as putative contaminants

(Supplementary Information, Supplementary Methods). Most samples were sequenced

to near-asymptote (Supplementary Information Fig. 1b, 1c).

Transit above oceanic or continental surfaces was significantly correlated with

bacterial and fungal community structures at the ocean-atmosphere interface above the

Great Barrier Reef (bacteria: PERMANOVA R^2 = 0.07233, pseudo F= 1.9494, P = 0.024;

manyglm LRT = 1650, P = 0.02; fungi PERMANOVA R^2 = 0.08413, pseudo F = 2.1125, P =

0.044, manyglm LRT = 6033, P = 0.034). Kendall's rank correlation tau identified a weak

but significant distance relationship (bacteria 0.19 P = <0.001 and fungi 0.10 P = 0.008)

(Fig. S5) which suggests air source but also distance can affect community similarities.

In terms of overall taxa richness, bacteria displayed similar richness but higher

phylogenetic diversity in continental versus ocean-derived samples (Fig. 1d;

Supplementary Information Fig. S2), and similarly the fungi displayed markedly greater

richness in continent-derived samples (Fig. 1e). Heatmaps of the most abundant ASVs

representing 95% of overall diversity illustrate the major taxonomic differences in

bacterial (Fig. 1f) and fungal (Fig. 1g) assemblages between ocean and continent-

derived air mass. Repeating this analysis with the 1,000 most abundant taxa from each

meta-library captured 99.98% of bacterial and 84.24% of fungal diversity in the study

and yielded similar results (Supplementary Information Fig. S3). Principal coordinate

analysis using Bray-Curtis dissimilarities of bacterial and fungal communities revealed

somewhat mixed trends in ordination plots (Supplementary Information Fig. S4),

however changes in relative abundance at the phylum level were striking between days

when air mass was sourced from continent or ocean (Fig. 2c, 2e).

Overall, the shared ASVs between samples from ocean and continental air

masses represented a large proportion of total bacterial (87.9% and 89.6%) and fungal

(64.9% and 69.1%) reads. However, the abundances for dominant genera was highly

variable between air masses. For example, the three most abundant bacterial genera

Burkholderia, *Ralstonia* and *Sphingomonas* were present at 28%, 14.4% and 5.5% in

oceanic air and 37%, 0.5% and 20% in continental air masses while the three most

abundant fungal genera (*Alternaria*, *Cladosporium* and *Mycosphaerella)* were present at

6.4%, 4.5% and 5.3% in oceanic air and 13.5%, 3.5% and 1.4% in continental air masses

respectively. A recent study characterizing bacteria aerosolized from a seawater

mesocosm concluded approximately 75% of taxa were alpha, beta and gamma

Proteobacteria [14], and notably identified *Bradyrhizobium* and *Ralstonia* as highly

abundant taxa thus establishing a potential marine source for some of the most

abundant airborne taxa in our study. While the core ASVs represented a high proportion

of reads, many less abundant ASVs were specific to either oceanic (538 bacteria, 1,335

fungi) or continental (395 bacteria, 1,810 fungi) air masses (Figs. 1f, 1g).

Shifts in relative abundance for bacteria mainly occurred within the phylum

Proteobacteria where the dominant genus recovered changed with air mass sampled

among *Acinetobacter*, *Alistipes*, *Bradyrhizobium* and *Ralstonia* (Fig. 2a, 2c). All have been

recovered as isolates or environmental rRNA gene sequences from both marine and

terrestrial sources, thus making any attempt at source tracking challenging. For the

Fungi terrestrial air sources supported higher diversity than marine sources and this

reflected the abundance of terrestrial fungal habitats. A striking observation was that 56% of recovered genera supported yeast-like taxa, and this provides support for a marine origin since oceanic waters are thought to support elevated abundance of yeasts over filamentous fungi [15]. Shifts in diversity were less pronounced overall for fungi (Fig. 2b, 2d), and were partially obscured by the high diversity relative to bacteria at the genus level. Major shifts in relative abundance occurred for *Aureobasidium, Cladosporium, Coprinopsis*, *Rhodosporidiobolus* and *Rhodotorula*, and all of these genera have known terrestrial and marine records, although it should be noted that terrestrial fungal spores have been recorded in many marine microbial diversity assessments but are unlikely to be active components of an ocean surface water microbiome.

We further interrogated the phylogenetic diversity of continental and oceanic air masses using Net Relatedness Index (NRI) analysis to estimate the level of phylogenetic structuring and putative recruitment from local (i.e. ocean or continental source only) and regional (all sources) pools [12]. The NRI analysis revealed that bacterial assemblages from both oceanic and continental origin displayed positive NRI values with effect sizes indicative of non-random assembly (Fig. 2d, 2f). Communities were thus phylogenetically highly structured, which is possibly due to environmental filtering of traits during transit over the different surface covers. The fungi from continental sources displayed a similar though more variable trend of phylogenetic structuring although in ocean-derived samples this clustering was relatively weak indicating they were more randomly assembled. Overall our analyses support the hypothesis that long-range transport of microbial taxa in air results in differential recruitment and selection during transit over oceanic or continental surfaces [12, 16, 17].

It has been proposed that airborne deposition of microorganisms may be a source of symbionts and pathogenic taxa to coral reefs [18, 19]. In order to establish the

potential recruitment of airborne microorganisms to coral reefs we compared our

sequence data to that obtained for coral microbiomes. We identified only one study with

directly comparable sequence data (Supplementary Information, Supplementary

Methods) recovered from *Porites lutea* coral microbiomes in the Andaman Sea and Gulf

of Thailand [20]. A total of 6.7% coral reads shared the same ASVs as our airborne

bacterial dataset. The air masses from oceanic sources had higher relative abundance of

ASVs in common with the coral dataset (8.8%) compared to air with recent continental

transit (2.2%) (Supplementary Information, Fig. S6a). Three of the six shared genera

(*Bradyrhizobium, Burkholderia* and *Sediminibacterium*) were identified as contributing

to the core coral microbiome for the Andaman Sea and Gulf of Thailand [20]. The class

Burkholdariales that supports *Burkholdaria* has been identified as a source of

ubiquitous coral endosymbionts globally [21]. Relaxing the stringency of this

comparison to the Genus level (i.e. ≥97% sequence matches) to account for possible

biogeographic variation and ecological variation resulted in higher relative abundance

of shared taxa between the datasets. Using this criteria 36% coral bacterial taxa were

shared with the atmospheric microbiome, and 77% of oceanic and 79% continental

derived atmospheric taxa matched those of the coral microbiome (Supplementary

Information, Fig. S6b).

We also compared our bacterial ASV sequences with a recent study of tropical

corals from the Singapore Straits, South China Sea [22] which shared significant

bacterial 16S rRNA sequence overlaps (Supplementary Information, Supplementary

Methods). We identified (with ≥97% sequence identities) 6.5% of coral taxa were

shared with the atmospheric microbiome, and 94.4% of ocean derived air and 87.8% of

continent derived air taxa were shared with the coral microbiome (Supplementary

Information, Fig. S6c). Finally, we broadly screened taxa from our sequence libraries at

taxon level (Genus) to known coral-associated taxa including putative symbionts and

pathogens identified using different rRNA loci or approaches (e.g. DGGE and Sanger

sequencing) (Supplementary Information, Supplementary Methods). Using this

approach we estimated the ocean derived air mass supported 8.4% bacterial genera

that may include coral associates whilst for continental sources this value was lower at

2.9% (Supplementary Information, Table S2). The most differentially abundant fungal

taxon from this comparison (*Ralstonia* sp.) showed greatest sequence similarity to a

previously identified coral endosymbiont (Supplementary Information, Table S3) thus

indicating any atmosphere-coral microbiome connectivity may be highly variable. For

our fungal library there were few coral-associated fungal sequences with which to make

comparisons and little phylogenetic information, but at a broad level we identified 8.7%

of ocean-derived taxa and 6.2% continent-derived taxa that have recorded coral

associates in the same Genus (Supplementary Information, Table S3).

Overall our study has provided a unique insight on the variability of airborne

microbial communities above the largest coral reef ecosystem on Earth and yielded

clues that atmospheric, oceanic and terrestrial biomes may be inter-connected via the

atmospheric microbiome. Our study indicates that the view of coral microbiomes as

harbouring few long term residents and instead comprising largely a "diverse transient

community that is responsive to surrounding environment"[23] may be explained in

part by the variability in airborne microbial diversity above reefs that may act as

recruitment reservoirs. Improved understanding of cross-biome microbial

biocomplexity and interaction will require coordinated and directly comparable

research effort from atmospheric, marine and terrestrial microbiologists.

Data availability

All sequence data generated by this study has been submitted to the EMBL European

Nucleotide Archive (ENL) under BioProject PRJEB31630 with sample accession

numbers ERS3215240 to ERS3215312.

Author contribution statement

S.D.J.A, K.C.L, T.C., M.H. and S.B.P. conceived the study; S.D.J.A. conducted ship-board

fieldwork; S.D.J.A and K.C.L. performed laboratory experiments; S.D.J.A., K.C.L., T.C., K.K-

M., B.J.W. and S.B.P. performed data analysis; S.D.J.A, M.H., D.H., B.J.W. and S.B.P.

analysed and interpreted the findings; S.B.P. wrote the manuscript.

Materials & Correspondence

Correspondence and requests for materials should be addressed to S.B.P. Email:

stephen.pointing@yale-nus.edu.sg.

Acknowledgements

Collaboration in this work was supported through ARC Discovery Project (grant

number DP150101649) and NIWA's Research Programme in Ocean-Climate

Interactions (grant number 2017/18 SCI). Financial support for laboratory research

was provided by the Yale-NUS College Start-up Fund. Author SDJA was supported by a

Postdoctoral Fellowship from Auckland University of Technology (AUT). We thank ARC

Discovery Project Principal Investigators Prof. Zoran Ristovski and Luke Cravigan

(Queensland University of Technology), the officers and crew of the RV Investigator

(IN2016_V05), and Tony Bromley and Sally Gray (NIWA) for their outstanding logistical

support and Timothy Lawrence (AUT) for his DNA sequencing support. Radon data was

provided by Scott Chambers and Alastair Williams (Australian Nuclear Science and

Technology Organisation).

Competing Interests

The authors declare no competing interests.

References

1. Hanson CA, Fuhrman JA, Horner-Devine MC, Martiny JBH. Beyond biogeographic patterns: processes shaping the microbial landscape. *Nat Rev Microbiol* 2012; **10**: 497–506.

2. Barberán A, Henley J, Fierer N, Casamayor EO. Structure, inter-annual recurrence, and global-scale connectivity of airborne microbial communities. *Sci Total Environ* 2014; **487**: 187–195.

3. Spracklen D V., Heald CL. The contribution of fungal spores and bacteria to regional and global aerosol number and ice nucleation immersion freezing rates. *Atmos Chem Phys* 2014; **14**: 9051–9059.

4. Caliz J, Triadó-Margarit X, Camarero L, Casamayor EO. A long-term survey unveils consistent seasonal trends in the airborne microbiome composition coupled to air masses circulation. *Proc Natl Acad Sci* 2019; **115**: 12229–12234.

5. Woo AC, Brar MS, Chan Y, Lau MCY, Leung FCC, Scott JA, et al. Temporal variation in airborne microbial populations and microbially-derived allergens in a tropical urban landscape. *Atmos Environ* 2013; **74**: 291–300.

6. Bowers RM, McCubbin IB, Hallar AG, Fierer N. Seasonal variability in airborne bacterial communities at a high-elevation site. *Atmos Environ* 2012; **50**: 41–49.

7. Bowers R, McLetchie S, Knight R, Fierer N. Spatial variability in airborne bacterial

communities across land-use types and their relationship to the bacterial communities of potential source environments. *ISME J* 2011; **5**: 601–612.

8. Favet J, Lapanje A, Giongo A, Kennedy S, Aung Y-Y, Cattaneo A, et al. Microbial hitchhikers on intercontinental dust: catching a lift in Chad. *ISME J* 2013; **7**: 850–67.

9. Mayol E, Arrieta JM, Jiménez MA, Martínez-Asensio A, Garcias-Bonet N, Dachs J, et al. Long-range transport of airborne microbes over the global tropical and subtropical ocean. *Nat Commun* 2017; **8**: 201.

10. Burrows SM, Butler T, Jöckel P, Tost H, Kerkweg A, Pöschl U, et al. Bacteria in the global atmosphere – Part 2: Modeling of emissions and transport between different ecosystems. *Atmos Chem Phys* 2009; **9**: 9281–9297.

11. Cropp R, Gabric A, van Tran D, Jones G, Swan H, Butler H. Coral reef aerosol emissions in response to irradiance stress in the Great Barrier Reef, Australia. *Ambio* 2018; **47**: 671–681.

12. Archer S, Lee K, Caruso T, Lee C, Maki T, Cowan D, et al. Microbial dispersal limitation to isolated soil microbial habitats of Antarctica. *Nat Microbiol* 2019; **4**: 925-932.

13. Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J* 2017; **11**: 2639–2643.

14. Michaud JM, Thompson LR, Kaul D, Espinoza JL, Richter RA, Xu ZZ, et al. Taxon-specific aerosolization of bacteria and viruses in an experimental ocean-atmosphere mesocosm. *Nat Commun* 2018; **9**: 2017.

15. Kutty SN, Philip R. Marine yeasts—a review. *Yeast* 2008; **25**: 465–483.

16. Mayol E, Jiménez MA, Herndl GJ, Duarte CM, Arrieta JM. Resolving the abundance

and air- sea fluxes of airborne microorganisms in the North Atlantic Ocean. *Front Microbiol* 2014; **5**: 557.

17. Šantl-Temkiv T, Gosewinkel U, Starnawski P, Lever M, Finster K. Aeolian dispersal of bacteria in southwest Greenland: Their sources, abundance, diversity and physiological states. *FEMS Microbiol Ecol* 2018; **94**: fiy031.

18. Weir-Brush JR, Garrison VH, Smith GW, Shinn EA. The Relationship Between Gorgonian Coral (Cnidaria: Gorgonacea) Diseases and African Dust Storms. *Aerobiologia* 2004; **20**: 119–126.

19. Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I. The role of microorganisms in coral health, disease and evolution. *Nat Rev Microbiol* 2007; **5**: 355–362.

20. Pootakham W, Mhuantong W, Yoocha T, Putchim L, Sonthirod C, Naktang C, et al. High resolution profiling of coral-associated bacterial communities using full-length 16S rRNA sequence data from PacBio SMRT sequencing system. *Sci Rep* 2017; **7**: 2774.

21. Ainsworth TD, Krause L, Bridge T, Torda G, Raina JB, Zakrzewski M, et al. The coral core microbiome identifies rare bacterial taxa as ubiquitous endosymbionts. *ISME J* 2015; **9**: 2261–2274.

22. Wainwright BJ, Afiq-Rosli L, Zahn GL, Huang D. Characterisation of coral-associated bacterial communities in an urbanised marine environment shows strong divergence over small geographic scales. *Coral Reefs* 2019; https://doi.org/10.1007/s00338-019-01837-1.

23. Leggat W, Ainsworth TD, Herrera C, Bongaerts P, Hernandez-Agreda A. Rethinking the Coral Microbiome: Simplicity Exists within a Diverse Microbial Biosphere. *MBio* 2018; **9**: 1–14.

Figure Legends

Figure 1. (a) HYSPLIT back trajectory analysis for modelled transit routes (3 d

residence time [12]) with colours representing air mass origin over continent (green)

or ocean (red) surface, scale bar = 500km; Rarefaction curves for samples by origin for

(b) bacteria (Continent n = 8, Ocean n = 19) and (c) fungi (Continent n = 8, Ocean n =

17). Chao1 diversity index by origin for (d) bacteria and (e) fungi. Individual data points

are shown as circles, boxplot whiskers represent 1.5 times the interquartile range from

the first-third quartiles; (f) Heatmap for bacterial amplicon sequence variants (ASVs)

that explain 95% of observed diversity by origin with control samples included

(Continent n = 8, Ocean n = 19, control n = 7); (g) Heatmap for fungal ASVs that explain

95% of observed diversity by origin with control samples included (Continent n = 8,

Ocean n = 17, control n = 5). All samples were sequenced to near-asymptote. More

detailed heatmaps that incorporate the 1,000 most abundant ASVs for bacteria and

fungi are shown in Supplementary information, Fig. S3.

Figure 2. Relative abundance and Genus level taxonomic assignment for ASV of all taxa

with >0.5% mean relative abundance for (a) bacteria and (b) fungi; Relative abundance

fold differences for taxa from ocean and continent derived air mass, for (c) bacteria and

(e) fungi; Net Relatedness Index (NRI) of phylogenetic structure for continental and

ocean-derived atmosphere taxa for (d) bacteria(Continent n = 8, Ocean n = 19, control n

= 7) and (f) fungal (Continent n = 8, Ocean n = 17, control n = 5) ASVs, error bars

indicate standard error of the mean for all samples. Bacterial analysis was based on a

phylogenetic tree derived using Maximum Likelihood analysis of 16S rRNA gene ASVs.

Fungal analysis utilised a Neighbour Joining tree that was compared with topology of

whole genome trees for fungi at high taxonomic ranks, in order to be conservative

regarding the phylogenetic information provided by ITS (Supplementary Information,

Supplementary Methods).