Air mass source determines airborne microbial diversity at the ocean-

atmosphere interface of the Great Barrier Reef marine ecosystem

Stephen D.J. Archer[1,2#], Kevin C Lee[2#], Katie King-Miaow[2], Tancredi Caruso[3], Mike

Harvey[4], Stephen Brian Pointing[1,5,6]

[1] *Yale-NUS College, National University of Singapore, Singapore 138527*

[2] *Institute for Applied Ecology New Zealand, Auckland University of Technology, Auckland*

1142, New Zealand

[3] *School of Biological Sciences and Institute for Global Food Security, Queen's University*

Belfast, Belfast BT9 7BL, Northern Ireland, UK

[4] *National Institute of Water and Atmospheric Research (NIWA), Wellington 6021, New*

Zealand

[5] *Department of Biological Sciences, National University of Singapore, Singapore 117558*

[6] *Institute of Nature and Environmental Technology, Kanazawa University, Kanazawa*

920-1151, Japan

# These authors contributed equally.

Running title: Airborne microbial diversity above Great Barrier Reef
18

Keywords: aerobiology, bio-aerosol, coral symbionts, Great Barrier Reef, marine

microbiology, microbial biogeography.

Abstract

The aerosphere is the least understood biome on Earth despite its critical role as a

microbial transport medium. The influence of surface cover on composition of airborne

microbial communities above marine systems is unclear. Here we report evidence for a

dynamic microbial presence at the ocean-atmosphere interface of a major marine

ecosystem, the Great Barrier Reef, and identify that an oceanic or continental trajectory

for aerosols correlates with observed shifts in bacterial and fungal diversity. A putative

airborne source for coral symbionts and pathogens was also demonstrated thus

highlighting biological connectivity between atmosphere and ocean.

Main text

Airborne microbial transport is central to dispersal outcomes [1] and several studies

have demonstrated diverse microbial biosignatures are recoverable from the

aerosphere. Microbial transport has been shown to occur across inter-continental

distances above terrestrial habitats [2–4]. Variation has been recorded seasonally [5, 6],

with underlying land use [7], and due to stochastic weather events such as dust storms

[8] . Above marine systems abundance of recoverable microorganisms decreases

exponentially with distance from land [9] and relatively little is known about potential

biogeography of airborne microorganisms above the oceans. Some studies have shown

that aerosolization of marine microorganisms to the aerosphere is an important

process, e.g. [10], but the reverse deposition of airborne taxa to the ocean is poorly

understood. This deposition is, however, a potential source of marine pathogens [11]

and symbionts which are crucial to the health of coral reefs [12]. Here we test the

hypothesis that persistent microbial inputs to the ocean-atmosphere interface of the

Great Barrier Reef ecosystem vary according to surface cover during transit of the air-

mass and that an atmospheric source of putative coral-associated microbial taxa occurs.

The Great Barrier Reef is an ideal model system for research on bio-aerosols

because incoming air mass during the average residence time for microorganisms in air

[13] arises from two distinct sources: a terrestrial continental source in Australia

transported across east and northeast dust paths and an oceanic source in the Coral Sea

(Fig. 1a). Our study took advantage of a persistent flat-calm sea state during September-

October 2016 that minimised interference from microorganisms that are aerosolised by

marine spray in heavier sea states at the Great Barrier Reef [14]. We recovered massive

bulk-phase air samples 25m above sea level using a high-volume liquid impinger

apparatus (Coriolis μ, Bertin Technologies, France) [15] during a voyage of the *RV*

Investigator to circumnavigate the reef and recovered approximately 3,000m^3 air

during daily from which environmental DNA was recovered (n = 53) [15]

(Supplementary Information, Supplementary Methods). We used the National Oceanic

and Atmospheric Administration (NOAA) HYSPLIT-WEB model

(https://ready.arl.noaa.gov/HYSPLIT.php) to identify back trajectories of air mass

during the average residence time for microbial cells in air [13] and this could be

delineated very clearly into those with recent transit over continental or oceanic

surfaces (Fig. 1a). The concurrent concentration of atmospheric radon gas measured in

real time was higher from back trajectories originating over continental Australia

compared with those that originated from the ocean (Mann-Whitney U Test, P = 0.003)

thus further verifying the binning of air mass into a continental or oceanic origin. High-

throughput sequencing of the bacterial 16S rRNA gene and fungal Internal Transcribed

Spacer region loci were performed as previously described [15] and phylogenetic

analysis of amplicon sequence variants (ASVs) were used to estimate diversity [16]

(Supplementary Information, Supplementary Methods). Sequencing of control samples

revealed very low recovery of putative contaminant microbial signatures, a total of only

17 out of 1403 bacterial and 5 out of 3775 fungal sequences were statistically classified

as putative contaminants (Supplementary Information, Supplementary Methods). With

few exceptions all samples were sequenced to asymptote thus allowing meaningful

ecological inference from these data (Supplementary Information Fig. 1a).

All air samples supported ultra-low biomass as inferred indirectly from DNA

yield (mean recovery 0.23 ng/m^3, SD 0.21 ng/m^3) with no significant difference

between sample groups (Mann-Whitney U Test, P = 0.37). Terrestrial or oceanic transit

significantly determined bacterial community structure at the ocean-atmosphere

interface above the great Barrier Reef (P = 0.001). In terms of overall taxa richness,

bacteria displayed similar richness in terrestrial and ocean-derived samples whereas

fungi were markedly more speciose in continent-derived samples (Supplementary

Information, Fig. S1b).

We interrogated the phylogenetic diversity of terrestrial and marine-sourced air

by generating heatmaps of distribution for the 100 most abundant taxa. This analysis

captured 98.4% bacterial and 66.9% fungal overall diversity in the libraries. Clear

taxonomic differences in bacterial (Fig. 1b) and fungal (Fig. 1c) assemblages between

ocean and continent-derived air mass occurred. The robust nature of these data was

further highlighted by the lack of recoverable taxa in experimental controls

(Supplementary Information, Fig. S2) and this is particularly important for

interrogation of low biomass environments such as the aerosphere. Overall the number

of shared bacterial taxa was high amongst samples derived from ocean (87.9%) and

terrestrial (89.6%) sources. Fungi displayed fewer shared taxa between ocean (64.9%)

and continental (69.1%) sources. Nonetheless a high number of taxa specific to either

ocean (538 bacteria, 1,335 fungi) or continental (395 bacteria, 1,810 fungi) aerosols

were recorded (Figs. 1b, 1c). We screened bacterial taxa from our sequence libraries at

genus level to known coral-associated taxa including symbionts and pathogens

(Supplementary Information, Supplementary Methods) [17]. The ocean derived air

mass supported 8.4% putative coral associates whilst for continental sources this value

was lower at 2.9% (Supplementary Information, Table S1). There were few fungal

sequences with which to make comparisons but at the genus level there were 8.7% of

ocean samples and 6.2% continental taxa that have been recorded as coral associates

(Supplementary Information, Table S2). The most differentially abundant ASV from our

study (*Ralstonia* sp.) showed high similarity to a previously identified coral

endosymbiont (Supplementary Information, Table S3).

Shifts in relative abundance of bacteria mainly occurred within the Phylum

Proteobacteria and followed a temporal pattern during the voyage where the dominant

genus *Alistipes* was gradually replaced by *Bradyrhizobium* and *Ralstonia*, and then in

turn by *Acinetobacter* (Fig. 2a, Supplementary Information, Fig. S3). All have been

recovered as isolates or environmental rRNA gene sequences from both marine and

terrestrial sources, thus making any attempt at source tracking challenging. For the

Fungi terrestrial air sources supported higher diversity than marine sources and this

reflected the large number of taxa associated with terrestrial habitats. A striking

observation was that 56% of recovered genera supported yeast-like taxa, and this

provides support for a marine origin of aerosols since oceanic waters are thought to

support elevated abundance of yeasts over filamentous fungi [18]. Shifts in diversity

were less pronounced overall for the Fungi (Fig. 2b, Supplementary Information, Fig.

S4), and were partially obscured by the high diversity relative to Bacteria at the genus

level. Major shifts in relative abundance occurred for *Aureobasidium, Cladosporium,*

Coprinopsis, *Rhodosporidiobolus* and *Rhodotorula*, and all of these genera have known

terrestrial and marine records, although it should be noted that terrestrial fungal spores

have been recorded in many marine microbial diversity assessments but are unlikely to

be active components of an ocean surface water microbiome.

We further interrogated the phylogenetic diversity of terrestrial and marine-

sourced air using Net Relatedness Index (NRI) analysis to estimate the level of

phylogenetic structuring and putative recruitment from local and regional pools [15].

The NRI analysis revealed that bacterial assemblages from both continental and marine

origin displayed positive NRI values with effect sizes indicative of non-random

assembly. Communities were thus phylogenetically highly structured, which is possibly

due to environmental filtering of traits during transit over the different surface covers

(Supplementary Information Fig. S5). The fungi from continental sources displayed a

similar though more variable trend of phylogenetic structuring although in ocean-

derived samples this clustering was relatively weak indicating they were more

randomly assembled (Supplementary Information Fig. S5). When considered in tandem

with our low diversity estimates for fungi in ocean derived samples these findings

suggest that fungi are severely depleted in marine aerosols. Community phylogenetics

and all our analyses support the hypothesis that long-range transport of microbial taxa

in air results in selection during transit over different surface covers [15, 19, 20].

Overall our findings demonstrate that the aerosphere provides connectivity

between the Great Barrier Reef marine ecosystem and exogenous microbial input from

terrestrial and marine sources. This may be important to coral symbioses and also to

pathogen input via persistent airborne deposition. Our study provided a unique insight

on variability of airborne microbial inputs to the largest coral reef ecosystem on Earth.

Data availability

All sequence data generated by this study has been submitted to the EMBL European

Nucleotide Archive (ENL) under BioProject PRJEB31630 with sample accession

numbers ERS3215240 to ERS3215312.

Author contribution statement

S.D.J.A, K.C.L, T.C., M.H. and S.B.P. conceived the study; S.D.J.A. conducted ship-board

fieldwork; S.D.J.A and K.C.L. performed laboratory experiments; S.D.J.A., K.C.L., T.C., K.K-

M. and S.B.P. performed data analysis; S.D.J.A, K.C.L, T.C., K.K-M., M.H. and S.B.P. analysed

and interpreted the findings; S.B.P. wrote the manuscript.

Materials & Correspondence

Correspondence and requests for materials should be addressed to S.B.P. Email:

stephen.pointing@yale-nus.edu.sg.

Acknowledgements

Collaboration in this work was supported through ARC Discovery Project (grant

number DP150101649) and NIWA's Research Programme in Ocean-Climate

Interactions (grant number 2017/18 SCI). Financial support for laboratory research

was provided by the Yale-NUS College Start-up Fund. Author SDJA was supported by a

Postdoctoral Fellowship from Auckland University of Technology (AUT). We thank ARC

Discovery Project Principal Investigators Prof. Zoran Ristovski and Luke Cravigan

(Qeensland University of Technology), the officers and crew of the RV Investigator

(IN2016_V05), and Tony Bromley and Sally Gray (NIWA) for their outstanding logistical

support and Timothy Lawrence (AUT) for his DNA sequencing support. Radon data was

provided by Scott Chambers and Alastair Williams (Australian Nuclear Science and

Technology Organisation).

Conflict of Interest

The authors declare no competing interests.

References

1.   Hanson CA, Fuhrman JA, Horner-Devine MC, Martiny JBH. Beyond biogeographic

patterns: processes shaping the microbial landscape. *Nat Rev Microbiol* 2012; **10**:

497–506.

2.   Barberán A, Henley J, Fierer N, Casamayor EO. Structure, inter-annual recurrence,

and global-scale connectivity of airborne microbial communities. *Sci Total*

Environ 2014; **487**: 187–195.

3.   Spracklen D V., Heald CL. The contribution of fungal spores and bacteria to

regional and global aerosol number and ice nucleation immersion freezing rates.

Atmos Chem Phys 2014; **14**: 9051–9059.

4.   Caliz J, Triadó-Margarit X, Camarero L, Casamayor EO. A long-term survey unveils

consistent seasonal trends in the airborne microbiome composition coupled to air

masses circulation. *Proc Natl Acad Sci* 2019; **115**: 12229–12234.

5.   Woo AC, Brar MS, Chan Y, Lau MCY, Leung FCC, Scott JA, et al. Temporal variation

in airborne microbial populations and microbially-derived allergens in a tropical

urban landscape. *Atmos Environ* 2013; **74**: 291–300.

6.   Bowers RM, McCubbin IB, Hallar AG, Fierer N. Seasonal variability in airborne

bacterial communities at a high-elevation site. *Atmos Environ* 2012; **50**: 41–49.

7.   Bowers R, McLetchie S, Knight R, Fierer N. Spatial variability in airborne bacterial

communities across land-use types and their relationship to the bacterial communities of potential source environments. *ISME J* 2011; **5**: 601–612.

8. Favet J, Lapanje A, Giongo A, Kennedy S, Aung Y-Y, Cattaneo A, et al. Microbial hitchhikers on intercontinental dust: catching a lift in Chad. *ISME J* 2013; **7**: 850–67.

9. Mayol E, Arrieta JM, Jiménez MA, Martínez-Asensio A, Garcias-Bonet N, Dachs J, et al. Long-range transport of airborne microbes over the global tropical and subtropical ocean. *Nat Commun* 2017; **8**: 201.

10. Aller JY, Kuznetsova MR, Jahns CJ, Kemp PF. The sea surface microlayer as a source of viral and bacterial enrichment in marine aerosols. *J Aerosol Sci* 2005; **36**: 801–812.

11. Weir-Brush JR, Garrison VH, Smith GW, Shinn EA. The Relationship Between Gorgonian Coral (Cnidaria: Gorgonacea) Diseases and African Dust Storms. *Aerobiologia (Bologna)* 2004; **20**: 119–126.

12. Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I. The role of microorganisms in coral health, disease and evolution. *Nat Rev Microbiol* 2007; **5**: 355–362.

13. Burrows SM, Butler T, Jöckel P, Tost H, Kerkweg A, Pöschl U, et al. Bacteria in the global atmosphere – Part 2: Modeling of emissions and transport between different ecosystems. *Atmos Chem Phys* 2009; **9**: 9281–9297.

14. Cropp R, Gabric A, van Tran D, Jones G, Swan H, Butler H. Coral reef aerosol emissions in response to irradiance stress in the Great Barrier Reef, Australia. *Ambio* 2018; **47**: 671–681.

15. Archer S, Lee K, Caruso T, Lee C, Maki T, Cowan D, et al. Microbial dispersal limitation to isolated soil microbial habitats of Antarctica. *Nat Microbiol* 2019; **4**:

doi: 10.1038/s41564-019-0370-4.

16.   Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace

operational taxonomic units in marker-gene data analysis. *ISME J* 2017; **11**:

2639–2643.

17.   Harvell D, Jordán-Dahlgren E, Merkel S, Rosenberg E, Raymundo L, Smith G, et al.

Coral disease, environmental drivers, and the balance between coral and

microbial associates. *Oceanography* 2007; **20**: 172–195.

18.   Kutty SN, Philip R. Marine yeasts—a review. *Yeast* 2008; **25**: 465–483.

19.   Mayol E, Jiménez MA, Herndl GJ, Duarte CM, Arrieta JM. Resolving the abundance

and air- sea fluxes of airborne microorganisms in the North Atlantic Ocean. *Front*

Microbiol 2014; **5**: 557.

20.   Šantl-Temkiv T, Gosewinkel U, Starnawski P, Lever M, Finster K. Aeolian dispersal

of bacteria in southwest Greenland: Their sources, abundance, diversity and

physiological states. *FEMS Microbiol Ecol* 2018; **94**: fiy031.

Figure Legends

Figure 1. (a) HYSPLIT back trajectory analysis for modelled transit routes (3 d

residence time [12]) with colours representing air mass origin over continent

(green) or ocean (red) surface; (b) distribution and relative abundance for the

1,000 most abundant bacterial amplicon sequence variants (ASVs), Continent n =

8, Ocean n = 19; (c) distribution and relative abundance for the 1,000 most

abundant fungal ASVs, Continent n = 8, Ocean n = 17.

Figure 2. (a) Distribution and relative abundance of the 100 most abundant bacterial

amplicon sequence variants (ASVs) with genus identified, Continent n = 8, Ocean

n = 19; (b) Distribution and relative abundance of the 100 most abundant fungal

ASVs with genus identified, Continent n = 8, Ocean n = 17.

a.

b.

Continent | Ocean

Relative Abundance

1e-01
1e-02
1e-03
1e-04
1e-05

Amplicon Sequence Variants

c.

Continent | Ocean

Relative Abundance

1e-01
1e-02
1e-03
1e-04
1e-05

Amplicon Sequence Variants