A significant role for the circadian clock in the long-term water use efficiency of

Arabidopsis

Noriane M. L. Simon, Nicholas E. Comben, Alistair M. Hetherington, Antony N. Dodd*

School of Biological Sciences, University of Bristol, Life Sciences Building, Bristol BS8 1TQ,

U.K.

* Corresponding author, antony.dodd@bristol.ac.uk

Word count (body text including methods): 5374

Short title: Circadian regulation and water use efficiency

Author contributions: N.M.L.S. and A.N.D. conceived experiments, N.M.L.S. and N.E.C.

performed experiments, N.M.L.S., A.M.H. and A.N.D. analysed and interpreted the data,

N.M.L.S., A.M.H. and A.N.D. wrote the paper. A.N.D. agrees to serve as the author

responsible for contact and ensures communication.

One-sentence summary: The circadian clock in Arabidopsis makes an important

contribution to long-term water use efficiency.

Abstract

In plants, water use efficiency is a complex trait derived from numerous physiological and

developmental characteristics. Here, we investigated the involvement of circadian regulation

in long-term water use efficiency. Circadian rhythms are generated by the circadian

oscillator, which provides a cellular measure of the time of day. In plants, the circadian

oscillator contributes to the regulation of many aspects of physiology, including stomatal

opening, the rate of photosynthesis, carbohydrate metabolism and developmental

processes. We investigated in Arabidopsis the impact upon whole plant, long-term water use

efficiency of the misregulation of genes encoding a large number of components of the

circadian oscillator, identifying a major role for the circadian oscillator in plant water use.

This appears to be due to contributions of the circadian clock to the control of transpiration

and biomass accumulation. We also identified that the circadian oscillator specifically within

guard cells contributes to both long-term water use efficiency and dehydration tolerance. Our

experiments indicate that knowledge of circadian regulation will be important for developing

future crops that use less water.

Introduction

World population growth is increasing the demand for fresh water for agriculture, with

climate change predicted to exacerbate this competition for water resources (Ruggiero et al.,

2017). One strategy to sustainably increase agricultural production involves the

improvement of crop water use (Condon et al., 2004; Xoconostle-Cazares et al., 2010; Hu

and Xiong, 2014; Ruggiero et al., 2017). Since up to 97% of water taken up from the soil by

plants is lost through stomatal transpiration (Yoo et al., 2009; Na and Metzger, 2014), the

manipulation of transpiration represents an excellent candidate for designing crops with

increased water use efficiency.

Plant water loss can be manipulated through changes in the regulation of stomatal opening

and by altering stomatal density and patterning (Pei et al., 1998; Hugouvieux et al., 2001;

Schroeder et al., 2001; Hetherington and Woodward, 2003; Yoo et al., 2010; Lawson and

Blatt, 2014; Franks et al., 2015; Caine et al., 2019). In addition to stomatal responses to

environmental cues such as light, temperature and phytohormones, there are circadian

rhythms of stomatal opening (Gorton et al., 1989; Hennessey and Field, 1991). Circadian

rhythms are self-sustaining biological cycles with a period of about 24 h. These rhythms are

thought to adapt plants to daily cycles of light and dark, by anticipating daily changes in the

environment and co-ordinating cellular processes. In higher plants, circadian rhythms are

generated by several interlocked transcription-translation feedback loops known as the

circadian oscillator (Hsu and Harmer, 2014). The phase of the circadian oscillator is adjusted

continuously to match the phase of the environment through the process of entrainment, in

response to light, temperature and metabolic cues (Somers et al., 1998; Millar, 2004;

Salomé and McClung, 2005; Haydon et al., 2013). Additionally, the circadian oscillator

communicates an estimate of the time of day to circadian-regulated features of the cell,

initially through transcriptional regulation (Harmer et al., 2000). The known circadian

oscillator controls circadian rhythms of stomatal opening because mutations that alter the

circadian period or cause circadian arrhythmia lead to equivalent alterations in the circadian

rhythm of stomatal opening (Somers et al., 1998; Dodd et al., 2004; Dodd et al., 2005). The

circadian oscillator is also involved in the responses of guard cells to environmental cues

such as drought and low temperature (Dodd et al., 2006; Legnaioli et al., 2009).

Circadian rhythms are often studied under conditions of constant light. However, the

circadian oscillator is also important for the regulation of stomatal opening under cycles of

light and dark. For example, overexpression of the circadian oscillator component CCA1

(CCA1-ox) alters the daily regulation of stomatal opening such that stomatal conductance

increases steadily throughout the photoperiod (Dodd et al., 2005). In comparison, in wild

type plants stomatal conductance remains relatively uniform during the photoperiod and is

substantially lower than CCA1-ox (Dodd et al., 2005). This suggests that misregulation of the

circadian oscillator might alter plant water use under cycles of light and dark.

Overexpression of CCA1 specifically within guard cells, using a guard cell specific promoter,

alters flowering time and drought response phenotypes under cycles of light and dark

(Hassidim et al., 2017). Like constitutive CCA1 overexpression (Dodd et al., 2005), CCA1

overexpression specifically within guard cells generally causes greater stomatal opening

during the light period (Hassidim et al., 2017). Therefore, the circadian oscillator within guard

cells is important for the daily regulation of stomatal opening (Hassidim et al., 2017).

Modelling suggests that under light/dark cycles, the circadian oscillator contributes at the

canopy scale to daily rhythms in stomatal aperture and carbon assimilation in bean and

cotton (Resco de Dios et al., 2016).

The contribution of the circadian oscillator to both stomatal opening and growth and biomass

accumulation (Dodd et al., 2005; Graf et al., 2010) suggests that the circadian oscillator

might make an important contribution to water use efficiency (WUE). WUE is the ratio of

carbon dioxide incorporated through photosynthesis into biomass to the amount of water lost

through transpiration. At the single leaf level, instantaneous, intrinsic WUE is often

measured with gas exchange techniques and expressed as net CO_2 assimilation per unit of

water transpired (Vialet-Chabrand et al., 2016; Ruggiero et al., 2017; Ferguson et al., 2018).

However, such measurements do not provide an accurate representation of WUE over the

plant lifetime, which is influenced by features such as leaf position, dark respiration, and time

of day changes in instantaneous WUE (Condon et al., 2004; Tomás et al., 2014; Medrano et

al., 2015; Ferguson et al., 2018). It is important to note that WUE is not a drought resistance

trait (Blum, 2009).

Given that the circadian oscillator affects stomatal opening and biomass accumulation

(Gorton et al., 1989; Hennessey and Field, 1991; Dodd et al., 2005; Edwards and Weinig,

2010; Graf et al., 2010; Edwards et al., 2012), we hypothesized that specific components of

the circadian oscillator might make an important contribution to long-term WUE of plants.

Therefore, we investigated the impact of the misregulation of parts of the circadian oscillator

upon the long-term WUE of Arabidopsis. We identified that the circadian oscillator has

profound effects upon the long-term WUE of plants. Importantly, some alterations in

oscillator function increase long-term WUE, suggesting potential targets for future

improvements of crop WUE.

Results

Circadian oscillator components contribute to water use efficiency

Each background accession had a distinct WUE (C24: 3.01 ± 0.07 mg ml^{-1}; Col-0: 2.22 ±

0.02 mg ml^{-1}; L. *er.*: 1.60 ± 0.04 mg ml^{-1}; Ws: 1.91 ± 0.06 mg ml^{-1}) (Fig. S1). These

differences between backgrounds are consistent with previous studies of WUE, stomatal

function and stomatal density in Arabidopsis (Nienhuis et al., 1994; Woodward et al., 2002;

Dodd et al., 2004; Masle et al., 2005; Karaba et al., 2007; Ruggiero et al., 2017; Ferguson et

al., 2018).

We identified that correct regulation of the circadian oscillator makes a substantial

contribution to WUE. 33 single mutants or overexpressors of genes associated with

circadian regulation, representing 22 circadian oscillator-associated components, were

screened for WUE (Fig. 1). Nearly half of the mutants or overexpressors examined had a

significantly different WUE from the wild type ($p < 0.05$; 16 of 33 genotypes). This

corresponded to mutants or overexpressors representing half of the circadian oscillator

components covered by our study (11 of 22 genes) (Fig. 1). The *cca1*-11, *elf3*-1, *prr5*-3,

prr9-1, *tps1*-11, *tps1*-12, and *ztl*-1 mutants, as well as the TOC1 and KIN10 (line 6.5)

overexpressors, had significantly lower WUE than the wild type (Fig. 1). The *gi*-2, *gi*-11,

grp7-1, *prr7*-11 and *tej*-1 mutants had significantly greater WUE than the wild type (Fig. 1).

This suggests that misregulating the expression of circadian clock components CCA1, ELF3,

GI, GRP7, PRR5, PRR7, PRR9, TEJ, TOC1 and ZTL changes whole plant long-term WUE

(Fig. 1). *tic*-2 had significantly greater WUE than the wild type, whereas *tic*-1 had

significantly lower WUE than the wild type, so the effect of *TIC* mutation upon WUE is

unclear (Fig. 1). We also included the *che*-2 mutant in our initial analysis, but inconsistency

of its WUE phenotype between experimental repeats led us to exclude the data. WUE was

also altered by changing the expression of the energy signalling components TPS1 and

KIN10 that participate in inputs to the circadian oscillator (Shin et al., 2017; Frank et al.,

2018) (Fig. 1). Therefore, correct expression of these circadian clock-associated genes

contributes to long-term WUE of Arabidopsis.

We were interested to determine whether the WUE alterations caused by misregulation of

circadian oscillator gene expression arose from changes in either biomass accumulation or

transpiration. There was no clear evidence that a change in one of these parameters alone

underlies the altered WUE phenotypes (Fig. 2A). This suggests that the altered WUE of lines

with misregulated circadian clock genes is due to the net effect of altered biomass

accumulation and altered transpiration in these genotypes (Fig. 2A).

We hypothesised that variations in WUE might be explained by specific circadian

phenotypes in the mutants and overexpressors that we tested. For example, mutations in

clock genes expressed with a particular set of phases might have a pronounced effect on

WUE. Likewise, the nature of the circadian period change or flowering time change resulting

from misexpression of each oscillator component might be associated with certain changes

in WUE. To test this, we related the data from our WUE screen to the circadian phase of

expression of each mutated or overexpressed gene. We also compared the direction of

change of WUE to the period and flowering time phenotypes that arise from each mutant or

overexpressor (Fowler et al., 1999; Schultz et al., 2001; Doyle et al., 2002; Nakamichi et al.,

2002; Yanovsky and Kay, 2002; Imaizumi et al., 2003; Más et al., 2003; Murakami et al.,

2004; Farré et al., 2005; Hazen et al., 2005; Baena-González et al., 2007; Streitner et al.,

2008; Wang et al., 2008; Baudry et al., 2010; Nakamichi et al., 2010; Rawat et al., 2011;

Wahl et al., 2013; Hsu and Harmer, 2014). We note that the phenotypes reported by these

studies were often identified under constant conditions, whereas our experiments occurred

under light/dark cycles.

There was no obvious relationship between the circadian phenotypes that are caused by

each mutant or overexpressor investigated and the WUE of each of these lines (Fig. 2B, C,

D). For example, mutating morning-phased circadian oscillator components can either

decrease or increase WUE (Fig. 2B). Mutants that cause long circadian periods and short

circadian periods can both increase and decrease WUE (Fig. 2C). Furthermore, mutants and

overexpressors that cause both early and delayed flowering can each increase and

decrease WUE (Fig. 2D).

Circadian regulation of water use efficiency combines multiple traits

Mutation or overexpression of components of the circadian oscillator can cause changes in

the development of Arabidopsis, such as alterations in rosette size, leaf shape and petiole

length (Fig. 3A) (Zagotta et al., 1992; Schaffer et al., 1998; Wang and Tobin, 1998; Dodd et

al., 2005; Ruts et al., 2012; Rubin et al., 2018). These changes are likely to have

implications for gas exchange because, for example, spatially separated leaves are

predicted to transpire more water (Bridge et al., 2013). We investigated whether the changes

in WUE that were identified by our screen might arise from differences in rosette architecture

between the circadian clock-associated mutants and overexpressors and the wild types.

There was a weak positive correlation between rosette leaf surface area and WUE (r =

0.400; r^2 = 0.160; p < 0.001) (Fig. 3B). Therefore, approximately 16% of variability in WUE

can be explained by the variations in rosette leaf surface area that arise from misregulation

of the circadian oscillator.

In comparison, rosette leaf surface area was strongly correlated with each of the individual

parameters of water used and dry biomass accumulated. The variation in rosette surface

area accounted for 83% of the variability in water transpired across the genotypes (Fig. 3C).

Furthermore, the variation in rosette surface area accounted for 73% of the variability in

biomass accumulation across the genotypes (Fig. 3D), which is unsurprising given that

larger leaves are likely to contain more biomass. This demonstrates that one way that

circadian regulation affects WUE is through the influence of the circadian oscillator upon

plant development and rosette architecture, but this variation in leaf area does not account

for the majority of the influence of circadian regulation upon WUE. It also further supports

the notion that the influence of the circadian oscillator upon WUE is complex, and cannot be

explained by variation in one of water use or biomass accumulation alone.

Circadian regulation within guard cells alone contributes to water use efficiency

Next, we identified that the circadian oscillator within guard cells contributes to WUE. There

is evidence that guard cells contain a circadian oscillator that regulates stomatal opening

(Gorton et al., 1989; Hassidim et al., 2017). To investigate the contribution of the guard cell

circadian oscillator to WUE, we overexpressed two circadian oscillator components (*CCA1*,

TOC1) in guard cells, using two guard cell-specific promoters (*GC1*, *MYB60*) for each of

CCA1 and *TOC1* (Fig. 4A) (Cominelli et al., 2005; Galbiati et al., 2008; Yang et al., 2008;

Nagy et al., 2009; Meyer et al., 2010; Cominelli et al., 2011; Bauer et al., 2013; Rusconi et

al., 2013). *GC1* is a strong guard cell-specific promoter that is relatively unresponsive to a

variety of environmental cues (cold, light, ABA, gibberellin) (Yang et al. 2008). We used the

full-length *MYB60* promoter sequence, because truncated and chimeric versions of this

promoter appear to have weaker activity and/or become rapidly downregulated by

dehydration and ABA (Francia et al., 2008; Cominelli et al., 2011; Rusconi et al., 2013). This

produced four sets of transgenic lines; *GC1::CCA1:nos* (*GC*), *GC1::TOC1:nos* (*GT*),

MYB60::CCA1:nos (*MC*) and *MYB60::TOC1:nos* (*MT*). We termed these guard cell specific

(GCS) plants. We confirmed the guard cell specificity of the *GC1* and *MYB60* promoters in

our hands, by driving green fluorescent protein (GFP) under the control of these promoters.

GFP accumulation was restricted to the guard cells (Fig. S2A, B). There was not a circadian

oscillation in the activity of either the *GC1* or *MYB60* promoter under our experimental

conditions (Fig. S2C), demonstrating that these promoters are appropriate for constitutive

overexpression of circadian oscillator components within guard cells under our experimental

conditions.

To further verify the guard cell-specific overexpression of *CCA1* and *TOC1* in the GCS

plants, we examined *CCA1* and *TOC1* transcript accumulation within guard cells. Under

constant light conditions, we measured *CCA1* transcript accumulation in epidermal peels at

dusk (when *CCA1* transcript abundance is normally low in the wild type) and *TOC1*

transcript accumulation at dawn (when *TOC1* transcript abundance is normally low in the

wild type). Guard cell *CCA1* overexpressors had greater *CCA1* transcript abundance in

epidermal peels at dusk than the wild type (*GC*: t_4 = - 2.233, *p*>0.05; *MC*: t_4 = -7.409, *p* =

0.002) (Fig. S2D), and guard cell *TOC1* overexpressors had greater *TOC1* transcript

abundance at dawn than the wild type (*GT*: t_4 = -6.636, *p* = 0.003; *MT*: t_4 = -2.736, *p* =

0.050) (Fig. S2D). These data indicate that *CCA1* and *TOC1* were overexpressed within the

guard cells of the guard cell-specific *CCA1* or *TOC1* overexpressor plants that we

generated, respectively.

We investigated the effect on WUE of overexpression of CCA1 and TOC1 within guard cells.

Two independent *GC1::CCA1* lines (*GC*-1 and *GC*-2) were significantly more water use

efficient than the wild type (*GC*-1: *p* < 0.001; *GC*-2: *p* = 0.002) (Fig. 4B). *GC*-1 and *GC*-2

were 8% and 4% more water use efficient than the wild type, respectively (Fig. 4B). Two

independent *MYB60::CCA1* lines also had numerically higher WUE than the wild type, but

this was not statistically significant (*p* > 0.05) (Fig. 4B). In contrast, overexpression of *TOC1*

in guard cells with both the *GC1* and *MYB60* promoters did not alter WUE (*p* > 0.05) (Fig.

4B). Together, these data suggest that overexpressing CCA1 in guard cells can increase

whole plant long-term WUE.

A previous study identified that constitutive overexpression of TOC1 (TOC1-ox) reduces the

dehydration tolerance of seedlings (Legnaioli et al., 2009). We wished to determine whether

this altered dehydration tolerance is due specifically to the circadian oscillator within guard

cells. Using a similar experimental system to Legnaioli et al. 2009, we found that

MYB60::CCA1 and *GC1::CCA1* increase dehydration survival (Fig. 4C). In contrast,

GC1::TOC1 and *MYB60::TOC1* had decreased dehydration survival relative to the wild type

(Fig. 4C). This suggests that overexpressing *CCA1* or *TOC1* in guard cells can increase or

decrease survival to dehydration under constant light conditions, respectively.

Like *MYB60::CCA1* and *GC1::CCA1*, more seedlings constitutively overexpressing CCA1

(CCA1-ox) survived dehydration under our experimental conditions (Fig. 4C). Similarly, like

GC1::TOC1 and *MYB60::TOC1*, more seedlings overexpressing TOC1 constitutively (TOC1-

ox) were killed by dehydration (Fig. 4C). Therefore, manipulation of the expression of these

clock genes in guard cell and whole plants causes similar phenotypes, with some

differences in magnitude (Fig. 4C). One interpretation is that altered dehydration survival in

CCA1-ox and TOC1-ox seedlings might be partly or wholly due to the circadian clock that is

specifically within guard cells. Because the stomatal density was unaltered relative to the

wild type in the guard cell overexpressors of CCA1 and TOC1 (Fig. 4D, E), the WUE and

dehydration survival phenotypes that we identified might be due to alterations in processes

within guard cells rather than due to altered stomatal density.

Discussion

Pervasive influence of the circadian oscillator upon water use efficiency

Our data indicate that the circadian oscillator is important for regulating the long-term WUE

of Arabidopsis. Misregulation of several functional subsections of the circadian oscillator

altered the WUE of Arabidopsis. Misexpression of morning (PRR7, PRR9, CCA1), late day

(GI, PRR5) and evening (TOC1, ZTL, ELF3) components of the circadian oscillator all

perturb WUE under our experimental conditions (Fig. 1, Fig. 2B). Additionally, altered

expression of TEJ and GRP7 also alters WUE (Fig. 1). Therefore, oscillator components that

impact WUE are not confined to a specific region or expression phase of the multi-loop

circadian oscillator. Misexpression of genes encoding some proteins that provide

environmental inputs to the circadian oscillator (ELF3, TPS1, ZTL, KIN10; (Covington et al.,

2001; Kim et al., 2007; Shin et al., 2017; Frank et al., 2018)) also alters WUE. Together, this

suggests that the entire circadian oscillator influences WUE, and that alterations in water

use that are caused by mutations to the circadian oscillator are not confined to a specific

sub-loop of the circadian oscillator or restricted to its input or output pathways. One

explanation for these circadian-system wide alterations in WUE relates to the nature of

feedback within the circadian oscillator. The complex feedback and interconnectivity of the

circadian oscillator means that individual components of the circadian oscillator that directly

influence stomatal function or water use are likely to be altered by mutations that are distal

to that component. Therefore, if correct circadian timing is required for optimum water use

efficiency, multiple components of the circadian oscillator are likely to influence water use

efficiency.

The sugar signalling proteins TPS1 and KIN10 influence a broad range of phenotypes, in

addition to participating in circadian entrainment (Baena-González et al., 2007; Gómez et al.,

2010; Paul et al., 2010; Delatte et al., 2011; Shin et al., 2017; Frank et al., 2018; Nietzsche

et al., 2018; Simon et al., 2018). The *tps1*-12 TILLING mutant of TPS1 decreases stomatal

aperture and increases the ABA sensitivity of guard cells (Gómez et al., 2010), whereas we

found that *tps1*-11 and *tps1*-12 had lower long-term WUE than the wild type (Fig. 1).

Reduced biomass accumulation in *tps1*-11 and *tps1*-12 (Fig. 2A) was consistent with slow

growth of these alleles (Gómez et al., 2010). Overall, this suggests that the decreased

stomatal aperture of *tps1*-12 mutants does not translate into an overall increase in WUE,

potentially due to slower growth of the *tps1* mutants (Fig. 2A) (Gómez et al., 2010). The

broad range of phenotypes that are altered in *tps1*-11, *tps1*-12 and KIN10-ox 6.5 indicates

that these genotypes might alter WUE through mechanisms other than circadian regulation.

Potential roles for the evening complex in WUE

Our finding that ELF3 is important for WUE (Fig. 1) is supported by previous evidence.

Under constant light conditions, wild type Arabidopsis has circadian rhythms of stomatal

aperture, whereas *elf3* stomata are constantly open and unresponsive to light and dark

(Kinoshita et al., 2011). Furthermore, ELF3 negatively regulates blue light-mediated stomatal

opening (Kinoshita and Hayashi, 2011). Therefore, perturbation of the anticipation of

day/night transitions or responses to environmental cues in *elf3* stomata might cause long-

term alterations in WUE.

ELF3 binds to the *PRR9* promoter and *elf3*-1 has elevated PRR9 transcript abundance

(Thines and Harmon, 2010; Dixon et al., 2011; Herrero et al., 2012). The low WUE of *elf3*-1

might potentially be caused by altered PRR9 expression, because misregulation of *PRR9*

also affected WUE (Fig. 1). In a similar fashion, ELF3/ELF4 signalling represses PRR7, and

elf3-1 has elevated *PRR7* transcript abundance (Herrero et al., 2012). Under light-dark

cycles, *elf3*-1 also has high and constitutive *GI* expression (Fowler et al., 1999), and *elf3*-1

and *gi* mutants have opposite WUE phenotypes (Fig. 1). Therefore, the WUE phenotype of

elf3-1 (Fig. 1) might be caused by disruption of ELF3 itself, or alterations in PRR7, PRR9

and/or GI expression.

Mutating further components of the evening complex (EC) (ELF4 and LUX) did not affect

WUE (Fig. 1). This is despite the way that these genes influence circadian oscillator function

and plant physiology (Hsu and Harmer, 2014; Huang and Nusinow, 2016), and nocturnal

regulation of stomatal aperture impacts WUE (Costa et al., 2015; Coupel-Ledru et al., 2016).

One possibility is that the impact of *elf3* on WUE may be greater than that of *elf4* or *lux*

because ELF3 is key to EC scaffolding, with ELF3 operating genetically downstream from

ELF4 and LUX (Herrero et al., 2012; Huang and Nusinow, 2016).

ELF4 appears to play a greater role in circadian regulation in the vascular tissue than

stomatal guard cells, with vasculature expression up to ten times higher than other tissues

(Endo et al., 2014). Because *elf3*-1 affects WUE differently from *elf4*-101 and *lux*-1 (Fig. 1), it

appears that ELF3 regulates WUE independently from ELF4 and LUX.

Multiple physiological causes of altered WUE in circadian oscillator mutants

Our data suggest that changes in WUE caused by misexpression of circadian clock

components might be due to a combination of physiological factors. Some mutants or

overexpressors tested alter biomass accumulation, whilst others predominantly alter water

loss (Fig. 2), so mutations to the circadian oscillator did not alter water use by specifically

altering one of carbon assimilation or transpiration. This is consistent with previous work

demonstrating that both stomatal opening and CO_2 fixation is perturbed in circadian

arrhythmic plants under light/dark cycles (Dodd et al., 2005), and with the findings that daily

carbohydrate management is dependent upon correct circadian regulation (Graf et al.,

2010). We speculate that delayed or advanced stomatal and photosynthetic responses to

the day-night cycle might occur in circadian period mutants, because period mutants

inaccurately anticipate the onset of dawn (Dodd et al., 2014). Circadian clock mutants might

also affect WUE by changing the sensitivity of stomatal movements and photosynthesis to

environmental transitions, because there is circadian gating of the responses of both

stomata and photosynthesis to environmental cues (Dodd et al., 2006; Kinoshita et al., 2011;

Litthauer et al., 2015; Joo et al., 2017; Cano-Ramirez et al., 2018). Some effects of the

circadian oscillator upon WUE arise from alterations in leaf size that occur in some circadian

oscillator mutants (Fig. 3A, B). This suggests that developmental alterations arising from

lesions in the circadian oscillator can lead to changes in WUE. Such developmental

alterations might alter WUE by changing airflow around the rosette, boundary layer

conductance, or internal leaf structure.

Conclusions

We show that circadian regulation contributes to whole plant long-term WUE under cycles of

day and night. This control occurs partly through the influence of the circadian oscillator

upon rosette architecture. Mutation or overexpression of CCA1, TOC1, ELF3, GI, GRP7,

PRR5, PRR7, PRR9, TEJ and ZTL altered WUE under our experimental conditions. The

roles of these genes in WUE may be independent or overlapping, and their WUE

phenotypes might be due to direct effects of these genes, or indirect effects on transcript

and/or protein abundance of other circadian clock gene(s). Misregulation of the expression

of CHE, FKF1, LKP2, RVE4, RVE8, PRR3, ELF4, LUX and WNK1 did not appear to alter

WUE under our experimental conditions.

Our results have a number of broad implications. Firstly, our data suggest that alterations in

circadian function that arise during crop breeding could have the potential to increase or

decrease WUE. Therefore, manipulation of the functioning of the circadian oscillator might

represent a pathway to tune the WUE of crops. Second, our results indicate that circadian

regulation in a single cell type can have implications for whole-plant physiology. Finally, our

findings suggest that circadian regulation can alter a single trait (WUE) by affecting many

aspects of physiology. In future, it would be informative to distinguish the contribution to

overall WUE of circadian regulation within additional cell types, such as the mesophyll,

vascular tissue, and root cell types.

Materials and methods

Plant material and growth conditions

Arabidopsis (*Arabidopsis thaliana* (L.) Heynh.) seeds were surface-sterilised as described

previously (Noordally et al., 2013). For experiments investigating stomatal density and index,

seeds were stratified for 3 days at 4 °C, then sown on compost mix comprising a 3:1 ratio of

coarsely sieved Levington Advance F2 seed compost (Everris) and horticultural silver sand

(Melcourt), supplemented with 0.4 g l^{-1} thiacloprid insecticide granules (Exemptor; Everris).

Plants were grown in controlled environment chambers (Reftech, Netherlands) under an 8 h

photoperiod at 70% humidity, 20 °C, and photon flux density of 100 µmol m^{-2} s^{-1} of overhead

lighting supplied by cool white fluorescent tubes (Reftech, Netherlands). For experiments

investigating long-term WUE, seeds were sown within a custom Falcon tube system then

stratified. Plants were cultivated in plant growth chambers (Snjider, Netherlands) under the

experimental conditions described above. The genotypes that were screened for WUE

alterations are identified in Table S1, and all have been described previously. For all

experiments, at least two completely independent experimental repeats were performed per

genotype and per treatment, with multiple replicate plants within each of the experimental

repeats.

Generation of transgenic lines

To create the *GC1::CCA1:nos* (*GC*), *GC1::TOC1:nos* (*GT*), *MYB60::CCA1:nos* (*MC*) and

MYB60::TOC1:nos (*MT*) constructs, the CaMV *nos* terminator sequence was ligated

between the SpeI and NotI restriction sites in the pGREENII0229 binary vector (Hellens et

al., 2000). The *GC1* upstream sequence (-1894 to -190) or *MYB60* upstream sequence (-

1724 to -429) was then ligated between the KpnI and ApaI restriction sites of

pGREENII0229. Finally, the *CCA1* coding sequence or *TOC1* coding sequence, obtained

using RT-PCR, was ligated between the restriction XhoI and XmaI sites. Primers used are

identified in Table S2. Constructs were transformed into Col-0 wild type Arabidopsis using

transformation with *Agrobacterium tumefaciens* strain GV3101. Transformants were

identified by screening for phosphinothricin resistance, then further validated using genomic

DNA PCR. Homozygous lines were identified via phosphinothricin (BASTA) resistance, and

two independently transformed homozygous lines were investigated in detail per genotype.

Guard cell specificity of promoter activity was investigated using *GC1::GFP:nos* and

MYB60::GFP:nos promoter-reporter lines (Sup. Fig. 3A-C), which were created as above

with the *GFP* coding sequence ligated between the XhoI and XmaI restriction sites. Leaf

discs (5 mm diameter) from seedlings or mature plants were mounted on microscope slides

with dH_2O, and examined for GFP fluorescence using confocal microscopy (Leica DMI6000).

The following settings were used: argon laser at 20% capacity, 488 nm laser at 48%

capacity with a bandwidth of 505 nm–515 nm, gain of 1250, offset at 0.2%, 20x or 40x

objective, zoom x1 to x4.

Measurement of water use efficiency

The WUE assay was adapted from Wituszynska et al. (2013) (Wituszyńska et al., 2013).

Plants were grown for 6 weeks in modified 50 ml Falcon tubes. The Falcon tube systems

consisted of a 50 ml Falcon tube filled with 37.5 ml of a 1:1 ratio of compost: perlite and 35

ml of Milli-Q water (Merck), with the remaining volume filled with a 1:1 ratio of compost: Milli-

Q water (Fig. S3). Each Falcon tube lid had a 2 mm diameter hole drilled in its centre to

allow plant growth. The lid was spray-painted black (Hycote) because we found that the

orange colour of the Falcon tube lid caused leaf curling (Fig. S3). The system was wrapped

in aluminium foil to exclude light (Fig. S3). 10-15 seeds were sown through the Falcon tube

lid using a pipette. Following stratification, Falcon tube systems were placed under growth

conditions using a randomised experimental design. 7 days after germination, seedlings

were trimmed to one per Falcon tube system, and initial Falcon tube weight was recorded.

After 6 weeks of growth, rosette leaf surface area was measured by photography (D50;

Nikon) and Fiji software, rosette dry weight was measured (4 d at 60°C), and final Falcon

tube weight was recorded. Negative controls (Falcon tube systems without plants) were

used to assess soil water evaporation over 18 experimental repeats, with an overall mean

weight loss of 0.513 g ± 0.004 g over 6 weeks for plant-free Falcon tubes.

Plant WUE was calculated as follows:

$$WUE = \frac{d}{(t_i - t_f) - e}$$

Where d is the rosette dry weight at the end of the experiment (mg), t_i and t_f are the falcon

tube weight at the start and end of the experiment, respectively (g), and e is the amount of

water evaporation directly from the compost (g). WUE is derived as mg biomass per ml^{-1}

water lost. These calculations assumed that 1 g of weight change was equivalent to a

change of 1 ml of water. For each of 3 independent experimental repeats, 15 plants were

screened per genotype. Due to variation between the WUE of each background (Fig. S1),

the WUE of each circadian oscillator genotype was normalized to its respective background

and expressed as a percentage of that background. Statistical comparisons with the wild

types were conducted before this normalization.

Dehydration tolerance assay

This assay was adapted from Legnaioli et al. (2009). For experiments investigating survival

to dehydration, surface-sterilized seeds were sown on Petri dishes containing half strength

Murashige & Skoog basal salt mixture (0.5x MS) (Duchefa), supplemented with 0.8% (w/v)

agar and 3% (w/v) sucrose, then stratified for 3 days at 4 °C before transfer to the growth

chamber. For these experiments, seedlings were cultivated in MLR-352 growth chambers

(Panasonic) at 19°C with photon flux density of 120 µmol m^{-2} s^{-1}. 14-day old seedlings were

dehydrated on a double layer of filter paper (Fisher Scientific) for 9 h under constant light

conditions, watered with sterile dH_2O, and kept under constant light conditions for a further

48 h before being scored for survival. Seedlings with a green apical meristematic region

were counted as survivors. 32 seedlings were treated per genotype and within each

experimental repeat.

Measurement of stomatal density

Plants were grown for 7-8 weeks on compost mix. Dental paste (Coltene) was applied to the

abaxial surface of fully expanded leaves. Transparent nail varnish (Rimmel) was applied to

these leaf moulds once they had set, and then peeled away from the mould using clear

adhesive tape (Scotch Crystal). Stomatal and pavement cells were counted within an

800 µm x 800 µm square at the centre of each leaf half, using an epifluorescence

microscope (HAL100; Zeiss) and Volocity (Perkin Elmer) and Fiji software. For each

experimental repeat, two leaves were sampled per plant and eight plants sampled per

genotype. Stomatal index was calculated as follows:

$$SI = \frac{s}{s + p} * 100$$

Where *SI* is the stomatal index, *s* the number of stomata in the field of view (800 µm x

800 µm), and *p* the number of pavement cells in the field of view.

RNA extraction and qRT-PCR

RNA extractions, cDNA synthesis, and qRT-PCR were performed according to (Simon et al.,

2018), except approximately 10 seedlings were used per RNA sample and analysis was

performed using an MXPro 3005 real time PCR system (Agilent) with 5x HOT FIREPol

EvaGreen qPCR mastermix (Solis Biodyne). qRT-PCR primers are provided in Table S3.

Rhythmic features within qPCR data were identified using the BioDare2 platform (Zielinski et

al., 2014), using the Fast Fourier Transform Non-Linear Least Squares method (FFT-NLLS).

One independently-transformed line of each guard cell-specific circadian clock gene

overexpressor was also investigated using qRT-PCR conducted on RNA isolated from

epidermal peels. Abaxial leaf epidermis was detached, then washed in 10 mM MES (pH

6.15, adjusted using 10 M KOH) to remove RNA derived from ruptured epidermal cells. Each

RNA sample was derived from 20 epidermal peels (five plants, four leaves per plant) that

were collated and flash-frozen in liquid nitrogen. Guard cell RNA was extracted using the

RNeasy UCP Micro Kit (Qiagen) according to manufacturer's instructions, with the following

modification: guard cell lysis was performed by adding glass beads (425 µm - 600 µm

diameter, acid washed, from Sigma-Aldrich) and 350 µl RULT buffer to the sample, then

vortexed for 5 min.

Accession numbers

Arabidopsis Genome Initiative identifiers for the genes mentioned in this study are: *CCA1*

(*CIRCADIAN CLOCK ASSOCIATED1*, At2g46830), *CHE* (*CCA1 HIKING EXPEDITION*,

At5g08330), *ELF3* (*EARLY FLOWERING3*, At2g25930), *ELF4* (*EARLY FLOWERING4*,

At2g40080), *FKF1* (*F BOX1*, At1g68050), *GI* (*GIGANTEA*, At1g22770), *GRP7* (*GLYCINE*

RICH PROTEIN7, At2g21660), *KIN10* (*SNF1-RELATED PROTEIN KINASE1.1*, At3g01090),

LKP2 (*LOV KELCH PROTEIN2*, At2g18915), *LUX* (*LUX ARRHYTHMO*, At3g46640),

MYB60 (*MYB DOMAIN PROTEIN60*, At1g08810), *PRR3* (*PSEUDO-RESPONSE*

REGULATOR3, At5g60100), *PRR5* (*PSEUDO-RESPONSE REGULATOR5*, At5g24470),

PRR7 (PSEUDO-RESPONSE REGULATOR7, At5g02810), *PRR9* (*PSEUDO-RESPONSE*

REGULATOR9, At2g46790), *RVE4* (*REVEILLE4*, At5g02840), *TEJ* (*POLY(ADP-*

RIBOSE)GLYCOHYDROLASE1, At2g31870), *TIC* (*TIME FOR COFFEE*, At3gt22380),

TOC1 (*TIMING OF CAB EXPRESSION1*, At5g61380), *TPS1* (*TREHALOSE-6-PHOSPHATE*

SYNTHASE1, At1g78580), *WNK1* (*WITH NO LYSINE KINASE1*, At3g04910), *ZTL*

(*ZEITLUPE*, At5g57360).

Acknowledgements

We thank Kester Cragg-Barber, James Chen, Ioanna Kostaki, Jean-Charles Isner, Deirdre McLachlan, Peng Sun, Ashutosh Sharma and Dora Cano-Ramirez for technical advice during experimentation. We thank Marc Knight and Tracy Lawson for helpful discussions concerning data interpretation. We are grateful to Keara Franklin, Alex Webb, Paloma Mas, Steve Kay, Isabelle Carre, Takato Imaizumi, Filip Rolland, Ian Graham, Stacey Harmer, and Steven Penfield for donating seed lines for this study. This research was funded the UK Biotechnology and Biological Sciences Research Council (BBSRC; grant BB/J014400/1) and the Wolfson Foundation.

 Figure legends

Figure 1. The circadian clock regulates long-term water use efficiency of Arabidopsis under

light/dark cycles. The WUE of circadian clock mutants and overexpressors is expressed as a

percentage of their respective background (normalized to 100%, red reference line) to

account for WUE variation between background accessions (n = 5 - 15). Data were analysed

using independent-samples t-tests and statistical significance is indicated relative to the

background using (* = p < 0.05; ** = p < 0.01; *** = p < 0.001). Statistical analysis was

performed on raw data, with data subsequently converted to a percentage of the wild type

for the purposes of comparison and presentation. Screens were repeated independently

three times per genotype, with one representative experimental repeat shown here for each

genotype.

Figure 2. Altered WUE of plants with mutations or overexpression of circadian clock

associated genes is not caused consistently by variation in one of dry weight, water use,

phase of expression of each gene, or resultant altered period or flowering time. Data are

derived from Fig. 1 and expressed as a percentage of the respective background (WT,

normalised to 100%, red reference line) (n = 5 - 15). (A) Altered WUE is not specifically due

to altered water use or altered dry weight of screened genotypes, but results from the

combination of both. (b-d) Variations in WUE are not explained by (B) phase expression of

mutated/overexpressed gene, nor by altered (C) period or (D) flowering time of the

mutated/overexpressor genotype. Genotypes reported to have no change (N/C) in period or

flowering time relative to the wild type are included on the left of panels (B) and (C), while

those for which period and/or flowering time are unknown are included on the right. Studies

describing the phase of expression, period and flowering time of the genotypes tested are

identified in the main text. We note that the phase of expression and period data used for

this analysis were often obtained under constant conditions, in contrast to our experiments

occurring under light/dark cycles.

Figure 3. The circadian oscillator alters WUE partially by changing rosette architecture. (A)

Altering circadian-associated gene expression can affect rosette architecture and size, as

illustrated for *elf3*-1, *lux*-1, and *gi*-2 in (Col-0 background). Image backgrounds removed for

clarity. Variation in rosette leaf surface area across the genotypes investigated explained (B)

16% of variation in WUE ($p < 0.001$, $r = 0.400$, $r^2 = 0.160$), (C) 83% of variation in

transpiration ($p < 0.001$, $r = 0.912$, $r^2 = 0.832$) and (D) 73% of variation in rosette dry

biomass ($p < 0.001$, $r = 0.857$, $r^2 = 0.734$). Data were analysed using Pearson correlation

tests.

Figure 4. Overexpressing *CCA1* or *TOC1* in guard cells affects WUE and survival of

dehydration by seedlings. (A) Constructs used to overexpress *CCA1* or *TOC1* coding

sequence under control of *GC1* or *MYB60* promoters. (B) Guard cell *CCA1* overexpression

can increase WUE. WUE expressed as a percentage of the wild type (normalised to 100%,

red reference line). Two to four independent experimental repeats were performed, with data

from one representative dataset shown ($n = 5$ - 15). Data for CCA1-ox and TOC1-ox are

derived from Fig. 1, for purposes of comparison. Data were analysed with independent

samples t-tests, and statistical significance compared to Col-0 is indicated using starring (**

$= p < 0.01$; *** $= p < 0.001$). (C) Guard cell *CCA1* or *TOC1* overexpression alters

dehydration survival of seedlings compared with the wild type. Data were obtained from

three independent experimental repeats (mean; $n = 32$ per experimental replicate; at least

two independent experimental repeats were performed for each genotype). A single

GC1::TOC1 line is shown here because other lines produced extremely variable data. (D,E)

Guard cell *CCA1* or *TOC1* overexpression does not affect (D) stomatal index nor (E)

stomatal density. Two independent experimental repeats were performed, with data from

one representative dataset shown ($n = 19$ - 32; mean ± S.E.M.). Data were analysed with

ANOVA and Tukey's post hoc tests (NS $= p > 0.05$). Bar colours identify the whole plant

overexpressor control (black), wild type control (dark grey), and guard cell-specific

overexpressor genotypes (light grey).

References

Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature **448:** 938

Baudry A, Ito S, Song YH, Strait AA, Kiba T, Lu S, Henriques R, Pruneda-Paz JL, Chua N-H, Tobin EM, Kay SA, Imaizumi T (2010) F-Box proteins FKF1 and LKP2 act in concert with ZEITLUPE to control Arabidopsis clock progression. The Plant Cell **22:** 606

Bauer H, Ache P, Lautner S, Fromm J, Hartung W, Al-Rasheid Khaled AS, Sonnewald S, Sonnewald U, Kneitz S, Lachmann N, Mendel Ralf R, Bittner F, Hetherington Alistair M, Hedrich R (2013) The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. Current Biology **23:** 53-57

Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. Field Crops Research **112:** 119-123

Bridge LJ, Franklin KA, Homer ME (2013) Impact of plant shoot architecture on leaf cooling: a coupled heat and mass transfer model. Journal of The Royal Society Interface **10:** 20130326

Caine RS, Yin X, Sloan J, Harrison EL, Mohammed U, Fulton T, Biswal AK, Dionora J, Chater CC, Coe RA, Bandyopadhyay A, Murchie EH, Swarup R, Quick WP, Gray JE (2019) Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. New Phytologist **221:** 371-384

Cano-Ramirez DL, Saskia de Fraine T, Griffiths OG, Dodd AN (2018) Photosynthesis and circadian rhythms regulate the buoyancy of marimo lake balls. Current Biology **28:** R869-R870

Cominelli E, Galbiati M, Albertini A, Fornara F, Conti L, Coupland G, Tonelli C (2011) DOF-binding sites additively contribute to guard cell-specificity of *AtMYB60* promoter. BMC Plant Biology **11:** article 162

Cominelli E, Galbiati M, Vavasseur A, Conti L, Sala T, Vuylsteke M, Leonhardt N, Dellaporta SL, Tonelli C (2005) A guard-cell-specific MYB transcription factor regulates stomatal movements and plant drought tolerance. Current Biology **15:** 1196-1200

Condon AG, Farquhar GD, Rebetzke GJ, Richards RA (2004) Breeding for high water-use efficiency. Journal of Experimental Botany **55:** 2447-2460

Costa JM, Monnet F, Jannaud D, Leonhardt N, Ksas B, Reiter IM, Pantin F, Genty B (2015) *OPEN ALL NIGHT LONG*: The dark side of stomatal control. Plant Physiology **167:** 289

Coupel-Ledru A, Lebon E, Christophe A, Gallo A, Gago P, Pantin F, Doligez A, Simonneau T (2016) Reduced nighttime transpiration is a relevant breeding target for high water-use efficiency in grapevine. Proceedings of the National Academy of Sciences **113:** 8963

Covington MF, Panda S, Liu XL, Strayer CA, Wagner DR, Kay SA (2001) ELF3 modulates resetting of the circadian clock in Arabidopsis. The Plant Cell **13:** 1305

Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg A, Primavesi LF, Paul MJ, Schluepmann H (2011) Growth arrest by trehalose-6-phosphate: An astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. Plant Physiology **157:** 160

Ding Z, Millar AJ, Davis AM, Davis SJ (2007) *TIME FOR COFFEE* encodes a nuclear regulator in the *Arabidopsis thaliana* circadian clock. The Plant Cell **19:** 1522

Dixon LE, Knox K, Kozma-Bognar L, Southern MM, Pokhilko A, Millar AJ (2011) Temporal repression of core circadian genes is mediated through EARLY FLOWERING 3 in Arabidopsis. Current Biology **21:** 120-125

Dodd AN, Dalchau N, Gardner MJ, Baek S-J, Webb AAR (2014) The circadian clock has transient plasticity of period and is required for timing of nocturnal processes in Arabidopsis. New Phytologist **201:** 168-179

Dodd AN, Jakobsen MK, Baker AJ, Telzerow A, Hou S-W, Laplaze L, Barrot L, Scott
Poethig R, Haseloff J, Webb AAR (2006) Time of day modulates low-temperature
Ca^{2+} signals in Arabidopsis. The Plant Journal **48:** 962-973
Dodd AN, Parkinson K, Webb AAR (2004) Independent circadian regulation of assimilation
and stomatal conductance in the *ztl*-1 mutant of Arabidopsis. New Phytologist **162:**
63-70
Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, Hibberd JM, Millar AJ, Webb
AAR (2005) Plant circadian clocks increase photosynthesis, growth, survival, and
competitive advantage. Science **309:** 630
Doyle MR, Davis SJ, Bastow RM, McWatters HG, Kozma-Bognár L, Nagy F, Millar AJ,
Amasino RM (2002) The ELF4 gene controls circadian rhythms and flowering time
in *Arabidopsis thaliana*. Nature **419:** 74
Edwards CE, Ewers BE, McClung CR, Lou P, Weinig C (2012) Quantitative variation in
water-use efficiency across water regimes and its relationship with circadian,
vegetative, reproductive, and leaf gas-exchange traits. Molecular Plant **5:** 653-668
Edwards CE, Weinig C (2010) The quantitative-genetic and QTL architecture of trait
integration and modularity in *Brassica rapa* across simulated seasonal settings.
Heredity **106:** 661
Endo M, Shimizu H, Nohales MA, Araki T, Kay SA (2014) Tissue-specific clocks in
Arabidopsis show asymmetric coupling. Nature **515:** 419
Eriksson ME, Hanano S, Southern MM, Hall A, Millar AJ (2003) Response regulator
homologues have complementary, light-dependent functions in the Arabidopsis
circadian clock. Planta **218:** 159-162
Farré EM, Harmer SL, Harmon FG, Yanovsky MJ, Kay SA (2005) Overlapping and
distinct roles of PRR7 and PRR9 in the Arabidopsis circadian clock. Current Biology
15: 47-54
Ferguson JN, Humphry M, Lawson T, Brendel O, Bechtold U (2018) Natural variation of
life-history traits, water use, and drought responses in Arabidopsis. Plant Direct **2:**
e00035
Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill
J (1999) *GIGANTEA*: a circadian clock- controlled gene that regulates photoperiodic
flowering in Arabidopsis and encodes a protein with several possible membrane-
spanning domains. The EMBO Journal **18:** 4679
Francia P, Simoni L, Cominelli E, Tonelli C, Galbiati M (2008) Gene trap-based
identification of a guard cell promoter in Arabidopsis. Plant Signaling & Behavior **3:**
684-686
Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D,
Yochikawa A, Cano-Ramirez DL, Chembath A, Cragg-Barber K, Haydon MJ,
Hotta CT, Vincentz M, Webb AAR, Dodd AN (2018) Circadian entrainment in
Arabidopsis by the sugar-responsive transcription factor bZIP63. Current Biology **28:**
2597-2606.e2596
Franks PJ, W. Doheny-Adams T, Britton-Harper ZJ, Gray JE (2015) Increasing water-use
efficiency directly through genetic manipulation of stomatal density. New Phytologist
207: 188-195
Galbiati M, Simoni L, Pavesi G, Cominelli E, Francia P, Vavasseur A, Nelson T, Bevan
M, Tonelli C (2008) Gene trap lines identify Arabidopsis genes expressed in
stomatal guard cells. The Plant Journal **53:** 750-762
Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA (2010) AtTPS1-mediated trehalose 6-
phosphate synthesis is essential for embryogenic and vegetative growth and
responsiveness to ABA in germinating seeds and stomatal guard cells. The Plant
Journal **64:** 1-13
Gorton HL, Williams WE, Binns ME, Gemmell CN, Leheny EA, Shepherd AC (1989)
Circadian stomatal rhythms in epidermal peels from *Vicia faba*. Plant Physiology **90:**
1329

Graf A, Schlereth A, Stitt M, Smith AM (2010) Circadian control of carbohydrate
availability for growth in Arabidopsis plants at night. Proceedings of the National
Academy of Sciences **107:** 9458
Hall A, Bastow RM, Davis SJ, Hanano S, McWatters HG, Hibberd V, Doyle MR, Sung S,
Halliday KJ, Amasino RM, Millar AJ (2003) The *TIME FOR COFFEE* gene
maintains the amplitude and timing of Arabidopsis circadian clocks. The Plant Cell
15: 2719
Harmer SL, Hogenesch JB, Straume M, Chang H-S, Han B, Zhu T, Wang X, Kreps JA,
Kay SA (2000) Orchestrated transcription of key pathways in *Arabidopsis* by the
circadian clock. Science **290:** 2110
Hassidim M, Dakhiya Y, Turjeman A, Hussien D, Shor E, Anidjar A, Goldberg K, Green
RM (2017) *CIRCADIAN CLOCK ASSOCIATED1* (*CCA1*) and the circadian control of
stomatal aperture. Plant Physiology **175:** 1864
Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR (2013)
Photosynthetic entrainment of the *Arabidopsis thaliana* circadian clock. Nature **502:**
689
Hazen SP, Schultz TF, Pruneda-Paz JL, Borevitz JO, Ecker JR, Kay SA (2005) *LUX*
ARRHYTHMO encodes a Myb domain protein essential for circadian rhythms.
Proceedings of the National Academy of Sciences of the United States of America
102: 10387
Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM (2000) pGreen: a
versatile and flexible binary Ti vector for Agrobacterium-mediated plant
transformation. Plant Molecular Biology **42:** 819-832
Hennessey TL, Field CB (1991) Circadian rhythms in photosynthesis. Plant Physiology **96:**
831
Herrero E, Kolmos E, Bujdoso N, Yuan Y, Wang M, Berns MC, Uhlworm H, Coupland
G, Saini R, Jaskolski M, Webb A, Gonçalves J, Davis SJ (2012) EARLY
FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the
Arabidopsis circadian clock. The Plant Cell **24:** 428
Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving
environmental change. Nature **424:** 901
Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. Trends in
Plant Science **19:** 240-249
Hu H, Xiong L (2014) Genetic engineering and breeding of drought-resistant crops. Annual
Review of Plant Biology **65:** 715-741
Huang H, Nusinow DA (2016) Into the evening: Complex interactions in the Arabidopsis
circadian clock. Trends in Genetics **32:** 674-686
Hugouvieux V, Kwak JM, Schroeder JI (2001) An mRNA cap binding protein, ABH1,
modulates early abscisic acid signal transduction in Arabidopsis. Cell **106:** 477-487
Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA (2005) FKF1 F-Box protein mediates
cyclic degradation of a repressor of CONSTANS in *Arabidopsis*. Science **309:** 293
Imaizumi T, Tran HG, Swartz TE, Briggs WR, Kay SA (2003) FKF1 is essential for
photoperiodic-specific light signalling in Arabidopsis. Nature **426:** 302
Joo Y, Fragoso V, Yon F, Baldwin IT, Kim S-G (2017) Circadian clock component, LHY,
tells a plant when to respond photosynthetically to light in nature. Journal of
Integrative Plant Biology **59:** 572-587
Karaba A, Dixit S, Greco R, Aharoni A, Trijatmiko KR, Marsch-Martinez N, Krishnan A,
Nataraja KN, Udayakumar M, Pereira A (2007) Improvement of water use
efficiency in rice by expression of *HARDY*, an *Arabidopsis* drought and salt tolerance
gene. Proceedings of the National Academy of Sciences **104:** 15270
Khanna R, Kikis EA, Quail PH (2003) *EARLY FLOWERING 4* functions in Phytochrome B-
regulated seedling de-etiolation. Plant Physiology **133:** 1530
Kikis EA, Khanna R, Quail PH (2005) ELF4 is a phytochrome-regulated component of a
negative-feedback loop involving the central oscillator components CCA1 and LHY.
The Plant Journal **44:** 300-313

Kim W-Y, Fujiwara S, Suh S-S, Kim J, Kim Y, Han L, David K, Putterill J, Nam HG,
Somers DE (2007) ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA
in blue light. Nature **449:** 356
Kinoshita T, Hayashi Y (2011) New insights into the regulation of stomatal opening by blue
light and plasma membrane H$^+$-ATPase. *In* KW Jeon, ed, International Review of Cell
and Molecular Biology, Vol 289. Academic Press, pp 89-115
Kinoshita T, Ono N, Hayashi Y, Morimoto S, Nakamura S, Soda M, Kato Y, Ohnishi M,
Nakano T, Inoue S-i, Shimazaki K-i (2011) FLOWERING LOCUS T regulates
stomatal opening. Current Biology **21:** 1232-1238
Lawson T, Blatt MR (2014) Stomatal size, speed, and responsiveness impact on
photosynthesis and water use efficiency. Plant Physiology **164:** 1556
Legnaioli T, Cuevas J, Mas P (2009) TOC1 functions as a molecular switch connecting the
circadian clock with plant responses to drought. The EMBO Journal **28:** 3745
Litthauer S, Battle MW, Lawson T, Jones MA (2015) Phototropins maintain robust
circadian oscillation of PSII operating efficiency under blue light. The Plant Journal
83: 1034-1045
Más P, Kim W-Y, Somers DE, Kay SA (2003) Targeted degradation of TOC1 by ZTL
modulates circadian function in *Arabidopsis thaliana*. Nature **426:** 567
Masle J, Gilmore SR, Farquhar GD (2005) The *ERECTA* gene regulates plant transpiration
efficiency in Arabidopsis. Nature **436:** 866
Medrano H, Tomás M, Martorell S, Flexas J, Hernández E, Rosselló J, Pou A, Escalona
J-M, Bota J (2015) From leaf to whole-plant water use efficiency (WUE) in complex
canopies: Limitations of leaf WUE as a selection target. The Crop Journal **3:** 220-228
Meyer S, Mumm P, Imes D, Endler A, Weder B, Al-Rasheid KAS, Geiger D, Marten I,
Martinoia E, Hedrich R (2010) AtALMT12 represents an R-type anion channel
required for stomatal movement in Arabidopsis guard cells. The Plant Journal **63:**
1054-1062
Michael TP, Salomé PA, Yu HJ, Spencer TR, Sharp EL, McPeek MA, Alonso JM, Ecker
JR, McClung CR (2003) Enhanced fitness conferred by naturally occurring variation
in the circadian clock. Science **302:** 1049
Millar AJ (2004) Input signals to the plant circadian clock. Journal of Experimental Botany
55: 277-283
Millar AJ, Carre IA, Strayer CA, Chua NH, Kay SA (1995) Circadian clock mutants in
Arabidopsis identified by luciferase imaging. Science **267:** 1161
Murakami M, Yamashino T, Mizuno T (2004) Characterization of circadian-associated
APRR3 pseudo-response regulator belonging to the APRR1/TOC1 quintet in
Arabidopsis thaliana. Plant and Cell Physiology **45:** 645-650
Na J-K, Metzger JD (2014) Chimeric promoter mediates guard cell-specific gene expression
in tobacco under water deficit. Biotechnology Letters **36:** 1893-1899
Nagy R, Grob H, Weder B, Green P, Klein M, Frelet A, Schjoerring JK, Brearley CA,
Martinoia E (2009) The Arabidopsis ATP-binding cassette protein
ATMRP5/ATABCC5 is a high-affinity inositol hexakisphosphate transporter involved
in guard cell signaling and phytate storage. Journal of Biological Chemistry
Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010) PSEUDO-
RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in *Arabidopsis*
circadian clock. The Plant Cell **22:** 594
Nakamichi N, Murakami-Kojima M, Sato E, Kishi Y, Yamashino T, Mizuno T (2002)
Compilation and characterization of a novel WNK family of protein kinases in
Arabidopsis thaliana with reference to circadian rhythms. Bioscience, Biotechnology,
and Biochemistry **66:** 2429-2436
Nienhuis J, Sills GR, Martin B, King G (1994) Variance for water-use efficiency among
ecotypes and recombinant inbred lines of *Arabidopsis thaliana* (Brassicaceae).
American Journal of Botany **81:** 943-947

Nietzsche M, Guerra T, Alseekh S, Wiermer M, Sonnewald S, Fernie AR, Börnke F (2018) STOREKEEPER RELATED1/G-element binding protein (STKR1) interacts with protein kinase SnRK1. Plant Physiology **176:** 1773

Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma M, Tanaka K, Hanaoka M, Dodd AN (2013) Circadian control of chloroplast transcription by a nuclear-encoded timing signal. Science **339:** 1316

Panda S, Poirier GG, Kay SA (2002) *tej* defines a role for poly(ADP-ribosyl)ation in establishing period length of the *Arabidopsis* circadian oscillator. Developmental Cell **3:** 51-61

Paul MJ, Jhurreea D, Zhang Y, Primavesi LF, Delatte T, Schluepmann H, Wingler A (2010) Up-regulation of biosynthetic processes associated with growth by trehalose 6-phosphate. Plant Signaling & Behavior **5:** 386-392

Pei Z-M, Ghassemian M, Kwak CM, McCourt P, Schroeder JI (1998) Role of farnesyltransferase in ABA regulation of guard cell anion channels and plant water loss. Science **282:** 287

Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. Science **323:** 1481

Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, Salemi MR, Phinney BS, Harmer SL (2011) REVEILLE8 and PSEUDO-REPONSE REGULATOR5 form a negative feedback loop within the Arabidopsis circadian clock. PLOS Genetics **7:** e1001350

Resco de Dios V, Gessler A, Pedro Ferrio J, Alday JG, Bahn M, del Castillo J, Devidal S, García-Muñoz S, Kayler Z, Landais D, Martín-Gómez P, Milcu A, Piel C, Pirhofer-Walzl K, Ravel O, Salekin S, Tissue DT, Tjoelker MG, Voltas J, Roy J (2016) Circadian rhythms have significant effects on leaf-to-canopy scale gas exchange under field conditions. GigaScience **5**

Rubin MJ, Brock MT, Baker RL, Wilcox S, Anderson K, Davis SJ, Weinig C (2018) Circadian rhythms are associated with shoot architecture in natural settings. New Phytologist **219:** 246-258

Ruggiero A, Punzo P, Landi S, Costa A, Van Oosten JM, Grillo S (2017) Improving plant water use efficiency through molecular genetics. Horticulturae **3**

Rusconi F, Simoni L, Simeoni F, Tonelli C, Galbiati M, Cominelli E, Conti L, Riboni M, Francia P, Martin CR (2013) The *Arabidopsis thaliana* MYB60 promoter provides a tool for the spatio-temporal control of gene expression in stomatal guard cells. Journal of Experimental Botany **64:** 3361-3371

Ruts T, Matsubara S, Wiese-Klinkenberg A, Walter A (2012) Aberrant temporal growth pattern and morphology of root and shoot caused by a defective circadian clock in *Arabidopsis thaliana*. The Plant Journal **72:** 154-161

Salomé PA, McClung CR (2005) *PSEUDO-RESPONSE REGULATOR 7* and *9* are partially redundant genes essential for the temperature responsiveness of the Arabidopsis circadian clock. The Plant Cell **17:** 791

Schaffer R, Ramsay N, Samach A, Corden S, Putterill J, Carré IA, Coupland G (1998) The *late elongated hypocotyl* mutation of Arabidopsis disrupts circadian rhythms and the photoperiodic control of flowering. Cell **93:** 1219-1229

Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) Guard cell signal transduction. Annual Review of Plant Physiology and Plant Molecular Biology **52:** 627-658

Schultz TF, Kiyosue T, Yanovsky M, Wada M, Kay SA (2001) A Role for LKP2 in the Circadian Clock of Arabidopsis. The Plant Cell **13:** 2659

Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. Plant, Cell & Environment **40:** 997-1008

Simon NML, Kusakina J, Fernández-López Á, Chembath A, Belbin FE, Dodd AN (2018) The energy-signaling hub SnRK1 is important for sucrose-induced hypocotyl elongation. Plant Physiology **176:** 1299

Simon NML, Sawkins E, Dodd AN (2018) Involvement of the SnRK1 subunit KIN10 in
sucrose-induced hypocotyl elongation. Plant Signaling & Behavior **13:** e1457913
Somers DE, Devlin PF, Kay SA (1998) Phytochromes and cryptochromes in the
entrainment of the *Arabidopsis* circadian clock. Science **282:** 1488
Somers DE, Schultz TF, Milnamow M, Kay SA (2000) *ZEITLUPE* encodes a novel clock-
associated PAS protein from *Arabidopsis*. Cell **101:** 319-329
Somers DE, Webb AA, Pearson M, Kay SA (1998) The short-period mutant, *toc1*-1, alters
circadian clock regulation of multiple outputs throughout development in *Arabidopsis*
thaliana. Development **125:** 485
Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Más P, Panda S, Kreps JA, Kay
SA (2000) Cloning of the *Arabidopsis* clock gene *TOC1*, an autoregulatory response
regulator homolog. Science **289:** 768
Streitner C, Danisman S, Wehrle F, Schöning JC, Alfano JR, Staiger D (2008) The small
glycine-rich RNA binding protein AtGRP7 promotes floral transition in *Arabidopsis*
thaliana. The Plant Journal **56:** 239-250
Thines B, Harmon FG (2010) Ambient temperature response establishes ELF3 as a
required component of the core Arabidopsis circadian clock. Proceedings of the
National Academy of Sciences**:** 200911006
Tomás M, Medrano H, Escalona JM, Martorell S, Pou A, Ribas-Carbó M, Flexas J
(2014) Variability of water use efficiency in grapevines. Environmental and
Experimental Botany **103:** 148-157
Vialet-Chabrand S, Matthews JSA, Brendel O, Blatt MR, Wang Y, Hills A, Griffiths H,
Rogers S, Lawson T (2016) Modelling water use efficiency in a dynamic
environment: An example using *Arabidopsis thaliana*. Plant Science **251:** 65-74
Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE,
Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate
signaling in *Arabidopsis thaliana*. Science **339:** 704
Wang Y, Liu K, Liao H, Zhuang C, Ma H, Yan X (2008) The plant WNK gene family and
regulation of flowering time in Arabidopsis. Plant Biology **10:** 548-562
Wang ZY, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK
ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own
expression. Cell **93:** 1207-1217
Wituszyńska W, Ślesak I, Vanderauwera S, Szechyńska-Hebda M, Kornaś A, Van Der
Kelen K, Mühlenbock P, Karpińska B, Maćkowski S, Van Breusegem F,
Karpiński S (2013) LESION SIMULATING DISEASE1, ENHANCED DISEASE
SUSCEPTIBILITY1, and PHYTOALEXIN DEFICIENT4 conditionally regulate cellular
signaling homeostasis, photosynthesis, water use efficiency, and seed yield in
Arabidopsis. Plant Physiology **161:** 1795
Woodward FI, Lake JA, Quick WP (2002) Stomatal development and CO2: ecological
consequences. New Phytologist **153:** 477-484
Xoconostle-Cazares B, Ramirez-Ortega FA, Flores-Elenes L, Ruiz-Medrano R (2010)
Drought tolerance in crop plants. American Journal of Plant Physiology **5:** 241-256
Yamamoto Y, Sato E, Shimizu T, Nakamich N, Sato S, Kato T, Tabata S, Nagatani A,
Yamashino T, Mizuno T (2003) Comparative genetic studies on the APRR5 and
APRR7 genes belonging to the APRR1/TOC1 quintet implicated in circadian rhythm,
control of flowering time, and early photomorphogenesis. Plant and Cell Physiology
44: 1119-1130
Yang Y, Costa A, Leonhardt N, Siegel RS, Schroeder JI (2008) Isolation of a strong
Arabidopsis guard cell promoter and its potential as a research tool. Plant Methods
4: 6
Yanovsky MJ, Kay SA (2002) Molecular basis of seasonal time measurement in
Arabidopsis. Nature **419:** 308
Yoo CY, Pence HE, Hasegawa PM, Mickelbart MV (2009) Regulation of transpiration to
improve crop water use. Critical Reviews in Plant Sciences **28:** 410-431

Yoo CY, Pence HE, Jin JB, Miura K, Gosney MJ, Hasegawa PM, Mickelbart MV (2010)
The Arabidopsis GTL1 transcription factor regulates water use efficiency and drought
tolerance by modulating stomatal density via transrepression of *SDD1*. The Plant
Cell **22:** 4128

Zagotta MT, Shannon S, Jacobs C, Meeks-Wagner DR (1992) Early-flowering mutants of
Arabidopsis thaliana. Functional Plant Biology **19:** 411-418

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) Strengths and limitations of
period estimation methods for circadian data. PLOS ONE **9:** e96462

Figure 1. The circadian clock regulates long-term water use efficiency of Arabidopsis under light/dark cycles. The WUE of circadian clock mutants and overexpressors is expressed as a percentage of their respective background (normalized to 100%, red reference line) to account for WUE variation between background accessions (n = 5 - 15). Data were analysed using independent-samples t-tests and statistical significance is indicated relative to the background using (* = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$). Statistical analysis was performed on raw data, with data subsequently converted to a percentage of the wild type for the purposes of comparison and presentation. Screens were repeated independently three times per genotype, with one representative experimental repeat shown here for each genotype.

Figure 2. Altered WUE of plants with mutations or overexpression of circadian clock associated genes is not caused consistently by variation in one of dry weight, water use, phase of expression of each gene, or resultant altered period or flowering time. Data are derived from Fig. 1 and expressed as a percentage of the respective background (WT, normalised to 100%, red reference line) (*n* = 5 - 15). (A) Altered WUE is not specifically due to altered water use or altered dry weight of screened genotypes, but results from the combination of both. (b-d) Variations in WUE are not explained by (B) phase expression of mutated/overexpressed gene, nor by altered (C) period or (D) flowering time of the mutated/overexpressor genotype. Genotypes reported to have no change (N/C) in period or flowering time relative to the wild type are included on the left of panels (B) and (C), while

those for which period and/or flowering time are unknown are included on the right. Studies describing the phase of expression, period and flowering time of the genotypes tested are identified in the main text. We note that the phase of expression and period data used for this analysis were often obtained under constant conditions, in contrast to our experiments occurring under light/dark cycles.

Figure 3. The circadian oscillator alters WUE partially by changing rosette architecture. (A) Altering circadian-associated gene expression can affect rosette architecture and size, as illustrated for *elf3*-1, *lux*-1, and *gi*-2 in (Col-0 background). Image backgrounds removed for clarity. Variation in rosette leaf surface area across the genotypes investigated explained (B) 16% of variation in WUE ($p < 0.001$, $r = 0.400$, $r^2 = 0.160$), (C) 83% of variation in transpiration ($p < 0.001$, $r = 0.912$, $r^2 = 0.832$) and (D) 73% of variation in rosette dry biomass ($p < 0.001$, $r = 0.857$, $r^2 = 0.734$). Data were analysed using Pearson correlation tests.

Figure 4. Overexpressing *CCA1* or *TOC1* in guard cells affects WUE and survival of dehydration by seedlings. (A) Constructs used to overexpress *CCA1* or *TOC1* coding sequence under control of *GC1* or *MYB60* promoters. (B) Guard cell *CCA1* overexpression can increase WUE. WUE expressed as a percentage of the wild type (normalised to 100%, red reference line). Two to four independent experimental repeats were performed, with data from one representative dataset shown (*n* = 5 - 15). Data for CCA1-ox and TOC1-ox are derived from Fig. 1, for purposes of comparison. Data were analysed with independent samples t-tests, and statistical significance compared to Col-0 is indicated using starring (** = *p* < 0.01; *** = *p* < 0.001). (C) Guard cell *CCA1* or *TOC1* overexpression alters dehydration survival of seedlings compared with the wild type. Data were obtained from three independent experimental repeats (mean; *n* = 32 per experimental replicate; at least

two independent experimental repeats were performed for each genotype). A single *GC1::TOC1* line is shown here because other lines produced extremely variable data. (D, E) Guard cell *CCA1* or *TOC1* overexpression does not affect (D) stomatal index nor (E) stomatal density. Two independent experimental repeats were performed, with data from one representative dataset shown (n = 19 - 32; mean ± S.E.M.). Data were analysed with ANOVA and Tukey's post hoc tests (NS = p > 0.05). Bar colours identify the whole plant overexpressor control (black), wild type control (dark grey), and guard cell-specific overexpressor genotypes (light grey).

Parsed Citations

Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature 448: 938

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Baudry A, Ito S, Song YH, Strait AA, Kiba T, Lu S, Henriques R, Pruneda-Paz JL, Chua N-H, Tobin EM, Kay SA, Imaizumi T (2010) F-Box proteins FKF1 and LKP2 act in concert with ZEITLUPE to control Arabidopsis clock progression. The Plant Cell 22: 606

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bauer H, Ache P, Lautner S, Fromm J, Hartung W, Al-Rasheid Khaled AS, Sonnewald S, Sonnewald U, Kneitz S, Lachmann N, Mendel Ralf R, Bittner F, Hetherington Alistair M, Hedrich R (2013) The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. Current Biology 23: 53-57

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. Field Crops Research 112: 119-123

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bridge LJ, Franklin KA, Homer ME (2013) Impact of plant shoot architecture on leaf cooling: a coupled heat and mass transfer model. Journal of The Royal Society Interface 10: 20130326

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Caine RS, Yin X, Sloan J, Harrison EL, Mohammed U, Fulton T, Biswal AK, Dionora J, Chater CC, Coe RA, Bandyopadhyay A, Murchie EH, Swarup R, Quick WP, Gray JE (2019) Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. New Phytologist 221: 371-384

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cano-Ramirez DL, Saskia de Fraine T, Griffiths OG, Dodd AN (2018) Photosynthesis and circadian rhythms regulate the buoyancy of marimo lake balls. Current Biology 28: R869-R870

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cominelli E, Galbiati M, Albertini A, Fornara F, Conti L, Coupland G, Tonelli C (2011) DOF-binding sites additively contribute to guard cell-specificity of AtMYB60 promoter. BMC Plant Biology 11: article 162

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cominelli E, Galbiati M, Vavasseur A, Conti L, Sala T, Vuylsteke M, Leonhardt N, Dellaporta SL, Tonelli C (2005) A guard-cell-specific MYB transcription factor regulates stomatal movements and plant drought tolerance. Current Biology 15: 1196-1200

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Condon AG, Farquhar GD, Rebetzke GJ, Richards RA (2004) Breeding for high water-use efficiency. Journal of Experimental Botany 55: 2447-2460

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Costa JM, Monnet F, Jannaud D, Leonhardt N, Ksas B, Reiter IM, Pantin F, Genty B (2015) OPEN ALL NIGHT LONG: The dark side of stomatal control. Plant Physiology 167: 289

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Coupel-Ledru A, Lebon E, Christophe A, Gallo A, Gago P, Pantin F, Doligez A, Simonneau T (2016) Reduced nighttime transpiration is a relevant breeding target for high water-use efficiency in grapevine. Proceedings of the National Academy of Sciences 113: 8963

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Covington MF, Panda S, Liu XL, Strayer CA, Wagner DR, Kay SA (2001) ELF3 modulates resetting of the circadian clock in Arabidopsis. The Plant Cell 13: 1305

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg A, Primavesi LF, Paul MJ, Schluepmann H (2011) Growth arrest by trehalose-6-phosphate: An astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. Plant Physiology 157: 160

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ding Z, Millar AJ, Davis AM, Davis SJ (2007) TIME FOR COFFEE encodes a nuclear regulator in the Arabidopsis thaliana circadian clock. The Plant Cell 19: 1522
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dixon LE, Knox K, Kozma-Bognar L, Southern MM, Pokhilko A, Millar AJ (2011) Temporal repression of core circadian genes is mediated through EARLY FLOWERING 3 in Arabidopsis. Current Biology 21: 120-125
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Dalchau N, Gardner MJ, Baek S-J, Webb AAR (2014) The circadian clock has transient plasticity of period and is required for timing of nocturnal processes in Arabidopsis. New Phytologist 201: 168-179
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Jakobsen MK, Baker AJ, Telzerow A, Hou S-W, Laplaze L, Barrot L, Scott Poethig R, Haseloff J, Webb AAR (2006) Time of day modulates low-temperature Ca2+ signals in Arabidopsis. The Plant Journal 48: 962-973
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Parkinson K, Webb AAR (2004) Independent circadian regulation of assimilation and stomatal conductance in the ztl-1 mutant of Arabidopsis. New Phytologist 162: 63-70
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, Hibberd JM, Millar AJ, Webb AAR (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. Science 309: 630
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Doyle MR, Davis SJ, Bastow RM, McWatters HG, Kozma-Bognár L, Nagy F, Millar AJ, Amasino RM (2002) The ELF4 gene controls circadian rhythms and flowering time in Arabidopsis thaliana. Nature 419: 74
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Edwards CE, Ewers BE, McClung CR, Lou P, Weinig C (2012) Quantitative variation in water-use efficiency across water regimes and its relationship with circadian, vegetative, reproductive, and leaf gas-exchange traits. Molecular Plant 5: 653-668
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Edwards CE, Weinig C (2010) The quantitative-genetic and QTL architecture of trait integration and modularity in Brassica rapa across simulated seasonal settings. Heredity 106: 661
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Endo M, Shimizu H, Nohales MA, Araki T, Kay SA (2014) Tissue-specific clocks in Arabidopsis show asymmetric coupling. Nature 515: 419
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Eriksson ME, Hanano S, Southern MM, Hall A, Millar AJ (2003) Response regulator homologues have complementary, light-dependent functions in the Arabidopsis circadian clock. Planta 218: 159-162
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Farré EM, Harmer SL, Harmon FG, Yanovsky MJ, Kay SA (2005) Overlapping and distinct roles of PRR7 and PRR9 in the Arabidopsis circadian clock. Current Biology 15: 47-54
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ferguson JN, Humphry M, Lawson T, Brendel O, Bechtold U (2018) Natural variation of life-history traits, water use, and drought responses in Arabidopsis. Plant Direct 2: e00035
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill J (1999) GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. The EMBO Journal 18: 4679
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Francia P, Simoni L, Cominelli E, Tonelli C, Galbiati M (2008) Gene trap-based identification of a guard cell promoter in Arabidopsis. Plant Signaling & Behavior 3: 684-686

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D, Yochikawa A, Cano-Ramirez DL, Chembath A, Cragg-Barber K, Haydon MJ, Hotta CT, Vincentz M, Webb AAR, Dodd AN (2018) Circadian entrainment in Arabidopsis by the sugar-responsive transcription factor bZIP63. Current Biology 28: 2597-2606.e2596
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Franks PJ, W. Doheny-Adams T, Britton-Harper ZJ, Gray JE (2015) Increasing water-use efficiency directly through genetic manipulation of stomatal density. New Phytologist 207: 188-195
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Galbiati M, Simoni L, Pavesi G, Cominelli E, Francia P, Vavasseur A, Nelson T, Bevan M, Tonelli C (2008) Gene trap lines identify Arabidopsis genes expressed in stomatal guard cells. The Plant Journal 53: 750-762
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA (2010) AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. The Plant Journal 64: 1-13
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gorton HL, Williams WE, Binns ME, Gemmell CN, Leheny EA, Shepherd AC (1989) Circadian stomatal rhythms in epidermal peels from Vicia faba. Plant Physiology 90: 1329
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Graf A, Schlereth A, Stitt M, Smith AM (2010) Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. Proceedings of the National Academy of Sciences 107: 9458
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hall A, Bastow RM, Davis SJ, Hanano S, McWatters HG, Hibberd V, Doyle MR, Sung S, Halliday KJ, Amasino RM, Millar AJ (2003) The TIME FOR COFFEE gene maintains the amplitude and timing of Arabidopsis circadian clocks. The Plant Cell 15: 2719
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Harmer SL, Hogenesch JB, Straume M, Chang H-S, Han B, Zhu T, Wang X, Kreps JA, Kay SA (2000) Orchestrated transcription of key pathways in Arabidopsis by the circadian clock. Science 290: 2110
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hassidim M, Dakhiya Y, Turjeman A, Hussien D, Shor E, Anidjar A, Goldberg K, Green RM (2017) CIRCADIAN CLOCK ASSOCIATED1 (CCA1) and the circadian control of stomatal aperture. Plant Physiology 175: 1864
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR (2013) Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. Nature 502: 689
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hazen SP, Schultz TF, Pruneda-Paz JL, Borevitz JO, Ecker JR, Kay SA (2005) LUX ARRHYTHMO encodes a Myb domain protein essential for circadian rhythms. Proceedings of the National Academy of Sciences of the United States of America 102: 10387
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM (2000) pGreen: a versatile and flexible binary Ti vector for Agrobacterium-mediated plant transformation. Plant Molecular Biology 42: 819-832
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hennessey TL, Field CB (1991) Circadian rhythms in photosynthesis. Plant Physiology 96: 831
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Herrero E, Kolmos E, Bujdoso N, Yuan Y, Wang M, Berns MC, Uhlworm H, Coupland G, Saini R, Jaskolski M, Webb A, Gonçalves J, Davis SJ (2012) EARLY FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the Arabidopsis circadian clock. The Plant Cell 24: 428
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving environmental change. Nature 424: 901

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. Trends in Plant Science 19: 240-249
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hu H, Xiong L (2014) Genetic engineering and breeding of drought-resistant crops. Annual Review of Plant Biology 65: 715-741
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Huang H, Nusinow DA (2016) Into the evening: Complex interactions in the Arabidopsis circadian clock. Trends in Genetics 32: 674-686
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hugouvieux V, Kwak JM, Schroeder JI (2001) An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in Arabidopsis. Cell 106: 477-487
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA (2005) FKF1 F-Box protein mediates cyclic degradation of a repressor of CONSTANS in Arabidopsis. Science 309: 293
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Imaizumi T, Tran HG, Swartz TE, Briggs WR, Kay SA (2003) FKF1 is essential for photoperiodic-specific light signalling in Arabidopsis. Nature 426: 302
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Joo Y, Fragoso V, Yon F, Baldwin IT, Kim S-G (2017) Circadian clock component, LHY, tells a plant when to respond photosynthetically to light in nature. Journal of Integrative Plant Biology 59: 572-587
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Karaba A, Dixit S, Greco R, Aharoni A, Trijatmiko KR, Marsch-Martinez N, Krishnan A, Nataraja KN, Udayakumar M, Pereira A (2007) Improvement of water use efficiency in rice by expression of HARDY, an Arabidopsis drought and salt tolerance gene. Proceedings of the National Academy of Sciences 104: 15270
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Khanna R, Kikis EA, Quail PH (2003) EARLY FLOWERING 4 functions in Phytochrome B-regulated seedling de-etiolation. Plant Physiology 133: 1530
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kikis EA, Khanna R, Quail PH (2005) ELF4 is a phytochrome-regulated component of a negative-feedback loop involving the central oscillator components CCA1 and LHY. The Plant Journal 44: 300-313
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kim W-Y, Fujiwara S, Suh S-S, Kim J, Kim Y, Han L, David K, Putterill J, Nam HG, Somers DE (2007) ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. Nature 449: 356
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kinoshita T, Hayashi Y (2011) New insights into the regulation of stomatal opening by blue light and plasma membrane H+-ATPase. In KW Jeon, ed, International Review of Cell and Molecular Biology, Vol 289. Academic Press, pp 89-115
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kinoshita T, Ono N, Hayashi Y, Morimoto S, Nakamura S, Soda M, Kato Y, Ohnishi M, Nakano T, Inoue S-i, Shimazaki K-i (2011) FLOWERING LOCUS T regulates stomatal opening. Current Biology 21: 1232-1238
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lawson T, Blatt MR (2014) Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. Plant Physiology 164: 1556
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Legnaioli T, Cuevas J, Mas P (2009) TOC1 functions as a molecular switch connecting the circadian clock with plant responses to drought. The EMBO Journal 28: 3745
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Litthauer S, Battle MW, Lawson T, Jones MA (2015) Phototropins maintain robust circadian oscillation of PSII operating efficiency under blue light. The Plant Journal 83: 1034-1045

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Más P, Kim W-Y, Somers DE, Kay SA (2003) Targeted degradation of TOC1 by ZTL modulates circadian function in Arabidopsis thaliana. Nature 426: 567

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Masle J, Gilmore SR, Farquhar GD (2005) The ERECTA gene regulates plant transpiration efficiency in Arabidopsis. Nature 436: 866

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Medrano H, Tomás M, Martorell S, Flexas J, Hernández E, Rosselló J, Pou A, Escalona J-M, Bota J (2015) From leaf to whole-plant water use efficiency (WUE) in complex canopies: Limitations of leaf WUE as a selection target. The Crop Journal 3: 220-228

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Meyer S, Mumm P, Imes D, Endler A, Weder B, Al-Rasheid KAS, Geiger D, Marten I, Martinoia E, Hedrich R (2010) AtALMT12 represents an R-type anion channel required for stomatal movement in Arabidopsis guard cells. The Plant Journal 63: 1054-1062

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Michael TP, Salomé PA, Yu HJ, Spencer TR, Sharp EL, McPeek MA, Alonso JM, Ecker JR, McClung CR (2003) Enhanced fitness conferred by naturally occurring variation in the circadian clock. Science 302: 1049

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Millar AJ (2004) Input signals to the plant circadian clock. Journal of Experimental Botany 55: 277-283

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Millar AJ, Carre IA, Strayer CA, Chua NH, Kay SA (1995) Circadian clock mutants in Arabidopsis identified by luciferase imaging. Science 267: 1161

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Murakami M, Yamashino T, Mizuno T (2004) Characterization of circadian-associated APRR3 pseudo-response regulator belonging to the APRR1/TOC1 quintet in Arabidopsis thaliana. Plant and Cell Physiology 45: 645-650

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Na J-K, Metzger JD (2014) Chimeric promoter mediates guard cell-specific gene expression in tobacco under water deficit. Biotechnology Letters 36: 1893-1899

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nagy R, Grob H, Weder B, Green P, Klein M, Frelet A, Schjoerring JK, Brearley CA, Martinoia E (2009) The Arabidopsis ATP-binding cassette protein ATMRP5/ATABCC5 is a high-affinity inositol hexakisphosphate transporter involved in guard cell signaling and phytate storage. Journal of Biological Chemistry

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010) PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in Arabidopsis circadian clock. The Plant Cell 22: 594

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nakamichi N, Murakami-Kojima M, Sato E, Kishi Y, Yamashino T, Mizuno T (2002) Compilation and characterization of a novel WNK family of protein kinases in Arabidopsis thaliana with reference to circadian rhythms. Bioscience, Biotechnology, and Biochemistry 66: 2429-2436

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nienhuis J, Sills GR, Martin B, King G (1994) Variance for water-use efficiency among ecotypes and recombinant inbred lines of Arabidopsis thaliana (Brassicaceae). American Journal of Botany 81: 943-947

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nietzsche M, Guerra T, Alseekh S, Wiermer M, Sonnewald S, Fernie AR, Börnke F (2018) STOREKEEPER RELATED1/G-element binding protein (STKR1) interacts with protein kinase SnRK1. Plant Physiology 176: 1773

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma M, Tanaka K, Hanaoka M, Dodd AN (2013) Circadian control of chloroplast transcription by a nuclear-encoded timing signal. Science 339: 1316

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Panda S, Poirier GG, Kay SA (2002) tej defines a role for poly(ADP-ribosyl)ation in establishing period length of the Arabidopsis circadian oscillator. Developmental Cell 3: 51-61

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Paul MJ, Jhurreea D, Zhang Y, Primavesi LF, Delatte T, Schluepmann H, Wingler A (2010) Up-regulation of biosynthetic processes associated with growth by trehalose 6-phosphate. Plant Signaling & Behavior 5: 386-392

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pei Z-M, Ghassemian M, Kwak CM, McCourt P, Schroeder JI (1998) Role of farnesyltransferase in ABA regulation of guard cell anion channels and plant water loss. Science 282: 287

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. Science 323: 1481

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, Salemi MR, Phinney BS, Harmer SL (2011) REVEILLE8 and PSEUDO-REPONSE REGULATOR5 form a negative feedback loop within the Arabidopsis circadian clock. PLOS Genetics 7: e1001350

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Resco de Dios V, Gessler A, Pedro Ferrio J, Alday JG, Bahn M, del Castillo J, Devidal S, García-Muñoz S, Kayler Z, Landais D, Martín-Gómez P, Milcu A, Piel C, Pirhofer-Walzl K, Ravel O, Salekin S, Tissue DT, Tjoelker MG, Voltas J, Roy J (2016) Circadian rhythms have significant effects on leaf-to-canopy scale gas exchange under field conditions. GigaScience 5

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rubin MJ, Brock MT, Baker RL, Wilcox S, Anderson K, Davis SJ, Weinig C (2018) Circadian rhythms are associated with shoot architecture in natural settings. New Phytologist 219: 246-258

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ruggiero A, Punzo P, Landi S, Costa A, Van Oosten JM, Grillo S (2017) Improving plant water use efficiency through molecular genetics. Horticulturae 3

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rusconi F, Simoni L, Simeoni F, Tonelli C, Galbiati M, Cominelli E, Conti L, Riboni M, Francia P, Martin CR (2013) The Arabidopsis thaliana MYB60 promoter provides a tool for the spatio-temporal control of gene expression in stomatal guard cells. Journal of Experimental Botany 64: 3361-3371

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ruts T, Matsubara S, Wiese-Klinkenberg A, Walter A (2012) Aberrant temporal growth pattern and morphology of root and shoot caused by a defective circadian clock in Arabidopsis thaliana. The Plant Journal 72: 154-161

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Salomé PA, McClung CR (2005) PSEUDO-RESPONSE REGULATOR 7 and 9 are partially redundant genes essential for the temperature responsiveness of the Arabidopsis circadian clock. The Plant Cell 17: 791

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schaffer R, Ramsay N, Samach A, Corden S, Putterill J, Carré IA, Coupland G (1998) The late elongated hypocotyl mutation of Arabidopsis disrupts circadian rhythms and the photoperiodic control of flowering. Cell 93: 1219-1229

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) Guard cell signal transduction. Annual Review of Plant Physiology and Plant Molecular Biology 52: 627-658

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schultz TF, Kiyosue T, Yanovsky M, Wada M, Kay SA (2001) A Role for LKP2 in the Circadian Clock of Arabidopsis. The Plant Cell 13: 2659

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. Plant, Cell & Environment 40: 997-1008

Simon NML, Kusakina J, Fernández-López Á, Chembath A, Belbin FE, Dodd AN (2018) The energy-signaling hub SnRK1 is important for sucrose-induced hypocotyl elongation. Plant Physiology 176: 1299

Simon NML, Sawkins E, Dodd AN (2018) Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation. Plant Signaling & Behavior 13: e1457913

Somers DE, Devlin PF, Kay SA (1998) Phytochromes and cryptochromes in the entrainment of the Arabidopsis circadian clock. Science 282: 1488

Somers DE, Schultz TF, Milnamow M, Kay SA (2000) ZEITLUPE encodes a novel clock-associated PAS protein from Arabidopsis. Cell 101: 319-329

Somers DE, Webb AA, Pearson M, Kay SA (1998) The short-period mutant, toc1-1, alters circadian clock regulation of multiple outputs throughout development in Arabidopsis thaliana. Development 125: 485

Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Más P, Panda S, Kreps JA, Kay SA (2000) Cloning of the Arabidopsis clock gene TOC1, an autoregulatory response regulator homolog. Science 289: 768

Streitner C, Danisman S, Wehrle F, Schöning JC, Alfano JR, Staiger D (2008) The small glycine-rich RNA binding protein AtGRP7 promotes floral transition in Arabidopsis thaliana. The Plant Journal 56: 239-250

Thines B, Harmon FG (2010) Ambient temperature response establishes ELF3 as a required component of the core Arabidopsis circadian clock. Proceedings of the National Academy of Sciences: 200911006

Tomás M, Medrano H, Escalona JM, Martorell S, Pou A, Ribas-Carbó M, Flexas J (2014) Variability of water use efficiency in grapevines. Environmental and Experimental Botany 103: 148-157

Vialet-Chabrand S, Matthews JSA, Brendel O, Blatt MR, Wang Y, Hills A, Griffiths H, Rogers S, Lawson T (2016) Modelling water use efficiency in a dynamic environment: An example using Arabidopsis thaliana. Plant Science 251: 65-74

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis thaliana. Science 339: 704

Wang Y, Liu K, Liao H, Zhuang C, Ma H, Yan X (2008) The plant WNK gene family and regulation of flowering time in Arabidopsis. Plant Biology 10: 548-562

Wang ZY, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. Cell 93: 1207-1217

Wituszyńska W, Ślesak I, Vanderauwera S, Szechyńska-Hebda M, Kornaś A, Van Der Kelen K, Mühlenbock P, Karpińska B, Maćkowski S, Van Breusegem F, Karpiński S (2013) LESION SIMULATING DISEASE1, ENHANCED DISEASE SUSCEPTIBILITY1, and

PHYTOALEXIN DEFICIENT4 conditionally regulate cellular signaling homeostasis, photosynthesis, water use efficiency, and seed yield in Arabidopsis. Plant Physiology 161: 1795

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Woodward FI, Lake JA, Quick WP (2002) Stomatal development and CO2: ecological consequences. New Phytologist 153: 477-484

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Xoconostle-Cazares B, Ramirez-Ortega FA, Flores-Elenes L, Ruiz-Medrano R (2010) Drought tolerance in crop plants. American Journal of Plant Physiology 5: 241-256

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yamamoto Y, Sato E, Shimizu T, Nakamich N, Sato S, Kato T, Tabata S, Nagatani A, Yamashino T, Mizuno T (2003) Comparative genetic studies on the APRR5 and APRR7 genes belonging to the APRR1/TOC1 quintet implicated in circadian rhythm, control of flowering time, and early photomorphogenesis. Plant and Cell Physiology 44: 1119-1130

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yang Y, Costa A, Leonhardt N, Siegel RS, Schroeder JI (2008) Isolation of a strong Arabidopsis guard cell promoter and its potential as a research tool. Plant Methods 4: 6

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yanovsky MJ, Kay SA (2002) Molecular basis of seasonal time measurement in Arabidopsis. Nature 419: 308

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoo CY, Pence HE, Hasegawa PM, Mickelbart MV (2009) Regulation of transpiration to improve crop water use. Critical Reviews in Plant Sciences 28: 410-431

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoo CY, Pence HE, Jin JB, Miura K, Gosney MJ, Hasegawa PM, Mickelbart MV (2010) The Arabidopsis GTL1 transcription factor regulates water use efficiency and drought tolerance by modulating stomatal density via transrepression of SDD1. The Plant Cell 22: 4128

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zagotta MT, Shannon S, Jacobs C, Meeks-Wagner DR (1992) Early-flowering mutants of Arabidopsis thaliana. Functional Plant Biology 19: 411-418

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) Strengths and limitations of period estimation methods for circadian data. PLOS ONE 9: e96462

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title