The circadian clock contributes to the long-term water use efficiency of Arabidopsis

Noriane M. L. Simon[1], Calum A. Graham[1,2], Nicholas E. Comben[1], Alistair M. Hetherington[1],

Antony N. Dodd[1,2]*

1. School of Biological Sciences, University of Bristol, Life Sciences Building, Bristol BS8

1TQ, U.K.

2. John Innes Centre, Norwich NR4 7UH, U.K.

* corresponding author, antony.dodd@jic.ac.uk

Word count (body text including methods): 6543

Short title: Circadian regulation and water use efficiency

Author contributions: N.M.L.S., A.M.H. and A.N.D. conceived experiments, N.M.L.S. and

N.E.C. performed experiments, N.M.L.S., C.A.G., A.M.H. and A.N.D. analysed and

interpreted the data, N.M.L.S., C.A.G., A.M.H. and A.N.D. wrote the paper. A.N.D. agrees to

serve as the author responsible for contact and ensures communication.

One-sentence summary: The circadian clock in Arabidopsis makes an important

contribution to long-term water use efficiency.

Abstract

In plants, water use efficiency is a complex trait derived from numerous physiological and

developmental characteristics. Here, we investigated the involvement of circadian regulation

in long-term water use efficiency. Circadian rhythms are generated by the circadian

oscillator, which provides a cellular measure of the time of day. In plants, the circadian

oscillator contributes to the regulation of many aspects of physiology, including stomatal

opening, the rate of photosynthesis, carbohydrate metabolism and developmental

processes. We investigated in Arabidopsis the impact of the misregulation of genes

encoding a large number of components of the circadian oscillator upon whole plant, long-

term water use efficiency. From this, we identified a role for the circadian oscillator in water

use efficiency. This appears to be due to contributions of the circadian clock to the control of

transpiration and biomass accumulation. We also identified that the circadian oscillator

within guard cells can contribute to long-term water use efficiency. Our experiments indicate

that knowledge of circadian regulation will be important for developing future crops that use

water more efficiently.

Introduction

World population growth is increasing the demand for fresh water for agriculture, with

climate change predicted to exacerbate this competition for water resources (Ruggiero et al.,

2017). One strategy to sustainably increase agricultural production involves the

improvement of crop water use (Condon et al., 2004; Xoconostle-Cazares et al., 2010; Hu

and Xiong, 2014; Ruggiero et al., 2017). Because up to 97% of water taken up from the soil

by plants is lost through stomatal transpiration (Yoo et al., 2009; Na and Metzger, 2014), the

manipulation of transpiration represents an excellent candidate for designing crops with

increased water use efficiency (Bertolino et al., 2019).

Plant water loss can be manipulated through changes in the regulation of stomatal opening

and by altering stomatal density and patterning (Pei et al., 1998; Hugouvieux et al., 2001;

Schroeder et al., 2001; Hetherington and Woodward, 2003; Yoo et al., 2010; Lawson and

Blatt, 2014; Franks et al., 2015; Caine et al., 2019). In addition to stomatal responses to

environmental cues such as light, temperature and phytohormones, there are circadian

rhythms of stomatal opening (Gorton et al., 1989; Hennessey and Field, 1991). Circadian

rhythms are self-sustaining biological cycles with a period of about 24 h. These rhythms are

thought to adapt plants to daily cycles of light and dark, by anticipating daily changes in the

environment and co-ordinating cellular processes. In higher plants, circadian rhythms are

generated by several interlocked transcription-translation feedback loops known as the

circadian oscillator (Hsu and Harmer, 2014). The phase of the circadian oscillator is adjusted

continuously to match the phase of the environment through the process of entrainment, in

response to light, temperature and metabolic cues (Somers et al., 1998; Millar, 2004;

Salomé and McClung, 2005; Haydon et al., 2013; Webb et al., 2019). Additionally, the

circadian oscillator communicates an estimate of the time of day to circadian-regulated

features of the cell, initially through transcriptional regulation (Harmer et al., 2000). The

known circadian oscillator controls circadian rhythms of stomatal opening because mutations

that alter the circadian period or cause circadian arrhythmia lead to equivalent alterations in

the circadian rhythm of stomatal opening (Somers et al., 1998; Dodd et al., 2004; Dodd et

al., 2005). The circadian oscillator is also involved in the responses of guard cells to

environmental cues such as drought and low temperature (Dodd et al., 2006; Legnaioli et al.,

2009).

Circadian rhythms are often studied under conditions of constant light. However, the

circadian oscillator is also important for the regulation of stomatal opening under cycles of

light and dark. For example, constitutive overexpression of the circadian oscillator

component CIRCADIAN CLOCK ASSOCIATED1 (CCA1; CCA1-ox) alters the daily

regulation of stomatal opening such that stomatal conductance increases steadily

throughout the photoperiod (Dodd et al., 2005). In comparison, stomatal conductance in wild

type plants remains relatively uniform during the photoperiod and is lower than in CCA1-ox

(Dodd et al., 2005). Similarly, guard cell-specific overexpression of CCA1 generally causes

greater stomatal opening during the light period, and alters drought response phenotypes

(Hassidim et al., 2017). Modelling suggests that under light/dark cycles, the circadian

oscillator contributes at the canopy scale to daily rhythms in stomatal aperture and carbon

assimilation in bean and cotton (Resco de Dios et al., 2016).

The contribution of the circadian oscillator to both stomatal opening and biomass

accumulation (Dodd et al., 2005; Graf et al., 2010) suggests that the circadian oscillator

might make an important contribution to long-term water use efficiency (WUE). WUE is the

ratio of carbon dioxide incorporated through photosynthesis into biomass to the amount of

water lost through transpiration. At the single leaf level, instantaneous, intrinsic WUE is often

measured with gas exchange techniques and expressed as net CO_2 assimilation per unit of

water transpired (Vialet-Chabrand et al., 2016; Ruggiero et al., 2017; Ferguson et al., 2018).

However, such measurements do not provide an accurate representation of WUE over the

plant lifetime, which is influenced by features such as leaf position, dark respiration, and time

of day changes in instantaneous WUE (Condon et al., 2004; Tomás et al., 2014; Medrano et

al., 2015; Ferguson et al., 2018). It is important to note that high WUE under well-watered

conditions is not the same as drought resistance, because drought resistance relates to the

capacity to maintain transpirational water supply under water-limited conditions through

strategies such as expanded root systems (Blum, 2009). This means that WUE does not

correlate reliably with drought resistance (Kobata et al., 1996).

Given that the circadian oscillator affects stomatal opening and biomass accumulation

(Gorton et al., 1989; Hennessey and Field, 1991; Dodd et al., 2005; Edwards and Weinig,

2010; Graf et al., 2010; Edwards et al., 2012), we hypothesized that specific components of

the circadian oscillator might make an important contribution to the long-term WUE of plants.

Although the circadian oscillator influences stomatal opening and biomass accumulation, the

influence of the circadian oscillator upon long-term WUE of plants remains unknown. This is

an important question for understanding roles for circadian regulation in crops, because the

long-term water use efficiency ultimately determines the amount of water that is required for

a given yield of the crop. Therefore, we investigated the impact of the misregulation of parts

of the circadian oscillator upon the long-term WUE of Arabidopsis. We identified that the

circadian oscillator has profound effects upon the long-term WUE of plants. Importantly,

some alterations in oscillator function increase long-term WUE, suggesting potential targets

for future improvements of crop WUE.

Results

Circadian oscillator components contribute to long-term water use efficiency

We identified that correct regulation of genes encoding circadian oscillator components

makes an important contribution to WUE. 32 single mutants or overexpressors of genes

associated with circadian regulation, representing 21 circadian oscillator-associated

components, were surveyed for WUE alterations (Fig. 1) using a previously-described

method (Wituszyńska et al., 2013). 44 % of the mutants or overexpressors examined had a

significantly different WUE from the wild type (Fig. 1, Table 1). The *elf3*-1, *prr9*-1, *tps1*-11,

tps1-12, and *ztl*-1 mutants, as well as CCA1, TOC1 and KIN10 overexpressors, had

significantly lower WUE than the wild type (Fig. 1, Table 1; Table S1). The *gi*-2, *gi*-11, *grp7*-

1, *tej*-1 and *tic*-2 mutants had significantly greater WUE than the wild type (Fig. 1, Table 1;

Table S1). This suggests that misregulating the expression of circadian clock components

CCA1, ELF3, GI, GRP7, PRR9, TEJ, TIC, TOC1 and ZTL can change whole plant long-term

WUE (Fig. 1, Table 1). WUE was also altered by changing the expression of the energy

signalling pathway components TPS1 and KIN10 that also provide inputs to the circadian

oscillator (Shin et al., 2017; Frank et al., 2018) (Fig. 1, Table 1). Overall, these data show

that correct expression of a variety of circadian clock-associated genes contributes to long-

term WUE of Arabidopsis.

Within this experiment, each background accession had a distinct WUE (C24: 3.01 ± 0.07

mg ml^{-1}; Col-0: 2.22 ± 0.02 mg ml^{-1}; L. *er.*: 1.60 ± 0.04 mg ml^{-1}; Ws: 1.91 ± 0.06 mg ml^{-1})

(Fig. S2). These differences between WUE of different Arabidopsis accessions are

consistent with previous studies of WUE, stomatal function and stomatal density in

Arabidopsis (Nienhuis et al., 1994; Woodward et al., 2002; Dodd et al., 2004; Masle et al.,

2005; Karaba et al., 2007; Ruggiero et al., 2017; Ferguson et al., 2018).

We hypothesised that variations in WUE might be associated with specific circadian

phenotypes in the mutants and overexpressors that we tested. For example, mutations in

circadian clock genes expressed with a particular phase (e.g. morning-expressed or

evening-expressed genes) might have a more pronounced effect on WUE. Likewise, the

nature of the circadian period change or flowering time change resulting from misexpression

of each oscillator component might be associated with certain changes in WUE. To test this,

we compared the data from our WUE screen with the circadian phase of expression of each

mutated or overexpressed gene. We also compared the direction of change of WUE to the

period and flowering time phenotypes that arise from each mutant or overexpressor (Hicks

et al., 1996; Fowler et al., 1999; Schultz et al., 2001; Doyle et al., 2002; Nakamichi et al.,

2002; Yanovsky and Kay, 2002; Imaizumi et al., 2003; Más et al., 2003; Murakami et al.,

2004; Farré et al., 2005; Hazen et al., 2005; Baena-González et al., 2007; Ding et al., 2007;

Niwa et al., 2007; Streitner et al., 2008; Wang et al., 2008; Baudry et al., 2010; Nakamichi et

al., 2010; Rawat et al., 2011; Wahl et al., 2013; Hsu and Harmer, 2014). We note that the

phenotypes reported by these studies were often identified under constant conditions, with

flowering time experiments performed under short or long photoperiods, whereas our

experiments occurred under cycles of 8 h light / 16 h darkness.

There was no obvious relationship between the circadian phenotypes reported to arise from

each mutant or overexpressor investigated and the WUE of each of these lines (Fig. 2A, B,

C). For example, mutating night-phased oscillator components can either decrease or

increase WUE (Fig. 2A). Mutants that cause long circadian periods and short circadian

periods can both increase and decrease WUE, although WUE was unaltered in the mutants

that are reported to not alter the period (Fig. 2B). Furthermore, mutants and overexpressors

that cause both early and delayed flowering can each increase and decrease WUE (Fig.

2C).

We were interested to determine whether the WUE alterations caused by misregulation of

circadian oscillator gene expression arose from changes in either biomass accumulation or

transpiration. Genotypes with low biomass accumulation generally had lower water use (Fig.

3A, B). The results for *tej*-1 (Fig. 3A, B) should be treated with caution because the model

fits were very poor (Table S1). No mutants or overexpressors tested increased the biomass

relative to the corresponding background genotype (Fig 3A). Together, this indicates that the

altered WUE phenotypes in some genotypes with misregulated circadian clocks was not due

to an alteration in just one of either water use or biomass accumulation (Fig. 3A, B). Instead,

the altered WUE of lines with misregulated circadian clock genes appears to be due to the

net effect of altered biomass accumulation and altered transpiration in these genotypes (Fig.

3A, B).

Circadian regulation of water use efficiency combines multiple traits

Mutation or overexpression of components of the circadian oscillator can cause changes in

the development of Arabidopsis, such as alterations in rosette size, leaf shape and petiole

length (Fig. 4A) (Zagotta et al., 1992; Schaffer et al., 1998; Wang and Tobin, 1998; Dodd et

al., 2005; Ruts et al., 2012; Rubin et al., 2018). These changes are likely to have

implications for gas exchange because, for example, spatially separated leaves are

predicted to transpire more water (Bridge et al., 2013). We investigated whether the changes

in WUE that were identified by our screen might arise from differences in rosette architecture

between the circadian clock-associated mutants and overexpressors and the corresponding

backgrounds. There was a weak positive correlation between rosette leaf surface area and

WUE ($r = 0.400$; $r^2 = 0.160$; $p < 0.001$) (Fig. 4B). Therefore, this suggests that approximately

16% of variability in WUE can be explained by the variations in rosette leaf surface area that

arise from misregulation of the circadian oscillator.

In comparison, rosette leaf surface area was strongly correlated with each of the individual

parameters of water used and dry biomass accumulated. The variation in rosette surface

area accounted for 83% of the variability in water transpired across the genotypes (Fig. 4C).

Furthermore, the variation in rosette surface area accounted for 73% of the variability in

biomass accumulation across the genotypes (Fig. 4D), which is unsurprising given that

larger leaves are likely to contain more biomass.

This demonstrates that one way that circadian regulation affects WUE is through the

influence of the circadian oscillator upon plant development and rosette architecture, but this

variation in leaf area does not account for the majority of the influence of circadian regulation

upon WUE. It also further supports the notion that the influence of the circadian oscillator

upon WUE is complex, and cannot be explained by variation in one of water use or biomass

accumulation alone.

Contribution of circadian regulation in guard cells to water use efficiency

Next, we investigated whether the circadian oscillator within guard cells contributes to long-

term WUE. There is evidence that guard cells contain a circadian oscillator that regulates

stomatal opening (Gorton et al., 1989; Hassidim et al., 2017). To investigate the contribution

of the guard cell circadian oscillator to WUE, we overexpressed two circadian oscillator

components (*CCA1, TOC1*) in guard cells, using two guard cell-specific promoters (*GC1,*

MYB60) for each of *CCA1* and *TOC1* (Fig. 5A) (Cominelli et al., 2005; Galbiati et al., 2008;

Yang et al., 2008; Nagy et al., 2009; Meyer et al., 2010; Cominelli et al., 2011; Bauer et al.,

2013; Rusconi et al., 2013). *GC1* is a guard cell-specific promoter that is relatively

unresponsive to a variety of environmental cues (cold, light, ABA, gibberellin) (Yang et al.

2008). We used the full-length *MYB60* promoter sequence, because truncated and chimeric

versions of this promoter appear to have weaker activity and/or become rapidly

downregulated by dehydration and ABA (Francia et al., 2008; Cominelli et al., 2011; Rusconi

et al., 2013). This produced four sets of transgenic lines; *GC1::CCA1:nos* (*GC*),

GC1::TOC1:nos (*GT*), *MYB60::CCA1:nos* (*MC*) and *MYB60::TOC1:nos* (*MT*) (Fig. 5A). We

termed these guard cell specific (GCS) plants. We confirmed the guard cell specificity of the

GC1 and *MYB60* promoters in our hands, by driving green fluorescent protein (GFP) under

the control of these promoters. GFP accumulation was restricted to the guard cells (Fig.

S3A, B). There was not a circadian oscillation in the activity of either the *GC1* or *MYB60*

promoter under our experimental conditions (Fig. S3C), demonstrating that these promoters

were appropriate for constitutive overexpression of circadian oscillator components within

guard cells in our experiments.

To further verify the guard cell-specific overexpression of *CCA1* and *TOC1* in the GCS

plants, we examined *CCA1* and *TOC1* transcript accumulation within guard cells. Under

constant light conditions, we measured *CCA1* transcript accumulation in epidermal peels at

dusk (when *CCA1* transcript abundance is normally low in the wild type) and *TOC1*

transcript accumulation at dawn (when *TOC1* transcript abundance is normally low in the

wild type). Guard cell *CCA1* overexpressors had greater *CCA1* transcript abundance in

epidermal peels at dusk than the wild type (Fig. S3D), and guard cell *TOC1* overexpressors

had greater *TOC1* transcript abundance at dawn than the wild type (Fig. S3D). These data

indicate that *CCA1* and *TOC1* were overexpressed within the guard cells of the guard cell-

specific *CCA1* or *TOC1* overexpressor plants that we generated, respectively.

We investigated the effect on WUE of overexpression of CCA1 and TOC1 within guard cells.

Two independent *GC1::CCA1* lines (*GC*-1 and *GC*-2) were significantly more water use

efficient than the wild type (*GC*-1: $p < 0.001$; *GC*-2: $p = 0.002$) (Fig. 5B). *GC*-1 and *GC*-2

were 8% and 4% more water use efficient than the wild type, respectively (Fig. 5B). In

comparison, two independent *MYB60::CCA1* did not have greater WUE than the wild type

($p > 0.05$) (Fig. 5B). This suggests that overexpressing *CCA1* in guard cells can increase

whole plant long-term WUE in a promoter-specific manner. Overexpression of *TOC1* in

guard cells with both the *GC1* and *MYB60* promoters did not alter WUE ($p > 0.05$) (Fig. 5B).

This suggests that decreased WUE in constitutive TOC1-ox plants (Fig. 1, Fig. 5B) might not

be explained by overexpression of TOC1 within the guard cells, and that this decreased

WUE might instead be due to TOC1 overexpression in other cell types. In this particular

experiment we did not see an alteration in CCA1-ox WUE relative to its background,

contrasting Fig. 1. This could suggest that the WUE alteration in CCA1-ox is fairly small or

variable. Because the stomatal density was unaltered relative to the wild type in the guard

cell overexpressors of *CCA1* and *TOC1* (Fig. 5C, D), the WUE phenotypes that we identified

from these lines might be caused by alterations in processes within guard cells, such as

those regulating stomatal aperture, rather than altered stomatal density.

Discussion

Pervasive influence of the circadian oscillator upon water use efficiency

Our data indicate that the circadian oscillator is important for regulating the long-term WUE

of Arabidopsis. Misregulation of several functional subsections of the circadian oscillator

altered the WUE. Misexpression of morning (PRR9, CCA1), late day (GI) and evening

(TOC1, ZTL, ELF3) components of the circadian oscillator all perturb WUE under our

experimental conditions (Fig. 1). Additionally, mutation of TEJ and GRP7 alters WUE (Fig.

1). Therefore, oscillator components that impact WUE are not confined to a specific

expression phase or architectural feature (e.g. morning loop) within the multi-loop circadian

oscillator. Misexpression of genes encoding some proteins that provide environmental inputs

to the circadian oscillator (ELF3, TPS1, ZTL, KIN10; (Covington et al., 2001; Kim et al.,

2007; Shin et al., 2017; Frank et al., 2018)) also alters WUE (Fig. 1). Together, this suggests

that the entire circadian oscillator can influence WUE, and that alterations in water use that

are caused by mutations to the circadian oscillator are not confined to a specific sub-loop of

the circadian oscillator, or restricted to its input or output pathways. One explanation for

these circadian-system wide alterations in WUE relates to the nature of feedback within the

circadian oscillator. The complex feedback and interconnectivity of the circadian oscillator

means that individual components of the circadian oscillator that directly influence stomatal

function or water use are likely to be altered by mutations that are distal to that component.

Therefore, if correct circadian timing is required for optimum water use efficiency, multiple

components of the circadian oscillator are likely to influence water use efficiency.

Alternatively, because mutation of a number of components of the circadian oscillator had no

effect upon WUE, it is possible that the oscillator components that influence WUE do so

through roles in directly regulating outputs of the circadian oscillator, such as by regulating

genes involved in stomatal function.

The sugar signalling proteins TPS1 and KIN10 influence a broad range of phenotypes, in

addition to participating in circadian entrainment (Baena-González et al., 2007; Gómez et al.,

2010; Paul et al., 2010; Delatte et al., 2011; Shin et al., 2017; Frank et al., 2018; Nietzsche

et al., 2018; Simon et al., 2018). The *tps1*-12 TILLING mutant of TPS1 decreases stomatal

aperture and increases the ABA sensitivity of guard cells (Gómez et al., 2010), whereas we

found that *tps1*-11 and *tps1*-12 had lower long-term WUE than the wild type (Fig. 1). Lower

biomass accumulation in *tps1*-11 and *tps1*-12 (Fig. 3A) was consistent with slow growth of

these alleles (Gómez et al., 2010). Overall, this suggests that the decreased stomatal

aperture of *tps1*-12 mutants (Gómez et al., 2010) does not translate into an overall increase

in WUE, perhaps due to slower growth or misregulated ABA signalling in the *tps* mutants

(Gómez et al., 2010). The broad range of phenotypes that are altered in *tps1*-11, *tps1*-12

and KIN10-ox indicates that these genotypes might alter WUE through mechanisms other

than circadian regulation.

Potential roles for the evening complex in WUE

Our finding that ELF3 can influence WUE (Fig. 1) is supported by previous evidence. Under

constant light conditions, wild type Arabidopsis has circadian rhythms of stomatal aperture,

whereas *elf3* stomata are constantly open and unresponsive to light and dark (Kinoshita et

al., 2011). Furthermore, ELF3 negatively regulates blue light-mediated stomatal opening

(Kinoshita and Hayashi, 2011). Therefore, perturbation of the anticipation of day/night

transitions or responses to environmental cues in *elf3* stomata might cause long-term

alterations in WUE.

ELF3 binds to the *PRR9* promoter and *elf3*-1 has elevated PRR9 transcript abundance

(Thines and Harmon, 2010; Dixon et al., 2011; Herrero et al., 2012). The low WUE of *elf3*-1

might potentially be caused by altered PRR9 expression, because misregulation of *PRR9*

also affected WUE (Fig. 1). In a similar fashion, ELF3/ELF4 signalling represses PRR7, and

elf3-1 has elevated *PRR7* transcript abundance (Herrero et al., 2012). Under light-dark

cycles, *elf3*-1 also has high and constitutive *GI* expression (Fowler et al., 1999), and *elf3*-1

and *gi* mutants have opposite WUE phenotypes (Fig. 1). Therefore, the WUE phenotype of

elf3-1 (Fig. 1) might be caused by disruption of ELF3 itself, or specific perturbations of

PRR7, PRR9 and/or GI expression.

Mutating further components of the evening complex (EC) (ELF4 and LUX) did not affect

WUE (Fig. 1). This is despite these genes influencing circadian oscillator function and plant

physiology (Hsu and Harmer, 2014; Huang and Nusinow, 2016), and nocturnal regulation of

stomatal aperture altering WUE (Costa et al., 2015; Coupel-Ledru et al., 2016). One

possibility is that the impact of *elf3*-1 on WUE may be greater than that of *elf4* or *lux*

because ELF3 is key to EC scaffolding, with ELF3 operating genetically downstream from

ELF4 and LUX (Herrero et al., 2012; Huang and Nusinow, 2016).

The EC binds upstream of and regulates a variety of other genes that might also underlie the

WUE alterations in *elf3*-1 mutants (Ezer et al., 2017). This includes regulators of growth,

components of the photosynthetic apparatus, and genes associated with phytohormone

signalling. This means that potential roles for the EC in WUE might occur through several

physiological mechanisms. There also appears to be a negative relationship between

temperature and EC promoter binding (Ezer et al., 2017), so it is possible that any influence

of the EC upon WUE might be temperature-sensitive.

ELF4 appears to play a greater role in circadian regulation in the vascular tissue than

stomatal guard cells, with vasculature expression up to ten times higher than other tissues

(Endo et al., 2014). Processes within the vasculature can affect WUE; for example,

mutations in *CELLULOSE SYNTHASE CATALYTIC SUBUNIT7* (*CESA7*) might impact

water use through effects of the collapse of the vasculature upon guard cell size (Liang et

al., 2010). Because *elf3*-1 affects WUE differently from *elf4*-101 and *lux*-1 (Fig. 1), ELF3

might regulate WUE independently from ELF4 and LUX.

Multiple physiological causes of altered WUE in circadian oscillator mutants

Our data suggest that changes in WUE caused by misexpression of circadian clock

components might be due to a combination of physiological factors. Many mutants or

overexpressors tested alter both biomass accumulation and water loss, often in the same

direction (Fig. 3A, B), so mutations to the circadian oscillator did not alter water use by

specifically altering either carbon assimilation or transpiration. This is consistent with

previous work demonstrating that both stomatal opening and CO_2 fixation is perturbed in

circadian arrhythmic plants under light/dark cycles (Dodd et al., 2005), and also with the

finding that daily carbohydrate management is dependent upon correct circadian regulation

(Graf et al., 2010). We speculate that delayed or advanced stomatal and photosynthetic

responses to the day-night cycle might occur in circadian period mutants, because period

mutants inaccurately anticipate the onset of dawn (Dodd et al., 2014). Circadian clock

mutants might also affect WUE by changing the sensitivity of stomatal movements and

photosynthesis to environmental transitions, because there is circadian gating of the

responses of both stomata and photosynthesis to environmental cues (Dodd et al., 2006;

Kinoshita et al., 2011; Litthauer et al., 2015; Joo et al., 2017; Cano-Ramirez et al., 2018).

Some effects of the circadian oscillator upon WUE arise from alterations in leaf size that

occur in some circadian oscillator mutants (Fig. 4A, B). This suggests that developmental

alterations arising from lesions in the circadian oscillator can lead to changes in WUE. Such

developmental alterations might alter WUE by changing airflow around the rosette, boundary

layer conductance, or internal leaf structure.

It has been reported previously that during the light period of light/dark cycles, CCA1-ox has

greater stomatal conductance than the wild type and decreased CO_2 assimilation and

biomass accumulation (Dodd et al., 2005; Graf et al., 2010). If these alterations in growth,

CO_2 fixation and transpiration persist throughout the vegetative growth phase, it might be

predicted that CCA1-ox would have lower long-term WUE than the wild type. We found that

both biomass accumulation and water loss were reduced significantly in CCA1-ox relative to

its background (Fig. 3A, B), with the ratio between the two parameters indicating also a

significant decrease in WUE of CCA1-ox (Fig. 1). This might be due to alterations in gas

exchange reported previously (Dodd et al., 2005), and also other developmental changes

caused by CCA1 overexpression.

Contribution of circadian regulation in guard cells to water use efficiency

We also investigated whether the circadian oscillator within guard cells contributes to long-

term WUE. This involved overexpressing two circadian clock genes in guard cells using two

different guard cell-specific promoters. Comparable approaches have been adopted to

investigate roles of specific cell types in the functioning of the circadian system and their

relationships with physiology and development (Endo et al., 2014; Shimizu et al., 2015;

Hassidim et al., 2017). Under our experimental conditions, we did not identify consistent

alterations in the long-term WUE of seedlings overexpressing *CCA1* or *TOC1* in stomatal

guard cells (Fig. 5B). This could indicate that decreased long-term WUE of TOC1-ox plants

(Fig. 1) arises from altered circadian regulation in cell types other than guard cells. Whilst

two lines harbouring a *GC1::CCA1* construct had greater WUE than the wild type, WUE was

unaltered in comparable lines harbouring *MYB60::CCA1* (Fig. 5B). The differing WUE

phenotype of *GC1::CCA1* and *MYB60::CCA1* might be explained by differences in promoter

strength, because the *GC1* promoter appears to have somewhat greater activity than the

MYB60 promoter (Fig. S3D, E). Although both promoters are guard cell-specific in our hands

(Fig. S3), we cannot exclude the possibility of ectopic promoter activity.

Interestingly, *GC1::CCA1* is reported to have greater drought sensitivity of long-term

biomass accumulation than the wild type (Hassidim et al., 2017), whereas we found that

GC1::CCA1 had greater WUE than the wild type (Fig. 5B). This might reflect the integration

of circadian regulation into ABA signalling (Legnaioli et al., 2009; Robertson et al., 2009), or

occur because guard cell circadian regulation is required for correct guard cell metabolism

and/or stomatal movements under conditions of abiotic stress. For example, circadian

regulation is proposed to participate in daily cycles of triacylglycerol mobilization that are

important for stomatal opening (McLachlan et al., 2016). Together, these findings suggest

that guard cell circadian regulation is important under both well-watered conditions and

conditions of environmental stress (Fig. 5C) (Robertson et al., 2009; Hassidim et al., 2017),

with circadian regulation in other tissues also contributing to overall WUE. It would be

informative in future to perform reverse genetic screening of the dehydration tolerance or

long-term drought tolerance of sets of circadian clock mutants. However, because well-

watered WUE is not a drought tolerance trait (Blum, 2009), it possible that different circadian

clock mutant alleles might confer dehydration or drought tolerance compared with those

alleles that alter WUE (Fig. 1).

Conclusions

We show that circadian regulation contributes to whole plant long-term WUE under cycles of

day and night. This control occurs partly through the influence of components of the

circadian oscillator upon rosette architecture. Mutation or overexpression of CCA1, TOC1,

ELF3, GI, GRP7, PRR9, TEJ, TIC and ZTL altered WUE under our experimental conditions.

The roles of these genes in WUE may be independent or overlapping, and their WUE

phenotypes might be due to direct effects of these genes, or indirect effects on transcript

and/or protein abundance of other circadian clock gene(s). Misregulation of the expression

of CHE, FKF1, LKP2, RVE4, RVE8, PRR3, PRR5, ELF4, LUX and WNK1 did not appear to

alter WUE under our experimental conditions.

Our results have broad implications. Firstly, our data suggest that alterations in circadian

function that arise during crop breeding could have the potential to alter WUE. Therefore,

manipulation of the functioning of the circadian oscillator might represent a pathway to tune

the WUE of crops. Second, our results indicate that circadian regulation in a single cell type

can have implications for whole-plant physiology. Third, our experiments with guard-cell

misregulation of circadian oscillator genes suggest that the circadian oscillator within guard

cells and other cell types influences WUE. Finally, our findings suggest that circadian

regulation potentially alters a single trait (WUE) by affecting many aspects of physiology,

along with leaf area. Overall, our study demonstrates that the circadian oscillator is important

for the water use efficiency of Arabidopsis plants during their entire vegetative growth period.

In future, it will be informative to distinguish the contribution to overall WUE of circadian

regulation within additional cell types, such as the mesophyll, vascular tissue, and root cell

types. It will also be important to identify specific mechanisms underlying the WUE

phenotypes, and determine the extent to which these findings scale to crop species.

Materials and methods

Plant material and growth conditions

Arabidopsis (*Arabidopsis thaliana* (L.) Heynh.) seeds were surface-sterilised as described

previously (Noordally et al., 2013). For experiments investigating stomatal density and index,

seeds were stratified for 3 days at 4 °C, then sown on compost mix comprising a 3:1 ratio of

coarsely sieved Levington Advance F2 seed compost (Everris) and horticultural silver sand

(Melcourt), supplemented with 0.4 g l^{-1} thiacloprid insecticide granules (Exemptor; Everris).

Seedlings were germinated in controlled environment chambers (Reftech, Netherlands)

under an 8 h photoperiod at 70% humidity, 20 °C, and photon flux density of 100 µmol m^{-2} s^{-}

[1] of overhead lighting supplied by cool white fluorescent tubes (Reftech, Netherlands). For

experiments investigating long-term WUE, seeds were sown within a custom Falcon tube

system then stratified. The genotypes that were screened for WUE alterations are identified

in Table S2, and all have been described previously. For all experiments, at least two

completely independent experimental repeats were performed per genotype and per

treatment, with multiple replicate plants within each of the experimental repeats.

Generation of transgenic lines

To create the *GC1::CCA1:nos* (*GC*)*, GC1::TOC1:nos* (*GT*)*, MYB60::CCA1:nos* (*MC*) and

MYB60::TOC1:nos (*MT*) constructs, the CaMV *nos* terminator sequence was ligated

between the SpeI and NotI restriction sites in the pGREENII0229 binary vector (Hellens et

al., 2000). The *GC1* upstream sequence (-1894 to -190) or *MYB60* upstream sequence (-

1724 to -429) was then ligated between the KpnI and ApaI restriction sites of

pGREENII0229. Finally, the *CCA1* coding sequence or *TOC1* coding sequence, obtained

using RT-PCR, was ligated between the restriction XhoI and XmaI sites. Primers used are

identified in Table S3. Constructs were transformed into Col-0 wild type Arabidopsis using

transformation with *Agrobacterium tumefaciens* strain GV3101. Transformants were

identified by screening for phosphinothricin resistance, and then further validated using

genomic DNA PCR. Homozygous lines were identified via phosphinothricin (BASTA)

resistance, and two independently transformed homozygous lines were investigated in detail

per genotype.

Guard cell specificity of promoter activity was investigated using *GC1::GFP:nos* and

MYB60::GFP:nos promoter-reporter lines (Fig. S3A-C), which were created as above with

the *GFP* coding sequence ligated between the XhoI and XmaI restriction sites. Leaf discs (5

mm diameter) from seedlings or mature plants were mounted on microscope slides with

dH$_2$O, and examined for GFP fluorescence using confocal microscopy (Leica DMI6000). The

following settings were used: argon laser at 20% capacity, 488 nm laser at 48% capacity

with a bandwidth of 505 nm–515 nm, gain of 1250, offset at 0.2%, 20x or 40x objective,

zoom x1 to x4.

Measurement of water use efficiency

The WUE assay was adapted from Wituszynska et al. (2013) (Wituszyńska et al., 2013).

Plants were grown for 6 weeks in modified 50 ml Falcon tubes, under an 8 h photoperiod at

70% humidity, 20 °C, and photon flux density of 100 µmol m^{-2} s^{-1} of overhead lighting

supplied by cool white fluorescent tubes (Reftech, Netherlands). The Falcon tube systems

consisted of a 50 ml Falcon tube filled with 37.5 ml of a 1:1 ratio of compost: perlite and 35

ml of Milli-Q water (Merck), with the remaining volume filled with a 1:1 ratio of compost: Milli-

Q water (Fig. S4). Each Falcon tube lid had a 2 mm diameter hole drilled in its centre to

allow plant growth. The lid was spray-painted black (Hycote) because we found that the

orange colour of the Falcon tube lid caused leaf curling (Fig. S4). The system was wrapped

in aluminium foil to exclude light (Fig. S4). 10-15 seeds were sown through the Falcon tube

lid using a pipette. Following stratification, Falcon tube systems were placed under growth

conditions using a randomised experimental design. 7 days after germination, seedlings

were thinned to one per Falcon tube system, and initial Falcon tube weight was recorded.

The seedling-thinning step was sensitive to seedling damage for genotypes with

substantially altered morphologies (e.g. *tps1* mutants), reducing the number of replicates

available for some genotypes. After 6 weeks of growth, rosette leaf surface area was

measured by photography (D50; Nikon) and Fiji software, rosette dry weight was measured

(4 d at 60°C), and final Falcon tube weight was recorded. All experiments were stopped

before flowering occurred, with the 8 h photoperiod being used to delay flowering as much

as possible. Plants were not obviously stressed during the experiment (e.g. leaves did not

become purple due to strong anthocyanin accumulation, and plants did not wilt or become

contaminated with mildew) (Fig. S4). Negative controls (Falcon tube systems without plants)

were used to assess soil water evaporation over 18 experimental batches, with an overall

mean weight loss of 0.513 g ± 0.004 g over 6 weeks for plant-free Falcon tubes.

Plant WUE was calculated as follows:

$$WUE = \frac{d}{(t_i - t_f) - e}$$

Where d is the rosette dry weight at the end of the experiment (mg), t_i and t_f are the falcon

tube weight at the start and end of the experiment, respectively (g), and e is the amount of

water evaporation directly from the compost (g). WUE is derived as mg biomass per ml^{-1}

water lost. These calculations assumed that 1 g of weight change was equivalent to a

change of 1 ml of water. For examination of individual batches of data (Fig. 1), the WUE of

each circadian oscillator genotype was normalized to its respective background to control for

variation in the WUE of each background accession and expressed as a percentage of that

background. Statistical comparisons with the background lines occurred before

normalization. For quantitative investigation of the entire dataset, linear mixed effects

models were used (below).

Measurement of stomatal density

Plants were grown for 7-8 weeks on compost mix. Dental paste (Coltene) was applied to the

abaxial surface of fully expanded leaves. Transparent nail varnish (Rimmel) was applied to

these leaf moulds once they had set, and then peeled away from the mould using clear

adhesive tape (Scotch Crystal). Stomatal and pavement cells were counted within an

800 μm x 800 μm square at the centre of each leaf half, using an epifluorescence

microscope (HAL100; Zeiss) and Volocity (Perkin Elmer) and Fiji software. For each

experimental repeat, two leaves were sampled per plant and eight plants sampled per

genotype. Stomatal index was calculated as follows:

$$SI = \frac{s}{s + p} * 100$$

Where SI is the stomatal index, s the number of stomata in the field of view (800 μm x

800 μm), and p the number of pavement cells in the field of view.

RNA extraction and qRT-PCR

RNA extractions, cDNA synthesis, and qRT-PCR were performed according to (Simon et al.,

2018), except approximately 10 seedlings were used per RNA sample and analysis was

performed using an MXPro 3005 real time PCR system (Agilent) with 5x HOT FIREPol

EvaGreen qPCR mastermix (Solis Biodyne). qRT-PCR primers are provided in Table S4.

Rhythmic features within qPCR data were identified using the BioDare2 platform (Zielinski et

al., 2014), using the Fast Fourier Transform Non-Linear Least Squares method (FFT-NLLS).

One independently-transformed line of each guard cell-specific circadian clock gene

overexpressor was also investigated using qRT-PCR conducted on RNA isolated from

epidermal peels. Abaxial leaf epidermis was detached, then washed in 10 mM MES (pH

6.15, adjusted using 10 M KOH) to remove RNA derived from ruptured epidermal cells. Each

RNA sample was derived from 20 epidermal peels (five plants, four leaves per plant) that

were collated and flash-frozen in liquid nitrogen. Guard cell RNA was extracted using the

RNeasy UCP Micro Kit (Qiagen) according to manufacturer's instructions, with the following

modification: guard cell lysis was performed by adding glass beads (425 µm - 600 µm

diameter, acid washed, from Sigma-Aldrich) and 350 µl RULT buffer to the sample, then

vortexed for 5 min.

Data analysis

Experiments were conducted in a series of 18 separate experimental batches, each

including a set of mutants and their corresponding backgrounds. This subdivision ensured

high quality experimental attention to each replicate within a large-scale study, and the

management of plant growth space. To investigate the nature of any differences between

the mutants and their backgrounds in a manner that accounted for between experimental

batch-variation in the backgrounds, we used a linear mixed effects modelling approach.

Because there are differences between the WUE of Arabidopsis background accessions

(Nienhuis et al., 1994; Woodward et al., 2002; Dodd et al., 2004; Masle et al., 2005; Karaba

et al., 2007; Ruggiero et al., 2017; Ferguson et al., 2018) (Fig. S2), separate models were

generated for each background to avoid comparing accessions with unequal underlying

WUE. WUE data were analysed by fitting a linear mixed models using the lme4 package (R

package version of lme4 v1.1-21) (Bates et al., 2015) within the R statistical computing

platform v3.6.2 (R Core Team, 2019). Using the lmer function, "Mutant" was assigned as a

fixed effect (xf) and the "Batch" as a random effect (xr) in order to test for the effect of the

mutations on the physiological parameters, whilst controlling for differences between

batches. Separate models for each background accession were created with R code:

527
```
model1 <- lmer(Physiological_parameter ~ Mutant + (1|Batch)
```

For the dataset from each Arabidopsis background, diagnostic residual plots suggested that

the model fits were appropriate for larger datasets (Col-0, C24, L. *er.* and Ws), and for

analytical consistency the same model was applied across the entire dataset. Genotypes

within two experimental batches (including *toc1*-101 and TOC1-ox) were analysed within a

separate model because the backgrounds had rather greater water loss and biomass

accumulation than the other experimental batches, so were incorporated into a separate

model to obtain the best possible fit. Conditional R^2 was obtained using the r.squaredGLMM

function in the MuMIn package v1.43.15 (https://CRAN.R-project.org/package=MuMIn)

(Nakagawa and Schielzeth, 2013). The emmeans function (previously lsmeans; (Searle et

al., 1980); https://CRAN.R-project.org/package=emmeans); v1.4.3.01) was used to

subsequently obtain an estimated marginal mean for each mutant, and conduct post-hoc

pairwise comparisons between mutant and its corresponding background, using the

Kenward-Roger method for determining degrees of freedom and Tukey method for P value

adjustment:

542
```
emmeans(model1, list(pairwise ~ Mutant), adjust = "tukey")
```

With this analysis, this output from group comparisons indicate the statistical significance of

any differences of the modelled mutants from the modelled means. This is indicated where

relevant on the figures.

Accession numbers

Arabidopsis Genome Initiative identifiers for the genes mentioned in this study are: *CCA1*

(*CIRCADIAN CLOCK ASSOCIATED1*, At2g46830), *CHE* (*CCA1 HIKING EXPEDITION*,

At5g08330), *ELF3* (*EARLY FLOWERING3*, At2g25930), *ELF4* (*EARLY FLOWERING4*,

At2g40080), *FKF1* (*F BOX1*, At1g68050), *GI* (*GIGANTEA*, At1g22770), *GRP7* (*GLYCINE*

RICH PROTEIN7, At2g21660), *KIN10* (*SNF1-RELATED PROTEIN KINASE1.1*, At3g01090),

LKP2 (*LOV KELCH PROTEIN2*, At2g18915), *LUX* (*LUX ARRHYTHMO*, At3g46640),

MYB60 (*MYB DOMAIN PROTEIN60*, At1g08810), *PRR3* (*PSEUDO-RESPONSE*

REGULATOR3, At5g60100), *PRR5* (*PSEUDO-RESPONSE REGULATOR5*, At5g24470),

PRR9 (*PSEUDO-RESPONSE REGULATOR9*, At2g46790), *RVE4* (*REVEILLE4*,

At5g02840), *TEJ* (*POLY(ADP-RIBOSE)GLYCOHYDROLASE1*, At2g31870), *TIC* (*TIME*

FOR COFFEE, At3gt22380), *TOC1* (*TIMING OF CAB EXPRESSION1*, At5g61380), *TPS1*

(*TREHALOSE-6-PHOSPHATE SYNTHASE1*, At1g78580), *WNK1* (*WITH NO LYSINE*

KINASE1, At3g04910), *ZTL* (*ZEITLUPE*, At5g57360).

Acknowledgements

We thank Kester Cragg-Barber, James Chen, Ioanna Kostaki, Jean-Charles Isner, Deirdre

McLachlan, Peng Sun, Ashutosh Sharma and Dora Cano-Ramirez for technical advice

during experimentation. We thank Marc Knight, Tracy Lawson and Steven Penfield for

discussions and advice concerning data interpretation and analysis. We thank Keara

Franklin, Alex Webb, Paloma Mas, Steve Kay, Isabelle Carre, Takato Imaizumi, Filip

Rolland, Ian Graham, Stacey Harmer, and Steven Penfield for donating seed lines. This

research was funded the UK Biotechnology and Biological Sciences Research Council

(SWBIO DTP awards BB/J014400/1 and BB/M009122/1 and Institute Strategic Programme

GEN BB/P013511/1) and the Wolfson Foundation.

Table 1. Genotypes with misregulated circadian clock-associated genes that have

altered long-term WUE. Findings are summarized from Fig. 1 and derived from statistical

significance within linear mixed effects models.

Significantly greater WUE than background	Significantly lower WUE than background	No significant WUE alteration compared to background
gi-2	CCA1-ox	*cca1*-11
gi-11	*elf3*-1	*che*-1
grp7-1	KIN10-ox 5.7	*che*-2
tej-1	KIN10-ox 6.5	CHE-ox 6
tic-2	*prr9*-1	CHE-ox 17
	TOC1-ox	*elf4*-101
	tps1-11	*fkf1*-2
	tps1-12	*lkp2*-1
	ztl-1	*lux*-1
		prr3-1
		prr5-3
		rve4-1
		rve8-1
		toc1-1

toc1-2

toc1-21

toc1-101

wnk1

 Figure legends

 Figure 1. The circadian clock regulates long-term water use efficiency of Arabidopsis under

light/dark cycles. Estimated WUE of each genotype relative to the estimated WUE of its

background, derived from linear mixed effects models that combine all 18 separate batches

of experimentation, each having 5 - 15 replicate plants per genotype. Statistical analysis

derived from pairwise post-hoc comparisons between mutants and corresponding

backgrounds, using the Kenward-Roger method for determining degrees of freedom and

Tukey method for P-value adjustment. s.e.m. reflects modelled data, rather than the range of

values sampled from the plants.

Figure 2. Relationship between circadian clock-associated characteristics and water use

efficiency. (A-C) Estimated WUE grouped according to (A) phase of expression of each

mutated or overexpressed gene, and the (B) period or (C) flowering time alteration caused

by mutation or overexpression of each gene indicated. Shading of bars on graphs indicates

statistical significance. Studies describing the phase of expression, period and flowering time

of the genotypes tested are identified in the main text. N/C indicates no change. We note

that the phase of expression and period data used for this analysis were often identified in

previous studies under constant conditions, in contrast to our experiments under light/dark

cycles. Estimated WUE relative to its corresponding background is derived from linear mixed

effects models combining all 18 separate batches of experimentation, each having 5 - 15

replicate plants per genotype. Statistical analysis derived from pairwise post-hoc

comparisons between mutants and corresponding backgrounds, using the Kenward-Roger

method for determining degrees of freedom and Tukey method for P-value adjustment.

Figure 3. Manipulating the expression of genes associated with circadian regulation alters

estimated WUE by changing both water use and biomass accumulation. (A) Biomass

accumulation loss and (B) water loss for each genotype relative to its respective background

over the course of the experiments. Genotypes are ordered according to their WUE

calculated in Fig. 1. Data are derived from linear mixed effects models that combine all 18

separate batches of experimentation, each having 5 - 15 replicate plants per genotype.

Statistical analysis derived from pairwise post-hoc comparisons between mutants and

corresponding backgrounds, using the Kenward-Roger method for determining degrees of

freedom and Tukey method for P-value adjustment. Note the model fit for the *tej*-1 mutant

was poor (Table S1). Genotypes are ordered from greatest (left) to lowest (right) estimated

WUE. s.e.m. reflects modelled data, rather than the range of values sampled from the

plants.

Figure 4. The circadian oscillator alters WUE partially by changing rosette architecture. (A)

Altering circadian-associated gene expression can affect rosette architecture and size, as

illustrated for *elf3*-1, *lux*-1, and *gi*-2 in the Col-0 background. Image backgrounds removed

for clarity. Variation in rosette leaf surface area across the genotypes investigated explained

(B) 16% of variation in WUE ($p < 0.001$, $r = 0.400$, $r^2 = 0.160$), (C) 83% of variation in

transpiration ($p < 0.001$, $r = 0.912$, $r^2 = 0.832$) and (D) 73% of variation in rosette dry

biomass ($p < 0.001$, $r = 0.857$, $r^2 = 0.734$). Data were analysed using Pearson correlation

tests.

Figure 5. Long-term WUE can be altered by misexpression of circadian oscillator genes

within stomatal guard cells. (A) Constructs used to overexpress *CCA1* or *TOC1* coding

sequence under control of *GC1* or *MYB60* promoters. (B) Guard cell *CCA1* overexpression

can increase WUE. WUE expressed as absolute WUE and percentage of the wild type

(normalised to wild type as 100%, red reference line). Data were analysed by one-way

ANOVA ($p < 0.001$) followed by pairwise post-hoc Tukey comparisons between transgenic

lines and their corresponding backgrounds (** = $p < 0.01$; *** = $p < 0.001$; N.S. = $p > 0.05$;

$n = 5$ - 15). (C, D) Guard cell *CCA1* or *TOC1* overexpression does not affect (C) stomatal

index nor (D) stomatal density. Data were analysed with ANOVA and Tukey's post hoc tests

(N.S. = $p > 0.05$; $n = 19$ - 32; mean ± S.E.M.). Bar colours identify the whole plant

overexpressor control (black), wild type control (dark grey), and guard cell-specific

overexpressor genotypes (light grey).

References

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK,
Zimmerman J, Barajas P, Chuek R, Gadrinab C, Heller C, Jeske A, Koesma E,
Meyers CC, Parker H, Prednis L, Ansari Y, Choy N, Deen H, Geralt M, Hazari N,
Hom E, Karnes M, Mulholland C, Ndubaku R, Schmidt I, Guzman P, Aguilar-
Henonin L, Schmid M, Weigel D, Carter DE, Marchand T, Risseeuw E, Brogden
D, Zeko A, Crosby WL, Berry CC, Ecker JR (2003) Genome-wide insertional
mutagenesis of *Arabidopsis thaliana*. Science **301:** 653-657
Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of
transcription networks in plant stress and energy signalling. Nature **448:** 938
Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using
lme4. Journal of Statistical Software **67:** 48
Baudry A, Ito S, Song YH, Strait AA, Kiba T, Lu S, Henriques R, Pruneda-Paz JL, Chua
N-H, Tobin EM, Kay SA, Imaizumi T (2010) F-Box proteins FKF1 and LKP2 act in
concert with ZEITLUPE to control Arabidopsis clock progression. The Plant Cell **22:**
606
Bauer H, Ache P, Lautner S, Fromm J, Hartung W, Al-Rasheid Khaled AS, Sonnewald
S, Sonnewald U, Kneitz S, Lachmann N, Mendel Ralf R, Bittner F, Hetherington
Alistair M, Hedrich R (2013) The stomatal response to reduced relative humidity
requires guard cell-autonomous ABA synthesis. Current Biology **23:** 53-57
Bertolino LT, Caine RS, Gray JE (2019) Impact of stomatal density and morphology on
water-use efficiency in a changing world. Frontiers in Plant Science **10:** article 225
Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is the
target of crop yield improvement under drought stress. Field Crops Research **112:**
119-123
Bridge LJ, Franklin KA, Homer ME (2013) Impact of plant shoot architecture on leaf
cooling: a coupled heat and mass transfer model. Journal of The Royal Society
Interface **10:** 20130326
Caine RS, Yin X, Sloan J, Harrison EL, Mohammed U, Fulton T, Biswal AK, Dionora J,
Chater CC, Coe RA, Bandyopadhyay A, Murchie EH, Swarup R, Quick WP, Gray
JE (2019) Rice with reduced stomatal density conserves water and has improved
drought tolerance under future climate conditions. New Phytologist **221:** 371-384
Cano-Ramirez DL, Saskia de Fraine T, Griffiths OG, Dodd AN (2018) Photosynthesis and
circadian rhythms regulate the buoyancy of marimo lake balls. Current Biology **28:**
R869-R870
Cominelli E, Galbiati M, Albertini A, Fornara F, Conti L, Coupland G, Tonelli C (2011)
DOF-binding sites additively contribute to guard cell-specificity of *AtMYB60* promoter.
BMC Plant Biology **11:** article 162
Cominelli E, Galbiati M, Vavasseur A, Conti L, Sala T, Vuylsteke M, Leonhardt N,
Dellaporta SL, Tonelli C (2005) A guard-cell-specific MYB transcription factor
regulates stomatal movements and plant drought tolerance. Current Biology **15:**
1196-1200
Condon AG, Farquhar GD, Rebetzke GJ, Richards RA (2004) Breeding for high water-
use efficiency. Journal of Experimental Botany **55:** 2447-2460
R Core Team (2019) R: A language and environment for statistical computing. R Foundation
for Statistical Computing.
Costa JM, Monnet F, Jannaud D, Leonhardt N, Ksas B, Reiter IM, Pantin F, Genty B
(2015) *OPEN ALL NIGHT LONG*: The dark side of stomatal control. Plant Physiology
167: 289
Coupel-Ledru A, Lebon E, Christophe A, Gallo A, Gago P, Pantin F, Doligez A,
Simonneau T (2016) Reduced nighttime transpiration is a relevant breeding target
for high water-use efficiency in grapevine. Proceedings of the National Academy of
Sciences **113:** 8963

Covington MF, Panda S, Liu XL, Strayer CA, Wagner DR, Kay SA (2001) ELF3 modulates resetting of the circadian clock in Arabidopsis. The Plant Cell **13:** 1305

Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg A, Primavesi LF, Paul MJ, Schluepmann H (2011) Growth arrest by trehalose-6-phosphate: An astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. Plant Physiology **157:** 160

Ding Z, Doyle MR, Amasino RM, Davis SJ (2007) A complex genetic interaction between *Arabidopsis thaliana* TOC1 and CCA1/LHY in driving the circadian clock and in output regulation. Genetics **176:** 1501

Ding Z, Millar AJ, Davis AM, Davis SJ (2007) *TIME FOR COFFEE* encodes a nuclear regulator in the *Arabidopsis thaliana* circadian clock. The Plant Cell **19:** 1522

Dixon LE, Knox K, Kozma-Bognar L, Southern MM, Pokhilko A, Millar AJ (2011) Temporal repression of core circadian genes is mediated through EARLY FLOWERING 3 in Arabidopsis. Current Biology **21:** 120-125

Dodd AN, Dalchau N, Gardner MJ, Baek S-J, Webb AAR (2014) The circadian clock has transient plasticity of period and is required for timing of nocturnal processes in Arabidopsis. New Phytologist **201:** 168-179

Dodd AN, Jakobsen MK, Baker AJ, Telzerow A, Hou S-W, Laplaze L, Barrot L, Scott Poethig R, Haseloff J, Webb AAR (2006) Time of day modulates low-temperature Ca2+ signals in Arabidopsis. The Plant Journal **48:** 962-973

Dodd AN, Parkinson K, Webb AAR (2004) Independent circadian regulation of assimilation and stomatal conductance in the *ztl*-1 mutant of Arabidopsis. New Phytologist **162:** 63-70

Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, Hibberd JM, Millar AJ, Webb AAR (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. Science **309:** 630

Doyle MR, Davis SJ, Bastow RM, McWatters HG, Kozma-Bognár L, Nagy F, Millar AJ, Amasino RM (2002) The ELF4 gene controls circadian rhythms and flowering time in *Arabidopsis thaliana*. Nature **419:** 74

Edwards CE, Ewers BE, McClung CR, Lou P, Weinig C (2012) Quantitative variation in water-use efficiency across water regimes and its relationship with circadian, vegetative, reproductive, and leaf gas-exchange traits. Molecular Plant **5:** 653-668

Edwards CE, Weinig C (2010) The quantitative-genetic and QTL architecture of trait integration and modularity in *Brassica rapa* across simulated seasonal settings. Heredity **106:** 661

Endo M, Shimizu H, Nohales MA, Araki T, Kay SA (2014) Tissue-specific clocks in Arabidopsis show asymmetric coupling. Nature **515:** 419

Eriksson ME, Hanano S, Southern MM, Hall A, Millar AJ (2003) Response regulator homologues have complementary, light-dependent functions in the Arabidopsis circadian clock. Planta **218:** 159-162

Ezer D, Jung J-H, Lan H, Biswas S, Gregoire L, Box MS, Charoensawan V, Cortijo S, Lai X, Stöckle D, Zubieta C, Jaeger KE, Wigge PA (2017) The evening complex coordinates environmental and endogenous signals in Arabidopsis. Nature Plants **3:** 17087

Farré EM, Harmer SL, Harmon FG, Yanovsky MJ, Kay SA (2005) Overlapping and distinct roles of PRR7 and PRR9 in the Arabidopsis circadian clock. Current Biology **15:** 47-54

Ferguson JN, Humphry M, Lawson T, Brendel O, Bechtold U (2018) Natural variation of life-history traits, water use, and drought responses in Arabidopsis. Plant Direct **2:** e00035

Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill J (1999) *GIGANTEA*: a circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. The EMBO Journal **18:** 4679

Francia P, Simoni L, Cominelli E, Tonelli C, Galbiati M (2008) Gene trap-based identification of a guard cell promoter in Arabidopsis. Plant Signaling & Behavior **3:** 684-686

Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D, Yochikawa A, Cano-Ramirez DL, Chembath A, Cragg-Barber K, Haydon MJ, Hotta CT, Vincentz M, Webb AAR, Dodd AN (2018) Circadian entrainment in Arabidopsis by the sugar-responsive transcription factor bZIP63. Current Biology **28:** 2597-2606.e2596

Franks PJ, W. Doheny-Adams T, Britton-Harper ZJ, Gray JE (2015) Increasing water-use efficiency directly through genetic manipulation of stomatal density. New Phytologist **207:** 188-195

Galbiati M, Simoni L, Pavesi G, Cominelli E, Francia P, Vavasseur A, Nelson T, Bevan M, Tonelli C (2008) Gene trap lines identify Arabidopsis genes expressed in stomatal guard cells. The Plant Journal **53:** 750-762

Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA (2010) AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. The Plant Journal **64:** 1-13

Gorton HL, Williams WE, Binns ME, Gemmell CN, Leheny EA, Shepherd AC (1989) Circadian stomatal rhythms in epidermal peels from *Vicia faba*. Plant Physiology **90:** 1329

Graf A, Schlereth A, Stitt M, Smith AM (2010) Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. Proceedings of the National Academy of Sciences **107:** 9458

Hall A, Bastow RM, Davis SJ, Hanano S, McWatters HG, Hibberd V, Doyle MR, Sung S, Halliday KJ, Amasino RM, Millar AJ (2003) The *TIME FOR COFFEE* gene maintains the amplitude and timing of Arabidopsis circadian clocks. The Plant Cell **15:** 2719

Harmer SL, Hogenesch JB, Straume M, Chang H-S, Han B, Zhu T, Wang X, Kreps JA, Kay SA (2000) Orchestrated transcription of key pathways in *Arabidopsis* by the circadian clock. Science **290:** 2110

Hassidim M, Dakhiya Y, Turjeman A, Hussien D, Shor E, Anidjar A, Goldberg K, Green RM (2017) *CIRCADIAN CLOCK ASSOCIATED1* (*CCA1*) and the circadian control of stomatal aperture. Plant Physiology **175:** 1864

Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR (2013) Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. Nature **502:** 689

Hazen SP, Schultz TF, Pruneda-Paz JL, Borevitz JO, Ecker JR, Kay SA (2005) *LUX ARRHYTHMO* encodes a Myb domain protein essential for circadian rhythms. Proceedings of the National Academy of Sciences of the United States of America **102:** 10387

Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM (2000) pGreen: a versatile and flexible binary Ti vector for Agrobacterium-mediated plant transformation. Plant Molecular Biology **42:** 819-832

Hennessey TL, Field CB (1991) Circadian rhythms in photosynthesis. Plant Physiology **96:** 831

Herrero E, Kolmos E, Bujdoso N, Yuan Y, Wang M, Berns MC, Uhlworm H, Coupland G, Saini R, Jaskolski M, Webb A, Gonçalves J, Davis SJ (2012) EARLY FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the *Arabidopsis* circadian clock. The Plant Cell **24:** 428

Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving environmental change. Nature **424:** 901

Hicks KA, Millar AJ, Carré IA, Somers DE, Straume M, Meeks-Wagner DR, Kay SA (1996) Conditional circadian dysfunction of the *Arabidopsis early-flowering 3* mutant. Science **274:** 790

Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. Trends in
Plant Science **19:** 240-249
Hu H, Xiong L (2014) Genetic engineering and breeding of drought-resistant crops. Annual
Review of Plant Biology **65:** 715-741
Huang H, Nusinow DA (2016) Into the evening: Complex interactions in the Arabidopsis
circadian clock. Trends in Genetics **32:** 674-686
Hugouvieux V, Kwak JM, Schroeder JI (2001) An mRNA cap binding protein, ABH1,
modulates early abscisic acid signal transduction in Arabidopsis. Cell **106:** 477-487
Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA (2005) FKF1 F-Box protein mediates
cyclic degradation of a repressor of CONSTANS in *Arabidopsis*. Science **309:** 293
Imaizumi T, Tran HG, Swartz TE, Briggs WR, Kay SA (2003) FKF1 is essential for
photoperiodic-specific light signalling in Arabidopsis. Nature **426:** 302
Joo Y, Fragoso V, Yon F, Baldwin IT, Kim S-G (2017) Circadian clock component, LHY,
tells a plant when to respond photosynthetically to light in nature. Journal of
Integrative Plant Biology **59:** 572-587
Karaba A, Dixit S, Greco R, Aharoni A, Trijatmiko KR, Marsch-Martinez N, Krishnan A,
Nataraja KN, Udayakumar M, Pereira A (2007) Improvement of water use
efficiency in rice by expression of *HARDY*, an *Arabidopsis* drought and salt tolerance
gene. Proceedings of the National Academy of Sciences **104:** 15270
Khanna R, Kikis EA, Quail PH (2003) *EARLY FLOWERING 4* functions in Phytochrome B-
regulated seedling de-etiolation. Plant Physiology **133:** 1530
Kikis EA, Khanna R, Quail PH (2005) ELF4 is a phytochrome-regulated component of a
negative-feedback loop involving the central oscillator components CCA1 and LHY.
The Plant Journal **44:** 300-313
Kim W-Y, Fujiwara S, Suh S-S, Kim J, Kim Y, Han L, David K, Putterill J, Nam HG,
Somers DE (2007) ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA
in blue light. Nature **449:** 356
Kinoshita T, Hayashi Y (2011) New insights into the regulation of stomatal opening by blue
light and plasma membrane H+-ATPase. *In* KW Jeon, ed, International Review of
Cell and Molecular Biology, Vol 289. Academic Press, pp 89-115
Kinoshita T, Ono N, Hayashi Y, Morimoto S, Nakamura S, Soda M, Kato Y, Ohnishi M,
Nakano T, Inoue S-i, Shimazaki K-i (2011) FLOWERING LOCUS T regulates
stomatal opening. Current Biology **21:** 1232-1238
Kobata T, Okuno T, Yamamoto Y (1996) Contributions of capacity for soil water extraction
and water use effficiency to maintenance of dry matter production in rice subjected to
drought. Japanese Journal of Crop Science **65:** 652-662
Lawson T, Blatt MR (2014) Stomatal size, speed, and responsiveness impact on
photosynthesis and water use efficiency. Plant Physiology **164:** 1556
Legnaioli T, Cuevas J, Mas P (2009) TOC1 functions as a molecular switch connecting the
circadian clock with plant responses to drought. The EMBO Journal **28:** 3745
Liang YK, Xie X, Lindsay SE, Wang YB, Masle J, Williamson L, Leyser O, Hetherington
AM (2010) Cell wall composition contributes to the control of transpiration efficiency
in *Arabidopsis thaliana*. Plant Journal **64:** 679-686
Litthauer S, Battle MW, Lawson T, Jones MA (2015) Phototropins maintain robust
circadian oscillation of PSII operating efficiency under blue light. The Plant Journal
83: 1034-1045
Más P, Kim W-Y, Somers DE, Kay SA (2003) Targeted degradation of TOC1 by ZTL
modulates circadian function in *Arabidopsis thaliana*. Nature **426:** 567
Masle J, Gilmore SR, Farquhar GD (2005) The *ERECTA* gene regulates plant transpiration
efficiency in Arabidopsis. Nature **436:** 866
McLachlan Deirdre H, Lan J, Geilfus C-M, Dodd Antony N, Larson T, Baker A, Hõrak H,
Kollist H, He Z, Graham I, Mickelbart Michael V, Hetherington Alistair M (2016)
The breakdown of stored triacylglycerols is required during light-induced stomatal
opening. Current Biology **26:** 707-712

Medrano H, Tomás M, Martorell S, Flexas J, Hernández E, Rosselló J, Pou A, Escalona
J-M, Bota J (2015) From leaf to whole-plant water use efficiency (WUE) in complex
canopies: Limitations of leaf WUE as a selection target. The Crop Journal **3**: 220-228
Meyer S, Mumm P, Imes D, Endler A, Weder B, Al-Rasheid KAS, Geiger D, Marten I,
Martinoia E, Hedrich R (2010) AtALMT12 represents an R-type anion channel
required for stomatal movement in Arabidopsis guard cells. The Plant Journal **63:**
1054-1062
Michael TP, Salomé PA, Yu HJ, Spencer TR, Sharp EL, McPeek MA, Alonso JM, Ecker
JR, McClung CR (2003) Enhanced fitness conferred by naturally occurring variation
in the circadian clock. Science **302:** 1049
Millar AJ (2004) Input signals to the plant circadian clock. Journal of Experimental Botany
55: 277-283
Millar AJ, Carre IA, Strayer CA, Chua NH, Kay SA (1995) Circadian clock mutants in
Arabidopsis identified by luciferase imaging. Science **267:** 1161
Murakami M, Yamashino T, Mizuno T (2004) Characterization of circadian-associated
APRR3 pseudo-response regulator belonging to the APRR1/TOC1 quintet in
Arabidopsis thaliana. Plant and Cell Physiology **45:** 645-650
Na J-K, Metzger JD (2014) Chimeric promoter mediates guard cell-specific gene expression
in tobacco under water deficit. Biotechnology Letters **36:** 1893-1899
Nagy R, Grob H, Weder B, Green P, Klein M, Frelet A, Schjoerring JK, Brearley CA,
Martinoia E (2009) The Arabidopsis ATP-binding cassette protein
ATMRP5/ATABCC5 is a high-affinity inositol hexakisphosphate transporter involved
in guard cell signaling and phytate storage. Journal of Biological Chemistry
Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R2 from
generalized linear mixed-effects models. Methods in Ecology and Evolution **4:** 133-
142
Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010) PSEUDO-
RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in *Arabidopsis*
circadian clock. The Plant Cell **22:** 594
Nakamichi N, Murakami-Kojima M, Sato E, Kishi Y, Yamashino T, Mizuno T (2002)
Compilation and characterization of a novel WNK family of protein kinases in
Arabidopsis thaliana with reference to circadian rhythms. Bioscience, Biotechnology,
and Biochemistry **66:** 2429-2436
Nienhuis J, Sills GR, Martin B, King G (1994) Variance for water-use efficiency among
ecotypes and recombinant inbred lines of *Arabidopsis thaliana* (Brassicaceae).
American Journal of Botany **81:** 943-947
Nietzsche M, Guerra T, Alseekh S, Wiermer M, Sonnewald S, Fernie AR, Börnke F
(2018) STOREKEEPER RELATED1/G-element binding protein (STKR1) interacts
with protein kinase SnRK1. Plant Physiology **176:** 1773
Niwa Y, Ito S, Nakamichi N, Mizoguchi T, Niinuma K, Yamashino T, Mizuno T (2007)
Genetic linkages of the circadian clock-associated genes, *TOC1*, *CCA1* and *LHY*, in
the photoperiodic control of flowering time in *Arabidopsis thaliana*. Plant and Cell
Physiology **48:** 925-937
Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma
M, Tanaka K, Hanaoka M, Dodd AN (2013) Circadian control of chloroplast
transcription by a nuclear-encoded timing signal. Science **339:** 1316
Panda S, Poirier GG, Kay SA (2002) *tej* defines a role for poly(ADP-ribosyl)ation in
establishing period length of the *Arabidopsis* circadian oscillator. Developmental Cell
3: 51-61
Paul MJ, Jhurreea D, Zhang Y, Primavesi LF, Delatte T, Schluepmann H, Wingler A
(2010) Up-regulation of biosynthetic processes associated with growth by trehalose
6-phosphate. Plant Signaling & Behavior **5:** 386-392
Pei Z-M, Ghassemian M, Kwak CM, McCourt P, Schroeder JI (1998) Role of
farnesyltransferase in ABA regulation of guard cell anion channels and plant water
loss. Science **282:** 287

Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. Science **323:** 1481

Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, Salemi MR, Phinney BS, Harmer SL (2011) REVEILLE8 and PSEUDO-REPONSE REGULATOR5 form a negative feedback loop within the Arabidopsis circadian clock. PLOS Genetics **7:** e1001350

Resco de Dios V, Gessler A, Pedro Ferrio J, Alday JG, Bahn M, del Castillo J, Devidal S, García-Muñoz S, Kayler Z, Landais D, Martín-Gómez P, Milcu A, Piel C, Pirhofer-Walzl K, Ravel O, Salekin S, Tissue DT, Tjoelker MG, Voltas J, Roy J (2016) Circadian rhythms have significant effects on leaf-to-canopy scale gas exchange under field conditions. GigaScience **5**

Robertson FC, Skeffington AW, Gardner MJ, Webb AAR (2009) Interactions beween circadian and hormonal signalling in plants. Plant Molecular Biology **69:** 419-427

Rubin MJ, Brock MT, Baker RL, Wilcox S, Anderson K, Davis SJ, Weinig C (2018) Circadian rhythms are associated with shoot architecture in natural settings. New Phytologist **219:** 246-258

Ruggiero A, Punzo P, Landi S, Costa A, Van Oosten JM, Grillo S (2017) Improving plant water use efficiency through molecular genetics. Horticulturae **3**

Rusconi F, Simoni L, Simeoni F, Tonelli C, Galbiati M, Cominelli E, Conti L, Riboni M, Francia P, Martin CR (2013) The Arabidopsis thaliana MYB60 promoter provides a tool for the spatio-temporal control of gene expression in stomatal guard cells. Journal of Experimental Botany **64:** 3361-3371

Ruts T, Matsubara S, Wiese-Klinkenberg A, Walter A (2012) Aberrant temporal growth pattern and morphology of root and shoot caused by a defective circadian clock in *Arabidopsis thaliana*. The Plant Journal **72:** 154-161

Salomé PA, McClung CR (2005) *PSEUDO-RESPONSE REGULATOR 7* and *9* are partially redundant genes essential for the temperature responsiveness of the Arabidopsis circadian clock. The Plant Cell **17:** 791

Schaffer R, Ramsay N, Samach A, Corden S, Putterill J, Carré IA, Coupland G (1998) The *late elongated hypocotyl* mutation of Arabidopsis disrupts circadian rhythms and the photoperiodic control of flowering. Cell **93:** 1219-1229

Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) Guard cell signal transduction. Annual Review of Plant Physiology and Plant Molecular Biology **52:** 627-658

Schultz TF, Kiyosue T, Yanovsky M, Wada M, Kay SA (2001) A Role for LKP2 in the Circadian Clock of Arabidopsis. The Plant Cell **13:** 2659

Searle SR, Speed FM, Milliken GA (1980) Population marginal means in the linear model: an alternative to least squares means. Journal of the American Statistical Association **34:** 216-221

Shimizu H, Katayama K, Koto T, Torii K, Araki T, Endo M (2015) Decentralized circadian clocks process thermal and photoperiodic cues in specific tissues. Nature Plants **1:** 15163

Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. Plant, Cell & Environment **40:** 997-1008

Simon NML, Kusakina J, Fernández-López Á, Chembath A, Belbin FE, Dodd AN (2018) The energy-signaling hub SnRK1 is important for sucrose-induced hypocotyl elongation. Plant Physiology **176:** 1299

Simon NML, Sawkins E, Dodd AN (2018) Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation. Plant Signaling & Behavior **13:** e1457913

Somers DE, Devlin PF, Kay SA (1998) Phytochromes and cryptochromes in the entrainment of the *Arabidopsis* circadian clock. Science **282:** 1488

Somers DE, Schultz TF, Milnamow M, Kay SA (2000) *ZEITLUPE* encodes a novel clock-associated PAS protein from *Arabidopsis*. Cell **101:** 319-329

Somers DE, Webb AA, Pearson M, Kay SA (1998) The short-period mutant, *toc1*-1, alters
circadian clock regulation of multiple outputs throughout development in *Arabidopsis*
thaliana. Development **125:** 485
Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Más P, Panda S, Kreps JA, Kay
SA (2000) Cloning of the *Arabidopsis* clock gene *TOC1*, an autoregulatory response
regulator homolog. Science **289:** 768
Streitner C, Danisman S, Wehrle F, Schöning JC, Alfano JR, Staiger D (2008) The small
glycine-rich RNA binding protein AtGRP7 promotes floral transition in Arabidopsis
thaliana. The Plant Journal **56:** 239-250
Thines B, Harmon FG (2010) Ambient temperature response establishes ELF3 as a
required component of the core Arabidopsis circadian clock. Proceedings of the
National Academy of Sciences**:** 200911006
Tomás M, Medrano H, Escalona JM, Martorell S, Pou A, Ribas-Carbó M, Flexas J
(2014) Variability of water use efficiency in grapevines. Environmental and
Experimental Botany **103:** 148-157
Vialet-Chabrand S, Matthews JSA, Brendel O, Blatt MR, Wang Y, Hills A, Griffiths H,
Rogers S, Lawson T (2016) Modelling water use efficiency in a dynamic
environment: An example using *Arabidopsis thaliana*. Plant Science **251:** 65-74
Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE,
Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate
signaling in *Arabidopsis thaliana*. Science **339:** 704
Wang Y, Liu K, Liao H, Zhuang C, Ma H, Yan X (2008) The plant WNK gene family and
regulation of flowering time in Arabidopsis. Plant Biology **10:** 548-562
Wang ZY, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK
ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own
expression. Cell **93:** 1207-1217
Webb AAR, Seki M, Satake A, Caldana C (2019) Continuous dynamic adjustment of the
plant circadian oscillator. Nature Communications **10:** 550
Wituszyńska W, Ślesak I, Vanderauwera S, Szechyńska-Hebda M, Kornaś A, Van Der
Kelen K, Mühlenbock P, Karpińska B, Maćkowski S, Van Breusegem F,
Karpiński S (2013) LESION SIMULATING DISEASE1, ENHANCED DISEASE
SUSCEPTIBILITY1, and PHYTOALEXIN DEFICIENT4 conditionally regulate cellular
signaling homeostasis, photosynthesis, water use efficiency, and seed yield in
Arabidopsis. Plant Physiology **161:** 1795
Woodward FI, Lake JA, Quick WP (2002) Stomatal development and CO2: ecological
consequences. New Phytologist **153:** 477-484
Xoconostle-Cazares B, Ramirez-Ortega FA, Flores-Elenes L, Ruiz-Medrano R (2010)
Drought tolerance in crop plants. American Journal of Plant Physiology **5:** 241-256
Yamamoto Y, Sato E, Shimizu T, Nakamich N, Sato S, Kato T, Tabata S, Nagatani A,
Yamashino T, Mizuno T (2003) Comparative genetic studies on the APRR5 and
APRR7 genes belonging to the APRR1/TOC1 quintet implicated in circadian rhythm,
control of flowering time, and early photomorphogenesis. Plant and Cell Physiology
44: 1119-1130
Yang Y, Costa A, Leonhardt N, Siegel RS, Schroeder JI (2008) Isolation of a strong
Arabidopsis guard cell promoter and its potential as a research tool. Plant Methods
4: 6
Yanovsky MJ, Kay SA (2002) Molecular basis of seasonal time measurement in
Arabidopsis. Nature **419:** 308
Yoo CY, Pence HE, Hasegawa PM, Mickelbart MV (2009) Regulation of transpiration to
improve crop water use. Critical Reviews in Plant Sciences **28:** 410-431
Yoo CY, Pence HE, Jin JB, Miura K, Gosney MJ, Hasegawa PM, Mickelbart MV (2010)
The Arabidopsis GTL1 transcription factor regulates water use efficiency and drought
tolerance by modulating stomatal density via transrepression of *SDD1*. The Plant
Cell **22:** 4128

Zagotta MT, Shannon S, Jacobs C, Meeks-Wagner DR (1992) Early-flowering mutants of
Arabidopsis thaliana. Functional Plant Biology **19:** 411-418
Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) Strengths and limitations of
period estimation methods for circadian data. PLOS ONE **9:** e96462

1014

1015

Figure 1. The circadian clock regulates long-term water use efficiency of Arabidopsis under light/dark cycles. Estimated WUE of each genotype relative to the estimated WUE of its background, derived from linear mixed effects models that combine all 18 separate batches of experimentation, each having 5 - 15 replicate plants per genotype. Statistical analysis derived from pairwise post-hoc comparisons between mutants and corresponding backgrounds, using the Kenward-Roger method for determining degrees of freedom and Tukey method for P-value adjustment. s.e.m. reflects modelled data, rather than the range of values sampled from the plants.

Figure 2. Relationship between circadian clock-associated characteristics and water use

efficiency. (A-C) Estimated WUE grouped according to (A) phase of expression of each

mutated or overexpressed gene, and the (B) period or (C) flowering time alteration caused

by mutation or overexpression of each gene indicated. Shading of bars on graphs indicates statistical significance. Studies describing the phase of expression, period and flowering time of the genotypes tested are identified in the main text. N/C indicates no change. We note that the phase of expression and period data used for this analysis were often identified in previous studies under constant conditions, in contrast to our experiments under light/dark cycles. Estimated WUE relative to its corresponding background is derived from linear mixed effects models combining all 23 separate batches of experimentation, each having 5 - 15 replicate plants per genotype. Statistical analysis derived from pairwise post-hoc comparisons between mutants and corresponding backgrounds, using the Kenward-Roger method for determining degrees of freedom and Tukey method for P-value adjustment.

Figure 3. Manipulating the expression of genes associated with circadian regulation alters

estimated WUE by changing both water use and biomass accumulation. (A) Biomass

accumulation loss and (B) water loss for each genotype relative to its respective background

over the course of the experiments. Genotypes are ordered according to their WUE calculated in Fig. 1. Data are derived from linear mixed effects models that combine all 23 separate batches of experimentation, each having 5 - 15 replicate plants per genotype. Statistical analysis derived from pairwise post-hoc comparisons between mutants and corresponding backgrounds, using the Kenward-Roger method for determining degrees of freedom and Tukey method for P-value adjustment. Note the model fit for the *tej*-1 mutant was poor (Table S1). Genotypes are ordered from greatest (left) to lowest (right) estimated WUE. s.e.m. reflects modelled data, rather than the range of values sampled from the plants.

Figure 4. The circadian oscillator alters WUE partially by changing rosette architecture. (A) Altering circadian-associated gene expression can affect rosette architecture and size, as illustrated for *elf3*-1, *lux*-1, and *gi*-2 in the Col-0 background. Image backgrounds removed

for clarity. Variation in rosette leaf surface area across the genotypes investigated explained

(B) 16% of variation in WUE ($p < 0.001$, $r = 0.400$, $r^2 = 0.160$), (C) 83% of variation in

transpiration ($p < 0.001$, $r = 0.912$, $r^2 = 0.832$) and (D) 73% of variation in rosette dry

biomass ($p < 0.001$, $r = 0.857$, $r^2 = 0.734$). Data were analysed using Pearson correlation

tests.

Figure 5. Long-term WUE can be altered by misexpression of circadian oscillator genes within stomatal guard cells. (A) Constructs used to overexpress *CCA1* or *TOC1* coding sequence under control of *GC1* or *MYB60* promoters. (B) Guard cell *CCA1* overexpression can increase WUE. WUE expressed as absolute WUE and percentage of the wild type (normalised to wild type as 100%, red reference line). Data were analysed by one-way ANOVA (p < 0.001) followed by pairwise post-hoc Tukey comparisons between transgenic

lines and their corresponding backgrounds (** = $p < 0.01$; *** = $p < 0.001$; N.S. = $p > 0.05$; n = 5 - 15). (C, D) Guard cell *CCA1* or *TOC1* overexpression does not affect (C) stomatal index nor (D) stomatal density. Data were analysed with ANOVA and Tukey's post hoc tests (N.S. = $p > 0.05$; n = 19 - 32; mean ± S.E.M.). Bar colours identify the whole plant overexpressor control (black), wild type control (dark grey), and guard cell-specific overexpressor genotypes (light grey).

Parsed Citations

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Chuek R, Gadrinab C, Heller C, Jeske A, Koesma E, Meyers CC, Parker H, Prednis L, Ansari Y, Choy N, Deen H, Geralt M, Hazari N, Hom E, Karnes M, Mulholland C, Ndubaku R, Schmidt I, Guzman P, Aguilar-Henonin L, Schmid M, Weigel D, Carter DE, Marchand T, Risseeuw E, Brogden D, Zeko A, Crosby WL, Berry CC, Ecker JR (2003) Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science 301: 653-657
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature 448: 938
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67: 48
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Baudry A, Ito S, Song YH, Strait AA, Kiba T, Lu S, Henriques R, Pruneda-Paz JL, Chua N-H, Tobin EM, Kay SA, Imaizumi T (2010) F-Box proteins FKF1 and LKP2 act in concert with ZEITLUPE to control Arabidopsis clock progression. The Plant Cell 22: 606
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bauer H, Ache P, Lautner S, Fromm J, Hartung W, Al-Rasheid Khaled AS, Sonnewald S, Sonnewald U, Kneitz S, Lachmann N, Mendel Ralf R, Bittner F, Hetherington Alistair M, Hedrich R (2013) The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. Current Biology 23: 53-57
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bertolino LT, Caine RS, Gray JE (2019) Impact of stomatal density and morphology on water-use efficiency in a changing world. Frontiers in Plant Science 10: article 225
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. Field Crops Research 112: 119-123
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bridge LJ, Franklin KA, Homer ME (2013) Impact of plant shoot architecture on leaf cooling: a coupled heat and mass transfer model. Journal of The Royal Society Interface 10: 20130326
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Caine RS, Yin X, Sloan J, Harrison EL, Mohammed U, Fulton T, Biswal AK, Dionora J, Chater CC, Coe RA, Bandyopadhyay A, Murchie EH, Swarup R, Quick WP, Gray JE (2019) Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. New Phytologist 221: 371-384
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cano-Ramirez DL, Saskia de Fraine T, Griffiths OG, Dodd AN (2018) Photosynthesis and circadian rhythms regulate the buoyancy of marimo lake balls. Current Biology 28: R869-R870
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cominelli E, Galbiati M, Albertini A, Fornara F, Conti L, Coupland G, Tonelli C (2011) DOF-binding sites additively contribute to guard cell-specificity of AtMYB60 promoter. BMC Plant Biology 11: article 162
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cominelli E, Galbiati M, Vavasseur A, Conti L, Sala T, Vuylsteke M, Leonhardt N, Dellaporta SL, Tonelli C (2005) A guard-cell-specific MYB transcription factor regulates stomatal movements and plant drought tolerance. Current Biology 15: 1196-1200
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Condon AG, Farquhar GD, Rebetzke GJ, Richards RA (2004) Breeding for high water-use efficiency. Journal of Experimental Botany 55: 2447-2460
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

R Core Team (2019) R: A language and environment for statistical computing. R Foundation for Statistical Computing.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Costa JM, Monnet F, Jannaud D, Leonhardt N, Ksas B, Reiter IM, Pantin F, Genty B (2015) OPEN ALL NIGHT LONG: The dark side of stomatal control. Plant Physiology 167: 289

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Coupel-Ledru A, Lebon E, Christophe A, Gallo A, Gago P, Pantin F, Doligez A, Simonneau T (2016) Reduced nighttime transpiration is a relevant breeding target for high water-use efficiency in grapevine. Proceedings of the National Academy of Sciences 113: 8963

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Covington MF, Panda S, Liu XL, Strayer CA, Wagner DR, Kay SA (2001) ELF3 modulates resetting of the circadian clock in Arabidopsis. The Plant Cell 13: 1305

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg A, Primavesi LF, Paul MJ, Schluepmann H (2011) Growth arrest by trehalose-6-phosphate: An astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. Plant Physiology 157: 160

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ding Z, Doyle MR, Amasino RM, Davis SJ (2007) A complex genetic interaction between Arabidopsis thaliana TOC1 and CCA1/LHY in driving the circadian clock and in output regulation. Genetics 176: 1501

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ding Z, Millar AJ, Davis AM, Davis SJ (2007) TIME FOR COFFEE encodes a nuclear regulator in the Arabidopsis thaliana circadian clock. The Plant Cell 19: 1522

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dixon LE, Knox K, Kozma-Bognar L, Southern MM, Pokhilko A, Millar AJ (2011) Temporal repression of core circadian genes is mediated through EARLY FLOWERING 3 in Arabidopsis. Current Biology 21: 120-125

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Dalchau N, Gardner MJ, Baek S-J, Webb AAR (2014) The circadian clock has transient plasticity of period and is required for timing of nocturnal processes in Arabidopsis. New Phytologist 201: 168-179

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Jakobsen MK, Baker AJ, Telzerow A, Hou S-W, Laplaze L, Barrot L, Scott Poethig R, Haseloff J, Webb AAR (2006) Time of day modulates low-temperature Ca2+ signals in Arabidopsis. The Plant Journal 48: 962-973

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Parkinson K, Webb AAR (2004) Independent circadian regulation of assimilation and stomatal conductance in the ztl-1 mutant of Arabidopsis. New Phytologist 162: 63-70

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, Hibberd JM, Millar AJ, Webb AAR (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. Science 309: 630

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Doyle MR, Davis SJ, Bastow RM, McWatters HG, Kozma-Bognár L, Nagy F, Millar AJ, Amasino RM (2002) The ELF4 gene controls circadian rhythms and flowering time in Arabidopsis thaliana. Nature 419: 74

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Edwards CE, Ewers BE, McClung CR, Lou P, Weinig C (2012) Quantitative variation in water-use efficiency across water regimes and its relationship with circadian, vegetative, reproductive, and leaf gas-exchange traits. Molecular Plant 5: 653-668

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Edwards CE, Weinig C (2010) The quantitative-genetic and QTL architecture of trait integration and modularity in Brassica rapa across simulated seasonal settings. Heredity 106: 661

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Endo M, Shimizu H, Nohales MA, Araki T, Kay SA (2014) Tissue-specific clocks in Arabidopsis show asymmetric coupling. Nature 515: 419

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Eriksson ME, Hanano S, Southern MM, Hall A, Millar AJ (2003) Response regulator homologues have complementary, light-dependent functions in the Arabidopsis circadian clock. Planta 218: 159-162

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ezer D, Jung J-H, Lan H, Biswas S, Gregoire L, Box MS, Charoensawan V, Cortijo S, Lai X, Stöckle D, Zubieta C, Jaeger KE, Wigge PA (2017) The evening complex coordinates environmental and endogenous signals in Arabidopsis. Nature Plants 3: 17087

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Farré EM, Harmer SL, Harmon FG, Yanovsky MJ, Kay SA (2005) Overlapping and distinct roles of PRR7 and PRR9 in the Arabidopsis circadian clock. Current Biology 15: 47-54

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ferguson JN, Humphry M, Lawson T, Brendel O, Bechtold U (2018) Natural variation of life-history traits, water use, and drought responses in Arabidopsis. Plant Direct 2: e00035

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill J (1999) GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. The EMBO Journal 18: 4679

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Francia P, Simoni L, Cominelli E, Tonelli C, Galbiati M (2008) Gene trap-based identification of a guard cell promoter in Arabidopsis. Plant Signaling & Behavior 3: 684-686

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D, Yochikawa A, Cano-Ramirez DL, Chembath A, Cragg-Barber K, Haydon MJ, Hotta CT, Vincentz M, Webb AAR, Dodd AN (2018) Circadian entrainment in Arabidopsis by the sugar-responsive transcription factor bZIP63. Current Biology 28: 2597-2606.e2596

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Franks PJ, W. Doheny-Adams T, Britton-Harper ZJ, Gray JE (2015) Increasing water-use efficiency directly through genetic manipulation of stomatal density. New Phytologist 207: 188-195

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Galbiati M, Simoni L, Pavesi G, Cominelli E, Francia P, Vavasseur A, Nelson T, Bevan M, Tonelli C (2008) Gene trap lines identify Arabidopsis genes expressed in stomatal guard cells. The Plant Journal 53: 750-762

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA (2010) AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. The Plant Journal 64: 1-13

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gorton HL, Williams WE, Binns ME, Gemmell CN, Leheny EA, Shepherd AC (1989) Circadian stomatal rhythms in epidermal peels from Vicia faba. Plant Physiology 90: 1329

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Graf A, Schlereth A, Stitt M, Smith AM (2010) Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. Proceedings of the National Academy of Sciences 107: 9458

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hall A, Bastow RM, Davis SJ, Hanano S, McWatters HG, Hibberd V, Doyle MR, Sung S, Halliday KJ, Amasino RM, Millar AJ (2003) The TIME FOR COFFEE gene maintains the amplitude and timing of Arabidopsis circadian clocks. The Plant Cell 15: 2719

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Harmer SL, Hogenesch JB, Straume M, Chang H-S, Han B, Zhu T, Wang X, Kreps JA, Kay SA (2000) Orchestrated transcription of key pathways in Arabidopsis by the circadian clock. Science 290: 2110

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hassidim M, Dakhiya Y, Turjeman A, Hussien D, Shor E, Anidjar A, Goldberg K, Green RM (2017) CIRCADIAN CLOCK ASSOCIATED1

(CCA1) and the circadian control of stomatal aperture. Plant Physiology 175: 1864
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR (2013) Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. Nature 502: 689
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hazen SP, Schultz TF, Pruneda-Paz JL, Borevitz JO, Ecker JR, Kay SA (2005) LUX ARRHYTHMO encodes a Myb domain protein essential for circadian rhythms. Proceedings of the National Academy of Sciences of the United States of America 102: 10387
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM (2000) pGreen: a versatile and flexible binary Ti vector for Agrobacterium-mediated plant transformation. Plant Molecular Biology 42: 819-832
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hennessey TL, Field CB (1991) Circadian rhythms in photosynthesis. Plant Physiology 96: 831
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Herrero E, Kolmos E, Bujdoso N, Yuan Y, Wang M, Berns MC, Uhlworm H, Coupland G, Saini R, Jaskolski M, Webb A, Gonçalves J, Davis SJ (2012) EARLY FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the Arabidopsis circadian clock. The Plant Cell 24: 428
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving environmental change. Nature 424: 901
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hicks KA, Millar AJ, Carré IA, Somers DE, Straume M, Meeks-Wagner DR, Kay SA (1996) Conditional circadian dysfunction of the Arabidopsis early-flowering 3 mutant. Science 274: 790
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. Trends in Plant Science 19: 240-249
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hu H, Xiong L (2014) Genetic engineering and breeding of drought-resistant crops. Annual Review of Plant Biology 65: 715-741
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Huang H, Nusinow DA (2016) Into the evening: Complex interactions in the Arabidopsis circadian clock. Trends in Genetics 32: 674-686
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hugouvieux V, Kwak JM, Schroeder JI (2001) An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in Arabidopsis. Cell 106: 477-487
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA (2005) FKF1 F-Box protein mediates cyclic degradation of a repressor of CONSTANS in Arabidopsis. Science 309: 293
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Imaizumi T, Tran HG, Swartz TE, Briggs WR, Kay SA (2003) FKF1 is essential for photoperiodic-specific light signalling in Arabidopsis. Nature 426: 302
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Joo Y, Fragoso V, Yon F, Baldwin IT, Kim S-G (2017) Circadian clock component, LHY, tells a plant when to respond photosynthetically to light in nature. Journal of Integrative Plant Biology 59: 572-587
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Karaba A, Dixit S, Greco R, Aharoni A, Trijatmiko KR, Marsch-Martinez N, Krishnan A, Nataraja KN, Udayakumar M, Pereira A (2007) Improvement of water use efficiency in rice by expression of HARDY, an Arabidopsis drought and salt tolerance gene. Proceedings of the National Academy of Sciences 104: 15270
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Khanna R, Kikis EA, Quail PH (2003) EARLY FLOWERING 4 functions in Phytochrome B-regulated seedling de-etiolation. Plant Physiology 133: 1530

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kikis EA, Khanna R, Quail PH (2005) ELF4 is a phytochrome-regulated component of a negative-feedback loop involving the central oscillator components CCA1 and LHY. The Plant Journal 44: 300-313

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kim W-Y, Fujiwara S, Suh S-S, Kim J, Kim Y, Han L, David K, Putterill J, Nam HG, Somers DE (2007) ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. Nature 449: 356

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kinoshita T, Hayashi Y (2011) New insights into the regulation of stomatal opening by blue light and plasma membrane H+-ATPase. In KW Jeon, ed, International Review of Cell and Molecular Biology, Vol 289. Academic Press, pp 89-115

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kinoshita T, Ono N, Hayashi Y, Morimoto S, Nakamura S, Soda M, Kato Y, Ohnishi M, Nakano T, Inoue S-i, Shimazaki K-i (2011) FLOWERING LOCUS T regulates stomatal opening. Current Biology 21: 1232-1238

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kobata T, Okuno T, Yamamoto Y (1996) Contributions of capacity for soil water extraction and water use effficiency to maintenance of dry matter production in rice subjected to drought. Japanese Journal of Crop Science 65: 652-662

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lawson T, Blatt MR (2014) Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. Plant Physiology 164: 1556

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Legnaioli T, Cuevas J, Mas P (2009) TOC1 functions as a molecular switch connecting the circadian clock with plant responses to drought. The EMBO Journal 28: 3745

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Liang YK, Xie X, Lindsay SE, Wang YB, Masle J, Williamson L, Leyser O, Hetherington AM (2010) Cell wall composition contributes to the control of transpiration efficiency in Arabidopsis thaliana. Plant Journal 64: 679-686

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Litthauer S, Battle MW, Lawson T, Jones MA (2015) Phototropins maintain robust circadian oscillation of PSII operating efficiency under blue light. The Plant Journal 83: 1034-1045

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Más P, Kim W-Y, Somers DE, Kay SA (2003) Targeted degradation of TOC1 by ZTL modulates circadian function in Arabidopsis thaliana. Nature 426: 567

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Masle J, Gilmore SR, Farquhar GD (2005) The ERECTA gene regulates plant transpiration efficiency in Arabidopsis. Nature 436: 866

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

McLachlan Deirdre H, Lan J, Geilfus C-M, Dodd Antony N, Larson T, Baker A, Hõrak H, Kollist H, He Z, Graham I, Mickelbart Michael V, Hetherington Alistair M (2016) The breakdown of stored triacylglycerols is required during light-induced stomatal opening. Current Biology 26: 707-712

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Medrano H, Tomás M, Martorell S, Flexas J, Hernández E, Rosselló J, Pou A, Escalona J-M, Bota J (2015) From leaf to whole-plant water use efficiency (WUE) in complex canopies: Limitations of leaf WUE as a selection target. The Crop Journal 3: 220-228

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Meyer S, Mumm P, Imes D, Endler A, Weder B, Al-Rasheid KAS, Geiger D, Marten I, Martinoia E, Hedrich R (2010) AtALMT12 represents an R-type anion channel required for stomatal movement in Arabidopsis guard cells. The Plant Journal 63: 1054-1062

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Michael TP, Salomé PA, Yu HJ, Spencer TR, Sharp EL, McPeek MA, Alonso JM, Ecker JR, McClung CR (2003) Enhanced fitness conferred by naturally occurring variation in the circadian clock. Science 302: 1049
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Millar AJ (2004) Input signals to the plant circadian clock. Journal of Experimental Botany 55: 277-283
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Millar AJ, Carre IA, Strayer CA, Chua NH, Kay SA (1995) Circadian clock mutants in Arabidopsis identified by luciferase imaging. Science 267: 1161
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Murakami M, Yamashino T, Mizuno T (2004) Characterization of circadian-associated APRR3 pseudo-response regulator belonging to the APRR1/TOC1 quintet in Arabidopsis thaliana. Plant and Cell Physiology 45: 645-650
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Na J-K, Metzger JD (2014) Chimeric promoter mediates guard cell-specific gene expression in tobacco under water deficit. Biotechnology Letters 36: 1893-1899
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Nagy R, Grob H, Weder B, Green P, Klein M, Frelet A, Schjoerring JK, Brearley CA, Martinoia E (2009) The Arabidopsis ATP-binding cassette protein ATMRP5/ATABCC5 is a high-affinity inositol hexakisphosphate transporter involved in guard cell signaling and phytate storage. Journal of Biological Chemistry
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R2 from generalized linear mixed-effects models. Methods in Ecology and Evolution 4: 133-142
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010) PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in Arabidopsis circadian clock. The Plant Cell 22: 594
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Nakamichi N, Murakami-Kojima M, Sato E, Kishi Y, Yamashino T, Mizuno T (2002) Compilation and characterization of a novel WNK family of protein kinases in Arabidopsis thaliana with reference to circadian rhythms. Bioscience, Biotechnology, and Biochemistry 66: 2429-2436
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Nienhuis J, Sills GR, Martin B, King G (1994) Variance for water-use efficiency among ecotypes and recombinant inbred lines of Arabidopsis thaliana (Brassicaceae). American Journal of Botany 81: 943-947
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Nietzsche M, Guerra T, Alseekh S, Wiermer M, Sonnewald S, Fernie AR, Börnke F (2018) STOREKEEPER RELATED1/G-element binding protein (STKR1) interacts with protein kinase SnRK1. Plant Physiology 176: 1773
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Niwa Y, Ito S, Nakamichi N, Mizoguchi T, Niinuma K, Yamashino T, Mizuno T (2007) Genetic linkages of the circadian clock-associated genes, TOC1, CCA1 and LHY, in the photoperiodic control of flowering time in Arabidopsis thaliana. Plant and Cell Physiology 48: 925-937
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma M, Tanaka K, Hanaoka M, Dodd AN (2013) Circadian control of chloroplast transcription by a nuclear-encoded timing signal. Science 339: 1316
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Panda S, Poirier GG, Kay SA (2002) tej defines a role for poly(ADP-ribosyl)ation in establishing period length of the Arabidopsis circadian oscillator. Developmental Cell 3: 51-61
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Paul MJ, Jhurreea D, Zhang Y, Primavesi LF, Delatte T, Schluepmann H, Wingler A (2010) Up-regulation of biosynthetic processes associated with growth by trehalose 6-phosphate. Plant Signaling & Behavior 5: 386-392

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pei Z-M, Ghassemian M, Kwak CM, McCourt P, Schroeder JI (1998) Role of farnesyltransferase in ABA regulation of guard cell anion channels and plant water loss. Science 282: 287

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. Science 323: 1481

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, Salemi MR, Phinney BS, Harmer SL (2011) REVEILLE8 and PSEUDO-REPONSE REGULATOR5 form a negative feedback loop within the Arabidopsis circadian clock. PLOS Genetics 7: e1001350

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Resco de Dios V, Gessler A, Pedro Ferrio J, Alday JG, Bahn M, del Castillo J, Devidal S, García-Muñoz S, Kayler Z, Landais D, Martín-Gómez P, Milcu A, Piel C, Pirhofer-Walzl K, Ravel O, Salekin S, Tissue DT, Tjoelker MG, Voltas J, Roy J (2016) Circadian rhythms have significant effects on leaf-to-canopy scale gas exchange under field conditions. GigaScience 5

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Robertson FC, Skeffington AW, Gardner MJ, Webb AAR (2009) Interactions beween circadian and hormonal signalling in plants. Plant Molecular Biology 69: 419-427

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rubin MJ, Brock MT, Baker RL, Wilcox S, Anderson K, Davis SJ, Weinig C (2018) Circadian rhythms are associated with shoot architecture in natural settings. New Phytologist 219: 246-258

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ruggiero A, Punzo P, Landi S, Costa A, Van Oosten JM, Grillo S (2017) Improving plant water use efficiency through molecular genetics. Horticulturae 3

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rusconi F, Simoni L, Simeoni F, Tonelli C, Galbiati M, Cominelli E, Conti L, Riboni M, Francia P, Martin CR (2013) The Arabidopsis thaliana MYB60 promoter provides a tool for the spatio-temporal control of gene expression in stomatal guard cells. Journal of Experimental Botany 64: 3361-3371

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ruts T, Matsubara S, Wiese-Klinkenberg A, Walter A (2012) Aberrant temporal growth pattern and morphology of root and shoot caused by a defective circadian clock in Arabidopsis thaliana. The Plant Journal 72: 154-161

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Salomé PA, McClung CR (2005) PSEUDO-RESPONSE REGULATOR 7 and 9 are partially redundant genes essential for the temperature responsiveness of the Arabidopsis circadian clock. The Plant Cell 17: 791

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schaffer R, Ramsay N, Samach A, Corden S, Putterill J, Carré IA, Coupland G (1998) The late elongated hypocotyl mutation of Arabidopsis disrupts circadian rhythms and the photoperiodic control of flowering. Cell 93: 1219-1229

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) Guard cell signal transduction. Annual Review of Plant Physiology and Plant Molecular Biology 52: 627-658

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schultz TF, Kiyosue T, Yanovsky M, Wada M, Kay SA (2001) A Role for LKP2 in the Circadian Clock of Arabidopsis. The Plant Cell 13: 2659

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Searle SR, Speed FM, Milliken GA (1980) Population marginal means in the linear model: an alternative to least squares means. Journal of the American Statistical Association 34: 216-221

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Shimizu H, Katayama K, Koto T, Torii K, Araki T, Endo M (2015) Decentralized circadian clocks process thermal and photoperiodic cues in specific tissues. Nature Plants 1: 15163

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. Plant, Cell & Environment 40: 997-1008

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Simon NML, Kusakina J, Fernández-López Á, Chembath A, Belbin FE, Dodd AN (2018) The energy-signaling hub SnRK1 is important for sucrose-induced hypocotyl elongation. Plant Physiology 176: 1299

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Simon NML, Sawkins E, Dodd AN (2018) Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation. Plant Signaling & Behavior 13: e1457913

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Somers DE, Devlin PF, Kay SA (1998) Phytochromes and cryptochromes in the entrainment of the Arabidopsis circadian clock. Science 282: 1488

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Somers DE, Schultz TF, Milnamow M, Kay SA (2000) ZEITLUPE encodes a novel clock-associated PAS protein from Arabidopsis. Cell 101: 319-329

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Somers DE, Webb AA, Pearson M, Kay SA (1998) The short-period mutant, toc1-1, alters circadian clock regulation of multiple outputs throughout development in Arabidopsis thaliana. Development 125: 485

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Más P, Panda S, Kreps JA, Kay SA (2000) Cloning of the Arabidopsis clock gene TOC1, an autoregulatory response regulator homolog. Science 289: 768

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Streitner C, Danisman S, Wehrle F, Schöning JC, Alfano JR, Staiger D (2008) The small glycine-rich RNA binding protein AtGRP7 promotes floral transition in Arabidopsis thaliana. The Plant Journal 56: 239-250

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Thines B, Harmon FG (2010) Ambient temperature response establishes ELF3 as a required component of the core Arabidopsis circadian clock. Proceedings of the National Academy of Sciences: 200911006

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tomás M, Medrano H, Escalona JM, Martorell S, Pou A, Ribas-Carbó M, Flexas J (2014) Variability of water use efficiency in grapevines. Environmental and Experimental Botany 103: 148-157

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Vialet-Chabrand S, Matthews JSA, Brendel O, Blatt MR, Wang Y, Hills A, Griffiths H, Rogers S, Lawson T (2016) Modelling water use efficiency in a dynamic environment: An example using Arabidopsis thaliana. Plant Science 251: 65-74

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis thaliana. Science 339: 704

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang Y, Liu K, Liao H, Zhuang C, Ma H, Yan X (2008) The plant WNK gene family and regulation of flowering time in Arabidopsis. Plant Biology 10: 548-562

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang ZY, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. Cell 93: 1207-1217

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Webb AAR, Seki M, Satake A, Caldana C (2019) Continuous dynamic adjustment of the plant circadian oscillator. Nature Communications 10: 550

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wituszyńska W, Ślesak I, Vanderauwera S, Szechyńska-Hebda M, Kornaś A, Van Der Kelen K, Mühlenbock P, Karpińska B, Maćkowski S, Van Breusegem F, Karpiński S (2013) LESION SIMULATING DISEASE1, ENHANCED DISEASE SUSCEPTIBILITY1, and PHYTOALEXIN DEFICIENT4 conditionally regulate cellular signaling homeostasis, photosynthesis, water use efficiency, and seed yield in Arabidopsis. Plant Physiology 161: 1795

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Woodward FI, Lake JA, Quick WP (2002) Stomatal development and CO2: ecological consequences. New Phytologist 153: 477-484

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Xoconostle-Cazares B, Ramirez-Ortega FA, Flores-Elenes L, Ruiz-Medrano R (2010) Drought tolerance in crop plants. American Journal of Plant Physiology 5: 241-256

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yamamoto Y, Sato E, Shimizu T, Nakamich N, Sato S, Kato T, Tabata S, Nagatani A, Yamashino T, Mizuno T (2003) Comparative genetic studies on the APRR5 and APRR7 genes belonging to the APRR1/TOC1 quintet implicated in circadian rhythm, control of flowering time, and early photomorphogenesis. Plant and Cell Physiology 44: 1119-1130

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yang Y, Costa A, Leonhardt N, Siegel RS, Schroeder JI (2008) Isolation of a strong Arabidopsis guard cell promoter and its potential as a research tool. Plant Methods 4: 6

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yanovsky MJ, Kay SA (2002) Molecular basis of seasonal time measurement in Arabidopsis. Nature 419: 308

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoo CY, Pence HE, Hasegawa PM, Mickelbart MV (2009) Regulation of transpiration to improve crop water use. Critical Reviews in Plant Sciences 28: 410-431

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoo CY, Pence HE, Jin JB, Miura K, Gosney MJ, Hasegawa PM, Mickelbart MV (2010) The Arabidopsis GTL1 transcription factor regulates water use efficiency and drought tolerance by modulating stomatal density via transrepression of SDD1. The Plant Cell 22: 4128

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zagotta MT, Shannon S, Jacobs C, Meeks-Wagner DR (1992) Early-flowering mutants of Arabidopsis thaliana. Functional Plant Biology 19: 411-418

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) Strengths and limitations of period estimation methods for circadian data. PLOS ONE 9: e96462

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title