An evaluation of pool-sequencing transcriptome-based exon capture for

population genomics in non-model species

Emeline Deleury[1], Thomas Guillemaud[1], Aurélie Blin[1] & Eric Lombaert[1]

[1] INRA, CNRS, Université Côte d'Azur, ISA, France

Corresponding author:

Emeline Deleury

Institut Sophia Agrobiotech - 400 Route des Chappes - BP 167 – 06 903 Sophia Antipolis cedex -

FRANCE

E-mail: emeline.deleury@inra.fr

Phone: +33 4 92 38 64 24

Keywords: target enrichment, non-model organism, population genomics, pool-sequencing,

Harmonia axyridis, intron-exon boundary prediction

Abstract

Exon capture coupled to high-throughput sequencing constitutes a cost-effective technical solution for addressing specific questions in evolutionary biology by focusing on expressed regions of the genome preferentially targeted by selection. Transcriptome-based capture, a process that can be used to capture the exons of non-model species, is use in phylogenomics. However, its use in population genomics remains rare due to the high costs of sequencing large numbers of indexed individuals across multiple populations. We evaluated the feasibility of combining transcriptome-based capture and the pooling of tissues from numerous individuals for DNA extraction as a cost-effective, generic and robust approach to estimating the variant allele frequencies of any species at the population level. We designed capture probes for ~5 Mb of randomly chosen *de novo* transcripts from the Asian ladybird *Harmonia axyridis* (5,717 transcripts). We called ~300,000 bi-allelic SNPs for a pool of 36 non-indexed individuals. Capture efficiency was high, and pool-seq was as effective and accurate as individual-seq for detecting variants and estimating allele frequencies. Finally, we also evaluated an approach for simplifying bioinformatic analyses by mapping genomic reads directly to targeted transcript sequences to obtain coding variants. This approach is effective and does not affect the estimation of SNP allele frequencies, except for a small bias close to some exon ends. We demonstrate that this approach can also be used to predict the intron-exon boundaries of targeted *de novo* transcripts, making it possible to abolish genotyping biases near exon ends.

Introduction

One of the core challenges in population genomics and phylogenetics is obtaining a reliable set of orthologous loci from a sufficient number of individuals across multiple populations and from species spanning a range of divergences, respectively. The targeted resequencing of a consistent subset of genomic regions represents a reproducible and cost-effective technical solution, with much lighter data handling than for whole-genome sequencing. Hybridization capture methods are among the available reduced representation methods available for the enrichment of genomic DNA samples in a preselected set of hundreds to thousands of genes or DNA fragments (Hodges *et al.*, 2007). The most common application of hybridization sequence capture is exon capture, in which only coding regions of the genome (and their flanking regions) are sequenced (Warr *et al.*, 2015). This approach provides focused information about gene function and adaptation, and, unlike transcriptome sequencing, is not biased by variations of gene expression.

Exome capture requires a knowledge of the exon sequences of the species studied, for the design of appropriate oligonucleotide probes. Consequently, its use has been largely limited to model species (i.e. species for which a reference genome is available). Absolute probe specificity is not required, however, and given the tendency of functional elements to be conserved even in after millions of years of divergence, exons can be captured from a non-model species (i.e. a species without a published genome sequence) with probes designed from the sequence of a related model species (e.g. Cosart *et al.*, 2011; Förster *et al.*, 2018). This approach has been used for the successful resolution of species phylogenies (e.g. Ilves & López-Fernández, 2014; Bossert & Danforth, 2018; Ilves *et al.*, 2018). It is also an effective method for capturing and enriching degraded and contaminated ancient DNA from extinct groups in paleontology (e.g. Castellano *et al.*, 2014).

An alternative approach to capture probe design for non-model species involves the use of

a *de novo* assembled transcriptome for the species studied or a related species (e.g. Bi *et al.*, 2012;

Neves *et al.*, 2013). One of the limitations of this approach is that the intron-exon structure is

unknown, and probes based on transcript sequences spanning two consecutive exons therefore

hybridize only partially to genomic DNA, reducing capture efficiency (e.g. Neves *et al.*, 2013). One

way of addressing this issue is to use orthologous genes from closely related model species to

predict intron positions (Bi *et al.*, 2012; Stephens *et al.*, 2015; Bragg *et al.*, 2016). However, the use

of data from closely related model species biases target choice through the retention of genes

displaying the highest degree of conservation throughout evolution, which is potentially

problematic in population genomics studies. Even when orthologous genes are targeted,

transcriptome-based exon capture studies have been found to have a lower capture efficiency close

to intron-exon boundaries, because it is not possible to design capture probes spanning exons and

their (unknown) non-coding flanking regions (Bi *et al.*, 2012; Neves *et al.*, 2013; Portik *et al.*, 2016).

Captured fragments are usually longer than probes. Transcriptome-based exon capture

therefore provides opportunities to capture the flanking regions of the exons, thereby expanding

gene models beyond the available coding sequence (Bi *et al.*, 2012; Neves *et al.*, 2013; Stephens *et*

al., 2015; Bragg *et al.*, 2016; Portik *et al.*, 2016). This requires the construction, in a fragmented way

(e.g. long introns not fully captured), of the genomic sequences corresponding to the targeted

transcripts (e.g. Neves *et al.*, 2013). This assembly (generally referred to as an in-target assembly) is

used as a reference for the mapping of genomic reads or for phylogeny construction. This assembly

is obtained by cleaning the raw genomic sequences and assembling them with various k-mer

combinations. The genomic contigs obtained are then compared with the target sequences for

ordering and scaffolding with each other when they correspond to the same targets. Such a complex

approach is necessary for phylogeny construction when probes are used to capture orthologous

targets from related species, but may be dispensable in the context of population genomics. Indeed,

several studies have shown that the assembly process does not completely rebuild all targets (e.g.

Bragg *et al.*, 2016; Portik *et al.*, 2016), and this may decrease the likelihood of detecting variants.

Bragg et al. (2016) tried to attenuate this problem by mapping the genomic reads of the species

used for probe design directly onto the transcript target sequences, thereby making it possible to

cover exons that were not successfully assembled but were nevertheless captured and sequenced.

Transcriptome-based exon capture has proved effective for the generation of thousands of

highly informative markers. Most studies to date have aimed to resolve the phylogenetic

relationships between various set of related species (Nicholls *et al.*, 2015; Stephens *et al.*, 2015;

Bragg *et al.*, 2016; Heyduk *et al.*, 2016; Ilves *et al.*, 2018), and applications in population genomics

remain rare (but see Bi *et al.*, 2013; Rellstab *et al.*, 2019). This is probably because population studies

traditionally analyze large numbers of individuals from multiple populations of non-model species,

which is complex, time-consuming and costly, even if sequencing capacity and costs are optimized

by indexing individuals for multiplexing (Shearer *et al.*, 2012). However, many research questions in

population genomics can be addressed by simply using the allele frequencies calculated at

population level (e.g. Gutenkunst *et al.*, 2009). In this context, pooling individuals within the

population before the extraction of DNA may be an efficient, cheap and time-saving strategy. Pool-

sequencing has been shown to estimate allele frequencies within populations as accurately as

individual genotyping, but with much less effort for library construction and sequencing (e.g. Gautier

et al., 2013; Schlötterer *et al.*, 2014). Allele frequencies have been estimated by a combination of

targeted capture methods and pool-sequencing strategies only in humans (Bansal *et al.*, 2011; Day-

Williams *et al.*, 2011; Ramos *et al.*, 2012; Ryu *et al.*, 2018) and, very recently, in pine trees (Rellstab

et al., 2019).

We propose here an efficient way of quantifying the single-nucleotide polymorphism (SNP)

of exomes at the population level in non-model species for which no genome sequence is available.

This approach is based on a combination of transcriptome-based exon capture and the pooling of

tissues from numerous individuals before DNA extraction. We report the development of capture

probes for the Asian ladybird *Harmonia axyridis* from its *de novo* transcriptome, without the use of

the available draft genome (HaxR v1.0; Gautier *et al.*, 2018) and, thus, in the absence of known gene

structure, as in most non-model species. We used these probes for two target enrichment

experiments with the same individuals, the first based on capture with indexed multiplexed

individuals, and the second based on capture with a pool of non-indexed individuals. We evaluated

the efficiency of capture and the accuracy of the estimates of allele frequencies obtained by pool

sequencing in this context. We also evaluated a rapid approach for the processing of read data

without their assembly, based on the mapping of reads directly to targeted *de novo* transcript

sequences for the detection of variant loci (Bragg *et al.*, 2016; Rellstab *et al.*, 2019). This direct

mapping also makes it possible to effectively predict the locations of the intron-exon boundaries of

target sequences.

Methods
123

Design of capture probes for target enrichment

We designed capture probes for ~5 Mb of randomly chosen coding exonic regions of *H. axyridis*.

Using a *de novo* transcriptome obtained by RNA-Seq on various tissues and conditions/stages of the

species (Vogel *et al.*, 2017), we sought putative peptide-coding sequences with FRAMEDP (v1.2.2,

Gouzy *et al.*, 2009), using bLASTX results (e-value ≤ 1e^{-7}) of transcripts against the insect proteomes

of *Tribolium castaneum*, *Anoplophora glabripennis, Dendroctonus ponderosae, Drosophila*

melanogaster and the SwissProt database (v2016-02-17) for training. The corresponding partial or

complete coding sequences (CDS) were filtered to eliminate (i) *Wolbachia* and other putative

endosymbiont sequences, (ii) CDS with more than 1% missing data (*Ns*) or with more than four

consecutive *Ns*, (iii) CDS with a GC% below 25 or above 75 and (iv) CDS with lengths below 400 bp

(to avoid the capture of too great a proportion of flanking regions and to allow better tiling) or above

3500 bp (to make it possible to target larger numbers of different genes). We identified 28,326

putative CDS, of which 12,739 were retained after the application of the filters. We then chose 5.5

Mb at random, corresponding to 5,736 CDS. We searched for sequence similarity with complete

proteins from the SwissProt database and the *D. melanogaster* and *T. castaneum* proteomes with

BlastX, and at least 895 CDS (15.6%) were found to be complete. Overall, 5,120 CDS (89.3%)

displayed hits with *T. castaneum* proteins (e-value $\leq 1e^{-7}$).

The final probes for the 5,736 selected CDS were designed and produced by NimbleGen. The

repetitiveness of the target sequences was assessed on the basis of the frequency of 15-mers in the

genomes of the coleopterans *T. castaneum* and *A. glabripennis*. Probes with more than one close

match, defined as no more than five single-base insertions, deletions or substitutions with the

SSAHA algorithm (Ning *et al.*, 2001), in the *H. axyridis de novo* transcriptome or in the draft genome

of *A. glabripennis* were discarded. Probes matching to sequences in the *H. axyridis* or *T. castaneum*

mitochondrial genome were also discarded. The few residual *Ns* in target sequences were replaced

with a random nucleotide to allow tile probing across a single unknown base. The final probe set

contained 6,400 regions of overlapping probes, corresponding to a targeted exome size of 5,347,461

bases   distributed   over   5,717   of   our   randomly   selected   CDS   (Zenodo

https://doi.org/10.5281/zenodo.2598388). This probe set was used for the preparation of a SeqCap

EZ Developer assay by Roche NimbleGen Technologies, which synthesized the probes as biotinylated

DNA oligomers. One capture reaction contained 2,100,000 overlapping probes of 50 to 99 bp in

length, with a mean length of 74.71 ± 4.92 bp.

Individual and pooled genomic DNA library preparation

Thirty-six individuals of *H. axyridis* from a wild population in China (Beijing; 40.057128°N,

116.53989°E) were collected in October 2015, immediately placed in RNAlater solution and stored

at -20°C. Based on the PIF tools proposed by Gautier *et al.* (2013), we would theoretically expect a

similar precision of allele frequency estimation for the sequencing of 23 individually indexed

individuals, each at 80X coverage, or a single pool of 36 individuals at 125X, assuming an

experimental error (i.e. ε, the coefficient of variation of the contribution of each individual to the

set of sequence reads obtained from the DNA pool) of 50%. We thus extracted DNA (i)

independently from 23 individuals, using half the body (without the hind legs), and (ii) from a pool

of 36 individuals, including the 23 individuals used for individual sequencing, using two hind legs per

individual. All DNA extractions were performed with the Qiagen DNeasy Blood & Tissue kit, in

accordance with the manufacturer's recommendations.

Genomic libraries were prepared with NimbleGen SeqCap EZ Library (v5.0). Briefly, for each

of the 24 samples (1 pool and 23 individuals), DNA (2 µg in 100 µl) was mechanically sheared to a

mean fragment length of 200 bp with a Covaris S2 E210 device (6x30s). Fragments were end-

repaired, A-tailed and indexed (one index per sample) with KAPA Library Preparation Kit Illumina

platforms. After the ligation of Illumina adaptors and indexes, only fragments with a length of 250

to 450 bp were retained. PCR was performed (7 cycles) with standard Illumina paired-end primers

and the amplicons were then purified with AMPure XP beads (Beckman). The length, quality and

concentration of the prepared DNA fragments were checked on a BioAnalyzer (Agilent High

Sensitivity DNA Assay) and a Qubit.

Sequence capture hybridization and sequencing

The 23 indexed individuals were pooled at equimolar concentrations before hybridization. For each capture (one with the pool and one with the indexed individuals), a total of 1 µg of amplified DNA was used for exome enrichments, with the SeqCap EZ Developer probes described above, strictly in accordance with the NimbleGen SeqCap EZ Library protocol (v5.0). For each capture, two parallel post-capture PCRs (14 cycles) were performed on elution solution, and the PCR products were then merged before purification with AMPure XP beads. The length, quality and concentration of the final DNA fragments were checked on a BioAnalyzer (Agilent High Sensitivity DNA Assay) and a Qubit. Each capture was sequenced on a half lane of an Illumina HiSeq3000 sequencer (on the same lane), in accordance with the manufacturer's instructions, in paired-end mode, for 150 cycles. The data were then demultiplexed and exported as FastQ files, and the 24 libraries were processed independently.

SNP calling by the mapping of reads to the CDS

Sequence quality was checked with FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Adapters were removed and low-quality base pairs were trimmed with Trimmomatic (v0.35, Bolger *et al.*, 2014), with the following parameters: ILLUMINACLIP:TruSeq-file.fa:2:30:10 LEADING:30 TRAILING:30 SLIDINGWINDOW:10:30 MINLEN:75, and FASTX (v0.0.14; http://hannonlab.cshl.edu/fastx_toolkit/) to remove the first five bases.

We chose to align the reads directly with the targeted CDS sequences, because target enrichment was performed on the same species as probe design, as suggested by Bragg *et al.* (2016; see also Rellstab *et al.*, 2019). Doing so, we avoid the complex and time-consuming steps of read

assembly and scaffolding/ordering the genomic contigs obtained to generate the in-target

assemblies for read mapping (see introduction). For each library, cleaned reads were aligned with

the 5,717 targeted CDS, with BOWTIE2 (v2.2.5; Langmead & Salzberg, 2012), using the default

parameters and *local* option to allow for genomic reads not aligning with the targeted CDS sequence

over their entire length, whilst ensuring a minimum alignment length of 20 bp. Reads were aligned

as unpaired, to allow for the mapping of orphan reads. Reads with multiple alignments, and those

for which the mate-pair was aligned to a different target were discarded after mapping. The aligned

data in SAM format were converted into BAM format and sorted with SAMtools (v0.1.19). Duplicate

reads were estimated with picard tools (v1.113) MarkDuplicates, using the default settings.

Coverage at each target base was calculated with mosdepth (v0.2.3; Pedersen & Quinlan, 2018).

Variant calling was performed with read alignments of more than 40 bases in length

containing fewer than 10 deletions, as deduced from the *cigar* format. SNP sites were called

separately for each of the 24 libraries with the SAMtools *mpileup* command, using option *-B*

followed by the varscan2 (v2.3; Koboldt *et al.*, 2012) *mpileup2snp* command with a minimum base

quality of 35 to count a read at a given position, and a *p*-value threshold of 0.01. For the other

parameters, we considered only positions with a minimum coverage of 40 reads for each individual,

and 250 reads for the pool (i.e. for these selected coverages, the same exome size was analyzed:

90.3% and 93% of target bases were covered by at least 40 and 250 reads for the individual and

pooled exomes, respectively; see Fig. S1). We also only included a variant if it was supported by 15

reads per individual, or by three reads for the single pool (given the pool's minimum coverage of

250X, we expected to have ~3 reads per haplotype if the 72 haplotypes in the pool were

homogeneously captured). Moreover to account for the variation in total read coverage across

positions, we required a minimum variant allele frequency threshold of 0.25 for each individual and

0.01 for the pool. For individual libraries, an individual was considered to be homozygous for the

variant if the variant allele frequency f(v) was above 0.75, and heterozygous if f(v) was between 0.25

and 0.75.

All 409,328 different SNP positions independently found in the 24 libraries were then

genotyped for each library with varscan2 *mpileup2cns,* using the same parameters as described

above. Several filters were then applied to all SNP positions, using either in-house PERL or R scripts

(R Development Core Team, 2015). We first retained only bi-allelic positions. We then retained only

positions genotyped for at least 20 individuals. Finally, we discarded SNPs with high levels of

coverage in the pool, which were likely to represent repetitive regions or paralogs of the genome:

maximum pool coverage was set at 6,960X, as calculated with the formula proposed by Li et al.

(2014): $c+4*\sqrt{c}$, where c is the mean coverage in the pool (c=6,634.8X). We retained 300,036 bi-

allelic SNPs after the application of all these filters.

At each site, and for each capture, variant allele frequency at population level was estimated

as the number of reads with the variant allele divided by total coverage. Individual genotypes were

used to calculate exact *p*-values for Hardy-Weinberg equilibrium tests with the "HardyWeinberg" R

package (Graffelman, 2015), with the correction of significance level by the false discovery rate

procedure (Benjamini & Hochberg, 1995). Finally, SNPs were annotated and categorized with the

SnpEff program (v4.3t; Cingolani *et al.*, 2012).

Evaluation of the approach based on read mapping directly onto CDS

Approaches based on the direct mapping of reads onto CDS have already been used to detect SNPs

(e.g. Rellstab *et al.*, 2019), but they have never been evaluated. We used an *H. axyridis* draft genome

sequence (HaxR v1.0; Gautier *et al.*, 2018) to evaluate this approach and that proposed below for

the prediction of intron-exon boundaries (IEBs). We selected CDS targets with a genomic match, as

proposed by Neves *et al.* (2013), with some adjustments. Briefly, CDS target sequences were first

aligned with genomic scaffolds with BLASTN (e-value cutoff of e^{-20} and identity cutoff of 90%), and

the best hit was retained for each target. The target sequences were then realigned with the

corresponding genomic scaffold, with both GMAP (version 2015-12-31; Wu & Watanabe, 2005) and

EXONERATE *est2genome* (v2.2.0;

http://www.genome.iastate.edu/bioinfo/resources/manuals/exonerate/). We isolated targets (i)

with genome matches over their entire length, (ii) for which intron/exon structure predictions were

the same for both alignment programs, (iii) which were covered by capture probes over their entire

length and (iv) for which we had access to at least 200 bp upstream and downstream from the

targeted CDS sequences. With these filters, we retained 3,161 CDS targets in total (3,220,718 bp,

i.e. 60.23% of our targeted exome). These 3,161 full or partial CDS corresponded to 11,754 exons,

with a mean of 3.72 exons per target (range: 1 to 17 exons) and a mean exon length of 274 bp

(range: 12 to 3,417 bp). These 11,754 exons included 6,295 that were unambiguously complete,

because they were surrounded by two others, and 8,593 reliable IEBs in total were identified

between consecutive exons.

All genomic regions corresponding to these 3,161 CDS targets were then isolated, including

200 bp upstream and downstream from the target sequences. Independently for each library, all

cleaned reads were aligned with these genomic regions, with BOWTIE2 (v2.2.5), using the default

parameters and the *end-to-end* option (i.e. reads must be aligned along their full length). Reads with

multiple alignments, and reads for which the mate-pair was aligned to another target were

discarded after mapping. Coverage calculation, SNP calling and variant allele frequency estimation

were performed as described above. We established the correspondence between the position of

a base in the genomic region and its position in the CDS sequence, using the *vulgar* format of

EXONERATE output. SNP positions for which there was a deletion or an insertion between the CDS

sequence and its associated genomic sequence (i.e. 559 positions) were excluded from the

comparison.

Intron-exon boundary detection

Our approach is based solely on the direct mapping of genomic reads to the CDS sequences, by

allowing reads to map over less than their full length (i.e. local alignment). Thus, a genomic read

with both part of an exon and its flanking sequences (intron or UTR) will start mapping to the

corresponding exon in the CDS sequence, and stop mapping at the exon end (Conklin *et al.*, 2013).

This will generate a signal of the intron-exon boundary in the number of reads that either begin or

end their local alignment at that position (hereafter referred to as *nbBE*). Along a covered exon,

nbBE should be much lower than the coverage rate, except at the end of the exon, where we should

observe a large increase in *nbBE* to a value theoretically equal to the coverage (Fig. S2). This strong

increase in *nbBE* should occur either at the base at the exact end of the exon, or a few bases away

if the start of the intron sequence is, by chance, highly similar to the start of the next exon sequence

on the CDS (Fig. S2). In this second case, it creates an abnormally inflated estimate of coverage

locally for a few bases at the boundary of the following exon (Fig. S2 and Fig. S3 for an example). At

an IEB (i.e. a junction of two exon ends in the CDS), depending on whether a signal offset occurs for

one or both exon ends, the size of the area with abnormally inflated local coverage is variable, but

generally less than 10 bp. In addition, the presence of a SNP at some bases at the end of the exon

may prevent reads with the variant from aligning up to the IEB position, potentially generating

another signal in addition to those of the two exon ends of the IEB. Combinations of these different

scenarios may also occur.

We developed and tested a method for predicting IEBs on the basis of the increase in *nbBE,*

by focusing on the 3,161 targeted transcripts for which genomic sequences were available, for which

intron-exon structure was known. We used the mapping output BAM file obtained with the pool

library. At each position in the transcript, *nbBE* was calculated from the start read alignment position

on the transcript and the *cigar* code, both present in the BAM file. The value of the *nbBE* parameter

at each position is a function of coverage (and read length). We took the possible effect of coverage

heterogeneity on *nbBE* into account by calculating the median *nbBE* over a window of 11 bases

centered on the position concerned. A signal (i.e. a predicted IEB), was detected at a position if *nbBE*

was *X* times larger than the median, *X* being a tool parameter defined by the user. For the definition

of *X*, it may be necessary to identify a subset of transcripts with orthologs in a close model species,

and then to determine which value of *X* provides the best prediction of IEBs for this subset. Signals

at the exact ends of the CDS were not considered. We took the possible signal deviation (i.e. a signal

that is slightly deported with respect to the IEB that generates it; Fig. S2) into account, by extending

signal detection to the three bases upstream, and three bases downstream from a position with

measured signal (default value for parameter *n*=3). A measured signal therefore predicts an IEB in a

window of seven bases in total. An IEB, composed of two exon ends, theoretically generates two

close signals with overlapping windows. If these two signals are found at the exact positions of the

exon ends (no deviation), then the predicted region measures 8 bp (Fig. S2). If these two signals

overlap (e.g. because of the deviation of one of the two signals) then the predicted region for the

IEB is 7 bp long. If these signals deviate by up to six bases from each other (i.e. cumulative deviation

of eight bases, the deviation being in opposite directions for the two signals) then the predicted

region measures 14 bp (Fig. S2). More rarely, if a third signal is detected in the same region, it

increases the length of the region by a few bases. If the two signals deviate by more than six bases

from each other, then the tool predicts two signal windows separated by a few bases for the IEB

concerned. Thus, each transcript was divided into short regions with prediction signals (+)

alternating with regions without prediction signals (-). Each region was then assessed as follows:

true positive (TP) if the + region contained an IEB; false positive (FP) if the + region did not contain

an IEB; true negative (TN) if the - region did not contain an IEB; and false negative (FN) if the - region

contained an IEB. We evaluated the performance of the method, by calculating the percentage of

true IEBs correctly predicted (i.e. sensitivity (SN)) as TP/(TP+FN) and the percentage of correct

predictions (i.e. specificity (SP)) as TP/(TP+FP) (Burset & Guigo, 1996). For a CDS with no introns (i.e.

862 of the 3,161 CDS), SN cannot be evaluated; SP was 100 if no IEB was predicted (good prediction)

and 0 otherwise. Scripts are available from https://github.com/edeleury/IEB-finder.

Results

Capture data and filtration

In the two target enrichment experiments, we obtained 186,998,614 and 185,457,659 *H. axyridis*

raw sequence read pairs for the pool library (SRA ERX3237193) and the 23 indexed individuals (SRA

ERX3237194 to ERX3237216), respectively. For each indexed individual library, we obtained a mean

of 8,063,376 raw read pairs (range: 6,912,561 to 9,176,292). After cleaning, 91.2% of the raw read

sequences were retained (see detailed information for all libraries in Table S1).

Capture efficiency, coverage and duplicates

Rather than assembling the cleaned reads, we chose to map them directly onto the target CDS

sequences used for probe design. Mean capture specificity, defined as the percentage of cleaned

reads mapping correctly (i.e. meeting our mapping parameters) onto the 5,717 targeted CDS, was

83.2% (range over libraries: 80.6%-89.2%; Table S2). Aligned bases accounted for 74.6% (range over

all libraries: 72.6%-75.7%) of all cleaned read bases. The global median read alignment length was

144 bp (range over libraries: 143-144 bp, see Table S2), with 62.7% of the mapped reads (range over

all libraries: 60.9%-64.3%) mapped over their entire length. A substantial proportion of the mapped

reads (22.2%; range over all libraries: 20%-25.9%) were 'orphaned' mates (i.e. only one of the two

mates aligned to the target sequences). This percentage strongly suggests that flanking regions of

targeted regions were also sequenced with our capture procedure (Yi *et al.*, 2010; Neves *et al.*,

2013), although quantification was not possible with this direct mapping approach.

The mean percentage of the 5,347,461 targeted bases (5,717 target CDS) covered by at least

one read (i.e. the capture sensitivity) was 93.5% (range over all libraries: 93.4%-93.9%, see Fig. S1

and Table S3). Overall, 93% of the targeted bases were covered in all 24 libraries, whereas 6% were

covered by no read in any of the libraries (Table 1). We identified 491 full targets (292,327 bp) with

no read aligned in the 24 libraries. For these uncovered targets, 487 (99.2%) had no match to the *H.*

axyridis draft genome either, whereas, for the 5,226 covered targeted CDS, only 74 (1.4%) had no

match to the genome (here partial alignments of targets with genome were also considered).

| Target | Target size | Percentage of samples with target sequence | |
		100%	0%
CDS level	5 717	5 226 (91.4%)	491 (8.6%)
Base level	5 347 461	4 967 690 (92.9%)	323 418 (6.0%)

Table 1: Reproducibility of capture sensitivity on the 24 libraries. 100% means that all 24 samples had at least one read

aligned with the target, whereas 0 % level indicates that no reads aligned with the target.

Considering all targeted bases, the median and mean base coverage for individual libraries

were 142X and 283X for the "23 individuals" libraries, and 3,361X and 6,635X for the pool libraries

(see Table S3), respectively. A small number of targeted bases had very high coverage, but around

90% had a coverage <300X for individuals or <7,000X for the pool (Table S3). For individuals, 90.3%

of the targeted bases on average had a mean coverage greater than 40X. For the pool, 93% had a

coverage greater than 250X (Fig. S1).

Base coverage varied along the targeted CDS (Fig. S4A). Using the known complete exons (≥

20 bp) identified on targeted CDS with a match over their entire length with the *H. axyridis* draft

genome, we confirmed that mean coverage increased steadily with exon size (Fig. S4B), and that

coverage decreased from the middle of an exon to its ends (Fig. S4C). Alignments of less than 20 bp

between the reads and the genome sequence were not permitted with this mapping procedure, so

coverage was probably underestimated at exon ends and for short exons. For all the targeted CDS

considered together, mean coverage differed between targeted CDS but was highly reproducible

between libraries (Fig. S5): the correlation coefficient of the target coverage between individual

libraries ranged from 0.96 to 0.99, with a mean of 0.99, and the coefficient of correlation between

the two capture processes (i.e. between pool and individuals) was 0.999.

For the 23 individual libraries, a mean of 66.1% of reads were scored as duplicates (range

over libraries: 63.6%-68.9%; Table S4). This high proportion was due to the very strong sequencing

effort regarding the target region size. Indeed, pooling the reads of the 23 individual libraries

increased the proportion of duplicates detected to >96%, as for the pool (Table S4), suggesting that

the reads scored as duplicates in our experiment probably mostly come from different individuals.

We did not therefore discard duplicated reads before SNP calling, so as to avoid artificially biasing

the estimation of allele frequencies.

SNP calling and accuracy of allele frequency estimation

SNP calling performed independently for the individuals and for the pool, following direct mapping

onto the targeted CDS sequences, led to the detection of 302,292 and 364,144 different putative

exonic SNPs, respectively, with 257,108 positions (62.8%) detected in both experiments. As

expected (because there were 13 more individuals in the pool), the number of SNPs called for the

pool was larger than that for the 23 individuals, with 26.2% of SNPs called for the pool only. The vast

majority (95.6%) of the 11% of SNPs called only for individuals had allele frequencies below 0.03.

All positions were genotyped in all libraries. After filtering, we finally selected a total of

300,036 exonic bi-allelic SNPs within 4,741 targeted CDS. Minor allele frequencies (MAFs) were

below 0.03 for 59.2% of these SNPs (Fig. 1); 92.75% were polymorphic in both the pool and the

individuals, 7.22% were polymorphic only in the pool, and 0.03% were polymorphic only in the

individuals. Only 1,007 of the total number of SNPs (i.e. 0.34%) displayed significant deviation from

Hardy-Weinberg equilibrium, as calculated with the individual genotypes. This number was more

than three times higher (i.e. 3,331 loci) before the removal of loci on the basis of the maximum

coverage filter applied to the pool.

Figure 1: Folded-site frequency spectrum of the 300,036 filtered exonic bi-allelic SNPs (mapping onto targeted CDS

sequences) according to their polymorphism in the pool and in individuals. For SNPs polymorphic in both the pool and

individuals, the MAF shown is the MAF calculated for the pool.

When all 300,036 loci were considered, allele frequency estimates were strongly correlated

between the pool and individuals (*r*=0.99; Fig. 2A). These results were not dependent on the large

fraction of SNPs with low allele frequency estimates (r=0.98, calculated for 102,494 loci with MAF ≥

0.03).

SNP annotation revealed that 74.75% of SNPs were synonymous, whereas 24.92% were non-

synonymous missense and 0.33% were non-synonymous nonsense SNPs.

Figure 2: Correlation of the SNP allele frequencies estimated for the population pool and individuals by two mapping

approaches: **(A)** mapping reads directly onto CDS and **(B)** mapping onto genomic regions available only for a subset of

target CDS. For exonic bi-allelic SNPs common to both mapping approaches, we also show **(C)** the correlation between

the allelic frequencies estimated for the pool by the two approaches.

Comparison of the SNPs detected with the two mapping approaches

We evaluated the ability of the method for mapping reads directly to CDS to call SNPs correctly by

comparing the SNPs identified with this approach and those found by mapping reads to the genome.

We used only the subset of 3,161 targeted CDS with a match over their entire length to sequences

in the draft genome of *H. axyridis* for this comparison. All trimmed reads were also mapped along

their entire length onto the regions of the genome corresponding to this subset. After application

of the same procedure and filters for SNP calling as were used for direct mapping onto CDS

sequences, 290,620 filtered bi-allelic SNPs were genotyped within the subset, including 187,591

(64.5%) in exons and 103,029 (35.5%) in the flanking regions (intron or UTR) of the exons. By

mapping reads onto the genome, we demonstrated that allele frequency estimates were strongly

correlated between the pool and individuals, for both exonic and flanking SNPs (r=0.99; Fig. 2B).

Of the bi-allelic exonic SNPs found by direct mapping onto the targeted CDS (Fig. 2A),

185,426 mapped onto the subset of 3,161 targeted CDS, and 94% were also identified as bi-allelic

SNPs by mapping to genomic regions. For these 174,307 exonic bi-allelic SNPs found with both

mapping approaches (mapping onto the genome or directly onto transcripts), the allele frequency

estimates obtained with the two mapping methods were highly correlated both for the pool

(r=0.998; Fig. 2C) and for the individuals (r=0.998).

Some of the private bi-allelic SNPs were variants identified with both mapping approaches

but discarded during filtering in one of the approaches. For example, 1,467 bi-allelic SNPs private to

the CDS mapping process had a coverage above the maximum threshold, and 714 were tri-allelic

SNPs when mapped onto the genome. If we exclude this latter group, 8,853 and 8,333 were found

to be private to CDS mapping and mapping onto genomic regions, respectively. A non-exhaustive

analysis of these positions highlighted two possible origins of private SNPs for mapping onto CDS.

Some targeted regions (the full CDS or part of the CDS) have homologous copies in the genome that

we were unable to identify during target selection and for which the SNPs found were not discarded

with the "maximum coverage" filter. We have thus identified 43 CDS with reads mapping onto their

genomic regions in a non-unique manner, and these CDS with paralogs displayed 3,141 SNPs, all

private to CDS mapping. The second origin was false SNPs occurring when the start of an intron

could be aligned with the start of the next exon due to similarities between the two sequences (Fig.

S2 and S6, for example). Of the SNPs private to the CDS mapping method, 6.9% (N=614) were

positioned close to IEB positions (i.e. at the exact end of the exon or on the three adjacent bases),

whereas only 0.6% of the 174,307 SNPs common to both mapping methods were found near IEBs.

Prediction of IEBs without read assembly

The method we developed for IEB detection based on the mapping of genomic reads directly onto

CDS sequences was evaluated on the 3,161 targeted CDS for which we had genomic information,

and therefore knew the positions of the IEBs. An IEB signal was predicted at a particular position in

the CDS if the number of reads either beginning or ending their local alignment at that position

(*nbBE*) was 20 times greater (X=20) than the median *nbBE* value within a window of 11 bases around

the position concerned, and the signal measured at that position extended to a short region of 7

bases (i.e. with a default parameter of n=3; see Methods section and Fig. S2). With parameters X=20

and n=3, 8,967 regions with lengths of 4 to 19 bp were predicted to be IEBs (Table 2). Regions of

less than seven bases in length were next to non-covered regions or regions with a measured signal

close to the end of the CDS for which signal extension was not possible. Of the 8,967 regions

predicted to be IEBs, 8,575 actually corresponded to true IEBs (i.e. method specificity = 95.63%). Of

the 8,632 regions truly containing an IEB, 8,575 were accurately predicted to be IEB regions (i.e.

sensitivity = 99.34%; Table 2). Of the 862 CDS with no intron, 763 (i.e. 88.51%) were correctly

predicted to have no IEBs. As expected, the use of a lower detection threshold (i.e. X=10) decreased

the specificity of the method and increased its sensitivity (Table 2). The regions predicted to have

no IEB (i.e. the - regions) were between 1 and 3,417 bp in length, with a mean length of 265.5 bp,

entirely consistent with the known distribution of exon lengths (see Methods section). The 153 very

small (<10 bp) - regions attested to the possible cumulative deviation of signals of more than eight

bases for a small number of IEBs. Increasing the length of the signal deviation (e.g. *n*=5 bases instead

of 3) made it possible to take this situation into account in the vast majority of cases. This slightly

improved performance but increased the length, and thus, the imprecision, of all predicted IEB

positions (Table 2).

Parameter		Regions +		Regions -		No prediction	Performance	
			Mean length		Mean length			
X	*n*	Number	[min;max] (bp)	Number	[min;max] (bp)	Number	Sensitivity	Specificity
20	3	8967	8.38 [4;19]	11849	265 [1;3417]	228	99.34	95.63
20	5	8856	12.44 [6;26]	11728	264.7 [1;3417]	228	99.88	97.01
10	3	9454	8.46 [4;26]	12254	255.8 [1;3417]	228	99.59	90.82
10	5	9306	12.55 [6;39]	12082	256.4 [1;3417]	228	99.94	92.34

Table 2: Evaluation of the IEB prediction method using pool data. The + region is a short region predicted to be an IEB.
The - region is a region with no signal (i.e. predicted to be an exon). Regions not covered by reads are regions without
prediction. Sensitivity is the percentage of regions with true IEBs correctly predicted to be + regions, and specificity is
the percentage of + regions that are true IEBs.

Discussion

Exome-targeted capture associated with next-generation sequencing has been mainly restricted to

model species for which genomic resources are available (Ng *et al.*, 2009). Transcriptome-based

exome capture has proved an efficient alternative for dealing with non-model species (e.g. Bi *et al.*,

2012), but is mostly used in phylogeny studies. Here, we propose an extension of this method with

a high capture efficiency, providing accurate estimates of the allele frequencies of SNPs for

population genomics studies. Our methodology is original because (i) it combines transcriptome-

based exome capture with a pool-based approach (pooling the tissues of numerous individuals

before DNA extraction), (ii) it does not require a genome sequence for the species itself or another

closely related species, and (iii) it does not involve the *de novo* assembly of genomic reads. This work

opens up interesting prospects for population genomics studies on non-model species because it is

cheap and less labor-intensive than other methods.

Capture efficiency

We found that capture efficiency was very high for the *H. axyridis* exome. The capture specificity

(83.2%) achieved was much higher than reported in more classical transcriptome-based studies

(range 25% - 73%), whereas capture sensitivity (93.5%) approached the highest reported values (see

Puritz & Lotterhos, 2018). Moreover, despite the differences in coverage observed between targets

and along the length of targets, reproducibility was high between libraries, whether in terms of

specificity, sensitivity or coverage. Our capture approach, based on a random choice of targets from

the *de novo* transcriptome of the species of interest itself, without the need to identify orthologous

genes in related model species, proved highly effective. This "blind" approach had only a minor

impact on capture performance: we only observed a slight decrease in coverage for small exons and

towards the ends of the exons. Similar patterns have been observed in classical transcriptome-based

exome capture in which IEBs are predicted by orthology, due to the use of less efficient probes

designed from the sequences of two consecutive exons and the absence of a probe straddling the

end of the exon and the beginning of the intron (e.g. Bi *et al.*, 2012; Neves *et al.*, 2013).

The random choice of targets was probably the cause of the slightly lower capture sensitivity

in this study than in previous studies (Bi *et al.*, 2012; Puritz & Lotterhos, 2018). Targets selected

without looking for orthology in related species are necessarily only partly known, and some may

belong to other species (e.g. symbionts or parasites) or be chimeras present in the transcriptome

used. This is a small price to pay given the efficiency of the method, a feature of considerable

interest for population genomics studies. The absence of searches for orthologous genes in related

model species prevents the analysis from focusing purely on the most conserved genes. Moreover,

our method can be applied to any species for which no closely related model species with genomic

resources is available.

SNP calling and the pool-based approach

With this method, we were able to genotype a very large number of SNPs (~300,000) over the exome

of *H. axyridis*. The study of the subset of CDS with complete genomic sequence matches further

confirmed that direct mapping onto the CDS has no major impact on SNP identification or on the

estimation of allele frequencies relative to mapping onto the genome. In addition to a few residual

paralogous sequences inherent to any approach dealing with non-model species, we observed

slightly larger numbers of SNPs called on the basis of mapping onto CDS in areas very close to the

IEBs, but the proportion of SNPs involved was minimal (0.33%).

For similar sequencing efforts, strong concordance was observed between the allele

frequencies for individual genotyping and those for the "pool-based" approach, whatever the

mapping method used. This finding demonstrates the feasibility of this approach, in situations in

which individual identification for many samples is practically impossible, technically difficult, too

costly and/or unnecessary. In population genomics studies, allele frequencies are sufficient for the

calculation of most summary statistics (e.g. nucleotide diversity, F_{ST}, site frequency spectrum) and,

thus, for addressing most of the questions posed (Schlötterer *et al.*, 2014). In the specific case of

exome sequencing, allele frequencies can be used for large population-scale screening of protein

mutations, or to address questions related to adaptation or to the evolution of the genetic load (e.g.

Gutenkunst *et al.*, 2009; Yi *et al.*, 2010; Casals *et al.*, 2013; Popp *et al.*, 2017; Castellano *et al.*, 2019).

One of the limitations of pool-seq compared to individual-based approaches is that

"populations" must be carefully defined before sequencing, as it will be difficult to detect a Wahlund

effect or an unforeseen breeding system afterwards. It is therefore important to be particularly

careful during the sampling step. A second drawback of pool-seq is that it makes it impossible to

remove SNPs that are not at Hardy-Weinberg equilibrium. Such disequilibrium may be caused by

copy number variations or paralogous sequences resulting in false-positive SNPs and local over-

coverage (Li, 2014). We show here that the application of our "maximum coverage" filter to the

pooled data made it possible to discard almost 70% of the SNPs not at equilibrium, as shown in

individuals. The extremely small remaining proportion (0.34%) of SNPs not at Hardy-Weinberg

equilibrium provides support for the feasibility of our pooling-based method. Finally, pool-seq is not

optimal for studies in which analyses rely on linkage phase information. However, a prerequisite in

this context is the availability of a genetic (or physical) map, which is not the case in most non-model

organisms.

One of the limitations of pool-seq compared to individual-based approaches is that it is not

optimal for studies in which analyses rely on linkage phase information. However, a prerequisite in

this context is the availability of a genetic (or physical) map, which is not the case in most non-model

organisms. Another weakness of pool-seq is that "populations" must be carefully defined before

sequencing, as it will be difficult to detect a Wahlund effect or an unforeseen breeding system

afterwards. It is therefore important to be particularly careful during the sampling step, and to be

as familiar as possible with the biology of the species being studied. Finally, one more drawback of

pool-seq is that it makes it impossible to remove SNPs that are not at Hardy-Weinberg equilibrium.

Such disequilibrium may be caused by copy number variations or paralogous sequences resulting in

false-positive SNPs and local over-coverage (Li, 2014). We show here that the application of our

"maximum coverage" filter to the pooled data made it possible to discard almost 70% of the SNPs

not at equilibrium, as shown in individuals. The extremely small remaining proportion (0.34%) of

SNPs not at Hardy-Weinberg equilibrium provides support for the feasibility of our pooling-based

method.

Method for predicting IEBs without read assembly

We have developed a method for detecting IEBs that is particularly useful in the case of non-model

species (or species for which genome sequences are being assembled). The proposed tool (available

from https://github.com/edeleury/IEB-finder) does not indicate the exact location of the IEBs,

instead identifying an area of a few bases (about eight, on average, in our case) predicted to contain

an IEB. This method, which requires only the genomic reads and the target CDS sequences, was

effective when evaluated on a subset of CDS for which we had a complete genomic match. Almost

all the true IEBs were identified, whereas false-positives accounted for less than 5% of the predicted

IEBs. Montes *et al.* (2013) proposed a similar approach, which they used to discard putative SNPs

found in predicted IEB areas. They obtained better genotyping results for the selected filtered SNPs

than other similar studies on non-model species. In this study, we went a step further by taking into

account the possible offset of the measured signal, and by evaluating the performance of the

approach on a set of transcripts for which the true intron-exon structure was known. We aimed to

present evidence that this approach is promising, although further improvement is required,

together with evaluation on larger dataset and at various level of coverage. A method for choosing

an appropriate value for parameter X for the correct identification of significant increases in *nbBE*

depending on local coverage must also be developed.

General recommendations, conclusion and perspectives

The combination of approaches proposed here — i.e. exome transcriptome-based capture, random

choice of targets, and direct mapping of reads onto targeted CDS sequences, coupled with a pool-

seq approach before DNA extraction — significantly decreases costs, the laboratory time required

and the complexity of the sequencing analyses required to obtain a large number of variants in

exome-scale coding regions for non-model species. However, direct mapping onto target sequences

has two disadvantages. First, it provides no information on flanking regions (including regulatory

regions), and it is therefore not possible to genotype SNPs in these areas. Nevertheless, detecting

SNP polymorphism at exonic level would be sufficient in many cases, particularly when the in-depth

analysis of a large number of synonymous and non-synonymous mutations is informative enough

to address the question posed. It is still feasible to assemble the genomic reads in situations in which

flanking regions appear to be of interest. Second, direct mapping onto CDS results in small biases in

the detection of SNPs in the vicinity of IEBs. As a means of limiting the risk of detecting false SNPs

further, we therefore suggest using the IEB detection method proposed here, followed by the

elimination of any SNPs found in the predicted IEB regions and in short regions ($\leq$ 20 bp) surrounded

by two predicted IEB regions. Using this approach on our data with the parameters X=20 and n=3,

we retained a final list of 296,736 SNPs. A large proportion of the 3,300 SNPs excluded (408 SNPs,

i.e. 12.36%) were not at Hardy Weinberg equilibrium in individuals, confirming the relevance of this

filter.

The allele frequencies measured for the pool with our method were consistent with those

obtained in the analysis of individuals. However, caution is nevertheless required with rare alleles

(i.e. SNPs with low MAFs), which may be confused with sequencing errors (Bansal, 2010). We

recommend the use of technical replicates. When possible, and depending on the biological

question addressed, the analysis could be restricted to SNPs with MAFs above a given threshold.

618   In conclusion, our study demonstrates the feasibility of obtaining, at low cost, for a non-

619 model species (i.e. without a reference genome), important and accurate information at population

620 level concerning (i) SNP polymorphism within the exome and (ii) the general structure of the exome

621 (intron-exon structure of the targeted genes). The results obtained with this method applied to the

622 Asian ladybird *Harmonia axyridis* were very promising, and we now need to investigate other

623 species with different genome sizes and complexities. Finally, the oversizing of these experiments,

624 resulting in very high levels of coverage, suggests that it should be possible to reduce costs further

625 by capturing multiple indexed pools in a single reaction.

628 ## Acknowledgments

630 We thank Stefan Toepfer for the *Harmonia axyridis* samples. We thank Thibaut Malausa, Arnaud

631 Estoup, Mathieu Gautier, Fabrice Legeai, and Bernhard Gschloessl for their helpful comments, which

632 helped us to improve this manuscript considerably. This work was funded by the INRA SPE

633 department and the INRA UMR Institut Sophia Agrobiotech. Sequencing was performed at the

634 GENOTOUL GeT platform (https://get.genotoul.fr/en/).

637 ## Author contributions

639 ED, EL and TG conceived the study. AB, ED and EL prepared the samples and the libraries. ED

640 developed the bioinformatics pipelines. ED and EL ran the analyses. ED, EL and TG wrote the paper.

Competing interests

645   The authors have no competing interests to declare.

Data accessibility

650   The exome sequence capture data reported here have been deposited in the European Nucleotide

651   Archive (accession no. PRJEB31592). The CDS sequences targeted and the description of the exon

652   positions on the subset of transcripts have been deposited at Zenodo

653   https://doi.org/10.5281/zenodo.2598388. The HaxR v1.0 genome sequence used in this study is

654   available from http://bipaa.genouest.org/sp/harmonia_axyridis/ (see Gautier *et al.*, 2018).

References

Bansal, V. 2010. A statistical method for the detection of variants from next-generation
resequencing of DNA pools. *Bioinformatics* **26**: 318–324.
Bansal, V., Tewhey, R., LeProust, E.M. & Schork, N.J. 2011. Efficient and cost effective population
resequencing by pooling and in-solution hybridization. *PLoS One* **6**: 1–6.
Benjamini, Y. & Hochberg, Y. 1995. Controlling the false discovery rate - a practical and powerful
approach to multiple testing. *J. R. Stat. Soc. Ser. B-Methodological* **57**: 289–300.
Bi, K., Linderoth, T., Vanderpool, D., Good, J.M., Nielsen, R. & Moritz, C. 2013. Unlocking the vault:
Next-generation museum population genomics. *Mol. Ecol.* **22**: 6018–6032.
Bi, K., Vanderpool, D., Singhal, S., Linderoth, T., Moritz, C. & Good, J.M. 2012. Transcriptome-based
exon capture enables highly cost-effective comparative genomic data collection at moderate
evolutionary scales. *BMC Genomics* **13**: 403.
Bolger, A.M., Lohse, M. & Usadel, B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**: 2114–2120.

Bossert, S. & Danforth, B.N. 2018. On the universality of target-enrichment baits for phylogenomic

research. *Methods Ecol. Evol.* **9**: 1453–1460.

Bragg, J.G., Potter, S., Bi, K. & Moritz, C. 2016. Exon capture phylogenomics: efficacy across scales

of divergence. *Mol. Ecol. Resour.* **16**: 1059–1068.

Burset, M. & Guigo, R. 1996. Evaluation of Gene Structure Prediction Programs. *Genomics* **34**: 353–

367.

Casals, F., Hodgkinson, A., Hussin, J., Idaghdour, Y., Bruat, V., de Maillard, T., *et al.* 2013. Whole-

Exome Sequencing Reveals a Rapid Change in the Frequency of Rare Functional Variants in a

Founding Population of Humans. *PLoS Genet.* **9**.

Castellano, D., Macià, M.C., Tataru, P., Bataillon, T. & Munch, K. 2019. Comparison of the full

distribution of fitness effects of new amino acid mutations across great apes. *Genetics* **213**:

953–966.

Castellano, S., Parra, G., Sanchez-Quinto, F.A., Racimo, F., Kuhlwilm, M., Kircher, M., *et al.* 2014.

Patterns of coding variation in the complete exomes of three Neandertals. *Proc. Natl. Acad.*

Sci. **111**: 6666–6671.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., *et al.* 2012. A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly (Austin).*

6: 80–92.

Conklin, D., Montes, I., Albaina, A. & Estonba, A. 2013. Improved conversion rates for SNP

genotyping of nonmodel organisms. *Proc. Iwbbio 2013 Int. Work. Bioinforma. Biomed. Eng.*

127–134.

Cosart, T., Beja-Pereira, A., Chen, S., Ng, S.B., Shendure, J. & Luikart, G. 2011. Exome-wide DNA

capture and next generation sequencing in domestic and wild species. *BMC Genomics* **12**:

347. BioMed Central Ltd.

Day-Williams, A.G., McLay, K., Drury, E., Edkins, S., Coffey, A.J., Palotie, A., *et al.* 2011. An

evaluation of different target enrichment methods in pooled sequencing designs for complex

disease association studies. *PLoS One* **6**: e26279.

Förster, D.W., Bull, J.K., Lenz, D., Autenrieth, M., Paijmans, J.L.A., Kraus, R.H.S., *et al.* 2018.

Targeted resequencing of coding DNA sequences for SNP discovery in nonmodel species. *Mol.*

Ecol. Resour. 1356–1373.

Gautier, M., Foucaud, J., Gharbi, K., Cezard, T., Galan, M., Loiseau, A., *et al.* 2013. Estimation of

population allele frequencies from next-generation sequencing data: pool-versus individual-

based genotyping. *Mol. Ecol.* **22**: 3766–3779.

Gautier, M., Yamaguchi, J., Foucaud, J., Loiseau, A., Ausset, A., Facon, B., *et al.* 2018. The Genomic

Basis of Color Pattern Polymorphism in the Harlequin Ladybird. *Curr. Biol.* **28**: 3296–3302.

Gouzy, J., Carrere, S. & Schiex, T. 2009. FrameDP: Sensitive peptide detection on noisy matured

sequences. *Bioinformatics* **25**: 670–671.

Graffelman, J. 2015. Exploring Diallelic Genetic Markers: The HardyWeinberg Package. *J. Stat.*

Softw. **64**: 1–23.

Gutenkunst, R.N., Hernandez, R.D., Williamson, S.H. & Bustamante, C.D. 2009. Inferring the joint

demographic history of multiple populations from multidimensional SNP frequency data.

PLoS Genet. **5**.

Heyduk, K., Trapnell, D.W., Barrett, C.F. & Leebens-Mack, J. 2016. Phylogenomic analyses of

species relationships in the genus *Sabal* (Arecaceae) using targeted sequence capture. *Biol. J.*

Linn. Soc. **117**: 106–120.

Hodges, E., Xuan, Z., Balija, V., Kramer, M., Molla, M.N., Smith, S.W., *et al.* 2007. Genome-wide *in*

situ exon capture for selective resequencing. *Nat. Genet.* **39**: 1522–1527.

Ilves, K.L. & López-Fernández, H. 2014. A targeted next-generation sequencing toolkit for exon-
based cichlid phylogenomics. *Mol. Ecol. Resour.* **14**: 802–811.

Ilves, K.L., Torti, D. & López-Fernández, H. 2018. Exon-based phylogenomics strengthens the
phylogeny of Neotropical cichlids and identifies remaining conflicting clades (Cichliformes:
Cichlidae: Cichlinae). *Mol. Phylogenet. Evol.* **118**: 232–243.

Koboldt, D.C., Zhang, Q., Larson, D.E., Shen, D., Mclellan, M.D., Lin, L., *et al.* 2012. VarScan 2 :
Somatic mutation and copy number alteration discovery in cancer by exome sequencing.
Genome Res. **22**: 568–576.

Langmead, B. & Salzberg, S.L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**:
357–359.

Li, H. 2014. Toward better understanding of artifacts in variant calling from high-coverage
samples. *Bioinformatics* **30**: 2843–2851.

Montes, I., Conklin, D., Albaina, A., Creer, S., Carvalho, G.R., Santos, M., *et al.* 2013. SNP Discovery
in European Anchovy (Engraulis encrasicolus, L) by High-Throughput Transcriptome and
Genome Sequencing. *PLoS One* **8**: e70051.

Neves, L.G., Davis, J.M., Barbazuk, W.B. & Kirst, M. 2013. Whole-exome targeted sequencing of the
uncharacterized pine genome. *Plant J.* **75**: 146–156.

Ng, S.B., Turner, E.H., Robertson, P.D., Flygare, S.D., Bigham, A.W., Lee, C., *et al.* 2009. Targeted
capture and massively parallel sequencing of 12 human exomes. *Nature* **461**: 272–276.
Nature Publishing Group.

Nicholls, J.A., Pennington, R.T., Koenen, E.J.M., Hughes, C.E., Hearn, J., Bunnefeld, L., *et al.* 2015.
Using targeted enrichment of nuclear genes to increase phylogenetic resolution in the
neotropical rain forest genus *Inga* (Leguminosae: Mimosoideae). *Front. Plant Sci.* **6**: 1–20.

Ning, Z., Cox, A.J. & Mullikin, J.C. 2001. SSAHA: a fast search method for large DNA databases. -
Abstract - UK PubMed Central. 1725–1729.

Pedersen, B.S. & Quinlan, A.R. 2018. Mosdepth: Quick coverage calculation for genomes and
exomes. *Bioinformatics* **34**: 867–868.

Popp, B., Ekici, A.B., Thiel, C.T., Hoyer, J., Wiesener, A., Kraus, C., *et al.* 2017. Exome Pool-Seq in
neurodevelopmental disorders. *Eur. J. Hum. Genet.* **25**: 1364–1376. Springer US.

Portik, D.M., Smith, L.L. & Bi, K. 2016. An evaluation of transcriptome-based exon capture for frog
phylogenomics across multiple scales of divergence (Class: Amphibia, Order: Anura). *Mol.*
Ecol. Resour. **16**: 1069–1083.

Puritz, J.B. & Lotterhos, K.E. 2018. Expressed exome capture sequencing: A method for cost-
effective exome sequencing for all organisms. *Mol. Ecol. Resour.* 1209–1222.

R Development Core Team. 2015. R: A language and environment for statistical computing. R
Foundation for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Ramos, E., Levinson, B.T., Chasnoff, S., Hughes, A., Young, A.L., Thornton, K., *et al.* 2012.
Population-based rare variant detection via pooled exome or custom hybridization capture
with or without individual indexing. *BMC Genomics* **13**: 683.

Rellstab, C., Dauphin, B., Zoller, S., Brodbeck, S. & Gugerli, F. 2019. Using transcriptome
sequencing and pooled exome capture to study local adaptation in the giga-genome of *Pinus*
cembra. *Mol. Ecol. Resour.* **19**: 536–551.

Ryu, S., Han, J., Norden-Krichmar, T.M., Schork, N.J. & Suh, Y. 2018. Effective discovery of rare
variants by pooled target capture sequencing: A comparative analysis with individually
indexed target capture sequencing. *Mutat. Res. - Fundam. Mol. Mech. Mutagen.* **809**: 24–31.
Elsevier.

Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. 2014. Sequencing pools of individuals — mining
genome-wide polymorphism data without big funding. *Nat. Rev. Genet.* **15**: 749–763. Nature
Publishing Group.
Shearer, A.E., Hildebrand, M.S., Ravi, H., Joshi, S., Guiffre, A.C., Novak, B., *et al.* 2012. Pre-capture
multiplexing improves efficiency and cost-effectiveness of targeted genomic enrichment.
BMC Genomics **13**.
Stephens, J.D., Rogers, W.L., Heyduk, K., Cruse-Sanders, J.M., Determann, R.O., Glenn, T.C., *et al.*
2015. Resolving phylogenetic relationships of the recently radiated carnivorous plant genus
Sarracenia using target enrichment. *Mol. Phylogenet. Evol.* **85**: 76–87. Elsevier Inc.
Vogel, H., Schmidtberg, H. & Vilcinskas, A. 2017. Comparative transcriptomics in three ladybird
species supports a role for immunity in invasion biology. *Dev. Comp. Immunol.* **67**: 452–456.
Elsevier Ltd.
Warr, A., Robert, C., Hume, D., Archibald, A., Deeb, N. & Watson, M. 2015. Exome Sequencing:
Current and Future Perspectives. *Genes|Genomes|Genetics* **5**: 1543–1550.
Wu, T.D. & Watanabe, C.K. 2005. GMAP: A genomic mapping and alignment program for mRNA
and EST sequences. *Bioinformatics* **21**: 1859–1875.
Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z.X.P., Pool, J.E., *et al.* 2010. Sequencing of Fifty
Human Exomes Reveals Adaptation to High Altitude. *Science (80-.).* **329**: 75–78.
783
Bansal, V., Tewhey, R., LeProust, E.M. & Schork, N.J. 2011. Efficient and cost effective population
resequencing by pooling and in-solution hybridization. *PLoS One* **6**: 1–6.
Benjamini, Y. & Hochberg, Y. 1995. Controlling the false discovery rate - a practical and powerful
approach to multiple testing. *J. R. Stat. Soc. Ser. B-Methodological* **57**: 289–300.
Bi, K., Linderoth, T., Vanderpool, D., Good, J.M., Nielsen, R. & Moritz, C. 2013. Unlocking the vault:
Next-generation museum population genomics. *Mol. Ecol.* **22**: 6018–6032.
Bi, K., Vanderpool, D., Singhal, S., Linderoth, T., Moritz, C. & Good, J.M. 2012. Transcriptome-based
exon capture enables highly cost-effective comparative genomic data collection at moderate
evolutionary scales. *BMC Genomics* **13**: 403.
Bolger, A.M., Lohse, M. & Usadel, B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**: 2114–2120.
Bossert, S. & Danforth, B.N. 2018. On the universality of target-enrichment baits for phylogenomic
research. *Methods Ecol. Evol.* **9**: 1453–1460.
Bragg, J.G., Potter, S., Bi, K. & Moritz, C. 2016. Exon capture phylogenomics: efficacy across scales
of divergence. *Mol. Ecol. Resour.* **16**: 1059–1068.
Burset, M. & Guigo, R. 1996. Evaluation of Gene Structure Prediction Programs. *Genomics* **34**: 353–
367.
Castellano, S., Parra, G., Sanchez-Quinto, F.A., Racimo, F., Kuhlwilm, M., Kircher, M., *et al.* 2014.
Patterns of coding variation in the complete exomes of three Neandertals. *Proc. Natl. Acad.*
Sci. **111**: 6666–6671.
Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., *et al.* 2012. A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly (Austin).*
6: 80–92.
Conklin, D., Montes, I., Albaina, A. & Estonba, A. 2013. Improved conversion rates for SNP
genotyping of nonmodel organisms. *Proc. Iwbbio 2013 Int. Work. Bioinforma. Biomed. Eng.*
127–134.
Cosart, T., Beja-Pereira, A., Chen, S., Ng, S.B., Shendure, J. & Luikart, G. 2011. Exome-wide DNA
capture and next generation sequencing in domestic and wild species. *BMC Genomics* **12**:

347. BioMed Central Ltd.

Day-Williams, A.G., McLay, K., Drury, E., Edkins, S., Coffey, A.J., Palotie, A., *et al.* 2011. An
evaluation of different target enrichment methods in pooled sequencing designs for complex
disease association studies. *PLoS One* **6**: e26279.

Förster, D.W., Bull, J.K., Lenz, D., Autenrieth, M., Paijmans, J.L.A., Kraus, R.H.S., *et al.* 2018.
Targeted resequencing of coding DNA sequences for SNP discovery in nonmodel species. *Mol.*
Ecol. Resour. 1356–1373.

Gautier, M., Foucaud, J., Gharbi, K., Cezard, T., Galan, M., Loiseau, A., *et al.* 2013. Estimation of
population allele frequencies from next-generation sequencing data: pool-versus individual-
based genotyping. *Mol. Ecol.* **22**: 3766–3779.

Gautier, M., Yamaguchi, J., Foucaud, J., Loiseau, A., Ausset, A., Facon, B., *et al.* 2018. The Genomic
Basis of Color Pattern Polymorphism in the Harlequin Ladybird. *Curr. Biol.* **28**: 3296–3302.

Gouzy, J., Carrere, S. & Schiex, T. 2009. FrameDP: Sensitive peptide detection on noisy matured
sequences. *Bioinformatics* **25**: 670–671.

Graffelman, J. 2015. Exploring Diallelic Genetic Markers: The HardyWeinberg Package. *J. Stat.*
Softw. **64**: 1–23.

Heyduk, K., Trapnell, D.W., Barrett, C.F. & Leebens-Mack, J. 2016. Phylogenomic analyses of
species relationships in the genus *Sabal* (Arecaceae) using targeted sequence capture. *Biol. J.*
Linn. Soc. **117**: 106–120.

Hodges, E., Xuan, Z., Balija, V., Kramer, M., Molla, M.N., Smith, S.W., *et al.* 2007. Genome-wide *in*
situ exon capture for selective resequencing. *Nat. Genet.* **39**: 1522–1527.

Ilves, K.L. & López-Fernández, H. 2014. A targeted next-generation sequencing toolkit for exon-
based cichlid phylogenomics. *Mol. Ecol. Resour.* **14**: 802–811.

Ilves, K.L., Torti, D. & López-Fernández, H. 2018. Exon-based phylogenomics strengthens the
phylogeny of Neotropical cichlids and identifies remaining conflicting clades (Cichliformes:
Cichlidae: Cichlinae). *Mol. Phylogenet. Evol.* **118**: 232–243.

Koboldt, D.C., Zhang, Q., Larson, D.E., Shen, D., Mclellan, M.D., Lin, L., *et al.* 2012. VarScan 2 :
Somatic mutation and copy number alteration discovery in cancer by exome sequencing.
Genome Res. **22**: 568–576.

Langmead, B. & Salzberg, S.L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**:
357–359.

Li, H. 2014. Toward better understanding of artifacts in variant calling from high-coverage
samples. *Bioinformatics* **30**: 2843–2851.

Montes, I., Conklin, D., Albaina, A., Creer, S., Carvalho, G.R., Santos, M., *et al.* 2013. SNP Discovery
in European Anchovy (Engraulis encrasicolus, L) by High-Throughput Transcriptome and
Genome Sequencing. *PLoS One* **8**: e70051.

Neves, L.G., Davis, J.M., Barbazuk, W.B. & Kirst, M. 2013. Whole-exome targeted sequencing of the
uncharacterized pine genome. *Plant J.* **75**: 146–156.

Ng, S.B., Turner, E.H., Robertson, P.D., Flygare, S.D., Bigham, A.W., Lee, C., *et al.* 2009. Targeted
capture and massively parallel sequencing of 12 human exomes. *Nature* **461**: 272–276.
Nature Publishing Group.

Nicholls, J.A., Pennington, R.T., Koenen, E.J.M., Hughes, C.E., Hearn, J., Bunnefeld, L., *et al.* 2015.
Using targeted enrichment of nuclear genes to increase phylogenetic resolution in the
neotropical rain forest genus *Inga* (Leguminosae: Mimosoideae). *Front. Plant Sci.* **6**: 1–20.

Ning, Z., Cox, A.J. & Mullikin, J.C. 2001. SSAHA: a fast search method for large DNA databases. -
Abstract - UK PubMed Central. 1725–1729.

Pedersen, B.S. & Quinlan, A.R. 2018. Mosdepth: Quick coverage calculation for genomes and

exomes. *Bioinformatics* **34**: 867–868.

Portik, D.M., Smith, L.L. & Bi, K. 2016. An evaluation of transcriptome-based exon capture for frog
phylogenomics across multiple scales of divergence (Class: Amphibia, Order: Anura). *Mol.*
Ecol. Resour. **16**: 1069–1083.

Puritz, J.B. & Lotterhos, K.E. 2018. Expressed exome capture sequencing: A method for cost-
effective exome sequencing for all organisms. *Mol. Ecol. Resour.* 1209–1222.

R Development Core Team. 2015. R: A language and environment for statistical computing. R
Foundation for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Ramos, E., Levinson, B.T., Chasnoff, S., Hughes, A., Young, A.L., Thornton, K., *et al.* 2012.
Population-based rare variant detection via pooled exome or custom hybridization capture
with or without individual indexing. *BMC Genomics* **13**: 683.

Rellstab, C., Dauphin, B., Zoller, S., Brodbeck, S. & Gugerli, F. 2019. Using transcriptome
sequencing and pooled exome capture to study local adaptation in the giga-genome of *Pinus*
cembra. *Mol. Ecol. Resour.* **19**: 536–551.

Ryu, S., Han, J., Norden-Krichmar, T.M., Schork, N.J. & Suh, Y. 2018. Effective discovery of rare
variants by pooled target capture sequencing: A comparative analysis with individually
indexed target capture sequencing. *Mutat. Res. - Fundam. Mol. Mech. Mutagen.* **809**: 24–31.
Elsevier.

Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. 2014. Sequencing pools of individuals — mining
genome-wide polymorphism data without big funding. *Nat. Rev. Genet.* **15**: 749–763. Nature
Publishing Group.

Shearer, A.E., Hildebrand, M.S., Ravi, H., Joshi, S., Guiffre, A.C., Novak, B., *et al.* 2012. Pre-capture
multiplexing improves efficiency and cost-effectiveness of targeted genomic enrichment.
BMC Genomics **13**.

Stephens, J.D., Rogers, W.L., Heyduk, K., Cruse-Sanders, J.M., Determann, R.O., Glenn, T.C., *et al.*
2015. Resolving phylogenetic relationships of the recently radiated carnivorous plant genus
Sarracenia using target enrichment. *Mol. Phylogenet. Evol.* **85**: 76–87. Elsevier Inc.

Vogel, H., Schmidtberg, H. & Vilcinskas, A. 2017. Comparative transcriptomics in three ladybird
species supports a role for immunity in invasion biology. *Dev. Comp. Immunol.* **67**: 452–456.
Elsevier Ltd.

Warr, A., Robert, C., Hume, D., Archibald, A., Deeb, N. & Watson, M. 2015. Exome Sequencing:
Current and Future Perspectives. *Genes|Genomes|Genetics* **5**: 1543–1550.

Wu, T.D. & Watanabe, C.K. 2005. GMAP: A genomic mapping and alignment program for mRNA
and EST sequences. *Bioinformatics* **21**: 1859–1875.

Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z.X.P., Pool, J.E., *et al.* 2010. Sequencing of Fifty
Human Exomes Reveals Adaptation to High Altitude. *Science (80-.).* **329**: 75–78.