Improving the accuracy of single turnover active fluorometry (STAF) for the estimation of phytoplankton primary productivity (PhytoPP)

Tobias G. Boatman[1], Richard J. Geider[2], Kevin Oxborough[3*]

[1]Department of Chemical Engineering, Imperial College, London, SW7 2AZ, UK

[2]School of Biological Sciences, University of Essex, Colchester, CO4 3SQ, UK

[3] Chelsea Technologies Group Ltd, 55 Central Avenue, West Molesey, Surrey, KT8 2QZ, UK

* Correspondence:
Kevin Oxborough
koxborough@chelsea.co.uk

Word count: 7389

Figures: 6

Tables: 7

Keywords: photosynthesis, primary productivity, Photosystem II, chlorophyll fluorescence,
package effect, ocean color.

Abstract

Photosystem II (PSII) photochemistry is the ultimate source of reducing power for phytoplankton
primary productivity (PhytoPP). Single turnover active chlorophyll fluorometry (STAF) provides a
non-intrusive method that has the potential to measure PhytoPP on much wider spatiotemporal scales
than is possible with more direct methods such as ^{14}C fixation and O_2 evolved through water
oxidation. Application of a STAF-derived absorption coefficient for PSII light-harvesting (a_{LHII})
provides a method for estimating PSII photochemical flux on a unit volume basis (JV_{PII}). Within this
study, we assess potential errors in the calculation of JV_{PII} arising from sources other than
photochemically active PSII complexes (baseline fluorescence) and the package effect. Although our
data show that such errors can be significant, we identify fluorescence-based correction procedures
that can be used to minimize their impact. For baseline fluorescence, the correction incorporates an
assumed consensus PSII photochemical efficiency for dark-adapted material. The error generated by
the package effect can be minimized through the ratio of variable fluorescence measured within
narrow wavebands centered at 730 nm, where the re-absorption of PSII fluorescence emission is
minimal, and at 680 nm, where re-absorption of PSII fluorescence emission is maximal. We conclude
that, with incorporation of these corrective steps, STAF can provide a reliable estimate of JV_{PII} and,
if used in conjunction with simultaneous satellite measurements of ocean color, could take us
significantly closer to achieving the objective of obtaining reliable autonomous estimates of PhytoPP.

1   Introduction

Phytoplankton contribute approximately half the photosynthesis on the planet (Field, 1998), thus
forming the base of marine food webs. Reliable assessment of Phytoplankton Primary Productivity
(PhytoPP) is crucial to an understanding of the global carbon and oxygen cycles and oceanic

ecosystem function. Consequently, PhytoPP has been recognized as an Essential Ocean Variable
(EOV) within the Global Ocean Observing System (GOOS). PhytoPP is a dynamic biological
process that responds to variability in multiple environmental drivers including light, temperature and
nutrients across spatial scales from meters to ocean basins, and time scales from minutes to tens of
years. This poses significant challenges for measuring and monitoring PhytoPP.

Historically, the most frequently employed method for assessing PhytoPP has been the fixation of
^{14}C within closed systems over several hours of incubation (Marra, 2002; Milligan et al. 2015).
Despite the widespread use of the ^{14}C method, which has led to measurements of PhytoPP by the ^{14}C
method providing the database against which remote sensing estimates of primary production are
calibrated (Bouman et al. 2018), there is considerable uncertainty in what exactly the ^{14}C method
measures and the accuracy of bottle-incubation based methods for obtaining PhytoPP in oligotrophic
ocean waters (Quay et al. 2010).

According to Marra (2002), the ^{14}C technique measures somewhere between net and gross carbon
fixation, depending on the length of the incubation. In this context, net carbon fixation is defined as
gross carbon fixation minus carbon respiratory losses and light-dependent losses due to
photorespiration and light-enhanced mitochondrial respiration (Milligan et al. 2015). Although it may
seem intuitive that short incubations should provide a good estimate of gross carbon fixation (and
closely match PhytoPP), several authors have reported that short-term ^{14}C fixation does not reliably
measure net or gross production (e.g. Halsey et al. 2013; Milligan et al. 2015). It should also be noted
that short-term, in the context of ^{14}C fixation, is several hours incubation. This clearly imposes major
limitations on the spatiotemporal scales at which PhytoPP can be assessed using this method.

Gross photosynthesis by phytoplankton is defined here as the rate at which reducing power is
generated by photosystem II (PSII) through the conversion of absorbed light energy (PSII
photochemistry). Within this study, gross photosynthesis is quantified by measuring the rate at which
O_2 is evolved through water oxidation by PSII photochemistry (Ferron et al. 2016) and is termed
PhytoGO. Although measurement of O_2 evolution provides some advantages over ^{14}C fixation, in
that both gross and net primary production can be obtained, the spatiotemporal limitations are
similar.

It is now widely accepted that active fluorometry can provide a non-intrusive method for measuring
PSII photochemistry on much wider spatiotemporal scales than either ^{14}C fixation or O_2 evolution.
Within oceanic systems, where optically thin conditions are the norm, the most appropriate form of
active fluorometry is the single turnover method (Kolber and Falkowski 1993; Kolber et al. 1998;
Moore et al. 2006; Suggett et al. 2001; Oxborough et al. 2012). One important parameter generated
by single turnover active fluorometry (STAF) is the absorption cross section of PSII photochemistry
(σ_{PII} in the dark-adapted state, σ_{PII}' in the light-adapted state, see Terminology) with units of m^2 PSII$^-$
^{1} (Kolber et al. 1998; Oxborough et al. 2012). This parameter allows for the calculation of PSII
photochemical flux through a single PSII center, as the product of σ_{PII}' and incident photon irradiance
(E, with units of photons m^{-2} s^{-1}). PSII photochemical flux has units of photons PSII^{-1} s^{-1} or
(assuming an efficiency of one stable photochemical event per photon) electrons PSII^{-1} s^{-1} (Equation
1).

$J_{PII} = \sigma_{PII}' \cdot E$ Equation 1

Both PhytoPP and PhytoGO can be reported per unit volume (SI units of C m^{-3} s^{-1} or O_2 m^{-3} s^{-1},
respectively). Given that J_{PII} provides the photochemical flux through the σ_{PII}' provided by a single

PSII, the PSII photochemical flux per unit volume (JV_{PII}, with units of electrons m^{-3} s^{-1}) can be
defined as the flux through the absorption cross section of PSII photochemistry provided by all open
PSII centers within the volume (Equation 2).

$$JV_{PII} = \sigma_{PII}' \cdot [PSII] \cdot (1 - C) \cdot E \qquad \text{Equation 2}$$

Where [PSII] is the concentration of photochemically active PSII complexes, with units of PSII m^{-3},
and (1 - C) is the proportion of these centers that are in the open state at the point of measurement
under actinic light. It follows that JV_{PII} can, in principle, provide a proxy for PhytoPP (Oxborough et
al. 2012).

An important caveat to using JV_{PII} as a proxy for PhytoPP is that there are a number of processes
operating within phytoplankton that can uncouple PhytoPP from PhytoGO and PhytoGO from PSII
photochemistry (Geider and MacIntyre 2002; Behrenfeld et al. 2004; Halsey et al. 2010; Suggett et
al. 2010; Lawrenz et al. 2013). It follows that JV_{PII} provides an upper limit for PhytoPP which is
defined by the release of each O_2 requiring a minimum of four photochemical events and each O_2
released resulting in the maximum assimilation of one CO_2.

Previous studies obtained a value for the [PSII] term within Equation 2 from discrete samples of
chlorophyll *a* by assuming that the number of PSII centers per chlorophyll *a* (n_{PSII}) is relatively
constant (Kolber & Falkowski, 1993; Suggett et al., 2001). A significant problem with this approach
is that n_{PSII} shows significant variability, both in laboratory-based cultures (Suggett et al. 2004) and
in natural phytoplankton communities (Moore et al. 2006; Suggett et al. 2006). In addition, the
derivation of nPSII requires a chlorophyll *a* extraction for each sample: a requirement that imposes
significant spatiotemporal limitations.

A STAF-based method for the determination of [PSII] was described by Oxborough at al. (2012).
This method operates on the assumption that the ratio of rate constants for PSII photochemistry (k_{PII})
and PSII fluorescence emission (k_{FII}) falls within a narrow range across all phytoplankton types. One
consequence of this assumption is illustrated by Equation 3.

$$[PSII] \propto \frac{k_{PII}}{k_{FII}} \cdot \frac{F_o}{\sigma_{PII}} \qquad \text{Equation 3}$$

Where F_o is the 'origin' of variable fluorescence from a dark-adapted sample (see Terminology). Data
from a follow-up study (Silsbe et al. 2015), were entirely consistent with Equation 3 and were used to
derive a sensor type-specific constant, termed K_a, for the FastOcean fluorometer (CTG Ltd, West
Molesey, UK). It follows that:

$$[PSII] = \frac{K_a}{E_{LED}} \cdot \frac{F_o}{\sigma_{PII}} \qquad \text{Equation 4}$$

Equation 4 incorporates both K_a and the photon irradiance of the LEDs used to generate the single
turnover (ST) pulse (E_{LED}, with units of photons m^{-2} s^{-1}). It is worth noting that K_a and [PSII] are
spectrally independent while, for a homogeneous population, F_o and σ_{PII} are expected to covary with
measurement LED intensity.

As noted by Oxborough et al. (2012), Equation 3 is only valid if a high proportion of the fluorescence
signal at F_o comes from PSII complexes that are photochemically active and in the open state. While
it is reasonable to expect that most photochemically active PSII complexes will be in the open state at
$t = 0$ during a STAF measurement, there are situations where a significant proportion of the

fluorescence signal at F_o may come from sources other than photochemically active PSII complexes.
This becomes a concern when the observed ratio of variable fluorescence (F_v) to maximum
fluorescence (F_m) from a dark-adapted sample is low: although the maximum observed F_v/F_m varies
among phytoplankton taxa, it is generally within the range of 0.5 to 0.6 for the fluorometers used
within this study.

One plausible explanation for sub-maximal F_v/F_m values is that PSII photochemistry is
downregulated by high levels of Stern-Volmer quencher within the PSII pigment matrix. As with
measurement LED intensity, F_o and σ_{PII} covary with Stern-Volmer quenching and Equation 4
remains valid. Light-dependent accumulation of Stern-Volmer quencher within the PSII pigment
matrix generates non-photochemical quenching of PSII fluorescence (NPQ) within a wide range of
phytoplankton groups (Olaizola and Yamamoto 1994; Demmig-Adams and Adams 2006; Goss and
Jakob 2010; Krause and Jahns, 2004). However, this form of quenching is generally reversed within
tens of seconds to a few minutes dark-adaptation and would therefore not be expected to significantly
decrease F_v/F_m.

A second plausible explanation for sub-maximal F_v/F_m values is that a proportion of the signal at F_o
is generated by PSII complexes that lack photochemically active reaction centers. Under the
assumption that these complexes are not energetically coupled to photochemically active PSII
complexes, their presence would increase Fo but have no impact on s_{PII}. Consequently, the value of
[PSII] generated by Equation 4 would increase in proportion to the increase in measured F_o. Within
this manuscript, the fraction of F_o that does not originate from open PSII complexes is termed
baseline fluorescence (F_b) and the fraction that does is termed baseline-corrected F_o (F_{oc}, see
Terminology).

Within Equation 2, JV_{PII} is proportional to the product of [PSII] and $(1 - C)$ during a STAF
measurement under actinic light. A value for the concentration of photochemically active PSII
centers can be generated from a STAF measurement made on a dark-adapted sample using Equation
4. The proportion of these complexes in the open state has routinely been estimated through the qP
parameter (Kolber et al. 1998) which is mathematically equivalent to the photochemical factor
(F_q'/F_v') defined by Baker and Oxborough (2004). This requires determination of F_o', using the
equation provided by Oxborough and Baker (1997) or through direct measurement after $1 - 2$ s dark-
adaptation following a STAF measurement under actinic light (Kolber et al. 1998).

As an alternative to Equation 2, Oxborough et al. (2012) include a method for calculating JV_{PII} that
does not require [PSII], $(I – C)$ or σ_{PII} (Equation 5).

$$JV_{PII} = a_{LHII} \cdot \frac{F_{q'}}{F_{m'}} \cdot E \quad \text{Equation 5}$$

Where a_{LHII} is the absorption coefficient for PSII light harvesting, with SI units of m^{-1}. A value for
a_{LHII} can be derived using Equation 6.

$$a_{LHII} = \frac{K_a}{E_{LED}} \cdot \frac{F_m - F_o}{F_m \cdot F_o} \quad \text{Equation 6}$$

The link between Equations 2 and 5 is illustrated by Equation 7 (Kolber et al. 1998) and Equation 8
(Oxborough et al. 2012) .

$$\sigma_{LHII} = \sigma_{PII} / \frac{F_v}{F_m} \qquad \text{Equation 7}$$

$a_{\mathrm{LHII}} = \sigma_{\mathrm{LHII}} \cdot [\mathrm{PSII}]$ Equation 8

The package effect is a consequence of the high concentration of chlorophyll a and other light-
absorbing pigments within phytoplankton cells. To put this in context, while the concentration of
chlorophyll a within the open ocean is often below 0.1 mg m^{-3}, the concentration within
phytoplankton cells is approximately a million times higher than this, at 0.1 kg m^{-3} (calculated from
data within Montagnes et al. 1994). It follows that while sea water with phytoplankton cells
suspended within it can be considered optically thin, the localized volume within each phytoplankton
cell is optically very thick.

Differences in the package effect due to pigment composition and morphology among species have
been identified (Berner et al., 1989; Bricaud et al. 1993; Morel and Bricaud, 1981). Even within
individual phytoplankton species, levels of pigment packing vary with eco-physiological condition
and life cycle (Berner et al. 1989; Falkowski and LaRoche, 1991). Increases in the magnitude of the
package effect will increase the absorption of photons generated by PSII fluorescence (FII) before
these photons leave the cell, and thus decrease the measured value of FII relative to PSII
photochemistry (PII). Given that a fundamental assumption of the absorption method is that the
relationship between PSII photochemistry (PII) and PSII fluorescence emission (FII) is reasonably
constant, variability in the level of package effect among samples clearly has the potential to
introduce significant errors.

The main objective for this study was to test the applicability of Equations 4 and 6. Because the
values generated by both equations are dependent on K_a, a comprehensive evaluation of the absolute
value and general applicability of K_a has been incorporated within the study. As a first step, a large
number of sample-specific values of K_a were generated by combining data from parallel STAF and
flash O$_2$ measurements from eleven phytoplankton species, grown under nutrient-replete and N-
limited conditions. This allowed an evaluation of the degree to which sub-maximal values of F_v/F_m
could be attributed to Stern-Volmer quenching or baseline fluorescence. It should be noted that the
term sample-specific K_a is used to define apparent K_a values that are not corrected for baseline
fluorescence. Each sample-specific K_a value referenced is the mean of all reps for a specific
combination of species and growth conditions (nutrient replete or N-starved).

In addition to the STAF and flash O$_2$ measurements used to generate sample-specific values of K_a,
the same measurement systems were used to run fluorescence light curves (FLCs) and oxygen light
curves (OLCs) on all eleven phytoplankton species. The data generated from these measurements
allowed for direct comparison of PhytoGO and JV$_{\mathrm{PII}}$ (from Equation 6) at multiple points through the
light curves.

This first set of experiments provided evidence for a wider range of K_a values across species and
environmental conditions than was evident in the earlier studies of Oxborough et al. (2012) and
Silsbe et al. (2015). Although the intra-species variance of K_a values (between values determined for
nutrient-replete and N-limited cultures) could confidently be linked to baseline fluorescence, the
inter-species variance was more easily explained in terms of the package effect. To test this
hypothesis, an additional set of measurements were made on 11 phytoplankton species, of which six
were common to the first set of experiments. The range of species was selected to cover a wide range
of cell sizes and optical characteristics. As before, sample-specific K_a values were generated from
parallel flash O$_2$ and STAF measurements. The STAF measurements were made using FastBallast
fluorometers (CTG Ltd, as before) fitted with narrow bandpass filters centered at 680 nm and 730
nm. These wavebands were chosen because chlorophyll a fluorescence is absorbed much more

strongly at 680 nm than at 730 nm. It follows that attenuation of fluorescence emission due to the
package effect will be much higher at 680 nm than at 730 nm and thus that variability of the package
effect among species should correlate with the ratio of fluorescence outputs measured at 730 nm and
680 nm. To allow for comparison with the existing FastOcean sensor, a third FastBallast sensor was
fitted with the bandpass filter used within FastOcean.

2   Materials and methods

2.1   Phytoplankton cultures (N-limited experiments)

Semi-continuous phytoplankton cultures were maintained and adapted to nutrient-replete conditions.
All cultures were grown in *f*/2 medium with silicates omitted where appropriate (Guillard, 1975).

The experimental work covered a period of several months. The initial work was conducted at the
University of Essex and incorporated six phytoplankton species (Table 1). Cultures were maintained
at 20 °C in a growth room (Sanyo Gallenkamp PLC, UK) and illuminated by horizontal fluorescent
tubes (Sylvania Luxline Plus FHQ49/T5/840, UK). The Light:Dark (L:D) cycle was set at 12 h:12 h.
Neutral density filters were used to generate low light and high light conditions (photon irradiances
of 30 and 300 μmol photons m^{-2} s^{-1}, respectively). 300 mL culture volumes were maintained within 1
L Duran bottles. Cultures were constantly aerated with ambient air and mixed using magnetic stirrers.

Additional experiments, incorporating the remaining five species, were conducted at CTG Ltd (as
before). Cultures were maintained as 30 mL aliquots within filter-capped tissue culture flasks (Fisher
Scientific, UK: 12034917). A growth temperature of 20 °C was maintained by placing the flasks
within a water bath (Grant SUB Aqua Pro 2 L, USA). Low light illumination (photon irradiance of
30 μmol photons m^{-2} s^{-1}) was provided from white LED arrays (Optoelectronic Manufacturing
Corporation Ltd. 1ft T5 Daylight, UK). The L:D cycle was set at 12 h:12 h.

The N-limited cultures were sub-cultured from the nutrient-replete cultures. High light-grown
cultures were used for the six species interrogated at the University of Essex. In all cases, the growth
photon irradiance of the original culture was maintained after sub-culturing. All N-limited cultures
were grown into the stationary growth phase using N-limiting *f*/2 medium before experimental
measurements were made.

2.2   Phytoplankton cultures (package effect experiments)

All package effect experiments were conducted at CTG Ltd. Cultures were maintained as 30 mL
aliquots within filter-capped tissue culture flasks (Fisher Scientific, UK: 12034917). A growth
temperature of 20 °C was maintained by placing the flasks within a water bath (Grant SUB Aqua Pro
2 L, USA). Low light illumination (photon irradiance of 30 μmol photons m^{-2} s^{-1}) was provided from
white LED arrays (Optoelectronic Manufacturing Corporation Ltd. 1ft T5 Daylight, UK). The L:D
cycle was set at 12 h:12 h.

2.3   Setup for OLCs and flash O$_2$ measurements

All OLCs and flash O2 measurements were made using an Oxygraph Plus system (Hansatech
Instruments Ltd, Norfolk, UK). The sample volume was always 1.5 mL and a sample temperature of
20 °C was maintained using a circulating water bath connected to the water jacket of the DW1
electrode chamber. The sample was mixed continuously using a magnetic flea (as supplied with the
Oxygraph Plus system). Illumination was provided from an Act2 laboratory system (CTG Ltd, as

before). The source comprised three blue Act2 LED units incorporated within an Act2 Oxygraph
head. Automated control of continuous illumination during OLCs or the delivery of saturating pulses
during flash O2 measurements was provided by an Act2 controller and the supplied Act2Run
software package.

2.4   Dilution of samples between flash O$_2$ and STAF measurements

245   The N-limited and dual waveband experiments included determination of sample-specific K_a values.
246   In all cases, the required dark STAF measurements of F_o and σ_{PII} were made after the flash O$_2$
247   measurements. In all cases, filtered medium was used to dilute the sample between Oxygraph and
248   STAF measurements.

2.5   Chlorophyll *a* extraction

In all cases, the concentrated sample used for flash O$_2$ or OLCs was normalized to the parallel dilute
STAF sample used to generate F_o and σ_{PII} or FLC data through direct measurement of chlorophyll a
concentration from both samples.

Chlorophyll was quantified by pipetting 0.5 mL of each sample into 4.5 mL of 90% acetone and
extracting overnight in a freezer at -20 °C (Welschmeyer, 1994). Samples were re-suspended and
centrifuged at approximately 12,000 x g for 10 minutes and left in the dark (~ 30 minutes) to
equilibrate to ambient temperature. Raw fluorescence from a 2 mL aliquot was measured using a
Trilogy laboratory fluorometer (Turner, UK). The chlorophyll a concentration was then calculated
from a standard curve.

2.6   Setup for dark STAF measurements and FLCs (N-limited experiments)

260   All STAF measurements for the N-limited experiments were made using a FastOcean sensor in
261   combination with an Act2 laboratory add-on (CTG Ltd, as before). The Act2 FLC head was
262   populated with blue LEDs. A water bath was used as a source for the FLC head water jacket,
263   maintaining the sample temperature at 20 °C.

2.7   Flash O$_2$ measurements for determining sample-specific K$_a$ values

The density of photochemically active PSII complexes within each sample was determined using the
flash O$_2$ method (Mauzerall & Greenbaum, 1989; Suggett et al., 2003; Silsbe et al. 2015). The
standard flash used was 120 µs duration on a 24 ms pitch at a photon irradiance of 22,000 µmol
photons m^{-2} s^{-1}.

The concentration of photochemically active PSII centers is proportional to the product of gross O$_2$
evolution rates (E_0) and the reciprocal of flash frequency (Hz). The basic theoretical assumptions are
that all photochemically active PSII centers undergo stable charge separation once during each flash,
that all photochemically active PSII centers re-open before the next flash and that four stable charge
separation events are required for each O2 released. In reality, small errors are introduced because
some centers do not undergo stable charge separation with each flash (misses) while some centers
will undergo more than one stable charge separation event with each flash (multiple hits).

The following checks were applied with all samples:

• The proportion of PSII centers closed during each flash was verified by comparison with
sequences of 120 µs flashes on a 24 ms pitch at a photon irradiance of 13,800 µmol photons
m^{-2} s^{-1}
• The default flash pitch of 24 ms was compared against 16 ms and 36 ms to assess the
accumulation of closed PSII centers, with 120 µs flashes of 22,000 µmol photons m^{-2} s^{-1}
being applied in all three cases
• Sequences of 180 and 240 µs flashes on a 24 ms pitch at a photon irradiance of 22,000 µmol
photons m^{-2} s^{-1} were applied to assess multiple hits

In all cases, a flash duration of 120 µs duration at a photon irradiance of 22,000 µmol photons m^{-2} s^{-1}
on a 24 ms pitch provided more than 96% saturation, with no evidence of a significant level of
multiple hits or the accumulation of closed PSII centers.

288 2.8 Parallel OLC and FLC measurements (N-limited experiments)

A series of parallel replicate OLC/FLC measurements were made on all nutrient-replete cultures, as
well as for the N-limited T. weissflogii culture (Table 1). The 10 to 12 light steps were identical
between the parallel OLC and FLC measurements. The sequences always started with a dark step,
with all subsequent steps lasting 180 s. Additional dark steps were incorporated after every third light
step. The dark respiration rate (Rd) was assessed before, during and after the OLC. The Rd values
measured during and after the OLC were always within 8% of the initial Rd (n = 65). The FastOcean
ST sequence comprised 100 flashlets on a 2 µs pitch. Each acquisition was an average of 16
sequences on a 100 ms pitch. The auto-LED and auto-PMT functions incorporated within the
Act2Run software were always active.

The reported gross O_2 evolution rates (E_0) were taken as the sum of measured net O_2 evolution (P_n)
and R_d (Equation 9).

$E_0 = P_n + R_d$ Equation 9

301 2.9 OLC and FLC curve fits (N-limited experiments)

OLCs and FLCs are variants of the widely used P-E (photosynthesis – photon irradiance) curve. For
OLCs, the metric for photosynthesis is the rate at which O_2 is evolved through water oxidation by
PSII. For FLCs, the metric for photosynthesis is the relative rate of PSII photochemistry, which is
assessed as the product of ϕ_{PII} and E. In the absence of baseline fluorescence (when $F_b = 0$), the
parameter F_q'/F_m' can be used to provide an estimate of ϕ_{PII}. It follows that FLC curves can be
generated by plotting E against the product of baseline-corrected F_q'/F_m' (F_q'/F_{mc}') and E.

There are three basic parameters derived from all P-E curve fits: α, E_k and P_m. The value of α
provides the initial slope of the relationship between E and P. E_k is an inflection point along the P-E
curve which is often described as the light saturation parameter (Platt and Gallegos, 1980). P_m is the
maximum rate of photosynthesis.

The FLC curve fits within this study were generated by the Act2Run software (CTG Ltd, as before).
The curve fitting routine within Act2Run is a two-step process which takes advantage of the fact that
the signal to noise within FLC data is highest during the initial part of the FLC curve. In the first step
(the Alpha phase), Equation 10 is used to generate values for α and E_k (Webb et al. 1974; Silsbe and
Kromkamp, 2012). The overall fit is an iterative process that minimizes the sum of squares of the
difference between observed and fit values. During the Alpha fit, a significant weighting on the initial

points (low actinic E values) is generated by multiplying each square of the difference by $(F_q'/F_{mc}')2$.
This approach normally generates a good fit up to E_k, but overshoots beyond this point.
Consequently, the P_m values generated by the Alpha phase are generally too high.

$$\frac{F_q'}{F_m'} = \alpha \cdot E_k \cdot \left(1 - e^{-E/E_k}\right) \cdot E^{-1} \qquad \text{Equation 10}$$

In the second step (the Beta phase) Equation 11 is used to improve the value of P_m. This step includes
a second exponential which is only applied to data points at E values above the E_k value generated by
the Alpha phase. The sum of squares of the difference between observed and fit values is not
weighted during the Beta phase. This approach forces ϕ_{PII} at E_k to be 63.2% of α.

$$\frac{F_q'}{F_m'} = \alpha \cdot E_k \cdot \left(1 - e^{-E/E_k}\right) \cdot E^{-1} \qquad \text{Equation 11}$$

The signal to noise for OLC data tends to increase with E (the opposite of what happens with FLC
data). Consequently, the fitting method used for the FLC data is not appropriate for OLC data as it is
highly dependent on having a good signal to noise during the early part of the curve. The iterative
OLC data fits used Equations 12 and 13 (Platt and Gallegos, 1980).

$$P = P_s \cdot \left(1 - e^{(-\alpha \cdot E/P_s)}\right) \cdot \left(e^{(-\beta \cdot E/P_s)}\right) \qquad \text{Equation 12}$$

$$P_m = P_s \cdot \left(\frac{\alpha}{\alpha+\beta}\right) \cdot \left(\frac{\beta}{\alpha+\beta}\right)^{\left(\frac{\beta}{\alpha}\right)} \qquad \text{Equation 13}$$

Within these equations, P_s and β improve some fits by incorporating a phase that accounts for
possible photoinactivation of PSII complexes and/or supra-optimal levels of PSII downregulation
(photoinhibition).

Direct comparison of α and β between the FLC and OLC is problematic because while $\square$ is
incorporated through the entire curve for both fits, β is only incorporated beyond E_k for the FLC and
through the entire curve for the OLC. For this reason, direct comparison between FLC and OLC data
was focused on P_m values.

2.10 Setup for the package effect STAF measurements

STAF measurements were made using three FastBallast sensors (CTG Ltd, as before). Each sensor
was fitted with one of the following bandpass filters:

730 nm bandpass, 10 nm FWHM (Edmund Optics, UK; part number 65-176)

680 nm bandpass, 10 nm FWHM (Edmund Optics, UK; part number 88-571)

682 nm bandpass, 30 nm FWHM (HORIBA Scientific, UK; part number 682AF30)

Where FWHM is Full Width at Half Maximum. The FastBallast units fitted with filters A, B and C
are, hereafter, termed B730, B680 and B682, respectively. Filter C is the standard bandpass filter
fitted within FastOcean and FastBallast fluorometers and was included here for comparison.

The emission peak for PSII fluorescence is cantered at 683 nm and is Stokes shifted from a strong
absorption peak cantered at 680 nm. Consequently, reabsorption of PSII fluorescence defined by

B680 is close to maximal and is also very high when PSII fluorescence is defined by B682. In
contrast, reabsorption of PSII fluorescence within the waveband defined by B730 is minimal.

Because the FastBallast sensor does not incorporate a water jacket, all measurements were made in a
temperature-controlled room at 20 °C. The FastBallast units were always switched on immediately
before each test and automatically powered down once a test had finished. This procedure prevented
any measurable increase in temperature within the FastBallast sample chamber during testing.

Calibration of FastBallast units does not include as assessment of ELED (Equation 4). Consequently,
there is no instrument-type specific K_a available for FastBallast. To get around this limitation, the
LED output was maintained at a constant level for all measurements. This allowed Equation 16 to be
used in place of Equation 4.

$$K_R = [PSII] \cdot \frac{J_{PII}}{F_o} \qquad \text{Equation 16}$$

Where K_R is the instrument-specific constant defined by Oxborough et al. (2012) with units of
photons m^{-3} s^{-1} and JPII is the initial rate of PSII photochemical flux during a STAF pulse, with units
of electrons $PSII^{-1}$ s^{-1}. As before, it is assumed that each photon used to drive PSII photochemistry
results in the transfer of one electron.

Samples used for the FastBallast STAF measurements were prepared by diluting a small aliquot from
the sample used for the associated flash O_2 measurement. 60 mL of cell suspension was prepared and
divided equally among the three FastBallast sample chambers. Samples were dark-adapted for at
least five minutes before measurements were made.

Each sample was run through the three FastBallast units simultaneously using the FaBtest software
supplied with the FastBallast fluorometer (CTG Ltd, as before). This test involves continuous
application of 400 µs saturating pulses at 10 Hz to a slowly stirred 20 mL sample for eight minutes.
Only 0.5 mL of the sample is illuminated by the saturating pulse at any point in time. Stirring the
sample ensures that the entire sample is interrogated during the test and prevents the accumulation of
closed PSII reaction centers. The mean value of F_v was extracted from each test result.

Following the initial measurement, a spike of Basic Blue 3 (BB3) was added to each sample to
increase the extracellular baseline fluorescence. BB3 is a water-soluble fluorescent dye, which
absorbs throughout the visible range and has a broad emission spectrum centered at approximately
690 nm (Sigma Aldrich, Saint Louis, MA, USA). As such, it can be used to simulate non-variable
fluorescence emission from any source, including CDOM and free chlorophyll a. The BB3 was
dissolved in MilliQ water to a final concentration of 118 µM. The volume of the BB3 spike was
never more than 30 µL within each 20 mL sample. After spiking with BB3, each sample was dark-
adapted for five minutes followed by a second test. In all cases, the F_b generated by addition of BB3
was at least three times the value of the original F_v and, consequently, decreased F_v/F_m by
approximately 65%.

2.11 Terminology

A structured approach has been taken in derivation of the parameters used within this manuscript. As
baseline fluorescence is central to this study, new fluorescence terms to describe baseline-corrected
values of existing fluorescence terms have been introduced. Otherwise, the parameters are structured
around root terms that are widely used within the fluorescence community.

Table T1 provides terms used to describe the fluorescence signal at any point. Table T2 provides
commonly used parameters derived from the terms in Table T1. Tables T3, T4 and T5 show the
derivation of terms used for the yields, rate constants, absorption cross sections and absorption
coefficients applied to PSII energy conversion processes. The remaining terms used are covered
within Table T6.

The root terms and subscripts provided in Tables T3 and T4, respectively, are very widely used
(examples include Butler and Kitajima, 1975; Kolber et al. 1998; Baker and Oxborough, 2004; and
Oxborough et al. 2012). These tables were constructed to introduce consistency and minimize
ambiguity: particularly with the distinction between absorption cross-sections and absorption
coefficients. It should also be noted that the 'optical absorption cross-section of PSII' and 'effective
absorption cross-section of PSII' (both unit area per photon) employed by Kolber et al. (1998) are, in
terms of usage, equivalent to the absorption cross-sections of PSII light harvesting and PSII
photochemistry (both unit area per PSII), respectively.

3   Results

3.1   Sample-specific K_a values under nutrient-replete and N-limited conditions

The sample-specific values of K_a for all nutrient-replete, low light grown cultures ranged from 7,822
m^{-1} for *C. vulgaris* to 25,743 m^{-1} for *T. pseudonana* (Figure 2A). Of the six species grown under both
low and high light, only *D. salina* exhibited a significant difference in the K_a values between light
treatments (Supplementary Table 1). In all cases, the N-limited sample-specific K_a values are
significantly lower than for the nutrient-replete samples they were sub-cultured from (Figure 2A).
These lower K_a values were matched to lower values of F_v/F_m (Supplementary Table 1).

As previously discussed, there are two mechanisms that could cause sub-maximal F_v/F_m values: dark-
persistent Stern-Volmer quenching and baseline fluorescence. Importantly, the absorption method is
insensitive to Stern-Volmer quenching while baseline fluorescence can introduce a significant error
in the calculation of JV_{PII}. In the context of these tests, the lower values of both sample-specific K_a
and F_v/F_m values observed within the N-limited cultures, when compared to the nutrient-replete
values, are entirely consistent with a baseline fluorescence-induced error being introduced by, for
example, the accumulation of photoinactivated PSII complexes. To test this possibility, Equation 14
(Oxborough, 2012) was used to derive a theoretical F_v/F_{mc} value that could be applied across all N-
limited cultures.

$$F_b = F_m - \frac{F_v}{(F_v/F_{mc})} \quad \text{Equation 14}$$

Where F_{mc} is the F_b-corrected value of the measured F_m (see Terminology). When using this
equation, F_m and F_v are measured from the sample and F_v/F_{mc} is an assumed baseline corrected value
of F_v/F_m for the photochemically active PSII complexes within the sample (see Figure 1). The single,
consensus value of F_v/F_{mc} used was generated iteratively, by minimizing the total sum of squares for
the differences in sample-specific K_a values from nutrient-replete cultures and corrected N-limited
cultures.

Applying an F_b-correction within Equation 14 brings the N-limited sample-specific K_a values for all
but one species (*T. pseudonana*) up to the point where they are not significantly different from the
matched nutrient-replete culture values and resulted in a consensus F_v/F_{mc} value of 0.518. Even with
T. pseudonana, the F_b-corrected value is much closer to the nutrient-replete value than is the

uncorrected N-limited value. The observation that the consensus F_v/F_{mc} value required for the F_b-
corrected values is slightly lower than most of the F_v/F_m values measured from the nutrient-replete
cultures (see Supplementary Table 1) may indicate that the photochemically active PSII complexes
within the N-limited cultures are operating at a slightly lower efficiency that the photochemically
active PSII complexes within the nutrient-replete cultures.

The dashed line within Figure 2A shows the default K_a value of 11,800 m^{-1} that is currently provided
for the FastOcean sensor (hereafter, K_a^{FO}). Although this value falls within the mid-range of K_a
values for the nutrient-replete cultures, there is considerable variability around this default value; for
example, K_a^{FO} is approximately 50% higher than the nutrient-replete, sample-specific K^a value for *C.*
vulgaris and less than 50% of the equivalent value for *T. pseudonana*.

3.2   Interspecific variability in K_a and Chl PSII^{-1}

A comparison between n_{PSII} and K_a is valid because they have a similar proportional impact in the
calculation of [PSII] within Equations 2 and 4, respectively. Part B of Figure 2 shows n_{PSII} values for
nutrient-replete cultures and N-limited cultures. The dashed line is at a widely used default value for
n_{PSII} of 0.002 Chl PSII-1 (Kolber & Falkowski 1993; Suggett et al. 2001). There are two noteworthy
features within this dataset. Firstly, the range of n_{PSII} values is very wide, at around 15:1: from less
than 0.0002 Chl PSII^{-1} for the N-limited T. pseudonana to more than 0.003 Chl PSII^{-1} for N-limited
E. huxleyi. Secondly, there is a lack of consistency between nPSII values from nutrient-replete
cultures and N-limited cultures: five species show higher n_{PSII} values for the N-limited cultures while
the remaining six species show lower n_{PSII} values for the N-limited cultures. Overall, these data
provide a good illustration of how an assumed value for n_{PSII} can introduce large errors in the
calculation of JV$_{PII}$, which can only be corrected through independent determination of PSII
concentration.

3.3   Comparison of OLC and FOC curves

Figure 3 shows OLC and FLC data from all eleven phytoplankton species used within this study. All
data are from the nutrient-replete, low light-grown cultures. The FLC values of PhytoGO (y-axes)
assume four electrons per O$_2$ released. Values from the STAF data were derived using either K_a^{FO} or
the sample-specific values shown in Figure 2A. Clearly, in most cases, the match between OLC and
FLC is greatly improved by using the sample-specific values of K_a in place of K_a^{FO}. The one
exception is Figure 3B (*D. salina*) where the sample-specific K_a value happens to be very close to
K_a^{FO}.

The data presented within Figure 4 have been extracted from the OLCs and FLCs within Figure 3 to
allow bulk comparison of the measured OLC and FLC PhytoGO values (A and C). Also shown is a
comparison of the P_m values (B and D) from the OLC and FLC curve fits. Values were generated
using either K_a^{FO} (A and C) or the sample-specific values of K_a (B and D).

Inevitably, the match between OLC and FLC data is improved significantly when the sample-specific
values of K_a (C and D) are used in preference to K_a^{FO} (A and B). The slope for the sample-specific K_a
data is very close to the 'ideal' of 1.0 and a high proportion of data points fall within the ±20% lines
included within the plot. In contrast, the K_a^{FO} data have a much lower slope of 0.6 and ±50% lines
are required to constrain a similar proportion of points. The sample-specific K_a values also generate a
much better correlation between the values of P_m derived from OLC and FLC curve fits than K_a^{FO} (D
and B, respectively). The slope for the sample-specific K_a data (D), at 0.778, is significantly lower

than the ideal of 1.0. This lower slope may be at least partly due to differences in the curve fits
applied to OLC and FLC data (see Methods).

3.4   The stability of F_b under actinic light

Clearly, the consensus F_v/F_{mc} (0.518) in Equation 14 generated a good match between K_a values for
all but one of the nutrient-replete and N-limited cultures in Figure 2A. In a wider context, it could
prove valid to use the consensus F_v/F_{mc} of 0.518 within Equation 14 when the measured F_v/F_m is
lower than this value and assume that F_b is zero when the measured F_v/F_m is above 0.518.

In situations where F_b is non-zero, the calculated value of a_{LHII} used within Equation 5 is decreased
while value of F_q'/F_m' used within the same equation is increased. The adjustment to a_{LHII} can largely
be justified by the fact that matched F_b and a_{LHII} values are derived from the same dark-adapted
STAF measurement. In contrast, the adjustment to F_q'/F_m' is potentially more complex, simply
because light-dependent NPQ can significantly decrease the maximum fluorescence level between
the dark-adapted F_m and light-adapted F_m' (see Introduction). Given that a proportion of F_b could be
from photoinactivated PSII complexes within the same membranes as the photochemically active
PSII complexes, it seems reasonable to consider the possibility that NPQ could also quench F_b.

To test the potential for a NPQ-dependent decrease in F_b, additional FLCs were run on the N-limited
cultures of *T. weissflogii* that had been sub-cultured from the low light-grown, nutrient-replete
cultures. The value of F_b for the original, nutrient-replete cultures was always assumed to be zero,
simply because the measured F_v/F_m from was always above 0.518. Conversely, the F_v/F_m values
measured from the N-limited cultures were always well below the consensus value, at 0.116 ± 0.006.

Figure 5A shows the maximum PhytoGO values ($PhytoGO_m$), measured as O_2-evolution (x-axis) or
calculated using the sample-specific K_a value from the nutrient-replete *T. weissflogii* of 15,868 m^{-1}.
For these values, F_b was set to zero for both the nutrient-replete cultures and the N-limited cultures.
Clearly, while there is good agreement between the measured and calculated values of $PhytoGO_m$
from the nutrient-replete cultures, most of the calculated $PhytoGO_m$ values from the N-limited
cultures are much higher than the measured values.

For Figure 5B, Equation 14 was used to generate a consensus F_v/F_{mc} specific to the N-limited
cultures. This consensus value was reached by minimizing the sum-of-squares for the regression line
through the N-limited data by allowing F_b to vary. The mean consensus F_v/F_{mc} from this fit (0.502) is
within 3% of the consensus value derived from the dark-adapted data presented in Figure 2. In
contrast, the average NPQ-dependent decrease from dark-adapted F_m to the light-adapted F_m'
measured at P_m was always more than 30% (data not shown). Consequently, these data do not imply
significant quenching of F_b between the dark-adapted state and P_m.

3.5   Dual waveband STAF measurements to correct for the package effect

We hypothesized that the variance of sample-specific K_a values within Figure 2A could be at least
partly due to variable package effect. As prevoiously noted within Materials and methods, three
FastBallast units (B730, B680 and B682) were used to measure fluorescence centered at 730 nm and
680 nm (both 10 nm FWHM) and 682 nm (30 nm FWHM), respectively.

To test the viability of a STAF-based approach to quantifying the package effect, we generated ratios
of the F_v measured by B730 as a proportion of the F_v measured by B680 ($F_v^{730/680}$) or B683 ($F_v^{730/683}$).
Within Figure 6, these values are plotted against sample-specific values of K_R (Figure 6A and D,

respectively). The F_v ratios from Figure 6A and D were used to generate F_v-derived values of K_R
(Figure 6B and E, respectively).

Calculated $K_R = \dfrac{F_v^{730/680} - Intercept}{Slope}$ Equation 15

Equation 15 provides the conversion between A and D. For the equivalent conversion between B and
E, $F_v^{730/680}$ was replaced with $F_v^{730/683}$. Slope and Intercept are the regression line values from A or D,
as appropriate. The calculated K_R values within C and F were generated by combining the + BB3
$F_v^{730/680}$ and $F_v^{730/683}$ data with the Slope and Intercept from A and D, as appropriate.

One feature that is immediately clear from these data is the much tighter grouping of points along the
regression lines for the $F_v^{730/680}$ data (A to C) than the $F_v^{730/683}$ data (D to F). This indicates that the 30
nm FWHM of the 682 nm bandpass filter is too broad to adequately isolate the fluorescence
generated close the 680 nm absorption peak and, consequently, that the 10 nm FWHM 680 nm
bandpass filter is the better choice for these measurements.

All 11 species used within the package effect tests were grown under nutrient-replete conditions and
exhibited F_v/F_m values that were above the consensus value of 0.518 generated from the first part of
this study. The addition of BB3 to each sample within the package effect tests was to simulate the
lower F_v/F_m values that are frequently observed under conditions of stress. The expectation was that
fluorescence from the added BB3 would increase F_b but have minimal impact F_v and, as a
consequence, that the slope of the relationship between calculated and measured K_R values would not
be significantly affected by a BB3-dependent increase in F_b. The absence of significant changes in
slope between B and C and E and F are consistent with this expectation.

3.6   Discussion

The absorption method described by Oxborough et al. (2012) provides a method for estimating
PhytoGO and PhytoPP on much wider spatiotemporal scales than O_2 evolution or ^{14}C fixation,
respectively, through determination of JV_{PII}. This study was undertaken to assess the extent to which
baseline fluorescence and the package effect could introduce errors into the calculation of JV_{PII}
(Equation 5).

With regard to baseline fluorescence (F_b), the underlying question was whether sub-maximal dark-
adapted value of F_v/F_m could be attributed to F_b or downregulation of PSII photochemistry by dark-
persistent Stern-Volmer quenching or some combination of the two. The data presented within Figure
2A provides strong evidence that, for the examples presented within this study, F_b is by far the
dominant contributor to sub-maximal F_v/F_m values. Although this interpretation may not hold for all
phytoplankton species and environmental conditions, this study provides a straightforward, practical
approach to addressing the question of how universally valid an F_b correction to low sub-maximal
dark-adapted F_v/F_m values might be.

We conclude that no correction for baseline fluorescence should be applied when the dark-adapted
F_v/F_m is above a certain consensus value. In situations where the dark-adapted F_v/F_m is below this
consensus value, Equation 14 should be used to calculate a value for F_b. From the data presented
here, a consensus value (F_v/F_{mc}) of between 0.50 and 0.52 seems an appropriate default value.

Clearly, the value of F_b generated by Equation 14 is dependent on a STAF measurement made on a
dark-adapted sample. The data presented in Figure 5 indicate that, for this specific example at least,
the dark-adapted F_b could be applied at the other end of the FLC scale, to correct the value of P_m.

With regard to the package effect, the wide range of K_a values within Figure 2A is entirely consistent
with a significant proportion of the fluorescence emitted from functional PSII complexes being
reabsorbed through this process. This interpretation is clearly supported by data presented in Figure
3, where use of the sample-specific K_a value in place of K_a^{FO} provides a much stronger match
between the FLC and OLC data. The dual waveband data presented in Figure 6 provide strong
evidence that the package effect-induced error could be decreased significantly through incorporation
of a $F_v^{730/680}$-derived correction factor applied to a default instrument-type specific K_a value such as
K_a^{FO}. From a practical point of view, routine implementation of this correction step will require either
two detectors with different filters or a single detector with switchable filtering. On balance, the latter
option is likely to prove more cost-effective and easier to calibrate.

Overall, the conclusions reached can be summarized by Equation 16

$$JV_{PII} = \frac{K_a^{TS} \cdot R_{PE}}{E_{LED}} \cdot \frac{F_{mc} - F_{oc}}{F_{mc} \cdot F_{oc}} \cdot \frac{F_{q'}}{F_{mc'}} \cdot E \qquad \text{Equation 16}$$

where K_a^{TS} is the instrument type-specific K_a value and R_{PE} is a dimensionless sample-specific
correction factor. All other terms are as before or are defined in Terminology.

For the species and conditions examined in this paper, the data presented provide strong evidence
that baseline correction and package effect correction can increase the accuracy of estimates of
PhytoGO from STAF. We anticipate that development and deployment of STAF instrumentation that
will allow Equation 16 to be applied will take us significantly closer to achieving the objective of
obtaining reliable autonomous estimates of PhytoGO. Such measurements, if used in conjunction
with simultaneous satellite measurements of ocean color, will likely lead to improved estimates of
local, regional or global pelagic PhytoPP.

3.7   Acknowledgements

The authors wish to thank Tania Cresswell-Maynard, James Fox and Philipp Siegel (University of
Essex, UK) for supplying the phytoplankton cultures. We would also like to thank Hoi Ga Chan
(University of Princeton, US) and Mark Moore and Anna Hickman (Southampton University, UK)
for helpful discussions. RJG acknowledges support from NERC grant NE/P002374/1.

3.8   Author contributions statement

KO conceived of the study and developed the software required to conduct the experiments. The
initial baseline experiments were designed by KO, RG and TB. All of the baseline experiments were
conducted by TB who also processed the primary data. The dual waveband experiments for
assessment of the package effect were conceived of by KO and conducted by TB. Package effect data
were processed by TB and jointly analyzed by KO and TB. Figure 1 was produced by KO, all
remaining Figures were produced by TB. The initial draft of the main text was produced by KO.
Iterations of the manuscript were implemented by KO, TB and RG. The submitted version of the
manuscript is approved for publication by KO, TB and RG.

3.9   Conflict of interest statement

The authors (KO, TB and RG) declare that the research was conducted in the absence of any
commercial or financial relationships that could be construed as a potential conflict of interest.

3.10 Contribution to the field statement

Phytoplankton photosynthesis is responsible for approximately half of the carbon fixed on the planet.
As a process, photosynthesis is responsive to variability in multiple environmental drivers including
light, temperature and nutrients across spatial scales from meters to ocean basins, and time scales
from minutes to tens of years. This poses significant challenges for measurement and monitoring.
While direct measurement of the carbon fixed by photosynthesis can only be applied on very limited
spatial and temporal scales, active chlorophyll fluorescence has enormous potential for the accurate
measurement of phytoplankton photochemistry, which provides the reducing power for carbon
fixation, on much wider spatiotemporal scales and with much lower operational costs. This study
identifies practical measures that can be taken to improve the accuracy of such measurements. We
are confident that these measures will have minimal impact on the frequency at which phytoplankton
photochemistry is assessed and that they will be suitable for application on autonomous measurement
platforms.

3.11 References

Baker NR, Oxborough K (2004). Chlorophyll Fluorescence as a Probe of Photosynthetic
Productivity. p. 65-82 In: G. C. Papageorgiou and Govindjee [eds.], Chlorophyll *a* Fluorescence: A
Signature of Photosynthesis. Springer

Behrenfeld MJ, Prasil O, Babin M, Bruyant F (2004). In search of a physiological basis for
covariations in light-saturated photosynthesis. Journal of Phycology 40(1): 4-25

Berner T, Dubinsky Z, Wyman K ,Falkoswki PG (1989). Photoadaptation and the "package effect" in
Dunaliella tertiolecta (*Chloropohyceae*). J. Phycol. 25: 70-78

Bouman HA et al. (2018). Photosynthesis-irradiance parameters of marine phytoplankton: synthesis
of a global data set. Earth System Science Data 10: 251-266 DOI: 10.5194/essd-10-251-2018

Bricaud A, Morel A, Prieur L (1983). Optical efficiency factors of some phytoplankters. Limnol.
Oceanogr. 28: 816–832

Butler WL, Kitajima M. (1975). Fluorescence quenching in photosystem II of chloroplasts.
Biochimica et Biophysica Acta 376: 116-125

Demmig-Adams B, Adams W (2006). Photoprotection in an ecological context: the remarkable
complexity of thermal energy dissipation. New Phytol 172: 11−21

Falkowski PG, LaRoche J (1991). Acclimation to spectral irradiance in algae. Journal of Phycology
27: 8-14

Ferrón S, del Valle A, Björkman KM, Church MJ, Karl DM (2016). Application of membrane inlet
mass spectrometry to measure aquatic gross primary production by the ^{18}O *in vitro* method.
Limnology and Oceanography 14: 610-622

Geider RJ, MacIntyre HL. (2002). Physiology and biochemistry of photosynthesis and algal carbon
acquisition. Phytoplankton productivity: Carbon assimilation in marine and freshwater ecosystems:
44-77

Goss R, Jakob T (2010). Regulation and function of xanthophyll cycle-dependent photoprotection in
algae. Photosynth Res 106: 103−122

Guillard RR (1975). Culture of phytoplankton for feeding marine invertebrates. Culture of Marine
Invertebrate Animals: Springer, 29-60

Halsey KH, Milligan AJ, Behrenfeld MJ (2010). Physiological optimization underlies growth rate-
independent chlorophyll-specific gross and net primary production. Photosynthesis research 103(2):
125-137

Kolber Z, Falkowski PG (1993). Use of active fluorescence to estimate phytoplankton photosynthesis
in situ. Limnology and Oceanography 38(8): 1646-1665

Kolber ZS, Prášil O, Falkowski PG (1998). Measurements of variable chlorophyll fluorescence using
fast repetition rate techniques: defining methodology and experimental protocols. Biochimica et
Biophysica Acta (BBA)-Bioenergetics 1367(1): 88-106

Krause GH, Jahns P (2004). Non-photochemical energy dissipation determined by chlorophyll
fluorescence quenching: characterization and function, p. 463-495. In G. C. Papageorgiou and
Govindjee [eds.], Chlorophyll a fluorescence: A signature of photosynthesis. Springer

Lawrenz E, Silsbe G, Capuzzo E, Ylöstalo P, Forster RM, Simis SG, Prášil O, Kromkamp JC,
Hickman AE, Moore CM (2013). Predicting the electron requirement for carbon fixation in seas and
oceans. PLoS One 8(3): e58137

Marra J (2002). Approaches to the measurement of plankton production. In: Williams, PJLB,
Thomas, DN and Reynolds CS (eds), Phytoplankton Productivity. Blackwell Science Ltd, Oxford,
pp. 78–108

Mauzerall D, Greenbaum NL (1989). The absolute size of a photosynthetic unit. Biochimica et
Biophysica Acta (BBA)-Bioenergetics 974(2): 119-140

Milligan AJ, Halsey, KH, Behrenfeld MJ (2015). Advancing interpretations of [14]C-uptake
measurements in the context of phytoplankton physiology and ecology. Journal of Plankton
Research: 37(4): 692–698

Montagnes DJS, Berges JA, Harrison PJ, Taylor FJR (1994). Estimating carbon, nitrogen, protein,
and chlorophyll a from volume in marine phytoplankton. Limnology and Oceanography 39(5) 1044-
1060

Moore CM, Mills MM, Milne A, Langlois R, Achterberg EP, Lochte K, Geider RJ, La Roche J
(2006). Iron limits primary productivity during spring bloom development in the central North
Atlantic. Global Change Biology 12(4): 626-634

Morel A, Bricaud A (1981). Theoretical results concerning light absorption in a discrete medium, and
application to specific absorption of phytoplankton. Deep-Sea Research 28A: 1375-1393

Olaizola M, Yamamoto HY (1994). Short-term response of the diadinoxanthin cycle and
fluorescence yield to high irradiance in Chaetocerus muelleri (Bacillariophyceae). Journal of
Phycology 30: 606−612

Oxborough K, Moore CM, Suggett DJ, Lawson T, Chan HG, Geider RJ. (2012). Direct estimation of
functional PSII reaction center concentration and PSII electron flux on a volume basis: a new
approach to the analysis of Fast Repetition Rate fluorometry (FRRf) data. Limnology and
Oceanography: Methods 10(3): 142-154

Platt T, Gallegos CL. (1980). Modelling primary production. Primary productivity in the sea:
Springer, 339-362

Quay PD, Peacock C, Björkman K, Karl DM (2010). Measuring primary production rates in the
ocean: Enigmatic results between incubation and non-incubation methods at Station ALOHA,Global
Biogeo-chem. Cycles 24 GB3014 doi:10.1029/2009GB003665

Silsbe GM, Kromkamp JC (2012). Modeling the irradiance dependency of the quantum efficiency of
photosynthesis. Limnology and Oceanography: Methods 10: 645 - 652

Silsbe GM, Oxborough K, Suggett DJ, Forster RM, Ihnken S, Komárek O, Lawrenz E, Prášil O,
Röttgers R, Šicner M (2015). Toward autonomous measurements of photosynthetic electron transport
rates: An evaluation of active fluorescence-based measurements of photochemistry. Limnology and
Oceanography: Methods 13(3): 138-155

Suggett D, Kraay G, Holligan P, Davey M, Aiken J, Geider R (2001). Assessment of photosynthesis
in a spring cyanobacterial bloom by use of a fast repetition rate fluorometer. Limnology and
Oceanography 46(4): 802-810

Suggett DJ, Maberly SC, Geider RJ (2006). Gross photosynthesis and lake community metabolism
during the spring phytoplankton bloom. Limnology and Oceanography 51(5): 2064-2076

Suggett DJ, MacIntyre HL, Geider RJ (2004). Evaluation of biophysical and optical determinations
of light absorption by photosystem II in phytoplankton. Limnology and Oceanography: Methods 2:
316-332

Suggett DJ, Moore CM, Geider RJ (2010). Estimating aquatic productivity from active fluorescence
measurements. Chlorophyll *a* fluorescence in aquatic sciences: methods and applications: Springer,
103-127

Suggett DJ, Oxborough K, Baker NR, MacIntyre HL, Kana TM, Geider RJ (2003). Fast repetition
rate and pulse amplitude modulation chlorophyll *a* fluorescence measurements for assessment of
photosynthetic electron transport in marine phytoplankton. European Journal of Phycology 38(4):
371-384

Webb WL, Newton M, Starr D (1974). Carbon dioxide exchange of *Alnus rubra*. Oecologia 17(4):
281-291

Welschmeyer NA (1994). Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and
pheopigments. Limnology and Oceanography 39(8): 1985-1992

Table 1. List of cultures used within each experiment. H = High Light, L = Low Light, N = N-limited.
UoE = University of Essex. (* = simultaneous N-limited OLC/FLC measurements made on *T.*
weissflogii cultures ($n = 10$).

Algal species (Symbol used within figures)	Clone	Site(s)	Media	F_b				Package effect
				Flash O_2			OLC	STAF/
				H	L	N	FLC	flash O_2
Calcidiscus leptoporus (*C. l*)	RCC1159	CTG	f/2	-	-	-	-	4
Chlorella vulgaris (*C. v*)	CCAP211 /12	CTG	BG11	-	8	4	5	4
Coccolithus pelagicus (*C. p*)	PCC182	CTG	f/2 (+Si)	-	8	4	5	-
Coscinodiscus granii (*C. g*)	CCAP1013 /10	CTG	f/2 (+Si)	-	-	-	-	4
Coscinodiscus sp. (*C.* sp.)	CCAP1013 /11	CTG	f/2 (+Si)	-	-	-	-	4
Dunaliella salina (*D. s*)	CCAP19 /18	UoE CTG	f/2	8	2	4	5	-
Dunaliella tertiolecta (*D. t*)	CCAP1320	UoE CTG	f/2	6	2	4	5	4
Emiliania huxleyi (*E. h*)	CCMP1516	UoE CTG	f/2	8	2	4	5	4
Isochrysis galbana (*I. g*)	CCMP1323	CTG	f/2	-	8	4	5	4
Pseudo-nitzschia fraudulenta (*P-n. f*)	CCAP1061 /46	CTG	f/2 (+Si)	-	-	-	-	4
Pycnococcus provasolii (*P. p*)	CCMP1199	CTG	f/2	-	8	4	5	4
Phaeodactylum tricornutum (*P. t*)	CCMP2561	UoE CTG	f/2 (+Si)	8	2	4	5	4
Thalassiosira pseudonana (*T. p*)	CCMP1335	CTG	f/2 (+Si)	-	8	4	5	-
Thalassiosira punctigera (*C. p*)	CCAP1085 /19	UoE CTG	f/2 (+Si)	8	2	4	5	-
Thalassiosira rotula (*T. r*)	CCAP1085 /20	CTG	f/2 (+Si)	-	-	-	-	4
Thalassiosira weissflogii (*T. w*)	CCMP1051	UoE/ CTG	f/2 (+Si)	8	2	4	5 (10*)	-

Table 2. The maximum phytoplankton gross photosynthesis rates (PhytoGO$_m$) from simultaneous OLC and FLC measurements of the 11 nutrient-replete phytoplankton cultures measured in Experiment 1. PhytoGO from the FLC data was calculated using K_a^{FO} (11,800 m^{-1}) and a sample-specific (K_a^S) values. Differences between OLC and FLC data was tested by a series of parametric One-Way ANOVA tests with a post hoc Tukey test (One-way ANOVA, Tukey post hoc test; $P < 0.05$). Letters show the significant differences between the maximum gross photosynthesis rate (PhytoGP$_m$) (O$_2$ RCII^{-1} s^{-1}); where [B] is significantly greater than [A], and [C] is significantly greater than [A] and [B].

Algal Species	OLC		K_a^{FO}		K_a^S	
C. vulgaris	17.9 (1.4)		28.0 (3.6)		18.6 (2.4)	
C. pelagicus	50.6 (2.6)	[B]	24.6 (2.5)	[A]	41.4 (4.3)	[B]
D. salina	20.3 (1.1)	[B]	16.2 (0.4)	[A]	16.6 (0.5)	[A]
D. tertiolecta	49.3 (2.0)	[B]	37.2 (1.9)	[A]	48.7 (2.4)	[B]
E. huxleyi	19.4 (2.1)	[B]	9.4 (1.1)	[A]	16.5 (2.0)	[B]
I. galbana	28.6 (0.8)	[B]	20.6 (0.4)	[A]	27.3 (0.5)	[B]
P. provasolii	28.6 (2.2)		25.4 (2.5)		30.2 (2.9)	
P. tricornutum	25.5 (0.6)	[B]	15.6 (0.4)	[A]	28.9 (0.7)	[C]
T. pseudonana	29.1 (2.2)	[B]	13.0 (0.9)	[A]	28.3 (2.1)	[B]
T. punctigera	34.1 (2.2)	[B]	16.7 (1.1)	[A]	30.9 (2.1)	[B]
T. weissflogii	44.8 (4.7)	[B]	27.5 (1.2)	[A]	37.0 (1.6)	

715   Table T1. The o, m and v subscripts define the origin (of variable fluorescence), maximum
716   fluorescence and variable fluorescence, respectively. The q subscript defines the proportion of
717   variable fluorescence that is quenched by PSII photochemistry. The b subscript defines baseline
718   fluorescence, which is assumed to contribute equally to F_o, F_m and F. In the interest of readability,
719   only dark-adapted values have been included in the Measurement or derivation column.

Dark term	Light term	Measurement or derivation
F_o	F_o'	Extrapolation to t = 0 from a ST pulse
F_m	F_m'	At the asymptote of a ST pulse
F_v	F_v'	$F_m - F_o$
F	F'	Any point between F_o and F_m
F_q	F_q'	$F_m - F$
F_b	F_b'	Fluorescence signal not attributable to functional PSII centers
F_b	F_b'	Proportion of the measured F_o that does not arise from functional PSII centers in the open state
F_{oc}	F_{oc}'	The baseline subtracted value of F_o such that $F_{oc} = F_o - F_b$
F_{mc}	F_{mc}'	The baseline subtracted value of F_m such that $F_{mc} = F_m - F_b$
F_c	F_c'	The baseline subtracted value of F such that $F_c = F - F_b$

720
721   Table T2. Fluorescence parameters derived using the terms within Table T1.

Dark parameter	Light parameter	Interpretation
F_v/F_m	F_v'/F_m'	Provides an estimate of PSII photochemical efficiency (ϕ_{PII}) when all PSII centers are in the open state and $F_b = 0$
F_v/F_{mc}	F_v'/F_{mc}'	Provides a baseline-corrected estimate of PSII photochemical efficiency (ϕ_{PII}) when all PSII centers are in the open state
F_q/F_m	F_q'/F_m'	Provides an estimate of PSII photochemical efficiency (ϕ_{PII}) when some centers are closed and $F_b = 0$
F_q/F_{mc}	F_q'/F_{mc}'	Provides a baseline-corrected estimate of PSII photochemical efficiency (ϕ_{PII}) when some centers are closed
F_q/F_v	F_q'/F_v'	Provides a value for the PSII photochemical factor
F_o/F_v	F_o'/F_v'	Provides an estimate of Stern-Volmer quenching within the PSII pigment matrix, normalized to PSII photochemistry (only valid when $F_b = 0$)
F_{oc}/F_v	F_{oc}'/F_v'	Provides a baseline-corrected estimate of Stern-Volmer quenching within the PSII pigment matrix, normalized to PSII photochemistry

722
723

724 Table T3. Root terms used in the derivation of parameter 'x' within Table T5.

Term	Meaning	Units
ϕ_x	Yield	dimensionless
k_x	Rate constant	photons s^{-1}
σ_x	Absorption cross-section	m^{-2} PSII^{-1}
a_x	Absorption coefficient	m^{-1}

725
726 Table T4. Subscripts used for derivation of parameters within Table T5.

Term	Usage
II	Photosystem II (PSII)
LH	Light-harvesting
P	Photochemistry
F	Fluorescence
D	Non-radiative decay

727
728 Table T5. Parameters derived from the root terms and subscripts within Tables T3 and T4, respectively.
729 Empty fields within the Light term column indicate an assumed lack of change for these quantities
730 between the dark and light-adapted states.

Dark term	Light term	Definition	SI units
ϕ_{PII}	ϕ_{PII}'	Yield of PSII photochemistry	dimensionless
ϕ_{FII}	ϕ_{FII}'	Yield of PSII fluorescence	dimensionless
ϕ_{DII}	ϕ_{DII}'	Yield of non-radiative decay within PSII	dimensionless
k_{PII}		Rate constant for photochemistry at PSII	photons s^{-1}
k_{FII}		Rate constant for fluorescence emission from PSII	photons s^{-1}
k_{DII}	k_{DII}'	Rate constant for non-radiative decay within PSII	photons s^{-1}
σ_{LHII}		Absorption cross-section of PSII light harvesting	m^2 PSII^{-1}
σ_{PII}	σ_{PII}'	Absorption cross-section of PSII photochemistry	m^2 PSII^{-1}
σ_{FII}	σ_{FII}'	Absorption cross-section of PSII fluorescence	m^2 PSII^{-1}
σ_{DII}	σ_{DII}'	Absorption cross-section of PSII non-radiative	m^2 PSII^{-1}
a_{LHII}		Absorption coefficient of PSII light harvesting	m^{-1}
a_{PII}	a_{PII}'	Absorption coefficient of PSII photochemistry	m^{-1}
a_{FII}	a_{FII}'	Absorption coefficient of PSII fluorescence	m^{-1}
a_{DII}	a_{DII}'	Absorption coefficient of PSII non-radiative decay	m^{-1}

731
732

Figure 1. STAF measurements from *E. huxleyi* illustrating the concept of baseline fluorescence (F_b).
(A) is from a nutrient-replete culture in log-growth phase. (B) is from a N-limited culture. Two
plausible explanations for the lower F_v/F_m in (B) are considered within the main text. The first
assumes that F_b is zero, as indicated by the black text in (B) and the second assumes that F_b has a
value that accounts for the entire difference in F_v/F_m between (A) and (B). All terms used are
described within Terminology.

Figure 2. Variability in K_a (A) and n_{PSII} (B) values measured across a range of phytoplankton species. In (A), the consensus F_v/F_{mc} value of 0.518 (see main text) was used to calculated F_b for each culture when the measured F_v/F_m was lower than 0.518. F_b was subtracted from the measured F_o and F_m. The dashed line represents K_a^{FO} (11,800 m^{-1}). Significant differences were tested by a series of parametric t-tests (t-test; $P < 0.05$); however, if normality was not achieved after data transformation a Mann-Whitney Rank Sum test was performed. All nutrient-replete and uncorrected N-limited K_a values were significantly different. In (B), the number of PSII reaction centers per chlorophyll a molecule was calculated from flash O_2 measurements and chlorophyll a extractions. More details are provided within Materials and methods.

Figure 3. A representative example of simultaneous oxygen light curve (OLC) and fluorescence light curve (FLC) measurements made across all phytoplankton species. The OLC and FLC measurements were made on cultures acclimated to ambient temperature (~ 20 °C) and low-light (LL = 30 µmol photons m^{-2} s^{-1}). FLC data were standardized to equivalent units of O$_2$, with both OLC and FLC data normalized to a derived concentration of functional PSII complexes (O$_2$ PSII^{-1} s^{-1}). FLC data were derived using K$_a^{FO}$ (11,800 m^{-1}) or a sample-specific (K$_a^S$) value. The solid lines represent the P-E curve fits (color-coded to match the data points). Panel (A) = *D. tertiolecta*; (B) = *D. salina*; (C) = *P. provasolii*; (D) = *C. vulgaris*; (E) = *P. tricornutum*; (F) = *T. pseudonana*; (G) = *T. punctigera*; (H) = *T. weissflogii*; (I) = *C. pelagicus*; (J) = *E. huxleyi*; (K) = *I. galbana*. Replicates from each species (*n* = 5) are presented in Supplementary Figures 3 to 13.

Figure 4. The relationship between the entire photosynthesis-photon irradiance (P-E) curve of PhytoGO, (A) and (C), and the maximum PhytoGO (PhytoGO$_m$) from simultaneous OLC and FLC measurements, (B) and (D). FLC data were standardized to equivalent units of O$_2$, with both OLC and FLC data normalized to a derived concentration of functional PSII complexes (O$_2$ PSII^{-1} s^{-1}). Within (A) and (B), FLC data were derived using K$_a^{FO}$ (11,800 m^{-1}). Within (B) and (D), sample-specific values of K$_a$ were used (see Materials and methods). Each species consisted of 5 biological replicates. The dashed line represents a 1:1 line, while the solid line is the linear regression used to generate r^2, slope and intercept values. A key for the symbols in (A) is incorporated within Table 1.

Figure 5. The relationship between simultaneous OLC and FLC measurements of maximum
PhytoGO (PhytoGO$_m$) within N-limited ($n = 10$) and nutrient replete ($n = 5$) *T. weissflogii* cultures.
The sample-specific K$_a$ value from the nutrient-replete cultures was applied throughout. In (A), no F$_b$
correction was applied. In (B), the N-limited values were F$_b$-corrected by applying a consensus
F$_v$/F$_{mc}$ value of 0.502. Further details are provided within Materials and methods.

Figure 6. The relationship between dual wavelength STAF measurements and the sample-specific K_a values using the flash O_2 method. The leftmost panels show the ratio of F_v measured by B730 to F_v measured by B680 (A) and F_v measured by B730 to F_v measured by B682 (D) against the measured K_a value from parallel measurements of flash O_2. The middle panels, (B) and (E) show the relationship between flash O_2-derived K_a and dual waveband-derived K_a values. The dual waveband K_a values within these panels were derived using the slopes and offsets reported in (A) and (D), as appropriate. The rightmost panels, (C) and (E), show the same relationships as reported in (B) and (E), respectively, but in the presence of extracellular baseline fluorescence (F_b) generated by a spike of BB3 (see Materials and methods). The dashed lines in each panel represents a 1:1 slope, the solid black line is the linear regression. The solid red lines represent ± 20% of the regression values.