Development of senescence biomarkers in the common woodlouse

Charlotte DEPEUX[1, 2] [*], Ascel SAMBA-LOUAKA[1], Thomas BECKING[1], Christine BRAQUART-VARNIER[1], Jérôme MOREAU[3], Jean-François LEMAÎTRE[2], Tiffany LAVERRE[1], Hélène PAUHLAC[1], François-Xavier DECHAUME-MONCHARMONT[3], Jean-Michel GAILLARD[2], Sophie BELTRAN-BECH[1]

[1] Université de Poitiers, Laboratoire Ecologie et Biologie des interactions, UMR CNRS 7267, 5 rue Albert Turpin, TSA 51106 86073 POITIERS Cedex 9, France.

[2] Université de Lyon, F-69000, Lyon; Université Lyon 1; CNRS, UMR5558, Laboratoire de Biométrie et Biologie Evolutive, F-69622, Villeurbanne, France

[3] Biogéosciences, UMR 6282 CNRS, Université Bourgogne Franche-Comté, 6 Boulevard Gabriel, 21000 Dijon, France.

* Corresponding author: charlotte.depeux@gmail.com

Declarations of interest: none

Abstract

During their lives, organisms accumulate, damages in their cells leading to a process called cellular senescence. Senescent cells display many cellular modifications that can be assessed through different biomarkers. Senescence biomarkers are highly studied in humans and are particularly useful to understand the processes involved in age-related diseases. However, while studies on invertebrate ageing are increasing, senescence biomarkers remain

poorly developed in these taxa. In order to develop useful senescence biomarkers in invertebrates and more particularly in the common woodlouse *Armadillidium vulgare*, we looked at the effect of age on three known biomarkers in vertebrates: immune cells (cell size, density and viability), β-galactosidase activity and Telomerase Reverse Transcriptase (TERT) (essential subunit of the telomerase protein) gene expression on this species. As expected, we observed that the size of immune cells was higher in older individuals while their density and viability decreased, the β-galactosidase activity increased with age while the Telomerase Reverse Transcriptase (TERT) gene expression decreased. These biomarkers classically used in vertebrates are thus correlated with age in our invertebrate model and make reliable biomarkers to highlight cell senescence in *A. vulgare*. A strong gender effect was also observed on senescence biomarkers that could indicate different resource allocation strategies. These biomarkers, allowing us to highlight differential cellular senescence between males and females, could be a new tool to test the impact of different factors influencing life history traits and related resources allocation and consequently help us to understand the diversity of senescence patterns observed between individuals, sexes, populations and species.

Keywords

Armadillidium vulgare, cellular senescence, immunosenescence, invertebrates, Telomerase Reverse Transcriptase (TERT), β-galactosidase activity.

1. Introduction

The cellular senescence corresponds to the progressive deterioration of the cell leading the stop of the cellular cycle (Campisi and di Fagagna, 2007). The main hypothesis proposed to explain this deterioration relies on the oxidative stress hypothesis, also known as free radical theory of ageing (Finkel and Holbrook, 2000; Wickens, 2001). According to this theory,

oxygenated reagents lead to an accumulation of damages, modifying the integrity and functions

of lipids, proteins as well as DNA the cell (Finkel and Holbrook, 2000; Terman and Brunk,

2006). Many studies have examined senescence and highlighted parameters related to this

phenomenon (De Jesus and Blasco, 2012).

One of the prominent characteristic of senescent cells is an increase in the cell size

which becomes larger than a non-senescent cell (Hayflick, 1965; Rodier and Campisi, 2011).

The evolution of the integrity and efficiency of immune cells is particularly relevant to study

cellular senescence as a diminution of the number of effective immune cells with age has been

reported in vertebrates (Cheynel et al., 2017) and invertebrates including mosquitoes *Aedes*

aegypti (Hillyer et al., 2004) and crickets *Gryllus assimilis* (Park et al., 2011). Another marker

used to study cellular senescence is the enzymatic activity of the β-galactosidase. This enzyme

is an hydrolase which transforms polysaccharides in monosaccharides, the lysosomal activity

of this enzyme is increased when the cell enters in senescence (Dimri et al., 1995; Itahana et

al., 2007). This phenomenon was observed in senescent cells of many organisms ranging from

humans (Gary and Kindell, 2005) to honeybees (Hsieh and Hsu, 2011). Another protein

involved in the cellular senescence process is the telomerase, a ribonucleoprotein complex

composed by two essential components, the telomerase reverse transcriptase (TERT) and the

telomerase RNA (TR) and others accessorial proteins (Podlevsky et al., 2007). Telomerase

lengthens the ends of telomeres, the latter are structures located at the end of chromosomes

which protect chromosome integrity and shorten after each cell division. Cell senescence arises

when the telomere size becomes critically short (Chiu and Harley, 1997; Shay and Wright,

2005).The telomerase activity depends on organism, age and also tissues. In humans,

telomerase is active during the development before the birth and after only in stem and germ

cells (Liu et al., 2007; Morgan, 2013) while in the *Daphnia pulicaria,* the telomerase activity

in all tissues of the body decreases with age (Schumpert et al., 2015). As the TERT is essential

in the telomerase protein complex it has been shown to be related to cell survival in humans (Cao et al., 2002) and was detected in numerous species including vertebrates, fungi, ciliates and insects (Podlevsky et al., 2007; Robertson and Gordon, 2006)

β-galactosidase activity, immune cell parameters (i.e. size, density and viability) and telomerase activity do not directly reflect the age of the organisms, but the senescence process itself; thus, they are classically considered as senescence biomarkers (Baker and Sprott, 1988). These biomarkers are often used to understand the processes involved in senescence and age-related diseases in humans (Hoos et al., 1998; Xia et al., 2017). Although their usefulness is not in doubt, they are never been used to understand variability of ageing patterns observed across and within species. Invertebrates are increasingly coveted in experimental approaches in senescence studies. Indeed, they share similarities with vertebrates in terms of senescence processes but also because it is easier to manipulate them experimentally and they can be monitored throughout their entire lifetime (Ram and Costa, 2018). These advantages make them models of choice for the development of senescence biomarkers.

In this context, we developed in a long-lived invertebrate (3-4 years old), the common woodlouse *Armadillidium vulgare,* the senescence biomarkers described above (β-galactosidase activity, immune cell parameters and the TERT gene expression) to be used in future studies on factors involved in ageing variability. According to the literature and over the time, we expected an increase in β-galactosidase activity, a decrease of both TERT gene expression and immune cell viability and density in *A. vulgare*. As at the adulthood, males have a greater survival than females (Paris and Pitelka, 1962) cellular senescence patterns are also expected to be sex-specific in *A. vulgare*.

2. Materials & Methods

2.1. Biological model

A. vulgare individuals used in the following experiments were derived from a wild

population collected in Denmark in 1982. These animals, able to live up to three years (Paris

and Pitelka, 1962) have been maintained on moistened soil under the natural photoperiod of

Poitiers (France 46.58°N, 0.34°E, 20°C) at 20°C fed (with dried linden leaves and carrots) *ad*

libitum. Crosses were monitored to control and promote genetic diversity. For each clutch

obtained, individuals were sexed, and brothers and sisters were separated to ensure virginity. In

common woodlouse, individuals molt throughout their lives, with approximately on moult per

month At 20°C (Steel, 1980), during this process all the cells of the concerned tissues are

renewed. However, the brain, the nerve cord and gonads are not renewed during molting and

are therefore relevant candidates for tissue-specific study of senescence in this species. Males

and females were tested separately to assess the impact of sex on our candidate biomarkers.

2.2 Measure of β-galactosidase activity

Animals

To test the impact of age on the on β-galactosidase activity, 180 individuals were used:

90 young (i.e. 6-months-old, 45 males and 45 females) and 90 old (2-years-old, 45 males and

45 females) individuals.

Protocol

Individuals were dissected separately in Ringer (Sodium Chloride 394 mM, Potassium

Chloride 2 mM, Calcium Chloride 2 mM, Sodium Bicarbonate 2 mM) and nerve cord was

removed. To obtain a sufficient amount of protein, we made pools of five nerve cords (from

five different individuals of the same age). The five nerve cords were filed in 500 µL of Lyse

Buffer 1X (CHAPS 5 mM, Citric acid 40 mM, Sodium Phosphate 40 mM, Benzamidine 0.5

mM, PMSF 0.25 mM, pH = 6) (Gary and Kindell, 2005), and then were centrifuged at 15000g

at 4°C for 30 minutes. The supernatant was taken and kept at -80°C until its utilization. The

protein concentration was determined by the BCA assay (Thermofisher) and were homogenized

at 0.1 mg/mL. The β-galactosidase activity was measured as previously described by Gary and

Kindell (2005). Briefly, 100 µL of extracted protein at the concentration of 0.1 mg/mL were

added to 100 µL of reactive 4-methylumbelliferyl-D-galactopyranoside (MUG) solution in a

96 well-microplate. The MUG reactive, in contact to β-galactosidase, leads by hydrolysis to the

synthesis of 4-methylumbelliferone (4-MU), which is detectable using fluorescent

measurements. Measures were performed by the multimode microplate reader Mithras LB940

HTS III, Berthold; excitation filter: 120 nm, emission filter 460 nm, for 120 minutes. Two

technical replicates were measured for each pool.

2.3 Measure of immune cell parameters

Animals

To test the impact of age on the immune cell parameters (i.e. density, viability, and size)

that were candidates for providing biomarkers of senescence in *A. vulgare,* 60 mature

individuals were used: 30 young (i.e. 1-year-old, 15 males and 15 females) and 30 old (3-years-

old, 15 males and 15 females) individuals.

Protocol

To study the impact of age on the immune parameters, a hole was bored in the middle

of the 6[th] segment and 3 µL of haemolymph were collected (per individual) with an eyedropper

and deposited promptly in 15 µL of anticoagulant solution of (MAS-EDTA (EDTA 9 mM,

Trisodium citrate 27 mM, NaCl 336 mM, Glucose 115 mM, pH 7, (Rodriguez et al., 1995))).

Then, 6 µL of Trypan blue at 0.4% (Invitrogen) were added to color the dead cells. Thereafter,

10 µL of this solution were deposed in (Invitrogen Coutness®) counting slide (Thermofisher).

The immune cell density, the immune cell viability and the immune cell size were evaluated

using an automated Cell Counter (Invitrogen Countess®).

2.4 Measure of TERT gene expression

The identification of the Telomerase Reverse Transcriptase (TERT) gene was firstly performed from the *A. vulgare* genome (Chebbi et al., 2019). In order to verify whether this gene was present and preserved in crustaceans, phylogenetic analyses were carried out upstream (see Supplementary materials 1, 2, 3 and 4). This gene has been found in crustacean transcriptomes and the topology of the TERT gene tree follows the phylogenetic relationships between the involved species, (Supplementary material 3), suggesting a significant and conserved role of the TERT gene.

2.4.1 Gene expression

Animals

We tested the effect of age on the expression of TERT gene within 4 different age groups: (1) 4-months-old, (2) 1-year-old, (3) 2-years-old and (4) 3-years-old. Females and males were tested separately by pools of 5 individuals in 1-, 2-, 3-years-old groups and by pools of 7 individuals in 4-months-old group. All conditions require 4 replicates for each sex. 176 individuals were used for this experiment. For each group we tested the expression level of the TERT gene in two different tissues: the nerve cord (somatic line) and gonads (germinal line).

Protocol

Animals were washed by immersion for 30s in a 30% sodium hypochlorite solution followed by two 30s immersion in distilled water. Tissues were dissected in Ringer solution (Sodium Chloride 394 mM, Potassium Chloride 2 mM, Calcium Chloride 2 mM, Sodium Bicarbonate 2 mM) and deposited by specific tissues pools of 5 on TRIzol reagent (Invitrogen) to extract RNA according to the manufacturer's protocol after a cell disintegration using a Vibra Cell 75,185 sonicator (amplitude of 30%). Total RNA was quantified by NanoDrop technology

and was stored at -80°C until used. Reverse transcriptions (RT) were made from 500ng of RNA

previously extracted and using the kit SuperScript[TM] IV Reverse Transcriptase (Thermo Fisher

Scientific) according to the supplier's instructions. Primers were designed using the identified

gene: primer TERT_F: 5'-AGGGAAAACGATGCACAACC-3' and primer TERT_R: 5'-

GTTCGCCAAATGTTCGCAAC- 3' (see Supplementary material 1). Quantitative RT-PCR

was performed using 0.6 μl of each primer (10 μM), 2.4 μl of nuclease-free water and 1.5 μl of

cDNA template and the LightCycler LC480 system (Roche) with the following program: 10

min at 95 °C , 45 cycles of 10 s at 95 ° C, 10 s at 60 °C, and 20 s at 72 °C. Expression levels of

target genes were normalized based on the expression level of two reference genes previously

established: the Ribosomal Protein L8 (RbL8) and the Elongation Factor 2 (EF2) (Chevalier et

al., 2011).

Statistics

All statistical analyses were performed using the R software (R. Core Team, 2016). The

β-galactosidase activity was analyzed with linear mixed effect models using the package lme4

(Bates et al., 2014). As two technical replicates were measured for each pool, the model

including the pools fitted as a random effect, age and sex and their two-way interactions as

fixed factors.

Concerning the immune parameters, linear models with Gaussian distribution were

fitted to analyze variations in the cell size and viability. For the cell density, a linear model of

the cell number (log-transformed, (Ives and Freckleton Robert, 2015) was fitted.

The level of TERT expression according to age in the two different tissues were

compared by a Kruskal–Wallis rank sum test in combination with Nemenyi's post hoc multiple

comparison test with the Tuckey correction using R package PMCMR.

3. Results

β-galactosidase activity

The β-galactosidase activity was higher in old individuals (i.e. 2-years-old) than in young ones (i.e. 6-months-old) (X^2_1=6.15, p=0.013, Figure 1). We also detected a significantly higher β-galactosidase activity in females than in males (X^2_1=7.26, p=0.007, Figure 1).

Figure 1: β-galactosidase activity according to age and sex in *A. vulgare* (F=females, M=males)

The thick line depicts the median, the box the interquartile range, and the whisker are bounded to the most extreme data point within 1.5 the interquartile range. The outliers outside this range are displayed as open circles. N = 24 pools of 5 individuals. * denotes *p<0.05*

Immune cells parameters

Cell size was larger in 3-years-old than in 1-year-old individuals ($F_{1,58}$=8.54, p=0.005, Figure 2A). Conversely, the cell density was higher in 1-year-old than in 3-years-old individuals ($F_{1,58}$=4.33, p=0.01, Figure 2B). Concerning the immune cell viability, a statistically significant

226   interaction occurred between age and sex, with a relatively lower immune cell viability in 3-

227   years-old females ($F_{3,56}=6.85$, p=0.01, Figure 2C). No sex effect was detected on cell size

228   ($F_{2,57}=0.76$, p=0.38, Figure 2A) or cell density ($F_{2,57}=0.32$, p =0.57, Figure 2B).

Figure 2: Immune cell size (A), density (B) and viability (C) according to age and sex in *A. vulgare* (F=females, M=males)

The thick line depicts the median, the box the interquartile range, and the whisker are bounded to the most extreme data point within 1.5 the interquartile range. The outliers outside this range are displayed as open circles. N= 60 individuals: 15 1-year-old females, 15 1-year-old males, 15 3-years-old females and 15 3-years-old males. * denotes p<0.05 and ** denotes p<0.01

TERT gene expression

The TERT gene expression decreased with age in nerve cords (X^2_3=23.30, p<0.001, Figure 3A). More precisely, the TERT gene expression was higher in 4-months-old individuals comparing to 2-years-old individuals and 3-years-old individuals (Respectively: p=0.001, p<0.001) and in 1-year-old individuals comparing to 3-years-old individuals (p=0.038), without sex effect (X^2_1=0.14, p=0.70, Figure 3A). In gonads, the TERT gene expression was strongly higher in females (X^2_1=17.81, p<0.001, Figure 3B) and tended to decrease with age (X^2_3=7.5, p=0.057, Figure 3B) as the TERT gene expression tended to be higher in 4-months-old females comparing to 3-years-old females (p=0.054). In males, a general tendency was also observed (X^2_1=7.34, p=0.061, Figure 3B), the TERT gene expression tending to be higher in 2-years-old individuals but not significantly comparing to 1-year-old individuals and 3-years-old individuals (Respectively: p=0.14, p=0.12).

Figure 3: Relative expression level of TERT in (A) nerve cords and (B) in gonads in *A. vulgare* (F=females, M=males).

Expression of each gene was normalized based on the expression of Ribosomal Protein L8 (RbL8) and Elongation Factor 2 (EF2) as reference genes. The thick line depicts the median, the box the interquartile range, and the whisker are bounded to the most extreme data point within 1.5 the interquartile range. N= 176 individuals: 28 4-months-old females, 28 4-months-old males, 20 1-year-old females, 20 1-year-old males, 20 2-years -old females, 20 2-years-old males, 20 3-years-old females, 20 3-years-old males.

. denotes p<0.10, ** denotes p<0.01

4. Discussion

In this study, we tested several effective senescence biomarkers on vertebrates to see if they could also be used in our invertebrate model: the woodlouse. As expected, immune cells showed an increase in their size and a decrease in their density and viability with age. In nerve cords, the activity of the β-galactosidase enzyme increased while the TERT gene expression decreased with age. These results support the presence of increasing cellular senescence in *A.*

vulgare with age. In the gonads, the TERT gene expression was too low in males and was not

sufficiently variable between sexes to provide good senescence biomarker in this tissue.

Our study is in line with previous studies that have already revealed the possibility of

using vertebrate biomarkers in invertebrates (Hsieh and Hsu, 2011; Park et al., 2011; Schumpert

et al., 2015). By testing a set of different biomarkers, often studied independently, our study

confirms the idea that routine biomarkers used in vertebrates can be adapted in invertebrates

and supports that the senescence process is quasi-ubiquitous in living world and can be

expressed in a similar way in very different organisms.

Previously study showed that the probabilities to survive decrease with age in *A. vulgare*

(Paris and Pitelka, 1962). The cellular damages accumulated during the animal's life could be

the cause of cell senescence and therefore the driving force behind actuarial senescence. (Barja,

2000; Barja and Herrero, 2000; Finkel and Holbrook, 2000; Harman, 1956). In *A. vulgare*, the

2- and 3-years-old individuals could have therefore accumulated more cellular damages during

their life leading the cellular senescence observed.

Our present study also showed a strong difference between sexes concerning the cellular

senescence patterns on biomarkers: females present higher β-galactosidase activity and lower

immune cell viability than males. Between-sex differences in lifespan have been reported in *A.*

vulgare with a longer lifespan in males than in females (Geiser, 1934; Paris and Pitelka, 1962).

Exact differences in actuarial senescence patterns remain to be quantified in *A. vulgare* but such

differences are quite common both in vertebrates and invertebrates (Tidière et al. 2015; Marais

et al. 2018 for reviews). One of the main theory proposed to explain sex differences in longevity

senescence patterns relies on different resource allocation strategies between sexes

(Bonduriansky et al., 2008; Vinogradov, 1998), which was already evoked to explain the shorter

lifespan observed in females *A. vulgare* as they invest more energy in reproduction than males

(Paris and Pitelka, 1962).

The resources allocation strategies proposed to explain sex differences could also be

driven by environment (Shertzer and Ellner, 2002). As our senescence biomarkers were able to

highlight differences between sexes, and as support in Depeux et al., 2019, they could constitute

useful tools to underline other factors (e.g. environment stressors) involved in the diversity of

senescence patterns observed between individuals, sexes, populations and species.

Acknowledgements

298   We would like to thank Sylvine Durand, Richard Cordaux, Isabelle Giraud and

299   Bouziane Moumen for our constructive discussions as well as Marius Bredon, Carine Delaunay,

300   Maryline Raimond and Alexandra Lafitte for technical assistance.

Funding

This work was supported by the 2015–2020 State-Region Planning Contract and

European Regional Development Fund and intramural funds from the Centre National de la

Recherche Scientifique and the University of Poitiers. J.F.L. and J.M.G. are supported by a

grant from the Agence Nationale de la Recherche (ANR-15-CE32-0002-01 to J.F.L.). This

work has also received funding from the "Appel à projets de recherche collaborative inter-

équipes 2016-2017" by the laboratory EBI.

Bibliography

312   Barja, G., 2000. The flux of free radical attack through mitochondrial DNA is related to aging
313   rate. Aging (Milano) 12, 342–355.
314   Barja, G., Herrero, A., 2000. Oxidative damage to mitochondrial DNA is inversely related to
315   maximum life span in the heart and brain of mammals. FASEB J. 14, 312–318.

Bonduriansky, R., Maklakov, A., Zajitschek, F., Brooks, R., 2008. Sexual selection, sexual
conflict and the evolution of ageing and life span. Functional Ecology 22, 443–453.
https://doi.org/10.1111/j.1365-2435.2008.01417.x
Campisi, J., di Fagagna, F. d'Adda, 2007. Cellular senescence: when bad things happen to good
cells. Nature reviews. Molecular cell biology 8, 729.
Cao, Y., Li, H., Deb, S., Liu, J.-P., 2002. TERT regulates cell survival independent of
telomerase enzymatic activity. Oncogene 21, 3130–3138.
https://doi.org/10.1038/sj.onc.1205419
Chebbi, M.A., Becking, T., Moumen, B., Giraud, I., Gilbert, C., Peccoud, J., Cordaux, R., 2019.
The Genome of *Armadillidium vulgare* (Crustacea, Isopoda) Provides Insights into Sex
Chromosome Evolution in the Context of Cytoplasmic Sex Determination. Molecular
Biology and Evolution. https://doi.org/10.1093/molbev/msz010
Chevalier, F., Herbinière-Gaboreau, J., Bertaux, J., Raimond, M., Morel, F., Bouchon, D.,
Grève, P., Braquart-Varnier, C., 2011. The Immune Cellular Effectors of Terrestrial
Isopod Armadillidium vulgare: Meeting with Their Invaders, Wolbachia. PLOS ONE
6, e18531. https://doi.org/10.1371/journal.pone.0018531
Cheynel, L., Lemaître, J.-F., Gaillard, J.-M., Rey, B., Bourgoin, G., Ferté, H., Jégo, M., Débias,
F., Pellerin, M., Jacob, L., Gilot-Fromont, E., 2017. Immunosenescence patterns differ
between populations but not between sexes in a long-lived mammal. Scientific Reports
7. https://doi.org/10.1038/s41598-017-13686-5
Chiu, C.-P., Harley, C.B., 1997. Replicative Senescence and Cell Immortality: The Role of
Telomeres and Telomerase. Experimental Biology and Medicine 214, 99–106.
https://doi.org/10.3181/00379727-214-44075
Clutton-Brock, T.., Isvaran, K., 2007. Sex differences in ageing in natural populations of
vertebrates. Proceedings of the Royal Society B: Biological Sciences 274, 3097–3104.
https://doi.org/10.1098/rspb.2007.1138
De Jesus, B.B., Blasco, M.A., 2012. Assessing cell and organ senescence biomarkers.
Circulation research 111, 97–109.
Depeux, C., Samba-Louaka, A., Braquart-Varnier, C., Moreau, J., Lemaître, J.-F., Laverre, T.,
Pauhlac, H., Dechaume-Moncharmont, F.-X., Gaillard, J.-M., Beltran-Bech, S., 2019.
Impact of temperature and photoperiod on survival and biomarkers of senescence in
common woodlouse. bioRxiv. https://doi.org/10.1101/433011
Dimri, G.P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., Medrano, E.E., Linskens,
M., Rubelj, I., Pereira-Smith, O., 1995. A biomarker that identifies senescent human
cells in culture and in aging skin in vivo. PNAS 92, 9363–9367.
Finkel, T., Holbrook, N.J., 2000. Oxidants, oxidative stress and the biology of ageing. Nature
408, 239–247.
Gary, R.K., Kindell, S.M., 2005. Quantitative assay of senescence-associated β-galactosidase
activity in mammalian cell extracts. Analytical biochemistry 343, 329–334.
Geiser, S.W., 1934. Further observations on the sex-ratios of terrestrial isopods. Field Lab 3,
7–10.
Harman, D., 1956. Aging: a theory based on free radical and radiation chemistry. Journal of
gerontology 11, 298–300.
Hayflick, L., 1965. The limited in vitro lifetime of human diploid cell strains. Experimental
Cell Research 37, 614–636. https://doi.org/10.1016/0014-4827(65)90211-9
Hillyer, J.F., Schmidt, S.L., Fuchs, J.F., Boyle, J.P., Christensen, B.M., 2004. Age-associated
mortality in immune challenged mosquitoes (Aedes aegypti) correlates with a decrease
in haemocyte numbers: Age-associated mortality in Aedes aegypti. Cellular
Microbiology 7, 39–51. https://doi.org/10.1111/j.1462-5822.2004.00430.x

Hoos, A., Hepp, H.H., Kaul, S., Ahlert, T., Bastert, G., Wallwiener, D., 1998. Telomerase activity correlates with tumor aggressiveness and reflects therapy effect in breast cancer. International Journal of Cancer 79, 8–12. https://doi.org/10.1002/(SICI)1097-0215(19980220)79:1<8::AID-IJC2>3.0.CO;2-5

Hsieh, Y.-S., Hsu, C.-Y., 2011. Honeybee trophocytes and fat cells as target cells for cellular senescence studies. Experimental Gerontology 46, 233–240. https://doi.org/10.1016/j.exger.2010.10.007

Itahana, K., Campisi, J., Dimri, G.P., 2007. Methods to detect biomarkers of cellular senescence: the senescence-associated β-galactosidase assay. Biological Aging: Methods and Protocols 21–31.

Ives, A.R., Freckleton Robert, 2015. For testing the significance of regression coefficients, go ahead and log-transform count data. Methods in Ecology and Evolution 6, 828–835. https://doi.org/10.1111/2041-210X.12386

Liu, L., Bailey, S.M., Okuka, M., Muñoz, P., Li, C., Zhou, L., Wu, C., Czerwiec, E., Sandler, L., Seyfang, A., Blasco, M.A., Keefe, D.L., 2007. Telomere lengthening early in development. Nature Cell Biology 9, 1436–1441. https://doi.org/10.1038/ncb1664

Morgan, G., 2013. Telomerase regulation and the intimate relationship with aging. Research and Reports in Biochemistry 71. https://doi.org/10.2147/RRBC.S28603

Paris, O.H., Pitelka, F.A., 1962. Population characteristics of the terrestrial isopod Armadillidium vulgare in California grassland. Ecology 43, 229–248.

Park, Y., Kim, Y., Stanley, D., 2011. Cellular immunosenescence in adult male crickets, Gryllus assimilis. Archives of Insect Biochemistry and Physiology 76, 185–194. https://doi.org/10.1002/arch.20394

Podlevsky, J.D., Bley, C.J., Omana, R.V., Qi, X., Chen, J.J.-L., 2007. The Telomerase Database. Nucleic Acids Research 36, D339–D343. https://doi.org/10.1093/nar/gkm700

Ram, J.L., Costa, A.J., 2018. Invertebrates as Model Organisms for Research on Aging Biology ∗, in: Conn's Handbook of Models for Human Aging. Elsevier, pp. 445–452. https://doi.org/10.1016/B978-0-12-811353-0.00032-4

Robertson, H.M., Gordon, K.H.J., 2006. Canonical TTAGG-repeat telomeres and telomerase in the honey bee, Apis mellifera. Genome Research 16, 1345–1351. https://doi.org/10.1101/gr.5085606

Rodier, F., Campisi, J., 2011. Four faces of cellular senescence. The Journal of Cell Biology 192, 547–556. https://doi.org/10.1083/jcb.201009094

Rodriguez, J., Boulo, V., Mialhe, E., Bachere, E., 1995. Characterisation of shrimp haemocytes and plasma components by monoclonal antibodies. Journal of Cell Science 108, 1043–1050.

Schumpert, C., Nelson, J., Kim, E., Dudycha, J.L., Patel, R.C., 2015. Telomerase Activity and Telomere Length in Daphnia. PLOS ONE 10, e0127196. https://doi.org/10.1371/journal.pone.0127196

Shay, J.W., Wright, W.E., 2005. Senescence and immortalization: role of telomeres and telomerase. Carcinogenesis 26, 867–874. https://doi.org/10.1093/carcin/bgh296

Shertzer, K.W., Ellner, S.P., 2002. State-dependent energy allocation in variable environments: life history evolution of a rotifer. Ecology 83, 2181–2193. https://doi.org/10.1890/0012-9658(2002)083[2181:SDEAIV]2.0.CO;2

Steel, C.G.H., 1980. Mechanisms of coordination between moulting and reproduction in terrestrial isopod crustacea. The Biological Bulletin 159, 206–218. https://doi.org/10.2307/1541019

Terman, A., Brunk, U.T., 2006. Oxidative Stress, Accumulation of Biological "Garbage", and
Aging. Antioxidants & Redox Signaling 8, 197–204.
https://doi.org/10.1089/ars.2006.8.197
Vinogradov, A.E., 1998. Male Reproductive Strategy and Decreased Longevity. Acta
Biotheoretica 46, 157–160. https://doi.org/10.1023/A:1001181921303
Wickens, A.P., 2001. Ageing and the free radical theory. Respiration Physiology 128, 379–391.
https://doi.org/10.1016/S0034-5687(01)00313-9
Xia, X., Chen, W., McDermott, J., Han, J.-D.J., 2017. Molecular and phenotypic biomarkers of
aging. F1000Research 6, 860. https://doi.org/10.12688/f1000research.10692.1