Visual area V4 encodes history dependent attentional effort and single-trial perceptual detection

Authors: Supriya Ghosh[*] and John H. R. Maunsell

Affiliations: Department of Neurobiology, The University of Chicago, Chicago, Illinois 60637, USA.

* Correspondence: sghosh5@uchicago.edu

ABSTRACT

Intense attentional effort improves performance. Neurons in visual area V4 exhibit changes in spike rates that are correlated with effort, selective attention and absolute reward expectation. How neuronal spikes in V4 encode attentional states and perceptual performance with reference to other task relevant motivational and decision signals is poorly understood. We recorded neuronal spikes from monkey visual area V4, and isolated attentional effort from reward expectancy and motor decisions in a visual orientation detection task using a statistical approach based on generalized linear models. We found that V4 spike responses exhibit hysteresis with reward changes. Single trial spiking activity encodes a combination of temporally overlapping sensory and cognitive signals. Neuronal spikes integrate contextual history of attentional effort on a longer time scale, and task related perceptual detection in a shorter timescale within a single trial. Notably, immediate reward expectation and behavioral choice probability are poor predictors of V4 spikes throughout the trial. Further, detection of task relevant perceptual signal is decoded from single trial population spikes with high accuracy. These results provide a detailed representation of perceptual and cognitive signals in V4 that are crucial for performance.

KEYWORDS

Attentional effort; reward; visual cortex; single-unit recording

INTRODUCTION

The modulation of neuronal responses by spatial or featural attention has been examined extensively in area V4 in monkey visual cortex. When a monkey's attention is directed towards the receptive field (RF) location of a V4 neuron, improvement in perceptual performance in that region is accompanied by several changes in neuronal activity. Typically, spike rates are greater (Luo and Maunsell, 2015; Moran and Desimone, 1985); individual response variance is reduced; and pairwise spike correlations (noise correlation) decrease (Cohen and Maunsell, 2009; Reynolds and Heeger, 2009). Similarly, responses of V4 neurons can be augmented when an animal increases it attentional effort (the intensive aspect of attention) when faced with a more challenging task (Boudreau et al., 2006; Spitzer et al., 1988).

However, interpreting neuronal correlates of attention is challenging because it is so closely intertwined with many cognitive and experimental covariates, such as stimulus features, action planning and reward parameters. Expectation of a potential reward is one of the primary external motivators for an animal engaged in a laboratory task. Information about rewards has been reported to exist in many brain structures, including visual areas that are affected by changes in attentional states. Spike rates of neurons in visual cortical area V4 and the lateral intraparietal cortex (LIP) can follow absolute reward size associated with a visual target inside their RFs in a manner independent of attentional performance (Baruni et al., 2015; Rorie et al., 2010). Neuronal encoding of expected reward or choice is well studied in the striatum (Schultz et al., 1992; Tremblay et al., 1998) and prefrontal cortex (Donahue and Lee, 2015; Padoa-Schioppa and Assad, 2006; Roesch and Olson, 2004), and is characterized by neuronal activity correlated with the absolute reward contingency associated with a stimulus while being independent of the relative reward value. Because attention and reward are both defined by their action on goal-directed behavior, their high

covariance in relation to performance makes it difficult to isolate the true nature of the

representation of attention states within the visual cortex.

The contribution of visual cortical neurons in encoding attention is typically examined by

instructing the animal to direct attention in one way within a block of trials using different, fixed

reward conditions, which vary between blocks. This allows animals to attain a stable cognitive

state of reward expectation as well as attention. In studies aimed at understanding neuronal

encoding of relative versus absolute reward sizes, different reward conditions are often interleaved

across trials and signaled by a cue at the start of each trial. Such task designs effectively rule out

contributions due to slowly varying or history-dependent attentional effort that changes over

several trials. In this framework, the cognitive state of attentional effort may not settle to a stable

level, limiting the animal's ability to take advantage of perceptual benefits of intense attentional

effort (or economic utilization of costly cognitive resources) in the case of large over small rewards

(Baruni et al., 2015; Rorie et al., 2010) (but see (Cicmil et al., 2015)). Thus, in order to define

attention quantitatively, it is crucial to understand how neurons represent attentional effort and its

transitions in relation to global physiological cognitive state. To do so, one must first define and

experimentally control the cognitive effort that manifests across neuronal, physiological and

perceptual performance levels. It is additionally crucial to identify how both steady-state and

transitions in attentional effort (e.g. high-to-low or low-to-high) correlate with neuronal,

physiological and behavioral responses. Finally, it is important to isolate the relative contributions

of different task-relevant cognitive and experimental covariates to visual cortical spikes within a

single trial as the trial progresses.

We manipulated the level of monkeys' attentional effort using a novel task in which their

overall attentional effort was switched between high and low levels with no change in attentional

selectivity for stimuli at one location over another. We found that attentional effort modulates V4

neurons independently of spatial or featural attention. Additionally, by varying reward size on each

trial around an average, we could produce a single trial estimates of reward expectation as well as

both the transient and stable (slow-varying) attentional effort. Using a generalized linear model

(GLM) on single trial spike-counts of individual V4 neurons, we quantitatively described

contributions of temporally overlapping neuronal response components associated with attentional

effort, reward expectation and perceptual detectability. This framework revealed both steady state

and time varying signals encoded in V4, enabling the assessment of a broader spectrum of

cognitive functions to which visual cortical areas can contribute dynamically.

RESULTS

Reward size modulates attentional effort

We trained two rhesus monkeys to distribute their visual spatial attention between the left and right

hemifields while doing an orientation detection task (**Figure 1A**). The animal held its gaze on a

central fixation spot throughout each trial. Two Gabor sample stimuli appeared briefly (200 ms)

after a randomly varying period of fixation and, after a subsequent delay (200-300 ms), a single

Gabor test stimulus appeared at one of the two sample locations (selected pseudo-randomly). If

the orientation of the test stimulus differed from the sample that had appeared in that location (a

target) the monkey had to rapidly make a saccade to the stimulus to earn a juice reward. On a

random half of the trials, the orientation of test stimulus was unchanged (a non-target) and the

monkey was required to maintain fixation. In that case, a second test stimulus that always had a

different orientation was presented after a short delay and monkeys needed to saccade to this

stimulus to earn a reward. Reward values for correct responses were always the same on both sides

and fixed, either small or large, within alternating blocks of 120 trials. Stimulus parameters

remained unchanged throughout all trial blocks within a session except the reward size. The

transitions between blocks were unsignaled and sequential (small to large or large to small, **Figure**

1B). The monkey therefore had to estimate reward expectancy based on previous trials to adjust

its behavioral strategy in allocating effort between blocks.

Varying rewards in different blocks of trials allowed us to behaviorally control the amount

of attention the animal directed to each stimulus location. Animals were encouraged to maintain a

Figure 1. Manipulation of attentional effort using differential reward outcomes. (**A**) Visual spatial attention task. Monkeys were required to fixate, attend to sample stimuli (Gabors) presented in both hemifields (inside and outside of recorded neurons' receptive field (RF)) and report an orientation change that occurred in one of the two test intervals by making a saccade to the stimulus location. (**B**) Unsignaled change in reward size between large and small values over blocks of 120 trials. (**C**) Attention operating characteristic (AOC) curve, indicating behavioral sensitivity (*d'*, circles) and criterion (*c*, triangles) on individual sessions and their average (solid markers) for test stimuli inside and outside RF during two reward conditions, large and small (24. Total sessions: 9 monkey P; 15 monkey S). Dotted colored lines indicate average *d'* in each hemifield. Lines connect two reward conditions within a session. Error bars, 95% confidence intervals. (**D**) Eye positions (*top*) and pupil areas (*bottom*) aligned to trial events in an example session. Each of 9 blocks, 120 trials/ block. Error bars, 95% confidence intervals. (**E**) Block averaged pupil area during pre-stimulus fixation and sample stimulus periods. Pupil area was normalized to mean of pre-stimulus fixation during small reward trial blocks. Error bars, 95% confidence intervals.

behavioral target/non-target bias (criterion, *c*) close to zero by providing rewards of different sizes

for hits and correct rejections (**Figure S1**). Because the reward sizes for targets were always the

same in both hemifields, animals typically allocated equal attention to both sides within either the

high and low reward condition. Figure 1c plots behavioral sensitivity (*d'*) and response criterion

(*c*) for behavioral performance on the first test stimuli on the left and right side for the large and

small reward conditions. Solid lines join average sensitivities from alternating high- and low-

reward blocks from a single day. Dashed arcs around the origin mark isopleths of constant total

effort across the two stimulus locations. Movements perpendicular to these lines correspond to

changes in attentional effort; movements along these lines correspond to changes in attentional

selectivity. Sensitivities on both locations are well balanced (no net attentional selectivity) during

most individual sessions. Larger rewards strongly motivated the animals to increase their

attentional effort compared to small reward trial blocks. A four-fold increase in reward size

(median small reward 131 µl, quartiles 108 and 150 µl; median large reward 522 µl, quartiles 469

and 570 µl) increased overall behavioral *d'* (Methods) by 71% (mean small 2.03, SEM 0.07; mean

large 3.48, SEM 0.08, $p < 10^{-17}$, paired t-test; **Figure 1C**, **Table S1**). Behavioral response criterion

remained near zero regardless of reward size (**Figure 1C**, **Table S1**).

Non-luminance mediated task-evoked modulation of pupil size is commonly considered a

proxy for mental effort across species (Beatty, 1982; Piquado et al., 2010) (for review (Laeng et

al., 2012)). Throughout the task, animals maintained a stable fixation (example session in **Figure**

1D, *top*), during which pupil area remain elevated for large reward trials relative to small reward

trials (example session in **Figure 1D**, *bottom*). Large reward increased pupil area during fixation

by 9% (all values normalized to fixation on small reward trials, mean large 1.09, SEM 0.02) and

during sample stimuli by 24% (mean small 1.05, SEM 0.02, mean large 1.31, SEM 0.02) (repeated

measures ANOVA, $F_{(1, 415)} = 45.5$, $p < 10^{-10}$; **Figure 1E**). While the within block trial-averaged

behavioral *d'* was strongly positively correlated with both reward size and pupil area during sample

stimuli, pupil area failed to exhibit any correlation with reward size (partial correlation, reward -

d', $\rho = 0.68$, p $< 10^{-55}$; pupil area - *d'*, $\rho = 0.23$, p $= 10^{-5}$; reward - pupil area, $\rho = 0.08$; p $= 0.09$; *N*

= 417 blocks of trials).

Responses of V4 neurons increase with increasing attentional effort

Some previous studies have shown that attention-related modulation of the responses of V4

neurons increases when tasks demand more cognitive effort (Boudreau et al., 2006; Spitzer et al.,

1988). However, in those studies increases in task difficulty always covaried with increases in

behavioral selectivity for one visual field location over another. A more recent study similarly

showed that the neurons in V4 respond more strongly when monkeys shift their attention to

increase their behavioral *d'* at the recorded neurons' receptive field location relative to another

location in the opposite hemifield (Luo and Maunsell, 2015), a manipulation that also covaried

selectivity and sensitivity. Our task design allowed us to change attentional effort with no

appreciable change in the selectivity for one spatial location over the other (**Figure 1C).**

We recorded 970 single units and small multi-unit clusters during 24 recording sessions

from the two monkeys (monkey P, 9 sessions, 407 units; monkey S, 15 sessions, 563 units) using

96 channel multielectrode arrays chronically implanted in V4 in the superficial prelunate gyrus of

one hemisphere. Trial averaged population spike rates during the sample stimuli period for

correctly completed trials (hits and correct rejections) increased in large reward trial-blocks when

the monkeys' attentional effort was higher (**Figure 2A-C**). Average responses were low because

the stimulus was patently suboptimal for most of the simultaneously recorded neurons. The peak-

normalized population spike rates in one monkey (animal S) show greater sustained activity than

the other (animals P, **Figure 2B, C**). This difference arose from differences in average absolute

peak firing rate (monkey P mean 28.7 SEM 1.1 spikes/s, monkey S mean 24.6 SEM 0.9 spikes/s,

A Example neuron

Figure 2. Higher attentional effort increased spiking of neurons in area V4. (**A**) Peri-stimulus time histograms (PSTH) of spike rates of correct trials in large and small reward blocks for an example neuron in V4. Single trial spike counts were binned at 2 ms, smoothed with $\sigma = 15$ ms half-Gaussian and then aligned at the onset of sample stimulus. Horizontal bar, sample stimulus presentation. (**B-C**) Population PSTHs for monkey P (B) and monkey S (C). For population average, spike rates of each neuron were normalized to its peak response within 60 - 260 ms from sample stimulus onset (monkey P, $n = 407$; monkey-S, $n = 563$). Error bars, 95% confidence intervals (bootstrap, $n = 10^4$). (**D**) Distribution of neuronal modulation indices (MI) of all units from both monkey P and S ($n = 970$). Red bars, neurons with MI values significantly different from 0 ($n = 603/970$, $p < 0.05$). White bars, non-significant MI. Solid triangle, population MI averaged across all units ($n = 970$, $p < 10^{-105}$, t-test).

B Monkey P
$n = 407$

C Monkey S
$n = 563$

D

p < 0.01, ranksum test) and average absolute sustained rates (monkey P mean 16.2 SEM 0.7

spikes/s, monkey S mean 18.5 SEM 0.6 spikes/s, p < 0.01, ranksum test). Principal component

(PC) analysis of spike peri-stimulus histograms of all neurons revealed that monkey P has higher

principal component scores associated with third PC, which captured transient peak response

(**Figure S2**). To quantify neuronal modulation by attentional effort, we computed a modulation

index (MI) as the difference of firing rates (60-260 ms from sample onset) between high effort

(large reward) and low effort (small reward) conditions divided by their sum. The mean MI was

significantly greater than zero, indicating higher firing rates during high effort condition (mean =

0.05, $p < 10^{-105}$, $n = 970$, t-test; **Figure 2D** or ~11.6% increase in firing rate). The attentional effort

effect was also significant for individual monkeys (monkey P, mean = 0.08, $p < 10^{-70}$; monkey S,

mean = 0.03, $p < 10^{-52}$; **Figure S3**). In addition, of 970 units recorded in V4 in the two monkeys,

significant effort modulations ($p < 0.05$) were observed in 603 (62%) units. Thus, an isolated

change in attentional effort affects responses in V4 in absence of behavioral selectivity or response

criterion changes.

Trial-by-trial physiological and behavioral dynamics in response to reward modulation

Many experimental covariates might closely follow the timing of changes in reward size. We

therefore examined the dynamics of neuronal, physiological and behavioral responses associated

with reward changes. Trial-by-trial values of behavioral d', sample-stimuli-evoked pupil area and

spike counts were aligned to the first correct trial after block transitions, and then averaged across

blocks (**Figure 3**). The first reward in a block unambiguously signaled a block transition, which

was otherwise unannounced. Block-averaged single trial reward values received by the animals

approximated step functions (single exponential fit: $\tau = 0.2$ and 0.5 trials respectively for

transitions from small-to-large ($n = 212$ blocks) and from large-to-small ($n = 205$ blocks), **Figure**

3A). Behavioral d' closely tracked reward changes for the transition from large-to-small ($\tau = 1$).

Figure 3. Directions of reward change have differential delayed effects on behavior and physiology. (A-D) Block averaged single received rewards (A), behavioral sensitivity (*d'*) (B), normalized mean pupil area during sample stimulus period (C) and normalized V4 spike counts across all neurons (*n* = 970; d) in large (*N* = 212) and small (*N* = 205) reward blocks. Circles, observed data. Lines, single exponential fits. *τ*, decay or rise constants. Trials are aligned with respect to the first correct trial following block transition. Dashed lines, 95% confidence intervals. **(E-G)** Hysteresis with transition of attentional effort. Behavioral *d'* (E), pupil area (F) and V4 spike counts (G) in response to reward changes from large to small and reverse. Horizontal error bars, mean ± SEM of reward size across blocks at the center value of y-axis variable. **(H)** V4 spike counts as a function of *d'* changes. Vertical lines, mean *d'* across blocks at the half-value of spike counts. **(I)** *Top*, Analysis widows for reward matched V4 spike counts in large and small reward blocks within first 10 trials at the beginning (*middle*, cyan) and last 60 trials at the end of blocks (*bottom*, purple). Bars, mean received rewards across trials binned between 0-1 (bin size = 0.2, overlap = 0.1; within session normalized) for two reward-block conditions. Error bars, 95% confidence intervals (bootstrap, *n* = 10^4). Circles, mean normalized spike counts (*n* = 970) on immediate next trial. **(J)** *Left*, Linear regression between mean pupil area (0 - 400

ms), spike counts (60 - 260 ms) and reward conditions (high and low efforts) across all trials were fit separately for each neuron. Population mean ± SEM of fitted coefficients (effort and spike counts) for neurons with significant fit (p < 0.05, $n = 625$). *Right*, Two separate linear regression fits between pupil area and spike counts for large and small reward trials. Mean ± SEM of fitted coefficients of spike counts in for the two reward conditions for neurons with significant fit (p < 0.05, $n = 334$).

However, the *d'* was much slower ($\tau = 12.2$) for transitions from small to large (**Figure 3B**). As

expected, trial constants of percent-correct followed a time course similar to *d'* (large-to-small, τ

$= 1$; small to large, $\tau = 9.0$; **Figure S4**). In contrast to task performance, pupil area and population

mean V4 spike responses (60-260 ms from sample onset) had the same slow decays for large to

small transitions ($\tau = 9$ for both) and similar, faster rises for small to large transitions ($\tau = 4$ and 2

respectively for pupil area and spikes, **Figure 3C-D**).

The dynamics of the effects of reward transitions on neuronal firing, pupil area and

behavior can be clearly seen in hysteresis plots (**Figure 3E-G**). Behavioral *d'* tracks transitions

from small-to-large reward size more slowly than transitions from large-to-small reward size

(**Figure 3E)**. In contrast, pupil area and neuronal responses track transitions from small-to-large

reward size more quickly than transitions from large-to-small. The normalized reward value at

which the values reach their mid-point differ significantly for high-to-low and low-to-high

transitions for all three measures (horizontal bars in Figure 3E-G, mean ± SEM, $p < 10^{-31}$, t-test).

Similar neuronal hysteresis is also seen in motion perception and contrast detection in humans

where stimulus history affects perceptual detection threshold (Kleinschmidt et al., 2002; Williams

et al., 1986). The different dynamics between *d'* and spike rate were almost complementary

(**Figure 3H**), with spike rate leading for low-to-high transitions and lagging for high-to-low

transitions of *d'*.

While the average spike rate in area V4 changed slowly following block transitions, the

averages might obscure trial-to-trial changes in neuronal activity that tracked the reward received

on the previous trial. Because rewards for hits and correct rejections (CR) typically differed

(**Figure S1**), we could measure of dependency of spike counts on the previous reward. We sorted

trials into different bins depending on the normalized reward size on the preceding trial and

measured spike counts of each neuron (Methods). We further separated the trials based on whether

they occurred immediately after block transitions (within first 1-10 trials from the first correct

response after the reward switch) or during steady-state within a block (final 60 trials, 61-120)

(*top*, **Figure 3I**). Reward value on the previous trial did not affect spike counts [$F_{(8, 12801)} = 1.46$, p

$= 0.164$ for repeated measured factor reward value (9 levels); $F_{(1, 12801)} = 3.72$, $p = 0.0537$ for

between group factor, block-reward condition (large and small)] (*middle*, **Figure 3I**). The lack of

reward dependency of spike responses remained unchanged even during the late, steady state

portion of blocks (late trials), when a large effect of block-reward size was seen [$F_{(8, 10752)} = 1.18$,

$p = 0.307$ for reward value; $F_{(1, 10752)} = 7.78$, $p = 0.005$ for block-reward condition] (*bottom*, **Figure**

3I). Thus, the strong V4 response modulations we saw were dominated by slow effects.

The similar dynamics of pupil area (**Figure 3C**) and spike rate (**Figure 3D**) suggest that a

larger pupil area associated with high effort might enhance effective retinal illumination, which

might elevate V4 neuronal spiking. To test this possibility, we quantified how well pupil area

correlates with V4 spike rates and effort states ('high' or 'low') for all neurons using linear

regression (Methods). Regression coefficients for effort were more than eight-fold higher than the

coefficients for spike rate (mean ± SEM, spike rate, $0.08 ± 0.005$; effort, $0.68 ± 0.01$; Wilcoxon

rank sum test, (WRS) $n = 625$, $p = 10^{-196}$; **Figure 3J**, *left*). Weak relationship between pupil area

and spike rate was similar for the two reward conditions, when mean pupil area was fit separately

for small and large reward trials (mean ± SEM, small reward, $0.094 ± 0.008$; large reward, 0.097

$± 0.01$; WRS test, $n = 334$, $p = 0.13$; **Figure 3J**, *right*). We additionally measured the temporal

relationship between single trial V4 spike trains and pupil area dynamics using cross-correlation

and spike triggered averaged (STA) pupil area for each neuron (Methods). None of the neurons

show any significant correlation between PSTHs of pupil area and spike rates (binned at 10 ms)

over the analysis window from -350 ms to 350 ms from sample on (**Figure S6**). Similarly, STA-

pupil area that measured the extent to which individual spikes were aligned with pupil area did not

show any significant relationship between single spikes of individual neurons and pupil area

(absent of STA-pupil area >0 at negative lags; **Figure S7**). Together, these results suggest that V4

spike activity and the autonomic sympathetic system that regulates pupil dilation are both strongly

modulated by common top-down cognitive states rather than a direct influence of changing retinal

illumination on V4 spike activity within the limit of observed pupil area change.

Encoding of attentional states in area V4

To understand better how V4 neurons encode a cognitive state associated with enhanced

attentional effort, we fitted single trial spike counts of every neuron using a generalized linear

model (GLM) as a function of experimental predictor variables (Methods). In a detailed GLM

(complete model; **Figure 4A**), we used 13 predictor variables: orientation of sample stimuli inside

RF; mean pupil area during the sample period; the saccadic choice; and a reward history that

consisted of reward outcomes in each of the previous 10 trials. We also compared the performance

of this complete model with two reduced GLMs (**Figure S10**) that included either reward history

(reward model) or the pupil area (pupil model) alone (**Figure 4A**). Fitted single trial spike counts

in small and large reward conditions of an example neuron are shown in **Figure 4C**. Model

performance was measured by pseudo-R^2 values (Methods). Most neurons (78%, 757/970) were

significantly fit (p < 0.05, F test) with the complete model. Fewer neurons had significant fits for

the reduced models (reward model, 12%, 114/970; pupil model, 50%, 483/970; **Figure 4B**).

Figure 4D shows population averaged spike counts of all fitted neurons estimated from the

complete GLM with overlying observed responses. We next compared the relative contributions

Figure 4. Encoding of cognitive states by single trial spike counts. (**A**) *Top*, Generalized linear models (GLM) on single trial spike counts during sample stimuli period (60 - 260 ms from sample onset). Spike counts were fit with three different GLMs. Two were reduced models, based on either reward history (10 immediate past received rewards; reward model) or mean pupil area over 400 ms following stimulus onset (pupil model). A more detailed model contained multiple experimental and cognitive state variables that included reward history (past 10 trials), pupil area, saccade response and Gabor stimulus orientation (complete model). (**B**) *Left*, Cumulative probability densities of pseudo R^2 values (Methods) of single

neurons measure goodness of fit for different GLMs. *Right*, fraction of the neuron population that were fitted significantly better ($p < 0.05$, F test) compared to a null model (a constant) [reward model, 114/ 970 (12%); pupil model, 483/ 970 (50%); complete, 757/ 970 (78%)]. (**C**) Observed and model predicted spike counts on single trials from cross-validation test-dataset for an example neuron in two representative reward blocks. (**D**) Population averaged observed and fitted (complete model) spike counts. Gray, observed; blue, complete model. (**E**) Comparing predictor importance (PI) that measures contributions of different predictor variables estimated by absolute standardized predictor coefficient values. Colormap represents PIs of all neurons that were fitted with the complete model ($n = 757$). Neurons were sorted based on the pseudo R^2 values. (**F**) Population averaged PIs. Error bars, 95% confidence intervals (bootstrap, $n = 10^4$).

of the experimental variables on spike responses for each neuron as measured by predictor

importance (PI), which is the normalized magnitude of fitted coefficients (Methods). Stimulus

orientation has the strongest contribution to the spike counts, which is expected because most V4

neurons are orientation selective (PIs of individual neurons, **Figure 4E**; mean PI across neurons,

Figure 4F). More interestingly, single trial pupil area (a physiological proxy of attentional effort)

also strongly predicts spike response compared to reward history and saccade choice.

Instantaneous pupil area remained a strong predictor when we compared with constant effort levels

('high' and 'low') as well as block-averaged single trial pupil area in two alternate models (PI,

mean ± SEM, constant effort, 2.3 ± 0.07; blocked-averaged pupil area, 2.7 ± 0.09, instantaneous

pupil area, 5.37 ± 0.19; ANOVA, $F_{(2, 2195)} = 166.31$, $p < 10^{-67}$; **Figure S11**). Together, these suggest

that even though reward history is the primary external motivator for the monkeys, V4 spike

responses to the sample stimuli encode instantaneous attentional effort more strongly than recent

reward outcome value or saccade choices.

It is possible that reward history has more weight around the time of block transitions, when

expectations change rapidly (Baruni et al., 2015). Because most of the trials within blocks in our

task occur long after a transition (steady-state), this weight would be diluted. To address this, we

also fit spike counts from only the first 20 trials after block transitions with the same complete-

GLM as shown in Figure 4A. (**Figure S12**). As with the complete data set, stimulus orientation

and pupil area contribute much more strongly to V4 spike counts than does reward history.

Although both the pupil area and spike counts are not directly related, they are linked to the top-

down attentional effort in a similar way. Thus, the attentional state as represented by pupil area is

encoded on single trial spike counts during the sample stimuli period.

Dynamic encoding of stimulus change-detection in area V4

Although reward representation in V4 is weak during the sample stimuli interval, stronger signals

might emerge closer to the behavioral response. We therefore separately fit spike counts during

the choice interval (test 1 interval) using a similar complete GLM. We used only those trials for

which test 1 stimulus appeared inside RFs and the animal did not initiate a saccade until after the

analysis period (which was 60-260 ms from test 1 stimulus onset; 44.4% (22238/50038) of the

trials) (**Figure S13**). The stimulus feature variable in this model was the product of orientation

tuning filter and test 1 stimulus orientation, as there were four different orientations of test 1

stimulus. The remaining variables were mean pupil area during the test 1 presentation, reward

outcomes in each of the previous 10 trials and the saccadic choice.

Reward history remained a poor predictor of single trial spike counts compared to pupil

area (**Figure S13**). Notably, the saccade choice variable is also a strong predictor of spike

responses for the same neuronal population. Previous reports have shown that spike responses of

V4 neurons enhance before initiating a saccade towards the RF location(Burrows et al., 2014;

Moore et al., 1998). Among all the saccade sensitive neurons ($n = 256$, significant saccade

coefficient, p < 0.05) about two-thirds were more active before saccades (saccade-preferred

neurons; saccade coefficient > 0; $n = 149/256$) and the rest were less active (saccade-anti-preferred

neurons, saccade coefficient < 0; $n = 107/256$). Further, based on behavioral outcomes (hit, miss,

CR and FA), we sorted observed spike counts on single trials during sample and test 1 stimulus

periods for these two neuron types, saccade preferred and saccade anti-preferred (**Figure S13**). A

true encoding of saccade choice by these neurons predicts indistinguishable spike response

between hits and FAs (and between CRs and misses). Instead, spike responses of saccade preferred

neurons increase for hits and misses (nonmatch trials) compared to CRs and FAs (match trials).

Similarly, responses of saccade anti-preferred neurons increase for CRs and FAs (match trials)

compared to hits and misses (nonmatch trials). This suggests that saccade choice is not a suitable

predictor variable in accounting single trial spike counts during the test 1 interval; rather 'stimulus

orientation-change' can be a better predictor parameter.

We next fit the same dataset of spike counts during test 1 stimulus with a complete-GLM

as mentioned above except, the saccadic choice variable was replaced with stimulus orientation

change (Δori) (*top*, **Figure 5A**). The variable Δori represents the trial type: match or non-match

trial. Covariance between single trial absolute test 1 stimulus orientation and Δori is small (mean

partial correlation 0.005) across sessions due to small Δori values ($16°$-$32°$). Further, we used only

the neurons with very weak correlation coefficients (< 0.2) between model-predicted spike counts

due to stimulus feature (product of orientation tuning filter and test 1 orientation) and Δori to rule

out stimulus orientation-related effects on estimates of Δori coefficient. V4 spike counts robustly

encode information of single trial Δori (**Figure 5A-B**). Based on significant Δori coefficients (p <

0.05), two populations of neurons were classified, Δori-preferred (Δori coefficient > 0, *n* = 130/

521) and Δori anti-preferred (Δori coefficient < 0, *n* = 109/ 521). As expected, orientation change

differentially modulates mean spike counts during the test 1 interval in these two neuron

subpopulations in accordance with the change in stimulus orientation [two factor repeated

measured ANOVA, $F_{(1, 237)} = 15.33$, $p = 10^{-4}$ for neuron types (Δori preferred, Δori anti-preferred);

$F_{(3, 711)} = 7.83$, $p < 10^{-4}$ for repeated measured factor response choices (hit, miss, CR and FA)]

(*right*, **Figure 5C**).

Figure 5. V4 spikes encode change in stimulus orientation immediately prior to response selection.
(**A**) *Top*, Δori model: GLM of spike counts during test 1 interval (60-260 ms from test 1 on). Predictor variables: product of test 1 stimulus orientation and neuron's orientation tuning filter, pupil area, reward history and orientation change (match or nonmatch trial). Only the trials when test 1 stimulus appeared inside RF of recorded neurons and monkey did not initiate any saccade before analysis period (44.4% of the total trails). *Bottom*, Contributions of predictor variables to spike counts of single neurons. (**B**) Population averaged predictor importance ($n = 521$). (**C**) Observed normalized mean V4 spike rates of Δori responsive neurons during sample and test 1 stimulus presentations for different behavior choice trials. H: hit, M: miss, R: correct rejection, F: false alarm. Neurons with significant Δori coefficient (p < 0.05) are classified as Δori-preferred (coefficient < 0, $n = 130$) or Δori-anti-preferred (coefficient < 0, $n = 109$) based on the GLM fits. (**D**) Decoded task variables on randomly selected pair of trials from cross-validation test-dataset in an example session. GLM fitted coefficients and observed single trial V4 spike counts are used to estimate probability of Δori (Δori model, Figure 5A) and probability of saccade (saccade model, Figure S11). Open red circle, observed response. Filled black circles, incorrect prediction. Dot, predicted probability of a response. (**E**) Decoding accuracy of response choice, saccade and Δori for cross validation test datasets across all sessions (N = 24) using GLM fits. Triangles, population means. Dashed line, 25% chance level. (**F**) Decoding accuracy as a function of number of neurons used for decoding across sessions (*N* = 24). Markers, individual session. Solid lines, second order polynomial fits. Dashed lines, 95% confidence intervals. Error bars, 95% confidence intervals.

Finally, we tested the detection accuracy of stimulus orientation change from observed V4 spiking

activity during the test 1 stimulus by using the fitted model on a single trial in a 10-fold cross-

validation test dataset (Methods). A random pair of trials, one from non-match (hit or miss) and

another from match (CR or FA) without replacement, were selected. For each of these two trials,

we evaluated the likelihoods of observed spike counts to be consistent with spike counts under the

two possible orientation changes for each fitted neuron within a session. Sum of the log-likelihood

ratios across neurons amounts to the predicted probability of a realization of orientation change on

that trial. Only if the decoded orientation changes on both trials of the selected pair match with the

observed data was the discrimination of an orientation-change from no-change considered correct.

Figure 5D shows model-fitted population-averaged spike counts and decoded task variables,

orientation change (*top*, using Δori GLM) and saccadic choice (*bottom*, using saccade GLM as

described in Figure S11) on 20 pairs of randomly selected trials in an example session. Similar to

saccade prediction, we also tested accuracy in discriminating response choices i.e. hit from miss

or FA from CR from observed spike counts during test 1 using the same saccade-GLM. Prediction

accuracy of orientation change (using Δori-GLM) is better than both saccade and choice (hit vs

miss or CR vs FA) predictions (using saccade-GLM) across all sessions (mean $\pm$ SEM, Δori, 68.5

$\pm$ 2.8%, saccade, 50.0 $\pm$ 1.4%, choice, 18.1 $\pm$ 1.8%; p = 10^{-25}, $F_{(2,69)}$ = 154.2, ANOVA; **Figure**

5E). Decoding performance of Δori is better compared to saccade irrespective of the number of

neurons used for decoding within a session (**Figure 5F**). Together, these encoding and decoding

models offer powerful methods in isolating task relevant variants and reading out perceptual or

decision signals from a population of neuronal spikes.

DISCUSSION

We found that isolated changes in attentional effort modulate the activity of neurons in V4. Previous reports on effort-related modulation in V4 (Boudreau et al., 2006; Spitzer et al., 1988) conflated changes in effort with changes in attentional selectivity (how much one location is favored over others), which strongly modulates V4 neuronal activity (Cohen and Maunsell, 2009; Luo and Maunsell, 2015; Moran and Desimone, 1985; Reynolds and Heeger, 2009). Compared to previous results, the current data is the first demonstration that V4 neurons are modulated to a comparable degree by either non-selective attentional effort or attentional selectivity. As a result of reward size changes, the block averaged cortical responses covary with physiological pupil area and behavioral performance (d'). However, these three measures dissociate at the block transitions and exhibit differential degrees of dependency on trial history. Stimulus features and pupil area strongly contribute to V4 spike counts throughout the trial duration as revealed by single trial information encoding. V4 spike counts contain very little information about immediate reward history. Additionally, perceptual detection of a task-relevant stimulus emerges as a strong contributor to V4 spiking during the decision period. This perceptual detection is independent of behavioral choice.

The rise of d' following an increase in reward is slow relative to the increase in V4 spike rate and pupil area. This may occur because performance in an attention demanding task depends not only on the state of sensory structures such as V4, but also on that in downstream decision and motor circuits. These downstream circuits could respond to changes in reward with their own dynamics. It remains to be determined how much subjects can control the dynamics of their responses to changes in reward expectation. Our design used no cueing other than reward size itself. It is possible that an explicit cue at the beginning of each trial might produce more coincident changes in neuronal spiking and d'. However, animal subjects typically achieve less pronounced

behavioral changes when trials of different reward contingencies are randomly interleaved on each

trial, and spatial cues at the start of the trial indicate the trial type (Baruni et al., 2015). Block-wise

control of animals' effort can be more effective for producing performance changes. Although a

limitation of the latter design is that the reward outcome and attentional state covary and make it

more challenging to dissociate true representation of neuronal modulation (for review (Maunsell,

2004)). Our task design of manipulating attentional states within small block of trials and using

single trial analysis was crucial for isolating relative contributions of attentional effort, reward

history and perceptual performance.

In contrast to reward increase, performance drops quickly with decreased reward even

though pupil area and neuronal response remain elevated (i.e. delayed decay, **Figure 3**). Dissimilar

effects of reward change directions on behavior are often seen across species (Kimchi and

Laubach, 2009; Saez et al., 2017). Independence of V4 neuronal activity from single trial

behavioral performance is consistent with the notion that transitions between states occur with

different dynamics in downstream structures that read out perceptual detection and weigh

anticipated outcomes against the cost of initiating and executing action (Baruni et al., 2015; Zénon

and Krauzlis, 2012).

The observed hysteresis in V4 spike responses with reward changes represents a form of

memory. Hysteresis at the input-output level of a single neuron can improve memory capacity and

retrieval properties in the presence of noise in a synchronous network of neurons with nonlinear

threshold (Wang and Ross, 1990). Indirect evidence of hysteresis in single cortical neurons comes

from the fact that spike train history is an important predictor of individual spikes (Park et al.,

2014; Pillow et al., 2008). Spike counts on reward-matched trials are indistinguishable between

the two reward conditions immediately after the reward switch and become different only in later

trials when the animal's effort attains a steady level (**Figure 3I**). These results together with

reward-spike hysteresis (**Figure 3G**) strongly support a dependency of V4 neuronal activity on

history of attentional effort state. Previous trial history of task relevant signals has also been

reported to strongly modulate PFC neurons (Donahue and Lee, 2015). Neuronal hysteresis during

visual perception has been suggested to arise from cooperativity between interconnected neurons

(Williams et al., 1986; You et al., 2011). Future experiments are required to dissociate the role of

single cell excitability and connectivity among neurons on neuronal hysteresis in V4. Absence of

reward modulation (immediate past trial outcome) of spike counts on reward-matched trials in

trials immediately after block transitions is inconsistent with immediate reward representation as

reported in previous studies (Baruni et al., 2015; Cicmil et al., 2015). Like V4, many other brain

areas show correlated spike activity with pupil area, including structures implicated in

attention(Joshi et al., 2016). A noradrenergic neuromodulation of V4 neurons might mediate such

brain-wide attentional state representation.

The relationships between reward expectation, behavioral d' and V4 spiking have also been

examined in a previous study (Baruni et al., 2015) that found the expectation of a large reward

increases spike rates of V4 neurons relative to expectation of a low reward, even when d' is the

same across those conditions. Consistent with this previous finding, we found that the d' was

slower to respond to changes in reward schedule than was V4 spiking. The early study randomly

interleaved different rewards conditions on a trial-by-trial basis, while our task contingencies were

entirely predictable over blocks of 120 trials. We would expect to see results qualitatively similar

to the previous report (Baruni et al., 2015) if we randomly interleaved trials. Unlike previously

observed positive correlation between absolute reward size expectation and V4 spiking (Baruni et

al., 2015), we found very weak representation of reward history in single-trial V4 spike counts

using both reward-matched spike count analysis and GLM-based encoding model, even

immediately after block transitions when the knowledge of reward expectancy was crucial for

guiding behavior. In our study, the reward on each trial varied around a mean within each reward

schedule (**Figure S1),** whereas a fixed reward was used for a given reward-schedule in the earlier

study. Our additional reward-variance provided negligible covariances between reward size and

other task variables, and was crucial for reliable isolation of multiplexed information on single-

trials using GLM. Prior application of GLM-based encoding in primate cortical areas has identified

dependencies of correlated spikes on multiplexed sensory, cognitive and motor signals (McGinty

et al., 2016; Park et al., 2014). Performance of our model fit can be further improved by taking

into account of spike history or interaction with other units within the network (Park et al., 2014).

Previous studies have described that V4 spiking increased when an animal intends to make

a saccade towards the neuron's RF (Burrows et al., 2014; Fischer and Boch, 1981a, b; Moore et

al., 1998). However, single trial encoding shows that the observed correlation between V4 spike

counts and saccade initiation is primarily due to perceptual signal detection and not saccade

preference. Spike rate (Shiozaki et al., 2012) and pairwise noise correlation (Cohen and Maunsell,

2009) between spike counts of V4 neurons have been suggested to represent a detection threshold

and predict choice probability in perceptual discrimination tasks. Ablation of V4 has revealed no

effects on saccade amplitude or latency to visual targets (Schiller and Lee, 1994). In partial

agreement with some of these findings, our results confirm the role of V4 in single trial perceptual

detection but rejects a direct contribution to saccade choice. High covariance between perceptual

detection and behavioral choice in previous tasks may have limited the detailed isolation of

information coding of perception from decision. In the present study, we identified two types of

response populations in V4 that are differentially modulated with perceptual signal changes. Future

work at the microcircuitry level is required for detailed characterization of these two cell

populations with respect to their inputs and recurrent connections. Understanding the finer

dynamics of cognitive task variants within a single trial would require modeling at the time scales

of single spikes (Park et al., 2014). A perceptual task with a continuous stream of visual stimuli

would be ideal to evoke a sufficient number of spikes throughout the task period for studying

visual areas like V4 that have low activity in the absence of visual stimulation. Overall, our

experimental framework provides new evidence revealing information encoding of higher brain

function that can be extended to other brain areas.

METHODS

Subjects and surgery

All experimental procedures were approved by the Institutional Animal Care and Use Committee

at the University of Chicago and were in compliance with US National Institutes of Health

guidelines. Two adult male rhesus monkeys (*Macaca mulatta*, 13 and 9 kg) were implanted with

a titanium head post using aseptic surgical techniques before training began. After the completion

of behavioral training (3 to 5 months), we implanted an 10x10 array microelectrodes with 400 μm

spacing (Blackrock Microsystems) into dorsal visual area V4 of one hemisphere, between lunate

and superior temporal sulci.

Behavioral task

During training and neurophysiological recording, the monkey was seated in a primate chair facing

a calibrated CRT display (1024 x 768 pixels, 100 Hz refresh rate) at 57 cm viewing distance inside

a darkened room. Binocular eye position and pupil area were recorded at 500 Hz using an infrared

camera (Eyelink 1000, SR Research). Trials started once the animal fixated within 1.5° of a central

white spot (0.1° square) presented on a mid-level gray background (**Figure 1A**). The animal had

to maintain fixation until its response at the end of the trial. After a fixation period of 400-800 ms,

two achromatic Gabor sample stimuli appeared for 200 ms, one in each visual hemifield. After a

variable delay of 200-300 ms, a Gabor test stimulus (test 1) appeared for 200 ms at one of the two

target locations, randomly selected with equal probability. The test stimulus was identical to the

preceding sample stimulus, except for its orientation. On half of the trials, the test 1 stimulus had

a different orientation (nonmatch trial) and the monkey had to make a saccade to that target

location to receive an apple juice reward. On the remaining half of the trials, the test 1 stimulus

had the same orientation as the corresponding sample stimulus (match trial), and the monkey had

to maintain fixation until a second test stimulus with a different orientation (test 2, 200 ms)

appeared in the same location after an additional delay interval of 200-300 ms. The monkey then

had to saccade to that target to get a reward. Inter-trial intervals varied from 2-3 s. Stimuli were

presented always in the lower hemifields at 2°-4° eccentricity. Gabors were odd-symmetric with

the same average luminance as the background. Spatial frequency, size and base orientation of

Gabor stimuli were optimized for one neuron recorded each day, and remained unchanged

throughout each session (left, azimuth –2.5° to –5.3°, elevation 0.0° to –3.5°, sigma, 0.35° to 0.60°,

spatial frequency 0.6 to 3.5 cycles/°; right, azimuth 2.8° to 5.5°, elevation 0.0° to –2.5°, sigma,

0.40° to 0.65°, spatial frequency 0.7 to 3.0 cycles/°). On every trial, the orientation of the sample

stimuli randomly took one of two values, base orientation or orthogonal. Orientation change

(difficulty) remained fixed within a session and varied across sessions between 24°-32° for monkey

S and 16°-20° for monkey P.

Attentional effort was controlled over blocks of trials by changing reward volumes for

correct responses over a 4-5-fold range. Every 120 trials, reward size alternated between large and

small values without any prior cue to the animal. Reward sizes for hits (correct response in

nonmatch trial) and CRs (correct rejections in match trial) were adjusted as needed to encourage

the animal to maintain a behavioral criterion close to zero(Luo and Maunsell, 2015). Although,

trial averaged criterion within a reward condition remained zero, there was a transient change in

criterion immediate after the block transition (reward switch) which then approached zero over a

few trials (**Figure S5**). Distributions of reward sizes are shown in **Figure S1**.

Electrophysiological Recording and Data Collection

Extracellular neuronal signals from the chronically implanted multielectrode array were amplified,

bandpass filtered (250–7,500 Hz) and sampled at 30 kHz using a Cerebus data acquisition system

(Blackrock Microsystems). We simultaneously recorded from multiple single units as well as

multiunits (563, monkey S; 407, monkey P) over 24 sessions (15 for monkey S, 9 for monkey P).

Before each experimental session, we mapped receptive fields and stimulus preferences of neurons

while the animal fixated. These fields were used to optimize the stimulus parameters. Spikes from

each electrode were sorted offline (Offline-Sorter, Plexon) by manually defining cluster

boundaries using principal component as well as waveform features. Well isolated clusters based

on J3 statistics were classified as single units (Nicolelis et al., 2003). J3 measures the ratio of the

average distance between points within clusters to the average distance between clusters. It takes

a maximum value for compact, well-separated clusters.

Data Analysis

Behavioral performance

All completed trials (120 trial per block) were included in our analysis. Behavioral sensitivity (*d'*)

and criterion (*c*) were measured from hit rates within nonmatch trials and FA rates within match

trials as:

$$d' = \Phi^{-1}(hit\ rate) - \Phi^{-1}(FA\ rate) \tag{1}$$

$$c = -\frac{1}{2}[\Phi^{-1}(hit\ rate) + \Phi^{-1}(FA\ rate)] \tag{2}$$

where Φ^{-1} is inverse normal cumulative distribution function. We measured average *d'* and *c*

within a session across all trials for large- and small-reward blocks separately. To examine the

dynamics of d', c, and percent-correct at i^{th} trial (i = 1 to 120), we measured block averaged values

for each reward conditions. Overall d' was measured by,

$$d'_{overall} = \sqrt{(d'^2_{InRF} + d'^2_{OutRF})} \tag{3}$$

Where, d'_{InRF} and d'_{OutRF} are the sensitivities in the two hemifields, inside and outside the recorded

neurons' receptive fields.

Pupil area

All pupil area measurements were measured binocularly at 500 Hz while monkeys maintained

fixation in absence of a luminosity change using infrared camera (EyeLink 1000, SR Research).

Raw pupil areas were normalized for each session and each eye separately. Mean pupil area was

measured by averaging the normalized pupil area over 400 ms from sample appearance.

Neuronal response

Only neurons with an average spike rate 60-260 ms after sample stimulus onset that was

significantly ($p < 0.01$) greater than the rate 0 to 250 ms before sample onset were used in the

analysis. To construct peri-stimulus time histograms (PSTHs) for figures, spike trains were

smoothed with a half Gaussian kernel (15 ms SD with only a rightward tail), aligned to sample

stimuli onset and averaged across trials. A spike rate modulation index (MI) for each neuron was

calculated as the difference in average spike counts (60–260 ms after sample onset) between large

and small reward blocks (correct trials) divided by their sum.

$$MI = \frac{\langle Spike\ count_{large}\rangle - \langle Spike\ count_{small}\rangle}{\langle Spike\ count_{large}\rangle + \langle Spike\ count_{small}\rangle} \tag{4}$$

For analyzing trial-by-trial dynamics of spike counts and GLM, absolute spike counts within 60 to

260 ms from sample stimuli onset (or test 1 onset) were used.

Principal component analysis (PCA) was done on spike rates from 0-400 ms separately for

each monkey in Matlab. Each neuron had two spike rates, one each for one type of reward block

(small or large reward). For monkey P, there were 407 neurons (814 spike trains) and for monkey

S, there were 563 neurons (1126 spike trains). Linear regression was fitted between PC scores of

spike histograms in small and large reward blocks for the first three principal components.

Pupil area-spike count-effort linear regression

Average pupil area over 0-400 ms from sample onset and spike counts from 60-260 ms on each

individual trial were used for linear regression analysis for pupil area *versus* spike count and

average effort states ("low" and "high") (**Figure 3J**, *left*). The pupil area was modeled as: *pupil*

area ~ *a**(*spike count*) + *b**(*effort*). Where *a* and *b* are the regression coefficients. All the trials

across small and large reward conditions that occurred after the first correct response (hit or CR)

following the reward switch were included in the analysis. In a second model: *pupil area* ~

a*(*spike count*), each neuron was fitted separately for two reward conditions, small and large

(**Figure 3J**, *right*). For this model, we used a subset of trials with matched pupil areas between the

two reward conditions. Standardized coefficients were compared across neurons that were fit at

significant level of $p < 0.05$ (F-test).

Pupil area-spike count cross-correlation

We measured cross-correlations between two time series, spike rates and pupil area as a function

of time lag over 700 ms period around the sample stimulus on (−350 ms to 350 ms) for each neuron.

Single trial spike trains were binned in a 10 ms sliding window (2 ms increments) and converted

to spike rates. Pupil area time series was sampled at 500 Hz and directly used for the cross-

correlation analysis. Single trial cross-corrrelations were averaged, and trial-shuffled values were

subtracted separately for trials with small and large reward conditions.

Spike triggered averaged pupil area

We measured a spike triggered averaged (STA) pupil area for each neuron within a time window

of 0-400 ms from sample stimulus onset, across all trials. The time series of pupil areas were

aligned to individual spikes, and averaged. Finally, trial-shuffled subtracted STA was subtracted

from this averaged STA-pupil area for each neuron.

Generalized linear model (GLM)

Generalized linear model (GLM) regression was used to estimate the relationship between single

trial spike responses and reward expectancies, attentional efforts, behavior choices and stimulus

parameters. Single trial stimulus evoked spike counts were modeled to follow a negative binomial

distribution. The negative binomial distribution is well suited for the purpose, as spike count

variances of cortical neurons are most often equal to or greater than their means (**Figure S8**)

(Churchland et al., 2010; Shadlen and Newsome, 1998). Reward expectancy was represented by

reward history on 10 trials preceding the trial being considered. Mean pupil area during 400 ms

following stimulus onset served as a proxy for attentional effort. A categorical saccade choice

variable could take two values: saccade (hit and FA) or no saccade (CR and miss). For fitting of

test 1 spike counts, an alternate categorical task variable "stimulus orientation change" (Δori) was

used instead (**Figure 5A**). This perceptual variable had two values: orientation change (nonmatch

trial) or no orientation change (match trial). The product of the absolute orientation of Gabor

stimulus and neuron's orientation tuning filter served as a test 1 stimulus feature variable. There

were four different orientations of test 1 stimulus. There were no correlations (partial) among

predictor variables (**Figure S9**). For the GLMs on test 1 spikes, in addition to the predictor variable

covariances, we also cross checked the correlation between model predicted spike counts due to

test 1 stimulus feature and Δori in order to isolate the orientation-tuning related effect on the

perceptual variable Δori. We considered only the neurons that showed correlation (Spearman) of

<0.2 for further analysis.

For a sample $y_1, y_2,....,y_n$ of independent response variable y from an exponential family, there

exists a linear predictor $\beta_0 + \beta_1 x_1 +....+\beta_k x_k$ of the response variable y. The mean response

$\left(\mu_y = E[y|x_1,.., x_k]\right)$ depends on the linear predictor through a link function $g(\cdot)$:

$$g\left(\mu_y\right) = \beta_0 + \beta_1 x_1 + \cdots + \beta_k x_k \tag{5}$$

Where, x_j ($j = 1,2,…,k$) is a set of independent predictor variables. Assuming that the spike count

responses in non-overlapping time intervals are independent and variance is equal or greater than

(over dispersion) the mean, they can be modeled as negative binomial variables. As the mean spike

count is always positive, it can be modeled as:

$$\mu_y = exp\left(\beta_0 + \beta_1 x_1 + \cdots + \beta_k x_k\right) \tag{6}$$

The probability density function for a random variable for which there are y successes of Bernoulli

trials until θ^{th} failure occurs follows a negative binomial distribution:

$$P\left(y; \mu_y, \theta\right) = \frac{\Gamma(y+\theta)}{\Gamma(\theta)y!} \frac{\mu_y{}^y \theta^\theta}{(\mu_y+\theta)^{y+\theta}} \tag{7}$$

with a mean μ_y and variance $V(y)$:

$$\mu_y = \frac{\theta p}{1-p} \tag{8}$$

$$V_y = \mu_y + \frac{\mu_y{}^2}{\theta} \tag{9}$$

where, p is probability of success of each Bernoulli trial. $1/\theta$ is the dispersion parameter (θ is also

called as shape parameter). Γ is gamma distribution. The likelihood for a negative binomial random

variable y_i ($i = 1,2,…,n$) that depends on a set of k predictors of $x_{i,j}$ ($j = 1,2,…,k$):

$$L(P) = \prod_{i=1}^{n} \frac{\Gamma(y_i+\theta)}{\Gamma(\theta)\, y_i!} \frac{\mu_{y_i}{}^{y_i} \theta^\theta}{\left(\mu_{y_i}+\theta\right)^{y_i+\theta}} \tag{10}$$

The log-likelihood is:

$$LL(\beta_0, \beta_1, …, \beta_k, \theta) = log\left(L(P)\right) \tag{11}$$

584 The predictor coefficients β_j ($j = 1,2,…,k$) and shape parameter θ are obtained from maximum

585 likelihood estimates by solving:

$$\frac{\partial LL}{\partial \beta_j} = 0, \ j = 1,2, …, k \tag{12}$$

$$\frac{\partial LL}{\partial \theta} = 0 \tag{13}$$

588 GLM was implemented in Matlab separately for each neuron. In order to compare different

589 predictor coefficients, they were converted post-hoc into standardized coefficients:

$$\beta_j' = \beta_j * \frac{\sigma_y}{\sigma_x}, \ j = 1,2, …, k \tag{14}$$

591 Standardized coefficient corresponds to the *log* ratios of the mean responses (in units of standard

592 deviation) due to one standard deviation change in the predictor variable, holding all other

593 variables constant.

594 <u>Goodness of GLM fit and model comparison:</u> Goodness of fit for a given GLM was measured by

595 residual deviance (D) and pseudo R-squared (Cragg & Uhler's method) value. Residual deviance

596 measures how close the predicted values from the fitted model match the actual values from the

597 raw data:

$$D = 2[LL(saturated\ model) - LL(model)] \tag{15}$$

599 The saturated model comprises a model for which the number of parameters equals the number of

600 samples. Thus, the predicted response in the saturated model is same as the observed response.

601 Deviance measures lack of fit in maximum log-likelihood estimations similar to the residual

602 variance in ordinary least square methods. The lower the deviance, better is the model fit. For a

603 large sample size and small deviance, it approximately follows chi-square distribution of $n–k–1$

604 degrees of freedom. A statistic that indicates value of a proposed model is based on how well the

605 model predicts data compared to a null model having a single predictor β_0:

$$\chi^2 = D(null\ model) - D(model) \tag{16}$$

607
$$= 2[LL(model) - LL(null\ model)] \tag{17}$$

608 The test statistic χ^2 follows an approximate chi-square distribution of degrees freedom n–k. We

609 tested at a level of significance $p < 0.05$.

610 As the ratio of the likelihoods reflects the improvement of the proposed model over the null model

611 (the smaller the ratio, the greater the improvement), a pseudo R-squared was computed based on

612 Cragg & Uhler's approach that falls between 0 and 1:

613
$$R^2 = \frac{1 - \left[\frac{L(null\ model)}{L(model)}\right]^{2/n}}{1 - L(null\ model)^{2/n}} \tag{18}$$

614 where, n is the number of observations in the data set. Different models were compared by their

615 R-squared values.

616 Predictor importance: The absolute value of the z-statistic of each estimated predictor coefficient

617 measures a relative importance of that predictor variable:

618
$$z_j = \left|\frac{\beta_j}{SE(\beta_j)}\right|, \ j = 1,2,\dots,k \tag{19}$$

619 where, SE is the standard error.

620 Cross validation: Predictive performance of the GLMs was measured by cross validation.

621 Observations in each neuron's dataset were split at random into K partitions. GLM fit on K-1

622 training partitions was repeated and the remaining partition was used for validation. This cross

623 validation was repeated K times at each time one of the partitions served as the validation set.

624
$$Error = \frac{1}{n}\sum_{k=1}^{K}\sum_{i=1}^{n_k}(y_{i,k} - \hat{y}_{i,k})^2 \ k \tag{20}$$

Decoding of saccade selection, response choice and orientation change detection: Saccade

selection between saccade (hit or FA) and no-saccade (miss or CR) was decoded from spike counts

during the test 1 stimulus period based on Bayesian inference of maximum posterior probability

using saccade-GLM (**Figure 5D**). According to Bayes' rule, the posterior probability density of

the estimated choice given a spike count response is:

$$P(x|Y) = \frac{P(Y|x)P(x)}{P(Y)} \, k \qquad\qquad (21)$$

where, $P(Y)$ is a normalization term independent of x. Also, assuming that there is no prior

knowledge on x, (P (x) = constant), maximizing the posterior is equivalent to maximizing the log-

likelihood function used in the GLM encoding model. A random pair of trials one from saccade

trial (hit or FA) and another from no-saccade trial (CR or miss) were selected. For each of these

two trials, we evaluated the likelihoods of observed spike counts to be consistent with spike counts

under the two possible saccade responses for each fitted neuron within a session. Sum of the log-

likelihood ratios across neurons amounts the predicted probability of a realization of saccade

choice on that trial. If the decoded saccade commitment on both trials of the selected pair match

with the observed data, it is only then considered as correct discrimination of a saccade from no-

saccade. For choice selection, a pair of trials were randomly selected either hit and miss

(nonmatch) or FA and CR (match trial). Prediction accuracy was measured for correct

discrimination of hit from miss or FA from CR using saccade-GLM. Similarly, prediction accuracy

for orientation change detection was estimated using Δori-GLM. The chance level of prediction is

0.25.

Statistical analysis

Unless otherwise specified, we used paired t-test and multifactor repeated measured ANOVA for

comparing normally distributed datasets. Normality was checked with Kruskal-Wallis test.

REFERENCES

Baruni, J.K., Lau, B., and Salzman, C.D. (2015). Reward expectation differentially modulates
attentional behavior and activity in visual area V4. Nat Neurosci 18, 1656.

Beatty, J. (1982). Task-evoked pupillary responses, processing load, and the structure of
processing resources. Psychol Bull 91, 276.

Boudreau, C.E., Williford, T.H., and Maunsell, J.H. (2006). Effects of task difficulty and target
likelihood in area V4 of macaque monkeys. J Neurophysiol 96, 2377-2387.

Burrows, B.E., Zirnsak, M., Akhlaghpour, H., Wang, M., and Moore, T. (2014). Global selection
of saccadic target features by neurons in area V4. J Neurosci 34, 6700-6706.

Churchland, M.M., Byron, M.Y., Cunningham, J.P., Sugrue, L.P., Cohen, M.R., Corrado, G.S.,
Newsome, W.T., Clark, A.M., Hosseini, P., and Scott, B.B. (2010). Stimulus onset quenches
neural variability: a widespread cortical phenomenon. Nat Neurosci 13, 369.

Cicmil, N., Cumming, B.G., Parker, A.J., and Krug, K. (2015). Reward modulates the effect of
visual cortical microstimulation on perceptual decisions. Elife 4, e07832.

Cohen, M.R., and Maunsell, J.H.R. (2009). Attention improves performance primarily by reducing
interneuronal correlations. Nat Neurosci 12, 1594-1600.

Donahue, C.H., and Lee, D. (2015). Dynamic routing of task-relevant signals for decision making
in dorsolateral prefrontal cortex. Nat Neurosci 18, 295-301.

Fischer, B., and Boch, R. (1981a). Enhanced activation of neurons in prelunate cortex before
visually guided saccades of trained rhesus monkeys. Exp Brain Res 44, 129-137.

Fischer, B., and Boch, R. (1981b). Selection of visual targets activates prelunate cortical cells in
trained rhesus monkey. Exp Brain Res 41, 431-433.

Joshi, S., Li, Y., Kalwani, R.M., and Gold, J.I. (2016). Relationships between pupil diameter and
neuronal activity in the locus coeruleus, colliculi, and cingulate cortex. Neuron 89, 221-234.

Kimchi, E.Y., and Laubach, M. (2009). Dynamic encoding of action selection by the medial
striatum. J Neurosci 29, 3148-3159.

Kleinschmidt, A., Büchel, C., Hutton, C., Friston, K.J., and Frackowiak, R.S. (2002). The neural
structures expressing perceptual hysteresis in visual letter recognition. Neuron 34, 659-666.

Laeng, B., Sirois, S., and Gredebäck, G. (2012). Pupillometry: a window to the preconscious?
Perspect Psychol Sci 7, 18-27.

Luo, T.Z., and Maunsell, J.H.R. (2015). Neuronal modulations in visual cortex are associated with
only one of multiple components of attention. Neuron 86, 1182-1188.

Maunsell, J.H.R. (2004). Neuronal representations of cognitive state: reward or attention? Trends
Cogn Sci 8, 261-265.

McGinty, V.B., Rangel, A., and Newsome, W.T. (2016). Orbitofrontal Cortex Value Signals
Depend on Fixation Location during Free Viewing. Neuron 90, 1299-1311.

Moore, T., Tolias, A.S., and Schiller, P.H. (1998). Visual representations during saccadic eye
movements. Proc Natl Acad Sci 95, 8981-8984.

Moran, J., and Desimone, R. (1985). Selective attention gates visual processing in the extrastriate
cortex. Front Cogn Neurosci 229, 342-345.

Nicolelis, M.A., Dimitrov, D., Carmena, J.M., Crist, R., Lehew, G., Kralik, J.D., and Wise, S.P.
(2003). Chronic, multisite, multielectrode recordings in macaque monkeys. Proc Natl Acad Sci
100, 11041-11046.

Padoa-Schioppa, C., and Assad, J.A. (2006). Neurons in the orbitofrontal cortex encode economic
value. Nature 441, 223.

Park, I.M., Meister, M.L., Huk, A.C., and Pillow, J.W. (2014). Encoding and decoding in parietal
cortex during sensorimotor decision-making. Nat Neurosci 17, 1395.

Pillow, J.W., Shlens, J., Paninski, L., Sher, A., Litke, A.M., Chichilnisky, E., and Simoncelli, E.P.
(2008). Spatio-temporal correlations and visual signalling in a complete neuronal population.
Nature 454, 995.

Piquado, T., Isaacowitz, D., and Wingfield, A. (2010). Pupillometry as a measure of cognitive
effort in younger and older adults. Psychophysiology 47, 560-569.

Reynolds, J.H., and Heeger, D.J. (2009). The normalization model of attention. Neuron 61, 168-
185.

Roesch, M.R., and Olson, C.R. (2004). Neuronal activity related to reward value and motivation
in primate frontal cortex. Science 304, 307-310.

Rorie, A.E., Gao, J., McClelland, J.L., and Newsome, W.T. (2010). Integration of sensory and
reward information during perceptual decision-making in lateral intraparietal cortex (LIP) of the
macaque monkey. PloS one 5, e9308.

Saez, R.A., Saez, A., Paton, J.J., Lau, B., and Salzman, C.D. (2017). Distinct roles for the amygdala
and orbitofrontal cortex in representing the relative amount of expected reward. Neuron 95, 70-
77. e73.

Schiller, P.H., and Lee, K. (1994). The effects of lateral geniculate nucleus, area V4, and middle
temporal (MT) lesions on visually guided eye movements. Vis Neurosci 11, 229-241.

Schultz, W., Apicella, P., Scarnati, E., and Ljungberg, T. (1992). Neuronal activity in monkey
ventral striatum related to the expectation of reward. J Neurosci 12, 4595-4610.

Shadlen, M.N., and Newsome, W.T. (1998). The variable discharge of cortical neurons:
implications for connectivity, computation, and information coding. J Neurosci 18, 3870-3896.

Shiozaki, H.M., Tanabe, S., Doi, T., and Fujita, I. (2012). Neural activity in cortical area V4
underlies fine disparity discrimination. J Neurosci 32, 3830-3841.

Spitzer, H., Desimone, R., and Moran, J. (1988). Increased attention enhances both behavioral and
neuronal performance. Science 240, 338-340.

Tremblay, L., Hollerman, J.R., and Schultz, W. (1998). Modifications of reward expectation-
related neuronal activity during learning in primate striatum. J Neurophysiol 80, 964-977.

Wang, L., and Ross, J. (1990). Synchronous neural networks of nonlinear threshold elements with
hysteresis. Proc Natl Acad Sci 87, 988-992.

Williams, D., Phillips, G., and Sekuler, R. (1986). Hysteresis in the perception of motion direction
as evidence for neural cooperativity. Nature 324, 253.

You, H., Meng, Y., Huan, D., and Wang, D.-H. (2011). The neural dynamics for hysteresis in
visual perception. Neurocomputing 74, 3502-3508.

Zénon, A., and Krauzlis, R.J. (2012). Attention deficits without cortical neuronal deficits. Nature
489, 434.

ACKNOWLEDGEMENTS

We thank Jackson J. Cone, Zaina Zayyad, Julian Day-Cooney and Thomas Z. Luo for discussion and comments on the manuscript. This work was supported by NIH grant R01EY005911.

AUTHOR CONTRIBUTIONS

S.G. and J.H.R.M. designed the experiments, performed the surgeries and wrote the paper. S.G. performed the experiments and analyzed the data.

DECLARATION OF INTERESTS

The authors declare no competing interests.