***N*-chlorination mediates protective and immunomodulatory effects of oxidized human**

plasma proteins

Running title: Effects of *N*-chlorinated plasma proteins

Agnes Ulfig, Anton V. Schulz, Alexandra Müller, Natalie Lupilov, and Lars I. Leichert

Ruhr University Bochum, Institute of Biochemistry and Pathobiochemistry – Microbial

Biochemistry, Universitätsstrasse 150, 44780 Bochum, Germany

Correspondence: Lars I. Leichert

12   Ruhr University Bochum

13   Institute of Biochemistry and Pathobiochemistry

14   Microbial Biochemistry

15   Universitätsstrasse 150

16   44780 Bochum, Germany

17   Phone: +49 234 3224585

18   Fax: +49 234 3214332

19   E-mail: lars.leichert@ruhr-uni-bochum.de

ABSTRACT

Hypochlorous acid (HOCl), a powerful bactericidal and fungicidal oxidant, is produced by

neutrophils to fight infections. To prevent tissue damage at sites of inflammation, excess

HOCl is scavenged by serum albumin and other plasma proteins. Here we show that plasma

proteins are not only an effective sink for HOCl, but that HOCl-induced *N*-chlorination

converts plasma proteins into chaperone-like holdases that protect other proteins from

aggregation. This chaperone-like conversion was reversible by antioxidants and was

abrogated by prior methylation of basic amino acids. Furthermore, reversible *N*-chlorination

of basic amino acid side chains is the major factor that converts plasma proteins into efficient

activators of immune cells. Finally, HOCl-modified serum albumin was found to act as a pro-

survival molecule that protects neutrophils from cell death induced by highly immunogenic

foreign antigens. We propose that activation and enhanced persistence of neutrophils

mediated by HOCl-modified plasma proteins, resulting in the increased and prolonged

generation of ROS, including HOCl, constitutes a potentially detrimental positive feedback

loop that can only be attenuated through the reversible nature of the modification involved.

INTRODUCTION

Recruitment and activation of neutrophils at sites of infection is considered one of the principal mechanisms by which the human body protects itself against diseases. The killing strategy of neutrophils involves the ingestion of pathogens into the phagosome, accompanied by the production of a diverse set of so-called reactive oxygen species (ROS), including superoxide anions ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2) and hypochlorous acid (HOCl) in a process known as the respiratory burst (recently reviewed in refs. [1,2]).

HOCl, a major inflammatory ROS, is produced from hydrogen peroxide and chloride ions by the heme enzyme myeloperoxidase (MPO) [3,4]. The antimicrobial properties of HOCl are well documented and numerous reports have provided strong evidence for severe damage to bacterial components within the neutrophil phagosome [5–7].

A particularly important target of HOCl are proteins. In proteins, HOCl exposure typically leads to side chain modification [8–12], fragmentation [13,14], misfolding/aggregation [15] or intermolecular di-tyrosine cross-linking, a hallmark of HOCl-oxidized proteins [16]. This often leads to their aggregation [17–19]. Consequently, various mechanisms to counter the accumulation of these misfolded proteins under HOCl stress exist in bacteria. These are mediated by chaperones that are activated by HOCl, the very same reactive species they protect against. One of the first protective proteins to be noticed was Hsp33, which gets reversibly activated by HOCl through oxidation of four critical cysteine residues [19]. This is not surprising, as cysteine thiols together with methionine residues react rapidly with HOCl [20,21]. More recently, we found that *E. coli* RidA, a member of the highly conserved, but functionally diverse YjgF/YER057c/UK114 protein family, also undergoes HOCl-based conversion into a chaperone holdase. This chaperone was highly active as protector of proteins from HOCl-induced aggregation. However in this case, instead of cysteine oxidation, *N*-chlorination of basic amino acids was the mechanism of its activation [17], although reactivity of HOCl with side chains of lysine and arginine is four to seven orders of

magnitude lower than that of cysteine [22]. But much like cysteine oxidation, *N*-chlorination is a

reversible oxidative modification that can be removed by cellular antioxidants such as

ascorbate, glutathione or thioredoxin and thus can switch the holdase function of RidA off

[17,23,24].

Similar observations were made more recently with *E. coli* CnoX (YbbN) that, when

activated via *N*-chlorination, binds to and prevents a variety of substrates from aggregation

and being irreversible oxidized [25].

While HOCl is highly bactericidal, generation of HOCl by immune cells is not without risk to

the human body itself (reviewed in refs. [26–29]). During inflammatory processes, up to 30% of

total cellular MPO is secreted by neutrophils into extracellular surroundings via

degranulation, leakage during phagocytosis, or by association with NETs [30]. Neutrophils,

accumulated in the interstitial fluid of inflamed tissues, have been reported to generate HOCl

at concentrations of up to 25-50 mM per hour [31]. It is thus not surprising that HOCl can also

drastically increase the activity of the extracellular chaperone α_2-macroglobulin (α_2M) in

human blood plasma to counteract the aggregation of host proteins under hypochlorous acid

stress [32]. However, the underlying molecular mechanism of HOCl-mediated conversion of

this plasma protein is still unclear.

And while neutrophils produce high amounts of HOCl, it does not accumulate at those levels,

as it reacts instantly with diverse biological molecules including proteins, DNA [33], cholesterol

[34] and lipids [35]. But due to their high abundance in blood and interstitial fluid, human serum

albumin (HSA) and other plasma proteins are considered the major targets of HOCl-mediated

damage and as such constitute the main sink for HOCl in the vicinity of inflammation [36–40].

The resulting products of the reaction of plasma proteins with hypochlorous acid, known as

advanced oxidation protein products (AOPPs), have, therefore, been employed as *in vivo*

markers of chronic inflammation [41]. Accumulation of AOPPs has been first discovered in

patients with chronic kidney disease [41] and later also found in a variety of other inflammatory

diseases, e.g. cardiovascular disease, neurodegenerative disorders, rheumatoid arthritis and

some cancers (recently reviewed in ref. [42]).

To date, a number of studies have been carried out to elucidate the role of AOPPs in

inflammatory processes [43–46]. Accumulating experimental evidence supports a critical

contribution of AOPPs to the progression of inflammation [44]. HOCl-modifed HSA

accumulates in inflammatory diseases and was found to act as proinflammatory mediator by

increasing oxidative stress and inflammation through stimulation of leukocytes [46,47].

Based on these findings we hypothesized that reversible *N*-chlorination in response to HOCl-

stress could be the principal chemical modification contributing to the observed physiological

properties of AOPPs. Furthermore, as we observed in previous studies that the HSA

homologue bovine serum albumin (BSA) could be transformed into a chaperone-like holdase

by HOCl-treatment [17], we speculated that *N*-chlorination serves as general rather than specific

mechanism to transform certain proteins into a chaperone-like state.

Here, we study the effects of reversible *N*-chlorination on the function of human plasma

proteins. We show that, upon *N*-chlorination, not only α_2M, but all plasma fractions tested,

exhibit chaperone-like activity and as such could prevent the HOCl-induced formation of

protein aggregates at the site of inflammation. Moreover, exposure to HOCl at concentrations

present in chronically inflamed tissues turned the majority of plasma proteins into efficient

activators of neutrophil-like cells. Previous studies revealed that HOCl-treated HSA can

stimulate leukocytes to produce more ROS during inflammation [47,48]. Now we find that

reversible *N*-chlorination is the main chemical modification that mediates the activation of

NADPH oxidase-dependent ROS generation by immune cells by AOPPs. Finally, we show

that HOCl-modified HSA is a pro-survival factor for immune cells and protects neutrophils

from cell death by highly immunogenic antigens.

Our data strongly suggest that *in vivo* reversible *N*-chlorination of human plasma proteins not

only converts them into effective chaperone-like holdases but is also the principal mechanism

that turns these proteins into mediators of the innate immune system.

MATERIAL AND METHODS

Preparation of plasma protein solutions

Albumin from human serum (HSA, Product # A9511), human γ-globulins (Product # G4386),

immunoglobulin G from human serum (IgG, Product # I4506) Cohn fraction IV (Product #

G3637) and human whole serum (sterile-filtered from male AB plasma, Product # H4522)

were purchased from Sigma-Aldrich, St. Louis, USA, and used without further purification.

Protein stock solutions were freshly prepared by dissolving or diluting varying amounts of the

protein fractions in 1xPBS buffer, pH 7.4 (Gibco Life Sciences).

Purification of α$_2$-macroglobulin

α$_2$-Macroglobulin was purified from human plasma (obtained from Zen-Bio, Inc., North

Carolina, USA, Product # SER-SLP, Lot # 11108) by the method of Imber and Pizzo, 1981 [49]

with slight modifications. Briefly, 880 mL fresh-frozen human plasma were thawed on ice

and dialyzed against frequent changes of deionized water for 72 hours at 4 °C using a

Spectra/Por dialysis membrane with a MWCO of 12-14 kDa (Spectrum Laboratories Inc.,

Rancho Dominguez, CA). Insoluble material was removed by centrifugation for 30 minutes at

10,000 x g and 4 °C. The supernatant plasma solution (200 mL) was then dialyzed for 24

hours at 4 °C against 5 L 1xPBS pH 6.0. Metal chelate chromatography was performed at 4

°C using an IMAC zinc-Sepharose 6 Fast Flow column (2.6 x 20 cm, GE Healthcare Life

Sciences, Amersham, UK) equilibrated with 1xPBS pH 6.0. Dialyzed plasma was applied to

the column and washed with 1xPBS pH 6.0 until the absorption at 280 nm of the eluant

(measured with a JASCO V-650 UV/VIS spectrophotometer (JASCO, Tokyo, Japan))

reached a value lower than 0.01. Bound protein was then eluted from the column with 0.01 M

NaOAc, 0.15 M NaCl, pH 5.0. Peak protein fractions were combined and concentrated using

Vivaspin 20 concentrators with a MWCO of 100 kDa. Gel filtration of the concentrated

protein pool fraction was performed at 4 °C on a HiPrep 26/60 Sephacryl S-300 High

resolution column (2.6 x 60 cm) equilibrated with 1xPBS pH 7.4. High molecular weight

peak fractions containing α_2-macroglobulin were combined, concentrated and dialyzed

against 1xPBS pH 7.4 containing 40% (v/v) glycerol. Aliquots were stored at -20° C.

Determination of protein concentrations

Protein concentration in g · L^{-1} of human serum was calculated using Pierce^{TM} Bicinchoninic

Acid (BCA) Protein Assay Kit (Thermo Fisher Scientific, Waltham, Massachusetts, USA)

with bovine albumin as standard carried out following the manufacturer's instructions. To

calculate a molar concentration, an average molar mass of proteins of 66,357.12 Da was

assumed (molar mass of human serum albumin).

For HSA, IgG, the γ-globulin fraction, and α_2-macroglobulin, the concentration was

determined by measuring the absorbance at 280 nm (A_{280nm}) using a JASCO V-650 UV/VIS

spectrophotometer. The molar extinction coefficient used for HSA was ε_{280}= 35,700 M^{-1} cm^{-1}

[50]. Concentration of the γ-globulin fraction was estimated using the extinction coefficient for

immunoglobulin G (IgG) of 1.36 cm^{-1} (mg · mL^{-1})$^{-1}$ [51]. Assuming a molecular weight of

150,000 Da [52], the molar extinction coefficient at 280 nm used for IgG was 210,000 M^{-1} cm^{-1}.

The molar extinction coefficient used for α_2-macroglobulin, ε_{280}= 145,440 M^{-1} cm^{-1}, was

calculated from amino acid sequence using ProtParam [53].

Concentration of Cohn fraction IV was determined using Pierce^{TM} Bicinchoninic Acid (BCA)

Protein Assay Kit and bovine serum albumin (BSA) as standard according to the

manufacturer's instructions. To calculate a molar concentration, a weighted average protein

mass of 80,000 Da was assumed based on the composition of the Cohn fraction IV.

Methylation of proteins

Proteins were dissolved in 1ml 1xPBS pH 7.4 to a concentration of 10 mg · mL^{-1} and the

solution cooled to 4 °C. 20 µl of 60 mg · mL^{-1} dimethylamine borane complex and 40 µl 1M

formaldehyde were then added. After 2 hours of incubation at 4 °C, this step was repeated. 2

hours later a final aliquot of 10 µl dimethylamine borane complex solution was added, before

incubation of the reaction mixture at 4 °C overnight. The next morning 125 µl of 1M Tris pH

7.5 were added to stop the reaction. The reacting agents were then separated from the now

methylated proteins by size-exclusion chromatography using Nap™-5 columns according to

the manufacturer's instructions (GE Healthcare Life Sciences, Amersham, UK).

HOCl-treatment of proteins and reduction

The concentration of the NaOCl stock solution of 0.64 M (Sigma-Aldrich, St. Louis, USA)

was confirmed regularly by measuring the absorbance at 292 nm using a JASCO V-650

UV/VIS spectrophotometer and the extinction coefficient ε_{292}= 350 M^{-1} cm^{-1}. When

necessary, NaOCl stock solution was diluted by mixing an adequate volume of NaOCl with

1x PBS solution pH 7.4 immediately prior to each chlorination reaction.

Varying amounts of a protein were then treated with a 10-fold, 50-fold and/or 150-fold molar

excess of NaOCl for 10 minutes at 30 °C (maximum NaOCl concentration used was 50 mM –

a concentration that can be produced by neutrophils per hour in chronically inflamed tissues

[31]). Excess HOCl was removed by size-exclusion chromatography using Nap™-5 columns

according to the manufacturer's instructions. Due to dilution during the Nap™-5 desalting

step, protein concentrations were re-determined as described above.

To reverse protein *N*-chlorination in HOCl-treated proteins, sodium ascorbate was dissolved

in 1xPBS pH 7.4 to a concentration of 1 M and the proteins were incubated with a 50-fold

molar excess of sodium ascorbate for 45 minutes at 37 °C. After removal of excess reductant

(see above), protein concentrations were again re-determined.

Protein aggregation assays with citrate synthase

200 μL Citrate synthase in ammonium sulfate solution (Sigma-Aldrich, St. Louis, USA) was

dialyzed overnight against 2 L 20mM Tris 2mM EDTA buffer at 4°C under constant stirring

using a Spectra/Por dialysis membrane with a MWCO of 6-8,000 Da. (Spectrum Laboratories

Inc., Rancho Dominguez, CA). This dialyzed citrate synthase preparation was then

chemically denatured in 4.5 M GdnHCl, 40 mM HEPES, pH 7.5 at room temperature

overnight. The final concentration of denatured citrate synthase was 12 μM. Aggregation was

induced by the addition of 20 µL denatured citrate synthase stock to 1580 µL 40 mM

HEPES/KOH buffer, pH 7.5. Final concentration of citrate synthase in the aggregation assay

was thus 0.15 µM. Untreated or treated plasma proteins were added to the assay buffer to a

final concentration of 0.5 µM (α_2-Macroglobulin) or 1.5 µM (HSA, IgG, γ-globulins, Cohn

fraction IV) (corresponding to a 3.3-fold and 10-fold molar excess over the dimeric citrate

synthase, respectively) prior to the addition of citrate synthase. The increase of light scattering

was monitored in a JASCO FP-8500 fluorescence spectrometer equipped with an EHC-813

temperature-controlled sample holder (JASCO, Tokyo, Japan) at 30 °C for 200-240 s under

continuous stirring. Measurement parameters were set to 360 nm (Ex/Em), 2.5 nm slit width

(Ex/Em) and medium sensitivity. Chaperone activity was expressed as the difference between

initial and final light scattering of an individual sample in arbitrary units. Aggregation of

citrate synthase in the absence of any other proteins was set to 100%. Depending on the batch

of citrate synthase, absolute maximum and minimum light scattering values may vary, thus

control experiments with the same batch were carried out for each individual aggregation

experiment.

Detection of accessible amino groups in proteins

Accessible amino groups in HSA were detected using fluorescamine (Sigma-Aldrich, St.

Louis, USA) as described [54]. Briefly, 334 µl of 3 mg · mL^{-1} fluorescamine stock in acetone

were added to 1 mL of 80 µg · mL^{-1} native HSA or the variously treated HSA solutions

described above. Emission spectrum of fluorescamine from 400 to 600 nm was measured

upon excitation with 388 nm using a JASCO FP-8500 fluorescence spectrometer. The relative

amount of accessible amino groups in the variously treated HSA samples was calculated by

setting the maximum fluorescence of native HSA to 100% representing the total relative

amino group content.

Nile red hydrophobicity assay

Nile red (Sigma-Aldrich, St. Louis, USA) was dissolved in dimethyl sulfoxide (DMSO) to a

final concentration of 30 μM. Varying amounts of native or HOCl-treated HSA (0-200 μM)

in 1xPBS were mixed with Nile red stock to a final dye concentration of 1.6 μM.

Fluorescence was measured using a JASCO FP-8500 fluorescence spectrometer with the

following parameters: 550 nm excitation, 570-700 nm emission, 5 nm slit width (Ex/Em) and

medium sensitivity. Concentrations of native or HOCl-treated HSA, at which the proteins

were half-saturated with dye were calculated by plotting the fluorescence intensity of Nile red

against the logarithm of the molar protein concentrations. Data were fitted with GraphPad

Prism 8 software using the sigmoid fit function.

PLB-985 culture and differentiation

The human myeloid leukemia cell line PLB-985 (certified mycoplasma negative, obtained

from DSMZ, German collection of microorganisms and cell culture) was cultured in RPMI-

1640 medium supplemented with 10% heat-inactivated FBS and 1% GlutaMAX (Life

Technologies, Carlsbad, CA) at 37°C with 5% CO_2. Cells were passaged twice weekly to

maintain a cell density between 2×10^5 and $1 \times 10^6 \cdot mL^{-1}$ and used until passage no. 10. For

differentiation into neutrophil-like cells, cells were seeded at a density of $2 \times 10^5 \cdot mL^{-1}$ and

cultured for 96 hours in RPMI-1640 medium with 10% FBS, 1% GlutaMAX and 1.25%

DMSO [55]. After 72 hours of incubation in the presence of DMSO, 2000 U $\cdot mL^{-1}$ interferon-γ

(ImmunoTools, Friesoythe, Germany) was added to the cell culture [56]. In a previous study,

differentiation was checked by detecting the expression of the associated surface markers

CD11b and CD64 (see ref. [7]). The viability of the cells was evaluated using trypan blue dye

and was typically >90%.

Chemiluminescence-based NADPH-oxidase activity assay

NADPH-oxidase-dependent   superoxide   production   was   selectively   measured   by

chemiluminescence (CL) using the chemiluminogenic substrate lucigenin (10,10'-Dimethyl-

9,9'-biacridinium dinitrate; Carl Roth, Karlsruhe, Germany) [57].

Differentiated PLB-985 cells were washed once with 1xPBS pH 7.4 and diluted in the same

buffer to a final concentration of 5×10^6 cells $\cdot$ mL^{-1}. 100 µL of this cell suspension were

placed in the wells of a non-transparent, white, clear-bottom 96-well plate (Nunc, Rochester,

NY). For some experiments, cells were preincubated with 10 µM diphenyleneiodium (DPI;

NADPH oxidase inhibitor), 100 nM wortmannin (phosphoinositide 3-kinase (PI3K)

inhibitor), 200 nM Gö 6983 (protein kinase C (PKC) inhibitor) or vehicle (1% DMSO) for 30

minutes at 37 °C. All inhibitors dissolved in DMSO were purchased from Sigma-Aldrich, St.

Louis, USA. The final concentration of DMSO in all wells was adjusted to 1%. The bottom of

the plate was covered using a white, non-transparent adhesive seal prior to measurement. 50

µL of either 1xPBS (resting CL) or agents to be tested, including the untreated and treated

plasma proteins were added to the respective wells. Final concentrations were: 2 mg $\cdot$ mL^{-1}

α_2-macroglobulin, 3 mg $\cdot$ mL^{-1} HSA, IgG, γ-globulins and Cohn fraction IV, 0.2 µM PMA

(Phorbol 12-myristate 13-acetate; Sigma-Aldrich, St. Louis, USA). Lucigenin was dissolved

in 1xPBS to a concentration of 400 µM immediately prior to measurement and 50 µL were

added to the wells. Chemiluminescence was measured every 1-2 minutes over 1.5 hours at 37

°C using the Synergy H1 multi-detection microplate reader (Biotek, Bad Friedrichshall,

Germany) in triplicates and chemiluminescence activity was expressed as integrated total

counts. In inhibition assays, NADPH-oxidase activation, as measured by chemiluminescence

activity, induced by PMA, HSA, treated with a 50-fold molar excess of HOCl, or IgG, treated

with a 150-fold molar excess of HOCl, was set to 100% and other relevant data as percentage

of this control.

Cloning of Ag85B

E. coli strains, plasmids and primers used in this study are listed in Table 1. Ag85B gene

(*fbpB*) of *Mycobacterium bovis* (differs by one base-pair from corresponding gene of *M.*

tuberculosis, resulting in a Leu to Phe replacement at position 100 [58]) was designed with

optimized codon usage for expression in *E. coli*, synthesized and cloned into pEX-A2

standard vector by Eurofins Genomics. *fbpB* was amplified from pEXA_fbpB by PCR using

primers fbpB-fw and fbpB-rv, purified according to the instructions of the NucleoSpin Gel

and PCR Clean-up Kit (Macherey-Nagel GmbH, Düren, Germany) and cloned into the

pET22b(+) expression vector via the restriction sites NdeI and XhoI with a hexahistidine

(His$_6$)-tag placed at the C-terminal end of the gene. *E. coli* DH5α cells were transformed with

the plasmid using a standard heat-shock method and plated on Luria Bertani (LB) agar plates

supplemented with 50 µg · mL^{-1} ampicillin. Screening for recombinant plasmids was

performed by colony PCR, followed by isolation of potentially correct plasmids from the

respective strains. Successful cloning of *fbpB* gene into pET22b(+) vector was verified by

sequencing.

Table 1: *E. coli* strains, plasmids and primers used in this study.

	Relevant properties [a] or genotype	Source or reference
***E. coli* strains**		
DH5α	*supE44, ΔlacU169 (φ80 lacZΔM15) hsdR17 recA1 endA1 hsdR gyrA relA thi*	Invitrogen
BL21(DE3)	*F⁻ ompT gal dcm lon hsdS$_B$(r$_B$⁻ m$_B$⁻) λ(DE3 [lacI lacUV5- T7 gene 1 ind1 sam7 nin5])*	Stratagene, Santa Clara, CA
Plasmids		
pEXA_fbpB	Amp^R pEX-A128 vector carrying synthesized *fbpB* gene from *M. bovis*	Eurofins Genomics
pET22b (+)	Amp^R, vector for overexpression of genes in *E. coli*	Novagen
pET22b-fbpB	Amp^R, vector for overexpression of *fbpB* gene in *E. coli* BL21(DE3)	This study
Primers	**Sequence (5' -> 3')**	
fbpB-fw	CCCCATATGTTCTCTCGTCCGG	
fbpB-rv	CCCCTCGAGACCAGCACCCAG	

281 *ᵃ* Amp^R, ampicillin resistance

Expression and purification of Ag85B

For heterologous expression and subsequent purification of His$_6$-tagged Ag85B, recombinant

plasmid pET22b-fbpB was transferred into the *E. coli* expression strain BL21 (DE3) (see

Table 1). The transformed cells were plated on LB agar plates containing 50 μg · mL^{-1}

ampicillin and incubated at 37 °C for 24 hours. 2 x 50 ml LB medium with ampicillin were

inoculated with a single colony from the agar plate and incubated at 37 °C overnight. These

overnight cultures were then used for inoculation of 5 x 1 L ampicillin-containing LB

medium to a starting optical density at 600 nm (OD$_{600}$) of 0.1. The bacteria were grown at 37

°C with shaking at 130 rpm until the OD$_{600}$ was ~ 0.5. Expression of Ag85B including a C-

terminal   hexahistidine   (His$_6$)-tag   was   induced   with   1   mM   isopropyl-beta-D-

thiogalactopyranoside (IPTG) and continued for ~ 12 hours (overnight) at 20 °C. Cells were

harvested by centrifugation at 6500 x g for 45 minutes at 4 °C. Pellets were washed once with

1xPBS and resuspended in lysis buffer (5 mM imidazole, 300 mM NaCl, 50 mM NaH$_2$PO$_4$,

pH 8.0). Cells were disrupted using a Constant Cell disrupter (Constant Systems Limited,

Daventry, England), and the obtained lysate was centrifuged for 45 minutes at 4 °C and

57,500 x g. Solid precipitate and the supernatant were separated and evaluated for detection of

recombinant protein by SDS-PAGE, followed by Coomassie Blue G-250 staining.

Overexpressed Ag85B protein was found predominantly in the pellet fraction. Pellet was thus

resuspended in lysis buffer containing 8 M urea and thoroughly homogenized. Suspension

was applied to a polystyrene column filled with nickel-nitrilotriacetic (Ni-NTA) resin, and

washed with urea-containing lysis buffer, followed by a washing step with wash buffer

supplemented with a small amount of imidazole (8M urea, 20 mM imidazole, 300 mM NaCl,

50 mM NaH_2PO_4, pH 8.0). Recombinant Ag85B was eluted from the column using 250 mM

imidazole buffer solution (8M urea, 250 mM imidazole, 300 mM NaCl, 50 mM NaH_2PO_4, pH

8.0). In order to remove imidazole, combined elution fraction was diluted 1:10 in sodium

phosphate buffer (300 mM NaCl, 50 mM NaH_2PO_4, pH 8.0) and mixed overnight at 4 °C.

Precipitated protein was collected by centrifugation for 45 minutes at 4 °C and 57,500 x g and

resuspended in urea-containing sodium phosphate buffer (8M urea, 300 mM NaCl, 50 mM

NaH_2PO_4, pH 8.0). Concentration of purified Ag85B protein was quantified

spectrophotometrically by measuring the absorbance at 280 nm using a JASCO V-650

UV/VIS spectrophotometer. The molar extinction coefficient used for Ag85B was $\varepsilon_{280}=$

75,860 $M^{-1} cm^{-1}$ [59].

Flow cytometry-based cell apoptosis assay

Differentiated PLB-985 cells were counted, washed once with 1xPBS pH 7.4 and diluted in

RPMI-1640 medium supplemented with 10% heat-inactivated FBS to a final concentration of

3.3×10^6 cells $\cdot$ mL^{-1}. 600 µL of this cell suspension were placed in the wells of a transparent,

flat-bottom 24-well plate (Sarstedt, Nümbrecht, Germany). The cells were preincubated with

100 nM wortmannin, 50 µM Z-VAD-FMK, a broad spectrum pan-caspase inhibitor

(dissolved in DMSO; Santa Cruz Biotechnology, Dallas, TX, USA), 250 µM cytochalasin D,

an inhibitor of actin polymerization (dissolved in DMSO; Sigma-Aldrich, St. Louis, USA),

155 μM native or modified HSA, that has been treated with a 50-fold molar excess of HOCl

as described above, or vehicle (1% DMSO) for 1 hour at 37 °C. The final concentration of

DMSO in all wells was adjusted to 1%. Subsequently, 1 μM Ag85B (85 μM stock in urea-

containing buffer (8 M urea, 300 mM NaCl, 50 mM NaH_2PO_4, pH 8.0)), 2 μM staurosporine

(dissolved in DMSO; Sigma-Aldrich, St. Louis, USA) or vehicle (94 mM urea and 1%

DMSO) were added. After 1 hour of incubation with Ag85B or 6 hours with staurosporine at

37°C, cells were washed once with cold 1xPBS, followed by one washing step with 1x

Annexin V binding buffer. For the analysis of cell viability, the variously treated cell

suspensions were stained with Annexin V-FITC and propidium iodide (PI) using the Dead

Cell Apoptosis Kit (Invitrogen, Thermo Fisher Scientific, Waltham, Massachusetts, USA)

according to the manufacturer's instructions and subsequently subjected to flow cytometry.

Samples were analyzed using a BD FACSCanto II flow cytometer (Becton, Dickinson and

Company). Fluorescence emitted by Annexin V-FITC and PI was measured through a

530/30-nm and 585/42-nm bandpass filter, respectively, upon excitation with an argon ion

laser operating at 488 nm. Single-stained compensation controls were used to calculate the

compensation matrix. 20,000 events were acquired and recorded per sample. Data were

analyzed using FlowJo (version 10) software.

Fluorescent labeling of Ag85B

For fluorescent labeling of Ag85B the green fluorescent dye CF^{TM} 488A succinimidyl ester

(Sigma-Aldrich, St. Louis, USA) was used. Labeling occurred via reaction of the

succinimidyl ester group of the dye with amine groups of Ag85B. Labeling was performed

according to the manufacturer's instructions with slight modifications. To prevent aggregation

of Ag85B, labeling reaction was performed in urea-containing buffer (8 M urea, 300 mM

NaCl, 50 mM NaH_2PO_4, pH 8.0) and carried out for 6 hours at room temperature under

continuous shaking. Excess dye was removed by size-exclusion chromatography using PD-10

desalting columns containing Sephadex G-25 resin according to the manufacturer's

instructions (GE Healthcare Life Sciences, Amersham, UK). Concentration of the conjugate

and degree of labeling (DOL) were calculated using the formula provided by the dye's

manufacturer. DOL was 3.1.

Ag85B-488 uptake assay

For the antigen uptake assay, differentiated PLB-985 cells were incubated with fluorescently-

labeled Ag85B (Ag85B-488) protein in the presence or absence of native and HOCl-modified

HSA in the same way as for the cell apoptosis assay described above. After 1 hour of

incubation at 37°C, cells were washed twice with 1xPBS, followed by fixation with 4%

paraformaldehyde (PFA) for 10 minutes on ice. Cells were washed with 1xPBS and

subsequently subjected to flow cytometry. Samples were analyzed using a BD FACSCanto II

flow cytometer. Fluorescence emitted by Ag85B-488 was measured through a 530/30-nm

bandpass filter upon excitation with the 488 nm argon laser line. 30,000 events were acquired

and recorded per sample. Data were analyzed using FlowJo (version 10) software.

Ag85B aggregation assay

Aggregation of Ag85B was induced by the stepwise addition of 75 µL Ag85B stock (5 x 15

µL every 20 seconds) to 1525 µl 40 mM HEPES/KOH buffer, pH 7.5. Final concentration of

Ag85B in the aggregation assay was 0.186 µM. Native or variously treated HSA as described

above, was added to the assay buffer to a final concentration of 14.88 µM (corresponding to a

80-fold molar excess over Ag85B) prior to the addition of Ag85B. The increase of light

scattering was monitored in a JASCO FP-8500 fluorescence spectrometer equipped with an

EHC-813 temperature-controlled sample holder at 30 °C for 200 s under continuous stirring.

Measurement parameters were set to 360 nm (Ex/Em), 2.5 nm slit width (Ex/Em) and medium

sensitivity. Chaperone activity was expressed as the difference between initial and final light

scattering of an individual sample in arbitrary units. Aggregation of Ag85B in the absence of

HSA was set to 100%.

RESULTS

HOCl-treated human serum prevents protein aggregation

In previous experiments we showed that the bacterial protein RidA is transformed into a competent holdase-type chaperone upon treatment with HOCl or *N*-chloramine [17]. In control experiments, we discovered that bovine serum albumin also shows increased chaperone activity in response to HOCl [17]. Since HOCl can be present in a physio-pathological context, most notably in the vicinity of inflammation, we wanted to test if human serum proteins can also be transformed into holdase-type chaperones through HOCl-treatment. Thus, we performed aggregation assays using chemically denatured citrate synthase and untreated and HOCl-treated human serum. Human serum was incubated with an estimated 10-fold molar excess of HOCl for 10 minutes at 30°C, a concentration that was sufficient to fully activate or markedly improve the chaperone function of RidA and BSA, respectively [17]. When chemically denatured citrate synthase was diluted into denaturant-free buffer, it formed aggregates that can be monitored by increased light scattering of the solution (Fig. 1). This aggregation of citrate synthase was not prevented by the addition of untreated human serum. However, when pre-incubated with a 10-fold molar excess of HOCl, serum significantly decreased aggregate formation. This suggested to us, that at least some serum proteins could be transformed to a holdase-type chaperone by *N*-chlorination in a mechanism similar to RidA. Because *N*-chlorination can be reduced by certain antioxidants, we used ascorbate to re-reduce HOCl-treated serum. Ascorbate is a mild antioxidant, which typically does not reduce native or HOCl-induced disulfide bonds. This ascorbate-treated serum lost its capability to bind denatured citrate synthase, suggesting an *N*-chlorination-based mechanism.

Figure 1: HOCl-treated human serum decreases protein aggregation. Human serum, when treated with a 10-fold molar excess of HOCl (Human serum $_{10xHOCl}$), significantly decreases aggregation of chemically-denatured citrate synthase as measured by light scattering at 360 nm. Reduction of HOCl-treated human serum with a 50-fold molar excess of the antioxidant ascorbate (Human serum $_{10xHOCl/Asc}$) reverses this chaperone-like conversion of the serum. **(a)** A representative measurement of citrate synthase aggregation in the presence of untreated (Human serum $_{UT}$), HOCl-treated (Human serum $_{10xHOCl}$) and re-reduced (Human serum $_{10xHOCl/Asc}$) human serum is shown. **(b)** Data are represented as means and standard deviations from three independent aggregation assays. Student's t-test: ***$p < 0.001$. Aggregation of citrate synthase in the absence of human serum was set to 100% and all the data are presented as percentage of this control. Labels of aggregation curves are written in the order of the final intensity of light scattering of the respective treatment.

Albumin, the major protein component of serum, shows HOCl-induced chaperone-like activity

The major protein component of human serum is human serum albumin (HSA). Previously, we showed that its bovine homologue bovine serum albumin (BSA) exhibits increased chaperone activity upon treatment with HOCl [17]. We therefore suspected that HSA could be a

major contributor to the observed decrease in protein aggregate formation in the presence of

HOCl-treated serum. To test whether HSA gains chaperone function upon exposure to HOCl,

HSA at a final concentration of 1 mM was treated with a 10-fold molar excess of HOCl for 10

minutes at 30 °C. An HOCl concentration of at least 10 mM has been shown to be required

for the generation of significant quantities of the so-called advanced oxidation protein

products (AOPPs) in plasma, associated with a number of inflammatory diseases [41]. The

localized concentration of HOCl generated by accumulated neutrophils in the interstitium of

chronically inflamed tissues, however, can be much higher and reach values of 25-50 mM per

hour [31]. To mimic a state of chronic inflammation, HSA was thus also incubated in the

presence of a 50-fold molar excess of HOCl corresponding to the maximum reported

concentration of 50 mM. HSA treated in both ways significantly reduced the aggregation of

citrate synthase when added at a 10-fold molar excess, showing that HSA acts as a potent

chaperone in serum upon modification by HOCl (Fig. 2 a, b). The chaperone activity of HSA,

however, was much higher upon exposure to a 50-fold molar excess of HOCl, suggesting a

dose-dependent activation of the HSA chaperone function by HOCl. In contrast to bacterial

RidA and Hsp33 [17,19], full activation of the chaperone function of HSA seems to require a

higher oxidation/chlorination level of the protein.

Chaperone-like conversion of serum albumin can be reversed by antioxidants

It is well known that exposure to high HOCl concentrations, such as those present at sites of

chronic inflammation, can lead to plasma protein unfolding and the formation of carbonylated

and di-tyrosine cross-linked protein aggregates that cannot return to the free, functional pool

of proteins upon reduction by antioxidants [16]. Such irreversibly misfolded plasma proteins

could principally act as chaperones and bind to and prevent the aggregation of other unfolding

substrates through hydrophobic interactions with the newly exposed hydrophobic protein

surfaces.

However, based on the observation that the chaperone activity of serum albumin increased in

a dose-dependent manner with the quantity of HOCl added, we asked whether the activation

of HSA chaperone function upon treatment with a 10- or 50-fold molar excess of HOCl could

be mediated by reversible *N*-chlorination instead. We have found this mechanism of action in

the bacterial protein RidA [17]. We thus exposed HOCl-treated HSA to a 50-fold molar excess

of the antioxidant ascorbate, which specifically removes *N*-chlorination. Indeed, ascorbate

rendered $HSA_{10xHOCl}$ and $HSA_{50xHOCl}$ unable to prevent citrate synthase aggregation when

added at the same molar excess as the HOCl-treated protein samples (Fig. 2 c, d). This result

strongly supports the idea that HOCl-mediated activation of HSA chaperone function

involves reversible chlorination of its side chain amines.

Methylation of basic amino acid residues in serum albumin inhibits HOCl-induced

activation of its chaperone function

Prompted by the observation that HOCl-mediated conversion of serum albumin into a potent

chaperone could be reversed by ascorbate, we assumed an *N*-chlorination-based activation

mechanism of its chaperone function. To further confirm this hypothesis, we blocked free

amino groups of lysine and nitrogens in the guanidino-moiety of arginine residues in HSA via

selective methylation. Exposure of methylated HSA to 10- or 50-fold molar excess of HOCl

did not convert HSA into a chaperone, suggesting that activation of the HSA chaperone

function indeed requires chlorination of its basic amino acids (Fig. 2 e, f).

Figure 2: Conversion of serum albumin into a potent chaperone upon HOCl exposure is based on reversible *N*-chlorination of its basic amino acids. (a, b) Serum albumin, when treated with a 10- or 50-fold molar excess of HOCl (HSA$_{10xHOCl}$ and HSA$_{50xHOCl}$, respectively), significantly decreases aggregation of chemically-denatured citrate synthase as measured by light scattering at 360 nm. **(c, d)** Reduction of HOCl-treated HSA with a 50-fold molar excess of the antioxidant ascorbate (HSA$_{10xHOCl/Asc}$ and HSA$_{50xHOCl/Asc}$) switches off its chaperone activity. **(d, e)** Methylation of basic amino acid side chains prior to HOCl treatment (HSA$_{Met/10xHOCl}$ and HSA$_{Met/50xHOCl}$) abrogates the chaperone-like conversion of HSA. In **a**, **c** and **e** representative measurements are shown. In **b**, **d** and **f** data are represented as means and standard deviations from three independent experiments. Student's t-test: **p < 0.01, ***p < 0.001. Aggregation of citrate synthase in the absence of HSA was set to 100% and all the data are presented as percentage of this control. Labels of aggregation curves are written in the order of the final intensity of light scattering of the respective treatment.

468 **Decreased amino group content of HSA upon HOCl treatment is accompanied by an**

469 **increased overall hydrophobicity of the protein**

470 Our combined data strongly suggested that the chaperone function of HSA is activated by

471 chlorination of its basic amino acids. HSA in its secreted form possesses 84 potential targets

472 for chlorination by HOCl: 59 lysine residues, 24 arginine residues and one amino group at the

473 N-terminus [60]. To investigate the extent to which HOCl decreases the total amount of

474 accessible, non-modified amino groups in HSA, we analyzed the free amino group content

475 before and after HOCl-treatment using fluorescamine [54]. Evidently, exposure to HOCl

476 resulted in some loss of free amino groups. Amino group content of HSA decreased by

477 approximately 10% after treatment with a 10-fold molar excess of HOCl (HSA$_{10xHOCl}$) and by

478 40% upon exposure to a 50-fold molar excess of HOCl (HSA$_{50xHOCl}$) (Fig. 3 a, b). Reduction

479 of both chlorinated HSA samples with ascorbate resulted in a full recovery of accessible

480 amino groups with concomitant loss of chaperone activity (Fig. 2 c, d). Activation of HSA

481 chaperone function by HOCl thus coincides with a decrease in free amino group content,

482 providing further evidence for an *N*-chlorination based-mechanism.

483 We argued that the reduction of positive charges on the protein's surface through HOCl-

484 induced *N*-chlorination should lead to an increase in surface hydrophobicity, thus allowing

485 high-affinity binding to unfolded proteins.

486 To detect changes in HSA's surface hydrophobicity upon HOCl treatment, we used the

487 uncharged hydrophobic dye Nile red and measured its fluorescence upon addition to 25 μM

488 native HSA or 25 μM HSA$_{50xHOCl}$ [61]. Absolute fluorescence of 1.6 μM Nile red was higher for

489 HSA$_{50xHOCl}$ when compared with untreated HSA with a maximum at 616.5 nm (621.5 nm for

490 native HSA) showing a blue shift in emission maximum consistent with a decreased polarity

491 of the protein solution (Fig. 3 c). The concentration of native HSA and HSA$_{50xHOCl}$, at which

492 the proteins have been half-saturated with dye were calculated (Fig. 3 d). This concentration

was significantly higher for untreated HSA compared to HSA$_{50xHOCl}$, pointing towards an

increased hydrophobicity of HSA$_{50xHOCl}$.

Figure 3: *N*-chlorination of serum albumin decreases accessible amino group content and increases

surface hydrophobicity. (**a**, **b**) Amino group content of variously treated HSA was analyzed using

fluorescamine. Treatment of HSA with HOCl resulted in a dose-dependent loss of free amino groups. Reduction

of chlorinated HSA with ascorbate fully restored the free amino group content. (**c**) Fluorescence of 1.6 μM Nile

red in the presence of native HSA or HSA, that has been treated with a 50-fold molar excess of HOCl. An

increased absolute fluorescence and a shift in maximum emission wavelength (from 621.5 nm to 616.5 nm) can

be observed for HSA$_{50xHOCl}$. (**d**) Absolute fluorescence of Nile red measured at the emission maximum at 621.5

nm and 616.5 nm, respectively, was plotted against the corresponding HSA$_{ut}$ and HSA$_{50xHOCl}$ concentrations,

respectively. The average concentrations, at which HSA$_{ut}$ and HSA$_{50xHOCl}$ have been half-saturated with Nile red

are marked by arrows. Means and standard deviations in **b** and **d** are based on three independent experiments.

(**b**) Student's t-test: *p < 0.05, **p < 0.01, ***p < 0.001. (**d**) Student's t-test: $^+$p < 0.05, $^#$p < 0.01, *p < 0.001.

For **a** and **c** representative measurements are shown.

Activation of neutrophil-like cells by HOCl-treated serum albumin is based on

reversible *N*-chlorination

Several lines of evidence point toward a key role of HOCl-modified serum albumin in the

progression of chronic inflammation, a hallmark of various degenerative diseases [46,47]. Upon

exposure to high doses of HOCl, as those present in chronically inflamed tissues, modified

HSA was found to induce neutrophil NADPH oxidase activation reflected by an increased

generation of reactive oxygen species and accompanied by degranulation [47].

Pathophysiological concentrations of HOCl are able to induce different modifications on

plasma proteins including carbonylation, *N*-chlorination, cysteine and methionine oxidation or

inter- and intramolecular di-tyrosine cross-linking [8,9,11,16], with most of them being considered

irreversible. So far, it has not been elucidated which HOCl-induced modification is sufficient

to convert HSA into a potent activator of leukocytes. We thus wondered whether this

functional conversion of HSA might be specifically mediated by *N*-chlorination and thus

reflects a reversible process, similar to the activation of its chaperone function in response to

HOCl stress.

To test this, we analyzed the effect of HOCl-treated HSA before and after reduction by

antioxidants on the activity of the phagocytic NADPH oxidase. For this purpose, we chose the

human myeloid cell line PLB-985 that acquires a neutrophil-like phenotype upon

differentiation with DMSO and IFNγ [55,56].

Generation of oxidants by the NADPH oxidase was assessed by lucigenin, a well

characterized and frequently used chemiluminescence probe which predominantly reacts with

superoxide anion radicals to form a light-emitting species [62]. Differentiated PLB-985 cells,

when incubated in buffer in the absence of any activating agents, showed a low basal level of

superoxide generation resulting from IFNγ-mediated enhancement of the NADPH oxidase

activity during the differentiation period [63] (Fig. 4 a). In line with expectations, treatment of

the cells with phorbol-12-myristate-13-acetate (PMA), a known activator of neutrophil

NADPH oxidase, led to a drastic increase in superoxide production and was thus used as

positive control [64] (Fig. 4 a, c). Addition of untreated HSA significantly increased the

generation of superoxide by $> 20\%$ compared to the mock control. No further enhancement of

the NADPH oxidase activity was observed with HSA after exposure to a 10-fold molar excess

of HOCl, indicating that the chosen HOCl concentration might have been too low for the

conversion of HSA into a potent activator of neutrophil-like cells. In contrast, cells subjected

to HSA that has been treated with a 50-fold molar excess of HOCl exhibited a substantial

increase in superoxide production within the first ~ 25 minutes of incubation. In comparison

to PMA, NADPH oxidase activation by $HSA_{50xHOCl}$ proceeded at a lower rate and was less

sustained, returning to basal level 40 minutes after the addition of $HSA_{50xHOCl}$ to the cells.

During our experiments we observed that *N*-chlorinated HSA can directly react with the

fluorescent dye, 2', 7'-dichlorodihydrofluorescein diacetate ($H_2DCF-DA$), which is the most

widely used probe for measuring intracellular ROS production [65], eliciting a positive signal in

the absence of cells (Supplementary Fig. S1). To confirm that the observed chemiluminescent

signal emitted by lucigenin derives from $HSA_{50xHOCl}$-induced activation of the phagocytic

NADPH-oxidase rather than from a similar direct reaction of lucigenin with $HSA_{50xHOCl}$ -

derived chloramines, we incubated lucigenin with the various agents in the absence of cells

and no significant chemiluminescence was observed (Fig. 4 b).

Importantly, when reduced by ascorbate, $HSA_{50xHOCl}$ lost its stimulatory effect, showing that

the functional conversion of HSA to an efficient activator of neutrophil-like cells upon HOCl

exposure reflects a reversible process. Since ascorbate specifically reduces *N*-chloramines

without affecting other HOCl-induced modifications, these results strongly support an *N*-

chlorination-based mechanism. The latter finding was further corroborated by methylation of

the basic amino acids of HSA prior to HOCl treatment, which completely prevented HOCl-

treated HSA from activating the neutrophil oxidative burst.

N-chlorinated HSA, formed upon exposure to high doses of HOCl, can be thus considered as

an inflammatory response modulator that contributes to the activation of neutrophils at sites

of inflammation.

Figure 4: Activation of neutrophil-like cells by HOCl-treated serum albumin is mediated by reversible *N*-chlorination. Treatment with a 50-fold molar excess of HOCl (HSA $_{50xHOCl}$) converted HSA into an efficient inducer of the neutrophil respiratory burst, reflected by the increased production and release of oxidants that induce lucigenin chemiluminescence. This activating function of HSA $_{50xHOCl}$ could be reversed by reduction with the antioxidant ascorbate (HSA $_{50xHOCl/Asc}$) and was abrogated by methylation of its basic amino acid side chains prior to HOCl exposure (HSA $_{Met/50xHOCl}$). **(a)** Extracellular oxidant production by neutrophil NADPH oxidase was measured in one- to two-minutes intervals over 90 minutes at 37 °C using lucigenin-enhanced chemiluminescence. Phorbol 12-myristate 13-acetate (PMA; final concentration, 0.2 µM), untreated and variously treated HSA samples (final concentration, 3 mg · mL^{-1}) from the previous citrate synthase aggregation assays (see above) or PBS buffer (basal oxidant production) were added to **(a)** differentiated PLB-985 cells in PBS buffer or **(b)** cell-free PBS puffer containing 400 µM lucigenin immediately prior to chemiluminescence measurement. **(c)** Results shown in **a** are expressed as integrated total counts (means and standard deviations of

three independent measurements) higher than buffer control. Student's t-test: **$p < 0.01$, ***$p < 0.001$. PMA-

induced activation of NADPH-oxidase was set to 100%.

All plasma fractions tested exhibit chaperone activity upon HOCl treatment

Our finding that the major protein in human plasma, serum albumin. can be converted into a

potent chaperone upon HOCl-mediated *N*-chlorination, prompted us to ask whether also other

plasma protein fractions exhibit similar chaperone activity upon treatment with HOCl.

We thus tested the γ-globulin fraction, the Cohn fraction IV (comprising α- and β-globulins),

and specifically α_2-macroglobulin, an important protease inhibitor in human plasma, for

chaperone activity upon exposure to various doses of HOCl.

Aside from its known interaction with proteases, native α_2-macroglobulin (α_2M) acts as an

extracellular chaperone that binds to and prevents the accumulation of misfolded proteins,

particularly during the innate immune response. Exposure to HOCl was found to further

improve the chaperone function of α_2M, but the mechanism remained unclear [32].

To test whether the previously observed effect is due to *N*-chlorination, we treated α_2M with a

10- and 50-fold molar excess of HOCl (corresponding to 0.3 mM and 1.5 mM HOCl,

respectively), followed by the addition of the antioxidant ascorbate for re-reduction. In line

with expectations, chaperone activity of α_2M increased with the amount of HOCl added (Fig.

5 a, b). Treatment with ascorbate, as well as methylation of basic amino acid residues prior to

HOCl exposure fully inhibited the chaperone activation of HOCl-modified α_2M. These results

strongly support the notion that HOCl-induced *N*-chlorination of basic amino acid side chains

also accounts for the increased chaperone activity of α_2M.

Intriguingly, we also observed similar chaperone-like conversion upon HOCl treatment for

Cohn fraction IV and the γ-globulin fraction (Fig. 5 c-f). In both cases, however, efficient

activation of the chaperone activity required higher HOCl concentrations. It is important to

note, however, that all HOCl concentrations used were below the range known to be

generated directly at sites of inflammation [31]. While exposure of Cohn fraction IV to an

estimated 10-fold molar excess of HOCl (corresponding to 2 mM HOCl) had only little effect

on citrate synthase aggregation, treatment with a 50-fold molar excess of HOCl strongly

activated chaperone-like properties (Fig. 5 c, d). This chaperone activity was markedly

reduced, but not completely inhibited, after reduction with ascorbate. Likewise, methylation

of amine side chains prior to HOCl exposure did not fully abrogate the chaperone-like

conversion of this protein fraction.

Similarly, activation of the chaperone function of the γ-globulin fraction occurred at higher

HOCl concentrations ranging from 4.3 mM to 13 mM (corresponding to an estimated 50- to

150-fold molar excess of HOCl). Strikingly, while chaperone activity increased with the

amount of HOCl added, reversibility of the chaperone function by ascorbate decreased (Fig. 5

e, f).

These results suggest that at least some proteins in both plasma fractions tested are

transformed to chaperones upon modification by HOCl. Efficient activation of their

chaperone function is strongly, but, unlike HSA, not exclusively linked to *N*-chlorination,

suggesting some other HOCl-induced modifications that cannot be removed by ascorbate in

these plasma proteins.

Figure 5: All plasma protein fractions tested exhibit reversible chaperone activity upon modification by HOCl. α_2-Macroglobulin (**a**, **b**), Cohn fraction IV (**c**, **d**) and the γ-globulin fraction (**e**, **f**) were analyzed for chaperone activity in a citrate synthase aggregation assay upon treatment with various doses of HOCl. Each plasma protein fraction, when treated with a 10-, 50- or 150-fold molar excess of HOCl, significantly decreased aggregation of chemically-denatured citrate synthase as measured by light scattering at 360 nm. Exposure of the various HOCl-treated plasma proteins to the reductant ascorbate significantly decreased or completely inhibited their chaperone function. Methylation of basic amino acid residues prior to HOCl treatment mostly prevented chaperone-like conversion of the plasma proteins. In **a**, **c** and **e** representative measurements are shown. In **b**, **d** and **f** data are depicting means and standard deviations from three independent experiments. Student's t-test: *p < 0.05, **p < 0.01, ***p < 0.001. Aggregation of citrate synthase in the absence of any plasma protein fraction was set to 100% and all the data are presented as percentage of this control. Labels of aggregation curves are written in the order of the final intensity of light scattering of the respective treatment.

HOCl-induced *N*-chlorination converts the majority of plasma proteins into activators

of neutrophil-like cells

Ours and others findings indicate that *N*-chlorinated serum albumin is a key factor for the

stimulation of leukocytes at sites of inflammation. To test if also other plasma proteins exert

this function upon HOCl exposure, we analyzed the NADPH oxidase-dependent generation of

superoxide by differentiated PLB-985 cells in the presence of various treated plasma protein

fractions, as described above.

Addition of α_2-macroglobulin, treated with a 50-fold molar excess of HOCl (i.e. 1.5 mM

HOCl), did not enhance superoxide production (Fig. 6 a, b). In contrast, exposure to the same

molar excess of HOCl, corresponding to 10 mM HOCl in this case, converted at least some

proteins of Cohn fraction IV into highly efficient stimulators of the neutrophil NADPH

oxidase (Fig. 6 c, d). Analysis of the kinetics showed that activation by this HOCl-treated

protein fraction occurred at a similar rate as the activation by PMA. Such a stimulatory effect

was also observed for the γ-globulin fraction upon treatment with an estimated 150-fold molar

excess of HOCl (i.e. 13 mM). The minimum HOCl concentration required for the activating

function of these plasma proteins was thus 10 mM and conformed to the range of HOCl

concentrations expected under inflammatory conditions. Importantly, the activating effect of

both HOCl-treated plasma fractions was completely abolished upon reduction with ascorbate

or by prior methylation of their amino groups, strongly suggesting that *N*-chlorination is the

responsible mechanism for the functional switch of these proteins into efficient activators of

neutrophil-like cells as well.

These findings demonstrate that HOCl-mediated *N*-chlorination constitutes a key mechanism

to increase the immunogenicity of plasma proteins under inflammatory conditions. Upon

modification, not only serum albumin, but the majority of the plasma fractions tested, form a

feed-forward inflammatory loop to amplify and sustain inflammatory responses which can

lead to accelerated pathogen clearance but could also contribute to chronic inflammation.

Figure 6: The majority of human plasma proteins stimulate neutrophil respiratory burst upon *N*-chlorination by HOCl. The effect of HOCl-treated α_2-macroglobulin (α_2M) (**a**, **b**), Cohn fraction IV (**c**, **d**) and the γ-globulin fraction (**e**, **f**) on the activity of the neutrophil NADPH oxidase was investigated. α_2M, when treated with various doses of HOCl, had no influence on ROS generation by the NADPH oxidase (**a**, **b**). Treatment with a 50- or 150-fold molar excess of HOCl converted at least some proteins of Cohn fraction IV (CohnIV $_{50xHOCl}$) and the γ-globulin fraction (γGlobulins $_{150xHOCl}$), respectively, into efficient inducers of the neutrophil respiratory burst, reflected by the increased production and release of oxidants that induce lucigenin chemiluminescence (**c-f**). The activating function of CohnIV $_{50xHOCl}$ and γGlobulins $_{150xHOCl}$ could be reversed by treatment with the reductant ascorbate (CohnIV $_{50xHOCl/Asc}$ and γGlobulins $_{150xHOCl/Asc}$) and was abrogated by

methylation of basic amine side chains prior to HOCl exposure (CohnIV $_{Met/50xHOCl}$ and γGlobulins $_{Met/150xHOCl}$).

(a, **c**, **e)** Extracellular oxidant production by neutrophil NADPH oxidase was measured in one- to two-minutes

intervals over 90 minutes at 37 °C using lucigenin-enhanced chemiluminescence. Phorbol 12-myristate 13-

acetate (PMA; final concentration (fc), 0.2 μM), untreated and the variously treated plasma fraction samples (fc,

2 mg · mL^{-1} for α$_2$-macroglobulin and 3 mg · mL^{-1} for Cohn fraction IV and the γ-globulin fraction) from the

previous citrate synthase aggregation assays (see above) or PBS buffer (basal oxidant production) were added to

differentiated PLB-985 cells in PBS buffer containing 400 μM lucigenin immediately prior to

chemiluminescence measurement. **(b**, **d**, **f)** Results are expressed as integrated total counts (means and standard

deviations of three independent measurements) higher than buffer control. Student's t-test: *p < 0.05, **p <

0.01, ***p < 0.001. PMA- induced activation of NADPH-oxidase was set to 100%.

Activation of NADPH oxidase by *N*-chlorinated serum albumin and immunoglobulin G

occurs predominantly via PI3K-dependent signaling pathways

Activity of the NADPH oxidase complex of neutrophils is regulated by several signaling

pathways downstream of cell surface receptors. Central amongst these are PLC/PKC-

(phospholipase C/protein kinase C) and PI3K- (phosphoinositide 3-kinase)-dependent

pathways, blockade of which severely lowers NADPH oxidase activation by various stimuli

such as chemotactic peptides, opsonized particles or phorbol esters [66–69].

To identify the predominant signaling mechanism through which the *N*-chlorinated serum

albumin and the major component of the γ-globulin fraction, immunoglobulin G, activate the

neutrophil respiratory burst, the effect of various inhibitors on the HSA$_{50xHOCl}$ and

IgG$_{150xHOCl}$-induced ROS generation was tested. PMA, a direct activator of conventional PKC

isoforms such as PKCα and PKCβ was used as a control for PKC-dependent activation of the

NADPH oxidase [64].

It is worth noting, that all cell suspensions contained DMSO at a final concentration of 1%.

As a consequence, NADPH oxidase activation by PMA, HSA$_{50xHOCl}$ or IgG$_{150xHOCl}$ proceeded

at a lower rate compared to the previously described experiments (see Figs. 4 and 6), peaking

at 35-50 minutes after the addition of these agents to the cells.

Pretreatment of neutrophil-like cells with 10 μM diphenyleneiodonium (DPI), a direct

inhibitor of the NADPH oxidase complex, for 30 minutes prior to stimulation with PMA,

HSA$_{50xHOCl}$ or IgG$_{150xHOCl}$ fully inhibited ROS generation confirming that the ROS production

induced by these stimulatory agents was NADPH-oxidase dependent (Fig. 7). In line with

expectations, presence of the PI3K inhibitor wortmannin (100 nM) had only little effect on

the PMA-mediated activation of the NADPH oxidase (Fig. 7 c, d). In contrast, wortmannin

strongly attenuated the HSA$_{50xHOCl}$- and IgG$_{150xHOCl}$-induced ROS generation by the immune

cells, suggesting a PI3K-dependent mechanism of NADPH oxidase activation (Fig. 7 a, b, d).

Along this line, pretreatment with 200 nM Gö 6983, that selectively inhibits several PKC

isozyme families, including the classical PKCα and PKCβII, the novel PKCδ and the atypical

PKCζ, strongly inhibited NADPH oxidase activation by PMA, but not by HSA$_{50xHOCl}$ and

IgG$_{150xHOCl}$.

These results suggest that PI3K is the key signaling component in the pathway that leads to

HSA$_{50xHOCl}$ – and IgG$_{150xHOCl}$- dependent activation of the neutrophil NADPH oxidase.

Figure 7: Activation of the NADPH oxidase of neutrophil-like cells by HOCl-treated serum albumin and immunoglobulin G occurs predominantly via a PI3K-dependent signaling pathway. Effect of 10 μM diphenyleneiodonium (DPI; NADPH oxidase inhibitor), 100 nM wortmannin (PI3K inhibitor) and 200 nM Gö 6983 (protein kinase C (PKC) inhibitor) on the NADPH oxidase activation mediated by **(a)** 3 mg · mL^{-1} HSA$_{50xHOCl}$, **(b)** 3 mg · mL^{-1} IgG$_{150xHOCl}$ and **(c)** 0.2 μM PMA was tested. **(d)** Results shown in **a**, **b** and **c** are expressed as integrated total counts (means and standard deviations of three independent measurements) higher than the respective buffer control. Student's t-test: **p < 0.01, ***p < 0.001.

HOCl-treated serum albumin promotes survival of neutrophil-like cells in the presence of the foreign protein antigen Ag85B, but not in the presence of staurosporine

As shown previously, activation of the NADPH oxidase by HOCl-modified serum albumin involves the action of PI3K. Since PI3K is a key component of the well-documented, anti-apoptotic PI3K/Akt signaling pathway [70], it was tempting to speculate that HOCl-modified HSA could promote cell survival in the presence of noxious stimuli. To test this hypothesis, we exposed neutrophil-like PLB-985 cells to the highly immunogenic mycobacterial protein antigen Ag85B and staurosporine in the presence of native or HOCl-treated HSA. The

mycolyltransferase Ag85B is the major antigen produced and secreted from all mycobacterial

species during infection [71] and has been shown to play an important role in the induction of

protective immunity [72,73] by inducing strong T cell proliferation and IFN-γ secretion [74,75].

Activation of neutrophils during mycobacterial infections is often accompanied by accelerated

apoptosis [76,77], but the mechanism by which mycobacterial species or their secreted antigens

induce apoptosis has not been elucidated in detail. Staurosporine, a protein kinase inhibitor,

has been characterized as an efficient inducer of apoptosis in various cell types via caspase-

dependent and -independent pathways [78,79] and thus, was also used as an cell death-promoting

agent in our experiments.

After one hour or six hours of incubation with Ag85B and staurosporine, respectively,

viability of the variously treated cells was evaluated by flow cytometry using Annexin V-

FITC/propidium iodide (PI) staining. During early apoptosis, phosphatidylserine is

translocated from the inner to the outer cell membrane leaflet with the plasma membrane left

intact and thus, available for binding of extrinsically applied annexin V protein [80]. In late

apoptosis/necrosis, the integrity of the plasma membrane is lost, allowing the normally

membrane-impermeable propidium iodide to enter and stain the DNA. Cells that are in late

apoptosis or necrotic are thus both Annexin V-FITC and PI positive. Accordingly, cells that

are viable are both Annexin V-FITC and PI negative.

Data plots were generated from analysis of ungated data (Fig. 8). Viable cells appear in the

lower left quadrant (Q4), early apoptotic cells in lower right quadrant (Q3) and late

apoptotic/necrotic cells in the upper right (Q2) quadrant.

Upon treatment with Ag85B, the ratio of viable cells to early/late apoptotic cells has markedly

decreased compared to the control cells, that were incubated in the absence of Ag85B,

pointing towards a lifetime-limiting effect of Ag85B. Pretreatment with the broad-spectrum

pan-caspase inhibitor Z-VAD-FMK did not significantly affect Ag85B-induced cell death,

suggesting that Ag85B exerts its lethal effect through a caspase-independent mechanism. To

test whether the toxicity of Ag85B depends on its uptake by the immune cells, we pretreated

the cells with cytochalasin D, an inhibitor of phagocytosis, prior to the addition of Ag85B.

Reduction of the cells' phagocytic capacity resulted in markedly improved cell survival in the

presence of Ag85B. This phenomenon is illustrated by a shift of the cells from Q3 to Q4 on

the Annexin V-FITC/PI plot with concomitant reduction of necrotic cells. Remarkably,

pretreatment with HOCl-modified HSA, added at a 155-fold molar excess over Ag85B,

completely prevented cell death upon addition of Ag85B. In contrast, native HSA had almost

no effect on Ag85B-induced cell death. We thus speculated that HOCl-treated HSA could

rescue immune cells from Ag85B-induced cell death by preventing or strongly reducing its

uptake, rather than by boosting PI3K/Akt signaling. In support of this conclusion, Z-VAD-

FMK, but not HOCl-treated HSA was able to significantly reduce staurosporine-induced

apoptosis. Instead, when combined with staurosporine, HOCl-modified HSA triggered

necrosis with the majority of the cells being Annexin V and PI positive.

Figure 8: HOCl-treated serum albumin improves survival of neutrophil-like cells in the presence of the major mycobacterial protein antigen Ag85B. Differentiated neutrophil-like PLB-985 cells were preincubated with 50 μM Z-VAD-FMK, 155 μM native (HSA$_{UT}$) or HOCl-treated HSA (HSA$_{50xHOCl}$) prior to the addition of 1 μM Ag85B or 2 μM staurosporine. After one hour (Ag85B) or six hours (staurosporine) of incubation, viability of the variously treated cells was assessed by flow cytometry using Annexin V/propidium iodide (PI) staining. Cells treated only with buffer served as control. **(a)** Annexin V-FITC vs. propidium iodide dot plots show all analyzed events. Staining of cells simultaneously with Annexin V- FITC (green fluorescence) and the non-vital dye PI (red fluorescence) allows the discrimination of viable, intact cells (FITC-, PI-; Q4), early apoptotic (FITC+, PI-; Q3) and late apoptotic/necrotic cells (FITC+, PI+; Q2). 20,000 events were acquired and recorded per sample. Data were analyzed using FlowJo (version 10) software. Results shown are representative of three

experiments. **(b)** Results of three independent experiments for viable cells are shown (means and standard

deviations). Student's t-test: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

HOCl-treated serum albumin effectively binds Ag85B and reduces its uptake by

neutrophil-like cells

Prompted by the finding that HOCl-treated HSA can act as a chaperone being highly effective

at preventing protein aggregation, we asked whether *N*-chlorinated HSA can also bind to

Ag85B and thus prevent its uptake by immune cells. To test this, Ag85B was diluted stepwise

in the presence of native or HOCl-treated HSA and aggregation of Ag85B was monitored by

light scattering. When denatured Ag85B was diluted into buffer, it readily formed aggregates

(Fig. 9 a, b). This aggregation could not be prevented by untreated HSA. Presence of HOCl-

modified HSA, however, significantly reduced the aggregation of Ag85B when added at a 80-

fold molar excess over the protein. Again, reduction with ascorbate fully abrogated

HSA_{HOCl}'s ability to bind Ag85B.

We argued that Ag85B's association with HOCl-modified HSA could decrease its propensity

to enter immune cells and thus, provide a possible explanation for the enhanced survival of

neutrophils in the presence of this mycobacterial antigen. To test whether this HSA_{HOCl} -

mediated cell survival is indeed due to a reduced uptake, we added recombinant fluorescently-

labeled Ag85B (Ag85B-488) to differentiated neutrophil-like cells and analyzed its uptake in

the presence of native or HOCl-treated HSA using flow cytometry. Cells conincubated with

native HSA accumulated around 30% less Ag85B compared to cells treated with Ag85B

alone (Fig. 9 c, d). But presence of HOCl-treated HSA reduced uptake of Ag85B by up to

80%. Sustained cell viability in the presence of Ag85B is thus most likely linked to decreased

antigen uptake by the immune cells.

Figure 9: HOCl-treated serum albumin binds to and prevents uptake of the major mycobacterial protein antigen Ag85B by neutrophil-like cells. (a, b) HSA, treated with a 50-fold molar excess of HOCl (HSA$_{50xHOCl}$) significantly decreased aggregation of denatured Ag85B as measured by light scattering at 360 nm. Reduction of HSA$_{50xHOCl}$ with a 50-fold molar excess of the antioxidant ascorbate (HSA$_{50xHOCl/Asc}$) reversed this chaperone activity. **(a)** A representative measurement of Ag85B aggregation in the presence of native HSA (HSA$_{UT}$), HSA$_{50xHOCl}$ and HSA$_{50xHOCl/Asc}$ is shown. Labels of aggregation curves are written in the order of the final intensity of light scattering of the respective treatment. **(b)** Data are represented as means and standard deviations from three independent aggregation assays. Aggregation of Ag85B in the absence of HSA was set to 100% and all the data are presented as percentage of this control. **(c, d)** Differentiated neutrophil-like PLB-985 cells were incubated in the absence or presence of the fluorescently-labeled Ag85B protein, Ag85B-488. In some cases, HSA$_{UT}$ or HSA$_{50xHOCl}$ was added at a 50-fold molar excess over Ag85B-488 to the cells. After one hour of incubation, uptake of fluorescent Ag85B-488 by the variously treated cells was assessed by flow cytometry. 30,000 events were acquired and recorded per sample. Data were analyzed using FlowJo (version 10) software. **(c)** Single-parameter histogram overlays of Ag85B-488 fluorescence of the various samples are shown. **(d)** Results of three independent experiments are shown (means and standard deviations). Student's t-test: **p <

0.01, ***p < 0.001. Uptake of Ag85B-488, reflected by the median fluorescence intensity, in the absence of

HSA was set to 100%.

DISCUSSION

Myeloperoxidase (MPO)-derived HOCl is one of the strongest oxidants produced and

released by neutrophils during infection and inflammation (recently reviewed in ref. [42]).

Because of its high reactivity with a wide variety of biomolecules, HOCl forms an efficient

weapon against a wide range of pathogens, but also contributes to host tissue injury associated

with diverse inflammatory diseases [81]. Due to their high abundance in blood and interstitial

fluid, plasma proteins constitute major targets for HOCl-mediated modification with thiol

oxidation and chlorination of amine side chains being the favored reactions [22], followed by

some irreversible modifications including tryptophan oxidation, peptide bond cleavage or di-

tyrosine cross-linking [82], when exposed to high doses of HOCl as those present at sites of

chronic inflammation. Formation and accumulation of such advanced oxidation protein

products (AOPPs) is indeed a hallmark of various inflammatory diseases [44,83].

Aside from their role as important scavengers of the majority of generated HOCl [39,84], there is

a growing body of evidence demonstrating protective and/or immunomodulatory function of

these AOPPs during inflammation. α_2-Macroglobulin (α_2M), a plasma glycoprotein, already

acts as an extracellular chaperone capable of suppressing the aggregation of a range of

proteins [85], but modification by HOCl has been reported to drastically increase its chaperone

activity by a yet unknown mechanism [32]. Previously, we observed a similar phenomenon in *E.*

coli, where the protein RidA, a member of the YjgF/YER057c/UK114 protein family, was

converted into an efficient chaperone holdase upon exposure to HOCl by a mechanism that

involved the reversible chlorination of its basic amino acids [17]. This finding was rather

surprising, since activation of other known redox-regulated chaperones, such as Hsp33 or 2-

cys peroxiredoxins [19,86], by HOCl occured via oxidation of their cysteine residues and not by

N-chlorination, a modification that before our study has been thought to have a mostly

deleterious effect on protein function. Since similar observations were made more recently for

E. coli CnoX (YbbN) [25] and the HSA homologue bovine serum albumin (BSA) [17], HOCl-

mediated *N*-chlorination can, therefore, be considered a novel, cysteine-independent and

reversible mechanism of chaperone activation in response to HOCl-stress.

Encouraged by these findings, we hypothesized that HOCl-mediated improvement of α_2M's

chaperone function may be also based on *N*-chlorination and suspected that α_2M could be not

the only plasma protein exhibiting chaperone activity upon modification by HOCl. In fact, we

found that not only α_2M but a number of proteins in all plasma fractions tested turned into

chaperone-like proteins, being highly effective at preventing formation of potentially toxic

protein aggregates *in vitro*. The role of plasma proteins in inflamed tissue thus goes beyond

that of a passive sink for HOCl.

Since exposure to high HOCl concentrations can lead to oxidative, irreversible protein

unfolding [16], one might speculate that the observed chaperone activity of such unfolded

plasma proteins could simply be the result of an increased affinity to other unfolded proteins.

Here, we provide direct evidence that HOCl-mediated conversion of plasma proteins into

potent chaperones depends primarily on reversible chlorination of their basic amino acids.

Along this line, treatment of HSA with HOCl led to dose-dependent reduction of the

accessible amino group content accompanied by an increase in surface hydrophobicity,

providing an obvious explanation for the increased affinity to unfolding proteins of *N*-

chlorinated plasma proteins. Since *N*-chlorination does not irreversibly alter the structural and

functional properties of a protein, this reversible post-translational modification provides a

new strategy to recruit new chaperone-like proteins on demand in response to HOCl-stress in

order to minimize self-damage associated with the formation of protein aggregates during

inflammation.

In recent years, other roles for HOCl-modified plasma proteins and lipoproteins in

inflammatory processes have also been described. For example, HOCl-oxidized low-density

lipoproteins [87] as well as HOCl-modified HSA [47,48] have been shown to elicit various

polymorphonuclear leukocyte (PMNL) responses such as the NADPH-dependent generation

of reactive oxygen species (ROS), degranulation or shape change [47], but the mechanism for

the HOCl-mediated functional conversion into a potent activator of human PMNLs remained

unclear.

As *N*-chlorination was the principal chemical modification responsible for the plasma

protein's switch to a chaperone-like holdase, we hypothesized that it could also be the reason

for the activation of immune cells. To mimic an *in vivo* situation, where plasma proteins are

located in immediate vicinity to accumulated neutrophils in inflamed tissue, we exposed

several plasma fractions to HOCl at pathophysiological concentrations [31] and investigated the

effect of the resulting products on ROS production by neutrophil-like cells. Remarkably, we

found that not only the main plasma protein HSA, but the majority of plasma fractions tested,

when treated with a sufficient amount of HOCl, were able to elicit a significant immune

response, as shown by the increased generation of ROS by the immune cells. This effect was

not seen upon re-reduction of the HOCl-modified proteins or prior methylation of their basic

amino acids, strongly supporting a *N*-chlorination based mechanism as well. To the best of

our knowledge, this is the first study which demonstrates that reversible HOCl-mediated *N*-

chlorination is the principal mechanism of turning plasma proteins into critical modulators of

the innate immune response.

Gorudko et al. reported that the phosphoinositide 3-kinase (PI3K) inhibitor wortmannin

inhibited the stimulating effect of HOCl-modified HSA on immune cells [47]. The same was

true in our model system: the immunomodulatory action of both HOCl-modified HSA and

immunoglobulin G could be fully inhibited or strongly attenuated by wortmannin. PI3K and

its downstream effectors, such as the serine/threonine kinase Akt, are indeed considered to

play a key role in the regulation of the neutrophil NADPH oxidase [88,89]. Since the PKC

inhibitor Gö 6983 showed only little effect on $HSA_{50xHOCl}$- and $IgG_{150xHOCl}$-mediated NADPH

oxidase activation, we thus propose that these proteins stimulate the neutrophil respiratory

burst predominantly via PI3K-dependent signaling pathways. However, further studies are

needed to investigate the exact mechanism by which these proteins trigger PI3K signaling. A

possible scenario might be, that *N*-chlorination confers higher affinity to a membrane

scavenger receptor allowing the binding of HOCl-modified HSA. In support of this theory, it

has been reported that *N*-chlorinated, but not native HSA can irreversibly bind to and block

the major high-density lipoprotein receptor, scavenger receptor class B, type 1 (SR-BI) [90]. It is

therefore tempting to speculate that *N*-chlorination could also increase the affinity of IgG to

Fc gamma receptors (FcγRs).

The PI3K/Akt pathway is also considered an anti-apoptotic pathway [91,92]. Thus, it was

tempting to hypothesize that HOCl-modified HSA may play a role as a pro-survival molecule

during inflammation. Remarkably, we found that HOCl-modified HSA indeed enhances

survival of neutrophils in the presence of the highly immunogenic mycobacterial protein

antigen Ag85B, however, not in the presence of staurosporine, a broad spectrum protein

kinase inhibitor that induces apoptosis in various cell types [78,79]. Looking for a mechanism by

which HOCl-modified HSA is able to rescue immune cells from Ag85B-induced cell death,

we found that it can effectively bind to and strongly decrease the phagocytosis of Ag85B.

Internalization of Ag85B by the cells proved to be the direct cause of cell death and could

also be prevented by other inhibitors of phagocytosis.

Neutrophils are typically short-lived, but their apoptosis can be delayed both by microbial

products and by various proinflammatory stimuli [93,94]. In this study, we describe HOCl-

modified HSA as a novel pro-inflammatory mediator, which can promote cell survival by

binding to highly immunogenic foreign antigens and reducing their phagocytosis at sites of

bacterial infection. A similar phenomenon was observed in other studies, where HOCl-

modified HSA was shown to bind and neutralize proteins from HIV and West Nile virus [95,96].

The immunomodulatory effects of *N*-chlorinated plasma proteins found in this and previous

studies constitute a double-edged sword. Although stimulation of neutrophil respiratory burst

and enhanced neutrophil survival may be beneficial for pathogen elimination at the initial

stage of infection, it can eventually perpetuate a positive feedback loop and contribute to the

development and progression of chronic inflammation (Fig. 10). Secretion of HOCl and other

oxidants by permanently activated neutrophils leads not only to the destruction of

neighboring, healthy cells resulting in tissue injury [26,97,98], but generates more *N*-chlorinated

plasma proteins. Similarly, the pro-survival effect of *N*-chlorinated HSA is not only positive,

as cell death and the subsequent recognition of dying neutrophils by macrophages has a

critical function in the resolution of the inflammatory response and is strictly required to

protect the surrounding tissue and prevent pathological sequelae [99]. Were all these properties

of AOPPs dependent on the numerous irreversible modifications reportedly caused by HOCl

exposure, this could lead to a spreading out-of-control immune reaction. The presence of high

concentrations of antioxidants such as ascorbate and glutathione in plasma, both of which can

reduce *N*-chlorination, provide a mechanism to contain the immune reaction at the site of

inflammation. Indeed, depletion of these antioxidants is often associated with chronic

inflammation and other diseases [100–103] and antioxidant therapy such as high-dose intravenous

vitamin C treatment leads to decrease of inflammation markers such as CRP [104–106]. Our data

explains important aspects of these effects and further highlights the role of antioxidant

homeostasis in inflammatory processes.

In summary, our data support a critical role for HOCl-mediated *N*-chlorination of plasma

proteins during inflammatory processes and suggest that it is the critical modification

mediating the physiological effects of so-called AOPPs. Although the reversible conversion

of HOCl-modified plasma proteins to effective chaperones confers protection against HOCl-

induced protein aggregation, the increase of their immunogenicity can potentially exacerbate

self-damage at sites of inflammation through a positive feedback loop. The fact that the

activation of immune cells is mediated through *N*-chlorination, a modification that is

reversible by antioxidants present in plasma, provides a mechanism to attenuate or deactivate

this positive feedback loop. These findings contribute importantly to our understanding of the

development and progression of chronic inflammation.

Figure 10: Proposed mechanism of the immunomodulatory role of *N*-chlorination of plasma proteins at a site of inflammation. At the site of inflammation neutrophils (and potentially other immune cells) are activated. Neutrophils then produce HOCl at concentrations of up to 25 to 50 mM per hour. Plasma proteins, such as HSA, α- and β-globulins (Cohn Fraction IV), γ-globulins (IgG), and $α_2$-Macroglobulin then act as an effective sink. Reversible *N*-chlorination of these proteins turns them into effective chaperons, which can prevent the formation of protein aggregates and their uptake by immune cells, enhancing the survival of neutrophils in the presence of toxic antigens. *N*-chlorinated plasma proteins also activate more immune cells, which then in turn produce more HOCl, leading to the formation of more *N*-chlorinated plasma proteins. This positive feedback loop can be attenuated and deactivated by antioxidants present in plasma, such as ascorbate and reduced glutathione (GSH).

ACKNOWLEDGMENTS

Funding for this study was provided by the German Research Foundation (DFG) through

grant LE2905/1-2 to LIL as part of the priority program 1710 'Dynamics of Thiol-based

Redox Switches in Cellular Physiology'. Parts of this manuscript were written during a

Writing Retreat funded by the Ruhr-Universität Bochum Research School RURS^plus.

AUTHOR CONTRIBUTIONS

AU, AVS, AM, and LIL designed the study. AU, AVS and AM performed the experiments.

AU and AVS analyzed the results. NL purified α_2-Macroglobulin and Ag85B. AVS consulted

on the manuscript and AU and LIL wrote the manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflict of interest.

REFERENCES

1. Mortaz E, Alipoor SD, Adcock IM, Mumby S, Koenderman L. Update on Neutrophil

Function in Severe Inflammation. *Front Immunol*. 2018;9:2171.

doi:10.3389/fimmu.2018.02171

2. Thomas DC. The phagocyte respiratory burst: Historical perspectives and recent

advances. *Immunol Lett*. 2017;192:88-96. doi:10.1016/j.imlet.2017.08.016

3. Furtmüller PG, Burner U, Obinger C. Reaction of myeloperoxidase compound I with

chloride, bromide, iodide, and thiocyanate. *Biochemistry*. 1998;37(51):17923-17930.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=99221

60&retmode=ref&cmd=prlinks.

4. van Dalen CJ, Whitehouse MW, Winterbourn CC, Kettle AJ. Thiocyanate and chloride

as competing substrates for myeloperoxidase. *Biochem J*. 1997;327 (Pt 2(Pt 2):487-

492. file:///pmc/articles/PMC1218820/?report=abstract.

5.   Chapman ALP, Hampton MB, Senthilmohan R, Winterbourn CC, Kettle AJ.

Chlorination of bacterial and neutrophil proteins during phagocytosis and killing of

Staphylococcus aureus. *J Biol Chem*. 2002;277(12):9757-9762.

doi:10.1074/jbc.M106134200

6.   Rosen H, Crowley JR, Heinecke JW. Human neutrophils use the myeloperoxidase-

hydrogen peroxide-chloride system to chlorinate but not nitrate bacterial proteins

during phagocytosis. *J Biol Chem*. 2002;277(34):30463-30468.

doi:10.1074/jbc.M202331200

7.   Degrossoli A, Müller A, Xie K, et al. Neutrophil-generated HOCl leads to non-specific

thiol oxidation in phagocytized bacteria. *Elife*. 2018;7:6290. doi:10.7554/eLife.32288

8.   Clark RA, Szot S, Williams MA, Kagan HM. Oxidation of lysine side-chains of elastin

by the myeloperoxidase system and by stimulated human neutrophils. *Biochem*

Biophys Res Commun. 1986;135(2):451-457.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=30087

20&retmode=ref&cmd=prlinks.

9.   Beck-Speier I, Leuschel L, Luippold G, Maier KL. Proteins released from stimulated

neutrophils contain very high levels of oxidized methionine. *FEBS Lett*. 1988;227(1):1-

4.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=28281

08&retmode=ref&cmd=prlinks.

10.   Heinecke JW. Mass spectrometric quantification of amino acid oxidation products in

proteins: insights into pathways that promote LDL oxidation in the human artery wall.

FASEB J. 1999;13(10):1113-1120.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10385

603&retmode=ref&cmd=prlinks.

11.   Domigan NM, Charlton TS, Duncan MW, Winterbourn CC, Kettle AJ. Chlorination of

tyrosyl residues in peptides by myeloperoxidase and human neutrophils. *J Biol Chem*.

1995;270(28):16542-16548.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=76224

59&retmode=ref&cmd=prlinks.

12.   Sivey JD, Howell SC, Bean DJ, McCurry DL, Mitch WA, Wilson CJ. Role of lysine

during protein modification by HOCl and HOBr: Halogen-transfer agent or sacrificial

antioxidant? *Biochemistry*. 2013;52(7):1260-1271.

http://pubs.acs.org/doi/10.1021/bi301523s.

13.   Vissers MC, Winterbourn CC. Oxidative damage to fibronectin. I. The effects of the

neutrophil myeloperoxidase system and HOCl. *Arch Biochem Biophys*.

1991;285(1):53-59.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=18467

32&retmode=ref&cmd=prlinks.

14.   Hawkins CL, Davies MJ. Hypochlorite-induced damage to proteins: formation of

nitrogen-centred radicals from lysine residues and their role in protein fragmentation.

Biochem J. 1998;332:617-625.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=96208

62&retmode=ref&cmd=prlinks.

15.   Hazell LJ, van den Berg JJ, Stocker R. Oxidation of low-density lipoprotein by

hypochlorite causes aggregation that is mediated by modification of lysine residues

rather than lipid oxidation. *Biochem J*. 1994;302:297-304.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=80680

18&retmode=ref&cmd=prlinks.

16.   Heinecke JW, Li W, Daehnke HL, Goldstein JA. Dityrosine, a specific marker of

oxidation, is synthesized by the myeloperoxidase-hydrogen peroxide system of human

neutrophils and macrophages. *J Biol Chem*. 1993;268(6):4069-4077.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=83826

89&retmode=ref&cmd=prlinks.

17.   Müller A, Langklotz S, Lupilova N, Kuhlmann K, Bandow JE, Leichert LIO.

Activation of RidA chaperone function by N-chlorination. *Nat Commun*. 2014;5:5804.

http://www.nature.com/doifinder/10.1038/ncomms6804.

18.   Chapman ALP, Winterbourn CC, Brennan SO, Jordan TW, Kettle AJ. Characterization

of non-covalent oligomers of proteins treated with hypochlorous acid. *Biochem J*.

2003;375(Pt 1):33-40.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12852

783&retmode=ref&cmd=prlinks.

19.   Winter J, Ilbert M, Graf PCF, Ozcelik D, Jakob U. Bleach activates a redox-regulated

chaperone by oxidative protein unfolding. *Cell*. 2008;135(4):691-701.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=19013

278&retmode=ref&cmd=prlinks.

20.   Storkey C, Davies MJ, Pattison DI. Reevaluation of the rate constants for the reaction

of hypochlorous acid (HOCl) with cysteine, methionine, and peptide derivatives using

a new competition kinetic approach. *Free Radic Biol & Med*. 2014;73:60-66.

doi:10.1016/j.freeradbiomed.2014.04.024

21.   Arnhold J, Hammerschmidt S, Arnold K. Role of functional groups of human plasma

and luminol in scavenging of NaOCl and neutrophil-derived hypochlorous acid.

Biochim Biophys Acta. 1991;1097(2):145-151.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=16550

46&retmode=ref&cmd=prlinks.

22.   Pattison DI, Davies MJ. Absolute rate constants for the reaction of hypochlorous acid

with protein side chains and peptide bonds. *Chem Res Toxicol*. 2001;14(10):1453-

1464.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=11599

938&retmode=ref&cmd=prlinks.

23.   Chesney JA, Eaton JW, Mahoney JR. Bacterial glutathione: a sacrificial defense

against chlorine compounds. *J Bacteriol*. 1996;178(7):2131-2135.

http://www.ncbi.nlm.nih.gov/pubmed/8606194.

24.   Peskin A V, Winterbourn CC. Kinetics of the reactions of hypochlorous acid and

amino acid chloramines with thiols, methionine, and ascorbate. *Free Radic Biol &*

Med. 2001;30(5):572-579.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=11182

528&retmode=ref&cmd=prlinks.

25.   Goemans C V, Vertommen D, Agrebi R, Collet J-F. CnoX Is a Chaperedoxin: A

Holdase that Protects Its Substrates from Irreversible Oxidation. *Mol Cell*.

2018;70(4):614-627.e7. doi:10.1016/j.molcel.2018.04.002

26.   Pechous RD. With Friends Like These: The Complex Role of Neutrophils in the

Progression of Severe Pneumonia. *Front Cell Infect Microbiol*. 2017;7:160.

doi:10.3389/fcimb.2017.00160

27.   Thieblemont N, Wright HL, Edwards SW, Witko-Sarsat V. Human neutrophils in auto-

immunity. *Semin Immunol*. 2016;28(2):159-173. doi:10.1016/j.smim.2016.03.004

28.   Klebanoff SJ. Myeloperoxidase: friend and foe. *J Leukoc Biol*. 2005;77(5):598-625.

http://www.jleukbio.org/cgi/doi/10.1189/jlb.1204697.

29.   Davies MJ. Myeloperoxidase-derived oxidation: mechanisms of biological damage and

its prevention. *J Clin Biochem Nutr*. 2011;48(1):8-19. doi:10.3164/jcbn.11-006FR

30.   Parker H, Albrett AM, Kettle AJ, Winterbourn CC. Myeloperoxidase associated with

neutrophil extracellular traps is active and mediates bacterial killing in the presence of

hydrogen peroxide. *J Leukoc Biol*. 2012;91(3):369-376. doi:10.1189/jlb.0711387

31.   Summers FA, Morgan PE, Davies MJ, Hawkins CL. Identification of plasma proteins

that are susceptible to thiol oxidation by hypochlorous acid and N-chloramines. *Chem*

Res Toxicol. 2008;21(9):1832-1840. doi:10.1021/tx8001719

32.   Wyatt AR, Kumita JR, Mifsud RW, Gooden CA, Wilson MR, Dobson CM.

Hypochlorite-induced structural modifications enhance the chaperone activity of

human α2-macroglobulin. *Proc Natl Acad Sci USA*. 2014;111(20):E2081-90.

http://www.pnas.org/cgi/doi/10.1073/pnas.1403379111.

33.   Prütz WA. Hypochlorous acid interactions with thiols, nucleotides, DNA, and other

biological substrates. *Arch Biochem Biophys*. 1996;332(1):110-120.

doi:10.1006/abbi.1996.0322

34.   Carr AC, van den Berg JJ, Winterbourn CC. Chlorination of cholesterol in cell

membranes by hypochlorous acid. *Arch Biochem Biophys*. 1996;332(1):63-69.

doi:10.1006/abbi.1996.0317

35.   Winterbourn CC, van den Berg JJ, Roitman E, Kuypers FA. Chlorohydrin formation

from unsaturated fatty acids reacted with hypochlorous acid. *Arch Biochem Biophys*.

1992;296(2):547-555.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=13215

89&retmode=ref&cmd=prlinks.

36.   Shao B, Oda MN, Bergt C, et al. Myeloperoxidase impairs ABCA1-dependent

cholesterol efflux through methionine oxidation and site-specific tyrosine chlorination

of apolipoprotein A-I. *J Biol Chem*. 2006;281(14):9001-9004.

doi:10.1074/jbc.C600011200

37.   Peng D-Q, Wu Z, Brubaker G, et al. Tyrosine modification is not required for

myeloperoxidase-induced loss of apolipoprotein A-I functional activities. *J Biol Chem*.

2005;280(40):33775-33784. doi:10.1074/jbc.M504092200

38. Colombo G, Clerici M, Altomare A, et al. Thiol oxidation and di-tyrosine formation in

human plasma proteins induced by inflammatory concentrations of hypochlorous acid.

J Proteomics. 2017;152:22-32. doi:10.1016/j.jprot.2016.10.008

39. Himmelfarb J, McMonagle E. Albumin is the major plasma protein target of oxidant

stress in uremia. *Kidney Int*. 2001;60(1):358-363. doi:10.1046/j.1523-

1755.2001.00807.x

40. Fogh-Andersen N, Altura BM, Altura BT, Siggaard-Andersen O. Composition of

interstitial fluid. *Clin Chem*. 1995;41(10):1522-1525.

http://www.ncbi.nlm.nih.gov/pubmed/7586528.

41. Witko-Sarsat V, Friedlander M, Capeillère-Blandin C, et al. Advanced oxidation

protein products as a novel marker of oxidative stress in uremia. *Kidney Int*.

1996;49(5):1304-1313.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=87310

95&retmode=ref&cmd=prlinks.

42. Aratani Y. Myeloperoxidase: Its role for host defense, inflammation, and neutrophil

function. *Arch Biochem Biophys*. 2018;640:47-52. doi:10.1016/j.abb.2018.01.004

43. Colombo G, Clerici M, Giustarini D, et al. A central role for intermolecular dityrosine

cross-linking of fibrinogen in high molecular weight advanced oxidation protein

product (AOPP) formation. *Biochim Biophys Acta*. 2015;1850(1):1-12.

doi:10.1016/j.bbagen.2014.09.024

44. Liu SX, Hou FF, Guo ZJ, et al. Advanced oxidation protein products accelerate

atherosclerosis through promoting oxidative stress and inflammation. *Arterioscler

Thromb Vasc Biol*. 2006;26(5):1156-1162. doi:10.1161/01.ATV.0000214960.85469.68

45. Descamps-Latscha B, Witko-Sarsat V, Nguyen-Khoa T, et al. Advanced oxidation

protein products as risk factors for atherosclerotic cardiovascular events in nondiabetic

predialysis patients. *Am J Kidney Dis*. 2005;45(1):39-47.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=15696

442&retmode=ref&cmd=prlinks.

46. Witko-Sarsat V, Gausson V, Nguyen A-T, et al. AOPP-induced activation of human

neutrophil and monocyte oxidative metabolism: a potential target for N-acetylcysteine

treatment in dialysis patients. *Kidney Int*. 2003;64(1):82-91. doi:10.1046/j.1523-

1755.2003.00044.x

47. Gorudko I V, Grigorieva D V, Shamova E V, et al. Hypohalous acid-modified human

serum albumin induces neutrophil NADPH oxidase activation, degranulation, and

shape change. *Free Radic Biol & Med*. 2014;68:326-334.

http://linkinghub.elsevier.com/retrieve/pii/S0891584913015645.

48. Witko-Sarsat V, Friedlander M, Nguyen-Khoa T, et al. Advanced oxidation protein

products as novel mediators of inflammation and monocyte activation in chronic renal

failure. *J Immunol*. 1998;161(5):2524-2532.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=97252

52&retmode=ref&cmd=prlinks.

49. Imber MJ, Pizzo S V. Clearance and binding of two electrophoretic "fast"

forms of human alpha 2-macroglobulin. *J Biol Chem*. 1981;256(15):8134-8139.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=61675

73&retmode=ref&cmd=prlinks.

50. Leggio C, Galantini L, Pavel NV. About the albumin structure in solution: cigar

Expanded form versus heart Normal shape. *Phys Chem Chem Phys*. 2008;10(45):6741-

6750. doi:10.1039/b808938h

51. Johnstone AP, Thorpe R. *Immunochemistry in Practice*. Wiley-Blackwell; 1996.

http://books.google.de/books?id=sM6YQgAACAAJ&dq=Immunochemistry+in+

Practice&hl=&cd=1&source=gbs_api.

52. Murphy K, Weaver C. *Janeway's Immunobiology, 9th Edition*. Garland Science;

2016.

http://books.google.de/books?id=GmPLCwAAQBAJ&printsec=frontcover&

dq=Immunobiology+The+Immune+System+in+Health+and+Disease&hl=&

cd=6&source=gbs_api.

53.   Wilkins MR, Gasteiger E, Bairoch A, et al. Protein identification and analysis tools in

1169   the ExPASy server. *Methods Mol Biol*. 1999;112:531-552.

1170   http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10027

1171   275&retmode=ref&cmd=prlinks.

1172   54.   Udenfriend S, Stein S, Böhlen P, Dairman W, Leimgruber W, Weigele M.

1173   Fluorescamine: a reagent for assay of amino acids, peptides, proteins, and primary

1174   amines in the picomole range. *Science*. 1972;178(4063):871-872.

1175   http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=50859

1176   85&retmode=ref&cmd=prlinks.

1177   55.   Pivot-Pajot C, Chouinard FC, El Azreq MA, Harbour D, Bourgoin SG.

1178   Characterisation of degranulation and phagocytic capacity of a human neutrophilic

1179   cellular model, PLB-985 cells. *Immunobiology*. 2010;215(1):38-52.

1180   http://linkinghub.elsevier.com/retrieve/pii/S0171298509000205.

1181   56.   Tlili A, Dupré-Crochet S, Erard M, Nüsse O. Kinetic analysis of phagosomal

1182   production of reactive oxygen species. *Free Radic Biol & Med*. 2011;50(3):438-

1183   447. http://linkinghub.elsevier.com/retrieve/pii/S0891584910013961.

1184   57.   Minkenberg I, Ferber E. Lucigenin-dependent chemiluminescence as a new assay for

1185   NAD(P)H-oxidase activity in particulate fractions of human polymorphonuclear

1186   leukocytes. *J Immunol Methods*. 1984;71(1):61-67.

1187   http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=67259

1188   61&retmode=ref&cmd=prlinks.

58.   De Wit L, Palou M, Content J. Nucleotide sequence of the 85B-protein gene of

Mycobacterium bovis BCG and Mycobacterium tuberculosis. *DNA Seq*.

1994;4(4):267-270. http://www.ncbi.nlm.nih.gov/pubmed/7987013.

59.   Lajiness DH. *Trehalose Monomycolate Biosynthesis and Utilization in Mycobacterium*

Tuberculosis. University of Toledo; 2011.

60.   Temple A, Yen T-Y, Gronert S. Identification of specific protein carbonylation sites in

model oxidations of human serum albumin. *J Am Soc Mass Spectrom*.

2006;17(8):1172-1180. doi:10.1016/j.jasms.2006.04.030

61.   Sackett DL, Wolff J. Nile red as a polarity-sensitive fluorescent probe of hydrophobic

protein surfaces. *Anal Biochem*. 1987;167(2):228-234.

http://www.ncbi.nlm.nih.gov/pubmed/3442318.

62.   Aasen TB, Bolann B, Glette J, Ulvik RJ, Schreiner A. Lucigenin-dependent

chemiluminescence in mononuclear phagocytes. Role of superoxide anion. *Scand J*

Clin Lab Invest. 1987;47(7):673-679. http://www.ncbi.nlm.nih.gov/pubmed/2825342.

63.   Ellison, M. A., Thurman, G., Gearheart, C. M., Seewald, R. H., Porter, C. C., &

Ambruso DR. INF-γ enhances Nox2 activity by upregulating phox proteins when

applied to differentiating PLB-985 cells but does not induce Nox2 activity by itself.

PLoS One. 2015;10(8). doi:10.1371/journal.pone.0136766

64.   Wolfson M, McPhail LC, Nasrallah VN, Snyderman R. Phorbol myristate acetate

mediates redistribution of protein kinase C in human neutrophils: potential role in the

activation of the respiratory burst enzyme. *J Immunol*. 1985;135(3):2057-2062.

http://www.ncbi.nlm.nih.gov/pubmed/3160785.

65.   Chen X, Zhong Z, Xu Z, Chen L, Wang Y. 2′,7′-Dichlorodihydrofluorescein as a

fluorescent probe for reactive oxygen species measurement: Forty years of application

and controversy. *Free Radic Res*. 2010;44(6):587-604.

http://www.tandfonline.com/doi/full/10.3109/10715761003709802.

66.   Combadière C, el Benna J, Pedruzzi E, Hakim J, Périanin A. Stimulation of the human

neutrophil respiratory burst by formyl peptides is primed by a protein kinase inhibitor,

staurosporine. *Blood*. 1993;82(9):2890-2898.

http://www.ncbi.nlm.nih.gov/pubmed/8219237.

67. Bissonnette SA, Glazier CM, Stewart MQ, Brown GE, Ellson CD, Yaffe MB.

Phosphatidylinositol 3-phosphate-dependent and -independent functions of p40phox in

activation of the neutrophil NADPH oxidase. *J Biol Chem*. 2008;283(4):2108-2119.

doi:10.1074/jbc.M706639200

68. Simonsen A, Stenmark H. PX domains: attracted by phosphoinositides. *Nat Cell Biol*.

2001;3(8):E179-82. doi:10.1038/35087112

69. Perisic O, Wilson MI, Karathanassis D, et al. The role of phosphoinositides and

phosphorylation in regulation of NADPH oxidase. *Adv Enzyme Regul*. 2004;44:279-

298. doi:10.1016/j.advenzreg.2003.11.003

70. Martelli AM, Cocco L, Capitani S, Miscia S, Papa S, Manzoli FA. Nuclear

phosphatidylinositol 3,4,5-trisphosphate, phosphatidylinositol 3-kinase, Akt, and PTen:

emerging key regulators of anti-apoptotic signaling and carcinogenesis. *Eur J*

Histochem. 2007;51 Suppl 1:125-131.

http://www.ncbi.nlm.nih.gov/pubmed/17703603.

71. Belisle JT, Vissa VD, Sievert T, Takayama K, Brennan PJ, Besra GS. Role of the

major antigen of Mycobacterium tuberculosis in cell wall biogenesis. *Science*.

1997;276(5317):1420-1422. http://www.ncbi.nlm.nih.gov/pubmed/9162010.

72. Andersen P. Effective vaccination of mice against Mycobacterium tuberculosis

infection with a soluble mixture of secreted mycobacterial proteins. *Infect Immun*.

1994;62(6):2536-2544. http://www.ncbi.nlm.nih.gov/pubmed/7910595.

73. Baldwin SL, D'Souza C, Roberts AD, et al. Evaluation of new vaccines in the mouse

and guinea pig model of tuberculosis. *Infect Immun*. 1998;66(6):2951-2959.

http://www.ncbi.nlm.nih.gov/pubmed/9596772.

74.   Launois P, DeLeys R, Niang MN, et al. T-cell-epitope mapping of the major secreted

1243   mycobacterial antigen Ag85A in tuberculosis and leprosy. *Infect Immun*.

1244   1994;62(9):3679-3687. http://www.ncbi.nlm.nih.gov/pubmed/7520418.

1245   75.   Takamura S, Matsuo K, Takebe Y, Yasutomi Y. Ag85B of mycobacteria elicits

1246   effective CTL responses through activation of robust Th1 immunity as a novel

1247   adjuvant in DNA vaccine. *J Immunol*. 2005;175(4):2541-2547.

1248   http://www.ncbi.nlm.nih.gov/pubmed/16081827.

1249   76.   Alemán M, García A, Saab MA, et al. Mycobacterium tuberculosis-induced activation

1250   accelerates apoptosis in peripheral blood neutrophils from patients with active

1251   tuberculosis. *Am J Respir Cell Mol Biol*. 2002;27(5):583-592. doi:10.1165/rcmb.2002-

1252   0038OC

1253   77.   Perskvist N, Long M, Stendahl O, Zheng L. Mycobacterium tuberculosis promotes

1254   apoptosis in human neutrophils by activating caspase-3 and altering expression of

1255   Bax/Bcl-xL via an oxygen-dependent pathway. *J Immunol*. 2002;168(12):6358-6365.

1256   http://www.ncbi.nlm.nih.gov/pubmed/12055253.

1257   78.   Chae HJ, Kang JS, Byun JO, et al. Molecular mechanism of staurosporine-induced

1258   apoptosis in osteoblasts. *Pharmacol Res*. 2000;42(4):373-381.

1259   doi:10.1006/phrs.2000.0700

1260   79.   Zhang XD, Gillespie SK, Hersey P. Staurosporine induces apoptosis of melanoma by

1261   both caspase-dependent and -independent apoptotic pathways. *Mol Cancer Ther*.

1262   2004;3(2):187-197. http://www.ncbi.nlm.nih.gov/pubmed/14985459.

1263   80.   Vermes I, Haanen C, Steffens-Nakken H, Reutelingsperger C. A novel assay for

1264   apoptosis. Flow cytometric detection of phosphatidylserine expression on early

1265   apoptotic cells using fluorescein labelled Annexin V. *J Immunol Methods*.

1266   1995;184(1):39-51. http://www.ncbi.nlm.nih.gov/pubmed/7622868.

81.   Pullar JM, Vissers MC, Winterbourn CC. Living with a killer: the effects of

hypochlorous acid on mammalian cells. *IUBMB Life*. 2000;50(4-5):259-266.

doi:10.1080/713803731

82.   Annibal A, Colombo G, Milzani A, Dalle-Donne I, Fedorova M, Hoffmann R.

Identification of dityrosine cross-linked sites in oxidized human serum albumin. *J*

Chromatogr B Analyt Technol Biomed Life Sci. 2016;1019:147-155.

doi:10.1016/j.jchromb.2015.12.022

83.   Cao W, Hou FF, Nie J. AOPPs and the progression of kidney disease. *Kidney Int*

Suppl. 2014;4(1):102-106. doi:10.1038/kisup.2014.19

84.   Pattison DI, Hawkins CL, Davies MJ. What are the plasma targets of the oxidant

hypochlorous acid? A kinetic modeling approach. *Chem Res Toxicol*. 2009;22(5):807-

817. doi:10.1021/tx800372d

85.   French K, Yerbury JJ, Wilson MR. Protease activation of alpha2-macroglobulin

modulates a chaperone-like action with broad specificity. *Biochemistry*.

2008;47(4):1176-1185. doi:10.1021/bi701976f

86.   Rhee SG, Kang SW, Netto LE, Seo MS, Stadtman ER. A family of novel peroxidases,

peroxiredoxins. *Biofactors*. 1999;10(2-3):207-209. doi:10.1002/biof.5520100218

87.   Kopprasch S, Leonhardt W, Pietzsch J, Kühne H. Hypochlorite-modified low-density

lipoprotein stimulates human polymorphonuclear leukocytes for enhanced production

of reactive oxygen metabolites, enzyme secretion, and adhesion to endothelial cells.

Atherosclerosis. 1998;136(2):315-324. http://www.ncbi.nlm.nih.gov/pubmed/9543103.

88.   Kennedy AD, DeLeo FR. PI3K and NADPH oxidase: a class act. *Blood*.

2008;112(13):4788-4789. doi:10.1182/blood-2008-09-179135

89.   Hawkins PT, Davidson K, Stephens LR. The role of PI3Ks in the regulation of the

neutrophil NADPH oxidase. *Biochem Soc Symp*. 2007;(74):59-67.

doi:10.1042/BSS0740059

90.   Binder V, Ljubojevic S, Haybaeck J, et al. The myeloperoxidase product hypochlorous

acid generates irreversible high-density lipoprotein receptor inhibitors. *Arterioscler*

Thromb Vasc Biol. 2013;33(5):1020-1027.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=23493

288&retmode=ref&cmd=prlinks.

91.   Zhuang Z, Zhao X, Wu Y, et al. The anti-apoptotic effect of PI3K-Akt signaling

pathway after subarachnoid hemorrhage in rats. *Ann Clin Lab Sci*. 2011;41(4):364-372.

http://www.ncbi.nlm.nih.gov/pubmed/22166507.

92.   Chang F, Lee JT, Navolanic PM, et al. Involvement of PI3K/Akt pathway in cell cycle

progression, apoptosis, and neoplastic transformation: a target for cancer

chemotherapy. *Leukemia*. 2003;17(3):590-603. doi:10.1038/sj.leu.2402824

93.   Lee A, Whyte MK, Haslett C. Inhibition of apoptosis and prolongation of neutrophil

functional longevity by inflammatory mediators. *J Leukoc Biol*. 1993;54(4):283-288.

http://www.ncbi.nlm.nih.gov/pubmed/8409750.

94.   Colotta F, Re F, Polentarutti N, Sozzani S, Mantovani A. Modulation of granulocyte

survival and programmed cell death by cytokines and bacterial products. *Blood*.

1992;80(8):2012-2020. http://www.ncbi.nlm.nih.gov/pubmed/1382715.

95.   Vossmann M, Kirst M, Ludolfs D, Schreiber M. West Nile virus is neutralized by

HOCl-modified human serum albumin that binds to domain III of the viral envelope

protein E. *Virology*. 2008;373(2):322-328.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=18191

981&retmode=ref&cmd=prlinks.

96.   Speth C, Brodde MF, Hagleitner M, et al. Neutrophils turn plasma proteins into

weapons against HIV-1. *PLoS One*. 2013;8(6):e66073.

http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0066073.

97.   Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med*. 1989;320(6):365-376.

doi:10.1056/NEJM198902093200606

98.   Margaroli C, Garratt LW, Horati H, et al. Elastase Exocytosis by Airway Neutrophils

1321   Associates with Early Lung Damage in Cystic Fibrosis Children. *Am J Respir Crit*

1322   *Care Med*. October 2018. doi:10.1164/rccm.201803-0442OC

1323   99.   Erwig L-P, Henson PM. Immunological consequences of apoptotic cell phagocytosis.

1324   *Am J Pathol*. 2007;171(1):2-8. doi:10.2353/ajpath.2007.070135

1325   100.   Bonham MJ, Abu-Zidan FM, Simovic MO, et al. Early ascorbic acid depletion is

1326   related to the severity of acute pancreatitis. *Br J Surg*. 1999;86(10):1296-1301.

1327   doi:10.1046/j.1365-2168.1999.01182.x

1328   101.   Ruan EA, Rao S, Burdick JS, et al. Glutathione levels in chronic inflammatory

1329   disorders of the human colon. *Nutr Res*. 1997;17(3):463-473. doi:10.1016/S0271-

1330   5317(97)00007-9

1331   102.   Huijskens MJAJ, Wodzig WKWH, Walczak M, Germeraad WT V, Bos GMJ.

1332   Ascorbic acid serum levels are reduced in patients with hematological malignancies.

1333   *Results Immunol*. 2016;6:8-10. doi:10.1016/j.rinim.2016.01.001

1334   103.   Rose S, Melnyk S, Pavliv O, et al. Evidence of oxidative damage and inflammation

1335   associated with low glutathione redox status in the autism brain. *Transl Psychiatry*.

1336   2012;2:e134. doi:10.1038/tp.2012.61

1337   104.   Mikirova N, Casciari J, Rogers A, Taylor P. Effect of high-dose intravenous vitamin C

1338   on inflammation in cancer patients. *J Transl Med*. 2012;10:189. doi:10.1186/1479-

1339   5876-10-189

1340   105.   Carr AC, Cook J. Intravenous Vitamin C for Cancer Therapy - Identifying the Current

1341   Gaps in Our Knowledge. *Front Physiol*. 2018;9:1182. doi:10.3389/fphys.2018.01182

1342   106.   Block G, Jensen CD, Dalvi TB, et al. Vitamin C treatment reduces elevated C-reactive

1343   protein. *Free Radic Biol Med*. 2009;46(1):70-77.

1344   doi:10.1016/j.freeradbiomed.2008.09.030

107.   Thomas EL, Grisham MB, Jefferson MM. Preparation and characterization of

chloramines. *Methods Enzymol*. 1986;132:569-585.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=38215

26&retmode=ref&cmd=prlinks.

108.   Carr AC, Hawkins CL, Thomas SR, Stocker R, Frei B. Relative reactivities of N-

chloramines and hypochlorous acid with human plasma constituents. *Free Radic Biol*

& Med. 2001;30(5):526-536.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=11182

523&retmode=ref&cmd=prlinks.

109.   Thomas EL, Grisham MB, Jefferson MM. Cytotoxicity of chloramines. *Methods*

Enzymol. 1986;132:585-593.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=35470

27&retmode=ref&cmd=prlinks.

110.   Bournazos S, Ravetch J V. Fcγ Receptor Function and the Design of Vaccination

Strategies. *Immunity*. 2017;47(2):224-233. doi:10.1016/j.immuni.2017.07.009

111.   Irani V, Guy AJ, Andrew D, Beeson JG, Ramsland PA, Richards JS. Molecular

properties of human IgG subclasses and their implications for designing therapeutic

monoclonal antibodies against infectious diseases. *Mol Immunol*. 2015;67(2 Pt A):171-

182. doi:10.1016/j.molimm.2015.03.255

112.   Scott AM, Allison JP, Wolchok JD. Monoclonal antibodies in cancer therapy. *Cancer*

Immun. 2012;12:14. http://www.ncbi.nlm.nih.gov/pubmed/22896759.

1366