1

Title:

Modular tissue-specific regulation of *doublesex* underpins sexually dimorphic development

in *Drosophila*

Running Title:

Tissue-specific regulation of *dsx*

Gavin R. Rice[1,2], Olga Barmina[1], David Luecke[1,4], Kevin Hu[1,5], Michelle Arbeitman[3],

Artyom Kopp[1]*

[1] Department of Evolution and Ecology, University of California–Davis, Davis, California,

United States of America,

[2]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA

[3]Department of Biomedical Sciences, Florida State University, Tallahassee, FA, USA

[4] Current address: Department of Integrative Biology, Michigan State University, East Lansing,

MI, USA

[5] Current address: Department of Pathology, University of Michigan, Ann Arbor, MI, USA

*Corresponding Author: akopp@ucdavis.edu

Key words:

Sexual Dimorphism

Drosophila

doublesex

Sex comb

Enhancers

Summary Statement:

We identify the modular *cis*-regulatory elements that direct expression of *doublesex* in sexually

dimorphic structures in *Drosophila* legs and genitalia. This regulatory landscape provides insight

into how cells obtain their sex-specific identity.

Abstract:

The ability of a single genome to produce distinct and often dramatically different male and

female forms is one of the wonders of animal development. In most animals, sex-specific

phenotypes are shaped by interactions between a sex determination pathway and spatial

patterning gene networks. In *Drosophila melanogaster,* most sexually dimorphic traits are

controlled by sex-specific isoforms of the *doublesex* (*dsx*) transcription factor, and *dsx*

expression is mostly limited to cells that give rise to sexually dimorphic traits. However, it is

unknown how this mosaic of "sex-naïve" and "sex-aware" tissues arises. Here, we characterize

the *cis*-regulatory sequences that control *dsx* expression in the foreleg, which contains multiple

types of sex-specific sensory organs. We find that separate modular enhancers are responsible for

dsx expression in each sexually dimorphic organ. Expression of *dsx* in the sex comb is co-

regulated by two enhancers with distinct spatial and temporal specificities that are separated by a

genitalia-specific enhancer. Thus, the mosaic of sexually dimorphic and monomorphic organs

depends on modular regulation of *dsx* transcription by dedicated cell type-specific enhancers.

Introduction:

Most animals are mosaics of sexually dimorphic and monomorphic tissues. Despite the multitude of traits that distinguish males from females, many tissues and organs lack overt sexual dimorphism. To understand how this mosaic pattern is produced, the action of sex determination pathways needs to be understood at the cellular level.

Sexually dimorphic morphologies are specified by diverse molecular mechanisms in different animal phyla (Kopp, 2012; Matson and Zarkower, 2012). One of the best studied mechanisms is found in the fruit fly, *Drosophila melanogaster*, where sex-specific development of most somatic cells is controlled by an alternative pre-mRNA splicing pathway (reviewed in Christiansen et al., 2002; Cline, 1993; Erickson and Quintero, 2007; McKeown, 1992). In *Drosophila*, the number of X chromosomes sets off a cell-autonomous cascade of sex-specific splicing that leads to the production of functional Transformer (Tra) protein in females, but not in males. Tra controls alternative splicing of *doublesex* (*dsx*) pre-mRNAs, such that the presence of functional Tra in females leads to a female-specific *dsx* isoform, and the absence of functional Tra leads to the production of a male-specific isoform (dsx^F and dsx^M, respectively). *dsx* encodes transcription factors involved in the establishment of almost all morphological traits that differ between males and females (Baker and Ridge, 1980; Hildreth, 1965). The Dsx^M and Dsx^F proteins share a common DNA binding domain and bind to the same target sequences, but have different and sometimes opposite effects on gene expression, leading to sex-specific cell differentiation (Arbeitman et al., 2004; Arbeitman et al., 2016; Burtis et al., 1991; Goldman and Arbeitman, 2007; Lebo et al., 2009; Li and Baker, 1998; Yang et al., 2008). For instance, the *yolk protein 1, bric-à-brac*, *Fmo-2,* and *desat-F* genes are directly regulated by Dsx and are expressed at higher levels in females compared to males. *dsx* mutants show intermediate expression of *yolk protein 1, bric-à-brac*, and *Fmo-2* in both sexes, indicating that Dsx^F is likely activating, and Dsx^M repressing, these genes (Coschigano and Wensink, 1993; Luo and Baker, 2015; Williams et al., 2008). *desat-F,* on the other hand, is activated by Dsx^F but is not affected by Dsx^M (Shirangi et al., 2009). At the morphological level, loss-of-function *dsx* mutants develop as intersexes with a mixture of male, female, and intermediate traits (Baker and Ridge, 1980; Hildreth, 1965).

The details of the *tra*/*dsx* splicing cascade were characterized in the 1980s (Boggs et al., 1987; Burtis and Baker, 1989; Butler et al., 2018; McKeown et al., 1987; McKeown et al., 1988;

Nagoshi et al., 1988), and are a textbook example of the role of alternative splicing in

development. In contrast, the role of transcriptional regulation of *dsx* in *Drosophila* sexual

differentiation was realized only recently. Although it may seem that every cell would need to

"know" its sex, *dsx* is only transcribed in cells that are associated with sexually dimorphic organs

(Camara et al., 2008; Rideout et al., 2010; Robinett et al., 2010; Tanaka et al., 2011). Thus, both

males and females are mosaics of sexually dimorphic and monomorphic cells. This mosaicism

makes it all the more remarkable that *dsx* regulates the development of so many different sex-

specific traits, from pigmentation and genital morphology to brain neural circuits and gene

expression in the gut (Goldman and Arbeitman, 2007; Keisman et al., 2001; Luo and Baker,

2015; Sanders and Arbeitman, 2008; Williams et al., 2008). To understand how this mosaic

pattern is specified, we need to elucidate the mechanisms that establish *dsx* transcription.

Among the best studied sexually dimorphic organs in *Drosophila* are the foreleg bristles.

The foreleg of *D. melanogaster* displays two distinct sex-specific features. First, the sex comb –

a row of strongly modified mechanosensory bristles – is present only in males, while in females

the homologous bristles retain the typical mechanosensory bristle morphology (Kopp, 2011;

Tokunaga, 1962). Second, males have a greater number of foreleg chemosensory bristles

compared to females (Mellert et al., 2012; Tokunaga, 1962). The sex comb and chemosensory

bristles are found in close proximity to each other, and both are involved in mating behavior (Fan

et al., 2013; Hurtado-Gonzales et al., 2015; Ng and Kopp, 2008; Spieth, 1952). The male-

specific chemosensory bristles are used early in the stereotypic courtship sequence to taste the

female cuticular pheromones (Fan et al., 2013; Spieth, 1952), while the sex comb assists in

grasping the female prior to copulation (Hurtado-Gonzales et al., 2015; Ng and Kopp, 2008;

Spieth, 1952).

dsx is required both for the distinctive sex comb morphology and for the higher number

of chemosensory bristles in males (Belote and Baker, 1982; Mellert et al., 2012). Expression of

dsx in the developing foreleg starts in the late 3rd instar larva (Tanaka et al., 2011). During

prepupal and early pupal development, this pattern resolves into several distinct clusters of cells

corresponding to chemosensory bristle precursors, sex comb bristle precursors, and surrounding

epithelial cells (Mellert et al., 2012; Robinett et al., 2010; Tanaka et al., 2011). This raises a

question – is *dsx* expression in the foreleg controlled by a different, modular enhancer for each

sex-specific bristle type, or by a common leg enhancer? More generally, is *dsx*, despite its

apparently global function as a binary sex switch, subject to modular transcriptional control, with

a dedicated enhancer for every cell population that expresses *dsx*?

To address this question, we identified and characterized the enhancers that control *dsx*

expression in the sex comb and the foreleg chemosensory bristles. We found that three separate

enhancers are responsible for regulating *dsx* expression in the foreleg. One enhancer drives broad

foreleg expression that encompasses both sex comb and chemosensory organ primordia, while

the other two have mutually exclusive activities – one in the sex comb, and the other in sex-

specific chemosensory bristles. The two enhancers that contribute to sex comb development are

separated by a genitalia-specific enhancer that has no activity in the leg. This complex *cis*-

regulatory architecture suggests that *dsx* transcription, like that of other developmentally

regulated transcription factors, is controlled by modular tissue-specific enhancers that can have

both overlapping and non-overlapping spatial activities.

Results:

Three enhancers with distinct spatial and temporal activities contribute to complex dsx *expression in the foreleg*

To identify the enhancers that specify the complex expression pattern of *dsx* during foreleg development, we generated a series of pPTGAL GAL4 reporter lines covering the nearly 50 kb of the upstream and intronic regions of the *dsx* locus (brown boxes, Fig. S1). In parallel, we examined an overlapping series of pBPGUw GAL4 reporters generated using a different vector and integration method (Pfeiffer et al., 2008) (grey boxes, Fig. S1). These lines were crossed to UAS-GFP.nls and screened 5 hr and 24 hr after puparium formation (APF). Using both sets of reporters, we identified three DNA fragments that drove expression patterns consistent with known *dsx* expression in the foreleg (Rideout et al., 2010; Robinett et al., 2010; Tanaka et al., 2011). We identified a chemosensory enhancer in a ~4 kb fragment of the first *dsx* intron (Fig. 1, F,G and Sup Fig. 1B, blue text), a sex comb enhancer in a ~3 kb section of the second intron (Sup Fig. 1C, red and green text), and an early foreleg enhancer in a ~4 kb section of the second intron downstream from the sex comb enhancer (Sup Fig. 1C, purple and green text). In parallel, another research group had identified a genital enhancer in the overlap between the 40F03 and 42D04 reporters (Sup Fig. 1C) (John Yoder, University of Alabama, personal communication), both of which showed foreleg expression in our screen. To test whether the sex comb and early foreleg enhancers were separate, and to reduce the temporal lag between enhancer activation and reporter detection, we cloned smaller sub-fragments of the *dsx* second intron into a nuclear GFP expression vector. We found that both the sex comb and the early foreleg enhancers drove the same expression patterns in this assay as in the larger GAL4 reporter constructs (Fig. 1 B-E).

The three enhancers direct distinct spatial and temporal expression of reporter genes, consistent with known *dsx* expression. *dsx* expression in the foreleg first becomes visible in wandering third instar larvae and is well developed in prepupal legs (Tanaka et al., 2011). Leg bristles are specified over several developmental periods. Most chemosensory bristles and the largest mechanosensory bristles are specified at the end of the 3^{rd} larval instar or early in prepupal development, while most mechanosensory bristles (including sex comb teeth) are specified between 6 and 15 hr APF (Belote and Baker, 1982; Held, 2002). At 5 hr APF, Dsx is present in an anterior-ventral region of tarsal segments 1 through 4 (ta1-4), where the sex comb and sex-specific chemosensory bristles will form (Tanaka et al., 2011). At this stage, the early foreleg

enhancer drives reporter expression in a pattern similar to Dsx protein (Fig 1B). Although Dsx expression is already becoming sexually dimorphic by 5 hr APF (Tanaka et al., 2011), the GFP reporter intensity at this stage is roughly monomorphic, possibly due to the higher stability of the GFP protein (Fig 1B). The sex comb enhancer is not active at 5 hrs APF (Fig 1D), while the chemosensory enhancer is expressed in both sexes in small clusters of cells that match the pattern of developing chemosensory bristles (Fig 1F).

By 28 hr APF, Dsx expression is strongly dimorphic; in males, high Dsx expression is seen in sex comb bristles and surrounding epithelial cells, whereas in females, the homologous region of ta1 shows much lower Dsx expression (Tanaka et al., 2011; Fig 1C). In the rest of the tarsus, epithelial expression of Dsx disappears by 28 hr APF. At this stage in males, the early foreleg enhancer still shows weak activity in epithelial cells around the sex comb, especially on the distal-ventral side; however, it is not visibly active in sex comb bristles (Fig 1C). The sex comb enhancer drives a complementary pattern, with strong expression in sex comb bristles and weak expression in the surrounding epithelial cells (Fig 1E). In females, neither of these enhancers shows significant expression at 28 hr APF, consistent with the sexually dimorphic expression of Dsx (Fig 1C,E). For the chemosensory enhancer, cell clusters that reflect the pattern of chemosensory bristles continue to be observed at 28 hr APF in both males and females (Fig 1G). Thus, each of the three enhancers directs a distinct expression pattern in the foreleg.

All three enhancers contribute to sex-specific sensory organ development:

The patterns of GFP reporter expression suggest that the sex comb enhancer controls sex comb development and the chemosensory enhancer controls sex-specific chemosensory bristle development, while the early foreleg enhancer could potentially contribute to the development of both types of sex-specific bristles. To test these hypotheses, we used the chemosensory (42C06), foreleg (42D04), and sex comb (40F03) pBPGUw-GAL4 lines to drive a UAS-RNAi construct targeting the *shaven/Pax2* gene (*sv*). Knockdown of *sv* in sensory organ precursors causes severe truncation of bristle shafts (Kavaler et al., 1999), allowing us to identify the bristles where each *dsx* enhancer is active during prepupal and pupal development. We collected and analyzed 8 male and 5 female samples for each of the three GAL4 lines. This approach also allowed us to test for enhancer activity at developmental stages that were not observed by visual inspection of GFP

reporters. The chaetotaxy of ta1 is particularly stereotypic and has been thoroughly mapped

(Tokunaga, 1962), allowing us to compare the male and female knockdown phenotypes.

First, to identify all *dsx*-expressing bristles, we expressed UAS-*sv*RNAi using a *dsx*-GAL4

knock-in line, which was generated by inserting the GAL4 coding sequence into the *dsx* locus and

reflects the full expression pattern of *dsx* (Robinett et al., 2010). We found that the ta1

chemosensory bristles fall into three classes (Fig 2). First, the dorsal-posterior chemosensory

bristles are not affected by *dsx*-GAL4/UAS-*sv*RNAi; in the wild type these bristles are sexually

monomorphic in size and position, suggesting they do not express or require *dsx* for their

development (Fig 2A, B). Second, on the dorsal-anterior side, males have four chemosensory

bristles on the distal half of ta1 (dark orange triangles in Fig 2A), compared to only one in females

(Fig 2B); these bristles were always strongly affected by *sv* RNAi in both sexes, indicating that

they express *dsx* (Fig 2A,B). The third class, two chemosensory bristles (yellow triangles in Fig

2A,B) were affected weakly and with more variability among individuals, suggesting that they

may express *dsx* at lower levels or for a briefer period.

The male sex comb was always severely affected by *dsx*-GAL4/UAS-*sv*RNAi, whereas the

homologous female bristles were affected weakly and with more variability (Fig 2A,B), consistent

with a lower level of *dsx* expression in females compared to males (Tanaka et al., 2011). With the

exception of the sex comb and homologous female bristles, no mechanosensory bristles were

affected by *dsx*-GAL4/UAS-*sv*RNAi (Fig 2A,B), confirming their sexually monomorphic nature.

Next, we used each of the three *dsx* leg enhancers to drive UAS-*sv*RNAi and compared the

resulting phenotypes in ta1 to that of the *dsx* knock-in. The early foreleg enhancer was similar to

the *dsx* knock-in in having a strong and consistent effect on the male sex comb, but only a weak

and variable effect on homologous bristles in females (Fig 2C,D). It also affected a single

chemosensory bristle in proximal ta1 in both sexes (Fig 2C,D). As expected, the sex comb

enhancer had a clear effect on sex comb development in males, but did not affect chemosensory

bristles or any bristles in females (Fig 2E,F). The chemosensory enhancer consistently affected all

anterior-ventral chemosensory bristles in males, including the two proximal bristles that were only

weakly affected by the *dsx* knock-in; it did not affect any mechanosensory bristles (Fig 2G).

Surprisingly, the chemosensory enhancer had no effect in females (Fig 2H), even though some

female chemosensory bristles express *dsx* (Fig 2B), and the chemosensory enhancer drives UAS-

GFP expression in ta1 in females (Fig 1C).

In addition to their function in ta1, the foreleg and chemosensory enhancers show
expression in the more distal segments (ta2-ta4) (Fig 1B, F). These segments also contain
chemosensory bristles, some of which are sexually dimorphic (Mellert et al., 2012; Tokunaga,
1962). In ta2 and ta3, the *dsx* knock-in and the chemosensory enhancer affect the same subset of
four chemosensory bristles, while the sex comb enhancer has no effect (Fig Sup 2). The early
foreleg enhancer affected some but not all of the bristles affected by the *dsx* knock-in, as well as
one of the chemosensory bristles that was not affected by the *dsx* knock-in or the chemosensory
enhancer (Fig Sup 2). Pupal expression of the early foreleg enhancer is epithelial, not bristle-
specific (Fig 1E); thus, it is difficult to say whether this phenotype reflects a true developmental
function of the early foreleg enhancer in sensory organs, or the perdurance of the GAL4 protein or
the RNAi effect from an earlier time in development.

Examination of adult bristle phenotypes caused by UAS-*sv*RNAi expression suggests that
the three leg enhancers of *dsx* make distinct contributions to sex-specific bristle development. The
foreleg and sex comb enhancers function in sex comb development, while the chemosensory
enhancer functions in the development of sexually dimorphic chemosensory organs. We also
cannot rule out some contribution of the early foreleg enhancer to chemosensory bristle
development.

234   *The foreleg and sex comb enhancers are modular and flank a separate genital enhancer*

The foreleg and sex comb enhancers are located in close proximity within the second intron
of *dsx* and are both active in the sex comb, although in different spatial patterns and at different
times (Fig 1B-E). The same broad region also contains an enhancer that is active in the third instar
larval genital disc (John Yoder, University of Alabama, personal communication). Mutations in
dsx have been previously shown to disrupt clasper (also known as the surstylus) bristles (Hildreth,
1965), which have a morphology similar to sex comb teeth. Through antibody staining, we found
that Dsx is expressed in several regions of the pupal male genitalia (Fig. S3). In the developing
clasper, Dsx is enriched in the bristle precursor cells, similar to its enrichment in sex comb bristles.
Analysis of the genital enhancer showed weak expression in the clasper as well as strong
expression in the dorsal postgonites (Fig. S3, S4). These observations led us to inquire whether
the foreleg, genital, and sex comb enhancers were fully modular and independent, or parts of a
broader pleiotropic enhancer with overlapping activities in multiple tissues. To distinguish

between these possibilities, we generated non-overlapping reporter constructs for each enhancer

and examined each reporter both in the foreleg and in the genitalia. In addition, we generated larger

constructs where either the sex comb or the early foreleg enhancer was excluded (Fig 3).

We first confirmed that a 5.5 kb genomic fragment containing all three candidate enhancers

(sex comb, genital, and foreleg) was able to drive both early and late expression in the foreleg, as

well as genital expression (Fig. 3 and Fig. S4). Each separate enhancer can be seen as driving a

distinct subset of that overall pattern. The late sex comb enhancer is active in the sex comb bristle

cells but not in the genitalia, while the genital enhancer drives expression in the genitalia but has

no activity in the foreleg (Fig. 3 and Fig. S4). The construct encompassing the early foreleg

enhancer is active in the epithelial cells surrounding the sex comb, as well as in the genitalia in a

pattern that resembles the genital enhancer (Fig. S4). However, further dissection of this region

shows that the foreleg and genital activities are largely separable. The sequences necessary for

accurate foreleg expression are located on the 3' side of the foreleg enhancer and do not drive

genital expression, while the sequences that activate expression in the genitalia are located on the

5' side and only drive ectopic expression in the foreleg (Fig. 4 G-H and Fig. S3). Pairwise

combinations of enhancers behave in a predictably modular fashion: a construct including the sex

comb and genital enhancers drives genital and late sex comb expression, but no early epithelial

expression in the foreleg, while the construct including the genital and early foreleg enhancers

drives early foreleg and genital expression but no expression in sex comb teeth (Fig. 3 and Fig.

S4). The chemosensory enhancer, located in a different intron and thus clearly distinct from the

foreleg and sex comb enhancers (Fig 3). In summary, we conclude that the three leg enhancers are

all fully separable, modular elements, but we cannot rule out some minor overlap between the

foreleg and genital enhancers.

Sequences in the foreleg and sex comb enhancers confine their activity to sexually dimorphic

primordia

To identify the minimal sequences required for correct activity, we subdivided the sex

comb and early foreleg enhancers into smaller reporter constructs (Fig 4A). While the ~3kb sex

comb enhancer drives specific expression in sex comb bristles and the epithelial cells distal to the

sex comb (Fig 1E), shorter sequences are unable to recapitulate this pattern. The 3' ~1.5 kb of the

sex comb enhancer (*sex comb enhancer sub1*) drives specific expression in sex comb bristles but

is not expressed in the epithelial cells (Fig 4B). A shorter ~1 kb 3' fragment (*sex comb enhancer*

sub2) retains expression in the sex comb bristles, but also causes ectopic expression in the most

distal TBR (Fig 4C). A core sequence of ~0.5 kb (*sex comb enhancer sub3*) expands expression to

all ta1 TBRs (Fig 4D). The TBRs are serially homologous to the sex comb (Kopp, 2011; Tokunaga,

1962), but are sexually monomorphic and do not express Dsx (Robinett et al., 2010; Tanaka et al.,

2011).

The 5' ~1.5kb of the sex comb enhancer (*sex comb enhancer sub4*) drives expression in

some of the epithelial cells surrounding the sex comb, but this expression does not encompass all

of the epithelial cells where the full sex comb enhancer is active (Fig. 4E). Reduction of this

sequence to a ~1kb 5' fragment (*sex comb enhancer sub5*) causes ectopic expression in proximal

epithelial cells (Fig 4F) that do not express Dsx (Robinett et al., 2010; Tanaka et al., 2011). Thus,

sequences within the sex comb enhancer are necessary to prevent its activity in sexually

monomorphic bristle and epithelial cells.

Dissection of the early foreleg enhancer identified a shorter 3' fragment (*early foreleg*

enhancer sub2) capable of driving expression in the correct region of the foreleg (Fig. 4H). A

partially overlapping fragment (*early foreleg enhancer sub1*) shows broad ectopic activity

throughout the tarsal segments (Fig 4G). Thus, similar to the late sex comb enhancer, both

activating and repressive sequences are needed to limit this enhancer's activity to sexually

dimorphic cell populations.

Discussion:

dsx expression in the foreleg is controlled by modular enhancers with both distinct and

overlapping activities

In this study, we have characterized the *cis*-regulatory architecture directing the precise

spatiotemporal and cell type-specific expression of *dsx* in the foreleg. Our results help explain the

origin of the developmental mosaicism that generates a mix of sex-specific and sexually

monomorphic organs in *Drosophila*, and possibly in other insects. Given the largely cell-

autonomous control of insect sexual differentiation, the essential function of *dsx* enhancers is to

provide spatial landmarks that establish which organs will become sexually dimorphic. In

principle, *dsx* expression in different types of foreleg bristles could be controlled either by a single,

broadly acting leg enhancer, or by multiple enhancers, each dedicated to a different bristle type.

For example, the proneural genes *achaete* and *scute*, which control bristle specification, have very

complex *cis*-regulatory regions with dozens of modular enhancers responsible for different,

stereotypically patterned bristles (Garcia-Bellido and de Celis, 2009). Other enhancers are

pleiotropic and active in multiple tissues, reflecting a history of evolutionary co-option from the

more ancient organs to the more recently evolved (Glassford et al., 2015; Nagy et al., 2018; Rice

and Rebeiz, 2019). In fact, some genes contain both modular (tissue-specific) and pleiotropic

(multi-tissue) enhancers (Preger-Ben Noon et al., 2018).

We observe a partial overlap in the temporal and spatial activities of the three leg enhancers

(Fig 1-3). However, these are not functionally redundant "shadow enhancers" of the type found in

some other genes (Frankel et al., 2010; Hong et al., 2008). During prepupal development, the early

foreleg enhancer is expressed in cells that give rise both to the sex comb and to the chemosensory

bristles (Fig 2, Sup Fig 2). Later, the early foreleg enhancer expression in the sex comb becomes

complementary to the activity of the sex comb enhancer: the former is expressed only in the

epithelial cells, and the latter predominantly in sex comb bristles and only weakly in the epithelial

cells. At pupal stages, when the early foreleg enhancer is no longer active in bristle cells, the late-

acting sex comb and chemosensory enhancers have completely non-overlapping expression

patterns. Overall, the existence of these three modular enhancers suggests that *dsx* expression in

the foreleg goes through two successive stages: relatively broad expression early in development,

followed by tightly limited, cell type specific expression later on.

These phases of *dsx* expression may have different developmental functions. For example, *dsx* knockdown in chemosensory bristles late in bristle development causes a defect in the axonal midline crossing by the bristle neurons (Mellert et al., 2012). Early *dsx* expression in the foreleg induces a male-specific increase in the number of bristles both in the sex comb and in the chemosensory system (Belote and Baker, 1982; Mellert et al., 2012). The broad early foreleg expression of *dsx* is established prior to the formation of mechanosensory bristles (Tanaka et al., 2011), suggesting that the early foreleg enhancer may control sex comb size by modulating the specification of sensory organ precursors. On the other hand, the later-acting sex comb enhancer is more likely to be involved in controlling the morphology of the individual sex comb teeth, which develop from mechanosensory bristle shafts but, in contrast to most such bristles, are thick, blunt, and darkly pigmented. Indeed, knockdown of *dsx* late in development affects the morphology but not the number of sex comb teeth, which become masculinized in females and feminized in males (Belote and Baker, 1982). Both the foreleg and the sex comb enhancers are active in the epithelial cells around the sex comb, suggesting that both could contribute to sex comb rotation – a coordinated sequence of cell shape changes that moves the sex comb from an initial transverse to the final longitudinal orientation (Atallah et al., 2009; Ho et al., 2018; Malagón et al., 2014; Tanaka et al., 2011). Since sex comb development involves so many different cellular processes spread over multiple developmental stages, it is perhaps not surprising that it requires multiple *dsx* enhancers.

Not all tissues and cell types behave in a modular fashion with respect to *dsx* expression. In the central nervous system, only one of the pPTGAL reporters tiling the *dsx* region showed an overlap with endogenous *dsx* expression (in 4 neurons in the PC2 cluster) (M. Arbeitman, unpublished). A separate study identified a *dsx* enhancer upstream of the transcription start with an activity overlapping Dsx protein expression in the brain (Zhou et al., 2014). However, most of the Dsx CNS expression pattern is not accounted for by any of the *dsx* reporter constructs (M. Arbeitman, unpublished). This suggests that for most *dsx*-expressing cells in the CNS, *dsx* regulation may not be due to modular enhancers acting independently, but rather to more complex, long-distance sequence interactions that are not captured by ~5 kb reporter fragments. If true, such "entangled" regulatory architecture could be more constraining than the modular enhancer organization we observe in the foreleg, which could in turn lead to slower evolution of *dsx* expression in the brain compared to other tissues.

cis-regulatory information limits dsx expression to sexually dimorphic cells

cis-regulatory information limits dsx expression to sexually dimorphic cells

In contrast to some enhancers where very short sequences are sufficient to convey correct

regulatory information (Grieder et al., 1997; Manak et al., 1994; Swanson et al., 2010), both the

sex comb and the early foreleg enhancers are quite large and cannot be reduced to a core region

without perturbing their activities. Interestingly, dissecting these enhancers further leads not to the

loss of activity, but rather to its expansion. This is similar to the abdominal enhancer of the *ebony*

gene, where a core 0.7 kb sequence is sufficient for abdominal activity, but a much broader

genomic region is necessary to prevent ectopic expression (Rebeiz et al., 2009). We find a similar

pattern in our study: both the foreleg and the late sex comb enhancers of *dsx* have core sequences

of less than 1 kb capable of driving expression in the foreleg, but require additional sequences to

repress ectopic activity.

The late sex comb enhancer is particularly illuminating in this respect. The sex comb

develops from the most distal transverse bristle row on ta1 (Atallah et al., 2009; Tanaka et al.,

2009; Tokunaga, 1962), so the bristles that comprise it are serially homologous to the more

proximal TBRs. However, while the distal TBR that gives rise to the sex comb undergoes dramatic

male-specific modification, the remaining TBRs retain their original, sexually monomorphic

morphology. Consistent with this, *dsx* is expressed, and the sex comb enhancer of *dsx* is active,

only in the most distal TBR. Loss of sequences at the 5' end of this enhancer leads to an expansion

of its activity to the proximal, sexually monomorphic TBRs. Thus, much of the regulatory

information contained in the sex comb enhancer is devoted to restricting its activity to sexually

dimorphic cells, from a broader population of similar cells. This may reflect a general trend for

dsx expression: in each tissue or body part (such as the brain, midgut, foreleg, body wall epidermis,

etc.), only a subset of cells express *dsx* and undergo sex-specific differentiation (Goldman and

Arbeitman, 2007; Luo and Baker, 2015; Rideout et al., 2010; Robinett et al., 2010; Sanders and

Arbeitman, 2008; Tanaka et al., 2011). Integrating both activating and repressive inputs in *dsx*

enhancers may be necessary to segregate these cells from otherwise similar but sexually

monomorphic cells.

Modular dsx enhancers may allow different sex-specific traits to evolve independently

While the sex combs are present in only a subset of *Drosophila* species, particularly in the

melanogaster and *obscura* species groups, most though not all *Drosophila* species have sex-

specific chemosensory bristles on their forelegs. Thus, one could imagine a more ancestral

chemosensory foreleg enhancer becoming co-opted for sex comb development in the *melanogaster*

and *obscura* groups. A more extreme scenario would involve a co-option of a genital enhancer for

sex-specific foreleg development. Male genitalia and legs are serially homologous and share

similar developmental programs involving some of the same transcription factors and signaling

pathways (Gorfinkiel et al., 1999; Keisman et al., 2001; Sánchez et al., 1997), and the male

genitalia of most *Drosophila* species have strongly modified clasper bristles reminiscent of sex

comb teeth. Indeed, we find that Dsx is expressed in the clasper during late genital development.

However, our results argue against a simple co-option scenario. The chemosensory bristle

enhancer of *dsx* is clearly distinct from both enhancers that contribute to sex comb development,

being located in a different intron. The situation is less clear with the early foreleg enhancer, where

we cannot rule out a minor overlap with the genital enhancer. However, the late sex comb

enhancer, which appears to play the dominant role in the development of sex comb teeth, is clearly

separate from the genital enhancer.

Both the sex comb and the chemosensory bristles play important roles during courtship

(Hurtado-Gonzales et al., 2015; Mellert et al., 2012; Ng and Kopp, 2008) and have undergone

extensive changes in number, size, morphology, and spatial distribution in *Drosophila* evolution.

The ability of all three foreleg enhancers to function and evolve independently of each other may

have been a key element driving this evolutionary diversification. Analysis of these enhancers in

other *Drosophila* species will provide important insights into the role of *dsx* in the evolution of sex

combs and chemosensory bristles. The new GAL4 drivers that mark different types of sex-specific

sensory organs may also prove valuable for dissecting the neural mechanisms of courtship

behavior.

413   Modular transcriptional regulation of *dsx* may reflect an evolutionarily ancient mode of

414   sexual development.

Dsx-related transcription factors (the *Dmrt* gene family) play important roles in sexual

differentiation in animals as different as vertebrates, insects, and nematodes (Kopp, 2012; Matson

and Zarkower, 2012). Outside of insects, however, *Dmrt* genes do not produce distinct male and

female isoforms, but are regulated instead at the transcriptional level. Sex-specific splicing of *dsx*,

and more generally the mode of sexual differentiation based on alternative splicing, is an insect

innovation (Kopp, 2012; Wexler et al.). In crustaceans and chelicerates, *dsx* homologs are

transcribed exclusively or predominantly in males, particularly in male-specific organs, and are

required for male but not for female sexual development (Kato et al., 2011; Li et al., 2018;

Pomerantz and Hoy, 2015). In *C. elegans*, the *Dmrt* genes *mab3, mab23,* and *dmd3* are expressed

in tightly restricted cell lineages and are essential for male-specific differentiation of these cells,

but are dispensable in females (Lints and Emmons, 2002; Mason et al., 2008; Ross et al., 2005;

Serrano-Saiz et al., 2017; Yi et al., 2000). In vertebrates, the *Dmrt1* gene is highly expressed in

the male but not in the female gonad, and is necessary to direct or maintain male-specific

differentiation of the initially bipotential gonad (Matson et al., 2011).

These similarities suggest that sexual differentiation based on male-specific transcription

of *Dmrt* genes in restricted cell populations is an evolutionarily ancient mode of establishing sexual

dimorphism, and that in insects, sex-specific splicing of *dsx* was overlaid on this ancestral

mechanism (Kopp, 2012; Wexler et al.). The finding that *dsx* transcription in *Drosophila* is

controlled by multiple modular enhancers may reflect this more ancestral mode of sexual

development. We predict that a similar modular control of *dsx* expression will be found in other

arthropods, from basal insects and crustaceans to chelicerates.

Materials and Methods:

Fly stocks:

Fly strains were obtained from the Bloomington Stock Center, including UAS-GFP.nls (Bloomington stock #4775, genotype: w[1118]; P{w[+mC]=UAS-GFP.nls}14), UAS-svRNAi P{TRiP.JF02582}attP2 (Bloomington stock #27269, genotype: y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02582}attP2), and *dsx*-GAL4 (Robinett et al., 2010) (Bloomington stock #66674, genotype: w[1118]; TI{GAL4}dsx[GAL4]/TM6B, Tb[1])

Transgenic reporter constructs:

Candidate enhancer regions were cloned into the GAL4 reporter vector pPTGAL (Sharma et al., 2002), and randomly integrated into the *D. melanogaster* genome by P-element transformation. Three independent lines from each of ten overlapping fragments were examined. In addition, we analyzed 15 independently generated GAL4 lines which span the same non-coding regions of *dsx* but have different fragment boundaries (Pfeiffer et al., 2008). These lines were constructed using the site-specific pBPGUw vector (Pfeiffer et al., 2008) and integrated into either the attP2 or attP40 genomic landing sites via the PhiC31 integrase system (Groth et al., 2004; Venken et al., 2006). Each line from both sets was crossed to a UAS-GFP.nls reporter and GFP expression was examined in the developing foreleg at 5 and at 24 hours after puparium formation (APF).

While the high level of GFP expression driven by the GAL4/UAS system was useful for the preliminary screening of genomic fragments, subsequent analysis was performed using direct GFP reporters, which enable more accurate temporal resolution. Genomic fragments were amplified by PCR, cloned into either pCR2 or pCR8 vectors (Invitrogen), transferred into the pGreenFriend GFP destination vector (Miller et al., 2014) using either restriction/ ligation or Gateway cloning, and integrated into the attP2 or attP40 genomic landing sites using the PhiC31 integrase system. Germline transformation was performed either in the Kopp lab or by BestGene (http://www.thebestgene.com/). In tests performed on a subset of constructs, we found little difference between attP2 and attP40; we subsequently used the attP40 site, which gave higher integration efficiency. In designing the sub-fragments of the sex comb and early foreleg enhancers, we used Evo-printer (Odenwald et al., 2005) and VISTA (Frazer et al., 2004) to identify sequences within these enhancers that were conserved between *D. melanogaster* and *D. pseudoobscura*. Whenever possible, we avoided breaking up highly conserved regions. The

boundaries of all reporter fragments and the primers used to amplify them are listed in

Supplementary Tables 1 and 2.

Light Microscopy:

Legs were dissected from adult flies and mounted in PVA-Mounting-Medium (BioQuip) or

Hoyer's media to clear the samples. Images were taken under Brightfield illumination using a

Leica DM500B microscope with a Leica DC500 camera.

Immunohistochemistry and Confocal Microscopy:

Samples were aged, dissected, fixed, stained and imaged as described in Tanaka et al 2009 and

Mellert, et al 2012. The primary mouse-Dsx[DBD] antibody (a gift from C. Robinett and B.

Baker, later obtained from the Developmental Studies Hybridoma Bank, Cat# DsxDBD,

RRID:AB_2617197) was used at 1:10. The secondary Goat anti-Mouse Alexa Fluor 594

antibody (Invitrogen Cat# A20185) was used at 1:200. Confocal images were taken using an

Olympus FV1000 laser scanning confocal microscope.

Acknowledgments:

We thank Dr. Carmen Robinett, Dr. Bruce Baker, and the Developmental Studies Hybridoma

Bank for Dsx antibodies, Dr. John Yoder and the Bloomington Stock Center for *Drosophila*

strains, and the FlyLight Project for pBPGUw GAL4 lines. We want to thank Dr. John Yoder

and Dr. Carmen Robinett for discussions of *dsx* enhancer expression, and Dr. Kohtaro Tanaka

and Dr. Benjamin Vincent for comments on the manuscript.

Competing interests:

No competing interests declared

Funding:

This work was funded by NIH grants R01GM105726 and R35GM122592 to AK, by a

Collaborative Supplement to NIH grant R01GM073039 to AK and MNA, and by the NSF GRFP

and DDIG 1501148 to GR.

References:

Arbeitman, M. N., Fleming, A. A., Siegal, M. L., Null, B. H. and Baker, B. S. (2004). A

genomic analysis of Drosophila somatic sexual differentiation and its regulation.

Development **131**, 2007–2021.

Arbeitman, M. N., New, F. N., Fear, J. M., Howard, T. S., Dalton, J. E. and Graze, R. M.

(2016). Sex Differences in Drosophila Somatic Gene Expression: Variation and

Regulation by doublesex. *G3* **6**, 1799–1808.

Atallah, J., Liu, N. H., Dennis, P., Hon, A., Godt, D. and Larsen, E. W. (2009). Cell dynamics

and developmental bias in the ontogeny of a complex sexually dimorphic trait in

Drosophila melanogaster. *Evol. Dev.* **11**, 191–204.

Baker, B. S. and Ridge, K. A. (1980). Sex and the single cell. I. On the action of major loci

affecting sex determination in Drosophila melanogaster. *Genetics* **94**, 383–423.

Belote, J. M. and Baker, B. S. (1982). Sex determination in Drosophila melanogaster:

analysis of transformer-2, a sex-transforming locus. *Proc. Natl. Acad. Sci. U. S. A.* **79**,

1568–72.

Boggs, R. T., Gregor, P., Idriss, S., Belote, J. M. and McKeown, M. (1987). Regulation of

sexual differentiation in D. melanogaster via alternative splicing of RNA from the

transformer gene. *Cell* **50**, 739–747.

Burtis, K. C. and Baker, B. S. (1989). Drosophila doublesex gene controls somatic sexual

differentiation by producing alternatively spliced mRNAs encoding related sex-specific

polypeptides. *Cell* **56**, 997–1010.

Burtis, K. C., Coschigano, K. T., Baker, B. S. and Wensink, P. C. (1991). The doublesex

proteins of Drosophila melanogaster bind directly to a sex-specific yolk protein gene

enhancer. *EMBO J.* **10**, 2577–2582.

Butler, B., Pirrotta, V., Irminger-Finger, I. and Nöthiger, R. (2018). The sex-determining

gene tra of Drosophila: molecular cloning and transformation studies. *EMBO J.* **5**,

3607–3613.

Camara, N., Whitworth, C. and Doren, M. Van (2008). *The Creation of Sexual Dimorphism*
in the Drosophila Soma. Elsevier Inc.

Christiansen, A., Keisman, E., Ahmad, S. and Baker, B. (2002). Sex comes in from the
cold: the integration of sex and pattern. *Trends Genet.* **18**, 510–516.

Cline, T. W. (1993). The Drosophila sex determination signal: how do flies count to two?
Trends Genet. **9**, 385–390.

Coschigano, K. T. and Wensink, P. C. (1993). Sex-specific transcriptional regulation by the
male and female doublesex proteins of Drosophila. *Genes Dev.* **7**, 42–54.

Erickson, J. W. and Quintero, J. J. (2007). Indirect effects of ploidy suggest X chromosome
dose, not the X:A ratio, signals sex in Drosophila. *PLoS Biol.* **5**, 2821–2830.

Fan, P., Manoli, D. S., Ahmed, O. M., Chen, Y., Agarwal, N., Kwong, S., Cai, A. G., Neitz, J.,
Renslo, A., Baker, B. S., et al. (2013). Genetic and neural mechanisms that inhibit
drosophila from mating with other species. *Cell* **154**, 89–102.

Frankel, N., Davis, G. K., Vargas, D., Wang, S., Payre, F. and Stern, D. L. (2010).
Phenotypic robustness conferred by apparently redundant transcriptional enhancers.
Nature **466**, 490–493.

Frazer, K. A., Pachter, L., Poliakov, A., Rubin, E. M. and Dubchak, I. (2004). VISTA:
Computational tools for comparative genomics. *Nucleic Acids Res.* **32**, 273–279.

Garcia-Bellido, A. and de Celis, J. F. (2009). The Complex Tale of the achaete–scute
Complex: A Paradigmatic Case in the Analysis of Gene Organization and Function
During Development. *Genetics* **182**, 631–639.

Glassford, W. J., Johnson, W. C., Dall, N. R., Smith, S. J., Liu, Y., Boll, W., Noll, M. and
Rebeiz, M. (2015). Co-option of an Ancestral Hox-Regulated Network Underlies a
Recently Evolved Morphological Novelty Article. *Dev. Cell* **34**, 520–531.

Goldman, T. D. and Arbeitman, M. N. (2007). Genomic and Functional Studies of
Drosophila Sex Hierarchy Regulated Gene Expression in Adult Head and Nervous

System Tissues. *PLoS Genet.* **3**, e216.

Gorfinkiel, N., Sánchez, L. and Guerrero, I. (1999). Drosophila terminalia as an appendage-like structure. *Mech. Dev.* **86**, 113–123.

Grieder, N. C., Marty, T., Ryoo, H. D., Mann, R. S. and Affolter, M. (1997). Synergistic activation of a Drosophila enhancer by HOM/EXD and DPP signaling. *EMBO J.* **16**, 7402–7410.

Groth, A. C., Fish, M., Nusse, R. and Calos, M. (2004). Construction of Transgenic Drosophila by Using the Site-Specific Integrase From Phage φC31. *Genetics* **166**, 1775–1782.

Held, L. I. (2002). *Imaginal Discs: The Genetic and Cellular Logic of Pattern Formation*.

Hildreth, P. E. (1965). Doublesex, Recessive Gene That Transforms Both Males and Females of Drosophila Into Intersexes. *Genetics* **51**, 659–678.

Ho, E. C. Y., Malagón, J. N., Ahuja, A., Singh, R. and Larsen, E. (2018). Rotation of sex combs in Drosophila melanogaster requires precise and coordinated spatio-temporal dynamics from forces generated by epithelial cells. *PLoS Comput. Biol.* **14**, 1–28.

Hong, J. W., Hendrix, D. A. and Levine, M. S. (2008). Shadow enhancers as a source of evolutionary novelty. *Science (80-.).* **321**, 1314.

Hurtado-Gonzales, J., Gallaher, W., Warner, A. and Polak, M. (2015). Microscale laser surgery demonstrates the grasping function of the male sex combs in drosophila melanogaster and drosophila bipectinata. *Ethology* **121**, 45–56.

Kato, Y., Kobayashi, K., Watanabe, H. and Iguchi, T. (2011). Environmental sex determination in the branchiopod crustacean Daphnia magna: Deep conservation of a Doublesex gene in the sex-determining pathway. *PLoS Genet.* **7**, 1–12.

Kavaler, J., Fu, W., Duan, H., Noll, M. and Posakony, J. W. (1999). An essential role for the Drosophila Pax2 homolog in the differentiation of adult sensory organs. *Development* **126**, 2261–2272.

Keisman, E. L., Christiansen, A. E. and Baker, B. S. (2001). The Sex Determination Gene

doublesex Regulates the A/P Organizer to Direct Sex-Specific Patterns of Growth in the

Drosophila Genital Imaginai Disc. *Dev. Cell* **1**, 215–225.

Kopp, A. (2011). Drosophila sex combs as a model of evolutionary innovations. *Evol. Dev.*

13, 504–522.

Kopp, A. (2012). Dmrt genes in the development and evolution of sexual dimorphism.

Trends Genet. **28**, 175–84.

Lebo, M. S., Sanders, L. E., Sun, F. and Arbeitman, M. N. (2009). Somatic, germline and

sex hierarchy regulated gene expression during Drosophila metamorphosis. *BMC*

Genomics **10**, 1–23.

Li, H. and Baker, B. S. (1998). hermaphrodite and doublesex function both dependently

and independently to control various aspects of sexual differentiation in Drosophila.

Development **125**, 2641–2651.

Li, S., Li, F., Yu, K. and Xiang, J. (2018). Identification and characterization of a doublesex

gene which regulates the expression of insulin-like androgenic gland hormone in

Fenneropenaeus chinensis. *Gene* **649**, 1–7.

Lints, R. and Emmons, S. W. (2002). Regulation of sex-specific differentiation and mating

behavior in C. elegans by a new member of the DM domain transcription factor family.

Genes Dev. **16**, 2390–2402.

Luo, S. D. and Baker, B. S. (2015). Constraints on the evolution of a *doublesex* target gene

arising from doublesex's pleiotropic deployment. *Proc. Natl. Acad. Sci.* **112**, E852–

E861.

Malagón, J. N., Ahuja, A., Sivapatham, G., Hung, J. and Lee, J. (2014). Evolution of

Drosophila sex comb length illustrates the inextricable interplay between selection

and variation. *Proc. Natl. Acad. Sci.* **111**, E4103–E4109.

Manak, J. R., Mathies, L. D. and Scott, M. P. (1994). Regulation of a decapentaplegic

midgut enhancer by homeotic proteins. *Development* **120**, 3605–19.

Mason, D. A., Rabinowitz, J. S. and Portman, D. S. (2008). dmd-3, a doublesex-related

gene regulated by tra-1, governs sex-specific morphogenesis in C. elegans.

Development **135**, 2373–2382.

Matson, C. K. and Zarkower, D. (2012). Sex and the singular DM domain: insights into

sexual regulation, evolution and plasticity. *Nat. Rev. Genet.* **13**, 163–74.

Matson, C. K., Murphy, M. W., Sarver, A. L., Griswold, M. D., Bardwell, V. J. and

Zarkower, D. (2011). DMRT1 prevents female reprogramming in the postnatal

mammalian testis. *Nature* **476**, 101–104.

McKeown, M. (1992). Sex differentiation: The role of alternative splicing. *Curr. Opin. Genet.*

Dev. **2**, 299–303.

McKeown, M., Belote, J. M. and Baker, B. S. (1987). A molecular analysis of transformer, a

gene in Drosophila melanogaster that controls female sexual differentiation. *Cell* **48**,

489–499.

McKeown, M., Belote, J. M. and Boggs, R. T. (1988). Ectopic expression of the female

transformer gene product leads to female differentiation of chromosomally male

drosophila. *Cell* **53**, 887–895.

Mellert, D. J., Robinett, C. C. and Baker, B. S. (2012). Doublesex Functions Early and Late

in Gustatory Sense Organ Development. *PLoS One* **7**,.

Miller, S. W., Rebeiz, M., Atanasov, J. E. and Posakony, J. W. (2014). Neural precursor-

specific expression of multiple *Drosophila* genes is driven by dual enhancer modules

with overlapping function. *Proc. Natl. Acad. Sci.* **111**, 17194–17199.

Nagoshi, R. N., McKeown, M., Burtis, K. C., Belote, J. M. and Baker, B. S. (1988). The

control of alternative splicing at genes regulating sexual differentiation in D.

melanogaster. *Cell* **53**, 229–236.

Nagy, O., Nuez, I., Savisaar, R., Peluffo, A. E., Yassin, A., Lang, M., Stern, D. L., Matute, D.

R., David, J. R. and Courtier-Orgogozo, V. (2018). Correlated Evolution of Two

Copulatory Organs via a Single cis-Regulatory Nucleotide Change. *Curr. Biol.* **28**, 1–8.

Ng, C. S. and Kopp, A. (2008). Sex combs are important for male mating success in

Drosophila melanogaster. *Behav. Genet.* **38**, 195–201.

Odenwald, W. F., Rasband, W., Kuzin, A. and Brody, T. (2005). EVOPRINTER, a

multigenomic comparative tool for rapid identification of functionally important DNA.

Proc. Natl. Acad. Sci. U. S. A. **102**, 14700–5.

Pfeiffer, B. D., Jenett, A., Hammonds, A. S., Ngo, T.-T. B., Misra, S., Murphy, C., Scully, A.,

Carlson, J. W., Wan, K. H., Laverty, T. R., et al. (2008). Tools for neuroanatomy and

neurogenetics in Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 9715–20.

Pomerantz, A. F. and Hoy, M. A. (2015). Expression analysis of Drosophila doublesex,

transformer-2, intersex, fruitless-like, and vitellogenin homologs in the parahaploid

predator Metaseiulus occidentalis (Chelicerata: Acari: Phytoseiidae). *Exp. Appl. Acarol.*

65, 1–16.

Preger-Ben Noon, E., Sabarís, G., Ortiz, D. M., Sager, J., Liebowitz, A., Stern, D. L. and

Frankel, N. (2018). Comprehensive Analysis of a cis-Regulatory Region Reveals

Pleiotropy in Enhancer Function. *Cell Rep.* **22**, 2809–2817.

Rebeiz, M., Pool, J. E., Kassner, V. A., Aquadro, C. F. and Carroll, S. B. (2009). Stepwise

modification of a modular enhancer underlies adaptation in a Drosophila population.

Science **326**, 1663–1667.

Rice, G. and Rebeiz, M. (2019). How Many Phenotypes Do Regulatory Mutations Affect ?

Curr. Biol. **29**, R21–R23.

Rideout, E. J., Dornan, A. J., Neville, M. C., Eadie, S. and Goodwin, S. F. (2010). Control of

sexual differentiation and behavior by the doublesex gene in Drosophila melanogaster.

Nat. Neurosci. **13**, 458–66.

Robinett, C. C., Vaughan, A. G., Knapp, J. M. and Baker, B. S. (2010). Sex and the single

cell. II. There is a time and place for sex. *PLoS Biol* **8**, e1000365.

Ross, J. M., Kalis, A. K., Murphy, M. W. and Zarkower, D. (2005). The DM domain protein

MAB-3 promotes sex-specific neurogenesis in C. elegans by regulating bHLH proteins.

Dev. Cell **8**, 881–892.

Sánchez, L., Casares, F., Gorfinkiel, N. and Guerrero, I. (1997). The genital disc of

Drosophila melanogaster. II. Role of the genes hedgehog, decapentaplegic and

wingless. *Dev. Genes Evol.* **207**, 229–241.

Sanders, L. E. and Arbeitman, M. N. (2008). Doublesex establishes sexual dimorphism in

the Drosophila central nervous system in an isoform-dependent manner by directing

cell number. *Dev. Biol.* **320**, 378–390.

Serrano-Saiz, E., Oren-Suissa, M., Bayer, E. A. and Hobert, O. (2017). Sexually Dimorphic

Differentiation of a C. elegans Hub Neuron Is Cell Autonomously Controlled by a

Conserved Transcription Factor. *Curr. Biol.* **27**, 199–209.

Sharma, Y., Cheung, U., Larsen, E. W. and Eberl, D. F. (2002). PPTGAL, a convenient Gal4

P-element vector for testing expression of enhancer fragments in drosophila. *Genesis*

34, 115–118.

Shirangi, T. R., Dufour, H. D., Williams, T. M. and Carroll, S. B. (2009). Rapid Evolution of

Sex Pheromone-Producing Enzyme Expression in Drosophila. *PLoS Biol.* **7**, e1000168.

Spieth, H. T. (1952). Mating behavior within the genus *Drosophila (Diptera)*. *Bull. AMNH*

99, 395–474.

Swanson, C., Evans, N. and Barolo, S. (2010). Structural Rules and Complex Regulatory

Circuitry Constrain Expression of a Notch- and EGFR-Regulated Eye Enhancer. *Dev.*

Cell **18**, 359–376.

Tanaka, K., Barmina, O. and Kopp, A. (2009). Distinct developmental mechanisms

underlie the evolutionary diversification of Drosophila sex combs. *Proc. Natl. Acad. Sci.*

106, 4764–4769.

Tanaka, K., Barmina, O., Sanders, L. E., Arbeitman, M. N. and Kopp, A. (2011). Evolution

of sex-specific traits through changes in HOX-dependent doublesex expression. *PLoS*

Biol. **9**, e1001131.

Tokunaga, C. (1962). Cell lineage and differentiation on the male foreleg of Drosophila

melanogaster. *Dev. Biol.* **4**, 489–516.

Venken, K. J. T., He, Y., Hoskins, R. a and Bellen, H. J. (2006). P[acman]: A BAC

Transgenic Platform Fragments in D . melanogaster. *Science (80-.).* **314**, 1747–1751.

Wexler, J., Delaney, E. K., Belles, X., Schal, C., Wada-Katsumata, A., Amicucci, M. and

Kopp, A. Hemimetabolous insects elucidate the origin of sexual development via

alternative splicing. *Submitted*.

Williams, T. M., Selegue, J. E., Werner, T., Gompel, N., Kopp, A. and Carroll, S. B. (2008).

The regulation and evolution of a genetic switch controlling sexually dimorphic traits

in Drosophila. *Cell* **134**, 610–23.

Yang, Y., Zhang, W., Bayrer, J. R. and Weiss, M. A. (2008). Doublesex and the Regulation

of Sexual Dimorphism in Drosophila melanogaster. *J. Biol. Chem.* **283**, 7280–7292.

Yi, W., Ross, J. M. and Zarkower, D. (2000). mab-3 is a direct tra-1 target gene regulating

diverse aspects of C. elegans male sexual development and behavior. *Development*

127, 4469–4480.

Zhou, C., Pan, Y., Robinett, C. C., Meissner, G. W. and Baker, B. S. (2014). Article Central

Brain Neurons Expressing doublesex Regulate Female Receptivity in Drosophila.

Neuron **83**, 149–163.

     Figures:

Figure 1: Three modular enhancers drive *doublesex* expression in the foreleg

A) A map of the *doublesex* locus with the positions of chemosensory (blue), sex comb (red), and
foreleg (purple) *dsx* enhancers. Thick grey lines represent the flanking genes *lds* and *CD98hc*,
thick black lines are the exons of *dsx*, and thin black lines are the *dsx* introns. Panels B – G show
the foreleg expression patterns of the three *dsx* enhancers, with enhancer-driven GFP in green
and magnifications of 28hr samples of the early foreleg and sex comb enhancers show Dsx

antibody staining in red. Hours are after puparium formation (APF). The early foreleg and sex

comb enhancers constructs are directly adjacent to GFP (Panels B-E), while the chemosensory

enhancer (42C06 (Pfeiffer et al., 2008)) construct is a GAL4 driver crossed to a UAS-GFP.nls

line. **B)** At 5 hours APF, the early foreleg enhancer recapitulates the Dsx expression pattern in

the leg epithelium (Tanaka et al., 2011) and shows a similar expression pattern in males (left)

and females (right). Both sexes show ectopic expression in the proximal first tarsal segment

(ta1). **C)** At 28 hrs APF, the early foreleg enhancer shows clear sexual dimorphism in the first

tarsal segment. Males show strong expression in the epithelial cells surrounding the sex comb

and weak to no expression in the sex comb bristle cells. Females show little to no expression in

the distal ta1. Both males and females show ectopic expression in the proximal ta1, in the same

region as at 5 hrs APF (B), that is not seen by Dsx antibody staining (Tanaka et al., 2011). **D)**

The sex comb enhancer is not active in either sex at 5 hrs APF, except for a proximal patch of

ectopic expression seen near the joint in some individuals **E)** At 28 hrs APF, the sex comb

enhancer is active in the bristle cells of the male sex comb (large nuclei), and weakly in the

epithelial cells ventral to the sex comb. Females show weak expression in the distal portion of

the segment. Both sexes show ectopic expression in the joint between the tibia and ta1. **F)** At 5

hrs APF, the chemosensory enhancer is active in small clusters of cells in ta1-ta5 in both sexes,

with more GFP-positive cells in males than in females. **G)** At 28 hrs APF, both sexes show small

clusters of expression in ta1-ta5; only ta1 is shown. (No Dsx antibody staining was performed in

F-G).

Figure 2: All three leg enhancers contribute to sex-specific bristle development

For each enhancer, a GAL4 reporter was used to drive a UAS-*shaven* RNAi construct, truncating

the bristles where that enhancer is active. Images of male ta1 are shown next to a schematic

representation of male and female ta1 chaetotaxy modified from (Tokunaga 1962). Circles

designate mechanosensory bristles, while triangles designate chemosensory bristles. Dark

orange symbols mark bristles that were affected in all individuals; bristles marked with yellow

varied among individuals of the same genotype raised under standard conditions. **A)** A GAL4

knock-in in the *dsx* gene, which reflects the full expression pattern of *dsx* (Robinett et al., 2010),

affects both sex comb and sexually dimorphic chemosensory bristles in males. **B)** In females, the

dsx-GAL4 knock-in has a variable effect on the distal Transverse Bristle Row (TBR), which is

homologous to the sex comb, and on some chemosensory bristles. **C)** In males, the early foreleg

enhancer affects sex comb bristles in all flies as well as a single chemosensory bristle in some

individuals. **D)** In females, the early foreleg enhancer affects one bristle of the distal TBR and

the same chemosensory bristle as in males. **E)** In males, the sex comb enhancer affects only the

sex comb. **F)** The sex comb enhancer does not affect any bristles in females. **G)** In males, the

chemosensory enhancer affects the same chemosensory bristles as the *dsx*-GAL4 knock-in but

has no effect on the sex comb. **H)** The chemosensory enhancer does not affect any bristles in

females.

Figure 3: The foreleg and sex comb enhancers are separate and modular

To test for overlap between the foreleg, genital, and sex comb enhancer activities, the region they encompass was split into six constructs: full (all three enhancers, green), genital (black), foreleg (purple), foreleg + genital (purple + black), sex comb (red), and sex comb + genital (red + black). In the drawings, green shading shows reporter expression that matches endogenous Dsx

expression, with light green indicating weak expression and white a lack of expression.

Representative images of each construct are shown in Fig. S3. Only the constructs that include

the early foreleg enhancer drive expression in the forelegs at 5 hrs APF and in the epithelial cells

surrounding the sex comb at 24 hrs. Only regions that include the sex comb enhancer drive

expression in the sex comb bristles at 24 hrs APF. Both the genital and the early foreleg enhancer

constructs are able to drive expression in the genitalia at 48 hrs. ta1-ta4: tarsal segments 1-4; h:

hypandrium; a: aedeagus; p: dorsal postgonites; s: aedeagal sheath.

GFP

Figure 4: Repressor elements are necessary to limit *dsx* expression to sexually dimorphic cells.

A) A map of the sex comb (red), genital (black), and foreleg (purple) enhancers. **B-E)** Representative images of male forelegs at 36 hrs APF for the *sex comb sub1-sub5* constructs; only the ta1 segment is shown. **B)** The *sex comb sub1* enhancer drives expression in the sex comb bristle cells; **C)** *sex comb sub2* is active in sex comb bristles but has additional ectopic expression in the central bristle and in the second most distal TBR; **D)** *sex comb sub3* is expressed in all TBRs bristle cells in the ta1 segment; **E)** *sex comb sub4* is expressed in some of

the epithelial cells that surround the sex comb, but these cells are not directly adjacent to the sex

comb bristles; **F)** *sex comb sub5* at 24 hrs AFP is expressed in the epithelial cells surrounding the

sex comb, but also shows ectopic expression in more proximal epithelial cells **G-H)** Foreleg

expression of the *foreleg sub1* and *sub2* constructs at 5 hrs APF. The *foreleg sub1* region drives

broad ectopic expression throughout the ta1-ta4; *foreleg sub2* shows expression in a pattern

similar to the complete early foreleg enhancer (Fig. 1B).