A peptide of a type I toxin-antitoxin system induces *Helicobacter pylori* morphological

transformation from spiral-shape to coccoids

Short title: *H. pylori* shape conversion induced by a TA toxin

Lamya El Mortaji [1], Stéphanie Marsin [2+], Aline Rifflet [3,4], Ivo G. Boneca [3,4], J. Pablo

Radicella [2], Gérard Péhau-Arnaudet [5] and Hilde De Reuse [1] *

[1] Institut Pasteur, Département de Microbiologie, Unité Pathogenèse de *Helicobacter*, ERL

CNRS 6002, 28 rue du Dr Roux 75724 PARIS Cedex 15 FRANCE.

[2] CEA, Institute of Molecular and Cellular Radiobiology, Institut François Jacob, INSERM

UMR967, Universités Paris Diderot et Paris Sud, Fontenay-aux-Roses, France.

+ Current affiliation: I2BC/UMR 9198 CNRS- CEA -Université Paris Sud, Bat. 430, Orsay,

FRANCE.

[3] Institut Pasteur, Département de Microbiologie, Unité Biologie et Génétique de la Paroi

Bactérienne, 75724 PARIS, FRANCE.

[4] INSERM, Equipe Avenir, 75015 Paris, France

[5] Institut Pasteur, Ultrastructural BioImaging (*UTechS* UBI) /UMR CNRS 3528, PARIS,

FRANCE.

* -Corresponding author: Hilde De Reuse (hdereuse@pasteur.fr)

Institut Pasteur, Unité Pathogenèse de *Helicobacter*

28, rue du Docteur Roux

75724 PARIS Cedex 15 FRANCE

Tel : +33 1 40 61 36 41

Summary

Toxin-antitoxin systems are found in many bacterial chromosomes and plasmids with roles ranging from plasmid stabilization to biofilm formation and persistence. In these systems, the expression/activity of the toxin is counteracted by an antitoxin, which in type I systems is an antisense-RNA. While the regulatory mechanisms of these systems are mostly well-defined, the toxins' biological activity and expression conditions are less understood. Here, these questions were investigated with a type I toxin-antitoxin system (AapA1-IsoA1) expressed from the chromosome of the major human pathogen *Helicobacter pylori*. We show that expression of the AapA1 toxin in *H. pylori* causes growth arrest associated with massive morphological transformation from spiral-shaped bacteria to round coccoid cells. Coccoids are observed in patients and during *in vitro* growth as a response to different conditions such as oxidative stress. The AapA1 toxin, first molecular effector of coccoids to be identified, targets *H. pylori* inner membrane without disruption, as visualized by Cryo-EM. The peptidoglycan composition of coccoids is modified as compared to spiral bacteria. No major changes in membrane potential or ATP concentration result from AapA1 expression, suggesting coccoid viability. Using single-cell live microscopy, we observed that shape conversion is associated with cell division interference. Oxidative stress represses antitoxin promoter activity and enhances processing of its transcript leading to an imbalanced ratio in favor of AapA1 toxin expression.

Our data are in favor of viable coccoids with characteristics of dormant bacteria that might be important in *H. pylori* infections refractory to treatment.

Significance Statement

Helicobacter pylori, a gastric pathogen responsible for 800,000 deaths in the world every

year, is encountered, both *in vitro* and in patients, as spiral-shaped bacteria and as round cells

named coccoids. We discovered that the toxin from a chromosomal type I toxin-antitoxin

system is targeting *H. pylori* membrane and acting as an effector of the morphological

conversion of *H. pylori* to coccoids. We showed that these round cells maintain their

membrane integrity and metabolism, strongly suggesting that they are viable dormant bacteria.

Oxidative stress was identified as a signal inducing toxin expression. Our findings reveal new

insights into a form of dormancy of this bacterium that might be associated with *H. pylori*

infections refractory to treatment.

Introduction

Toxin-antitoxin (TA) systems are small genetic elements that are widely distributed on bacterial mobile genetic elements and among archaeal and bacterial genomes, often in multiple copies (1-3). They code for a small stable toxic protein, designated toxin, whose action or expression is counteracted by an unstable antitoxin molecule that can either be an RNA or a protein (3, 4). Under conditions in which the toxins can act, they target essential cellular processes or components (transcription, DNA replication, translation, cell wall or membrane) resulting in growth arrest or cell death (3, 4). The molecular mechanism of the toxins' activities and their regulation are often known in detail while the biological function remains elusive. TA systems take part in plasmid maintenance and protection from phage infection. Less is known on the biological functions and targets of the chromosomally encoded TA systems. Some are important for bacterial survival in their mammalian host (5) or biofilm formation (6). There is accumulating evidence that upon stress conditions, some TA systems play a role in the switch of actively growing bacteria to persisters or dormant cells (7, 8). Persister bacteria constitute a subpopulation that is metabolically active but slow-growing and highly tolerant to antibiotics and stress conditions (9). Persistence is considered as an important cause of recalcitrance of chronic bacterial infections to therapy. However, stress-induced activation of TA-encoded toxins does not necessarily cause persister formation (10) and it was more generally found that decreased intracellular ATP concentration is a landmark of persister formation (11).

In the present work, we explored the role of TA systems in *Helicobacter pylori*, a bacterium that colonizes the stomach of half of the human population worldwide and causes the development of gastritis. In some cases, gastritis evolves into peptic ulcer disease or gastric carcinoma that causes about 800,000 deaths in the world every year (12, 13). This microaerophilic bacterium is unique in its capacity to persistently colonize the stomach despite its extreme acidity and intense immune response (13). The molecular mechanisms at

the origin of this exceptional adaptation capacity of *H. pylori* remain only partially

understood and their elucidation is crucial to understand its virulence and to improve its

eradication in patients with recurrent peptic ulcer disease. *H. pylori* is a Gram-negative

bacterium with a helical shape. Upon stress conditions (antibiotics, aerobic growth) or

prolonged culture, the shape of *H. pylori* progressively evolves into a U-shape followed by a

spherical form designated coccoid (14, 15). *H. pylori* coccoids are non-culturable bacteria

proposed to be dormant forms (15). Recently, dormant non culturable cells were associated

wit a deeper state of dormancy when compared to persister cells (16)

Coccoids were observed in human gastric biopsies and like spirals adhere to gastric

epithelial cells. Despite numerous reports on coccoid forms of *H. pylori*, no cellular effector

of this conversion has been reported so far.

In type I TAs, the antitoxin is a small regulatory RNA inhibiting the synthesis of the toxin

by base pairing the toxin-encoding mRNA (3, 4, 17, 18). Four families (A-B-C-D) of

conserved type I TA systems are expressed from the chromosome of *H. pylori,* only the A

family was studied as it is highly expressed and conserved among *H. pylori* strains (19, 20).

For the A1 TA system, the detailed mechanism by which transcription of the IsoA1 antitoxin

RNA impairs AapA1 toxin synthesis by base-pairing with its primary transcript, ensuring

both translation inhibition of the AapA1 active message and leading to rapid degradation of

the duplex by RNase III has been recently published (20). The *H. pylori* type I toxins are

typically small hydrophobic peptides of 30-40 amino acids predicted to form alpha-helices.

No clues on the mode of action or the physiological role of these systems have been reported.

Here we show that the AapA1 toxin induces a rapid and massive morphological

transformation of *H. pylori* from spirals to coccoids by targeting the inner membrane and

interfering with cell division, and that oxidative stress triggers imbalanced expression of the

TA components in favor of toxin production.

Results

The AapA1 toxic peptide, expressed by the AapA1/IsoA1 TA system, triggers rapid transformation of *H. pylori* into coccoids

Recently, the study of the AapA1/IsoA1 TA system of *H. pylori* (**Fig. 1A**) showed that expression of the AapA1 toxin (a 30 amino acids-long hydrophobic peptide) leads to bacterial growth arrest (20). Given the genetic organization of the type I AapA1/IsoA1 locus in B128 strain, we decided to investigate the mechanism underlying the activity of AapA1 by using a strain in which AapA1 is under the control of an inducible promoter. We thus transformed *H. pylori* strain B128 deleted of the AapA1/IsoA1 locus with each of the three plasmids constructed by Arnion *et al.* (20) (**Fig. 1A**). The first, pA1-IsoA1, derived of the pILL2157 vector (21) contains a functional TA locus in which the AapA1 gene is under the control of an IPTG inducible promoter (20). In plasmid pA1, derived from this first construct, the antitoxin IsoA1 promoter has been inactivated by point mutations that do not interfere with transcription or translation of the AapA1 ORF (20). Derived from this second construct, plasmid pA1* contains an additional mutation that inactivates the start codon of the AapA1 peptide (20). Under our experimental conditions, we observed, in agreement with data of (20), that addition of IPTG did not significantly influence the growth rate of strains harboring pA1-IsoA1 or pA1*. In contrast, addition of the inducer causes a rapid and immediate growth arrest of *H. pylori* with pA1 indicating a toxic effect of AapA1 expression (**Fig. 1B**). The growth arrest was accompanied by loss of culturability of more than 10^4-fold 8h after induction.

In parallel to growth, we investigated the consequences of AapA1 expression on *H. pylori* morphology. Samples from bacterial cultures with these plasmids were grown in the presence or absence of IPTG, stained with a membrane-specific dye (TMA-DPH) and analyzed by fluorescence microscopy at different time points (**Fig. 1C**). *H. pylori* cells expressing the wild-type AapA1/IsoA1 locus (pA1-IsoA1) or the pA* mutated locus present a classical helical rod shape phenotype upon IPTG exposure (**Fig. 1C**). Under the same conditions, cells

carrying the pA1 plasmid, and thus expressing the AapA1 toxin exhibit a rapid and dramatic

morphological conversion to spherical cells also known as coccoid forms. Eight hours after

IPTG-induction, some bacteria containing pA1-IsoA1 start to convert to coccoids, this is

probably due to accumulation of the AapA1 toxin under these conditions. No bacterial lysis

was detected even at 24h post-induction. With strains carrying the pA1* plasmid, coccoids

were only observed after 48h of culture, similarly to the control wild type strain containing an

empty plasmid. These data show for the first time that the AapA1 toxin induces a fast

conversion of *H. pylori* cells from spiral-shaped to coccoid forms.

Inner membrane targeting by the AapA1 toxin

To investigate the mode of action of the AapA1 toxin, we first analyzed its subcellular

localization in *H. pylori* using three reporter strains. Two constructs corresponded to C-

terminal fusions of the toxin with a SPA-tag (pA1-SPA) (20) or with GFP (pA1-GFP)

expressed from the IPTG-inducible promoter of vector pILL2157 (21) (**Fig. S1**). These

plasmids were introduced into *H. pylori* strain B128 deleted of the chromosomal

AapA1/IsoA1 module. After IPTG induction none of these fusions affected *H. pylori* growth

indicating that a C-terminal fusion prevents the toxicity of the toxin. Attempts to construct N-

terminal tagged-AapA1 were unsuccessful. In addition, a B128 derivative expressing a

chromosomal AapA1-GFP fusion under control of its native promoter was constructed (**Fig.**

S1).

Cellular fractionation was then performed on the three strains, and the subcellular

localization of the fused toxins was analyzed by immunoblotting. **Figure 2A** shows that upon

induction by IPTG, the large majority of the AapA1-SPA is present at the inner membrane of

H. pylori with no peptide detected in the culture medium. Identical results were obtained with

the A1-GFP fusion expressed either from the inducible plasmid or the chromosome (**Fig. S2**).

Fluorescence microscopy of live bacteria after induction of the AapA1-GFP fusion revealed a

weak patchy pattern in the periphery of the cell (**Fig. 2B**), different from a GFP-alone control

cell (**Fig. S2**), and compatible with membrane localization. This observation was confirmed

by measuring the overlap of the fluorescence intensity profiles of the toxin and TMA-PH

measured perpendicular to the length axis of *H. pylori* (**Fig. 2B**).

From these results, we conclude that the AapA1 toxin is specifically targeted to the inner

membrane of *H. pylori*.

AapA1 toxin has a minor impact on *H. pylori* membrane potential and intracellular ATP

content

Considering the membrane localization of the toxin, we hypothesized that AapA1

expression could affect the membrane potential and by consequence the ATP content of the

cells. To explore this possibility, *H. pylori* cells harboring pA1-IsoA1, pA1* or pA1 plasmids

were taken at different culture time points, stained with a membrane potential sensitive dye

(MitoRed CMXROS) and analyzed by microscopy (**Fig. 2C**). Bacteria harboring either an

empty plasmid (B128), pA1-IsoA1 or pA1* exhibit a uniform MitoRed CMXROS

fluorescence pattern indicative of membrane integrity. For *H. pylori* cells overexpressing

AapA1 and converting to coccoids, the fluorescence pattern progressively evolves from

uniform to that of discrete foci, suggesting local changes in membrane permeability. No such

foci were observed with coccoids from aging WT strain cultures (72h) or with cells treated

with 3,5,3',4'-tetrachlorosalicylanilide (TCS), an effective *H. pylori* protonophore (22) that

causes an overall MitoRed CMXROS fluorescence loss due to massive dissipation of

membrane potential (**Fig. 2C**). Similar results were obtained with the membrane potential

sensitive dye DIOC-5, confirming the local effect of AapA1 on *H. pylori* inner membrane

(**Fig. S3**). These data indicate that when the AapA1 toxin is expressed in *H. pylori,* the

membrane permeability is not globally compromised but probably locally perturbed.

Next, the ATP content of *H. pylori* strains carrying either pA1-IsoA1 or pA1 was

measured from total metabolites extracted at different culture time points and quantification

by a luciferase based-assay (**Fig. 2D**). After 6h of culture, IPTG induction had a minor effect

on cellular ATP content of both strains as compared to the drastic consequences observed

with the TCS control. Note that at 6h of culture with IPTG, most of the pA1-containing cells

have already transformed into coccoids. After 16h, the overall ATP content strongly drops in

both strains and IPTG induction causes an additional weak decrease that is poorly changed

upon toxin expression by pA1 (**Fig. 2D).**

Taken together, these results show that the AapA1 toxin locally perturbs the *H. pylori*

membrane potential with no major consequences on the cellular ATP content. We conclude

that dissipation of membrane potential is not the major cause of toxin-induced bacterial

growth arrest.

Ultrastructural analysis of toxin-induced coccoids

For the first time, cryo-electron microscopy (cryo-EM) was used to visualize *H. pylori*

coccoids in near-native states. Using this method, we compared exponentially growing

bacteria, toxin-induced coccoids and aging coccoids (70h growth) (**Fig. 3**). In exponential

phase, the characteristic helicoidal shape of *H. pylori* (strains B128 WT, 4h and pA1-IsoA1,

4-8h) was perfectly visible with a uniformly contrasted cytoplasm surrounded by two dense

layers of membranes and multiple flagella. The periplasm of variable thickness surrounding

the cell is distinguishable as the low-density space between the inner and outer membrane.

Strains expressing the toxin (pA1 at 4-8h) have intact flagella and are visible in two major

morphotypes. In the first "U" shaped morphology, a round intact outer membrane surrounds a

bent intact inner membrane. The second morphotype corresponds to round coccoids of a

diameter of approximately 1µm, with visible integrity of both membranes and a central

uniformly stained dense cytoplasm. This second morphotype is similar to that of aging

coccoids from the WT strain (70h).

In conclusion, CryoEM images revealed that the toxin-induced *H. pylori* coccoids have no

visible defect in membrane integrity and are ultrastructurally similar to aging coccoid forms.

Modification of peptidoglycan composition upon transformation of *H. pylori* into

coccoids

Our data showed that the AapA1 toxin targets *H. pylori* inner membrane without causing

significant membrane potential collapse and without visible loss of integrity. Therefore, we

tested whether the toxin-induced cell shape transformation could be associated with

modifications in the cell wall and thus in peptidoglycan (PG) composition. PG extracted from

B128 WT strain in exponential phase or after 72h growth (aging coccoids) was compared to

PG from the pA1 strain induced during 6 and 16h (toxin-induced coccoids). Samples were

digested with mutanolysin and subjected to HPLC/MS analyses. The relative abundance of

muropeptides in each sample was calculated according to Glauner *et al.* (23) (**Table S1**).

When compared to PG of helicoidal *H. pylori,* both "aging" and "toxin-induced" coccoids

present similar changes in dipeptide and tripeptide monomers with a stronger effect of toxin-

induction at 16h. Coccoids after 16h induction present a 3-fold increase in dipeptide

monomers concomitant with tri- and pentapeptide monomers reduction. These data indicate

that toxin-induced and aging coccoids undergo similar PG modifications, compatible with a

looser PG macromolecule, and suggest that they might be generated by similar mechanisms.

Kinetics of the toxin-induced *H. pylori* morphological transformation assessed by live

cell imaging

To progress in our understanding of the morphological conversion of *H. pylori*, we

monitored the entire process of *H. pylori* toxin-induced conversion by live cell time-lapse

microscopy under physiological microaerobic conditions in a temperature-controlled chamber. The *H. pylori* B128 strain tested constitutively expressed cytoplasmic GFP, was impaired in motility (by deletion of *flaA* encoding the major flagellar protein) and carried the inducible toxin-expressing plasmid pA1. Individual live *H. pylori* bacteria were monitored in agarose pad by both phase contrast and fluorescence microscopy.

First, the general growth features of the strain were established in the absence of the IPTG inducer. Under these conditions, statistics of the data collected showed that the strain has a mean doubling time of 165 min (indicative of optimal growth conditions) with an initial bacterial length of 2.1μm after division and a maximum length prior to division of 3.7μm (**Fig. S4**). In a second round of experiments, the inducer was added to the pad and snapshots of the cells were taken at intervals of 10 or 15min during 10h (**Fig. 4**). A total number of 63 cells were analyzed from 4 independent experiments, the data are statistically significant. Two representative movies can also be viewed in **Fig. S5**. Only 14 cells did not show visible cell shape modification while the majority (49 bacteria) went through sequential morphological transformations. We classified the transformation steps into 3 categories: "bending" (U-shaped), "pseudo-coccoid" (quasi-round bacteria) and "donut" (fully developed coccoids) and quantified them (**Fig. 4**). At 185±8.0 min post-induction, 77% of the cells analyzed had undergone bending. One hundred minutes later, only 12 cells were still at this stage while the others had turned into pseudo-coccoids. Finally, one third of the analyzed cells (21 cells) ended up in the round "donut" shape at 420±25.2 min after toxin-induction.

Most interestingly, the length of bacteria just before they entered the first morphological transformation (bending) was 3.48±0.08μm, which precisely corresponds to the maximum length we measured before *H. pylori* division. In addition, we observed many examples of two bacteria that were dividing but not yet separated and from which one out of the two underwent morphological transformation (see movie in **Fig. S5**). This analysis gave us a dynamic picture of the morphological transformation of *H. pylori.* Importantly, this analysis

strongly suggests that the morphological transformation is consecutive to a toxin-induced

perturbation of cell division, through a mechanism that is still to be defined.

Toxin-antitoxin imbalanced expression upon oxidative stress

We then searched what physiological signal could trigger the induction of the AapA1 toxin

expression in *H. pylori*. First, the respective activities of individual *aapA1* and *IsoA1*

promoters were measured with chromosomally expressed transcriptional *lacZ* fusions (**Fig.**

S1). Under normal growth conditions, ß-galactosidase activities indicated that the antitoxin

promoter has a 10-fold stronger activity than the toxin promoter (**Fig. 5A**). This result is

consistent with type I TA features where the antitoxin RNA is strongly expressed (but highly

labile) compared to the more stable toxin transcript that is expressed at a lower level (7, 8, 18).

The *lacZ* fusions were then used to follow the activity of the two promoters during *H. pylori*

growth and to search for conditions relevant to *H. pylori* life-style that could lead to

imbalanced expression of the two promoters. Only a slight increase in activity of both

promoters was observed over time of growth (**Fig. 5A**). No significant differential expression

of the promoters was observed upon acid, antibiotic stress or during exposure to high nickel

concentrations (**Fig. S6**). In contrast, oxidative stress induced by hydrogen peroxide (H_2O_2)

resulted in a strong specific decrease in the *IsoA1* promoter activity while, at the same

concentration, the toxin promoter activity was marginally reduced (**Fig. 5B**). Exposure to

paraquat, another oxidative stress agent, resulted in a comparable strong decrease of both

promoters. The expression patterns and stability of the *aapA1* and *IsoA1* transcripts were then

analyzed by Northern Blot with total RNA extracted at different time points after rifampicin

addition. As shown in **Fig. 5C** and **Fig. S7**, the *aapA1* transcript was detected as a 175

nucleotide-long band with an estimated half-life of approx. 120min. In contrast, the 75 nt

IsoA1 full length transcript declines much faster, with an estimated 25min half-life, and the

production of an approx. 50 nt short processed form. Upon hydrogen peroxide exposure, the

half-life of the AapA1 transcript was not significantly modified while we observed a more

rapid depletion of the full-length *IsoA1* transcript with an almost immediate accumulation of

the processed form (**Fig. 5C**). From three independent experiments, we quantified the relative

amounts of the transcripts under normal growth conditions versus H_2O_2 exposure during 120

min after rifampicin addition (**Fig. 5D**). No significant change was observed for the *aapA1*

transcript while H_2O_2 exposure resulted in a diminished half life of the *IsoA1* full length

transcript and in an imbalanced ratio of *IsoA1* in favor of its processed form (**Fig. 5D**). Taken

together, these results show that H_2O_2 causes both diminished *IsoA1* transcription and

increased degradation of *IsoA1* full-length transcript. These observations suggest that, by

decreasing the amounts of antitoxin transcript, exposure to oxidative stress favors translation

of the AapA1 mRNA and thus toxin production.

Oxidative stress induction of coccoids: are class A Type I TA the only effectors ?

Since we established that oxidative stress results in imbalanced expression in favor of the

expression AapA1 toxin, we examined the consequences of exposure to this condition on *H.*

pylori morphology and viability. Upon H_2O_2 oxidative stress, we observed a rapid

transformation into coccoids (approx. 6h) in strain B128. This strain contains six class A type

I TA systems, the A2 locus being inactive (**Fig. S8**). To assess their role, we constructed

mutants carrying deletions of each of its active AapA/IsoA TA loci, Δ*A1*, Δ*A3*, Δ*A4*

(including deletion of both *A4-1* and *A4-2* tandem systems), Δ*A5*, Δ*A6* or of the five loci

collectively (Δ*5*) using a non-marked deletion strategy (24). Under normal conditions, the

growth rate, culturability and kinetics of occurrence of coccoids of the different mutants was

similar to that of the parental strain (**Fig. S9**). Upon H_2O_2 oxidative stress, the Δ5 mutant

strain displayed similar kinetics of conversion to coccoids when compared to the parental

strain.

This shows that oxidative stress is triggering transformation into coccoid but that the type I

toxins are not the only effectors. TA systems have been shown to play a role in persister cell

formation under stress conditions (7, 8). Therefore, the number of persisters after exposure to

hydrogen peroxide was measured for the ΔA1 and Δ5 mutants and compared to the parental

strain (**Fig. S9**). No significant difference was observed, suggesting that, upon hydrogen

peroxide stress, the class A TA systems are not promoting viable persisters in *H. pylori*.

Discussion

Here we established that the expression of a small toxin from a toxin-antitoxin system triggers a massive and rapid morphological transformation of the spiral-shaped *H. pylori* bacterium into round coccoid cells. Transformation of spirals into coccoids has been observed after extended *H. pylori* growth (>70h, aging coccoids) or upon stress conditions (14) but the process itself was poorly characterized. The toxin (AapA1) is a small hydrophobic peptide of a type I TA system that we showed here to target *H. pylori* inner membrane without being secreted. Toxins of type I TA system from other organisms such as *Escherichia coli*, *Bacillus subtilis* or *Staphylococcus aureus* have been shown to localize to the inner membrane but there are no reports on induction of major cell shape modifications (8, 25-27). The *E. coli* TisB (28) and the *S. aureus* PepA1 (29) toxins act by creating membrane pores (similar to phage holins) thereby disrupting the membrane potential, drastically impairing ATP synthesis and depending on the toxin concentration, either leading to the formation of persisters or to cell death (8, 27). One notable exception is the BsrG/SR4 Type I TA system of *B. subtilis* (26). The BsrG toxin was found to target the membrane without causing destruction or affecting the Proton Motive Force but it rather directly interferes with the cell envelope biosynthesis, indirectly delocalizes the cell-wall synthesis machinery and ultimately triggers bacterial autolysis (26). There are now two examples of type I TA hydrophobic toxins (BsrG and AapA1 reported here) that do not act by permeabilizing the membrane and probably reflect a novel mode of action for small peptidic toxins.

Using for the first time cryo-EM, we observed that toxin-induced *H. pylori* coccoids present neither visible membrane disruption nor pores even 8h post toxin induction and are ultrastructurally comparable to 70h-old aging coccoids. The membrane potential of coccoids was not dissipated despite the fact that the toxin induces local perturbation of the membrane. This is in contrast with a previous study that reported a total loss of membrane potential in *H. pylori* aging coccoids, associated with a complete loss of membrane integrity (30). Unlike the

effects of the TisB toxin in *E. coli* (28), we found that the ATP content in our *H. pylori* toxin-

induced coccoids is marginally affected and drops only after prolonged culture, as it is the

case for spiral-shaped cells (31).

Toxins expressed by TA systems cause growth arrest by interfering with conserved vital

cellular processes such as translation, division or peptidoglycan (PG) synthesis (8, 27). In

bacteria, PG cell-wall dictates cell shape. We found that coccoid transformation was

accompanied by changes in PG composition. Indeed, when compared to spirals, coccoids

have a three-fold increase in dipeptide monomers concomitant with a tri- and pentapeptide

monomers reduction. These changes are comparable to those previously reported for "aging"

H. pylori coccoids (32) and were confirmed here. Our results point to a reduction in the

potential for generating new cross-bridges in the PG; this is compatible with a looser PG

macromolecule and could explain the morphological transition to coccoids. The PG

metabolism has been extensively studied in *H. pylori* (33). The modifications we observed in

coccoids suggest activation of at least three sequential *H. pylori* PG hydrolase activities. A

D,D-carboxypeptidase that transforms pentapeptides into tetrapeptides; HdpA/Csd3 is the

only reported such enzyme (34). A L,D-carboxypeptidase that transforms tretapeptides into

tripeptides carried out by Csd6, a member of the L,D-transpeptidases (33). Finally, a (g)-

glutamyl-diaminopimelate carboxypeptidase generating dipeptides from tripeptides carried

out by Cds4, a member of the M14-zinc metallopeptidases (33). We previously showed that

these three PG modifications are blocked in a mutant deficient in AmiA, a putative *H. pylori*

amidase and that an *amiA* mutant fails to transform into coccoids (35). Therefore, we propose

that AmiA could be a central hub for the functionality of a PG hydrolase complex including

AmiA, HdpA, Csd4 and Csd6. In this case, the AapA1 toxin would interfere with the function

of this complex by causing abnormal activation of their enzymatic activities.

Life imaging analysis of the toxin-induced conversion of *H. pylori* spirals into coccoids

suggested that the AapA1 toxin interferes with cell division. This is also consistent with a

central role of AmiA in the conversion, indeed this enzyme was shown to be essential for the

separation of daughter cells (35). In *E. coli*, the CbtA toxin of a type IV TA system was found

to inhibit cell division and cell elongation via direct and independent interactions with FtsZ

and MreB (36). More work is needed to identify the molecular target(s) of the AapA1 toxin.

Under normal conditions, as for all TA systems, the *IsoA1* antitoxin potently inhibits the

AapA1 toxin, by preventing its expression through a multilayered mechanism (20). We

searched for physiological conditions that could result in toxin expression. We found that

oxidative stress, generated by H_2O_2, causes a rapid and specific decline in the levels of IsoA1

antitoxin full-length transcript by reducing both its promoter activity and enhancing its

degradation through a mechanism that remains to be precisely defined. Imbalanced expression

of the AapA1/IsoA1 system in favor of the toxin mRNA should lead to toxin expression.

Accordingly, we observed that H_2O_2 causes a rapid conversion of *H. pylori* into coccoids

resembling those induced by the AapA1 toxin. Prolonged exposure to aerobic conditions was

previously reported to lead to the coccoid formation, however, in that case these cells lost

their membrane integrity (37). Regulation of type I TA systems in response to stress had been

observed in other species. For the *bsrE*/SR5 type I system of *B. subtilis*, the SR5 antitoxin is

affected by pH, anoxia and iron limitation while the BsrE toxin is sensitive to temperature

shock and alkaline stress (38). In *S. aureus*, the SprA1/SprA1$_{AS}$ system is induced in response

to acidic or oxidative stresses (29). In *E. coli*, several type I toxins are induced by the SOS

response (7), a system that does not exist in *H. pylori*.

The biological function of *H. pylori* coccoids is still a matter of debate. Coccoids are

observed in prolonged *in vitro* cultures and induced by stress conditions, mostly aerobic,

anaerobic culture or exposure to antibiotic or oxidative stress. However, these reports are

difficult to compare since no standard procedure was used and analysis was most of the time

performed after more than a week stress exposure (14, 30, 39). *H. pylori* coccoids are non-

culturable cells under standard laboratory growth conditions. Our analysis of both toxin-

induced and 70h aging coccoids are in favor of their viability despite their "unculturability".

We argue that contradictory conclusions on viability refer to "damaged" or so-called

"fragmented" coccoid forms corresponding to membrane-less bacteria in prolonged *H. pylori*

cultures (30). In favor of coccoid viability are previous reports showing that coccoids express

virulence factors (14) and are capable, like spirals, of binding to host cells and inducing

cellular changes (40). *H. pylori* coccoids were also visualized in human gastric biopsies (41),

probably as part of biofilms (39). In mice, coccoids induce gastritis (42) and can revert to

colonizing spiral bacteria (43). Taken together, these observations suggest that coccoids are

"dormant" viable forms of *H. pylori* that recover during host infection and might play a role

in transmission or in treatment failure. In agreement with this view, Chaput *et al.* (35) showed

that coccoids present decreased activation of NF-kB and might allow the bacterium to escape

the host immune response.

We showed that *H. pylori* coccoids are induced by a class A type I TA toxin and by

oxidative stress probably through the imbalanced expression of TA systems. During infection,

H. pylori is indeed exposed to harsh oxidative stress as a consequence of the chronic

inflammation it generates (44). Deletion of the five "class A" Type I TA clusters of *H. pylori*

did not preclude the oxidative stress induction of coccoids nor did it change the number of

persister cells. We concluded that these toxins are probably not the only triggers of coccoid

transformation and suggest that the two other classes of chromosomally-encoded type I TA

systems (B and C representing 9 systems) might also be important.

Dormant/persister bacteria present enhanced tolerance to antibiotics and it has been shown

that they can be induced by TA systems (7, 8), although other mechanisms resulting in

lowered ATP content are also associated with their occurrence (11). We hypothesize that TA-

induced *H. pylori* coccoids are dormant bacteria and, as previously proposed, that non-

culturable cells and "classical" dormant bacteria/persisters are part of a shared "dormancy

continuum" (45). In our model, this continuum would depend on the intracellular toxin

concentration as already suggested (46).

Stress-induced morphological transformation of bacteria into non-culturable coccoid-like

cells has been reported in at least 85 bacterial species among which important bacterial

pathogens such as *Campylobacter jejuni, Vibrio cholerae* or *Salmonella typhimurium* (16).

For these organisms, the inducing trigger and function remains to be characterized. We

postulate that these forms are dormant bacteria and that our findings on *H. pylori* might be

more generally relevant and play important roles in recurrent and persistent bacterial

infections.

Figure legends

Figure 1: Expression of the AapA1 toxin from the AapA1/IsoA1 TA system induces growth arrest and transformation of *H. pylori* strain B128 into coccoids.

A) Genetic organization of AapA1/IsoA1 locus in *H. pylori* B128 strain with the AapA1 transcript encoding the toxin and the *IsoA1* antitoxin RNA. Below is the representation of the inserts of the three plasmids constructed by Arnion *et al* (20) and used in this study, each derived from the pILL2157 *E. coli/H. pylori* vector (21). These inserts are expressed under the control of an IPTG inducible promoter. Plasmid pA1-IsoA1 expresses both the toxin and the first 30 nucleotides of the *IsoA1* RNA antitoxin. In pA1, the promoter region of IsoA1 was mutated without affecting the amino acid sequence of AapA1. Plasmid pA1* was derived from pA1 and contained an additional mutation inactivating the start codon of AapA1.

B) Expression of the AapA1 toxin causes rapid growth arrest of *H. pylori* strain B128. Growth kinetics of *H. pylori* B128 strains carrying each of the three plasmids illustrated in panel A, grown in the absence (black symbols) or in the presence of 1mM IPTG (empty symbols) that was added at the time indicated by an arrow (16h).

C) Expression of the AapA1 toxin induces transformation of *H. pylori* into coccoids. Microscopy analysis of the *H. pylori* B128 strains carrying each of the three plasmids after growth in the absence of IPTG or at 4 and 8h following IPTG addition. *H. pylori* membranes were stained with the lipophilic dye TMA-DPH. Phase-contrast and fluorescence images are respectively represented on the upper and bottom panel. The scale bar corresponds to 1μm.

Figure 2: AapA1 toxin targets *H. pylori* inner membrane, weakly disturbs the bacterial membrane potential and poorly affects intracellular ATP content.

A) AapA1 toxin localizes to the *H. pylori* inner membrane. Western blot analysis of total extracts (T), soluble extracts (SE), inner membrane (IM) and outer membrane (OM) fractions

464 prepared from *H. pylori* B128 strain expressing either a SPA tagged-AapA1 toxin (pA1-SPA)

465 under control of an IPTG-inducible promoter or carrying plasmid pA1* as a negative control.

466 The fractionation procedure was validated with antibodies against the following control

467 proteins, PBP2 for the inner membrane, AlpA for the outer membrane and NikR for the

468 cytoplasm. In **Fig. S2**, we show that a A1-GFP fusion expressed either from the chromosome

469 of B128 or from a plasmid also localizes at the inner membrane.

470 **B) In live cells, AapA1-GFP localizes as discrete foci around the *H. pylori* membrane.**

471 Strain B128 expressing the AapA1-GPF fusion protein was analyzed on agarose pads by

472 fluorescence microscopy 6h after IPTG-induction. Membranes were stained with TMA-DPH

473 lipophilic dye. The membrane association of AapA1-GPF was quantified by measuring the

474 fluorescence intensity profile perpendicular to the length axis of *H. pylori*. The graph shows

475 the average fluorescence intensity profiles with standard deviation (n=25). The fluorescence

476 maxima separated by 0.5 μm, correlate with the *H. pylori* cell width.

477 **C) Analysis of the effect of the AapA1 toxin on *H. pylori* membrane potential.**

478 MitoRed CMXROS, a membrane potential reactive dye was used to analyze live *H. pylori*

479 B128 strains expressing each of the three plasmids illustrated in **Fig. 1**. Cells were grown in

480 presence or absence of 1mM IPTG and analyzed on agarose pads at 0, 4, 8 and 24 h after

481 IPTG addition. While uniformly stained by the MitoRed CMXROS in the absence of IPTG,

482 cells expressing the toxin (pA1) present discrete foci of stronger fluorescence suggesting a

483 local disturbance of the membrane potential. For comparison, a 72h-old culture of B128 WT

484 strain forming "aging" coccoids was also analyzed. As a positive control of membrane

485 potential loss, 4h-old B128 WT cells were analyzed after treatment or not with TCS (3,3',4',5-

486 Tetrachlorosalicylanilide) an *H. pylori* active protonophore. Bar scale represents 1μm.

487 **D) Measurement of intracellular ATP content.**

488 Intracellular ATP was extracted from B128 strains harboring pA1-IsoA1 or pA1 after 6 or

489 16h growth in the presence or absence of IPTG. B128 WT strain treated with 500μM of TCS

protonophore was used as a positive control. ATP concentrations were determined using a

luciferase-based assay (BacTiter-Glo™, Promega). Results from 3 independent experiments

performed in triplicates are shown. Error bars represent the standard deviation, with * and **

indicating that the mean value is significantly different, * corresponds to $P < 0.05$, ** to $P <$

0.01, *** to $P < 0.001$ and NS corresponds to non-significant, ($P > 0.05$).

Figure 3: Ultrastructural analysis of toxin-induced coccoids and "aging" coccoids by

cryo-electron microscopy.

Cryo-electron microscopy was used to compare "aging" coccoids (70 h-old cultures) and

exponentially growing B128 strains carrying plasmid pA1-IsoA1 or pA1 grown for 4 or 8h in

the presence of IPTG. Expression of the toxin results in the formation of U-shaped bacteria at

4h and round coccoids cells at 8h with visible intact bacterial cell envelop similar to the aging

coccoids. Scale bar represents 1μm.

Figure 4: Time-lapse microscopy of *H. pylori* conversion to coccoid upon toxin

expression.

Live cell time lapse microscopy was used to record, during 10 hours, the morphological

modifications of individual *H. pylori* B128 strain cells after IPTG-induction of the AapA1

toxin. Phase-contrast and fluorescence images of one example of these kinetics are presented.

Scale bar represents 2μm. Three major phenotypes were observed; "bending", "pseudo-

coccoids" and "donut". Statistics (median and mean ± SEM) on the total number of analyzed

cells and the precise timing to reach the phenotypes are presented. SEM corresponds to

standard error of the mean. Representative movies of the transformation can be seen in **Fig.**

S5.

Figure 5: Oxidative stress generated by hydrogen peroxide results in decreased antitoxin promoter activity and promotes *IsoA1* transcript processing.

A) Activity of the AapA1 and IsoA1 promoters as a function of growth.

β-galactosidase activities (expressed in Miller units) measured with *H. pylori* B128 strain carrying chromosomal P*aapA1-lacZ* and P*IsoA1-lacZ* fusions after 6, 24 and 48h growth. Results from 3 independent experiments performed in duplicates are shown. Error bars represent the standard deviation, with * ($P < 0.05$) and ** ($P < 0.01$) indicating that the mean values are significantly different and NS indicating that they are not significantly different ($P > 0.05$). The activity of the toxin fusion (*aapA1-lacZ*) is about ten times that of the antitoxin fusion (*IsoA1-lacZ*) and the activity of both promoters slightly increases as a function of growth.

B) Oxidative stress generated by H_2O_2 decreases antitoxin promoter activity.

β-galactosidase activities expressed by P*aapA1-lacZ* and P*IsoA1-lacZ* fusions were measured after 6h treatment with oxidative stress generators, paraquat (5, 50 and 500 μM) and H_2O_2 (0.03 and 0.3%). β-galactosidase activities are presented as ratio (expressed in %) of activities measured with stress versus activities of untreated samples. Hydrogen peroxide strongly decreases the expression of the P*IsoA1-lacZ* fusion. Results from 3 independent experiments performed in duplicates are shown. Error bars represent the standard deviation, with * ($P < 0.05$), ** ($P < 0.01$), *** (P < 0.001) indicating that the mean values are significantly different and NS that they are not significantly different ($P > 0.05$).

C) Determination of AapA1 mRNA and IsoA1 RNA half-lives in *H. pylori* strain B128 under normal conditions and upon hydrogen peroxide exposure.

Northern blots of total RNA from B128 WT strain grown under normal conditions or exposed to 1% hydrogen peroxide, extracted at the indicated times after addition of 80 μg/ml rifampicin. Five μg of RNA were loaded in each lane and the membranes were probed with

[γ-^{32}P] ATP-labeled oligonucleotides specific to the following RNAs, *aapA1*, *IsoA1* and 5S

rRNA as a loading control. In the presence of H_2O_2, the half-live of *IsoA1* transcript is

diminished in favor of the processed *IsoA1* form.

D) Hydrogen peroxide promotes *IsoA1* transcript processing.

Graphic representation of the effects of hydrogen peroxide exposure on the amounts of *aapA*

and *IsoA1* transcripts as well as on *IsoA1* processing during 120 min after rifampicin addition.

The relative amounts of *aapA1* (upper graph) and full length (FL) *IsoA1* (middle graph)

RNAs versus 5S rRNA are shown. The lower graph presents the relative amounts of the two

forms of *IsoA1*, full length (FL) versus the processed form. The results of 3 independent

experiments are shown, error bars represent the standard deviation analyzed by two-way

ANOVA multiple comparisons.

Acknowledgments

Acknowledgments

This work was funded by the Agence National de la Recherche (ANR 09 BLAN 0287 01 and

ANR 12 BSV5-0025-02), the Laboratoire d'Excellence IBEID (Integrative Biology of

Emerging Infectious Diseases) Grant ANR-10-LABX-62-IBEID from the French

government's Investissement d'Avenir program and the Pasteur-Weizmann Consortium of

"The Roles of Noncoding RNAs in Regulation of Microbial Life Styles and Virulence". LEM

was funded by a Roux fellowship of the Institut Pasteur and by a Carnot MI fellowship. AR

was supported by post-doctoral fellowship from the Labex IBEID (10-LABX-62-IBEID). We

also thank Janssen for financial support. We are grateful to F. Darfeuille and H. Arnion for

the gift of plasmids and to W. Fischer for the gift of anti-AlpA antibodies. We also appreciate

the expertise and help of A. Chery-Faleme, M. Fromont-Racine and A. Jacquier with

Northern blotting and of R. Wheeler for peptidoglycan extraction. We thank M. Denic, S.

Kumar and A. Tejada-Arranz for comments on the manuscript and J. Berry for discussions.

Finally, we thank M. Nilges and the Equipex CACSICE for providing the Falcon II direct

detector.

Material and Methods

Bacterial strains and growth conditions

The *H. pylori* strains used in this study (Suppl **Table S2**) were all derived of strain B128 (47) (48). Plasmids (suppl **Table S3**) used to create or complement *H. pylori* mutants were constructed and amplified using *E. coli* One-Shot TOP10 or DH5α strains (Thermofisher). *H. pylori* strains were grown on Blood Agar Base 2 (Oxoid) plates supplemented with 10% defibrinated horse blood and with the following antibiotics-antifungal cocktail: amphotericin B 2.5 $\mu g.ml^{-1}$, polymyxin B 0.31 $\mu g.ml^{-1}$, trimethoprim 6.25 $\mu g.ml^{-1}$ and vancomycin 12.5 $\mu g.ml^{-1}$. Selection of *H. pylori* mutants was performed using kanamycin 20 $\mu g.ml^{-1}$, Streptomycin 10$\mu g.ml^{-1}$, Apramycin 10$\mu g.ml^{-1}$ or chloramphenicol 8 $\mu g.ml^{-1}$. For liquid cultures, we used Brain Heart Infusion (BHI) broth (Oxoïd) supplemented with 10% Fetal Calf Serum (FCS) (Eurobio), the antibiotics-antifungal cocktail and the selective antibiotic when necessary. *H. pylori* cells were grown at 37°C under microaerophilic atmosphere conditions (6% O_2, 10% CO_2, 84% N_2) using an Anoxomat (MART Microbiology) atmosphere generator. When indicated, 1mM of isopropyl ß-D-1-thiogalactopyranoside (IPTG, EuroMedex) was added to agarose pads or culture media.

Molecular techniques

Molecular biology experiments were performed according to standard procedures and the supplier (Fermentas) recommendations. NucleoBond Xtra Midi Kit (Macherey-Nagel) and QIAamp DNA Mini Kit (Qiagen) were used for plasmid preparations and *H. pylori* genomic DNA extractions, respectively. PCR were performed either with Taq Core DNA polymerase (MP Biomedicals), or with Phusion Hot Start DNA polymerase (Finnzymes) when the product required high fidelity polymerase. The pGEM-T easy vector systems (Promega) was

used to construct in *E. coli*, the suicide plasmids that served for markerless deletions of TA

systems in *H. pylori*.

Construction of H. pylori strains carrying mutation or plasmids

Mutations of *H. pylori* were introduced into strain B128 either WT or a streptomycin

resistant variant B128 *rpsL1* for markerless mutagenesis (24, 49). Chromosomal deletions of

the *AapA1/*IsoA1 locus, of the *flaA* gene, or of the different class A TA loci were performed

either by insertion of a selectable antibiotic resistance marker to disrupt or replace the gene of

interest or when necessary, by the markerless counter-selected mutagenesis strategy (24, 49).

Plasmids were derived from the pILL2157 *E. coli/H. pylori* shuttle vector that contains an

IPTG-inducible promoter (21). Deletions were introduced by allelic exchange using suicide

plasmids (see suppl **Table S3**) or PCR fragments. Introduction of plasmids and construction

of *H. pylori* mutants were obtained by natural transformation and selection with the

corresponding antibiotic as described previously (24). PCR and sequencing of the regions of

interest were used to validate the introduction of plasmids, deletion of genes of interest and

correct insertion of fusions. Primers used for these constructs or their validation are listed in

Suppl **Table S4.**

Confocal Fluorescence Microscopy

Fluorescence microscopy was performed with an Axio Observer microscope (Zeiss)

equipped with an Axiocam camera under an X100 magnification. Acquisition images was

performed using the axiovision software. Images were cropped and adjusted using ImageJ

1.47v software. *H. pylori* B128 strains were grown in BHI medium with chloramphenicol (8

μg/ml) to maintain the plasmids. IPTG was added to the culture and samples were taken at

different time-points, concentrated by 2 min centrifugation at 3000 x g and washed twice with

PBS buffer. Cells were immobilized using 2% (wt/vol) agarose pads containing PBS before

being imaged. In **Fig. 1C**, cell membranes were stained with 0.01 mM TMA-DPH (1-(4-trimethylammoniumphenyl)-6-phenyl-1,3,5-hexatriene p-toluenesulfonate, Euromedex). Membrane potential was revealed with 25nM MitoTracker®Red CMXROS (Invitrogen) in **Fig. 2C** and with 1µM DIOC-5 (3) (3,3-Dipentyloxacarbocyanine iodide, Interchim) in **Fig. S3**. For the membrane potential analysis, the negative control was obtained by treatment with the protonophore 500µM TCS (3,3',4',5-Tetrachlorosalicylanilide; Fisher scientific).

Fractionation and Immunoblotting

The protocol of *H. pylori* fractionation was adapted from (50). *H. pylori* strains were grown in the absence or presence of 1mM IPTG. When cultures reached OD_{600} 0.8, cells were harvested by centrifugation, washed twice in phosphate-buffered saline medium (PBS) prior to their disruption by sonication in Buffer 1, 10mM Tris-HCl, pH7.5 containing 5mM ß-mercaptoethanol and proteases inhibitors (cOmplete™, EDTA-free Protease Inhibitor Cocktail, Roche). Cell debris were removed by centrifugation (10 min, 16,000 x *g*, 4°C) and supernatants containing the soluble extract and membrane fractions were collected as total extracts. Samples of total extracts were frozen and the remaining supernatants were subjected to ultracentrifugation (125,000 x *g*, 45 min, 4°C). Samples from the supernatant containing the cytoplasm and periplasm were collected and frozen while pellets containing total membranes were resuspended in buffer 1 supplemented with 1% N-lauroylsarcosine (Buffer 2) prior to be ultracentrifuged (125,000 x *g*, 45 min, 4°C). Samples of inner membrane fractions from supernatants were frozen. Outer membrane pellets were washed twice in buffer 2 to avoid inner membrane fractions contamination prior to freezing. In each fraction, protein amounts were calibrated using the Bradford DC Protein Assay (Biorad) with bovine serum albumin (BSA) as a standard. For each cellular compartment, equal protein amounts were loaded and separated on a 4-20% Mini-Protean TGX precast protein gel (BioRad) and subsequently electrotransferred on a polyvinylidene difluoride (PVDF) membrane (Biorad).

The *H. pylori* proteins PPB2, AlpA and NikR as well as the fusion protein A1-SPA or A1-

GFP were detected with rabbit polyclonal antibodies α-PBP2 (51), α-AlpA (50), α-NikR

(52); with mouse monoclonal anti-FLAG M2 antibody (Sigma-Aldrich) or with goat anti-

GFP-HRP antibody (Abcam) at the respective dilutions of 1:2,000, 1:1,000, 1:100 and

1:1,000 or 1:5,000. Goat anti-rabbit IgG-HRP (Santa Cruz) and ECL anti-mouse IgG

Horseradish Peroxydase (Amersham) were used as secondary antibodies and the detection

was achieved with the ECL reagent (Pierce).

ATP extraction and assay

Exponentially growing *H. pylori* cells (6h of culture OD_{600nm} 0.35) and stationary growing

cells (16h, OD_{600nm} 3) were harvested by centrifugation at room temperature for 4 min at

5000g. Metabolites from the resulting cell pellets were extracted immediately using 300μL of

a solvent mixture of Acetonitrile/Methanol/H_2O (40/40/20) for 15min at 4°C and 10min at

95°C. Mixtures were subsequently spun in a microfuge for 5 min at maximum speed and 4°C

to separate insoluble materials from the extracted metabolites. The resulting pellets were then

re-extracted twice with 200 μL of solvent at 4°C. The supernatants were pooled to yield 700

μL of final extract. Metabolites were lyophilized and subsequently diluted in water for ATP

assays. ATP content was determined by a luciferase based ATP bioluminescence assay kit

(BacTiter-Glo™ Microbial cell viability assay, Promega). Luminescence values were

determined using a 10 sec RLU signal integration time and measured using a Centro XS^3

LB960 Luminometer (Berthold Technologies). ATP concentrations were calculated based on

values determined using serial dilutions of known amounts of ATP and expressed as a

function of the OD_{600nm} of the corresponding culture.

Cryo-electron microscopy

671 Four μL of *H. pylori* bacteria were spotted on glow-discharged lacey grids (S166-3, EMS) or

672 Quantifoil R2/2 (Quantifoil, Germany). The samples were cryo-fixed by plunge freezing at -

673 180°C in liquid ethane using a Leica EMGP (Leica, Austria). Grids were observed at 200kV

674 with a Tecnai F20 (Thermo Fisher Scientific). Images were acquired under low-dose

675 conditions using the software EPU (Thermo Fisher Scientific) and a direct detector Falcon II

676 (Thermo Fisher Scientific).

678 **Time-lapse microscopy**

679 Confocal analysis of live cells was performed as previously described (53). Twenty μL of

680 exponentially grown bacteria (10^5 cells) suspended in BHI with or without 1mM IPTG were

681 deposited on 35mm glass bottom Petri dishes. The suspension was covered with BHI medium

682 in 1.5% low melting agarose supplemented with Chloramphenicol (8μg/μl) with or without

683 IPTG (1mM). Live-cell imaging was performed with a Nikon A1R confocal laser scanning

684 microscope system attached to an inverted ECLIPSE Ti (Nikon Corp., Tokyo, Japan) and

685 equipped with an environmental chamber allowing the control of temperature (37°C),

686 humidity (90%) and gas mixture (10% CO_2, 3% O_2). GFP fluorescence images were captured

687 through a Plan APO 60X objective (NA: 1.40) by using optimal spatial resolution settings.

688 The cytoplasm compartment volume was defined by using GFP staining (excitation with 488

689 nm laser, emission collected with an 500/50 nm filter set). Image captions were performed

690 every 10min during 10h. Statistics were performed on a total of 63 cells from 4 independent

691 experiments. Image treatment and analysis were performed using NIS elements (Nikon Corp.,

692 Tokyo, Japan) and ImageJ software.

694 **ß-galactosidase activity assays**

695 B128Δ*AapA1-IsoA1::PaapA1-lacZ-Kan* and B128Δ*AapA1-IsoA1::PIsoA1-lacZ-Kan*

696 strains were grown to OD_{600nm} 0.3 in BHI liquid medium, then divided into two samples, one

of them being submitted to stress conditions during 6h with 5, 50 and 500 μM paraquat

(Sigma), 0.03 and 0.3% hydrogen peroxide (Sigma), 20 and 200 mM NiCl$_2$ (Sigma), 0.1 and

1 mg/mL tetracycline, or 0.05 and 0.5 mg/ml rifampicin. Then 0.5ml samples were taken,

washed twice with 1X PBS (Phosphate Buffered Saline) and further permeabilized with

100μL Chloroform and 50μl SDS 0.1% in Z buffer containing 70mM Na$_2$HPO$_4$.12H$_2$O,

30mM NaH$_2$PO$_4$. H$_2$O, 1mM MgSO$_4$ and 0.2mM MnSO$_4$ (54). Samples were briefly vortexed

and incubated at 28°C for 2 min. The ß-galactosidase assay was started by adding 0.5mL

ONPG (ThermoFischer) at 4mg/ml and stopped by the addition of 0.5mL 1M Na$_2$CO$_3$ when

sufficient yellow color was reached. The ß-galactosidase activity is expressed in Miller units

(54) and represented in **Fig. 5** and **S6** as percentage of activities relative to the control activity

with the corresponding strains not exposed to stress and was calculated from 3 independent

experiments performed in duplicates.

Total RNA extraction and Northern blotting

Total RNAs were extracted from 10 mL *H. pylori* cultures (OD$_{600}$ 0.5 -0.9) exposed or not to

1% H$_2$O$_2$ using the NucleoSpin miRNA kit (Macherey Nagel) at different time points after

80 μg/mL rifampicin addition (to stop transcription). For Northern blotting, 5 μg of total RNA

were separated on 10% Mini protean TBE Urea gel (Biorad) and transferred to Hybond N+

(Amersham Biosciences) membrane susing a Trans-Blot Turbo system (Biorad). A DNA

marker was used for size estimation.

Transferred RNA was fixed to the membrane by UV irradiation for 2 min. The membrane

was blocked for 45 min at 42°C with ULTRAhyb Hybridization Buffer (Ambion), then 20 μl

of 5'-labeled (γ^{32}P) oligodeoxynucleotides (**Table S4**) were added and the membrane was

further incubated overnight at the same temperature. After three washes for 10 min at 65°C

with 2x SSC 0.2 % SDS, the membrane was exposed to a phosphorimager screen (KODAK)

and scanned with FLA-9000 Phospho Imager (Fujifilm).

Peptidoglycan extraction and Muropeptides analysis by HPLC/HRMS

Samples (1L) of *H. pylori* B128 WT strain (16h, exponential phase and 72h, "aging"

coccoids) and of B128 Δ*aapA1-IsoA1* pA1 strain, induced by IPTG during 6h and 16h, were

taken and chilled in an ice-ethanol bath. Crude murein sacculi were immediately extracted in

boiling sodium dodecyl sulfate (SDS 4% final). The resulting purified peptidoglycan was

digested overnight at 37°C in 12.5 mM sodium phosphate buffer (pH 5.6) supplemented with

100 UI of mutanolysin from *Streptomyces globisporus* ATCC 21553 (Sigma). The reaction

was stopped by heat inactivation of the enzyme and insoluble material was removed by

centrifugation. Soluble muropeptides were then reduced with sodium borohydride in borate

buffer (pH 9). After centrifugation, reduced muropeptides were diluted 20-fold in water-

formic acid 0.1% (v/v; solvent A). HPLC/HRMS (High Performance Liquid Chromatography

/ High Resolution Mass Spectrometry) was performed on an Ultimate 3000 UHPLC (Ultra

High Performance Liquid Chromatography) system coupled to a quadrupole-Orbitrap mass

spectrometer (Q-Exactive Focus, Thermo Fisher Scientific). Muropeptides were separated on

an Hypersil Gold aQ analytical column (1.9 μm, 2.1x150 mm) at 200 μL/min, column

temperature at 50°C. Applied linear gradient from 0 to 12.5% acetonitrile + 0,1% (v/v) formic

acid (solvent B), followed by increasing to 20% B at 25 min for 5 min, hold 20% B for min

and additional 10 min with 100% A for column re-equilibration. Eluted muropeptides were

introduced into Q-Exactive instrument, operating in positive ion mode, and then were

analyzed in the data-dependent acquisition mode (FullMSddMS2). Data were then processed

with the software TraceFinder 3.3 (Thermo Fisher Scientific) for peak areas determination,

the relative abundance of muropeptides in each sample was calculated according to Glauner *et*

al. (23).

Statistical analysis

The Student's t test was used to determine significance of the means of the data. Error bars

represent the standard deviation, with * ($P < 0.05$), ** ($P < 0.01$), *** (P <0.001) indicating

that the mean values are significantly different and NS that they are not significantly different

(P >0.05). SEM corresponds to standard error of the mean. The two-way ANOVA multiple

comparisons was used to compare Northern blots bands intensities under different conditions.

755   References

1.   Yamaguchi Y, Park J-H, & Inouye M (2011) Toxin-Antitoxin Systems in Bacteria and

Archaea. *Annual Review of Genetics* 45(1):61-79.

2.   Goeders N & Van Melderen L (2014) Toxin-Antitoxin Systems as Multilevel

Interaction Systems. *Toxins* 6(1):304-324.

3.   Page R & Peti W (2016) Toxin-antitoxin systems in bacterial growth arrest and

persistence. *Nat Chem Biol* 12(4):208-214.

4.   Yang QE & Walsh TR (2017) Toxin–antitoxin systems and their role in disseminating

and maintaining antimicrobial resistance. *FEMS Microbiology Reviews* 41(3):343-353.

5.   Helaine S*, et al.* (2014) Internalization of *Salmonella* by Macrophages Induces

Formation of Nonreplicating Persisters. *Science* 343(6167):204-208.

6.   Wang X*, et al.* (2011) Antitoxin MqsA helps mediate the bacterial general stress

response. *Nat Chem Biol* 7(6):359-366.

7.   Berghoff BA & Wagner EGH (2017) RNA-based regulation in type I toxin–antitoxin

systems and its implication for bacterial persistence. *Current Genetics*.

8.   Harms A, Brodersen DE, Mitarai N, & Gerdes K (2018) Toxins, Targets, and

Triggers: An Overview of Toxin-Antitoxin Biology. *Molecular Cell*.

9.   Harms A, Maisonneuve E, & Gerdes K (2016) Mechanisms of bacterial persistence

during stress and antibiotic exposure. *Science* 354(6318).

10.   Goormaghtigh F*, et al.* (2018) Reassessing the Role of Type II Toxin-Antitoxin

Systems in Formation of *Escherichia coli* Type II Persister Cells. *mBio* 9(3).

11.   Shan Y*, et al.* (2017) ATP-Dependent Persister Formation in *Escherichia coli*. *mBio*

8(1):e02267-02216.

12.   Yamaoka Y (2010) Mechanisms of disease: *Helicobacter pylori* virulence factors. *Nat*

Rev Gastroenterol Hepatol 7:629–641.

13.   Amieva M & Peek Jr RM (2016) Pathobiology of *Helicobacter pylori*–Induced

782   Gastric Cancer. *Gastroenterology* 150(1):64-78.

783   14.   Loke MF, Ng CG, Vilashni Y, Lim J, & Ho B (2016) Understanding the dimorphic

784   lifestyles of human gastric pathogen *Helicobacter pylori* using the SWATH-based

785   proteomics approach. *Scientific Reports* 6:26784.

786   15.   Reshetnyak VI & Reshetnyak TM (2017) Significance of dormant forms of

787   *Helicobacter pylori* in ulcerogenesis. *World Journal of Gastroenterology*

788   23(27):4867-4878.

789   16.   Ayrapetyan M, Williams T, & Oliver JD (2018) The relationship between the viable

790   but nonculturable state and antibiotic persister cells. *Journal of Bacteriology*.

791   17.   Brantl S & Jahn N (2015) sRNAs in bacterial type I and type III toxin-antitoxin

792   systems. *FEMS Microbiol Rev.* 39:413-427.

793   18.   Masachis S & Darfeuille F (2018) Type I Toxin-Antitoxin Systems: Regulating Toxin

794   Expression via Shine-Dalgarno Sequence Sequestration and Small RNA Binding.

795   *Microbiology Spectrum* 6(4).

796   19.   Sharma CM*, et al.* (2010) The primary transcriptome of the major human pathogen

797   *Helicobacter pylori*. *Nature* 464:250-255.

798   20.   Arnion H*, et al.* (2017) Mechanistic insights into type I toxin antitoxin systems in

799   *Helicobacter pylori*: the importance of mRNA folding in controlling toxin expression.

800   *Nucleic Acids Research* 45(8):4782-4795.

801   21.   Boneca IG*, et al.* (2008) Development of inducible systems to engineer conditional

802   mutants of essential genes of *Helicobacter pylori*. *Appl. Environ. Microbiol.*

803   74(7):2095-2102.

804   22.   Stingl K, Uhlemann E-M, Schmid R, Altendorf K, & Bakker EP (2002) Energetics of

805   *Helicobacter pylori* and Its Implications for the Mechanism of Urease-Dependent

806   Acid Tolerance at pH 1. *Journal of Bacteriology* 184(11):3053-3060.

23.   Glauner B (1988) Separation and quantification of muropeptides with high-
performance liquid chromatography. *Anal Biochem.* 172:451-464.

24.   Fischer F*, et al.* (2016) A novel essential ABC transporter and the NixA permease are
the two sole nickel uptake systems of *Helicobacter pylori. PLoS Pathog* 12:e1006018.

25.   Brantl S (2012) Bacterial type I toxin-antitoxin systems. *RNA Biology* 9(12):1488-
1490.

26.   Jahn N, Brantl S, & Strahl H (2015) Against the mainstream: the membrane‐
associated type I toxin BsrG from *Bacillus subtilis* interferes with cell envelope
biosynthesis without increasing membrane permeability. *Molecular Microbiology*
98(4):651-666.

27.   Brielle R, Pinel-Marie M-L, & Felden B (2016) Linking bacterial type I toxins with
their actions. *Current Opinion in Microbiology* 30:114-121.

28.   Unoson C & Wagner EGH (2008) A small SOS‐induced toxin is targeted against the
inner membrane in *Escherichia coli*. *Molecular Microbiology* 70(1):258-270.

29.   Sayed N, Nonin-Lecomte S, Réty S, & Felden B (2012) Functional and Structural
Insights of a *Staphylococcus aureus* Apoptotic-like Membrane Peptide from a Toxin-
Antitoxin Module. *The Journal of Biological Chemistry* 287(52):43454-43463.

30.   Kusters JG, Gerrits MM, Van Strijp JA, & Vandenbroucke-Grauls CM (1997)
Coccoid forms of *Helicobacter pylori* are the morphologic manifestation of cell death.
Infection and Immunity 65(9):3672-3679.

31.   Enroth H*, et al.* (1999) In Vitro Aging of *Helicobacter pylori*: Changes in Morphology,
Intracellular Composition and Surface Properties. *Helicobacter* 4(1):7-16.

32.   Costa K*, et al.* (1999) The Morphological Transition of *Helicobacter pylori* Cells from
Spiral to Coccoid Is Preceded by a Substantial Modification of the Cell Wall. *J.*
Bacteriol 181(12):3710-3715.

33. Yang DC, Blair KM, & Salama NR (2016) Staying in Shape: the Impact of Cell Shape

on Bacterial Survival in Diverse Environments. *Microbiology and Molecular Biology*

Reviews : MMBR 80(1):187-203.

34. Bonis M, Ecobichon C, Guadagnini S, Prévost M-C, & Boneca IG (2010) A M23B

family metallopeptidase of *Helicobacter pylori* required for cell shape, pole formation

and virulence. *Molecular Microbiology* 78(4):809-819.

35. Chaput C*, et al.* (2006) Role of AmiA in the Morphological Transition of

Helicobacter pylori and in Immune Escape. *PLoS Pathogens* 2(9):e97.

36. Heller DM, Tavag M, & Hochschild A (2017) CbtA toxin of *Escherichia coli* inhibits

cell division and cell elongation via direct and independent interactions with FtsZ and

MreB. *PLoS genetics* 13(9):e1007007.

37. Zeng H, Guo G, Mao XH, De Tong W, & Zou QM (2008) Proteomic Insights into

Helicobacter pylori Coccoid Forms Under Oxidative Stress. *Current Microbiology*

57(4):281.

38. Müller P*, et al.* (2016) A multistress responsive type I toxin-antitoxin system:

bsrE/SR5 from the *B. subtilis* chromosome. *RNA Biology* 13(5):511-523.

39. Cellini L (2014) *Helicobacter pylori*: A chameleon-like approach to life. *World*

Journal of Gastroenterology 20(19):5575-5582.

40. Segal ED, Falkow S, & Tompkins LS (1996) *Helicobacter pylori* attachment to gastric

cells induces cytoskeletal rearrangements and tyrosine phosphorylation of host cell

proteins. *Proceedings of the National Academy of Sciences of the United States of*

America 93(3):1259-1264.

41. Goldstein NS (2002) Chronic Inactive Gastritis and Coccoid *Helicobacter pylori* in

Patients Treated for Gastroesophageal Reflux Disease or With H pylori Eradication

Therapy. *American Journal of Clinical Pathology* 118(5):719-726.

42.   Nagai S*, et al.* (2007) Role of Peyer's patches in the induction of *Helicobacter pylori*-

induced gastritis. *Proceedings of the National Academy of Sciences of the United*

States of America 104(21):8971-8976.

43.   Cellini L*, et al.* (1994) Coccoid *Helicobacter pylori* not culturable *in vitro* reverts in

mice. *Microbiology and Immunology* 38(11):843-850.

44.   Butcher LD, den Hartog G, Ernst PB, & Crowe SE (2017) Oxidative Stress Resulting

From *Helicobacter pylori* Infection Contributes to Gastric Carcinogenesis. *Cellular*

and Molecular Gastroenterology and Hepatology 3(3):316-322.

45.   Ayrapetyan M, Williams TC, & Oliver JD (2015) Bridging the gap between viable but

non-culturable and antibiotic persistent bacteria. *Trends in microbiology* 23(1):7-13.

46.   Rotem E*, et al.* (2010) Regulation of phenotypic variability by a threshold-based

mechanism underlies bacterial persistence. *Proceedings of the National Academy of*

Sciences of the United States of America 107(28):12541-12546.

47.   Farnbacher M*, et al.* (2010) Sequencing, annotation and comparative genome analysis

of the gerbil-adapted *Helicobacter pylori* strain B8. *BMC Genom.* 11:335.: doi:

10.1186/1471-2164-1111-1335.

48.   McClain MS, Shaffer CL, Israel DA, Peek RMJ, & Cover TL (2009) Genome

sequence analysis of *Helicobacter pylori* strains associated with gastric ulceration and

gastric cancer. *BMC Genom.* 10:3:doi: 10.1186/1471-2164-1110-1183.

49.   Debowski AW*, et al.* (2012) Xer-cise in *Helicobacter pylori*: one-step transformation

for the construction of markerless gene deletions. *Helicobacter* 17:435-443.

50.   Kutter S*, et al.* (2008) Protein Subassemblies of the *Helicobacter pylori* Cag Type IV

Secretion System Revealed by Localization and Interaction Studies. *Journal of*

Bacteriology 190(6):2161-2171.

51.   El Ghachi M*, et al.* (2011) Characterization of the elongasome core PBP2: MreC

complex of *Helicobacter pylori*. *Molecular Microbiology* 82(1):68-86.

52.   Muller C*, et al.* (2011) Hierarchical regulation of the NikR-mediated nickel response

884   in *Helicobacter pylori*. *Nucleic Acids Res*. 39:7564-7575.

885   53.   Corbinais C, Mathieu A, Kortulewski T, Radicella JP, & Marsin S (2016) Following

886   transforming DNA in *Helicobacter pylori* from uptake to expression. *Mol Microbiol*

887   101(6):1039-1053.

888   54.   Miller JH (1992) *A short Course in Bacterial Genetics : A laboratory manual and*

889   *handbook for Escherichia coli and related bacteria. Cold Spring Harbor, N. Y* (Cold

890   Spring Harbor Laboratory).

Figure 2

Figure 3

A

| | Time before «bending» phenotype (min)

Time before «pseudo-coccoid» phenotype (min)	Time before «donut» phenotype (min)	Maximum length before morphological change (µm)
Number of cells	49	37	21	63
Median	180	280	420	3.58
Mean ± SEM	185 ± 8.0	277 ± 16	420 ± 25.2	3.48 ± 0.08

B

Figure 5