Relationship between cardiac cycle and the timing of actions during action execution and observation.

Authors: Palser, E. R.[1,2,3]*, Glass, J.[2], Fotopoulou, A.[1], & Kilner, J. M[2].

Author Affiliations: 1 Psychology and Language Sciences, University College London, London UK 2 Department of Clinical and Movement Neurosciences, Institute of Neurology, University College London, London UK 3 Department of Neurology, University of California San Francisco, San Francisco, USA.

*Correspondence: eleanor.palser.14@ucl.ac.uk

Abstract

Previous research suggests that there may be a relationship between the timing of motor events and phases of the cardiac cycle. However, this relationship has thus far only been researched using simple isolated movements such as key-presses in reaction-time tasks and only in a single subject acting alone. Other research has shown both movement and cardiac coordination among interacting individuals. Here, we investigated how the cardiac cycle relates to ongoing self-paced movements in both action execution and observation using a novel dyadic paradigm. We recorded electrocardiography (ECG) in 26 healthy adult subjects who formed 19 dyads (7 comprised of two subjects and 12 subjects paired with an experimenter). Each dyad contained an action executioner and observer as they performed a self-paced sequence of movements. We demonstrated that heartbeats are timed to movements during both action execution and observation. Specifically, movements were less likely to culminate synchronously with the heartbeat, around the time of the R-peak of the ECG. The same pattern was observed for action observation, with the observers' heartbeats occurring off-phase with movement culmination. These findings demonstrate that there is coordination between an action executioner's cardiac cycle and the timing of their movements, and that the same relationship is mirrored in an observer. This suggests that previous findings of interpersonal coordination may be caused by the mirroring of a phasic relationship between movement and the heart.

Introduction

A growing body of research, comprising physiological and psychological investigations, is consistent with the hypothesis that the central nervous system has access to cardiac information, and uses this information to guide behavior. One such instance is the relationship between the cardiac cycle and the timing of movement. One of the earliest observations in this vein was that walking rate and heart rate typically show a one-to-one mapping[1,2]. It was proposed that the rhythm of the heart might represent a pacemaker, guiding the timing of movement.

Crucially, the effect of the heart on movement is bidirectional, with both effects of the cardiac cycle on movement and effects of movement on cardiac output observed. First, anticipatory cardiac slowing in the fore period of a reaction-time task while the subject prepares to move was reported[3, 4]. Secondly, the motor response was also found to vary depending on when in the cardiac cycle it occurred [5,6], with faster reaction times during atrial contraction. A similar effect has recently been observed for movement inhibition, with faster responses to stop cues during systole[7]. Much of this previous physiological work on the relationship between the cardiac cycle and movement has employed simple prescribed movements within reaction-time and stop signal paradigms.

Cardiac-movement effects have been attributed to baroreceptors[8]. Located within the aortic arch, the carotid sinuses and the coronary arteries, these stretch receptors represent one of the key communication channels between the heart and the brain. Baroreceptors are active during systole, conveying information on the timing and strength of the heartbeat to the central nervous system via the vagal and glossopharyngeal nerves, but are quiescent during diastole when there is no such information to convey[9,10]. The heart's rhythmicity is modified by humoral and neural factors and the heart period can be increased or decreased according to

signals conveyed by vagal and sympathetic efferents[9-12]. There is therefore a closed loop

communication reflex mediated by baroreceptors that allows the timing of the heart cycle to

be controlled.

Interoceptive signals, particularly those originating from the heart, have been purported to

play a role in social cognition[13,14]. Simulation theories of action understanding posit that in

order to understand another's behavior, we simulate their perceptual, motor and bodily states

in ourselves[15]. As such, an action is understood based on the resonance of an observer's

motor system[16]. Such ideas have since been formalized by hierarchical predictive models,

wherein it is argued that action observation results in an observer generating an internal

model of how they would perform it[17]. Such models are proposed to contain not only

exteroceptive and proprioceptive predictions[18], but also interoceptive predictions about how

such an action might be performed[19]. That is, a good model contains information pertaining

not only to the expected position of the body in space, and afferent tactile feedback from the

effector, but also the somatovisceral conditions that would be generated by the execution of

that action. These predicted conditions can then be used to infer the causes of another's

behavior, based on the most likely causes of those conditions in the self. Including

interoceptive information in predictive models may provide unique information about others'

affective states, not otherwise accessible through exteroceptive and prioproceptive inference

alone, due to the ubiquitous role of somatovisceral states in emotion and feeling states[20-23].

Findings of resonance beyond the central nervous system suggest that this mirroring effect

may be more widespread than previously thought, extending into the peripheral nervous

system[28-31]. Both external stimuli such as observing someone perform an action, and internal

stimuli, such as mentally rehearsing an action, can elicit autonomic system activity[32,33].

Watching an actor lift a weight or run on a treadmill produces changes in the respiration rate

of an observer, such that it increased linearly with the treadmill speed[34]. These findings

support the idea that bodily systems involved in interoception may also be candidates for

resonance. The direct matching mechanism proposes a match between the conditions of

action execution and observation[16]. Therefore, if this mirroring effect does extend to the

interoceptive system, we would predict that any relationship between the cardiac cycle and

movement timing found during action execution should also be found during action

observation.

One consequence of resonance might be interpersonal coordination. Research in the field of

social neuroscience has found both movement and cardiac coordination between interacting

individuals. In the movement literature, interpersonal coordination may refer to either

mimicry or synchronization[29]. A distinction is generally made based on the temporal relations

of the interacting individuals, with mimicry reflecting non-periodic matched movements and

synchronicity used to describe movements temporally matched in time[30,31]. Early studies

reported that listeners to a speaker tend to move in time with the rhythms of a speaker's

speech[32,33] and mirror the postures of the speaker[34]. In other interactions, the direction of the

effect is less apparent, with partners mimicking each other's postural sway[35] and linguistic

forms[36]. In a recent study, groups of subjects (numbering between four and five individuals)

constructed models out of Lego blocks[37]. They either worked collectively (where subjects

took turns to work on one construction project) or individually (where subjects worked on

their own construction projects side by side). More stable shared heart rate dynamics were

observed in collective trials than individual trials, increasing across the course of collective

trials[37], suggesting that convergence was tied to the social nature of the task. Indeed, heart

rate entrainment has been suggested as a mechanism underlying emotional commonality in

social interaction, facilitating a sense of community between individuals[38], empathy[39,40] and

team performance[41-44]. In support of this, a study in which pairs of expert improvisers were

asked to mirror each other's movements found that periods of both increased kinematic and

subjective togetherness were characterized by higher heart rates and greater inter-player

correlation of heart rate[45]. Findings of interpersonal coordination and its influence on

measures of social competence are compatible with the hypothesis that resonance supports

action understanding.

The existing literature can largely be summarized as reflecting two broadly different

methodological approaches to studying the relationship between the heart and movement.

The first approach has studied the relationship between cardiac events and evoked

movements such as those elicited by paradigms designed to gauge reaction time [5,6] and

response inhibition[7]. These studies tested discrete, single movements. The second has taken

the opposite approach and analyzed cardiac and movement parameters in a number of

unconstrained self-paced environments, typically within a social context[37,38].

The current study was designed to answer the previously neglected question of whether

cardiac-motor timing and inter-individual resonance are linked, bridging the gap between the

two previous approaches, by investigating cardiac-motor synchronicity in both action

execution and observation. Naturalistic actions are more likely to comprise a series of goal-

directed self-paced movements than discrete, cued movements. Therefore, in the present

study, we sought to investigate the bidirectional relationship between the cardiac cycle and

movement during a sequence of meaningful self-paced movements. Subjects were divided

into dyads and took turns to perform and observe a sequence of semi-controlled movements

under experimental conditions, while electrocardiography (ECG) was recorded from both

(Figure 1). In brief, the task required both subjects within a dyad to memorize a sequence of

six movement locations and then take it in turn to replicate the sequence. In this way, subjects

both executed and observed the movements. The timing of movement was recorded using

touch-sensitive pads.

Firstly, we tested for coordination between execution and observation conditions within a

dyad, in both heart rate and movement speed. Secondly, we investigated whether there was

any relationship between the timing of the movements made, operationalized as the endpoint

of the action, and the location of this event within the cardiac cycle, both in terms of its phase

and time relative to R-peak. The endpoint of each action corresponded to each memorized

location in the movement sequence. Thirdly, we tested the prediction that any phasic or time-

based relationship observed between the timing of movement events and the cardiac cycle

during action observation would also be observed during action observation.

Results

Prior to testing our main hypothesis that there would be a relationship between movement timing and the period of the cardiac cycle, we tested for a significant difference in the length of the cardiac cycle (operationalized as the R-R interval) between observation and execution conditions. We also tested whether there was a relationship between the movement time of both subjects in a dyad and if there was a relationship between the length of the cardiac cycle and movement time.

There was no significant difference in the duration of the mean R-R interval between action execution and action observation (825.3 and 822.9 ms respectively; $t(24) = 0.64$, $p = 0.53$). There was no significant linear correlation between mean R-R interval and mean movement time. This was non-significant for both action execution ($r = -0.37$, $p = 0.07$) (Figure 2A) and action observation ($r = -0.12$, $p = 0.58$) (Figure 2B). However, there was a significant linear correlation between the movement time of subjects within a dyad ($r = 0.61$ $p = 0.007$ (Figure 2C). This significant relationship is consistent with the hypothesis that a common movement speed was adopted by the dyad but that this was not intrinsically tied to the length of the cardiac cycle.

The main aim of this study was to test whether there was evidence of a consistent relationship between the timing of movement events and the cardiac cycle phase both when performing actions and when observing someone else perform them. This was tested both in the time and the phase domains.

Time domain analysis

The data were divided into four time windows, centered around 0 (-100 – 100 ms), 200 (100 – 300 ms), 400 (300 – 500 ms) and 600 (500 – 700 ms) milliseconds post R-peak (Figure 3A). The repeated measures ANOVA with factors Condition (execution and observation) and

Time (0, 200, 400 and 600) revealed a significant main effect of Time ($F(2.21, 53.1) = 21.4$,

$p < 0.001$) and no significant effects of condition or interaction between the two factors

($F(1,24) = 0.01$, $p = 0.92$; $F(2.39, 57.24) = 0.10$, $p = 0.93$) respectively. Post hoc t-tests

revealed that the significant effect of Time was driven by there being significantly less

movement events in the 200 ms time period around the R-peak than the other three time

periods (Figure 3B). For execution, the number of events in the -100–100 ms time window

was significantly less than those in the 100–300 ms, and 300–500 ms time windows

(Execution: $t(24) = -4.59$; $d = -0.918$, $p < 0.05$ & $t(24) = -4.04$, $d = -0.81$, $p < 0.001$). For

observation the number of events in the -100–100 ms time window was significantly less

than those in the 100–300 ms, 300–500 ms and 500–700 ms time windows $t(24) = -7.404$, $d =$

-1.48, $p < 0.05$, $t(24) = -4.91$, $d = -0.98$, $p < 0.001$ & $t(24) = -3.65$, $d = -0.73$, $p = 0.001$). All

other within Condition post hoc comparisons were non-significant when corrected for

multiple comparisons.

Phase domain analysis

In contrast to the time-domain analysis, there was no evidence that the mean phase of each

subject's movements differed significantly from the uniform distribution in either the

Execution or Observation conditions (Rayleigh test $z = 0.07$, $p = 0.93$ & $z = 0.84$, $p = 0.43$

respectively (Figure 4A & B)). To investigate the phase data further we analyzed the mean

distribution of the phase of all events across subjects during action execution and observation

conditions. If there was no consistent phase relationship these values should be close to the

excepted value for a uniform distribution (Figure 4C & D). Here, with 20 phase bins, the

proportion in each bin with a uniform distribution would be 0.05 (1/20). However, in both

conditions there were more motor events than expected at particular phases of the cardiac

cycle (Figure 4C–F). For the Execution condition there were significantly more events than

expected in the phase bin centred on $81°$ ($t(24) = 2.07$, $p = 0.049$). For the Observation

condition this occurred later in the cycle, in the phase bin centred on 153° ($t(24) = 2.32$, $p =$

0.029). However, it should be noted that these were small effects that did not survive

correction for multiple comparisons. There was a reduction in the proportion of events that

occurred either before and slightly after the R-peak during Execution, or after the R-peak

during Observation. However, these reductions were not significantly less than expected.

Discussion

Here, we showed that the timing of self-paced movements and phases of the cardiac cycle are linked in a similar way for both action execution and observation. We showed that movement events were significantly less likely to occur at the time of the heartbeat, synchronous with the R-peak, and significantly more likely to occur between heartbeats. This same relationship was also observed between the motor events of an actor and the cardiac cycle of an observer. That is, the observer was significantly less likely to experience a heartbeat when observing movement endpoints. Additionally, there was a significant positive correlation between movement time in execution and observation conditions, suggesting dyads adopted a common movement speed. These results build on previously separate literatures that find both a relationship between discrete motor events and cardiac timing, and cardiac and movement coordination between interacting individuals.

In this study we investigated the relationship between movements and the cardiac cycle in both the time and phase domains. The effect of a relationship between the two was largest in the time domain analysis. Indeed, despite there being a clear and significant reduction in the proportion of the events around the time of the R-peak in the time domain analysis the group level phase analysis showed no evidence that the distribution was non-uniform. The time domain analysis assumes that any cardiac effects are time locked to the R-peak and are independent of the R-R interval. Whereas the phase analysis assumes that any cardiac effects are locked to a particular phase of the R-R interval and are not time locked. This difference in the effect sizes reported here most likely reflects these different sensitivities of the two analyses. Future work needs to focus on studying the different specificities and sensitivities of the different analysis pathways to understand when they should be used and how to understand the differences between them.

Many have suggested that an internal central pacemaker guides the timing of motor

actions[1,2,49,50]. The heart, with its regular rhythmicity that modulates with arousal level,

intuitively represents a good candidate for this role. Information on the strength and timing of

heartbeats is conveyed to the central nervous system via the vagal and glossopharyngeal

nerves. Such information could thus be capitalized on by the motor system when timing the

occurrence of actions. Indeed, there is evidence that stimulation of afferent vagal and

glossopharyngeal pathways is associated with modulations of efferent motor pathways. For

example, pressure applied to the carotid sinus, which modulates baroreceptor firing[51], inhibits

spontaneous movements and reduces muscle tone in anaesthetized animals[52]. Efferent

projections, which can delay the onset of the next heartbeat, could further optimize control

over timing relationships.

The benefits of timing behavior to the cardiac cycle, however, remain unknown. Global

perceptual attenuation during heartbeats has been proposed, as sensations are inferred as

consequences of internal cardiac activity and therefore down-weighted[53]. Indeed, a recent

study purports that exteroceptive sensory confidence is greatest during diastole and at its

lowest concomitant with the heartbeat[54], providing a neat explanation for the preference to

time movements in between heartbeats observed here.

It is possible that actions are timed to the point in the cardiac cycle where the ability to

balance autonomic and behavioural demands is greatest. Previous research has shown that as

an individual attends to and prepares to interact with a stimulus, vagal activation lengthens

the cardiac cycle[55]. In turn, vagal withdrawal, when the individual initiates a response,

shortens the cardiac cycle[56,57]. However, the timing of the behavioral response also impacts

the cardiac consequences. Responses made at or just after the R-wave impact the length of

the current cardiac cycle, whereas responses made later in the R-R interval impact the length

of subsequent cycles. Furthermore, the sensitivity of the sinoatrial node to input from the

vagus is not uniform across the cardiac cycle. Instead, a period of heightened sensitivity is

estimated to occur between 250 and 400 milliseconds post R-wave, followed by a period of

relative insensitivity[58,59]. This sensitive period broadly corresponds with the period of

increased movement events in the current study. Considering that sympathetic efferents

typically impact heart rate relatively slowly, on a time–scale of between 1,300 and 2000

milliseconds[60,61,62], timing movement events may capitalize on these effects to more rapidly

adjust autonomic outflow to match demand.

Additionally, it is possible that as agents, we act upon the environment such that relevant

signals appear during optimal periods of the cardiac cycle[63]. It may be that stimuli whose

processing would benefit from information on the state of cardiovascular arousal are more

acutely perceived at systole, when baroreceptor feedback occurs, while stimuli that would not

benefit from such signalling are better processed during diastole. As such, during systole,

fearful faces are more readily detected, and result in greater amygdala activation[64], and race-

related threat appraisal is heightened[65]. Experimentally increasing heartrate (and therefore the

amount of time spent in systole) heightens fear responses but not disgust responses[66].

Meanwhile, during diastole, target shooting[67, 68] and memory for words[69] is more accurate.

The current results suggest we may act upon our environment in such a way as to maximize

this interoceptive distinction.

Respiration was not measured in the current study, but could play a role in the current

findings. A recent study identified a correspondence between the timing of voluntary

movement and the respiratory cycle, with movement initiated more frequently during

expiration[70]. Expiration coincides with a decrease in heart rate, compared to the increase

observed during inspiration. Therefore, movements made during inspiration are more likely

to coincide with heartbeats than those made during expiration. We have greater voluntary

control over respiratory than cardiac events, control that could be leveraged to better govern

cardiac timing. Holding one's breath or breathing out slowly when performing a precise

movement may be one way of decreasing the chances of a heartbeat occurring at the crucial

moment.

In the present study, we not only observed a relationship between the cardiac cycle and action

execution, but also observed a similar relationship for action observation. Observation of

another may produce resonance of the mechanisms involved in executing that action.

Activation may remain sub-threshold for actually producing the motor output, however

associated somotovisceral changes may be produced[28]. Afferent interoceptive feedback

resulting from these changes may in turn support inferences about the causes of the other's

behavior. As such, it is possible that inference about the mental states of others is supported

by a form of interoceptive resonance or mirroring. Such a mechanism might explain how, in

the present data, action observation resulted in a similar movement-cardiac rhythm to that

seen in the individual executing the action. As the observer's motor system predicts the

actor's next movements, comparable interoceptive-conditions are simulated, much like action

observation results in an increase in excitability in the muscles required to perform the

observed action[15].

The mutual prediction of each other's' actions is thought to facilitate joint coordinated action

and the achievement of shared goals[71-73]. There is evidence to suggest that shared

representations of action form automatically even when no joint action is required and it

would be more effective to ignore the other person[74]. The results found here suggest the

possibility that cardiac-movement synchronicity may support the interpersonal coordination

that underlies joint action, through physiological attunement. Such coordination may serve as

"social glue", fostering feelings of group belonging and closeness[75]. A number of findings

appear to support this. Professional interviewers' judgements of trainees' competence have

been found to correlate positively with the trainees' convergence of speech rate and response

latency[76]. The degree of convergence appears to vary based on personality traits and social

standing. For example, subjects who scored higher on a measure of the need for social

approval converged more to their partner's speech intensity and pause length than subjects

who scored lower[77]. Moreover, greater language convergence is observed towards

occupational superiors than subordinates, in that foremen converge more to the language of

managers. Increased physiological covariation, in the form of galvanic skin response (GSR)

has been observed in dyads who either like or dislike each other, compared to those who hold

neutral opinions of each other[78]. Future work should seek to investigate whether inter-dyad

factors such as inter-individual liking and physical similarity to one's partner, or individual

differences such as personality traits, are predictive of the cardiac-movement coordination

observed here.

Previous investigations have suggested that the influence of interoceptive signals on behavior

may be mediated by the individual's degree of conscious awareness of those signals[69,79].

However, it is unclear whether the present effects rely on conscious control of behavior, or

reflexive processes below the level of conscious awareness. In a previous study investigating

the effect of the cardiac cycle on timing preferences for actively sampling the environment,

operationalized as the time subjects chose to press a key prompting a visual stimulus relative

to their cardiac phase, no effect of inter-individual differences in awareness of interoceptive

signals was observed[63]. If the action observation effects observed here are consciously

mediated, it raises the possibility that a greater awareness of cardiac signals might facilitate

greater social mimicry. Future research could test the possibility that inter-individual

variability in the action observation effect reported here is related to social competencies and

is mediated by interoceptive awareness.

Important questions remain regarding precisely under what conditions cardiac-movement

coordination occurs. Understanding this may aid in interpreting previous findings that report

no relationship between the heart and movement[80]. At the level of a single subject, it is likely

that not all centrally mediated sensorimotor events show a phased relationship with the heart.

It is perhaps noteworthy that in the present study, many of the subjects reported during

debriefing that they found replicating the sequence of movements challenging. The cognitive

demands of the task may be an essential difference between the present design and other

studies that report no relationship between heart and movement. In dyads, possible mediators

include the history of interpersonal interaction, speech coordination, the structure of the task

and the dynamics of turn-taking, and the degree of behavioral coordination between the

agents[39]. Different patterns of kinematic and autonomic coordination have been reported.

Under some circumstances, a leader-follower pattern is observed[32,34], while in other

scenarios, convergence appears to be more bi-directional[35-37]. It would be interesting for

future research to explore how the heart rates and kinematics of individuals in a dyad change

over time. A plausible prediction is that over the course of a trial, the observer's heartbeats

become increasingly linked to the timing of the actor's movements. Finally, research suggests

an association between elevated blood pressure and reduced social competence[81,82]. As

baroreceptor activity is tied to blood pressure, the current findings raise the possibility that

reduced cardiac mirroring may underlie these findings.

In summary, these findings demonstrate a relationship between the timing of motor events

and the cardiac cycle in both action execution and observation. These results suggest that

interoceptive signals from the heart and self-paced movement are intrinsically linked. The

observation of this phasic relationship both when executing and observing action raises the

possibility that this relationship may be driving interpersonal coordination effects, which

have been found to foster successful social interaction.

Materials and Methods

Subjects

A total of twenty-six healthy adult subjects with normal or corrected-to-normal vision were recruited. Subjects were aged between 19 and 36 years (mean = 24.92, *SD* = 4.78). Fourteen were female and 12 were male. Twenty-four subjects reported being right-handed, and two reported being left-handed. Data was collected in two stages. In the first stage, due to practical reasons, subjects performed the dyadic behavioural turn-taking task with an experimenter as their partner (n = 12). In stage two, the remaining subjects were recruited in pairs and performed the task together (n = 14). For analysis, data were pooled (see Supplementary Materials for the results reported separately). Ethical approval of the experimental protocol was granted by the Ethics Committee of University College London, and the methods were carried out in accordance with the Declaration of Helsinki. All subjects gave written informed consent before testing.

Materials

Subjects were seated in pairs at a table in front of a Dell laptop computer. Stimuli were presented using MATLAB (version 7.8.0, Mathworks Inc., MA, USA) with the Cogent 2000 toolbox (www.vislab.ucl.ac.uk/cogent_2000.php). Three electrodes were affixed to the abdomens of subjects as per the standard Lead II configuration, to record their ECG. On the table were four touch-sensitive pads and a marble, which the subjects used to execute a series of memorized movements.

Procedure

Once seated comfortably, the experimenter affixed three disposable or washable electrodes to each subject. They were instructed that they would view movement sequences on the screen (see Figure 1A) and would have to replicate them in turns afterwards using the marble and

touch-sensitive pads in front of them. They were also instructed not to talk to or interact with

their partner for the duration of the experiment. On the screen, subjects viewed an animation

of a sequence of six movements (see Figure 1B). The screen showed three white circles

arranged in a triangle. One circle briefly turned red before returning to white to indicate the

sequence. The order of the sequence was pseudo-randomized with the only constraint being

that the same circle could not appear two or more times in a row. Subjects viewed the

sequence twice and were instructed to memorize it. Subjects were informed that the ECG

measures their heart, but were not made aware of the purpose of the experiment or that the

timing of their movements was being recorded.

On the table were four touch-sensitive pads and a marble. Three of the pads were arranged in

a triangle and corresponded to the circles that appeared on the screen. The fourth pad was

placed in the middle and was the home pad where the marble was to be returned to at the end

of each movement sequence and from where each movement sequence commenced. On each

trial both subjects in the pair had to attempt to replicate the sequence, one after the other. To

do this, subjects were instructed to move the marble between the touch sensitive pads in the

order they believed to be the same as in the initial video (see Figure 1C). Subjects were

instructed to make the series of movements at a speed that was comfortable to them.

The order of which subject executed first and which second was counter-balanced across

blocks for each pair. That is, turn-taking occurred by block, where the same subject took the

first turn for a whole block, and then they swapped. Which subject was instructed to go first

was alternated, so that no subject took the first turn for more than one block in a row. For 12

subjects there were two blocks of ten trials in total. In one block one subject would go first

for all ten trials, and in the other the other subject would go first for all ten trials. For the

remaining 14 subjects there were four blocks of ten trials in total. In two blocks one subject

would go first for all ten trials, and in the other two the other subject would go first for all ten

trials. ECGs were recorded throughout the duration of each block.

For 18 subjects the ECG was recorded using an Active 2 Biosemi amplifier. The ECG was

recorded at 2048 Hz. For the remaining seven subjects, the ECG was recorded using a CED

1401 in Spike2 at a sampling rate of 1000 Hz. The time at which the marble was placed on

the touch-sensitive pad was also recorded. The experiment lasted approximately 40 minutes.

Afterwards, subjects were debriefed about the purpose of the experiment and their experience

of taking part, and were given the opportunity to ask the experimenter any questions.

Analysis

For all subjects the three electrode ECG recordings were transformed offline by linear

subtraction into two sets of bipolar recordings, which in turn were averaged to produce a

single ECG recording for each subject. For all subjects, the time of the peak of the R-wave of

each heartbeat was calculated from the ECG. First, the ECG data were high pass filtered at

0.01 Hz to remove any linear drift. Second, a threshold was determined from the data to

isolate the R-wave and the time point of local maximum for each suprathreshold peak was

calculated. For each subject, this time point was determined at 1000 Hz (i.e., to millisecond

accuracy) and in this way all data were then encoded at the same sampling rate. For each

block the time the touch-sensitive pads were contacted was extracted from the event channel

and also resampled at 1000 Hz where necessary. For one subject there was a failure of the

touch-sensitive pads and this subject was excluded from further analysis, leaving 25 subjects

for data analysis.

Analysis of movements and phase of the cardiac cycle

To address whether or not there was a statistical relationship between the time the marble was

placed on the touch-sensitive pad and the phase of cardiac cycle two different analyses were

performed, one in the time domain and one in the phase domain.

Time domain analysis

In the time domain the proportion of motor events that occurred in four time windows

relative to the R-peak were calculated. The four time windows were -100–100 ms, 100 –300

ms, 300 –500 ms and 500 –700 ms, that is, centered around 0, 200, 400 and 600 ms

respectively (Figure 3A). The proportion of motor events that occurred in these windows was

calculated for both executed and observed events. To determine if there was any effect of

time on the proportion of events that occurred a repeated measures ANOVA was run with

two factors, Condition (execution and observation) and Time 0, 200, 400, & 600). Significant

effects were further investigated with post-hoc t-tests, Bonferroni corrected for multiple

comparisons).

Phase domain analysis

In the phase domain, we calculated the phase of the movement event as a function of the R-R

interval[65]. Circular statistics were employed in order to exploit the repeating nature of the

cardiac cycle. To this end, for each action endpoint the time of both the preceding and

proceeding R-wave was calculated and the phase of the action endpoint was calculated as a

function of the R-R interval in which it occurred. For example, for an R-R interval of time t_R,

where the movement event occurred at time t_e the phase, in degrees, of that event was

calculated as t_e/T_R x 360. For each subject, the mean phase was then calculated for the

execution and observation blocks separately using circular averaging. This resulted in two

mean phases per subject, one for each condition (execution and observation). We then tested,

separately for each condition, whether these phases differed from uniformity using Rayleigh

tests. In addition, we also calculated the distribution of phases in a circular histogram, where

each bin was a 20^{th} of a circle. These data were normalized so each bin represented the

proportion of total events for each subject for each condition. These were smoothed by

averaging over three neighbouring bins, the central bin and the two neighbouring bins. The

first analysis shows the uniformity of the mean phase across subjects but gives no indication

of the degree of uniformity of the phases for each subject. This is shown by the second

analysis.

References

1. Coleman, W. M. The psychological significance of bodily rhythms. *J. Comp. Psychol.*, **1,** 213. (1921).

2. Anders, P. Über den individuellen Eigenrhythmus beim menschlichen Gange und seine Beziehungen zum Rhythmus der Herz- und Atemtätigkeit. *Pflügers Archiv*, **220,** 287 -299. (1928).

3. Lacey, J. I., Kagan, J., Lacey, B. C., & Moss, H. A. The visceral level: Situational determinants and behavioral correlates of autonomic response patterns. *Expression of the Emotions in Man*, **9**. (1963).

4. Lacey, J. I., & Lacey, B. C. Some autonomic-central nervous system interrelationships. *Physiological Correlates of Emotion*, 205-227. (1970).

5. Birren, J. E., Cardon, P. V., & Phillips, S. L. Reaction time as a function of the cardiac cycle in young adults. *Science*, **140**. (1963).

6. Jennings, J. R., van der Molen, M. W., Brock, K., & Somsen, R. J. M. On the synchrony of stopping motor responses and delaying heartbeats. *J. Exp. Psychol. Hum. Percept. Perform.*, **18,** 422-436. (1992).

7. Rae, C. L. et al. Response inhibition on the stop signal task improves during cardiac contraction. *Sci. Rep.*, **8,** 9136. (2018).

8. Lacey, B. C., & Lacey, J. I. Two-way communication between the heart and the brain: Significance of time within the cardiac cycle. *Am. Psychol.*, **33,** 99. (1978).

9. Brown, G. L., & Eccles, J. C. The action of a single vagal volley on the rhythm of the heart beat. J. Physiol., **82,** 211-241. (1934).

10. Levy, M. N., & Martin, P. J. Neural control of the heart. In *Physiology and Pathophysiology of the Heart* (pp. 337-354). Boston, MA: Springer. (1984).

11. Dong Jr, E., & Reitz, B. A. Effect of timing of vagal stimulation on heart rate in the dog. *Circ. Res.*, **27,** 635-646. (1970).

12. Levy, M. N., Martin, P. J., Iano, T., & Zieske, H. Effects of single vagal stimuli on heart rate and atrioventricular conduction. *Am. J. Physiol.*, **218,** 1256-1262. (1970).

13. Palmer, C. E., & Tsakiris, M. Going at the heart of social cognition: is there a role for interoception in self-other distinction?. *Curr. Opin. Psychol.*, **24,** 21-26. (2018).

14. Shah, P., Catmur, C., & Bird, G. From heart to mind: linking interoception, emotion, and theory of mind. *Cortex*, **93,** 220. (2017).

15. Gallese, V., & Goldman, A. Mirror neurons and the simulation theory of mind-reading. *Trends Cogn. Sci.*, **2,** 493-501. (1998).

16. Rizzolatti, G., Fogassi, L., & Gallese, V. (2001). Neurophysiological mechanisms underlying the understanding and imitation of action. *Nat. Rev. Neurosci.*, **2**, 661. (2001).

17. Press, C., Heyes, C., & Kilner, J. M. Learning to understand others' actions. *Biol. Lett.*, **7,** 457-460. (2011).

18. Kilner, J. M., Friston, K. J., & Frith, C. D. Predictive coding: an account of the mirror neuron system. *Cogn. Process.*, **8,** 159-166. (2007).

19. Ondobaka, S., Kilner, J., & Friston, K. The role of interoceptive inference in theory of mind. *Brain Cogn.,* **112,** 64-68. (2015).

20. Joffily, M., & Coricelli, G. Emotional valence and the free-energy principle. *PLoS Comp. Biol.*, **9,** e1003094. (2013).

21. Seth, A. K., & Critchley, H. D. Extending predictive processing to the body: emotion as interoceptive inference. *Behav. Brain Sci.*, **36,** 227-228. (2013).

22. Barrett, L. F., & Simmons, W. K. Interoceptive predictions in the brain. *Nat. Rev. Neurosci.*, **16,** 419. (2015).

23. Seth, A. K., & Friston, K. J. Active interoceptive inference and the emotional brain. *Philos. Trans. R. Soc. B: Biol. Sci.*, **371,** 20160007. (2016).

24. Montagna, M., Cerri, G., Borroni, P., & Baldissera, F. Excitability changes in human corticospinal projections to muscles moving hand and fingers while viewing a reaching and grasping action. *Eur. J. Neurosci.*, **22,** 1513-1520. (2005).

25. Borroni, P., & Baldissera, F. Activation of motor pathways during observation and execution of hand movements. *Soc. Neurosci.*, **3,** 276-288. (2008).

26. Decety, J., Jeannerod, M., Germain, M., & Pastene, J. Vegetative response during imagined movement is proportional to mental effort. *Behav. Brain Res.*, **42,** 1-5. (1991).

27. Collet, C., Di Rienzo, F., Hoyek, N., & Guillot, A. Autonomic nervous system correlates in movement observation and motor imagery. *Front. Hum. Neurosci.*, **7,** 415. (2013).

28. Paccalin, C., & Jeannerod, M. Changes in breathing during observation of effortful actions. *Brain Res.*, **862**, 194-200. (2000).

29. Hove, M. J., & Risen, J. L. It's all in the timing: Interpersonal synchrony increases affiliation. *Soc. Cogn.*, **27**, 949-960. (2009).

30. Bernier, F. J., & Rosenthal, R. (1991). Interpersonal coordination: Behavior matching and interpersonal synchrony. In R. S. Feldman & B. Rime (Eds), *Fundamentals of nonverbal behavior* (pp. 401-432). Cambridge, England: Cambridge University Press.

31. Semin, G. R., & Cacioppo, J. T. Grounding social cognition: Synchronization, coordination, and co-regulation. In G. R. Semin & E. R. Smith (Eds.), *Embodied grounding: Social, cognitive, affective and neuroscientific approaches* (pp. 119-147). Cambridge, England: Cambridge University Press. (2008).

32. Condon, W. S., & Ogston, W. D. Speech and body motion synchrony of the speaker-hearer. *The Perception of Language*, 150-184. (1971).

33. Hadar, U., Steiner, T. J., & Rose, F. C. Head movement during listening turns in conversation. *J. Nonverbal Behav.*, **9**, 214-228. (1985).

34. LaFrance, M. Posture mirroring and rapport. In M. Davis (Ed.), *Interaction Rhythms: Periodicity in Communicative Behavior,* (pp. 279-298). New York, NY: Human Scientific Press. (1982).

35. Shockley, K., Santana, M. V., & Fowler, C. A. Mutual interpersonal postural constraints are involved in cooperative conversation. *J. Exp. Psychol. Hum. Percept. Perform.*, **29,** 326. (2003).

36. Dale, R., Fusaroli, R., Duran, N. D., & Richardson, D. C. Chapter 2 - The self-organization of human interaction. *Psychol. Learn. Motiv*., **59,** 43-95. (2013).

37. Fusaroli, R., Bjørndahl, J. S., Roepstorff, A., & Tylén, K. A. Heart for Interaction: Shared Physiological Dynamics and Behavioral Coordination in a Collective, Creative Construction Task. *J. Exp. Psychol. Hum. Percept. Perform.*, **42,** 1297. (2016).

38. Konvalinka, I et al. Synchronized arousal between performers and related spectators in a fire-walking ritual. *Proc. Natl. Sci. U. S. A.*, **108,** 8514-8519. (2011).

39. Levenson, R. W., & Gottman, J. M. Marital interaction: physiological linkage and affective exchange. *J. Pers. Soc. Psychol.*, **45,** 587. (1983).

40. Levenson, R. W., & Ruef, A. M. Empathy: A physiological substrate. *J. Pers. Soc. Psychol.*, **63,** 234. (1992).

41. Henning, R. A., Boucsein, W., & Gil, M. O. Social-physiological compliance as a determinant of team performance. *Int. J. Psychophysiol.*, **40,** 221-232. (2001).

42. Elkins, A. N., Muth, E. R., Hoover, A. W., Walker, A. D., Carpenter, T. L., & Switzer, F. S. Physiological compliance and team performance. *Appl. Ergon.*, **40,** 997-1003. (2009).

43. Henning, R. A., & Korbelak, K. T. Social-psychophysiological compliance as a predictor of future team performance. *Psychologia*, **48,** 84-92. (2005).

44. Strang, A. J., Funke, G. J., Russell, S. M., Dukes, A. W., & Middendorf, M. S. Physio-behavioral coupling in a cooperative team task: Contributions and relations. *J. Exp. Psychol: Hum. Percept. Perform.*, **40,** 145-158. (2014).

45. Noy, L., Levit-Binun, N., & Golland, Y. Being in the zone: physiological markers of togetherness in joint improvisation. *Front. Hum. Neurosci.*, **9,** 187. (2015).

46. Stafford, R. W., Harris, W. S., & Weissler, A. M. Left ventricular systolic time intervals as indices of postural circulatory stress in man. *Circulation*, **41**, 485-492. (1970).

47. Hill, D. W., & Merrifield, A. J. (1976). Left ventricular ejection and the Heather index measured by non□invasive methods during postural changes in man. *Acta Anaesthesiol. Scand.*, **20**, 313-320. (1976).

48. Ring, C., Liu, X., & Brener, J. Cardiac stimulus intensity and heartbeat detection: Effects of tilt☐induced changes in stroke volume. *Psychophysiol.*, **31**, 553-564. (1994).

49. Treisman, M., Faulkner, A., Naish, P. L., & Brogan, D. The internal clock: Evidence for a temporal oscillator underlying time perception with some estimates of its characteristic frequency. *Percept.*, **19,** 705-742. (1990).

50. McCraty, R., Atkinson, M., Tomasino, D., & Bradley, R. T. (2009). The Coherent Heart Heart-Brain Interactions, Psychophysiological Coherence, and the Emergence of System-Wide Order. *Integral Review: A Transdisciplinary & Transcultural Journal for New Thought, Research, & Praxis*, **5**. (2009).

51. Mancia, G., & Mark, A. L. Arterial baroreflexes in humans. In J. T. Shepherd & F. M. Abboud (Eds.), *Handbook of Physiology, Section 2: The Cardiovascular System* (pp. 755-793). Bethesda, MD: American Physiological Society. (1983).

52. Koch, E. Die reflektorische selbststeuerung des kreislaufes. In *Ergebnisse der Kreislaufforschung*. E. Kisch (Ed.). Dresden: Steinkopff. (1931).

53. Salomon, R. et al. The insula mediates access to awareness of visual stimuli presented synchronously to the heartbeat. *J. Neurosci.*, **36**, 5115-5127. (2016).

54. Allen, M., Levy, A., Parr, T., & Friston, K. J. In the Body's Eye: The Computational Anatomy of Interoceptive Inference. *BioRxiv*, 603928. (2019).

55. Van der Molen, M. W., R. J. M. Somsen, and J. F. Orlebeke. The rhythm of the heart beat in information processing. *Advance. Psychophysiol.*, **1**, 1-88. (1985).

56. van der Molen, M. W., Bashore, T. R., Halliday, R., & Callaway, E. Chronopsychophysiology: Mental chronometry augmented by psychophysiological time markers. J. R. Jennings & M. G. H. Coles (Eds.). *Handbook of cognitive psychophysiology: Central and autonomic nervous system approaches*. NJ, USA: John Wiley & Sons. (1991).

57. Somsen, R.J.M., Van der Molen, M.W., Jennings, J.R. & Orlebeke, J.F. Response initiation, not completion, seems to alter cardiac cycle length. *Psychophysiol.*, **22**, 319-325. (1985).

58. Jennings, J. R., Van der Molen, M. W., & Terezis, C. Primary bradycardia and vagal inhibition as two manifestations of the timing of vagal influence on the heart beat. *J. Psychophysiol.*, **1**, 361–374. (1987).

59. Jennings, J.R., &Wood, C.C. Cardiac cycle time effects on performance, phasic cardiac responses, and their intercorrelation in choice reaction time. *Psychophysiol.*, **14**, 297-307. (1977).

60. Berger, R.D., Saul, J.P., & Cohen, R.J. Transfer function analysis of autonomic regulation. I. Canine atrial rate response. *Am. J. Physiol.-Heart C.*, **256**, 142-152. (1989).

61. Salata, J.J., & Zipes, D.P. Autonomic nervous system control of heart rate and atrioventricular nodal conduction. *Reflex control of the circulation.* FL, USA: CRC Press. (1991).

62. Warner, H.R., & Russell Jr, R.O. Effect of combined sympathetic and vagal stimulation on heart rate in the dog. *Circ. Res.*, **24**, 567-573. (1969).

63. Kunzendorf, S. et al. Active information sampling varies across the cardiac cycle. *Psychophysiology*, e13322. (2019).

64. Garfinkel, S. N. et al. Fear from the heart: Sensitivity to fear stimuli depends on individual heartbeats. *J. Neurosci.*, **34,** 6573-6582 (2014).

65. Azevedo, R. T., Garfinkel, S. N., Critchley, H. D., & Tsakiris, M. Cardiac afferent activity modulates the expression of racial stereotypes. *Nat. Commun.*, **8,** 13854. (2017).

66. Pezzulo, G., Iodice, P., Barca, L., Chausse, P., Monceau, S., & Mermillod, M. Increased heart rate after exercise facilitates the processing of fearful but not disgusted faces. *Sci. Rep.*, *8*, 398. (2018).

67. Bothwell, S., Donne, B., & Andrews, J. F. Does heart rate or phase of cardiac cycle at triggering affect the performance of elite marksmen? *Journal of Physiology – London,* **501,** 6. (1997).

68. Bellamy, M., Collins, D., Holmes, P., & Loze, G. Shot patterns in ECG recordings for elite air-pistol shooters. *Journal of Sports Sciences, Conference Communications*, **17,** 48-49. (1999).

69. Garfinkel, S. N. et al. What the heart forgets: Cardiac timing influences memory for words and is modulated by metacognition and interoceptive sensitivity. *Psychophysiology*, **50,** 505-512. (2013).

70. Park, H.D., Barnoud, C., Trang, H., Kannape, O.A., Schaller, K. & Blanke, O. Breathing is coupled with voluntary action and the cortical readiness potential. *Nat. Commun.*, **11**, 1-8. (2020).

71. Verfaillie, K., & Daems, A. Representing and anticipating human actions in vision. *Vis. Cogn.*, **9,** 217-232. (2002).

72. Flanagan, J. R., & Johansson, R. S. Action plans used in action observation. *Nat.*, **424,** 769. (2003).

73. Kilner, J. M., Vargas, C., Duval, S., Blakemore, S. J., & Sirigu, A. Motor activation prior to observation of a predicted movement. *Nat. Neurosci.*, **7,** 1299. (2004).

74. Sebanz, N., Knoblich, G., & Prinz, W. Representing others' actions: just like one's own? *Cogn.*, **88,** B11-B21. (2003).

75. Lakin, J. L., Jefferis, V. E., Cheng, C. M., & Chartrand, T. L. The Chameleon Effect as Social Glue: Evidence for the Evolutionary Significance of Nonconscious Mimicry. *J. Nonverbal Behav.*, **27,** 145–162. (2003).

76. Street Jr, R. L. Speech convergence and speech evaluation in fact-finding interviews. *Hum. Commun. Res.*, **11,** 139-169. (1984).

77. Natale, M. Convergence of mean vocal intensity in dyadic communication as a function of social desirability. *J. Pers. Soc. Psychol.,* **32,** 790. (1975).

78. Kaplan, H. B., Burch, N. R., & Bloom, S. W. Physiological covariation and sociometric relationships in small peer groups. In P. H. Leiderman & D Shapiro (Eds.), *Psychobiological Approaches to Social Behavior*. Stanford, CA: Stanford University Press. (1964).

79. Dunn, B. D. et al. Listening to your heart: How interoception shapes emotion experience and intuitive decision making. *Psychol. Sci.*, **21**, 1835-1844. (2010).

80. Wallot, S., Mitkidis, P., McGraw, J. J., & Roepstorff, A. Beyond synchrony: joint action in a complex production task reveals beneficial effects of decreased interpersonal synchrony. *PloS one*, **11,** e0168306. (2016).

81. Diamond, E.L. The role of anger and hostility in essential hypertension and coronary heart disease. *Psychol. Bull.*, **92**, 410. (1982).

82. Sommers-Flanagan, J. & Greenberg, R.P. Psychosocial variables and hypertension: a new look at an old controversy. *J. Nerv. Ment. Dis,* **177**, 15-24. (1989).

Acknowledgements

This work was supported by a PhD studentship awarded to E. R. Palser from the Economic and Social Research Council (ESRC) and the Medical Research Council (MRC). A. Fotopoulou was supported by a 'European Research Council Starting Investigator Award' [ERC-2012-STG GA313755].

Author Contributions

Designed the research ERP, AF, JMK; collected the data ERP, JG; analyzed the data ERP, JMK; wrote the manuscript ERP, JMK; provided comments on a draft and approved the final manuscript AF, JG.

Competing interests

The authors declare that they have no competing interests

Figures

Figure 1: Task design. Panel A: Subjects were seated side by side in front of a laptop computer; Panel B: an example of the sequence of movements subjects were required to learn; Panel C: how they would accurately replicate the sequence in (B) using the touch-sensitive containers.

Figure 2: Relationship between cardiac and movement parameters. Panels A & B show the relationship between the mean movement time and the mean R-R interval for each subject, during action execution (A) and action observation (B). In each panel the circles show the data from each subject and the solid line shows the line of best fit. Panel C shows the relationship between the mean movement time for the two people that comprised the dyad. The significant positive relationship is consistent with the hypothesis that subjects imitated the movement speed of the other. In each panel, each circle shows the data from one dyad.

Figure 3: Relationship between time of movement culmination and cardiac cycle in the time domain. Panel A shows one cycle of the cardiac cycle (R-R) and shows the four time windows used for the analysis. Panel B shows the mean proportion of events that occurred in each of the four windows shown in A. The dots show the mean data for each subject, the bar is the mean across subjects.

Figure 4: Relationship between time of movement culmination and cardiac cycle in the phase domain. Panels A-F show the relationship between the time of the movement event (action end point) with respect to the phase of the R-R interval. In each panel the R peaks occurred at 0^o and the phase through the R-R interval cycles clockwise around the circle. On the left-hand side of figure, panels A, C & E show the data for executed actions and on the right-hand side, panels B, D & F show the data for observed actions. In A & B the circles show the mean phase for each subject. White circles are from those dyads where the ECG was recorded from both subjects   and the grey circles show the data where ECG was recorded from only one subject. The circular histogram show the frequency of the data as a function of phase. The solid black arrow shows the circular mean phase across subjects. Panels C & D depict the distribution of the mean normalized circular histograms across subjects. The solid black line

depicts the mean phase distribution and the dotted lines show the standard error around this mean (SEM). Panels E & F show the same data in C & D but unwrapped. The solid grey line shows the null distribution (a uniform distribution). Error bars represent the SEM. * show phases where the proportion of events are significantly different from a uniform distribution.