Genomic data and multi-species demographic modelling uncover past

hybridization between currently allopatric freshwater species

Sofia L. Mendes[1], Miguel P. Machado[1], Maria M. Coelho[1*], Vitor C. Sousa[1*&]

[1] cE3c – Centre for Ecology, Evolution and Environmental Changes, Faculdade de

Ciências da Universidade de Lisboa, Lisbon, Portugal

[*]These authors contributed equally to this work

[&]corresponding author: vmsousa@fc.ul.pt

Abstract

Evidence for ancient interspecific gene flow through hybridization has been reported in
many animal and plant taxa based on genetic markers. The study of genomic patterns
of closely related species with allopatric distributions allow to assess the relative
importance of vicariant isolating events and past gene flow. Here, we investigated the
role of gene flow in the evolutionary history of four closely related freshwater fish
species with currently allopatric distributions in western Iberian rivers - *Squalius*
carolitertii, *S. pyrenaicus*, *S. torgalensis* and *S. aradensis* - using a population
genomics dataset of 23 562 SNPs from 48 individuals, obtained through genotyping by
sequencing (GBS). We uncovered a species tree with two well differentiated clades: (i)
S. carolitertii and *S. pyrenaicus*; and (ii) *S. torgalensis* and *S. aradensis.* By using D-
statistics and demographic modelling based on the site frequency spectrum, comparing
alternative demographic scenarios of hybrid origin, secondary contact and isolation, we
found that the *S. pyrenaicus* North lineage is likely the result of an ancient hybridization
event between *S. carolitertii* (contributing ~84%) and *S. pyrenaicus* South lineage
(contributing ~16%), consistent with a hybrid speciation scenario. Furthermore, in the
hybrid lineage we identify outlier loci potentially affected by selection favouring genes
from each parental lineage at different genomic regions. Our results suggest that
ancient hybridization can affect speciation and that freshwater fish species currently in
allopatry are useful to study these processes.

Introduction

How populations diverge and ultimately originate new species has always intrigued evolutionary biologists. Speciation is assumed to occur due to a continuous reduction in gene flow until reproductive isolation is achieved and populations maintain phenotypic and genetic distinctiveness (Coyne and Orr, 2004). This process might occur between geographically isolated populations in a strictly allopatric scenario, or in sympatry driven by ecological factors (Bush, 1975; Schluter, 2009). In either case (and the continuum in-between), genetic differences are expected to accumulate so that reproduction between the initial populations is no longer possible (Coyne and Orr, 2004).

Understanding the role of gene flow is key to explain how biological diversity is generated and species are formed. Historically, gene flow was considered a homogenizing force, mixing gene pools and counteracting the accumulation of divergence between populations. However, based on population genetics data, evidence for divergence in the presence of gene flow has accumulated (reviewed in Feder *et al.*, 2012). For example, ecologically driven divergent selection has been shown to promote population divergence in the absence of barriers to gene flow in several animal taxa, including insects, fish and mammals (Linn *et al.*, 2003; Gagnaire *et al.*, 2013; Pfeifer *et al.*, 2018).

While the role of hybridization between species in evolution has been discussed for quite some time (e.g. Anderson and Stebbins, 1954; Stebbins, 1959; Grant, 1981), advances in our ability to generate population genomics data and perform fine-scale analysis (Davey *et al.*, 2011; da Fonseca *et al.*, 2016; Payseur and Rieseberg, 2016) have shown that hybridization is more widespread than previously thought, especially in the animal kingdom (reviewed in Taylor and Larson, 2019). Hybrid individuals might be unfit in their parentals' habitat or have reduced viability and fertility due to hybrid incompatibilities (Bateson-Dobzhansky-Muller incompatibilities (Dobzhansky, 1937; Muller, 1942)). However, mixing old alleles into new combinations through hybridization can fuel adaptation and speciation (e.g. Rieseberg *et al.*, 2003; Hermansen *et al.*, 2014; Wallbank *et al.*, 2016; Richards and Martin, 2017; Svardal *et al.*, 2020; reviewed in Marques *et al.*, 2019). In this context, it is crucial to determine the timing and mode of gene flow. If hybridization between two species occurs at the time of origin of a third new reproductively isolated lineage, it is likely a case of hybrid speciation (reviewed in Mallet, 2007; Abbott *et al.*, 2013; Vallejo-Marín and Hiscock, 2016). Instead, if hybridization occurs without producing a new lineage but results in introgression of genetic material through backcrossing, it is a case of secondary contact.

Due to their outstanding diversification and remarkable adaptive radiations, freshwater fish have been widely used as model systems to study speciation (e.g. reviewed in Bernardi, 2013 and Seehausen and Wagner, 2014). Different events fuelled this diversity, including transitions from marine to freshwater habitats (e.g. Jones et al. 2012; Terekhanova et al. 2014), adaptation to extreme environments (e.g. Pfenninger et al. 2015), differentiation along water depth clines (e.g. Barluenga et al. 2006; Gagnaire et al. 2013) and changes in the configuration of rivers and lakes over geological time (e.g. Sousa-Santos *et al.*, 2019). Freshwater fish also stand out as a vertebrate group with high hybridization rates (Hubbs, 1955; Wallis *et al.*, 2017). For example, hybridization events that promoted rapid adaptive radiations within lake

systems have been documented (e.g. Meier *et al.*, 2017; Svardal *et al.*, 2020), as well
as several instances of hybridization between native and invasive species (e.g. Nolte *et*
al., 2006; Meraner *et al.*, 2013).

In this work, we analyse the role of gene flow during the speciation process of
four species of Iberian endemic chubs found in Western Iberian rivers: *Squalius*
carolitertii, *Squalius pyrenaicus*, *Squalius torgalensis* and *Squalius aradensis* (Figure
1). As obligatory freshwater fish, their evolutionary history is intertwined with the
geomorphological rearrangements of the river systems they inhabit. These species are
distributed along a latitudinal environmental cline with increasing temperatures and
propensity for drought from north to south, reflecting the transition from an Atlantic to a
Mediterranean climate type (Gasith and Resh, 1999; Jesus *et al.*, 2017). Two of them
have rather wide distribution ranges: *Squalius carolitertii* (Doadrio, 1988) inhabits
northern rivers down to the Mondego basin, while *Squalius pyrenaicus* (Gunther, 1868)
has a more southern distribution (Tagus, Sado, Guadiana and south-eastern basins)
(Coelho *et al.*, 1995, 1998). Contrastingly, the two other species are confined to small
basins in the southwest: *Squalius torgalensis* (Coelho *et al.*, 1998) inhabits the Mira
basin and *Squalius aradensis* (Coelho *et al.*, 1998) basins in the extreme southwestern
area (e.g. Arade) (Coelho *et al.*, 1998). These two last species are "Critically
Endangered" and *S. pyrenaicus* is "Endangered" in the Portuguese Red List (Cabral *et*
al., 2005) and "Vulnerable" in the Spanish Red List (Doadrio, 2002).

Although these species have allopatric distributions (Figure 1) associated with
different river basins, studies based on mitochondrial and nuclear markers found
phylogenetic incongruences between the two types of markers, which have been
interpreted as possible past hybridization but remain unresolved (Brito *et al.*, 1997;
Waap *et al.*, 2011; Sousa-Santos *et al.*, 2019; Perea *et al.*, 2020). Thus, this work had
three major goals: (i) to characterize the genome-wide patterns of genetic
differentiation and reconstruct the species tree for these four *Squalius* species; (ii) to
investigate the role of gene flow in the evolution of these species, particularly to assess
if previously reported incongruent mitochondrial and nuclear phylogenies can be
explained by incomplete lineage sorting; and (iii) to perform demographic modelling to
compare alternative scenarios, and to date and quantify past gene flow. To achieve
these goals, we obtained genome-wide single nucleotide polymorphisms (SNPs)
across the four species through Genotyping by Sequencing (GBS) and applied
methods to quantify genetic differentiation, reconstruct the evolutionary relationship
between species and test for evidence of gene flow. Furthermore, we performed
demographic modelling to compare alternative models of isolation, hybrid origin and
secondary contact, accounting for drift and shared ancestry (i.e., incomplete lineage
sorting). We found support for a potential case of hybrid speciation in *S. pyrenaicus*,
uncovering genomic regions that might have been under selection in the hybrid.

Methods

Sampling and sequencing

We sampled 65 individuals from 8 different locations, as displayed in Figure 1,
including at least one sampling location from a representative river basin per species.
S. carolitertii individuals were collected from the Mondego basin (n=10). In the northern
part of *S. pyrenaicus* distribution individuals were collected from the Ocreza river
(n=10) and Canha stream (n=10), both tributaries of the Tagus basin. Specimens were
also collected in the Lizandro basin (n=10). From here on, we use "*S. pyrenaicus*
North" to refer to *S. pyrenaicus* from Ocreza, Canha and Lizandro. In the southern part
of the distribution, *S. pyrenaicus* was sampled in the Guadiana (n=2) and Almargem
(n=8) basins, which we refer to as "*S. pyrenaicus* South". *S. torgalensis* individuals
were collected in the Mira basin (n=10) and *S. aradensis* individuals from the Arade
basin (n=5). GPS coordinates of the sampling locations and fishing licenses from the
Portuguese authority for conservation of endangered species [ICNF - Instituto de
Conservação da Natureza e das Florestas] can be found in Table S1.

Fish were collected by electrofishing (300V, 4A) and total genomic DNA was
extracted from fin clips using a phenol-chloroform protocol adapted from Taggart *et al.*,
1992 and quantified using Qubit® 2.0 Fluorometer (Live Technologies). Samples were
subjected to a paired-end Genotyping by Sequencing (GBS) protocol (adapted from
Elshire *et al.*, 2011), performed at Beijing Genomics Institute (BGI, www.bgi.com). DNA
was sent to the facility mixed with DNAstable Plus (Biomatrica) to preserve it at room
temperature during shipment. Briefly, upon arrival, DNA was fragmented using the
restriction enzyme ApeKI and the fragments were amplified after adaptor ligation
(Elshire *et al.*, 2011). The resulting library was sequenced using Illumina Hiseq2000.

Multispecies SNP dataset

We assessed the quality of the sequences using FastQC
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and MultiQC (Ewels *et al.*,
2016). We then used *process_radtags* from Stacks v2.2 (Rochette *et al.*, 2019) to trim
all reads to 82 base pairs and discard reads with uncalled bases and low quality
scores, using the default settings for window size (0.15x read length) and base quality
threshold (10 Phred score). Given the absence of a reference genome for any of the
species in this study, we built a catalog of all loci using the *denovo* approach from
Stacks v2.2 (Rochette *et al.*, 2019). We first tested different M and n parameter (see
definition below) values for catalog construction using a subset of representative
individuals (Figure S1), following the approach recommended by Paris et al. 2017. We
based our parameter selection on accounting for differences between four species with
an estimated divergence time of 14Mya (Sousa-Santos *et al.*, 2019) and low genetic
diversity (Almada and Sousa-Santos, 2010). Using all individuals for catalog
construction, we required a minimum depth of coverage of 4x for every stack (set of
exactly matching reads) (m=4) and allowed a maximum of 2 mismatches between
initially formed stacks to be considered putative alleles of the same locus within each
individual (M=2). We then allowed a maximum of 4 differences between stacks from
different individuals (n=4) for them to be considered one locus on the catalog. Given

the possibility that forward and reverse sequences of the same fragment were treated
as different loci, similar reads within the catalog were clustered using CD-HIT-EST from
the CD-HIT v4.7 package (Li and Godzik, 2006; Fu *et al.*, 2012) with a word length of 6
and a sequence identity threshold of 0.85. The reads from each individual were aligned
against the catalog using BWA-MEM from BWA v0.7.17-r1188 (Li, 2013) with default
parameters. We sorted the output alignments and removed unmapped reads using
Samtools v1.10 (Li and Durbin, 2009) and evaluated the alignments for each individual
using Picard v2.18.13 (http://broadinstitute.github.io/picard/) (Table S2). To call
genotypes for each individual at each site and identify SNPs we used the method
implemented in Freebayes v1.2.0 (Garrison and Marth, 2012), discarding reads and
bases with low quality and without using Hardy-Weinberg equilibrium priors (-p 2 --min-
mapping-quality 30 --min-base-quality 20 --hwe-priors-off).

To discard sites and genotypes that are likely due to sequencing or mapping
errors and to maximize the number of individuals with data, we applied a series of
filters using a combination of options from VCFtools v0.1.15 (Danecek *et al.*, 2011) and
BCFtools v1.6 (Li *et al.*, 2009). First, we kept only SNPs present in all sampling sites in
at least 50% of the individuals. Second, genotypes with a depth of coverage (DP)
outside of ¼ to 4 times the individual median DP and SNPs with excess of
heterozygotes when pooling all individuals were removed. Third, individuals with more
than 50% missing data were removed. Finally, only SNPs with a minor allele frequency
(MAF) larger than 0.01 were kept (MAF≥0.01).

Global patterns of genetic differentiation

To quantify the levels of differentiation between sampling locations, we
calculated the pairwise F_{ST} using the Hudson estimator (Hudson *et al.*, 1992). We
investigated fine population structure using individual based methods: principal
component analysis (PCA) and ADMIXTURE v1.3 (Alexander *et al.*, 2009). PCA was
performed using package LEA (Frichot and François, 2015) in RStudio v1.1.383 and R
v3.4.4. We used ADMIXTURE to determine the ancestry proportion of each individual
from a specified number of clusters (K), testing values of K between 1 and 8,
performing 100 independent runs for each value of K. We identified the best value of K
as the one with the lowest 5-fold cross-validation error. For 2≤K≤5, we assessed
similarity across the 100 replicates using the Greedy algorithm implemented in
CLUMPP v1.1.2 (Jakobsson and Rosenberg, 2007).

Relationship between species and populations

We reconstructed a graph describing the relationships between the populations
using TreeMix v1.13 (Pickrell and Pritchard, 2012). We explored models with no
migration and up to two migration events. Since there is only one individual from *S.*
pyrenaicus Guadiana in the final dataset (see Results), *S. pyrenaicus* South is
represented only by *S. pyrenaicus* Almargem. The position of the root was not
specified, thus the resulting trees are unrooted.

We also constructed a maximum likelihood phylogeny of the individuals using
IQ-TREE v1.6.12 (Nguyen *et al.*, 2015). We used vcf2phylip v2.0 (Ortiz, 2019) to
convert the VCF file into PHYLIP format. We used ModelFinder (Kalyaanamoorthy *et*
al., 2017) implemented on IQ-TREE to determine the best substitution model, limiting
the search to models with ascertainment bias correction (+ASC). According to the
corrected Akaike Information Criterion (AIC), the best model was GTR+F+ASC+R2,
which was used to construct the phylogeny with 5 000 standard non-parametric
bootstrap replicates. The resulting best tree was visualized with midpoint-rooting in
FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Effect of linked SNPs

To verify if the results were influenced by potential linkage between SNPs, we
produced a dataset minimizing the number of linked SNPs by dividing the catalog into
blocks of 200 base pairs, which is larger than the mean size of GBS loci, and sampling
one SNP per block. We selected the SNP with less missing data per block. Using this
"single SNP" dataset, we repeated analyses that might be affected when SNPs are not
independent (PCA, ADMIXTURE and TreeMix).

Detection of introgression between *S. carolitertii* and *S. pyrenaicus*

To test for past introgression between *S. carolitertii* and *S. pyrenaicus* we used
the D-statistic (Durand *et al.*, 2011) with four different combinations of populations,
using both *S. aradensis* and *S. torgalensis* as outgroup in all cases. First, we tested
either *S. carolitertii* or S. *pyrenaicus* South as the possible sources of introgression into
S. pyrenaicus North. To test for geographical cline in admixture proportions in *S.*
pyrenaicus North, we tested if the northernmost sampling site of *S. pyrenaicus* (Ocreza
- Figure 1) showed more shared alleles with *S. carolitertii* than the other *S. pyrenaicus*
North (Lizandro and Canha). Finally, we also considered all *S. pyrenaicus* North as
sister populations and *S. carolitertii* as the potential source of introgressed genes.

As in the TreeMix analysis, we used *S. pyrenaicus* Almargem as the *S.*
pyrenaicus South population. Significance of D-statistic values was assessed using a
block jackknife approach (Soraggi *et al.*, 2018), dividing the dataset into 25 blocks with
a similar number of SNPs. These computations were performed in RStudio v1.1.383
and R v3.4.4 using custom scripts.

If introgression between populations occurred in the relatively recent past, we
would expect individuals within the same population to show different degrees of
introgression. To test this hypothesis, we calculated the D-statistic for each individual of
S. pyrenaicus North for the same scenarios as above.

Demographic modelling of the divergence of *S. carolitertii* and *S. pyrenaicus*

We compared alternative divergence scenarios of *S. pyrenaicus* and *S.*
carolitertii using the composite likelihood method based on the site frequency
spectrum (SFS) implemented in *fastsimcoal2* v2.6 (Excoffier *et al.*, 2013).

First, we compared the fit of three models with and without admixture to the
observed SFS (see Figure S2 for parameters inferred in each model). We compared a
model that assumes *S. pyrenaicus* North received a contribution alpha (α) from *S.*
pyrenaicus South and 1-alpha (1-α) from *S. carolitertii* at the time of the split (Figure
S2A) with models without admixture (Figure S2B and Figure S2C). Importantly, models
without admixture account for incomplete lineage sorting as they consider that
populations share a common ancestry (different split times), and that each population
has a specific effective size. To ensure models have the same number of parameters,
in models without admixture we allowed for the possibility of a bottleneck associated
with the split of *S. pyrenaicus* North, mimicking a founder effect. Models B and C
without the possibility of bottlenecks were also considered. Second, we compared
three other models to distinguish between a hybrid origin of *S. pyrenaicus* North and
two models of secondary contact (Fig. S2D-F).

To obtain the observed minor allele frequency spectrum without missing data,
we built the pairwise 2D-SFS by sampling 3 individuals from *S. carolitertii* and *S.*
pyrenaicus South, and 4 individuals from *S. pyrenaicus* North sampled from Ocreza
and Canha. We excluded Lizandro since it is an independent river basin, and its
removal maximized the number of SNPs in the SFS. To sample individuals without
missing data, we used the initial dataset without the MAF filter and divided it into blocks
of 200bp. For each block, we sampled the individuals from each population with the
least missing data keeping only the sites with data across all individuals. Since the SFS
is affected by the depth of coverage, only genotypes with a depth of coverage >10x
were used (Nielsen *et al.*, 2011). This resulted in an observed SFS with 8 758 SNPs.
For each model we performed 100 independent runs with 100 cycles, approximating
the SFS with 100 000 coalescent simulations. Given that we used the SFS without
monomorphic sites, all parameters were scaled in relation to a reference effective size,
which was arbitrarily set to be the effective size (*Ne*) of *S. carolitertii*. To convert the
relative divergence times estimated into absolute time in million years (Mya), we
assumed a generation time of 3 years for these species (Magalhães *et al.*, 2003;
Almada and Sousa-Santos, 2010).

We then used the AIC to compare models. Since composite likelihoods
computed with linked sites and pairwise 2D-SFS tend to overestimate the likelihood
due to non-independence of data points (Excoffier *et al.*, 2013), we built a 3D-SFS
minimizing linked sites. We obtained 1 000 bootstrap joint 3D-SFS MAF by resampling
one site from each GBS locus, assuming they are independent. We obtained the
expected 3D-SFS for each model based on 100 000 coalescent simulations according
to the parameters that maximized the composite likelihood. To account for noise of the
approximation, we averaged the expected 3D-SFS of 10 runs. We then computed the
likelihood of the 1 000 bootstrap unlinked 3D-SFS for each model. For each bootstrap
replicate, we computed the AIC and relative likelihood of each model, following
Excoffier *et al.*, 2013.

Detection of outlier loci in *S. pyrenaicus* North

To identify variation from the parental species (*S. carolitertii* or *S. pyrenaicus*
South) that was likely selected in the introgressed *S. pyrenaicus* North we scanned for

outlier loci using pairwise F_{ST} (Hudson *et al.*, 1992). We assumed that selection
favouring alleles from a target parental (P1) in the potential hybrid population (H) would
lead to SNPs where (i) the target parental and hybrid have low differentiation ($F_{ST(H,P1)}$ <
quantile 0.05 $F_{ST(H,P1)}$); (ii) the hybrid and the other parental (P2) have high
differentiation ($F_{ST(H,P2)}$ > quantile 0.95 $F_{ST(H,P2)}$); and (iii) the differentiation between the
hybrid and the other parental is higher than between the two parental populations
($F_{ST(H,P2)}$ > quantile 0.95 $F_{ST(P1,P2)}$ between parentals). This is because under neutrality
we would expect the same level of differentiation between the minor parental (i.e., *S.*
pyrenaicus South) and both the major parental (i.e., *S. carolitertii*) and the hybrid
lineage (*S. pyrenaicus* North). Focusing on *S. carolitertii* and *S. pyrenaicus* (n=34), we
applied a MAF filter of MAF>0.05 and kept only sites with more than 20% of data per
population. We used only catalog loci with more than 2 SNPs and a mean distance
between SNPs higher than 9. This analysis was repeated considering individuals from
the three sampling locations of *S. pyrenaicus* North separately to identify outlier regions
shared between them. Calculations were done in in RStudio v1.1.383 and R v3.4.4
using custom scripts. Finally, we blasted the identified outlier catalog loci against the
NCBI database using the default settings of BLASTN (Zhang *et al.*, 2000) and BLASTX
(Altschul, 1997) v2.11.0+.

Results

Multispecies SNP dataset

After the initial processing, we obtained a mean of 5 759 264 high-quality reads
per individual (Table S2). The catalog comprised 524 911 loci with a mean length of
160bp. Mean depth of coverage per sample was 59.7x. The filters applied resulted in
17 individuals with more than 50% of missing data, which were removed. The final
dataset comprised 23 562 SNPs with 36.98% missing data and 48 individuals, as
follows: *S. carolitertii* (n=10), *S. pyrenaicus* Ocreza (n=6), *S. pyrenaicus* Lizandro
(n=4), *S. pyrenaicus* Canha (n=7), *S. pyrenaicus* Almargem (n=6), *S. pyrenaicus*
Guadiana (n=1), *S. torgalensis* (n=9), and *S. aradensis* (n=5).

Global patterns of genetic differentiation

Overall, the highest levels of genetic differentiation are between the two
southwestern species (*S. torgalensis* and *S. aradensis*) and the two more widely
distributed species (*S. carolitertii* and *S. pyrenaicus*) (F_{ST}≥0.294; Table 1). The lowest
levels of genetic differentiation were found between *S. pyrenaicus* North sampling
locations and between them and *S. carolitertii* (F_{ST}≤0.139). The latter were lower than
between *S. pyrenaicus* North and *S. pyrenaicus* South (F_{ST}≥0.171). Interestingly, the
levels of differentiation found between both *S. carolitertii* and *S. pyrenaicus* North and
S. pyrenaicus South are comparable to those found between *S. torgalensis* and *S.*
aradensis.

The first three principal components (PCs) of the PCA explain ~25% of the
variation (Figure S3). PC1 (Figures 2A.1 and S3C) explains ~14% of the variance and
clearly separates two groups: (i) *S. carolitertii* and *S. pyrenaicus*, and (ii) *S. aradensis*

and *S. torgalensis*. This is consistent with the higher pairwise F_{ST} values obtained
between these two groups. PC2 explains ∼6% of the variance and separates *S.*
aradensis from *S. torgalensis* (Figure 2A). Finally, PC3 explains ∼5% separates *S.*
pyrenaicus South from a cluster formed by *S. carolitertii* and *S. pyrenaicus* North
(Figure 2A.2 and S3C).

Regarding the ADMIXTURE analysis, K=2 achieved the lowest cross-validation
(Figure S4). The results were consistent across the 100 runs for 2≤K≤5 (mean G'=1.0
for K=2, K=4 and K=5 and mean G'≥0.9974 for K=3). For the best value of K (K=2),
one of the clusters includes the two southwestern species (*S. torgalensis* and *S.*
aradensis), while the other comprises *S. carolitertii* and *S. pyrenaicus* (Figure 2B.1).
This clustering mimics the separation created by PC1 in the PCA. For K=3, a third
cluster formed by *S. pyrenaicus* South appears (Figure 2B.2). For K=4 the two
southwestern species (*S. torgalensis* and *S. aradensis*) are differentiated into different
clusters (Fig 2B.3). Interestingly, in these last two K values, one of the individuals from
the Canha sampling location of *S. pyrenaicus* North shows a significant proportion from
the *S. pyrenaicus* South cluster. Two *S. pyrenaicus* North individuals also show a small
proportion of the *S. aradensis* cluster in K=4, which may be due to ancestral
polymorphism. K=5 does not fit the data as well (mean cross validation much higher
than for 2≤K≤4, Figure S4), and fails to separate *S. carolitertii* from *S. pyrenaicus*
North.

Relationship between species and populations

The unrooted population tree obtained with TreeMix (Figure 3) shows a clear
separation between two groups: (i) *S. aradensis* and *S. torgalensis,* and (ii) *S.*
carolitertii and *S. pyrenaicus*. Within the group of *S. carolitertii* and *S. pyrenaicus*, we
found two main lineages: *S. pyrenaicus* South (here represented by *S. pyrenaicus*
Almargem) and *S. carolitertii* and *S. pyrenaicus* North, which is consistent with pairwise
F_{ST}, PCA and ADMIXTURE results. One of the two migration edges inferred is between
S. pyrenaicus North and *S. pyrenaicus* South.

Regarding the maximum likelihood phylogeny of individuals (Figure S5), the
topology of the tree is consistent with the one obtained for the populations with
TreeMix, showing two well supported clades. It strongly supports the paraphyly of *S.*
pyrenaicus with respect to *S. carolitertii*, with two well supported subclades with
individuals of: (i) *S. carolitertii* and *S. pyrenaicus* North, and (ii) *S. pyrenaicus* South.

Effect of linked SNPs

The dataset with one SNP per block of 200bp comprised 2 607 SNPs with
∼32.4% missing data. The results of PCA, ADMIXTURE and TreeMix analysis were
overall consistent with those from the initial dataset of 23 562 SNPs (Figures S6-S8).

Detection of introgression between *S. carolitertii* and *S. pyrenaicus*

When considering *S. carolitertii* as the potential source of introgression, we found significant positive D-statistic values indicating an excess of sites where *S. pyrenaicus* North shares the same allele with *S. carolitertii*, independently of the outgroup used (Figure 4A). This could indicate that *S. carolitertii* and *S. pyrenaicus* North share a more recent common ancestor, in agreement with TreeMix and phylogeny results (Figures 3 and S5). However, when considering *S. pyrenaicus* South as the potential source of introgression we also found significant positive D-statistic values, indicating an excess of sites where both *S. pyrenaicus* share the same allele (Figure 4B), in agreement with the TreexMix migration edge (Figure 3). This indicates that the relationship between the species is not described by a bifurcating tree with a more recent common ancestry of *S. carolitertii* and *S. pyrenaicus* North, as we would expect D=0 if that was the case.

We find no evidence of a geographical cline of admixture with *S. carolitertii* along the *S. pyrenaicus* North distribution (Figure S9), with its northern most sampling location (Ocreza – location 2 on Figure 1) showing no signs of sharing more alleles with *S. carolitertii* than with other *S. pyrenaicus* North populations. Thus, the inexistence of a geographical cline of admixture and the consistent signal across different *S. pyrenaicus* North sampling locations in Figure 4 suggest a scenario of introgression between *S. carolitertii* and the ancestor of *S. pyrenaicus* North prior to the divergence of the different *S. pyrenaicus* North populations.

In the case of recent introgression events, we would expect to find differences in the D-statistic among individuals from a given population. However, when we computed the D-statistic by individual (Figure S10 and Table S4), we found limited variation among different individuals from the same population, suggesting that introgression events are likely pre-dating the divergence of populations.

Demographic modelling of divergence of *S. carolitertii* and *S. pyrenaicus*

First, we compared models with admixture (Figure 5A) to models of bifurcating trees without gene flow accounting for incomplete lineage sorting, i.e. different topologies, times of split and effective population sizes (Figures 5B-C). The model with admixture reached a higher likelihood (Table S5), approximately 9.00 log10 units higher than the best bifurcating tree model with the topology supported by the individual-based phylogeny (Figure S5), i.e., with a more recent ancestor of *S. carolitertii* and *S. pyrenaicus* North (Figure 5B). This result was not affected by the number of parameters, as bifurcating tree models with or without an extra effective size parameter to allow for a potential bottleneck (see Methods) had lower likelihood values than the admixture model (Tables S5, S6A, S7 and S8). This supports that *S. pyrenaicus* North likely experienced introgression and that its relationship with *S. carolitertii* and *S. pyrenaicus* South cannot be explained by a simple bifurcating tree, in agreement with TreeMix and D-statistics (Figures 3-4). Next, we explored models to distinguish between a scenario of hybrid origin of *S. pyrenaicus* North (Figure 5D) and secondary contact (Figure 5D-F). The parameter estimates were similar and consistent across models with admixture (Figure 5A, D-F), indicating (i) a major contribution of 80-84% from *S. carolitertii* into *S. pyrenaicus* North; (ii) a relatively old divergence of the parental lineages (~15 times older than admixture times); and (iii) relatively larger

effective sizes in ancestral lineages (Figures 5A, 5D-F). To convert relative parameters
into years we considered different effective sizes of *S. carolitertii*, which was
considered as a reference (Tables S6A-B). To compare models, we computed the
relative likelihoods based on AIC for 1000 bootstrap datasets with a single SNP per
GBS locus. Model D achieved the higher relative likelihood, irrespective of the
threshold used to pool SFS entries (Figure S11). The best model D is compatible with a
"hybrid origin", inferring that at time of admixture *S. pyrenaicus* North received ~84%
from *S. carolitertii* (major parental) and the remaining ~16% from *S. pyrenaicus* South
(minor parental).

Detection of outlier loci in *S. pyrenaicus* North

We identified 10 outlier loci with low F_{ST} between *S. carolitertii* and *S.*
pyrenaicus North and high F_{ST} between *S. pyrenaicus* North and *S. pyrenaicus* South
(Figure 6A – red triangles). These correspond to alleles from *S. carolitertii* which might
have been selected in *S. pyrenaicus* North. However, when separating *S. pyrenaicus*
North into the three sampling locations, none of these was shared between them
(Figure 6B). Moreover, none produced relevant BLAST hits.

We also identified 6 loci with high F_{ST} between *S. carolitertii* and *S. pyrenaicus*
North and low F_{ST} between *S. pyrenaicus* North and *S. pyrenaicus* South (Figure 6A –
blue triangles). These correspond to alleles from *S. pyrenaicus* South that might have
been selected in *S. pyrenaicus* North. One of the identified catalog loci, shared
between the individuals from the locations "Ocreza" and "Canha" and also identified
when all locations were pooled together (Fig 6C), revelated an alignment match to
vitellogenin, mostly from other cyprinid fish (Table S9). The other 5 outlier loci
produced no relevant BLAST hits.

Discussion

Contrasting role of gene flow in the two species' pairs

The genomic analysis presented here indicates a species tree comprising two
main groups: (i) the one of *S. torgalensis* and *S. aradensis* and (ii) that of *S. carolitertii*
and *S. pyrenaicus*. This is in agreement with phylogenies previously obtained for
mitochondrial (Brito *et al.*, 1997; Sanjur *et al.*, 2003; Mesquita *et al.*, 2007) and different
nuclear markers (Almada and Sousa-Santos, 2010; Waap *et al.*, 2011; Sousa-Santos
et al., 2019; Perea *et al.*, 2020). The results also show strong genetic differentiation
between the two species' pairs (Table 1, Figures 2-3), as formerly reported (Coelho *et*
al., 1995; Almada and Sousa-Santos, 2010), which likely reflects their old divergence
time, estimated to be approximately 14Mya based on phylogenies of mitochondrial and
nuclear genes (Perea *et al.*, 2010; Sousa-Santos *et al.*, 2019).

Gene flow appears to have played a very different role on the evolutionary
history of the two specie's pairs. We found no evidence of gene flow involving *S.*
torgalensis and *S. aradensis*. While we could not calculate D-statistics due to our
sampling scheme, the results of the analyses including the four species (Table 1,
Figures 2-3) did not support interspecific gene flow involving *S. torgalensis* and/or *S.*

aradensis. Thus, for these two species with more restricted distribution areas our
results are compatible with divergence of isolated lineages in allopatry. In contrast, for
the two species with wider distributions (*S. carolitertii* and *S. pyrenaicus*) our results
indicate gene flow was paramount in their evolutionary history, specifically in the *S.*
pyrenaicus North lineage (Figures 4-5). This contrasting role of gene flow between the
two more restricted and the two more widely distributed species might reflect their
historical dispersal routes into different river basins. Our results are consistent with the
hypothesis that the ancestor of the two southwestern species (*S. torgalensis* and *S.*
aradensis) became isolated in southwestern Iberia around 11.6-7.2Ma, while *S.*
carolitertii and *S. pyrenaicus* only reached their current ranges in the last 2.6Ma
(Sousa-Santos *et al.*, 2019), the latter as a consequence of geological changes that led
to the current conformation of the major Iberian river basins (Cunha *et al.*, 2005;
Casas-Sainz and de Vicente, 2009; Pais *et al.*, 2012; Antón *et al.*, 2014). Thus, the
older isolation of river basins in southwestern Iberia probably created less opportunity
for gene flow involving *S. torgalensis* and *S. aradensis*. This is in agreement with
results from other freshwater fish species (Sousa *et al.*, 2010) and amphibians
(Martínez-Solano *et al.*, 2006; Gonçalves *et al.*, 2009), indicating higher genetic
differentiation in this geographic region.

Evidence of hybridization between *S. carolitertii* and *S. pyrenaicus*

Our demographic modelling results indicates that the *S. pyrenaicus* North
lineage is likely the result of hybridization – the best model estimates indicate 84%
contribution from *S. carolitertii* and the remaining 16% from *S. pyrenaicus* South
(Figure 5D and Figure S12). A hybridization scenario is also supported by other
analyses (Figures 3-4). For example, TreeMix infers a common ancestor for *S.*
carolitertii and *S. pyrenaicus* North, together with a migration event between northern
and southern *S. pyrenaicus*, indicating allele sharing that cannot be explained by a
bifurcating tree (Figure 3). The uncovered hybridization provides an explanation for the
incongruences between mitochondrial and nuclear markers reported in previous
studies: while in mtDNA phylogenies all *S. pyrenaicus* had a more recent common
ancestor (Brito *et al.*, 1997; Mesquita *et al.*, 2007), in nuclear phylogenies individuals of
river basins corresponding to *S. pyrenaicus* North shared a more recent ancestor with
S. carolitertii (Waap *et al.*, 2011; Sousa-Santos *et al.*, 2019; Perea *et al.*, 2020).

The D-statistics results indicate that the hybridization event predates the
differentiation of the three *S. pyrenaicus* North sampling locations, since the
introgression signal is shared between them (Figure 4). Given that Lizandro (Figure 1 –
location 3) is an independent river basin with no connection to the Tagus basin (to
which the other two *S. pyrenaicus* North sampling locations belong), the most likely
explanation is that the hybridization event predates the colonization of this independent
river. Thus, the combination of the D-statistics with the best demographic model
indicates that *S. pyrenaicus* North likely originated through hybridization and posteriorly
dispersed to its current range. Geological data indicates that prior to reaching their
current conformation, the major Iberian river basins were endorheic, with the transition
to exorheism occurring in the last 2.6Ma (Cunha *et al.*, 2005; Casas-Sainz and de
Vicente, 2009; Pais *et al.*, 2012; Antón *et al.*, 2014). Thus, it is possible that this
transition increased the chances of contact between previously isolated lineages.

Another possibility is that these "lake like" endorheic basins allowed for lineages to mix,
with posterior dispersal of resulting hybrids.

The origin of *S. pyrenaicus* North through a single hybridization event followed
by isolation, as modelled in Figure 5D, is consistent with a scenario of hybrid
speciation. Most authors agree that homoploid hybrid speciation (HHS), or
recombinational speciation, occurs when hybridization plays a key role in establishing a
new reproductively isolated lineage without changes in ploidy (Anderson and Stebbins,
1954; Grant, 1981; Mallet, 2007; Mavárez and Linares, 2008; Abbott *et al.*, 2013).
Schumer *et al.*, 2014 establish three criteria to demonstrate HHS: (1) hybrid lineages
must be reproductively isolated from the parental ones that originated them; (2) the
genome of the hybrids must show evidence of hybridization; (3) reproductive isolation
must be demonstrated to be a consequence of hybridization. Our results demonstrate
S. pyrenaicus North meets the second criterion. Additionally, it has a non-overlapping
distribution with *S. carolitertii* and *S. pyrenaicus* South, thus being currently
reproductively isolated from its parentals through their allopatric distributions, and
ploidy is maintained between the three lineages (2n=50) (Collares-Pereira *et al.*, 1998).
Yet, we currently lack information on the possible role of hybridization on the
establishment of reproductive isolation to meet the three criteria. Indeed, very few
biological systems clearly meet all the criteria outlined above. A notable example are
the *Helianthus* sunflowers, in which hybridization between two species produced
independent linages reproductively isolated from the parentals, with clear evidence of
hybridization in their genomes, that can occupy dry and saline environments that the
parents cannot, as a direct result of hybridization (Rieseberg *et al.*, 1995, 1996, 2003).
In freshwater fish, some taxa have been proposed as being the result of HHS (e.g.
DeMarais *et al.*, 1992), although prior to the definition of the above criteria.

Potential implications of hybridization in *S. pyrenaicus* North

We estimated relative times indicating that the parental *S. carolitertii* and *S.*
pyrenaicus South lineages diverged much earlier than the hybridization event that
originated the *S. pyrenaicus* North lineage, inferring that the time of hybridization is
0.07 of the divergence time (Figure 5D, Table S6B). In this scenario, *S. pyrenaicus*
North might have benefited from the combination of different alleles from the two
parentals, already filtered out by selection over many generations. Yet, the
accumulation of genetic differentiation between the two parental lineages may also
have led to genetic incompatibilities when the two genomes mixed. Our estimates
indicate an asymmetric and larger contribution from *S. carolitertii* at the time of
hybridization (~84%). Such asymmetries in admixture proportions have been reported
in other freshwater fish, albeit in cases of recent hybridization, e.g., swordtail fish
(Schumer *et al.*, 2018) and sticklebacks (Marques, Lucek, *et al.*, 2019). Here, this
asymmetry could reflect the relative proportions of *S. carolitertii* and *S. pyrenaicus*
South in the initial reproductive pool that originated *S. pyrenaicus* North. Another
possibility is that incompatibilities might have been resolved towards the *S. carolitertii*
lineage. Notably, at mitochondrial markers *S. pyrenaicus* North is more closely related
with *S. pyrenaicus* South (Brito *et al.*, 1997; Mesquita *et al.*, 2007), despite the larger
nuclear contribution of *S. carolitertii*. This can be due to neutral stochasticity, but it

might also be related to cito-nuclear incompatibilities or behavioural mechanisms
favouring mating between *S. caralitertii* males and *S. pyrenaicus* South females.

560   Despite the asymmetry in the contribution of both parental lineages, there is
variation in differentiation along the genome between the hybrid and parental lineages
(Figures 6 and S12). This variation reflects the stochasticity of neutral processes (e.g.,
drift and gene flow), but it also provides indirect evidence for the action of selection
removing incompatibilities and/or favouring specific parental alleles (adaptive
introgression). We identified 6 outlier regions compatible with selection of alleles from
the minor parental lineage in *S. pyrenaicus* North, one of them corresponding to
vitellogenin. Vitellogenin is an egg yolk precursor that is usually only expressed in
females during oogenesis (Tyler *et al.*, 1996), although its expression can also be
triggered in males of different fish species in response to exogenous oestrogen
exposure (Harries *et al.*, 1997; Flammarion *et al.*, 2000; Van Den Belt *et al.*, 2003).
Given its role in egg yolk formation, *vitellogenin* is of great importance to egg-laying
organisms like fish. One possibility is that this genomic region was selected and
maintained from the minor parental (*S. pyrenaicus* South) due to adaptive
introgression, although the underlying selective pressures remain unclear. However,
we cannot discard the hypothesis that genetic incompatibilities were involved and thus
constrained the maintenance of this region from *S. pyrenaicus* South.

577   While we could not annotate other outlier genomic regions of interest in *S.*
pyrenaicus North, the geographic distribution of this lineage at an intermediate between
the Atlantic and Mediterranean climate types provides future opportunity to investigate
the potential adaptive role of hybridization. In particular, increasing temperatures and
propensity for seasonal drought from north to south (the latter leading to isolation of the
fish in the deeper water sections during the summer, when large portions of the
riverbed dry out) (Gasith and Resh, 1999; Magalhães *et al.*, 2003; Jesus *et al.*, 2017)
might impose strong selective pressures. This is consistent with previous studies that
found (i) signatures of positive selection based on dN/dS on circadian genes likely
associated with thermal adaptation in *Squalius* species (Moreno *et al.*, 2021), and (ii)
differences in gene expression between species inhabiting Atlantic (*S. caralitertii*) and
Mediterranean climates (*S. torgalensis*) when exposed to increasing temperatures
(Jesus *et al.*, 2016, 2017). Thus, its hybrid origin might confer some advantage to *S.*
pyrenaicus North at intermediate environmental conditions, as described in other
organisms. For example, in *Saccharomyces* yeast laboratory crosses between *S.*
cerevisiae and *S. paradoxus* produced transgressive F2 hybrids that slightly increased
their environmental range and outcompeted their parentals across the cline (Stelkens
et al., 2014). In freshwater fish, hybrids between low and high elevation swordtail
species experience higher fitness at intermediate conditions (Culumber *et al.*, 2012).

Implications for conservation

598   *S. pyrenaicus* populations, like other endemic Iberian cyprinids, have been
impacted by habitat degradation, due to the construction of dams and water extraction
for irrigation, and the introduction of exotic species. Consequently, *S. pyrenaicus* is
listed as "Endangered" in the Portuguese Red List (Cabral *et al.*, 2005) and as
"Vulnerable" in the Spanish Red List (Doadrio, 2002). Our work provides evidence of

two very distinct gene pools, each with its own evolutionary history, within the analysed
S. pyrenaicus distribution: the one inhabiting the Tagus and small Atlantic coastal
basins (*S. pyrenaicus* North) and the one distributed along the Guadiana and small
Mediterranean basins (*S. pyrenaicus* South). Thus, this difference should be
considered in future management efforts.

Conclusions

Our study revealed a species tree with two well differentiated groups in which
gene flow had a contrasting role: (i) *S. torgalensis* and *S. aradensis* and (ii) *S.*
carolitertii and *S. pyrenaicus*. We find no evidence of past gene flow in the first,
consistent with divergence in allopatry. In contrast, we uncover past hybridization
between *S. carolitertii* and *S. pyrenaicus*, originating the *S. pyrenaicus* North lineage,
with ~84% contribution from *S. carolitertii* and ~16% from *S. pyrenaicus* South. Our
estimates indicate that a single hybridization event followed by isolation, consistent with
a scenario of hybrid speciation, is more likely than a secondary contact scenario.
However, the scenario might be more complex, e.g., involving further changes in the
past effective sizes. In the future, whole genome sequencing data and more extensive
sampling would be helpful to explore the sources of the unbalanced estimated
admixture proportions in *S. pyrenaicus* North and evaluate the adaptive potential of this
hybridization, given the contrasting environments of the parental lineages.

This work adds to the growing list of examples where hybridization has been
uncovered, describing a study system suitable for future work on the processes and
consequences of hybridization. Much remains to be understood regarding how
hybridization shapes the genomes of species and genetic incompatibilities are
resolved, its relation to the evolution of reproductive barriers between hybrid and
parental lineages, and its role in adaptation.

Acknowledgements

We thank Tiago F. Jesus for the help in the preparation of the samples. This work was
funded by the strategic projects UID/BIA/00329/2013 (2015-2018) and
UIDB/00329/2020 granted to cE3c from the Portuguese National Science Foundation –
Fundação para a Ciência e a Tecnologia (FCT). SLM is funded by an FCT scholarship
(SFRH/BD/145153/2019). VCS was funded by FCT (CEECIND/02391/2017 and
CEECINST/00032/2018/CP1523/CT0008) and by EU H2020 program (Marie
Skłodowska-Curie grant 799729). We thank the INCD (https://incd.pt/) for use of their
computing infrastructure, which is funded by FCT and FEDER (project 01/SAICT/2016
nº 022153).

Conflict of interest

The authors declare that they have no conflict of interest.

References

Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJE, Bierne N, *et al.* (2013).
Hybridization and speciation. *J Evol Biol* **26**: 229–246.

Alexander DH, Novembre J, Lange K (2009). Fast model-based estimation of ancestry
in unrelated individuals. *Genome Res* **19**: 1655–1664.

Almada V, Sousa-Santos C (2010). Comparisons of the genetic structure of Squalius
populations (Teleostei, Cyprinidae) from rivers with contrasting histories, drainage
areas and climatic conditions based on two molecular markers. *Mol Phylogenet*
Evol **57**: 924–931.

Altschul S (1997). Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Res* **25**: 3389–3402.

Anderson E, Stebbins GL (1954). Hybridization as an Evolutionary Stimulus. *Evolution*
(N Y) **8**: 378.

Antón L, De Vicente G, Muñoz-Martín A, Stokes M (2014). Using river long profiles and
geomorphic indices to evaluate the geomorphological signature of continental
scale drainage capture, Duero basin (NW Iberia). *Geomorphology* **206**: 250–261.

Barluenga M, Stölting KN, Salzburger W, Muschick M, Meyer A (2006). Sympatric
speciation in Nicaraguan crater lake cichlid fish. *Nature* **439**: 719–723.

Van Den Belt K, Verheyen R, Witters H (2003). Comparison of vitellogenin responses
in zebrafish and rainbow trout following exposure to environmental estrogens.
Ecotoxicol Environ Saf **56**: 271–281.

Bernardi G (2013). Speciation in fishes. *Mol Ecol* **22**: 5487–5502.

Brito RM, Briolay J, Galtier N, Bouvet Y, Coelho MM (1997). Phylogenetic
Relationships within Genus Leuciscus (Pisces , Cyprinidae) in Portuguese Fresh
Waters , Based on Mitochondrial DNA Cytochrome b Sequences. *Mol Phylogenet*

Evol **8**: 435–442.

Bush GL (1975). Modes of Animal Speciation. *Annu Rev Ecol Syst* **6**: 339–364.

Cabral M (coord. ., Almeida J, Almeida P, Dellinger T, Ferrand de Almeida N, Oliveira
M, *et al.* (2005). *Livro Vermelho dos Vertebrados de Portugal*. Instituto da
Conservação da Natureza: Lisboa.

Casas-Sainz AM, de Vicente G (2009). On the tectonic origin of Iberian topography.
Tectonophysics **474**: 214–235.

Coelho MM, Bogutskaya NG, Rodrigues JA, Collares-Pereira MJ (1998). Leuciscus
torgalensis, and L. aradensis, two new cyprinids for Portuguese fresh waters. *J
Fish Biol* **52**: 937–950.

Coelho MM, Brito RM, Pacheco TR, Figueiredo D, Pires AM (1995). Genetic variation
and divergence of Leuciscus pyrenaicus and L. carolitertii (Pisces, Cyprinidae). *J
Fish Biol* **47**: 243–258.

Collares-Pereira MJ, Próspero MI, Biléu RI, Rodrigues EM (1998). Leuciscus (Pisces,
Cyprinidae) karyotypes: Transect of Portuguese populations. *Genet Mol Biol* **21**:
63–69.

Coyne JA, Orr HA (2004). *Speciation*. Sinauer Associates, Inc., Sunderland,
Massachusetts.

Culumber ZW, Shepard DB, Coleman SW, Rosenthal GG, Tobler M (2012).
Physiological adaptation along environmental gradients and replicated hybrid zone
structure in swordtails (Teleostei: Xiphophorus). *J Evol Biol* **25**: 1800–1814.

Cunha PP, Martins AA, Daveau S, Friend PF (2005). Tectonic control of the Tejo river
fluvial incision during the late Cenozoic, in Ródão - Central Portugal (Atlantic
Iberian border). *Geomorphology* **64**: 271–298.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, *et al.* (2011). The
variant call format and VCFtools. *Bioinformatics* **27**: 2156–2158.

Davey JW, Hohenlohe PA, Etter PD, Boone JQ, Catchen JM, Blaxter ML (2011).
Genome-wide genetic marker discovery and genotyping using next-generation
sequencing. *Nat Rev Genet* **12**: 499–510.

DeMarais BD, Dowling TE, Marsh PC, Douglas ME, Minckley WL (1992). Origin of Gila
seminuda (Teleostei : Cyprinidae) through introgressive hybridization :
Implications for evolution and conservation. *Evolution (N Y)* **89**: 2747–2751.

Doadrio I (ed. . (2002). *Atlas y Libro Rojo de los Peces Continentales de España*.
Dirección General de Conservación de la Naturaleza.: Museo Nacional de
Ciencias Naturales, Madrid.

Dobzhansky T (1937). *Genetics and the origin of species*. Columbia University Press:
New York.

Durand EY, Patterson N, Reich D, Slatkin M (2011). Testing for ancient admixture
between closely related populations. *Mol Biol Evol* **28**: 2239–2252.

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, *et al.* (2011). A
robust, simple genotyping-by-sequencing (GBS) approach for high diversity
species. *PLoS One* **6**: 1–10.

Ewels P, Magnusson M, Lundin S, Käller M (2016). MultiQC: Summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**: 3047–
3048.

Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M (2013). Robust
Demographic Inference from Genomic and SNP Data. *PLoS Genet* **9**.

Feder JL, Egan SP, Nosil P (2012). The genomics of speciation-with-gene-flow. *Trends
Genet* **28**: 342–350.

Flammarion P, Brion F, Babut M, Garric J, Migeon B, Noury P, *et al.* (2000). Induction
of fish vitellogenin and alterations in testicular structure: Preliminary results of
estrogenic effects in chub (Leuciscus cephalus). *Ecotoxicology* **9**: 127–135.

da Fonseca RR, Albrechtsen A, Themudo GE, Ramos-Madrigal J, Sibbesen JA,
Maretty L, *et al.* (2016). Next-generation biology: Sequencing and data analysis
approaches for non-model organisms. *Mar Genomics* **30**: 3–13.

Frichot E, François O (2015). LEA: An R package for landscape and ecological
association studies. *Methods Ecol Evol* **6**: 925–929.

Fu L, Niu B, Zhu Z, Wu S, Li W (2012). CD-HIT: Accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**: 3150–3152.

Gagnaire PA, Pavey SA, Normandeau E, Bernatchez L (2013). The genetic
architecture of reproductive isolation during speciation-with-gene-flow in lake
whitefish species pairs assessed by rad sequencing. *Evolution (N Y)* **67**: 2483–
2497.

Garrison E, Marth G (2012). Haplotype-based variant detection from short-read
sequencing. *http://arxiv.org/abs/12073907*.

Gasith A, Resh VH (1999). Streams in Mediterranean Climate Regions: Abiotic
Influences and Biotic Responses to Predictable Seasonal Events. *Annu Rev Ecol
Syst* **30**: 51–81.

Gonçalves H, MartÍnez-Solano I, Pereira RJ, Carvalho B, GarcÍa-ParÍs M, Ferrand N
(2009). High levels of population subdivision in a morphologically conserved
Mediterranean toad (Alytes cisternasii) result from recent, multiple refugia:
Evidence from mtDNA, microsatellites and nuclear genealogies. *Mol Ecol* **18**:
5143–5160.

Grant V (1981). *Plant Speciation*, Second Edi. Columbia University Press.

Harries JE, Sheahan DA, Jobling S, Matthiessen P, Neall P, Sumpter JP, *et al.* (1997).
Estrogenic activity in five United Kingdom rivers detected by measurement of
vitellogenesis in caged male trout. *Environ Toxicol Chem* **16**: 534–542.

Hermansen JS, Haas F, Trier CN, Bailey RI, Nederbragt AJ, Marzal A, *et al.* (2014).
Hybrid speciation through sorting of parental incompatibilities in Italian sparrows.

Mol Ecol **23**: 5831–5842.

Hubbs CL (1955). Hybridization between fish species in nature. *Syst Zool* **4**: 1–20.

Hudson RR, Slatkint M, Maddison WP (1992). Estimation of Levels of Gene Flow From
DNA Sequence Data. *Genetics* **589**: 583–589.

Jakobsson M, Rosenberg NA (2007). CLUMPP: A cluster matching and permutation
program for dealing with label switching and multimodality in analysis of
population structure. *Bioinformatics* **23**: 1801–1806.

Jesus TF, Grosso AR, Almeida-Val VMF, Coelho MM (2016). Transcriptome profiling of
two Iberian freshwater fish exposed to thermal stress. *J Therm Biol* **55**: 54–61.

Jesus TF, Moreno JM, Repolho T, Athanasiadis A, Rosa R, Almeida-Val VMF, *et al.*
(2017). Protein analysis and gene expression indicate differential vulnerability of
Iberian fish species under a climate change scenario (S Rutherford, Ed.). *PLoS*
One **12**: e0181325.

Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, *et al.* (2012). The
genomic basis of adaptive evolution in threespine sticklebacks. *Nature* **484**: 55–
61.

Kalyaanamoorthy S, Minh BQ, Wong TKF, Von Haeseler A, Jermiin LS (2017).
ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nat*
Methods **14**: 587–589.

Li H (2013). Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. *http://arxiv.org/abs/13033997*.

Li H, Durbin R (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**: 1754–1760.

Li W, Godzik A (2006). Cd-hit: A fast program for clustering and comparing large sets
of protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, *et al.* (2009). The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Linn C, Feder JL, Nojima S, Dambroski HR, Berlocher SH, Roelofs W (2003). Fruit
odor discrimination and sympatric host race formation in Rhagoletis. *Proc Natl*
Acad Sci U S A **100**: 11490–11493.

Magalhães MF, Schlosser IJ, Collares-Pereira MJ (2003). The role of life history in the
relationship between population dynamics and environmental variability in two
Mediterranean stream fishes. *J Fish Biol* **63**: 300–317.

Mallet J (2007). Hybrid speciation. *Nature* **446**: 279–283.

Marques DA, Lucek K, Sousa VC, Excoffier L, Seehausen O (2019). Admixture
between old lineages facilitated contemporary ecological speciation in Lake
Constance stickleback. *Nat Commun* **10**.

Marques DA, Meier JI, Seehausen O (2019). A Combinatorial View on Speciation and
Adaptive Radiation. *Trends Ecol Evol* **34**: 531–544.

Martínez-Solano I, Teixeira J, Buckley D, García-París M (2006). Mitochondrial DNA
phylogeography of Lissotriton boscai (Caudata, Salamandridae): Evidence for old,
multiple refugia in an Iberian endemic. *Mol Ecol* **15**: 3375–3388.

Mavárez J, Linares M (2008). Homoploid hybrid speciation in animals. *Mol Ecol* **17**:
4181–4185.

Meier JI, Marques DA, Mwaiko S, Wagner CE, Excoffier L, Seehausen O (2017).
Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat Commun* **8**:
1–11.

Meraner A, Venturi A, Ficetola GF, Rossi S, Candiotto A, Gandolfi A (2013). Massive
invasion of exotic Barbus barbus and introgressive hybridization with endemic
Barbus plebejus in Northern Italy: Where, how and why? *Mol Ecol* **22**: 5295–5312.

Mesquita N, Cunha C, Carvalho GR, Coelho MM (2007). Comparative phylogeography
of endemic cyprinids in the south-west Iberian Peninsula: Evidence for a new
ichthyogeographic area. *J Fish Biol* **71**: 45–75.

Moreno JM, Jesus TF, Coelho MM, Sousa VC (2021). Adaptation and convergence in
circadian-related genes in Iberian freshwater fish. *BMC Ecol Evol* **21**: 38.

Muller H (1942). Isolating mechanisms, evolution, and temperature. *Biol Symp* **6**: 71–
125.

Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015). IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol
Biol Evol* **32**: 268–274.

Nielsen R, Paul JS, Albrechtsen A, Song YS (2011). Genotype and SNP calling from
next-generation sequencing data. *Nat Rev Genet* **12**: 443–451.

Nolte AW, Freyhof J, Tautz D (2006). When invaders meet locally adapted types:
Rapid moulding of hybrid zones between sculpins (Cottus, Pisces) in the Rhine
system. *Mol Ecol* **15**: 1983–1993.

Ortiz EM. (2019). vcf2phylip v2.0: convert a VCF matrix into several matrix formats for
phylogenetic analysis.

Pais J, Cunha PP, Pereira D, Legoinha P, Dias R, Moura D, *et al.* (2012). The
Paleogene and Neogene of Western Iberia (Portugal): A Cenozoic Record in the
European Atlantic Domain. In: *The Paleogene and Neogene of Western Iberia
(Portugal): A Cenozoic record in the European Atlantic domain*, Springer Berlin
Heidelberg: Berlin, Heidelberg, pp 1–138.

Paris JR, Stevens JR, Catchen JM (2017). Lost in parameter space: a road map for
stacks. *Methods Ecol Evol* **8**: 1360–1373.

Payseur BA, Rieseberg LH (2016). A genomic perspective on hybridization and
speciation. *Mol Ecol* **25**: 2337–2360.

Perea S, Böhme M, Zupancic P, Freyhof J, Sanda R, Ozuluğ M, *et al.* (2010).
Phylogenetic relationships and biogeographical patterns in Circum-Mediterranean
subfamily Leuciscinae (Teleostei, Cyprinidae) inferred from both mitochondrial and

nuclear data. *BMC Evol Biol* **10**: 265.

Perea S, Sousa-Santos C, Robalo J, Doadrio I (2020). Multilocus phylogeny and
systematics of Iberian endemic Squalius (Actinopterygii, Leuciscidae). *Zool Scr*
49: 440–457.

Pfeifer SP, Laurent S, Sousa VC, Linnen CR, Foll M, Excoffier L, *et al.* (2018). The
evolutionary history of Nebraska deer mice: Local adaptation in the face of strong
gene flow. *Mol Biol Evol* **35**: 792–806.

Pfenninger M, Patel S, Arias-Rodriguez L, Feldmeyer B, Riesch R, Plath M (2015).
Unique evolutionary trajectories in repeated adaptation to hydrogen sulphide-toxic
habitats of a neotropical fish (Poecilia mexicana). *Mol Ecol* **24**: 5446–5459.

Pickrell JK, Pritchard JK (2012). Inference of Population Splits and Mixtures from
Genome-Wide Allele Frequency Data. *PLoS Genet* **8**.

Richards EJ, Martin CH (2017). Adaptive introgression from distant Caribbean islands
contributed to the diversification of a microendemic adaptive radiation of trophic
specialist pupfishes (J Mallet, Ed.). *PLOS Genet* **13**: e1006919.

Rieseberg LH, Van Fossen C, Desrochers AM (1995). Hybrid speciation accompanied
by genomic reorganization in wild sunflowers. *Nature* **375**: 313–316.

Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, *et al.*
(2003). Major ecological transitions in wild sunflowers facilitated by hybridization.
Science (80-) **301**: 1211–1216.

Rieseberg LH, Sinervo B, Linder CR, Ungerer MC, Arias DM (1996). Role of gene
interactions in hybrid speciation: Evidence from ancient and experimental hybrids.
Science (80-) **272**: 741–745.

Rochette NC, Rivera-Colón AG, Catchen JM (2019). Stacks 2: Analytical methods for
paired-end sequencing improve RADseq-based population genomics. *Mol Ecol*
28: 4737–4754.

Sanjur OI, Carmona JA, Doadrio I (2003). Evolutionary and biogeographical patterns
within Iberian populations of the genus Squalius inferred from molecular data. *Mol*
Phylogenet Evol **29**: 20–30.

Schluter D (2009). Evidence for ecological speciation and its alternative. *Science (80-)*
323: 737–741.

Schumer M, Rosenthal GG, Andolfatto P (2014). How common is homoploid hybrid
speciation? *Evolution (N Y)* **68**: 1553–1560.

Schumer M, Xu C, Powell DL, Durvasula A, Skov L, Holland C, *et al.* (2018). Natural
selection interacts with recombination to shape the evolution of hybrid genomes.
Science (80-) **360**: 656–660.

Seehausen O, Wagner CE (2014). Speciation in Freshwater Fishes. *Annu Rev Ecol*
Evol Syst **45**: 621–651.

Soraggi S, Wiuf C, Albrechtsen A (2018). Powerful inference with the D-statistic on low-
coverage whole-genome data. *G3 Genes, Genomes, Genet* **8**: 551–566.

Sousa-Santos C, Jesus TF, Fernandes C, Robalo JI, Coelho MM (2019). Fish
diversification at the pace of geomorphological changes: evolutionary history of
western Iberian Leuciscinae (Teleostei: Leuciscidae) inferred from multilocus
sequence data. *Mol Phylogenet Evol* **133**: 263–285.

Sousa V, Penha F, Pala I, Chikhi L, Coelho MM (2010). Conservation genetics of a
critically endangered Iberian minnow: Evidence of population decline and
extirpations. *Anim Conserv* **13**: 162–171.

Stebbins GL (1959). The Role of Hybridization in Evolution. *Proc Am Philos Soc* **103**:
231–251.

Stelkens RB, Brockhurst MA, Hurst GDD, Miller EL, Greig D (2014). The effect of
hybrid transgression on environmental tolerance in experimental yeast crosses. *J*
Evol Biol **27**: 2507–2519.

Svardal H, Quah FX, Malinsky M, Ngatunga BP, Miska EA, Salzburger W, *et al.* (2020).
Ancestral hybridization facilitated species diversification in the lake malawi cichlid
fish adaptive radiation. *Mol Biol Evol* **37**: 1100–1113.

Taggart JB, Hynes RA, Prodöuhl PA, Ferguson A (1992). A simplified protocol for
routine total DNA isolation from salmonid fishes. *J Fish Biol* **40**: 963–965.

Taylor SA, Larson EL (2019). Insights from genomes into the evolutionary importance
and prevalence of hybridization in nature. *Nat Ecol Evol* **3**: 170–177.

Terekhanova N V., Logacheva MD, Penin AA, Neretina T V., Barmintseva AE, Bazykin
GA, *et al.* (2014). Fast Evolution from Precast Bricks: Genomics of Young
Freshwater Populations of Threespine Stickleback Gasterosteus aculeatus. *PLoS*
Genet **10**.

Tyler CR, Van Der Eerden B, Jobling S, Panter G, Sumpter JP (1996). Measurement of
vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety
of cyprinid fish. *J Comp Physiol - B Biochem Syst Environ Physiol* **166**: 418–426.

Vallejo-Marín M, Hiscock SJ (2016). Hybridization and hybrid speciation under global
change. *New Phytol* **211**: 1170–1187.

Waap S, Amaral AR, Gomes B, Coelho MM (2011). Multi-locus species tree of the
chub genus Squalius (Leuciscinae: Cyprinidae) from western Iberia: New insights
into its evolutionary history. *Genetica* **139**: 1009–1018.

Wallbank RWR, Baxter SW, Pardo-Diaz C, Hanly JJ, Martin SH, Mallet J, *et al.* (2016).
Evolutionary Novelty in a Butterfly Wing Pattern through Enhancer Shuffling. *PLoS*
Biol **14**: 1–16.

Wallis GP, Cameron-Christie SR, Kennedy HL, Palmer G, Sanders TR, Winter DJ
(2017). Interspecific hybridization causes long-term phylogenetic discordance
between nuclear and mitochondrial genomes in freshwater fishes. *Mol Ecol* **26**:
3116–3127.

Zhang Z, Schwartz S, Wagner L, Miller W (2000). A greedy algorithm for aligning DNA
sequences. *J Comput Biol* **7**: 203–214.

Figure 1 – Distribution range of the four *Squalius* species in Portuguese rivers and sampling locations: (1) Mondego; (2) Ocreza; (3) Lizandro; (4) Canha; (5) Guadiana; (6) Almargem; (7) Mira; (8) Arade. Triangles correspond to the GPS coordinates of the sampling locations.

Figure 2 – Global patterns of genetic differentiation. (A) Principal Component Analysis. (A.1) PC1 and PC2; (A.2) PC2 and PC3. Each point corresponds to one individual. **(B) Individual ancestry proportions inferred with ADMIXTURE.** (B.1) K=2; (B.2) K=3; (B.3) K=4; (B.4) K=5. Each vertical bar corresponds to one individual and the proportion of each colour corresponds to the estimated ancestry proportion from a given cluster. Individuals are grouped from north to south. Sampling locations separated by black lines.

Figure 3 - Unrooted species tree graph obtained with TreeMix. Branch lengths are
represented in units of genetic drift, i.e., the longer a given branch the stronger the genetic drift
experienced in that lineage. Higher genetic drift can be due to older divergence times or smaller
effective population sizes of a given lineage. Arrows represent migration events.

Figure 4- D-statistics calculated for different topologies. For each topology, the results are shown for the different *S. pyrenaicus* North sampling locations (Ocreza, Lizandro, Canha) used. "S. carol" stands for *S. carolitertii* and "S.pyr Almargem" stands for *S. pyrenaicus* Almargem. Results obtained with each outgroup are represented by a different symbol (circles for *S. torgalensis* and triangles for *S. aradensis*). Full symbols represent significant D values (p<0.05).

Figure 5 – Schematic representation of the likelihood of the models tested with *fastsimcoal2*. The name given to each model is indicated above the schematic representation, as well the difference to maximum likelihood (Dif. To Max. Likelihood) which is the difference in log10 units between the estimated likelihood and the maximum likelihood if there was a perfect fit to the observed site frequency spectrum. The closer to zero (less negative values), the better the fit. The estimated admixture proportion is the parameter α. Models (A) to (C) have 8 parameters and therefore are directly comparable. Models (D) to (F) have 9 parameters and are also directly comparable. All inferred parameters are indicated in relation to the Ne of *S. carolitertii* (NeC)). T_A – divergence from the ancestral; T_P – divergence of the northern *S. pyrenaicus*; T_S – secondary contact. Population sizes are not to scale. See supplementary Figure S11 for relative likelihoods based on AIC, and Supplementary Tables S5-S8 for estimated likelihoods and parameter values.

Figure 6 – (A) Outlier loci detected in _S. pyrenaicus_ North. Red triangles correspond to
alleles from _S. carolitertii_ potentially selected for in _S. pyrenaicus_ North and blue triangles to
alleles from _S. pyrenaicus_ South potentially selected in _S. pyrenaicus_ North. **(B). Outlier loci**
from _S. carolitertii_ shared between the different _S. pyrenaicus_ North sampling locations
(C). Outlier loci from _S. pyrenaicus_ South shared between the different _S. pyrenaicus_
North sampling locations. Asterisk indicates the catalog loci identified as _vitellogenin_.

Table 1 – Pairwise F$_{ST}$ calculated between the different sampling locations. *S. pyrenaicus* Guadiana
was deliberately left out as there is only one individual from this sampling location.
958

	S. carolitertii	*S. pyrenaicus* Ocreza	*S. pyrenaicus* Lizandro	*S. pyrenaicus* Canha	*S. pyrenaicus* Almargem	*S. torgalensis*	*S. aradensis*
S. carolitertii	-	0.101	0.139	0.060	0.186	0.320	0.310
S. pyrenaicus Ocreza	-	-	0.138	0.054	0.204	0.344	0.331
S. pyrenaicus Lizandro	-	-	-	0.075	0.238	0.372	0.357
S. pyrenaicus Canha	-	-	-	-	0.171	0.309	0.294
S. pyrenaicus Almargem	-	-	-	-	-	0.351	0.338
S. torgalensis	-	-	-	-	-	-	0.184
S. aradensis	-	-	-	-	-	-	-