Error, noise and bias in de novo transcriptome assemblies

Adam H. Freedman*, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

Michele Clamp, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

Timothy B. Sackton, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

* Corresponding author: adamfreedman@fas.harvard.edu

ABSTRACT

De novo transcriptome assembly is a powerful tool, widely used over the last decade for making

evolutionary inferences. However, it relies on two implicit, untested assumptions: that the

assembled transcriptome represents an unbiased, if incomplete, representation of the underlying

expressed transcriptome, and that expression estimates from the assembly are good, if noisy

approximations of the relative abundance of expressed transcripts. Using publicly available data

for model organisms, we demonstrate that, across assembly algorithms, species, and data sets,

these assumptions are consistently violated. Using standard filtering approaches, coverage of

annotated genes by transcriptome assemblies falls far below complete coverage, even at the

highest expression levels, and hundreds of expressed genes are not assembled by any of the

algorithms we evaluated. Assemblies are compositionally biased towards non-coding elements

relative to their frequency in reads. Bias also exists at the nucleotide level, with genotyping error

rates ranging from 30-87%. As a result, diversity is underestimated in transcriptome assemblies,

with consistent under-estimation of heterozygosity in all but the most inbred sample we assayed.

Finally, even at the gene level and when one sums across contigs annotated to those genes,

expression estimates show wide deviations from map-to-reference estimates, and positive bias at

lower expression levels. We demonstrate a computational method to partly alleviate noise and

bias in expression estimates. Genome assembly, where feasible, may alleviate some of these

problems for non-model organisms, although genome assemblies are themselves imperfect in

many ways. Researchers should consider ways to minimize the impact of bias in transcriptome

assemblies.

1. INTRODUCTION

Although early advances fueled by RNA-seq were restricted to genome-enabled species, the advent of computational methods for de novo transcriptome assembly (Birol et al., 2009; Grabherr et al., 2011a) has empowered a wide array of evolutionary analyses in the absence of a reference genome. Their application has led to significant advances in fields as diverse as phylogenetics (Schwentner, Combosch, Pakes Nelson, & Giribet, 2017), evolutionary responses to climate change (Veilleux et al., 2015), the molecular basis of convergent evolution (Berens, Hunt, & Toth, 2015; Pankey, Minin, Imholte, Suchard, & Oakley, 2014; Velotta et al., 2017), speciation (K. Li et al., 2016), and local adaptation (De Wit & Palumbi, 2013).

All studies utilizing de novo transcriptome assemblies rely on two implicit assumptions: that the assembled transcriptome represents an unbiased, if incomplete, representation of the true underlying expressed transcriptome, and that the expression estimates from the assembly are good, if noisy approximations of the relative abundance of expressed transcripts. These assumptions are rarely tested. When tests are carried out, they typically employ reads simulated from curated transcript annotations that represent idealized data (e.g. Vijay, Poelstra, Künstner, & Wolf, 2013) . Similarly, empirical investigations typically rely on metrics of contiguity such as N_{50}, completeness with respect to the proportion of conserved orthologs recovered, i.e. BUSCOs (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015), and dimensionless quality metrics used to guide assembly parameter tuning using tools such as DETONATE (B. Li et al., 2014) or TRANSRATE (Smith-Unna, Boursnell, Patro, Hibberd, & Kelly, 2016). While these are standard metrics used to benchmark new assembly tools, they do not explicitly measure or test the robustness of expression estimates derived from the assembled contigs, or whether

genotypes, allele frequencies and population genetic parameters that are important for

evolutionary inference can be accurately estimated from a transcriptome assembly.

Indirect evidence suggests that violations of the assumptions noted above may be

frequent. Comparisons of different assembly algorithms indicate that there is a degree of

complementarity in missingness—one assembler will recover a gene that another cannot,

motivating methods to combine assemblies, e.g. the Oyster River Protocol (MacManes, 2018).

While gene absence is often attributed to the difficulties of assembling lowly expressed

transcripts, little is known about that kinds of genes that are missing, whether expression level is

their only important unifying feature, and whether they are biased towards functional categories

of interest to evolutionary ecologists. Second, highly fragmented assemblies comprised of

hundreds of thousands of relatively short contigs are pervasive in the literature, presumably

representing transcript fragments, haplotypes erroneously designated as separate transcripts, or

both. Intuitively, the challenge of mapping reads to fragmented gene models should diminish the

robustness of expression estimates. Unknown is whether the compositional aspects of

transcriptome assemblies negatively impact SNP calls and the allele frequency estimates and

other population genetic parameters derived from them.

Below, we evaluate the robustness of the assumptions inherent to de novo transcriptome

assembly, how their violation might distort evolutionary inferences, and identify particular

features of transcriptome assemblies that should be considered during the study design phase of a

project. Where possible, we demonstrate methods that ameliorate assumption violations and

augment the robustness of downstream results. While phylogenomic studies can discard the

majority of contigs using aggressive filtering schemes and still obtain recover a large number of

phylogenetically informative sites in conserved orthologs, these aggressive filtering options are

less appropriate for studies that seek to understand patterns of genetic variation or gene

expression across populations or closely related species. Therefore, we focus on methodological

considerations for this class of investigations.

2. MATERIALS AND METHODS

2.1 Reference genomes

For the purpose of benchmarking de novo transcriptome assemblies, and comparing assembly

and reference-based expression estimates, we downloaded from ENSEMBL genome and gtf

annotation files for the following organisms: house mouse, *Mus musculus* C57BL/6J

(GRCm38); clawed frog, *Xenopus tropicalis* (JGI_4.2); pufferfish, *Tetraodon nigroviridis*

(TETRAODON8); and fly, *Drosophila melanogaster* (BDGP6).

2.2 RNA-seq data

Brain tissue is both transcriptionally complex and expected to express a large proportion of an

organism's overall transcriptional profile. For this reason, the bulk of our analyses focus on de

novo assemblies for data generated for experiments involving mouse (*Mus* spp.) brain. These

data sets include the *Mus musculus* (C57BL/6J) dendritic cell data used in the original Trinity

paper (MDC), a pool of six whole brains from albino inbred *Mus* (BALB/c), and 8-sample pools

of whole brain samples from wild *M. musculus domesticus* from Massif Central, France (FRA),

Iran (IRN), Kazakhstan (KZK), and Germany (DEU). To assess the generality or particular

results with respect to assembly composition, we also generated assemblies for pufferfish

(*Tetraodon nigroviridis*) whole brain, clawed frog (*Xenopus tropicalis*) kidney, and fly

(*Drosophila melanogaster*) heads. Data set SRA accessions, sequencing strategy and depth are

summarized in Supplementary Table 1. All libraries except for MDC were sequenced on an

Illumina HiSeq 2000; MDC was sequenced on an Illumina GAII.

2.3 Short read processing

After an initial assessment of sequences reads with FASTQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), we performed kmer-based error

correction with RCORRECTOR (Song & Florea, 2015), followed by adapter and quality trimming

with TRIMGALORE! (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), setting

both the minimum phred quality score and adapter filtering stringency parameter to 5, and the

minimum post-trimming read length to 36. We based our quality trimming threshold upon

previous findings that performing stricter quality filtering can negatively impact assemblies

(MacManes, 2014). A second run of FASTQC was used to confirm that there was no remaining

adapter sequence in the filtered reads.

2.4 De novo transcriptome assembly

We built de novo transcriptome assemblies with TRINITY v. 2.1.1 (Grabherr et al., 2011b) using

default settings, and did not perform *in silico* normalization for any of the data sets. For the *Mus*

data sets, to evaluate the robustness of some key findings to the algorithm employed, we also

built assemblies using two promising, recently develop assemblers: SHANNON v. 0.0.2 (Kannan,

Hui, Mazooji, Pachter, & Tse, 2016) and BINPACKER v. 1.0 (Liu et al., 2016). Where strand-

specific reads were available, we ran TRINITY and BINPACKER with the appropriate strand-

specific options. Because the strand-specific mode of SHANNON is in experimental phase, on the

recommendation of its author (Kannan, personal communication), we treated reads as un-

stranded in all SHANNON assemblies.

To perform an initial assessment of assembly quality, we calculated standard metrics

typically reported in de novo transcriptome studies. First, we quantified the total number of

contigs ("transcripts" in TRINITY's nomenclature), the number of TRINITY-defined "genes"

comprising contigs deriving from the same graph, N_{50}, and the percent of contigs that were

successfully mapped to the genome. We then assessed read support for the assembly, by

mapping reads back to the assembly to assess the proportion that mapped concordantly, using

methods specified by the developers of TRINITY

(https://github.com/trinityrnaseq/trinityrnaseq/wiki/RNA-Seq-Read-Representation-by-Trinity-

Assembly). Finally, we quantified the number of completely covered, fragmented, and missing

single copy conserved orthologs using the program BUSCO v. 3.0.2 (Simão et al., 2015), using the

vertebrate ortholog model set for all assemblies but the fly, for which we used the arthropod set.

2.5 Assembly and read functional composition

For each contig within an assembly and for the assembly as a whole, we quantified the

proportion of unique assembly nucleotides that were CDS, UTR, intronic, start or stop codon,

noncoding elements (including pseudogenic transcripts of genes with

"polymorphic_pseudogene" gene biotype), introns of non-coding transcripts, and

intergenic/unnanotated sequence. We first mapped contigs to their respective reference genomes

with GMAP v. 2016-06-30 (Wu & Watanabe, 2005). We then filtered out unmapped contigs with

SAMTOOLS v. 1.5. (H. Li et al., 2009). For the analysis of whole assemblies, we then used

BEDTOOLS v. 2.25 (Quinlan & Hall, 2010) to sort, and merge intervals such that each reported

interval consistent of unique genomic nucleotides. Mapped intervals for GMAP alignments were

converted to bed format with the bamToBed function, with the '-splitD' argument such that each

mapped interval for a contig—excluding alignment gaps corresponding to putative splicing

events—was converted into a separate row.

To calculate the proportions of transcriptome assemblies corresponding to particular

functional elements, we employed a hierarchical classification of genomic nucleotides

(Supplementary Text). In order to determine if the functional composition of RNA-seq reads

differed from that of the transcriptome assemblies, we randomly sub-sampled one million paired-

end reads, and aligned them to their respective genomes using HISAT2 v. 2.0.4 (Kim, Langmead,

& Salzberg, 2015). We then performed the same hierarchical classification as was done for

assembled contigs, calculating frequencies for individual sequence fragments, i.e. across both

sequence reads when both were properly mapped. For multi-mapped reads, we calculated a

weighted average across all mappings. Visualization of functional composition was obtained by

randomly sub-sampling 100,000 reads that successfully aligned to the genome.

2.6 Coverage of reference transcripts and genes

After extracting reference transcripts from the genome annotation with RSEM v. 1.2.22 (B. Li &

Dewey, 2011a) , we mapped transcriptome assembly contigs to these transcripts using BLAT v.

3.1.6 (Kent, 2002). We required at least 95% identity in the alignment (but no minimum

alignment length in the query contig) to consider a hit valid. We quantified transcript coverage in

two ways. First, for each reference transcript, we identified the contig with the best alignment to

it, and calculated coverage as the sum of the BLAT alignment intervals. Given the minimal

evolutionary distance between query and target sequences, and because we required a minimum

of 95% identity for alignments, we did not penalize coverage for mismatches. Second, we

calculated the coverage of a reference transcript by summing coverage across all contigs with

alignments that passed our 95% percent identity criterion. Because a contig map produce valid

mappings to multiple transcripts of a gene, in calculating summed coverage we allowed a contig

to be used in coverage calculations for >1 reference transcript. Finally, we estimated gene-level

coverage as a TPM-weighted average of coverage for individual transcripts belonging to that

ENSEMBL gene id. We calculated these three coverage metrics for 1) an unfiltered transcriptome

assembly, and 2) for only protein coding transcripts with a minimum length of 200bp—the

default minimum contig length for TRINITY—using only contigs with transcripts per million

(TPM) $\geq$ 1, and with a predicted peptide sequence with a BLASTP hit against Uniref90

(downloaded 2017-04-05), We used TRANSDECODER v. 3.0.0

(https://github.com/TransDecoder/TransDecoder/wiki) to predict peptide sequences. For BLASTP

alignment, we set the maximum number of target sequences to one, and with a maximum

allowable e-value of 1×10^{-5}.

2.7 Expression estimation

For nine combinations of assembly method (TRINITY, SHANNNON, BINPACKER) and *Mus* sample

(MDC, BALB/c, Massif Central), we generated expression estimates with RSEM v. 1.2.22 (B. Li &

Dewey, 2011b) and BOWTIE2, v. 2.2.4 (Langmead & Salzberg, 2012). For TRINITY-based

analyses, isoform and gene level expression estimates were initially calculated using TRINITY-

specified gene and isoform identities for each contig. We also estimated "true" (reference-based)

expression by mapping reads directly onto reference transcripts, using the same RSEM/BOWTIE2

pipeline but with the ENSEMBL GTF annotation.

We assessed concordance between assembly and referenced-derived approaches in three

ways. First, we compared estimates for reference isoforms and their best hit BLAT contig in units

of counts per million (CPM), transcripts per million (TPM), and a corrected CPM that adjusts for

differences in effective length estimates between methods (see below, hereafter termed

weighted-adjusted CPM). We examine CPMs rather than raw counts because, despite using the

same RNA-seq data, differences in read mapping rates lead to differences in total mapped

counts, i.e. the estimates of library size by which differential expression algorithms normalize

individual samples. Second, we evaluated concordance between reference gene expression and

our best estimate of gene-level expression derived from the transcriptome assembly. Specifically,

we assigned assembly contigs to ENSEMBL genes based upon their BLAT alignments to ENSEMBL

isoforms, complied these assignments into a contig-to-gene map file, and provided this file to

RSEM when building the mapping index. RSEM *genes.results output files produced by this

protocol contain expression estimates summed across contigs aggregated by reference label; for

TRINITY, contigs without BLAT hits were aggregated by TRINITY components, i.e. "genes". For

SHANNON assemblies, the collection of contigs assembled from an individual graph do not

necessarily belong to the same gene (Kannan, personal communication). For all analyses

summing across contigs, Because preliminary analyses indicated that there were differences in

effective length estimates between the denovo assembly and map-to-reference approaches, and

because these lengths influence RNA abundance estimators, we assessed whether correcting for

effective length differences between reference transcripts and assembled contigs would improve

concordance between de novo assembly and map-to-reference based abundance estimates. We

did this by rescaling CPM as follows:

$CPM_{wt-adj} \dfrac{map-to-ref\ effective\ gene\ length}{denovo\ effective\ gene\ length} \times \dfrac{denovo\ gene\ length}{map-to-ref\ gene\ length} \times 10^6$

The first scalar adjusts for the difference in estimated effective length, while the second scalar

downweighs the correction as the denovo-based effective length approaches that obtained by

mapping to the reference annotation. In order to focus directly on structural aspects of

transcriptome assemblies (such as fragmentation and mis-assembly) that might negatively impact

expression estimate robustness, we only evaluated reference features to which at least one

transcriptome assembly contig could be aligned. In other words, we did not consider the trivial

case where expression is, practically speaking, zero when no contigs exist that can be used to

obtain counts.

Because it was suggested to us that RSEM might not perform as well when aligning to

transcriptome assemblies (N. Davidson, personal communication), we also compared gene-level

expression analyses between map-to-reference and map-to-denovo using expression values

estimated with KALLISTO v. 0.43.1 (Bray, Pimentel, Melsted, & Pachter, 2016). Because

KALLISTO operates on the transcript/contig level, we generated gene-level estimates by

aggregating to the gene level using the R package TXIMPORT (Soneson, Love, & Robinson,

2015).

If biases in expression estimates derived from transcriptome assemblies are correlated

across samples, this would mitigate downstream effects on differential expression testing. We

examined the distribution of differences in such biases among samples, measured as effect sizes.

Bias can be quantified as:

$$log_2\left(\frac{sample\ 1\ CPM_{denovo}}{sample\ 1\ CPM_{map-to-ref}}\right) - log_2\left(\frac{sample\ 2\ CPM_{denovo}}{sample\ 2\ CPM_{map-to-ref}}\right),$$

which can be re-arranged as:

$$log_2\left(\frac{sample\ 1\ CPM_{denovo}}{sample\ 2\ CPM_{denovo}}\right) - log_2\left(\frac{sample\ 1\ CPM_{ma-to-ref}}{sample\ 2\ CPM_{map-to-ref}}\right),$$

which is the bias observed when comparing log-fold change derived from different methods. We

can estimate an effect size as:

$$2^{EXP}\left[abs\left(log_2\left(\frac{sample\ 1\ CPM_{denovo}}{sample\ 2\ CPM_{denovo}}\right) - log_2\left(\frac{sample\ 1\ CPM_{ma-to-ref}}{sample\ 2\ CPM_{map-to-ref}}\right)\right)\right]$$

For example, given de novo CPM values for samples 1 and 2 of 15 and 20, and for map-to-ref 5

and 10, respectively, we would estimate that the magnitude of difference between samples would

be 1.5 times as great as if one had used the map-to-reference approach.

2.8 Analysis of missing genes

We investigated how often expressed genes are missing from transcriptome assemblies and

whether those genes are enriched for particular ontologies. For three of the *Mus* brain samples

(MDC, BALB/c, FRA), we first identified the set of expressed transcripts with TPM > 0 based

upon mapping to their respective reference genomes. We considered a gene present if any of its

transcripts were expressed. Conversely, we considered a gene absent from a de novo assembly if

no contig with TPM > 0 had a BLAT alignment to mapped to any of a gene's constituent

transcripts.

We performed two separate missingness analyses for each sample-assembler

combination. These involve differing levels of a) filtering on the assembly, and b) which

reference transcripts are considered as the "background" for enrichment analyses. For the first

analysis, we included all assembled contigs and their respective BLAT alignments, and used all

genes with at least one expressed transcript as background; this includes reference transcripts <

200 bp that Trinity, with default settings, will not assemble, as well as non-coding transcripts

(hereafter termed "background_unfiltered"). Second, we only retained contigs with TPM $\geq$1, and

restricted our background to protein-coding reference transcripts $\geq$ 200 bp ("background_filtered").

In both analyses, we only included in the background gene set those genes with a reasonable

probability of detection, which we defined as TPM ≥ 1 for at least one of a gene's constituent

isoforms.

To determine if particular functional categories were over-represented in genes that were

missing, we conducted a statistical overrepresentation test using the PANTHER classification

system (Mi, Muruganujan, Casagrande, & Thomas, 2013), via http://www.pantherdb.org/. We

used the ENSEMBL gene ids associated with the expressed reference transcripts as our "reference

list" and the ids of the missing genes as the "analyzed list", calculating statistical significance of

over/under-representation using a Fisher's Exact test.

To determine whether genes that are missing from assemblies are less conserved than

those that are present, we downloaded from the UCSC Genome Browser

(http://hgdownload.cse.ucsc.edu/goldenpath/mm10/phastCons60way/) the phastCons (Siepel,

2005) score bigWig track derived from multiple alignments of 59 vertebrate genomes to the

mouse genome (mm10). We then converted from bigWig format to bedGraph with the

KENTUTILS (https://genome-source.gi.ucsc.edu/gitlist/kent.git/blob/master/README)

bigWigToBedGraph tool. We calculated mean phastCons scores per ENSEMBL gene (defined as

the set of non-redundant CDS, start codon, 3` UTR, 5` UTR, stop codon, and exons associated

with an Ensembl gene ID). We compared phastCons scores of genes missing from the assembly

to actually expressed genes using a non-parametric Wilcoxon rank sum test.

2.9 Evaluating assembly redundancy

To evaluate contig redundancy, we first quantified the frequency and degree of overlaps between

contigs whose best BLAT hits are known, single-isoform genes in *Mus*. For four of these data

sets, there are independent estimates of genetic diversity obtained from genome-sequencing

(Harr et al., 2016), so we predicted that, if false assembly of haplotypes increased with genetic

variation, that the frequency of overlaps would correlate with diversity. We also calculated the

redundancy from BLAT all-by-all alignments of individual assemblies. If haplotypes are

assembled into different contigs, or contigs are assembled that only differ by a small number of

bases, e.g. by differences in assembled intronic sequence, this should manifest in an excess of

cases where alignments cover nearly all of the constituent contigs in those alignments, with few

or no mismatches.

2.10 Evaluation of variant calls from transcriptome assemblies

For these analyses, we focused on the *Mus* samples, and compared the map-to-reference

genotypes (hereafter MR) to those derived from TRINITY assemblies. We called variants off of

the reference genome using the GATK v. 4 (McKenna et al., 2010), and the Broad Institute's best

practices for RNA-seq data

(https://software.broadinstitute.org/gatk/documentation/article.php?id=3891; last updated 2017-

01-21). For the *Mus* samples representing pools, following the GATK developers'

recommendations, we also used the --sample_poidy argument, with a value set to two times the

number of individuals in the pool. For TRINITY assemblies, we first constructed

SuperTranscripts, hereafter (ST), (Davidson, Hawkins, & Oshlack, 2017), which attempt to

collapse overlapping contigs within Trinity "genes", and then called variants with the GATK

pipeline, based upon python scripts provided with TRINITY v. 2.5.1. Similar to our reference-based

approach to obtain our truth set, we edited the variant calling script to include the --

sample_ploidy argument. We did not use either base quality score or variant quality score

recalibration, to reflect the fact that the truth sets needed for these methods are rarely, if ever,

available for ecological genomics data. We applied the same strand bias, depth, and quality-by-

depth filters to raw variant calls from both methods to produce a final filtered set. To define

callable sites for the MR approach, we extracted read alignments to annotated exonic regions,

and then calculated the number of exonic sites with a coverage depth ≥ 5. For ST, we performed

similar operations, defining exonic coordinates from mappings of ST to the *Mus* genome using

GMAP, and callable sites within exons in the same way as with MR, i.e. coverage depth ≥ 5. We

defined heterozygous calls in our filtered vcf files as all bi-allelic SNP sites that passed filtering

criteria. In cases where sample ploidy was greater than diploid, we collapsed genotypes into the

union of all unique recorded alleles.

We obtained exonic regions for ST by mapping them to the *Mus* genome using GMAP, then

intersected these mappings with a bed file of exonic intervals with BEDTOOLS. We used a

slightly newer version of GMAP (v. 2018-07-04) than we did for the contig composition, which

fixed minor issues with CIGAR string handling that had the potential to generate spurious

genotype errors. Python scripts

(https://github.com/harvardinformatics/TranscriptomeAssemblyEvaluation) were then used make

transformations between genomic and ST coordinates, for the purpose of extracting ST

genotypes and writing genotype bed files back to genomic coordinates. ST alignments to the

genome were used to define unique polymorphic site ids that included cases where (a) both

methods called genotypes, (b) only MR called a genotype (including sites with and without valid

ST alignments), or (c) genotypes were only called from ST. We recorded cases where multiple

ST were aligned to the same genomic position, taking the union of all reported alleles as the ST

genotype for that position. We restricted our comparison of the two methods by filtering out MR

sites that were indels or contained > 2 alleles. Using genotype calls based upon the reference

annotation as the gold standard, we calculated 15 different metrics, defined as follows:

• **Error rate**: the proportion of ids where the set of alleles recorded by the two methods

were not the same, including cases where one method did not call a genotype; *total # of*

incorrect ST calls aligned to genomic positions÷ total # of genomic positions to which

STs were aligned

• **Error-MrefAlleleIncluded**: the proportion of genotyping errors (in ST) where the

reported alleles included at least one of those reported using MR; *total # of incorrect ST*

calls relative to MR, but that include on MR allele ÷ total # MR calls to which STs were

aligned

• **Error-Snv2Indel**: the proportion of errors, where a biallelic SNV in MR was called as an

indel in ST; *# of ST indel calls at biallelic MR SNV sites ÷ total # bi-allelic MR SNV*

calls

• **Error-MultipleSuperTranscripts**: the proportion of errors where multiple STs were

aligned to the genomic position; *total # of ST genotyping errors at MR sites with multiple*

ST alignments ÷ total # MR genotype calls with ST alignments

• **FN**: false negative rate, the proportion of MR variable sites with no ST genotype; *# MR*

variable sites with either no ST alignment or with an alignment but no genotype call ÷ #

MR variable sites

• **False negatives-noalign**: the proportion of false negatives that are due to the absence of

ST alignments to the genome at the variable position; *# MR variable sites with no ST*

alignment ÷ total # MR bi-allelic SNV sites

•   **FN-het**: the proportion of heterozygous MR genotype calls with either no ST genotype, a

fixed alternative (relative to ST) genotype, an indel heterozygote call, or a multi-allelic

ST genotype. ST SNV heterozygous genotypes that specify the wrong alleles are not

considered FN, which is relevant for comparisons of transcriptome-level heterozygosity

between methods*; # of bi-allelic SNV MR genotype calls with no corresponding ST SNV*

biallelic genotype ÷ total # bi-allelic SNV MR genotype calls

•   **FP**: false positive rate, the proportion of ST genotypes with either no associated MR

genotype, or where ST is polymorphic and MR is fixed for an alternative allele; *# MR*

invariant sites called as polymorphic with ST ÷ total # invariable MR sites

•   **FP-het**: the proportion of invariant MR sites that are represented by heterozygous bi-

allelic ST genotypes, i.e. with either no associated MR genotype or a fixed alternative

MR genotype; *# invariant MR sites with ST heterozygote genotypes ÷ total # invariant*

MR sites

•   **FP-NAtoFixed**: the proportion of MR sites with no genotype call that are called as a

fixed variant in ST (relative to the ST reference); *# fixed-alternative ST genotypes at MR*

sites with no genotype call ÷ total # sites with no MR genotype call

•   **FP-NAtoHet**: the proportion of MR sites with no genotype call that are called as

heterozygotes in ST ; *# heterozygote ST genotypes at MR sites with no genotype call ÷*

total # of sites with no MR genotype call

•   **FP-indel**: the proportion of ST genotype calls at MR invariant sites (both missing and

MR-fixed) that are indel; *# invariant MR sites called as indels with ST ÷ # MR invariant*

sites with ST genotype calls

• **Recall:** the proportion of MR genotypes recovered by ST, including matching invariant

sites; (# *genomic positions with MR genotype calls with matching ST calls + matching*

invariant MR and ST calls) ÷ (total # MR called genotypes +sites with matching invariant

calls), i.e. *#TP/(#TP+#FN), where TP = true positives.*

• Recall-het: the proportion of MR heterozygous genotypes that are correctly recovered in

ST; a genotype is not considered recovered if ST genotype is a heterozygous call

comprised of the wrong nucleotides; *# MR bi-allelic heterozygote calls with matching ST*

heterozygote calls ÷ total # MR bi-allelic heterozygous calls

• Precision: the proportion of ST genotypes that match the MR genotype; *# ST genotype*

calls with a matching MR genotype call ÷ total # ST genotype calls, i.e. TP/TP+FP

2.11 Analysis reproducibility

Software command lines, scripts and accessory files used to implement analyses described above

are available at https://github.com/harvardinformatics/TranscriptomeAssemblyEvaluation.

3. RESULTS

3.1 Assemblies

We assembled de novo a total of 21 transcriptomes—six mouse brain data sets using TRINITY,

SHANNON, and BINPACKER, as well as TRINITY assemblies for frog kidney, pufferfish brain and

fly head. Consistent with previously published reports, assemblies were highly fragmented, with

median contig sizes ranging from just a few hundred bases to slightly over 1kb (Supplementary

Table 2). Yet, typically employed measures of accuracy and completeness indicate that all

assemblies were of reasonably high quality (Supplementary Tables 2-4), particularly those

assembled from the *Mus* data sets comprising the majority of our analyses. Important in the

context of gene expression studies, large fractions of reads used to build assemblies mapped back

to them in a concordant fashion, usually exceeding 90% of the input library (Supplementary

Table 3)

3.2 Transcript and gene coverage

Without filtering on the reference transcriptome or the de novo assembly, transcript coverage is

increased when one sums coverage across orthologous assembly contigs (Figure 1a-b,

Supplementary Figure 1-3). In both methods, and at the level of genes (Figure 1c, Supplementary

Figures 1-3), coverage generally tracks reference expression level, although these correlations

are weaker at the transcript level, and weakest for coverage of a single best-hit contig. For

summed coverage of transcripts and gene coverage, while there is an expectation of complete

coverage above TPM=10, there are still many transcripts and, to a lesser extent, genes that fall

far below complete coverage. This "unfiltered" approach represents an idealized scenario in

which contigs are mapped directly to a reference annotation originating from the same species. If

one considers only protein-coding genes and transcripts ≥ 200bp—which should favor assembly

coverage—but takes a standard approach of filtering out lowly expressed contigs, and only

considering contigs with hits to UNIREF90 (a standard approach for assigning orthology without

an annotated genome)—coverage decreases substantially at both the transcript and gene level

(Figure 1d-f, Supplementary Figures 1-3). The expectation of coverage never reaches 100%,

regardless of expression level of the underlying reference structures, and there is far more

variance around the coverage-TPM trend. The decreases observed in these trends at the highest

expression levels result from poorly covered annotations making up an increasing fraction of the

most highly expressed transcripts and genes.

3.3 Missing genes

Only considering genes that are expressed at a level to permit a reasonable chance of being

detected (TPM ≥ 1), depending upon the sample, 17-78 (0.015-0.052%) of genes were missing

from all methods' unfiltered assemblies, and 25-115 (0.23-0.77%) genes were unrecovered by 1-

2 assembly methods (Table 1, Supplementary Figure 4a). Practically, unfiltered assemblies are

rarely, if ever, useable, so to test a more realistic scenario we also estimated the number of genes

missing from filtered assemblies. Missing from TRINITY assemblies were 3.5% (293), 2.6%

(277), and 4.7% (529) protein coding genes for MDC, BALB/c and FRA, respectively. Numbers

were substantially worse for SHANNON (MDC, 19.1% (1585); BALB/c, 9.9% (1039); FRA, 7.5%

(836)), and better in two of three samples for BINPACKER (MDC, 3.0% (251); BALB/c 2.9%

(306); FRA 3.0% (340)). Overall, any one assembler-sample combination led to hundreds to over

1,000 expressed genes being missed, at least after standard assembly filtering, which

unavoidably removes both spurious contigs and contigs representing real transcripts. In general,

genes not recovered by one method tended to be missing from all assemblers (Table 1,

Supplementary Figure 4b); SHANNON tended to show greater missingness for protein coding

genes than TRINITY and BINPACKER. With the exception of MDC, genes missed by all or some

assemblers had significantly lower reference gene expression and conservation (Figure 2,

Supplementary Figure 5, Table 2). Even so, there were plenty of cases where conserved genes

expressed at a relatively high level (TPM ≥ 100) were not recovered by any assembler (Figure

2). Correlations between TPM and phastCons scores were, while statistically significant,

relatively weak (MDC, Spearman's ρ = 0.082, p = 2.2 × 10^{-16}; BALB/c, Spearman's ρ = 0.18, p

= 2.2 × 10^{-16}; FRA, Spearman's ρ = 0.23, p = 2.2 × 10^{-16}). Missing genes were not enriched for

any GO-Slim Biological Process terms.

448
3.4 Assembly vs. read functional composition

For a gene to be recovered in an assembly, it has to be annotated. And, because transcriptome

assembly is motivated by the absence of a closely related reference genome, in nearly all cases

transcriptome annotation involves predicting open reading frames and searching protein

databases such as UNIREF90 and SWISSPROT. The greater the fraction of non-coding bases in a

contig, the harder it will be to align it to an annotated peptide. Thus, the functional composition

of contigs sheds light on patterns of missingness at both the transcript and gene levels.

Across all sample-assembler combinations, large fractions of contigs are dominated by intronic,

UTR, and intergenic sequence (Figure 3, Supplementary Figure 9). The extent of intronic

sequence is particularly surprising for *Mus* and *Drosophila* assemblies, as current annotations

should be inclusive of the majority of alternative splice variants. That the excess of intronic

sequence in contigs is not reflected in the composition of reads (Figure 3, Supplementary Figure

6) can be attributed to the sensitivity of the assembly algorithms, which require minimal kmer

coverage from just a few intron-containing reads to extend contigs across exon boundaries. The

prevalence of intronic and intergenic sequence in assembly contigs indicate that a substantial

proportion of assembled bases are derived from some combination of transcriptional read-

through across exon junctions, assembly of low-abundance pre-mRNA, and low levels of

spurious genome-wide transcription (Supplementary figure 7). The mean contig fraction

comprised of intron sequence for TRINITY assemblies ranged from 7.5% (*Xenopus*) to 56.5%

(FRA) (Supplementary table 5). Mean fractions for intergenic sequence ranged from 8.1%

(MDC) to 39.0% (*Xenopus*). Results for SHANNON and BINPACKER assemblies of MDC,

BALB/c and FRA were similar to those observed in TRINITY.

471   Redundant assembly of identical contigs from and assembly of different haplotypes as

unique contigs can create complications for downstream analyses. Contig multiplicity can

influence downward expression estimates of contig copies—particularly if redundant contigs are

not properly assigned to the same gene-level annotation—and decrease the signal-to-noise ratio

for differential expression analyses. Assembling haplotypes as contigs can lead to false positive

differential expression results, e.g. if frequencies of haplotypes differ between populations. It can

also distort allele frequency spectra that are critical for analyses of allele-specific expression.

Across all three assemblers, the frequency of contig overlaps within known single-isoform genes

was bimodal. Most genes were overlapped at few or no positions by ≥ 1 contig, but a large

number of genes were also covered by multiple contigs at most or all of their positions (Figure

4). SHANNON, in particular, showed a troubling increasing trend in frequency with proportional

reference gene coverage (Figure 4b), indicating it may be more prone to redundancy than either

TRINITY or BINPACKER. Overall, 11.9%-12.2%, 33.2%-35.3%, and 16.9%-18.7% of single

isoform genes were covered by more than one contig at all transcript positions for TRINITY,

SHANNON, and BINPACKER, respectively. The frequency of contig overlaps has no relationship

with estimates of genetic diversity obtained from whole genome sequencing (Figure 4), a pattern

not to be expected to occur if redundancy is driven solely by the assembly of polymorphic

haplotypes into separate contigs. Considering the variation in CDS and intron content among

contigs that cover a single-isoform gene, while there are plenty of cases with variable assembly

of CDS among contigs, the event of highest frequency across all assemblers appears to be one

where there is no variation in CDS among contigs but variable intron retention, due to either

transcription of intron sequence or the inclusion of un-spliced pre-mRNA in sequencing libraries

(Supplemental Figure 8). Looking across all contigs, regardless of their genomic position of

origin, all assemblies contain large numbers of contigs that are nearly identical (Figure 5a,

Supplemental Figure 9). For TRINITY assemblies that putatively report "genes" as components,

while the redundancy is greater for contigs assembled from the same graph, there are a

substantial number of contigs with high similarity that originate from different TRINITY "genes"

(Supplementary Figure 10). From different component graphs, Trinity will erroneously assemble

haplotypes that differ only by a few SNVs (Figure 5b).

3.5 SNP calling and heterozygosity

SNP calls and allele frequencies inferred from them and that are derived from transcriptome

assemblies have been used in a variety of contexts, including estimation genetic diversity in

evolutionary analyses (Romiguier et al., 2014), population structure (Tepolt & Palumbi, 2015),

reconstruction of demographic history and gene flow (De Wit & Palumbi, 2013; Le Gac et al.,

2016; Roda, Mendes, Hahn, & Hopkins, 2017), estimation of dN/dS (Cokus, Gugger, & Sork,

2015), F_{ST} outlier analysis (Bankers et al., 2017) , and tests for allele-specific expression (Singh

et al., 2018). Our ST mapped at extremely high rates to the *Mus* genome (MDC, 99.9%;

BALB/c, 99.8%; FRA, 99.8%; DEU, 99.8%; IRN, 99.8%; KZK, 99.7%), such that any

discrepancies between genotyping methods cannot be attributed to low alignment efficiency. In

principle, redundancy through assembly of haplotypes as distinct contigs should lead to

erroneous genotype calls and biased estimates of allele frequency spectra. For example, when

two contigs represent haplotypes: at a variant position, the true heterozygous position is instead

represented as two separate genotypes, one homozygous for the reference allele and one

homozygous for the alternative allele. The opposite problem can happen when paralogous loci

are collapsed, leading to a heterozygous genotype call where there would otherwise be two

invariant sites, or perhaps a mixture of alleles due to, for example, collapsing of an invariant and

a heterozygous position. We compared genotype calls and derived diversity metrics between the

map-to-reference and map-to-transcriptome approaches using 15 different metrics, in order to

determine whether such genotypic distortions occur in de novo transcriptome assemblies. The

map-to-reference (MR) approach involves mapping sequence reads to the genome, then calling

genotypes with GATK based upon those genomic alignments. In contrast, the transcriptome

assembly-based approach involves aligning reads to the ST collapse of a TRINITY assembly, then

calling variants with GATK relative to the ST. Comparisons between MR and ST genotypes are

based upon orthology determined by aligning the ST to the reference genome. To see example

cases of how particular errors are flagged, see Figure 6.

The frequency of errors, in which the allele set recovered by the two methods differed or

in which only one of the two methods called a genotype at all, was high, ranging from 30-38%

for the wild samples, and surprisingly, $\geq 50\%$ for the inbred samples (Table 3). While the median

per-base error rate per ST was 0.2% or less, maximum per ST rates typically exceeded 10%

(Supplementary Table 6). For five of six samples, the median number of errors per ST was 2,

but, depending upon the sample, as many as 56-187 errors could occur within a single ST

(Supplementary Table 6). On a per-sample basis, errors show a modest yet statistically

significant bias towards the distal end of a ST (sample-level Wilcoxon 1-sided tests, $H_0 = 0.5$, all

tests, $p \leq 0.002$; Supplementary Figure 11). Considering only contigs with an ORF hit to

UNIREF90 changed error rates in a predictable direction, e.g. reduced false positives and recall,

and increased false negatives (Table 3).

In 90-100% of all erroneous ST genotype calls, at least one MR allele was included in the

ST allele set (Table 3, Error-MrefAlleleIncluded). False negatives (FN; 24-66%) occurred far

more frequently than false positives (FP; 8-61%). Anywhere from 57-75% of FN were due to

there being no ST aligned at a genomic position where a MR genotype call was made (Table 3,

FN-noalign). Given that the ST mapping rates to the genome were extremely high, these FP

instances are primarily due to a failure to assembly transcripts overlapping polymorphic genomic

positions. The FP rate includes both sites where there is no genotype call in MR, or where MR is

fixed for an alternative allele and ST is polymorphic. Looking more closely, paralog collapsing

should lead to FP heterozygous ST calls, and FP ST fixed-alternative genotypes will be due local

assembly errors; in the latter case, mis-assembly is the most parsimonious explanation, given that

the ST reference sequences were derived from the data used to call genotypes. FP-NAtoHet was

5-38 times higher than FP-NAtoFixed, indicating that paralog collapsing plays a larger role in FP

than local assembly errors within ST. FP-NAtoHet includes ST indel heterozygotes, which are

not unexpected in cases of paralog collapse; The frequencies of cases where MR SNVs are called

as indels in ST (Error-Snv2Indel) approach 50% in the wild samples. This class of error as well

as the somewhat less frequent occurrence of false positive indel calls (FP-indel) collectively

suggest that paralog collapses are a general problem, if not as large a contributor to overall error

rates as FN. When genotypes were called in STs, the frequency of calls that were correct (i.e.

precision) ranged from 81-90% for the wild samples, but precision for the inbred samples,

particularly MDC, was substantially lower. Recall ranged from 73-76% in the wild samples; as

with precision, recall for the inbred MDC sample was extremely low.

Because read depth influences the ability of tools such as GATK to call genotypes, we

compared read depth at homologous positions, comparing MR depth and median ST depth (to

account for the possibility of > 1 ST mapping to a genomic position). Concordant genotypes

displayed a high degree of concordance in depth-of-coverage (Supplementary Figure 12). In

contrast, on average FN sites had very small reductions in ST read depth relative to MR at sites

where STs mapped to the genome (Supplementary Figure 13, blue lines). FN at sites with modest

loss of read depth is likely caused by multiple interacting factors that are beyond the scope of

this study to investigate in detail. In contrast, FN due to failure to assemble transcripts leads to an

effective ST read depth of zero, and including these sites leads to a much more dramatic

reduction in ST depth relative to MR (Supplementary Figure 13, yellow lines). With the

exception of a relatively small number of cases of zero MR coverage, MR and ST read depth

were largely co-linear (Supplementary Figure 14); the causes of noise about those correlations,

as with FNs, remain unclear.

One prediction resulting from such genotyping bias is that heterozygosity rates in ST will be

negatively biased relative to MR, because alterative alleles will be partitioned amongst different

assembly components (i.e. "genes") and reads overlapping these alleles will preferentially map to

their corresponding contigs, leading to separate homozygous reference genotype calls rather than

one heterozygous call. Consistent with this prediction, in all but the inbred (MDC, BALB/c)

strain samples, we observe reduced heterozygosity in the exonic regions of ST relative to MR

(Figure 7). This bias does not result from ST being preferentially assembled in regions of lower

heterozygosity, given that, for the wild samples, MR heterozygosity is actually higher in ST-

overlapping regions (Supplementary Figure 15). Furthermore, the bias cannot be due to some

unique feature of genotype calls in exons, as the same pattern holds when considering all

genotype calls (Supplementary Figure 16). Another possible explanation may be lower power to

detect low frequency alternative alleles with ST. However, in our five samples consisting of

pooled individuals, minor allele frequency at MR heterozygous sites did not vary between

correct ST heterozygous calls, FN, and FP (Supplementary Figure 17). The driving factor

appears to be FN at heterozygous sites; as already noted, FN are dominated by cases where ST

are not assembled (Table 3). An additional contributing factor may be erroneous indel

polymorphism genotypes in ST. Given that indel genotypes are harder to call correctly than

SNVs, a standard approach is to filter out indel sites. Because we perform the same filtering

when estimating heterozygosity, we effectively filter out many true heterozygous SNV positions,

further reducing heterozygosity estimates in ST. Paradoxically, FP genotype calls may actually

mask the magnitude of this bias.

Gene and isoform expression

The compositional deviations of de novo transcriptome assemblies relative to the true set of

expressed transcripts identified by mapping reads to a reference annotation results have a net

effect on expression estimates. Whether in terms of CPM or TPM, expression estimates derived

from the contigs that are the best BLAT hit relative to the reference transcripts are only weakly

correlated with map-to-reference estimates for those transcripts, and show positive bias for lowly

expressed reference transcripts (Supplementary Figure 18ab, Supplementary table 8). Weighted-

adjusted CPM correlations with reference isoform CPM led to virtually no change in

performance relative to other metrics (Supplementary tabler 8). Surprisingly, for TRINITY and

BINPACKER, contig weighted-adjusted CPM led to a substantially stronger correlation with

reference gene level expression (Supplementary Figure 18c, Supplementary Table 8), but fell

way below gene level correlations (see below and Table 4). Performance of SHANNON was

generally poor, and weighted-adjusted CPM correlation with reference gene expression was

similar to or even worse than for other abundance estimators.The practice of using single contigs

as proxies for gene-level expression is problematic, as contig-based estimates are also poor

estimators of reference gene expression. Since individual contigs typically represent a fraction of

the mappable sequence of a set of transcripts belonging to a gene, one expects smaller counts on

contigs relative to the full set of mappable positions encompassed by transcripts produced by a

gene locus. Given this fact, it is surprising that, for lowly expressed genes, their associated best-

hit contigs on average have larger counts (Supplementary Figure 18c). It is also surprising that

TPM, a metric that should account for effective length differences, shows a similar pattern of

bias (Supplementary Figure 18d).

Expression estimates derived from de novo transcriptome assemblies are far more robust

at the gene level, when one is able to group contigs into genes via annotation (Figure 8, Table 4).

It is reassuring that the correlation between the methods is quite strong, although there is sizeable

positive bias at lower expression levels for TPM, which is substantially reduced for CPM.

Furthermore, for many genes there are substantial deviations from estimates obtained with

mapping to the reference. Bias and deviations from the map-to-reference estimates are reduced

when one rescales CPMs to account for differences in effective length estimates (Figure 8c).

This indicates that effective length (EL) differences—that is, the difference in the number of start

sites for read alignments, given a particular sequenced fragment size, between a full-length

transcript, and a set of transcript sub-sequences represented by contigs—are part of the problem,

and in particular, that effective length is not properly estimated on the fragmented gene models

that are typical in denovo transcriptome assemblies. Correlations between reference gene

expression and gene-level summing across assembly contigs obtained with KALLISTO were very

similar to those obtained with RSEM (Table 5). Interestingly, adjusting for effective length had

little effect on KALLISTO CPM, and in some cases reduced the correlation between methods. This

suggests that KALLISTO may be doing a better job at estimating effective length on transcriptome

assemblies, such that it's inferior performance relative to RSEM must be due to other reasons.

As demonstrated earlier, high gene coverage is most frequently obtained for highly

expressed genes, but transcriptome assembly-derived expression estimates can still deviate

substantially from the map-to-reference gold standard at high coverage (Supplementary figure

21). Such deviation is well explained by the disparity in EL, with estimates showing the least

deviation where EL values are identical or nearly so (Figure 9), EL disparities persist at high

expression levels, even if those disparities are not as variable as at lower expression levels

(Figure 9). There are positive and negative trends in bias for CPM (Figure 9a) and TPM (Figure

9b), respectively, with the clearest bias pattern observed for TPM. Adjusting CPM for effective

length differences effectively eliminates biases that would otherwise be produced by differences

in EL (Figure 9c). Nevertheless, negative bias at lower expression levels and large deviations

from map-to-reference expression estimates are ubiquitous.

Biases in expression estimates derived from transcriptome assemblies might not have

large effects on downstream differential expression analyses if biases are consistent across

samples. While the pooled *Mus* data sets we used did not permit us to look at inter-individual

variation amongst individual libraries used to build a single assembly, we looked across the wild

Mus datasets to see if biases in gene-level expression were concordant, i.e. were biases for a

particular gene always of similar magnitude and in the same direction. We focus on results for

TRINITY assemblies of wild *Mus* samples, as TRINITY consistently showed stronger correlations

with map-to-reference based estimates, such that our findings are conservative with respect to

assemblers showing weaker correlations. Reassuringly, log-fold change of transcriptome

assembly vs. map-to-reference expression estimators (LFC$_{METHOD}$) showed broad patterns of

moderate concordance at the gene level indicating that gene-level features, influence the

magnitude of bias (Supplementary Figure 22). Inter-sample correlations were, for CPM, 0.625-

0.727; TPM, 0.790-0.830; and wt.-adj. CPM, 0.605-0.671. These correlations are weaker for

CPM and wt.-adj. CPM, in part, because biases are reduced overall, compressing LFC$_{METHOD}$

toward zero. Despite these broad correlations, a number of genes may show excessive bias. We

measured this as a bias effect size that indicates how much greater a difference in expression

would be detected using a denovo transcriptome approach, relative to mapping to reference

annotation. Comparing the FRA and DEU samples, for CPM, TPM, and wt-adj-CPM, 5.24%,

5.16% and 3.35% of genes showed bias effect sizes $\geq$ 5, with more than 2% of genes showing an

order of magnitude of difference between methods of $\geq$ 10 for CPM and TPM (Figure 10).

Correcting for effective length differences substantially reduced the frequency of such bias

outliers, but did not eliminate them entirely.

DISCUSSION

De novo transcriptome assemblies have been an important entry point for functional and

population genomics in non-model organisms, but the two key assumptions underlying their

application are consistently violated, albeit in unexpected ways. Below, we explain how

particular compositional aspects of de novo transcriptome assemblies may impact the power of

evolutionary inference.

While some genes not assembled by one algorithm are by another, dozens of genes will

consistently fail to get assembled. While this seems like a modest fraction of genes, this occurs

only in the idealized case where one can perform nucleotide-level searches against a transcript

database from a closely related organism, or in our case, the same organism. Using a common

approach that filters out lowly expressed contigs, and only retains those with a hit to a protein

database, we found that hundreds of protein coding genes (from 1-2%) would not be recovered

by any assembly method, and some assemblers missed several hundred to over 1,000 genes.

Thus, while almost all expressed genes may be in the assembly somewhere, annotation

challenges amplify missingness. Clearly, combining outputs from multiple assembly

algorithms—a non-trivial challenge in its own right—will not prevent missing a sizeable fraction

of genes. The efficiency of annotation will be further exacerbated by mis-assembly errors, such

as those revealed by indel errors within SuperTranscripts. As *Mus* proteins are well represented

in UNIREF90, our finding is overly optimistic relative to the typical predicament molecular

ecologists find themselves in. Annotation problems should be greatest for more rapidly evolving

genes as they will be more divergent from currently archived protein and nucleotide sequences.

Previous studies have confirmed that increasing divergence to annotation databases increases

annotation errors (Hornett & Wheat, 2012; Ockendon et al., 2016; Vijay et al., 2013), and these

issues should be more precisely considered with respect to both taxon divergence and the

evolutionary rate of genes within pathways of interest.

The difficulty in assembling genes that show less conservation is due, in part, to their

being lowly expressed. Whether due to the difficulty assembling contigs from few reads, or

polymorphism in more rapidly evolving genes, the fact remains that genes tied to phenotypes of

interest may be missing from transcriptome assemblies. To the extent that expression level drives

missingness, conserved, lowly expressed genes such as transcription factors that produce large

phenotypic effects may also be missing from transcriptome assemblies more often than expected

by chance. Molecular ecologists must not confuse expression level with the capacity to drive

phenotypic evolution and should find means to estimate the probability that genes of interest are

likely to be recovered in a de novo transcriptome assembly.

While it is unsurprising that lowly expressed genes might be harder to assemble, the

correlation between expression level and conservation is relatively weak, and in many cases

highly expressed genes were missing from annotation and expression-filtered assemblies. Thus,

factors correlated with conservation, such as polymorphism, may influence the probability an

expressed gene is recovered. Future investigation aimed at quantifying their relative effects could

shed important insights relevant to researchers interested in particular genes. Despite our

incomplete understanding of all the factors that may lead to genes being missing from an

assembly, the superficially good news is that missing genes do not represent a biased subset

enriched for particular GO categories. The bad news is that, beyond the reduced probability of

assembly for lowly expressed genes, it is not possible to predict in advance and with reasonable

certainty that all genes relevant to a particular study being proposed will be recovered.

The nucleotide composition of contigs sheds light on how genes may wind up missing

from a de novo transcriptome assembly. We found a relatively high frequency of non-coding

bases in assembly contigs, which by definition, places constraints on the utility of standard

protein database searches. To the extent that an underlying CDS is divided into a series of

contigs, power to discover a protein homolog will be reduced; for contigs dominated by non-

coding bases containing only short strings of CDS, successful ORF prediction and annotation is

unlikely.

We expect that missingness both actual and annotation-based, combined with the high

frequency of sequence reads dominated by UTR bases, will also reduce power for differential

expression studies in a previously unappreciated way. Practically speaking, one throws away all

reads that map to unannotated contigs in addition to unmapped reads. Rules of thumb for

sequencing depth sufficient for differential expression analyses are largely derived from map-to-

reference studies, and do not consider this additional source of annotation-conditional filtering.

Loss of coverage depth is most problematic for evolutionary studies of non-model organisms,

given that they frequently employ small numbers of biological replicates (Todd, Black, &

Gemmell, 2016) which places an additional constraint upon statistical power. We suggest that

researchers employing transcriptome assemblies determine the number of reads per sample that

remain after those mapping to unannotated contigs have been discarded and consider additional

sequencing if read counts far well below desired targets. A second possibility is to "rescue"

unannotated contigs, by finding ways to cluster them with annotated contigs. Standard clustering

approaches such as CD-HIT will not perform well in this context, as they operate on sequence

similarity. By definition, a contig derived mostly from UTR will share little sequence with an

annotated contig comprised mostly of coding sequence. Underlying this type of phenomenon is

the fact that assembly graphs will often get broken into pieces deriving from different functional

compartments. For example, a short repeat in a UTR can split a transcript into an exclusively

UTR contig, and a contig comprised of a few UTR bases plus the down or upstream CDS.

Recently developed methods that leverage information from equivalence classes obtained from

pseudo-alignments of paired-end reads, such as CORSET (Davidson & Oshlack, 2014) and

GROUPER (Davidson & Oshlack, 2014), may help mitigate the loss of unannotated contigs. In

particular, GROUPER utilizes "orphan" read pairs that map to different contigs, which should

enable clustering of contigs representing transcript fragments that share little or no sequence.

Integration of this type of approach with annotation-based clustering is a promising avenue that

requires further investigation.

In contrast to missingness, two kinds of artefactual redundancy create complications for

downstream analyses. First, for any given genomic region from which an actual transcript is

generated, transcriptome assembly algorithms will call multiple isoforms that differ only by the

proportion of retained intron sequence. This adds an additional obstacle to performing isoform-

level expression analyses beyond those imposed by the incomplete and fragmented recovery of

transcripts, as counts will be allocated incorrectly across multiple spurious splice variants when

they should be aggregated to one transcript, potentially obscuring true underlying inter-isoform

variations in expression level. A second form of redundancy occurs in that haplotypes regularly

get called as isoforms, and, at least for the commonly used Trinity SuperTranscript-based

pipeline, with those haplotypes often being produced by different DeBruijn graphs that are

supposed to represent different genes. This added source of false multiplicity makes robust

inference of isoform-level expression even more challenging, although gene-level analyses might

not be as seriously impacted, assuming that clustering approaches such as GROUPER yielded

robust groupings of contigs.

Even when annotation-based groupings of contigs are performed in an optimal setting,

such as our analyses of *Mus* transcriptomes, there is substantial noise in gene-based expression

estimates derived from transcriptome assemblies relative to a map-to-reference based approach,

and there is positive bias at lower expression levels. Furthermore, the practice of selecting a

single best hit contig to represent a gene seems ill-advised given that correlations between contig

and map-to-reference gene-level expression are poor, and subject not only to positive bias at

lower expression levels, but negative bias at higher levels. Incorrect estimation of effective

length metrics used in RNA abundance calculations play a substantial role in the observed

patterns of bias. Nevertheless, it may be surprising to researchers that biases persist for standard

length-adjusted metrics such as TPM. CPM has less bias compared to TPM. Bias and noise can

be reduced by rescaling CPM to account for differences in effective length, such that the rescaled

CPM more faithfully recapitulates map-to-reference based expression estimates, even if

substantial noise persists. Adjusting expression estimates to account for effective length

differences can be beneficially used in two different applications. First, one could conduct

differential expression estimates across different transcriptome assemblies when it is necessary

to generate different assemblies for different conditions, such as divergent species or sub-species.

Because one of the first steps differential expression tools execute is to normalize by library size,

one can simply apply the correction directly on the counts themselves. CPM and count scale are

directly proportional, with the former effectively being the latter multiplied by a sample-specific

constant, such that all benefits we report on weighted-adjusted CPM will apply to similarly

transformed couts. Adjusting counts across a set of species would require treating one species as

the "reference", and rescale counts for other species relative to it. Outside of the realm of formal

tests of differential expression, we suggest that wt.-adj.-CPM could be useful in phylogenetic

comparative analyses of expression at the gene level. One of the big challenges of these methods

is accounting for length differences between transcript/contig sets belonging to orthologous

genes, and the adjusted CPM appears to substantially outperform other metrics in this regard. We

should note that our adjustment method is relatively simple, at that it may be possible to adopt

more sophisticated approaches that leverage such information as sequence composition and

complexity for additional improvements.

We do not test here datasets amenable to differential expression analyses, and a counter-

argument to our findings of bias and noise is that neither should have a substantial effect, so long

as bias and deviations from map-to-reference expectations are consistent across replicates and

conditions. While we did find broad congruence in bias across samples, large differences in bias

among samples were frequent enough such that they will likely elevate false positive rates for

differential expression tests derived from de novo transcriptome assemblies. This effect can be

minimized by utilizing metrics that adjust for effective length differences. Finally, our results

clearly indicate that isoform-level analyses should be avoided.

By far the most unexpected findings from our study were the extremely high genotyping

error rates derived from transcriptome assemblies, specifically ST representations of TRNITY

assemblies. False negatives were due, in part, to the absence of an ST aligned to a putatively true

MR polymorphic site, which is not unexpected given the known fragmented nature of many *de*

novo transcriptome assemblies. As our mapping rates of ST to genomes was extremely high,

such FN are due to either the failure to assembly any contigs at all for a gene (i.e. missing genes),

or the incomplete coverage of genes not extending over some fraction of polymorphic sites. That

genotyping errors were biased towards the distal end of contigs suggest that assembly graphs are

getting broken at hard to assemble features containing low complexity and/or increased

polymorphism. When such graph-splitting occurs, the intervening polymorphic sites will, by

definition, not be covered by ST, leading to downwardly biased estimates of heterozygosity in *de*

novo assemblies. False negatives may have also been caused by the slightly reduced read depth

at homologous positions in read alignments to ST. That the reduction in read depth in the latter,

was, on average, small suggests that other factors may play a role. It may be that genotype

inference methods such at the GATK HAPLOTYPE CALLER that leverage haplotypic information,

may have poorer performance when iterating over alignments to relatively short assembly

contigs. Another possibility is that assembly errors may generate patterns that get flagged by

standard genotyping filters. In contrast, false positives appear to be driven by paralog collapses

given the bias towards heterozygous false positive sites. In the absence of a genome sequence, it

is hard to correct FN biases for any particular transcriptome assembly. In contrast, sophisticated

filtering methods might permit the learning of sequence features that produce false positives, and

to derive metrics that could mitigate the false positive rate.

While we document overall high error rates, the range of precision we observed was

broadly consistent with what was observed in the original paper that describes the LACE

algorithm (~ 80%) for converting transcriptome assemblies into SuperTranscripts (Davidson et

al., 2017). It is important to note that, in that paper, recall and false negative rates are not

reported, and genotyping performance is restricted to SuperTranscript genotype calls relative to

known variants in Genome in a Bottle Consortium high confidence regions for humans (Zook et

al., 2016). Such well-curated, filtered benchmark SNV sets are, by definition, not available for

non-model organisms with neither genomes nor extensive resequencing data. Results reported on

them are optimistic relative to what can be expected for organisms without such genomic

resources. For this reason and to create more realistic expectations for molecular ecologists, we

use genotypes obtained from mapping to reference genome as our truth set.

Even in the face of a high occurrence of false positive heterozygous genotype calls, false

negatives with respect to true heterozygous sites led to an overall negative bias in estimates of

heterozygosity estimates derived from ST relative to the gold standard MR approach, most

prominently in the wild *Mus* samples. This finding strongly suggests that the underlying site-

frequency spectrum derived from an ST assembly will be biased in ways that distort other

population genetic summary statistics typically reported in molecular ecological investigations

that utilize genotypes called from transcriptome assemblies. More broadly, the error rates we

document suggest that several lines of transcriptome-assembly based research are problematic.

First, erroneous heterozygous calls, when added to the redundancy problems described above,

will substantially reduce power and elevate false positive rates in allele-specific expression

analyses. Second, levels of genome-wide patterns of genetic diversity derived from

transcriptome assemblies will likely be negatively biased, at least for diploid species. The

distinction concerning ploidy is an important one because, in the absence of a genome or

karyotypic information, there can be no a priori expectation of how events such as whole-

genome or even chromosome-level duplication will impact genotyping inference. Thus, both

integration of diversity metrics across transcriptome assemblies with differing, unknown levels

of ploidy, and comparing transcriptome and genome assembly-derived metrics will confound

statistical method with biology and will lead to spurious inferences. Even when applied solely to

diploid transcriptome assemblies, as haplotypes will get assembled into different ST, the

expectation is that negative bias will be greater for more heterozygous genomes, such that

estimated levels of diversity across assemblies will artificially appear more similar than they are

in actuality, impeding researchers' ability to detect underlying patterns of variation.

While it is possible that a combination of algorithmic and annotation-based clustering

techniques can be employed to better collapse assembly redundancy, further research on this

topic is necessary. Whether or not more robust clustering techniques ameliorate bias, naïve

application of genotyping to ST derived from Trinity component "genes" will lead to high false

positive and false negative rates. Finally, the distortion of allele frequencies derived from

transcriptome assemblies strongly suggests that they will inject bias into analyses of population

structure, demographic inference, and efforts to detect positive selection.

Conclusion

Given the challenges associated with eco-evolutionary analyses that rely on de novo

transcriptome assemblies and the rapidly falling costs of long read sequencing, an unavoidable

recommendation for research teams launching new studies may be to generate a provisional

genome assembly, e.g. using a combination of Oxford Nanopore or PacBio long reads and

Illumina paired-end short reads (Austin et al., 2017; Marcionetti, Rossier, Bertrand, Litsios, &

Salamin, 2018; Michael et al., 2018; Nowoshilow et al., 2018), at least in some cases. Such an

assembly can be annotated with *ab initio* gene prediction tools and Illumina RNA-seq paired-end

reads. Given a reasonable quality annotated genome, fewer reads per sample would have to be

allocated for downstream expression analyses, and the cost savings from lighter RNA

sequencing would partially offset costs of genomic sequencing and *de novo* assembly.

Nevertheless, there are three reasons whole de novo transcriptome assembly might still be

attractive. First, genome assembly remains challenging and expensive species with large

genomes or with unusual genome structure precluding standard methods. Second, thousands of

RNA-seq data sets now exist for species without reference genomes, and it is highly unlikely that

genome assemblies will be carried out any time soon for anything but a small fraction of these

species. Third, analyses based upon genome assemblies will depend strongly on the quality of

the assembly. Even for genomes of reasonably high quality, protein coding genes may be

missing, either through problems with the annotation process or due to the fact that genes fall

into assembly gaps (Peona, Weissensteiner, & Suh, 2018). In other words, genome assembly in

not necessarily a panacea for all problems related to expression analyses. However analytically

challenging, there is great potential for employing de novo transcriptome assemblies in

interesting comparative analyses, so long as the conducted in an informed fashion—avoiding

analyses modes that suffer from fundamental biases, mitigating biases where possible for others,

and, always, placing appropriate caveats given the biases that will inevitably remain.

ACKNOWLEDGMENTS

The computations in this paper were run on the Odyssey cluster supported by the FAS Division

of Science, Research Computing Group at Harvard University. We thank members of the FAS

Informatics Group for discussion. We also thank Nadia Davidson and Alicia Oshlack for

feedback on an earlier version of this manuscript. This work was conducted on the traditional

territory of the Wampanoag and Massachusett peoples.

COMPETING INTERESTS

The authors have indicated that no competing interests exist.

DATA AVAILABILITY

Upon acceptance for publication, all transcriptome assemblies, SuperTranscript assemblies, and

vcf-format genotype files will be made available on Dryad.

AUTHOR CONTRIBUTIONS

The research project was designed by A. H. F., M. C., and T. B. S. Analyses were conducted by

A. H. F. with input from M.C. and T. B. S. A. H. F. wrote the paper with T. B. S. All authors

read and approved the final manuscript

REFERENCES

Austin, C. M., Tan, M. H., Harrisson, K. A., Lee, Y. P., Croft, L. J., Sunnucks, P., … Gan, H. M.

(2017). De novo genome assembly and annotation of Australia's largest freshwater fish,

the Murray cod (Maccullochella peelii), from Illumina and Nanopore sequencing read.

GigaScience, *6*(8), 1–6. doi: 10.1093/gigascience/gix063

Bankers, L., Fields, P., McElroy, K. E., Boore, J. L., Logsdon, J. M., & Neiman, M. (2017). Genomic

evidence for population-specific responses to co-evolving parasites in a New Zealand

freshwater snail. *Molecular Ecology*, *26*(14), 3663–3675. doi: 10.1111/mec.14146

Berens, A. J., Hunt, J. H., & Toth, A. L. (2015). Comparative transcriptomics of convergent

evolution: different genes but conserved pathways underlie caste phenotypes across

lineages of eusocial insects. *Molecular Biology and Evolution*, *32*(3), 690–703. doi:

10.1093/molbev/msu330

Birol, I., Jackman, S. D., Nielsen, C. B., Qian, J. Q., Varhol, R., Stazyk, G., … Jones, S. J. M. (2009).

De novo transcriptome assembly with ABySS. *Bioinformatics*, *25*(21), 2872–2877. doi:

10.1093/bioinformatics/btp367

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq

quantification. *Nature Biotechnology*, *34*(5), 525–527. doi: 10.1038/nbt.3519

Cokus, S. J., Gugger, P. F., & Sork, V. L. (2015). Evolutionary insights from de novo transcriptome

assembly and SNP discovery in California white oaks. *BMC Genomics*, *16*, 552. doi:

10.1186/s12864-015-1761-4

Davidson, N. M., Hawkins, A. D. K., & Oshlack, A. (2017). SuperTranscripts: a data driven

reference for analysis and visualisation of transcriptomes. *Genome Biology*, *18*(1). doi:

10.1186/s13059-017-1284-1

Davidson, N. M., & Oshlack, A. (2014). Corset: enabling differential gene expression analysis for

de novoassembled transcriptomes. *Genome Biology*, *15*(7), 410. doi: 10.1186/s13059-

014-0410-6

De Wit, P., & Palumbi, S. R. (2013). Transcriptome-wide polymorphisms of red abalone (Haliotis

rufescens) reveal patterns of gene flow and local adaptation. *Molecular Ecology*, *22*(11),

2884–2897. doi: 10.1111/mec.12081

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., … Regev, A.

(2011a). Full-length transcriptome assembly from RNA-Seq data without a reference

genome. *Nature Biotechnology*, *29*(7), 644–652. doi: 10.1038/nbt.1883

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., … Regev, A.

(2011b). Full-length transcriptome assembly from RNA-Seq data without a reference

genome. *Nature Biotechnology*, *29*(7), 644–652. doi: 10.1038/nbt.1883

Harr, B., Karakoc, E., Neme, R., Teschke, M., Pfeifle, C., Pezer, Ž., … Tautz, D. (2016). Genomic

resources for wild populations of the house mouse, Mus musculus and its close relative

Mus spretus. *Scientific Data*, *3*, 160075. doi: 10.1038/sdata.2016.75

Hornett, E. A., & Wheat, C. W. (2012). Quantitative RNA-Seq analysis in non-model species:

assessing transcriptome assemblies as a scaffold and the utility of evolutionary

divergent genomic reference species. *BMC Genomics*, *13*, 361. doi: 10.1186/1471-2164-

13-361

Kannan, S., Hui, J., Mazooji, K., Pachter, L., & Tse, D. (2016). Shannon: An Information-Optimal

de Novo RNA-Seq Assembler. *BioRxiv*, 039230. doi: 10.1101/039230

Kent, W. J. (2002). BLAT--the BLAST-like alignment tool. *Genome Research*, *12*(4), 656–664. doi:

10.1101/gr.229202. Article published online before March 2002

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low memory

requirements. *Nature Methods*, *12*(4), 357–360. doi: 10.1038/nmeth.3317

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, *9*(4), 357–359. doi: 10.1038/nmeth.1923

Le Gac, M., Metegnier, G., Chomérat, N., Malestroit, P., Quéré, J., Bouchez, O., … Chapelle, A.

(2016). Evolutionary processes and cellular functions underlying divergence in

Alexandrium minutum. *Molecular Ecology*, *25*(20), 5129–5143. doi: 10.1111/mec.13815

Li, B., & Dewey, C. N. (2011a). RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome. *BMC Bioinformatics*, *12*, 323. doi: 10.1186/1471-2105-

12-323

Li, B., & Dewey, C. N. (2011b). RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome. *BMC Bioinformatics*, *12*, 323. doi: 10.1186/1471-2105-

12-323

Li, B., Fillmore, N., Bai, Y., Collins, M., Thomson, J. A., Stewart, R., & Dewey, C. N. (2014).

Evaluation of de novo transcriptome assemblies from RNA-Seq data. *Genome Biology*,

15(12). doi: 10.1186/s13059-014-0553-5

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Genome Project Data

Processing, S. (2009). The Sequence Alignment/Map format and SAMtools.

Bioinformatics (Oxford, England), *25*(16), 2078–2079. doi:

10.1093/bioinformatics/btp352;

Li, K., Wang, H., Cai, Z., Wang, L., Xu, Q., Lövy, M., … Nevo, E. (2016). Sympatric speciation of

spiny mice, *Acomys* , unfolded transcriptomically at Evolution Canyon, Israel.

Proceedings of the National Academy of Sciences, *113*(29), 8254–8259. doi:

10.1073/pnas.1608743113

Liu, J., Li, G., Chang, Z., Yu, T., Liu, B., McMullen, R., … Huang, X. (2016). BinPacker: Packing-

Based De Novo Transcriptome Assembly from RNA-seq Data. *PLOS Computational*

Biology, *12*(2), e1004772. doi: 10.1371/journal.pcbi.1004772

MacManes, M. D. (2014). On the optimal trimming of high-throughput mRNA sequence data.

Bioinformatics and Computational Biology, *5*, 13. doi: 10.3389/fgene.2014.00013

MacManes, M. D. (2018). The Oyster River Protocol: a multi-assembler and kmer approach for

de novo transcriptome assembly. *PeerJ*, *6*. doi: 10.7717/peerj.5428

Malik, L., Almodaresi, F., & Patro, R. (2018). Grouper: graph-based clustering and annotation for

improved de novo transcriptome analysis. *Bioinformatics (Oxford, England)*, *34*(19),

3265–3272. doi: 10.1093/bioinformatics/bty378

Marcionetti, A., Rossier, V., Bertrand, J. A. M., Litsios, G., & Salamin, N. (2018). First draft

genome of an iconic clownfish species (Amphiprion frenatus). *Molecular Ecology*

Resources, *18*(5), 1092–1101. doi: 10.1111/1755-0998.12772

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo, M. A.

(2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Res*, *20*(9), 1297–1303. doi:

10.1101/gr.107524.110

Mi, H., Muruganujan, A., Casagrande, J. T., & Thomas, P. D. (2013). Large-scale gene function

analysis with the PANTHER classification system. *Nature Protocols*, *8*(8), 1551–1566. doi:

10.1038/nprot.2013.092

Michael, T. P., Jupe, F., Bemm, F., Motley, S. T., Sandoval, J. P., Lanz, C., … Ecker, J. R. (2018).

High contiguity Arabidopsis thaliana genome assembly with a single nanopore flow cell.

Nature Communications, *9*(1), 541. doi: 10.1038/s41467-018-03016-2

Nowoshilow, S., Schloissnig, S., Fei, J.-F., Dahl, A., Pang, A. W. C., Pippel, M., … Myers, E. W.

(2018). The axolotl genome and the evolution of key tissue formation regulators.

Nature, *554*(7690), 50–55. doi: 10.1038/nature25458

Ockendon, N. F., O'Connell, L. A., Bush, S. J., Monzón-Sandoval, J., Barnes, H., Székely, T., …

Urrutia, A. O. (2016). Optimization of next-generation sequencing transcriptome

annotation for species lacking sequenced genomes. *Molecular Ecology Resources*, *16*(2),

446–458. doi: 10.1111/1755-0998.12465

Pankey, M. S., Minin, V. N., Imholte, G. C., Suchard, M. A., & Oakley, T. H. (2014). Predictable

transcriptome evolution in the convergent and complex bioluminescent organs of squid.

Proceedings of the National Academy of Sciences, *111*(44), E4736–E4742. doi:

10.1073/pnas.1416574111

Peona, V., Weissensteiner, M. H., & Suh, A. (2018). How complete are "complete" genome

assemblies?—An avian perspective. *Molecular Ecology Resources*, *18*(6), 1188–1195.

doi: 10.1111/1755-0998.12933

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics*, *26*(6), 841–842. doi: 10.1093/bioinformatics/btq033

Roda, F., Mendes, F. K., Hahn, M. W., & Hopkins, R. (2017). Genomic evidence of gene flow

during reinforcement in Texas Phlox. *Molecular Ecology*, *26*(8), 2317–2330. doi:

10.1111/mec.14041

Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., … Galtier, N.

(2014). Comparative population genomics in animals uncovers the determinants of

genetic diversity. *Nature*, *515*(7526), 261–263. doi: 10.1038/nature13685

Schwentner, M., Combosch, D. J., Pakes Nelson, J., & Giribet, G. (2017). A Phylogenomic

Solution to the Origin of Insects by Resolving Crustacean-Hexapod Relationships. *Current

Biology*, *27*(12), 1818-1824.e5. doi: 10.1016/j.cub.2017.05.040

Siepel, A. (2005). Evolutionarily conserved elements in vertebrate, insect, worm, and yeast

genomes. *Genome Research*, *15*(8), 1034–1050. doi: 10.1101/gr.3715005

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics*, *31*(19), 3210–3212. doi: 10.1093/bioinformatics/btv351

Singh, R., Jones, T., Wai, C. M., Jifon, J., Nagai, C., Ming, R., & Yu, Q. (2018). Transcriptomic

analysis of transgressive segregants revealed the central role of photosynthetic capacity

and efficiency in biomass accumulation in sugarcane. *Scientific Reports*, *8*(1), 4415. doi:

10.1038/s41598-018-22798-5

Smith-Unna, R., Boursnell, C., Patro, R., Hibberd, J. M., & Kelly, S. (2016). TransRate: reference-

free quality assessment of de novo transcriptome assemblies. *Genome Research*, *26*(8),

1134–1144. doi: 10.1101/gr.196469.115

Soneson, C., Love, M. I., & Robinson, M. D. (2015). Differential analyses for RNA-seq: transcript-

level estimates improve gene-level inferences. *F1000Research*, *4*, 1521. doi:

10.12688/f1000research.7563.1

Song, L., & Florea, L. (2015). Rcorrector: efficient and accurate error correction for Illumina

RNA-seq reads. *GigaScience*, *4*, 48. doi: 10.1186/s13742-015-0089-y

Tepolt, C. K., & Palumbi, S. R. (2015). Transcriptome sequencing reveals both neutral and

adaptive genome dynamics in a marine invader. *Molecular Ecology*, *24*(16), 4145–4158.

doi: 10.1111/mec.13294

Todd, E. V., Black, M. A., & Gemmell, N. J. (2016). The power and promise of RNA-seq in ecology

and evolution. *Molecular Ecology*, *25*(6), 1224–1241. doi: 10.1111/mec.13526

Veilleux, H. D., Ryu, T., Donelson, J. M., van Herwerden, L., Seridi, L., Ghosheh, Y., … Munday, P.

L. (2015). Molecular processes of transgenerational acclimation to a warming ocean.

Nature Climate Change, *5*(12), 1074–1078. doi: 10.1038/nclimate2724

Velotta, J. P., Wegrzyn, J. L., Ginzburg, S., Kang, L., Czesny, S., O'Neill, R. J., … Schultz, E. T.

(2017). Transcriptomic imprints of adaptation to fresh water: parallel evolution of

osmoregulatory gene expression in the Alewife. *Molecular Ecology*, *26*(3), 831–848. doi:

10.1111/mec.13983

Vijay, N., Poelstra, J. W., Künstner, A., & Wolf, J. B. W. (2013). Challenges and strategies in

transcriptome assembly and differential gene expression quantification. A

comprehensive *in silico* assessment of RNA-seq experiments. *Molecular Ecology*, *22*(3),

620–634. doi: 10.1111/mec.12014

Wu, T. D., & Watanabe, C. K. (2005). GMAP: a genomic mapping and alignment program for

mRNA and EST sequences. *Bioinformatics*, *21*(9), 1859–1875. doi:

10.1093/bioinformatics/bti310

Zook, J. M., Catoe, D., McDaniel, J., Vang, L., Spies, N., Sidow, A., … Salit, M. (2016). Extensive

sequencing of seven human genomes to characterize benchmark reference materials.

Scientific Data, *3*. doi: 10.1038/sdata.2016.25

1061

 TABLES

1063   **Table 1**. Counts of missing genes by assembler for three *Mus* samples

| | Sample | | | | | |
| | MDC | | BALB/c | | FRA | |
Missingness pattern[a]	Unfiltered	Filtered[b]	Unfiltered	Filtered[b]	Unfiltered	Filtered[b]
PPP	11055 (99.6)	6646 (79.8)	14777 (98.7)	9167 (87.2)	16350 (99.3)	10170 (90.6)
MPP	1 (0.0)	46 (0.6)	47 (0.3)	197 (1.9)	7 (0.0)	106 (0.9)
PMP	2 (0.0)	1341 (16.1)	15 (0.1)	638 (6.1)	12 (0.1)	379 (3.4)
PPM	20 (0.2)	31 (0.4)	27 (0.2)	52 (0.5)	54 (0.3)	81 (0.7)
PMM	0 (0)	18 (0.2)	6 (0.0)	40 (0.4)	3 (0.0)	33 (0.3)
MPM	2 (0.0)	21 (0.3)	13 (0.1)	58 (0.6)	5 (0.0)	28 (0.2)
MMP	0 (0)	45 (0.5)	7 (0.0)	205 (1.9)	0 (0)	226 (2.0)
MMM	17 (0.2)	181 (2.2)	78 (0.5)	156 (1.5)	32 (0.2)	198 (1.8)

[a] Present (P) and missing (M) codes reflect the following assembler order: Trinity, Shannon, BinPacker.
[b] Restricted to protein-coding genes with minimum transcript length of 200 and $TPM \geq 1$, based upon contigs with $TPM \geq 1$ and a hit to Uniref90.

1064

1065

1066   **Table 2.** For filtered assemblies, median genomic alignment-based transcript per million (TPM)

1067   and phastCons scores for ENSEMBL genes that are recovered by all de novo transcriptome

1068   assembly methods (All), recovered by 1-2 of the 3 methods (Some), and recovered by no

1069   assembler (None). Flagged tests are significant after Bonferroni correction; * $P < 0.005$, ** P

1070 <0.0005, and *** $P < 1.0 \times 10^{-10}$.

| | TPM | | | | | | phastCons | | | | | |
Sample	All	Some	None	All vs. Some	All vs. None	Some vs. None	All	Some	None	All vs. Some	All vs. None	Some vs. None
MDC	16.8	13.1	15.8	***	NS	NS	0.41	0.41	0.30	NS	***	***
BALB/c	19.0	3.2	2.0	***	***	**	0.44	0.36	0.34	***	***	*
FRA	24.5	2.4	4.2	***	***	**	0.43	0.37	0.29	***	***	***

1071

1072   **Table 3**. Selected performance metrics comparing genotypes derived from SuperTranscript

1073   collapse of TRINITY de novo transcriptome assemblies with those derived from a map-to-

1074   reference genome approach. See **METHODS** for information on metric calculations. Values in

1075   parentheses are after converting SuperTranscripts with no protein hit to UNIREF90 to NA. The

1076   full table of metrics is available as Supplementary Table 7.

Metric	Sample					
	MDC	BALB/c	FRA	DEU	IRN	KZK
Error rate	0.870 (0.892)	0.500 (0.477)	0.359 (0.453)	0.376	0.336	0.301
Error-MrefAlleleIncluded	1.000 (1.000)	0.898 (0.898)	0.969 (0.971)	0.977	0.969	0.959
False negative (FN)	0.662 (0.754)	0.389 (0.412)	0.260 (0.396)	0.265	0.246	0.239
FN-noalign	0.728 (0.868)	0.745 (0.881)	0.585 (0.860)	0.585	0.572	0.668
FN-het	0.623 (0.752)	0.742 (0.885)	0.259 (0.451)	0.259	0.242	0.268
False positive (FP)	0.609 (0.558)	0.179 (0.110)	0.129 (0.091)	0.147	0.115	0.078
FP-NAtoFixed	0.118 (0.094)	0.053 (0.051)	0.027 (0.019)	0.028	0.026	0.024
FP-NAtoHet	0.882 (0.906)	0.919 (0.928)	0.916 (0.923)	0.909	0.917	0.915
FP-indel	0.188 (0.196)	0.063 (0.048)	0.184 (0.146)	0.225	0.194	0.070
Recall	0.334 (0.243)	0.609 (0.587)	0.735 (0.602)	0.731	0.750	0.757
Precision	0.176 (0.161)	0.735 (0.826)	0.829 (0.856)	0.805	0.849	0.897

1077

Table 4. Correlation between map-to-reference and map-to-denovo-transcriptome expression at

the gene level, for both RSEM and KALLISTO, based upon CPM, TPM, and weighted-adjusted

CPM. See equation in methods for details regarding the last of these. Correlations exclude

expressed reference genes where there are not associated denovo assembly contigs, i.e. genes

that are missing from the assembly. Correlations are calculated for entries with TPM > 0 for

either map-to-reference or map-to-denovo assembly. Bold indicates strongest correlation across

metrics and estimation algorithms for a particular sample-assembler combination. Bold italics

indicate the strongest correlation across all estimators and assembly algorithms for a particular

sample.

1087

Sample	CPM		TPM		Weighted EL-adjusted CPM	
	RSEM	kallisto	RSEM	kallisto	RSEM	kallisto
MDC						
Trinity	0.853	0.858	0.756	0.675	***0.901***	0.823
Shannon	0.629	**0.833**	0.567	0.662	0.680	0.784
BinPacker	0.824	0.825	0.796	0.754	**0.877**	0.783
BALB/c						
Trinity	0.869	0.881	0.818	0.819	***0.897***	0.863
Shannon	0.819	0.859	0.790	0.815	0.854	0.826
BinPacker	0.835	0.846	0.801	0.803	0.870	0.824
Massif Central						
Trinity	0.850	0.861	0.809	0.790	**0.903**	0.841
Shannon	0.849	0.857	0.812	0.785	***0.906***	0.819
BinPacker	0.815	0.821	0.808	0.786	0.872	0.793
Germany						
Trinity	0.848	0.859	0.804	0.788	**0.903**	0.838
Shannon	0.847	0.855	0.812	0.784	***0.905***	0.813
BinPacker	0.811	0.818	0.808	0.784	0.873	0.789
Iran						
Trinity	0.853	0.862	0.806	0.785	***0.906***	0.842
Shannon	0.847	0.854	0.804	0.777	**0.904**	0.817
BinPacker	0.809	0.816	0.799	0.773	0.869	0.787
Kazakhstan						
Trinity	0.852	0.862	0.810	0.791	**0.903**	0.841
Shannon	0.852	0.858	0.814	0.793	***0.907***	0.824
BinPacker	0.816	0.821	0.809	0.788	0.872	0.795

1088

 FIGURE LEGENDS

Figure 1. Coverage of individual *Mus* reference transcripts and genes by the TRINITY assembly.

Coverage of all expressed reference sequences by the unfiltered assembly is defined by BLAT

alignments with a minimum percent identity of 95% for **(a)** individual contigs with the highest

scoring alignment, **(b)** summed coverage of all BLAT hits and **(c)** gene-level coverage as the

expression-weighted average of summed coverage across isoforms. Comparable coverage

estimates for only protein-coding transcripts/genes of minimum length 200bp, by contigs with

TPM $\geq$ 1 and an open-reading frame alignment to UNIREF90 are represented by **(d-f)**. Blue lines

indicate smoothed, fitted trends on the coverage-TPM relationship, while purple lines indicate

the empirical cumulative frequency distribution on TPM.

1099

1100 **Figure 2.** Violin plots of **(a)** TPM and **(b)** phastCons scores for protein coding genes with a

1101 minimum transcript length of 200bp, which are recovered in all three transcriptome assemblers

1102 (TRINITY, SHANNON, BINPACKER), 1-2 of three assemblers, or that are missing from all three

1103 assemblers. Missingness is determined using filtered transcriptome assemblies, that only include

1104 contigs with TPM $\geq$ 1 and an open-reading frame alignment to UNIREF90.

1105

Figure 3. Base composition by TRINITY contig (left panels), and sequence reads (right panels)

for **(a)** *M. musculus* dendritic cells, **(b)** *M. musculus* BALB/c strain brain, **(c)** wild outcrossed *M.*

musculus brain, **(d)** *D. melanogaster* head, **(e)** *T. nigroviridis* brain, and **(f)** *X. tropicalis* kidney.

For each panel, each y-axis row represents a single contig or read. Sequence panels for reads are

based upon 100,000 randomly sampled mapped reads. Intersections of contigs and reads with

annotations are based upon mapping with GMAP and HISAT2, respectively (see **Methods** for

details). For visualization purposes, sequence proportions are arranged from the bottom to the

top of a panel, sorted in descending order, consecutively by CDS, UTR, non-coding, non-coding

intron, intron, and intergenic proportions.

1115

1116   **Figure 4.** Frequency of proportional overlaps between contigs, based upon BLAT all-by-all hits,

1117   for known *Mus* single isoform genes, for three 8-individual pools of wild *M. m. domesticus* (Iran;

1118   Massif Central, France and Germany) and one 8-individual pool of *M. m. musculus*

1119   (Kazakhstan), for **(a)** TRINITY, **(b)** SHANNON and **(c)** BINPACKER. Estimates of Watterson's θ are

1120   those reported from whole genome sequence data in Harr et al. (2016).

1121

Figure 5. (a) For FRA TRINITY assembly, hexbin 2D histogram of the proportion of contig

lengths comprising BLAT pairwise alignments resulting from all-by-all searches. **(b)** Example

alignment to the *Mus* genome, to a protein-coding transcript of the GDA gene, of nearly

identically sized (1220 bp) FRA TRINITY contigs. These contigs have a BLAT-based 99.8 %

identity in an alignment equal to the entirety of the contig length, and which are assembled from

different TRINITY components (i.e. "genes"), as indicated by the red box around their labels. The

contigs differ by only two adjacent SNPs (A and T), and also share fixed C (far left) and G (far

right) differences relative to the reference genome. Thus, different Trinity components have

assembled different haplotypes.

1131

1132   **Figure 6.** Examples of map-to-reference and SuperTranscript derived genotype combinations

1133   that produce particular error classes.

1134

1135 **Figure 7.** Heterozygosity per base for exonic biallelic SNV variants called relative to the *Mus*

1136 reference genome using the GATK best practices pipeline versus those obtained from

1137 supertranscripts constructed from TRINITY assemblies using scripts provided with the TRINITY v.

1138 2.6.5 package. See **Methods** for details concerning how exonic coordinates were obtained for

1139 SuperTranscripts.

1140

Figure 8. For the FRA data set, comparison of gene-level expression summed across BLAT

alignments of TRINITY contigs to reference transcripts (grouped by gene), to reference gene

expression in terms of **(a)** CPM, **(b)** TPM, and **(c)** weighted-adjusted CPM that accounts for

differences in effective length estimates between methods. For CPM, TPM, and weighted-

adjusted CPM, 0.05, 0.01 and 0.05 are added to values prior to log-transformation, in order to

visualize instances where expression = 0 for one of the methods being compared. The red line

represents unity, the blue line is the smoothed fit to data. We exclude all cases where an

expressed reference gene has no TRINITY contigs aligned to it, such that it is missing from the de

novo transcriptome assembly.

1150

1151 **Figure 9.** For the FRA data set, for genes with non-zero expression in both transcriptome

1152 assembly and map-to-reference, log-fold change between TRINITY and map-to-reference

1153 expression estimates as a function of difference in effective length (EL) for **(a)** CPM, **(b)** TPM,

1154 and **(c)** weighted-adjusted CPM.

1155

1156 **Figure 10.** Frequency of bias effect sizes for gene-level expression estimated by CPM, TPM,

1157 and weighted-adjusted CPM for a TRINITY-based comparison of the FRA and DEU samples.

(a)

(b)

Genome sequence	**G**	**G**	**G**	**G**	**G**	**G**	**G**	**G**
MR genotype	**G/G**	**G/G**	**G/G**	**A/G**	**A/A**	**A/G**	**A/G**	**A/A**
ST sequence	**G**	**G**	**G**	**AT**	**A**	**G**	**_**	**G**
ST genotype	**A/G**	**G/GT**	**T/T**	**A/AT**	**A/T**	**G/G**	**_**	**G/T**
STs aligned to genome	**1**	**1**	**1**	**1**	**1**	**1**	**0**	**2**
Error-MrefAlleleIncluded	●	●		●	●			
Error-MultipleSuperTranscripts								●
Error-Snv2Indel		●		●				
FN						●	●	
FN-noalign							●	
FN-het						●	●	
FP	●	●	●					
FP-het	●	●			●			
FP-NAtoFixed			●					
FP-NAtoHet	●	●						
FP-indel		●						