Classification: Biological Sciences, Evolution

Genetic structure of prey populations underlies the geographic mosaic of arms race

coevolution

Authors: Michael T.J. Hague[a*], Amber N. Stokes[b], Chris R. Feldman[c], Edmund D. Brodie, Jr.[d],

Edmund D. Brodie III[e]

Affiliations:

[a] Division of Biological Sciences, University of Montana, Missoula, MT, 59812.

[b] Department of Biology, California State University, Bakersfield, CA, 93311.

[c] Department of Biology, University of Nevada, Reno, NV, 89557.

[d] Department of Biology, Utah State University, Logan, UT, 84322.

[e] Department of Biology, University of Virginia, Charlottesville, VA, 22904.

[*] Corresponding author:

Michael T.J. Hague

Division of Biological Sciences, University of Montana

32 Campus Dr. HS 104, Missoula, MT 59812

Tel: (406) 243-2182

Email: michael.hague@mso.umt.edu

Keywords: coevolution, arms race, geographic mosaic, tetrodotoxin, $Na_V1.4$

ABSTRACT

Reciprocal adaptation is the hallmark of arms race coevolution, but the symmetry of

evolutionary change between interacting species is often untested, even in the best-studied

battles of natural enemies. We tested whether prey and predator exhibit symmetrical local co-

adaptation in the example of a geographic mosaic of coevolution between toxic newts (*Taricha*

granulosa) and resistant garter snakes (*Thamnophis sirtalis*). Prior work showing a tight

correlation between levels of newt toxin and snake resistance is regarded as textbook evidence of

the intense arms race between natural enemies. Here, we similarly found that toxin and resistance

are functionally matched in prey and predator populations, further suggesting that mosaic

variation in the armaments of both species results from the local pressures of reciprocal

selection. Contrary to conventional wisdom, however, we found that local variation in newt toxin

is best predicted by neutral population divergence rather than the resistance of co-occurring

predators. Snake resistance, on the other hand, is clearly explained by local levels of prey toxin.

Prey populations seem to structure variation in defensive toxin levels across the geographic

mosaic, which in turn determines selection on predator resistance. Exaggerated armaments

suggest that coevolution occurs in certain hotspots, but our results imply that neutral processes

like gene flow—rather than reciprocal adaptation—structure the greatest source of variation

across the landscape. This pattern supports the predicted role of "trait remixing" in the

geographic mosaic of coevolution, the process by which non-adaptive forces dictate spatial

variation in the interactions among species.

SIGNIFICANCE STATEMENT

When the weapons of natural enemies like prey toxins and predator resistance are

matched across the geographic landscape, they are usually presumed to result from arms race

coevolution. In the textbook example of an arms race, matched levels of newt toxin and garter

snake resistance have long been regarded as evidence of such local co-adaptation. To the

contrary, we found that local variation in newt toxicity is best explained by the neutral

geographic structure of newt populations. This spatial variation of prey in turn dictates local

selection on garter snakes, structuring the geographic pattern of predator resistance. These results

demonstrate how landscape patterns of phenotypic variation are determined by a mixture of

natural selection, historical biogeography, and gene flow that comprise the geographic mosaic of

coevolution.

INTRODUCTION

Coevolutionary dynamics result from the reciprocal selection generated by ecological interactions between species, which drives adaptation and counter-adaptation in the traits mediating interactions (1–4). Because the nature of species interactions and their fitness consequences vary spatially, heterogeneity in the form of reciprocal selection is expected to generate a geographic mosaic of coevolution, such that traits at the phenotypic interface of coevolution covary due to local conditions (2, 5–7). Coevolving species are expected to exhibit matched trait variation across the landscape, for example, in seed traits and their predators (5, 7) or host and pathogen genotypes (8, 9). As such, local co-adaptation is often presumed when a pair of interacting species share matched abilities throughout their shared range (1, 2, 4).

However, a geographic mosaic of matched phenotypes may not result solely from reciprocal co-adaptation (10, 11). A simpler, non-adaptive explanation for matched trait variation involves the spatial population structure and ancestry of each species. Common barriers to dispersal or a shared biogeographic history, for example, could structure phenotypic divergence congruently in co-occurring species (12). Populations are subject to selectively neutral processes, such as drift and gene flow, that continually alter the spatial distribution of alleles and traits at the phenotypic interface, a process termed "trait remixing" in the geographic mosaic theory of coevolution (2, 10, 13). Only when trait variation deviates from the neutral expectations of population structure in both species can we infer local co-adaptation in the geographic mosaic of coevolution (10). Otherwise, phylogeography, drift, and gene flow provide a parsimonious explanation for patterns of divergence across the landscape.

In the geographic mosaic of arms coevolution, populations of rough-skinned newts (*Taricha granulosa*) and common garter snakes (*Thamnophis sirtalis*) tend to have matched

levels of prey toxin and predator resistance, a pattern generally interpreted as evidence of local

co-adaptation (11, 14–16). We tested whether this inferred signature of coevolution can instead

be explained by the population structure of either species. In western North America, rough-

skinned newts secrete the deadly neurotoxin tetrodotoxin (TTX), which binds to the outer pore of

voltage-gated sodium channels (Na_V) and prevents the initiation of action potentials (17, 18).

Common garter snakes exhibit resistance to TTX that is largely due to specific amino acid

substitutions in the fourth domain pore-loop (DIV p-loop) of the skeletal muscle sodium channel

($Na_V1.4$) that disrupt toxin-binding (Fig. 1) (19, 20). Channel-level TTX resistance conferred by

each allele in the DIV p-loop is tightly correlated with muscle and whole-animal levels of

phenotypic resistance (19–21). TTX-resistant alleles occur at high frequency in coevolutionary

"hotspots" with highly toxic newts, but are largely absent in surrounding "coldspots" where

newts are non-toxic (22). Because quantitative levels of newt toxin and snake resistance are

functionally matched in most populations, the geographic mosaic of coevolution is thought to

primarily consist of co-adaptations to local pressures in the arms race (11, 14). If true, we predict

that population divergence in both prey toxin and predator resistance represent adaptive

deviations from the selectively neutral expectations of population structure and ancestry.

We conducted fine-scale population sampling of newts (*n*=138) and garter snakes

(*n*=169) along a latitudinal transect of nine locations on the Pacific Coast in Washington and

Oregon (USA) that spans the geographic mosaic (Fig. 1, Supplemental Table S1), ranging from

low ancestral levels of newt toxin and snake resistance (northern Washington) to a hotspot of

extreme escalation in both species (central Oregon). We focused our sampling in Washington

and Oregon, because other regions of sympatry, particularly California, contain additional TTX-

bearing newt species (e.g., *Ta. torosa*) and resistant garter snakes (e.g., *Th. atratus*) that

confound inferences of coevolution between discrete pairs of species. At each sampling site in

the mosaic, we characterized levels of TTX in newt populations and TTX resistance in garter

snakes, including whole-animal phenotypic resistance and $Na_V1.4$ channel genotypes. We then

tested whether trait divergence in the weaponry of each species deviates from the neutral

expectations of geographic population structure using genome-wide single nucleotide

polymorphisms (SNPs). If the geographic mosaic of prey toxin and predator resistance results

from co-adaptation to local pressures in the arms race, then trait divergence for each species

should be predicted solely by the local armaments of its natural enemy. Alternatively, if mosaic

variation in toxin and/or resistance is a selectively neutral result of population structure, then trait

divergence should be better predicted by genomic SNP variation (23, 24).

RESULTS AND DISCUSSION

Matched trait variation implies local co-adaptation. Geographic patterns of newt TTX and

snake resistance were broadly consistent with previous work indicating arms race coevolution

has led to the closely matched phenotypes in each species (11, 14). TTX levels ($\mu g/cm^2$) of newts

varied by population (ANOVA; $F[8,114]=37.43$, $p<0.001$) and by sex ($F[1,114]=4.37$, $p=0.039$)

along the latitudinal transect (Fig. 1; Supplemental Table S1). TTX resistance (50% MAMU

dose) of snakes also varied by population (according to non-overlapping 95% confidence

intervals; Fig. 1; Supplemental Table S1) and was closely correlated with prey toxin (regression

of distance matrices, $p=0.021$; Table 1). The presence of TTX-resistant alleles in the $Na_V1.4$

channel co-varied with phenotypic resistance in garter snakes, such that pairwise F_{ST} divergence

at the DIV p-loop was correlated with population divergence in phenotypic resistance (Mantel

test, $r=0.47$, $p=0.032$).

126   Quantitative estimates of prey toxin and predator resistance at each locality showed

127 functional overlap in the armaments of each species, implying that locally varying reciprocal

128 selection occurs throughout the geographic mosaic (Fig. 2) (11). For each locality, we estimated

129 whole-newt levels of TTX (mg) and the dose of TTX (mg) required to reduce the performance of

130 co-occurring snakes by 15%, 50%, and 85% (see Methods). The TTX dose required to reduce

131 snake performance by 50% is considered a perfect functional match between newt TTX and

132 snake resistance. The 15% and 85% doses delimit the range of functional overlap between toxin

133 and resistance, outside of which prey and predator are considered so mismatched that variable

134 fitness outcomes and reciprocal selection are unlikely to occur (11, 15, 16). Under this

135 framework, each locality we sampled exhibited some degree of overlap in the phenotypic

136 distributions of newt TTX and snake resistance, indicating potential for reciprocal selection

137 between prey and predator. Interestingly, all point estimate comparisons of TTX and resistance

138 fell below the 50% line, suggesting that garter snakes on average tend to have higher levels of

139 resistance than the toxins of co-occurring newts, a pattern consistent with the prediction that

140 predators experience intense selection when prey are deadly (see below) (3). Nonetheless, under

141 this framework of previous work, our results imply that matched trait variation in the geographic

142 mosaic of coevolution may be a result of reciprocal selection. Next, we characterized neutral

143 genomic variation for each species to test whether population genetic structure can alternatively

144 explain this observed pattern of matched weaponry.

146 **Prey and predator populations differ in geographic structure.** We used neutral SNPs

147 generated from double digest restriction-site associated DNA sequenced (ddRADseq; see

148 Methods) to characterize the population structure of prey and predator, which served as a neutral

expectation for trait divergence in the geographic mosaic of coevolution. Global F_{ST} values for

newts (F_{ST}=0.068, 95% CI [0.065, 0.069]) and snakes (F_{ST}=0.070, 95% CI [0.067, 0.075]) were

similar, and both species exhibited a pattern of isolation-by-distance (IBD) along the transect

(distance-based redundancy analysis; newts, F=-38.528, p=0.002; snakes, F=22.021, p=0.001).

This pattern is consistent with previous work suggesting limited population structure and a recent

northward post-glacial expansion of both species (25–27). Pairwise estimates of F_{ST}, however,

revealed subtle differences in the geographic population structure of prey and predator

(Supplementary Table S4), a pattern that was also supported by principal coordinate (PCoA) and

Bayesian clustering (STRUCTURE) analyses (Fig. 3, Supplemental Fig. S1). In particular, the

major axis of variation from the PCoA (PCo1) and STRUCTURE revealed limited structure

among newt populations along the transect, whereas garter snakes showed evidence of

population subdivision near the Washington-Oregon border. For example, the STRUCTURE plot

for snakes indicates that subdivision between the two most likely genetic clusters (K=2) occurs

just south of the Columbia River near the Warrenton locality in Oregon. Because pairwise F_{ST}

estimates, the PCoAs, and STRUCTURE produced very similar results, yet rely on different

assumptions, we feel confident in our characterization of population structure for each species.

Finally, we tested whether these patterns of population structure provide a more parsimonious

explanation than local co-adaptation for the geographic mosaic of matched trait variation

between prey and predator.

Asymmetric evidence of local adaptation in prey and predator. Trait divergence in newt TTX

was best predicted by population genetic structure, not predator resistance, suggesting that

reciprocal co-adaptation cannot fully explain the geographic mosaic of matched toxin and

resistance. We generated distance matrices to test whether phenotypic divergence in one species

(e.g., newt TTX levels) is best explained by (1) neutral genomic divergence (pairwise F_{ST};

Supplemental Table S4) or (2) phenotypic divergence in the natural enemy (snake resistance). In

univariate regressions, population divergence in the TTX level of newts was strongly predicted

by neutral F_{ST} divergence ($R^2=0.414$), as well as TTX resistance of garter snakes ($R^2=0.274$;

Table 1). F_{ST} divergence remained significant in the multiple regression, indicating that

population structure of newts predicts TTX levels, even after controlling for TTX resistance of

the predator (which was at best marginally significant; p=0.065). Unlike newt TTX, garter snake

resistance was strictly predicted by the prey toxin and not neutral F_{ST} divergence. Both

phenotypic resistance and F_{ST} divergence at the DIV p-loop (the site of toxin-binding in $Na_V1.4$)

were uncorrelated with neutral F_{ST} values in garter snakes (Table 1). These results indicate that

neutral genetic divergence is a key determinant of population differences in prey toxin levels,

which in turn predicts differences in TTX resistance among predator populations. The

geographic structure of newt populations appears to be so influential to spatial dynamics that

divergence in garter snake phenotypic resistance and F_{ST} at the DIV p-loop are both significantly

predicted by neutral F_{ST} divergence of newts (Table 1).

Clinal variation in TTX levels of newts is highly congruent with neutral genomic

structure based on the major axis of variation from the PCoA (PCo1; Fig. 4) and Bayesian

clustering analyses (Supplemental Fig. S2). TTX resistance of garter snakes, in contrast, clearly

deviates from neutral expectations to track variation in prey toxin. Cline-fitting analyses show

that prey toxin levels and predator resistance are tightly matched along the 611 km transect; the

geographic center points of each cline are located just 64 km apart and do not differ statistically.

The cline center of TTX-resistant alleles in snakes is also located nearby, although it differed

statistically from the center of newt TTX. Despite similar phenotypic clines for both species,

variation in levels of newt toxin showed an even tighter match to clinal variation in neutral

population structure. The center points of the TTX and neutral clines were located only 19 km

apart. PCo1 from the PCoA was a strong predictor of variation in TTX levels (linear model; t-

value=5.682, p<0.001), even after controlling for the effect of TTX resistance of garter snakes

(Supplemental Table S6, model 3). Conversely, variation in phenotypic resistance and TTX-

resistant alleles in snakes both deviated significantly from the neutral cline (Fig. 4), such that

resistance was not predicted by PCo1 or 2 from the PCoA (Supplemental Table S6, model 3).

The center points of the snake phenotypic resistance and neutral clines were located a distant 310

km apart.

Implications for the geographic mosaic of arms race coevolution. Levels of prey toxin and

predator resistance are tightly matched across the landscape, but this pattern does not appear to

primarily result from local arms race co-adaptation. Although predator resistance is

geographically structured by a signature of local adaptation to prey, levels of the prey toxin are

structured in tight accordance with neutral population divergence. These results imply that local

variation in the newt toxin across the geographic mosaic is best explained by non-adaptive

processes, such as drift and historical biogeography, rather than local variation in selection from

predators. For example, latitudinal patterns of newt TTX and neutral divergence are both

consistent with a signature of isolation-by-distance along the transect (25, 26).

The asymmetric signatures of adaptation we observed in prey and predator may reflect

differences in the mechanisms that underlie phenotypic variation in each species. Genes

associated with TTX biosynthesis have yet to be discovered; however, toxin production in newts

likely requires a complicated biosynthetic pathway. For example, biosynthesis of a similar

neurotoxin found in puffer fish, saxitoxin (STX), involves gene expression in a cluster of up to

26 genes (28). Some researchers suggest exogenous factors, like environmentally-derived

precursors, may also affect the ability of newts to synthesize or sequester TTX (29, 30). On the

other hand, TTX resistance in garter snakes is largely due to a small number of amino acid

changes to the p-loops of the $Na_V1.4$ channel (19–22). These large-effect mutations could make

TTX resistance more evolutionarily labile than the defensive toxins in newts, permitting rapid

local adaptation in predator populations (11, 21).

Asymmetric evolutionary changes could also arise from a selective imbalance associated

with the interactions between prey and predator. In antagonistic interactions, the species under

more intense selection is generally expected to be better adapted to local conditions (31). While

prey are typically thought to experience stronger selection than their predators (the "life-dinner

principle") (32), this asymmetry may be reversed when prey contain deadly toxins like TTX (3).

In fact, populations in central Oregon are the most toxic newts known (11), so non-resistant

predators should experience severe fitness consequences. Moreover, we found that garter snakes

tend to have greater functional estimates of resistance than the levels of TTX in co-occurring

newts (Fig. 2), implying intense selection on predators.

Non-adaptive processes of drift and gene flow provide a parsimonious explanation for the

overall pattern of variation in newt TTX across the geographic mosaic, but the extreme levels of

prey toxin and predator resistance at phenotypic hotspots such as central Oregon are likely a

result of arms race coevolution. After controlling for the effects of population structure,

escalation of TTX in newts was still marginally predicted by the TTX resistance of local garter

snakes in MRM analyses (p=0.065; Table 1) and significantly predicted by TTX resistance in a

linear model (t-value=2.803, p=0.006; Supplemental Table S6, model 3). This relationship seems

to be primarily driven by the southern portion of the transect, where intense reciprocal selection

has presumably led to the escalation of armaments in both species. Taken together, the overall

pattern suggests that co-adaptation can explain escalated TTX levels in specific hotspots of

coevolution, but global variation in TTX across the geographic mosaic is dictated by processes

like drift and gene flow. For example, TTX may be favored in specific hotspots of coevolution,

but not in surrounding regions like northern Washington (i.e., "coldspots"), and the tight spatial

correlation we observed between newt TTX and neutral SNPs reflects gene flow homogenizing

variation between hot- and coldspots.

The underlying importance of prey population structure in the geographic mosaic of

coevolution points to an influential role of "trait remixing", a largely untested component of the

geographic mosaic theory thought to generate spatial variation in the phenotypic interface of

coevolution (2, 10). The neutral processes of drift and gene flow are predicted to continually

alter the spatial distribution of allelic and phenotypic variation, potentially interfering with local

selection. Gene flow outwards from hotspots of coevolution is predicted to alter dynamics in

surrounding populations (13, 33), and if gene flow is high, the population with the strongest

reciprocal effects on fitness is expected to dictate broader landscape patterns of trait variation

(31, 33, 34). The homogenizing effects of gene flow may be less influential in snake populations

due to the simple genetic basis of TTX resistance and/or strong selection on predators.

Our results highlight how not all landscape patterns of phenotypic matching in natural

enemies may be the inherent result of coevolution (10). External factors such as abiotic

conditions (35), evolutionary constraints (36), or interactions with other species (37) are likely to

have unique effects on the evolution of prey and predator. In the geographic mosaic of arms race

coevolution, it appears that phenotypic divergence in newt toxin in disproportionally the result of

population structure, a pattern that underlies landscape-wide variation in the armaments of both

species. Ultimately, the evolutionary response to selection at the phenotypic interface is almost

certain to differ in two interacting species—so much so that coevolution may not always be the

most parsimonious explanation for observed patterns of phenotypic divergence and trait

matching across the geographic mosaic.

ACKNOWLEDGEMENTS

We thank the Departments of Fish and Wildlife in Washington and Oregon for scientific

collecting permits to MTJH (18-082 and 063-18, respectively). We also thank IACUC for the

protocol to EDB, Jr. at USU (1008). We are grateful for T. St. Pierre's help with fieldwork. J.

McGlothlin and K. Gendreau helped with sex-linked analyses of $Na_V1.4$ and provided valuable

feedback that improved the quality of this manuscript. R. Cox, D. Taylor, A. Bergland, D. Carr,

and the Brodie and Feldman lab groups also provided helpful input. This work was supported by

a Doctoral Dissertation Improvement Grant from the National Science Foundation to MTJH and

EDB III (DEB 1601296) and NSF support to EDB III (DEB 0922251).

AUTHOR CONTRIBUTIONS

MTJH designed the project, collected specimens, generated genetic data, and performed

statistical analyses. ANS collected phenotypic data on newt TTX levels. CRF collected

specimens and phenotypic data on snake resistance. EDB, Jr. collected phenotypic data on snake

resistance and provided leadership on the project. EDB III designed the project and provided

leadership. All authors prepared the manuscript.

DECLARATION OF INTERESTS

The authors declare no conflict of interest with this manuscript.

METHODS

We sampled phenotypic and genomic data from *Ta. granulosa* (*n*=138) and *Th. sirtalis*

(*n*=169) at nine locations along a latitudinal transect in the states of Washington and Oregon

(Fig. 1; Supplemental Table S1) and compared mosaic patterns of escalation in the arms race to

neutral population genomic structure across the landscape.

TTX levels of *Taricha granulosa*

We estimated levels of TTX in *Ta. granulosa* using a Competitive Inhibition Enzymatic

Immunoassay (CIEIA) and TTX-specific antibodies (38, 39). We quantified the amount of TTX

in a 5mm circular skin punch from the dorsum of each newt using a human skin-biopsy punch

(Acu-Punche, Acuderm Inc.) (26, 40, 41). These data were used to estimate the dorsal skin

concentration of TTX ($\mu g/cm^2$) in each individual. TTX is uniformly distributed throughout the

dorsum and levels of TTX in the dorsal skin are tightly correlated with toxicity in other regions

(41). We conducted a two-way ANOVA to test whether TTX differed by population and by sex,

because past work suggests toxin levels may vary by sex (40). Distribution and leverage analyses

indicated that a log(x + 0.1) transformation of TTX was needed. Transformed mean and variance

values were used in the cline-fitting analysis of TTX along the transect.

Although a considerable body of work has described the genetic basis of TTX resistance

in *Th. sirtalis* (see below), similar information regarding TTX in *Ta. granulosa* is unavailable.

Although we were unable to characterize genetic variation underlying prey toxin, we assume that

TTX production probably has a polygenic basis. For example, biosynthesis of saxitoxin (STX), a

similar neurotoxin found in marine species, involves gene expression in a cluster of up to 26

genes in cyanobacteria (42).

TTX resistance of *Thamnophis sirtalis*

We measured phenotypic TTX resistance using a well-established bioassay of whole animal performance (14, 22, 43, 44). Briefly, each individual was assayed on a 4 m racetrack to characterize its "baseline" crawl speed, then injected intraperitoneally with a known dose of TTX and assayed for "post-injection" speed. Population estimates of phenotypic TTX resistance are reported on a scale of mass-adjusted mouse units (MAMUs) to control for differences in body size (14). Resistance was estimated as the relative performance after injection: the MAMU dose of TTX that reduces performance by 50% of baseline speed. We incorporated racetrack data from previously published estimates of resistance from the same sampling locations to generate precise population estimates of phenotypic resistance in this study (see Supplemental Table S1) (14, 45).

The 50% MAMU dose was estimated separately for each population from a dose-response curve using curvilinear regression and the general transform $y´=\ln(1/y-1)$ (14). Individuals from each population received a series of TTX doses, with an average of 2.5 different doses per individual. At $y=0.5$ (i.e., 50%), $y´=0$ and the 50% dose is estimable $\hat{x}=-\alpha/\beta$ (where α is the intercept and β the slope from the curvilinear regression). Because $\hat{x}$ takes the form of a ratio, the standard error for the estimated 50% dose is calculated using standard methods for the variance of a ratio (14, 46). Confidence intervals of 95% were calculated as ±1.96 SE. Regression was performed in R with the "lmer" function implemented in the lme4 package (47). The individual ID of each snake was included as a random effect to account for the fact that each snake received multiple injections. Distribution and leverage analysis indicated that a transformation of the *x* variable (MAMU of TTX) was needed, so we transformed the data

using $x'=\ln(x + 1)$ (14). Differences among populations in phenotypic TTX resistance were

deemed significant if 95% confidence intervals did not overlap by more than half of a one-sided

error bar (48).

We genotyped snakes for their amino acid sequence in the DIV p-loop of the $Na_V1.4$

channel. Methods for Sanger sequencing are described in Hague et al. (2017) (22). For each

individual, we sequenced a 666 bp fragment that includes the DIV p-loop region of $Na_V1.4$. The

$Na_V1.4$ protein is encoded by the *SCN4A* gene located on the Z sex chromosome of *Th. sirtalis*

(Gendreau et al., *in prep*). Colubrid snakes, including garter snakes, have heteromorphic sex

chromosomes (ZZ males, ZW females) that are non-recombining (49, 50), and females appear to

be hemizygous for the Z-linked *SCN4A* gene. We used sex-specific PCR primers to confirm the

sex of each individual (ZW.734.F, 5'-TAAGGTCCTGGGCATGTCCT-3'; ZW.2468.R, 5'-

ATGGCTTGGAATGAGGTGGG-3'; Gendreau et al., *in prep*). In DIV p-loop sequences from

males, heterozygous positions on chromatograms were identified by eye and confirmed in both

directions with sequencing. The haplotype phase of the DIV p-loop sequence for each male was

inferred computationally with the program PHASE (51).

We translated the aligned DIV p-loop coding sequences into the amino acids and tested

for departures from Hardy-Weinberg Equilibrium (HWE) using a joint test for HWE and

equality of allele frequencies (EAF) using the *HWTriExact* function in the HardyWeinberg

package in R (52–54). Standard tests for HWE rely on the assumption of EAF in males and

females. The joint exact test for HWE and EAF accounts for the hemizygous sex and tests for

joint departures of HWE and EAF (53–55). We also calculated pairwise F_{ST} divergence at the

DIV p-loop in the program Arlequin (56) and used a Mantel test to test for a relationship

between pairwise F_{ST} divergence at the DIV p-loop and phenotypic divergence in whole-animal

TTX resistance. Significance was tested with 1000 permutations.

Functional analysis of trait matching

Following Hanifin et al. (2008) (11), we estimated functional levels of newt TTX and

snake resistance to visualize whether prey and predator exhibit matched levels of escalation

along the transect. The model provides a rough estimate of functional interactions between newt

TTX and snake resistance based on an extensive body of work (4, 14, 41, 43, 44, 57–59).

Localities are considered "matched" if a sympatric interaction between prey and predator could

potentially result in variable fitness outcomes for both species, leading to reciprocal selection

between newt TTX and snake resistance (11). For each locality, we estimated whole-newt levels

of TTX (mg) and the dose of TTX (mg) required to reduce performance of co-occurring snakes

to 15%, 50%, and 85% of their baseline performance. The TTX dose required to reduce snake

performance by 50% is considered a perfect functional match between newt TTX and snake

resistance. The 15% and 85% doses delimit the range of functionally relevant doses for snakes.

At performance levels <15%, all snakes that ingest newts are fully immobilized or killed and

newts escape, whereas at performance levels >85% all snakes are unaffected and captured newts

die (11, 15, 16). Localities where the full range of TTX doses found in newts fall outside the 15-

85% region of phenotypic space are considered phenotypic "mismatches", such that variable

fitness outcomes and reciprocal selection are unlikely to occur.

Methods for estimating functionally comparable values of newt TTX and snake

resistance are described in Hanifin et al. (2008) (11). Briefly, we extrapolated our measures of

newt TTX in skin samples ($\mu g/cm^2$) to the whole animal (mg of TTX/newt) using standard

methods. For snakes, the 15, 50, and 85% MAMU doses were extracted from the dose response

curves of each population (as described above). We converted snake resistance from MAMUs

based on intraperitoneal (IP) injections to mg of TTX in oral doses using the following equation:

$$\text{IPdose(mg)} = (\theta * 0.00001429) * \text{snake mass(g)}$$

where θ is the MAMU dose and 0.0001429 is the conversion factor (1 MAMU= 0.01429μg TTX

per gram of snake). The effects of TTX are dependent on body size, so we estimated the oral

dose of TTX required to slow the average adult *Th. sirtalis* (mean adult mass = 52.84 g). We

then converted the IP dose (mg) to the oral dose required to achieve the same performance

reduction by multiplying the IP dose by 40. Estimates of orally ingested doses of TTX (mg)

required to reduce snake performance by 15, 50, and 85% were used as a functionally equivalent

metric to compare predator phenotypes to those of the prey (Supplemental Table S2).

ddRADseq library preparation

393   We prepared ddRADseq libraries separately for *Ta. granulosa* and *Th. sirtalis* using the

protocol described in Peterson et al. (2012) (60). Genomic DNA was extracted using the DNeasy

Blood & Tissue kit (Qiagen Inc., Valencia, CA.). We digested 600 ng of genomic DNA for each

sample using the restriction enzymes *EcoRI* and *SbfI* for *Ta. granulosa* and *MfeI* and *SbfI* for *Th.*

sirtalis. Unique combinations of individual P1 and P2 barcoded adapters were annealed to the

digested genomic DNA of each sample. Each barcode was six base pairs long and differed by at

least two nucleotides. After barcoding, *Ta. granulosa* and *Th. sirtalis* samples were pooled

separately, purified with AmpureXP beads (Beckman Coulter, Inc., Brea, CA, USA), and size

selected for 500 to 600 bp fragments using a Pippin Prep (Sage Science, Inc., Beverly, MA,

USA). We enriched the adapter-ligated fragments in the size-selected libraries using 16 PCR

cycles and then purified the product with AmpureXP beads. The *Ta. granulosa* and *Th. sirtalis*

libraries were each run on two lanes of the Illumina HiSeq 2500 platform (Illumina, Inc., San

Diego, CA, USA) to generate 125 bp paired-end reads.

Data processing of Illumina sequencing

Read quality of the raw sequence data was assessed using FastQC 0.11.5 (61). We used

process_radtags in Stacks 1.46 (62) on both the *Ta. granulosa* and *Th. sirtalis* datasets to

demultiplex reads and removed sequences with low-quality scores or uncalled bases. For the *Ta.*

granulosa dataset, we used the *denovo_map.pl* pipeline in Stacks, because a reference genome is

not currently available. We used a minimum depth of three (-*m*), a distance of three between

stacks (-*M*), and a distance of three between catalog loci (-*n*). The *Th. sirtalis* reads were aligned

to the *Th. sirtalis* genome using Bowtie2 2.2.9 (63). We discarded reads that did not align or had

more than one match to the genome. We used *ref_map.pl* in Stacks to assemble the reference-

aligned sequences into loci, with a minimum depth of three (-*m*). For both species, we used

populations in Stacks to select loci with a minimum depth of 10x coverage. To avoid linkage

among sites within the same locus, we only retained one single nucleotide polymorphism (SNP)

per locus.

We used the dartR package in R (64, 65) to remove loci and individuals with >30%

missing data. We also removed loci with a minor allele frequency (MAF) <5%, including those

that were invariant. Finally, we removed putative loci under selection. The program BayeScan

2.1 was used to search for loci with F_{ST} coefficients that were significantly different than those

expected under neutrality (66). The Bayesian analysis used 20 pilot runs with 5,000 iterations

followed by an additional burn-in of 50,000 and 50,000 output iterations. An outlier analysis

with FDR-corrected p-values (q-values) <0.05 was used to identify and remove outlier loci

putatively under selection.

Illumina HiSeq sequencing yielded an average of 1,093,529 raw reads per sample of *Ta.*

granulosa (n=137). After initial quality control in *process_radtags*, the *denovo_map* pipeline in

Stacks identified an average of 32,469 loci per individual, with a mean of 19x coverage. After

further filtering steps in *populations*, dartR, and Bayescan, we retained 3,634 unlinked neutral

SNPs in 123 individuals. For *Th. sirtalis*, sequencing yielded an average of 2,451,623 raw reads

per sample (n=143). The *ref_map* pipeline identified an average of 13,501 loci per individual,

with a mean of 80x coverage. After additional filtering, we retained 1,027 unlinked neutral SNPs

in 132 individuals. The final datasets comprised a larger number of SNPs for *Ta. granulosa*

(3,634) than *Th. sirtalis* (1,027), so we reran analyses with a random subsample of 1,027 SNPs

for *Ta. granulosa* to generate equivalent datasets for each species. Results from the subsample

were highly similar to the full dataset, so only results from the full analysis are presented herein.

Analysis of geographic population structure

The filtered SNPs for each species were used to calculate observed heterozygosity (H_O)

and gene diversity (H_S) for each population (67) using the hierfstat package in R (68). To

estimate neutral population genomic structure, we calculated global and pairwise F_{ST} values (69).

Confidence intervals were estimated by running 1000 bootstraps over loci using the hierfstat and

stAMPP packages in R (70). Estimates of population genetic diversity from the final SNP dataset

of each species are reported in Supplementary Table S3.

We tested for a pattern of isolation-by-distance (IBD) along each transect by performing

Mantel tests on matrices of linearized pairwise F_{ST} and geographic distance (71). We also

conducted distance-based redundancy analyses (dbRDA), which are thought to be more reliable

than Mantel tests at detecting spatial patterns like IBD (72, 73). We conducted dbRDA analyses

in the vegan package in R (74) to test for a relationship between pairwise F_{ST} values and the

geographic coordinates (latitude and longitude) of sampling locations. We assessed significance

of IBD tests with 1000 permutations.

We visualized population structure of each species using a principal coordinate analysis

(PCoA) in the dartR package in R. The first axis (PCo1) from the PCoA captured latitudinal

variation along the transect for both newts and snake (Fig. 3); therefore, we used the PCo1 values

for each individual as a neutral expectation in the cline-fitting analyses (see below). We assessed

population structure using a Bayesian assignment approach implemented in the program

STRUCTURE (75, 76). We estimated the optimal number of genetic clusters (K) ranging for one

to nine, without populations included as priors. The model assumed population admixture and

correlated allele frequencies (76). The analysis first ran for 100,000 iterations as burn-in and then

we collected data from the following 1,000,000 interactions in 10 different independent runs.

STRUCTURE HARVESTER (77) was used to detect the most probable K using the Evanno's

method (78). Ancestry proportion (Q) values of the 10 runs for each value of the most probable

K we averaged using CLUMPP (79) and visualized using the pophelper package in R (80). We

calculated the average Q value for each population, which represent the fraction of membership

to each genetic cluster (K). These Q estimates were used as a neutral expectation in cline-fitting

analyses (see below) and compared to the cline results of PCo1 from the PCoA.

Multiple regression of distance matrices (MRMs)

MRMs are an extension of Mantel tests that involve multiple regression of a response

distance matrix on explanatory matrices (81–84). Partial regression coefficients can be used to

understand the relationship between two matrices while controlling for the effects of a third

matrix (7, 85). For each species, we generated two distance matrices: (1) pairwise phenotypic

divergence in the coevolutionary trait (e.g., log-transformed TTX of *Ta. granulosa*) and (2)

pairwise genomic divergence (F_{ST}) from neutral SNPs. We then tested whether population

patterns of phenotypic escalation in a focal species (e.g., TTX of *Ta. granulosa*) are best

explained by neutral genomic divergence (pairwise F_{ST}) or escalation in the natural enemy (TTX

resistance of *Th. sirtalis*). MRM analyses can be confounded when explanatory variables are

spatially autocorrelated, so we compared results when the two explanatory variables were

analyzed separately and together (85–87).

Cline analyses

We performed ML cline-fitting analyses on the phenotypic and genetic data from each

species (88, 89). For *Ta. granulosa*, we fit a cline to TTX levels using the mean and variance of

the log-transformed TTX data ($\mu g/cm^2$). For *Th. sirtalis*, we fit (1) a cline to phenotypic TTX

resistance using the 50% MAMU dose and variance from the ln-transformed MAMU data and

(2) a genetic cline to the frequency of TTX-resistant alleles in each population. Two different

TTX-resistant DIV alleles occur in the Pacific Northwest: $Na_V1.4^V$ is generally found at high

frequency in the center of the transect, whereas $Na_V1.4^{VA}$ occurs predominately in southern

populations (Fig. 1). We generated separate clines for each allele (data not shown), but the cline

fit to the combined frequency of both TTX-resistant alleles (Fig. 4) was the most representative

of variation along the entire transect.

For each species, we compared clinal variation in coevolutionary traits to variation at the

neutral SNPs. We first fit clines to the PCo1 values from the PCoA of each species, and

secondly, we fit clines to the ancestry proportions (Q) from the most likely value of K for each

species (K=2). Both analyses of population structure (PCo1 from the PCoA and K=2 from

STRUCTURE) produced concordant results (Supplemental Fig. S2; Supplemental Table S5), so

only the analysis of PCo1 is presented in the main text. PCoAs have no explicit population

genetic assumptions, whereas STRUCTURE assumes that all loci are unlinked at linkage

equilibrium and in HWE. Because the PCoA and STRUCTURE analyses produced similar

results despite different underlying assumptions, we feel confident that the cline-fitting analyses

provided a representative depiction of neutral population structure in each species. Finally, we

reran cline-fitting analyses with the Elk River population removed from the *Ta. granulosa*

dataset, because the PCoA (Fig. 3) and STRUCTURE plot of K=3 (Supplemental Fig. S1) both

suggest the population is genetically distinct from all others; however, this produced the same

qualitative result as when the population was included.

We fit clines using the HZAR package in R (90). We calculated distances along the cline

as kilometers (km) from the northernmost sampling site (Clallam). We ran 15 separate models

that varied in the number of cline shape parameters estimated. All models estimated the cline

center (distance from sampling location 1, c) and width (1/maximum slope, w), but could

additionally estimate combinations of exponential decay curve (tail) parameters (neither tail,

right tail only, left tail only, mirrored tails, or both tails separately), which represent the distance

from the cline center to the tail (δ) and the slope of the tail (τ). The genetic models varied as to

whether they estimated allele frequencies at the cline ends (p_{min} and p_{max}) or fixed them at 0 and

1. All models were then compared using AIC corrected for small sample sizes (AICc) and

maximum likelihood parameters were extracted for the best-fitting model. We considered cline

centers with non-overlapping two log-likelihood unit support limits (confidence intervals; CIs) to

occur in significantly different geographic locations (91, 92).

The cline-fitting analyses revealed a tight correlation between the TTX levels of *Ta.*

granulosa and PCo1 from the PCoA of neutral SNPs (Fig. 4). To further assess this relationship,

we fit linear models that tested whether clinal variation in TTX is best explained by neutral

population structure from PCo1 and 2 of the PCoA (model 1) or escalation of TTX resistance in

Th. sirtalis (model 2). We then fit a combined model including all explanatory variables (model

3) to test whether neutral structure from the PCoA predicts TTX levels even after controlling for

the TTX resistance of *Th. sirtalis* (Supplemental Table S6).

We conducted an analogous set of tests (models 1-3) for *Th. sirtalis* to assess whether

TTX resistance is best predicted by neutral structure from the PCoA or the TTX levels of *Ta.*

granulosa; however, the analyses were more complicated because population estimates of TTX

resistance (50% MAMU doses) are extracted from a dose response curve. TTX resistance of

individual snakes is measured as a proportion (i.e., the reduction in crawl speed after a given

dose of TTX), with each individual receiving multiple injections. We accounted for these data by

fitting linear mixed models using the "lmer" function in the R package *lme4* (47). We treated

"post-injection" crawl speed (after a given dose of TTX) as the response variable, and included

"baseline" crawl speed and the dose of TTX as covariates in each model. We also included

individual ID as a random effect to account for the fact that individuals received multiple doses.

Post-injection crawl speed was log-transformed to account for non-normality. Statistical

significance of fixed effects was determined by an ANOVA using a Wald Chi-Square test with

type III sum of squares and one degree of freedom, implemented in the *car* package in R (93).

We reported the marginal R^2 for each model, which describes the proportion of variance

explained by the fixed factors (not including random effects), implemented in the *MuMIn*

package (94, 95).

Data availability

DNA sequence alignments for the DIV p-loop of Na$_V$1.4 and the ddRADseq data will be

made available on GenBank upon manuscript acceptance. All phenotypic data and the code for

statistical analyses will be submitted to Dryad upon acceptance.

REFERENCES

1. Janzen DH (1980) When is it coevolution? *Evolution*.

2. Thompson JN (2005) *The Geographic Mosaic of Coevolution* (University of Chicago Press, Chicago).

3. Brodie III ED, Brodie, Jr. ED (1999) Predator-prey arms races. *BioScience* 49(7):557–568.

4. Brodie III ED, Ridenhour BJ (2003) Reciprocal selection at the phenotypic interface of coevolution. *Integr Comp Biol* 43(3):408–418.

5. Benkman CW, Parchman TL, Favis A, Siepielski AM (2003) Reciprocal selection causes a coevolutionary arms race between crossbills and lodgepole pine. *Am Nat* 162(2):182–194.

6. Zangerl AR, Berenbaum MR (2003) Phenotype matching in wild parsnip and parsnip webworms: causes and consequences. *Evolution* 57(4):806–815.

7. Toju H, Ueno S, Taniguchi F, Sota T (2011) Metapopulation structure of a seed-predator weevil and its host plant in arms race coevolution. *Evolution* 65(6):1707–1722.

8. Lively CM, Dybdahl MF (2000) Parasite adaptation to locally common host genotypes. *Nature* 405(6787):679.

9. Thrall PH, Burdon J, Bever JD (2002) Local adaptation in the *Linum marginale—Melampsora lini* host-pathogen interaction. *Evolution* 56(7):1340–1351.

10. Gomulkiewicz R, et al. (2007) Dos and don'ts of testing the geographic mosaic theory of coevolution. *Heredity* 98(5):249–258.

11. Hanifin CT, Brodie, Jr. ED, Brodie III ED (2008) Phenotypic mismatches reveal escape from arms-race coevolution. *PLoS Biol* 6(3):e60.

12. Swenson NG, Howard DJ, Hellberg AEME, Losos EJB (2005) Clustering of contact zones, hybrid zones, and phylogeographic breaks in North America. *Am Nat* 166(5):581–591.

13. Thompson JN, Nuismer SL, Gomulkiewicz R (2002) Coevolution and maladaptation. *Integr Comp Biol* 42(2):381–387.

14. Brodie, Jr. ED, Ridenhour BJ, Brodie III ED (2002) The evolutionary response of predators to dangerous prey: hotspots and coldspots in the geographic mosaic of coevolution between garter snakes and newts. *Evolution* 56(10):2067–2082.

15. Williams BL, Hanifin CT, Brodie, Jr. ED, Brodie III ED (2010) Tetrodotoxin affects survival probability of rough-skinned newts (*Taricha granulosa*) faced with TTX-resistant garter snake predators (*Thamnophis sirtalis*). *Chemoecology* 20(4):285–290.

16. Williams BL, Brodie, Jr. ED, Brodie III ED (2003) Coevolution of deadly toxins and
predator resistance: self-assessment of resistance by garter snakes leads to behavioral
rejection of toxic newt prey. *Herpetologica* 59(2):155–163.

17. Fozzard HA, Lipkind GM (2010) The tetrodotoxin binding site is within the outer vestibule
of the sodium channel. *Mar Drugs* 8(2):219–234.

18. Tikhonov DB, Zhorov BS (2012) Architecture and pore block of Eukaryotic voltage-gated
sodium channels in view of NavAb bacterial sodium channel structure. *Mol Pharmacol*
82(1):97–104.

19. Geffeney S, Brodie, Jr. ED, Ruben PC, Brodie III ED (2002) Mechanisms of adaptation in a
predator-prey arms race: TTX-resistant sodium channels. *Science* 297(5585):1336–1339.

20. Geffeney SL, Fujimoto E, Brodie III ED, Brodie, Jr. ED, Ruben PC (2005) Evolutionary
diversification of TTX-resistant sodium channels in a predator-prey interaction. *Nature*
434(7034):759–763.

21. Feldman CR, Brodie, Jr. ED, Brodie III ED, Pfrender ME (2010) Genetic architecture of a
feeding adaptation: garter snake (*Thamnophis*) resistance to tetrodotoxin bearing prey. *Proc*
R Soc B Biol Sci 277(1698):3317–3325.

22. Hague MTJ, Feldman CR, Brodie, Jr. ED, Brodie III ED (2017) Convergent adaptation to
dangerous prey proceeds through the same first-step mutation in the garter snake
Thamnophis sirtalis. *Evolution* 71(6):1504–1518.

23. Langerhans RB, DeWitt TJ (2004) Shared and unique features of evolutionary
diversification. *Am Nat* 164(3):335–349.

24. Keller SR, Sowell DR, Neiman M, Wolfe LM, Taylor DR (2009) Adaptation and
colonization history affect the evolution of clines in two introduced species. *New Phytol*
183(3):678–690.

25. Ridenhour BJ, Brodie, Jr. ED, Brodie III ED (2007) Patterns of genetic differentiation in
Thamnophis and *Taricha* from the Pacific Northwest. *J Biogeogr* 34(4):724–735.

26. Hague MTJ, et al. (2016) Toxicity and population structure of the Rough-Skinned Newt
(*Taricha granulosa*) outside the range of an arms race with resistant predators. *Ecol Evol*
6(9):2714–2724.

27. Janzen FJ, Krenz JG, Haselkorn TS, Brodie, Jr. ED, Brodie III ED (2002) Molecular
phylogeography of common garter snakes (*Thamnophis sirtalis*) in western North America:
implications for regional historical forces. *Mol Ecol* 11(9):1739–1751.

28. Moczydlowski EG (2013) The molecular mystique of tetrodotoxin. *Toxicon* 63:165–183.

29. Yotsu M, Iorizzi M, Yasumoto T (1990) Distribution of tetrodotoxin, 6-epitetrodotoxin, and
11-deoxytetrodotoxin in newts. *Toxicon* 28(2):238–241.

30. Yasumoto T, Yotsu-Yamashita M (1996) Chemical and etiological studies on tetrodotoxin
and its analogs. *Toxin Rev* 15(2):81–90.

31. Gandon S (2002) Local adaptation and the geometry of host–parasite coevolution. *Ecol Lett*
5(2):246–256.

32. Dawkins R, Krebs JR (1979) Arms races between and within species. *Proc R Soc Lond B*
Biol Sci 205(1161):489–511.

33. Gomulkiewicz R, Thompson JN, Holt RD, Nuismer Scott L, Hochberg ME (2000) Hot spots,
cold spots, and the geographic mosaic theory of coevolution. *Am Nat* 156(2):156–174.

34. Gandon S, Michalakis Y (2002) Local adaptation, evolutionary potential and host–parasite
coevolution: interactions between migration, mutation, population size and generation time.
J Evol Biol 15(3):451–462.

35. Johnson SG, Hulsey CD, León FJG (2007) Spatial mosaic evolution of snail defensive traits.
BMC Evol Biol 7(1):1–11.

36. Hague MTJ, et al. (2018) Large-effect mutations generate trade-off between predatory and
locomotor ability during arms race coevolution with deadly prey. *Evol Lett* 2(4):406–416.

37. Benkman CW, Holimon WC, Smith JW (2001) The influence of a competitor on the
geographic mosaic of coevolution between crossbills and lodgepole pine. *Evolution*
55(2):282–294.

38. Gall BG, et al. (2011) Tetrodotoxin levels in larval and metamorphosed newts (*Taricha*
granulosa) and palatability to predatory dragonflies. *Toxicon* 57(7–8):978–983.

39. Stokes AN, Williams BL, French SS (2012) An improved competitive inhibition enzymatic
immunoassay method for tetrodotoxin quantification. *Biol Proced Online* 14(1):3.

40. Hanifin CT, Brodie III ED, Brodie, Jr. ED (2002) Tetrodotoxin levels of the rough-skin
newt, *Taricha granulosa*, increase in long-term captivity. *Toxicon* 40(8):1149–1153.

41. Hanifin CT, Brodie III ED, Brodie, Jr. ED (2004) A predictive model to estimate total skin
tetrodotoxin in the newt *Taricha granulosa*. *Toxicon* 43(3):243–249.

42. Murray SA, Mihali TK, Neilan BA (2010) Extraordinary conservation, gene loss, and
positive selection in the evolution of an ancient neurotoxin. *Mol Biol Evol* 28(3):1173–
1182.

43. Brodie III ED, Brodie, Jr. ED (1990) Tetrodotoxin resistance in garter snakes: an
evolutionary response of predators to dangerous prey. *Evolution* 44(3):651–659.

44. Ridenhour BJ, Brodie III ED, Brodie, Jr. ED (2004) Resistance of neonates and field-
collected garter snakes (*Thamnophis* spp.) to tetrodotoxin. *J Chem Ecol* 30(1):143–154.

45. Ridenhour B (2004) The coevolutionary process: The effects of population structure on a
predator-prey system. Dissertation (Indiana University, Bloomington, IN).

46. Lynch M, Walsh B, others (1998) *Genetics and analysis of quantitative traits* (Sinauer
Sunderland, MA).

47. Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using
lme4. *J Stat Softw* 67(1):1–48.

48. Cumming G, Finch S (2005) Inference by eye: Confidence intervals and how to read pictures
of data. *Am Psychol* 60(2):170–180.

49. Vicoso B, Emerson J, Zektser Y, Mahajan S, Bachtrog D (2013) Comparative sex
chromosome genomics in snakes: differentiation, evolutionary strata, and lack of global
dosage compensation. *PLoS Biol* 11(8):e1001643.

50. Augstenová B, et al. (2018) ZW, XY, and yet ZW: Sex chromosome evolution in snakes
even more complicated. *Evolution* 72(8):1701–1707.

51. Stephens M, Smith NJ, Donnelly P (2001) A new statistical method for haplotype
reconstruction from population data. *Am J Hum Genet* 68(4):978–989.

52. Graffelman J, Morales-Camarena J (2008) Graphical tests for Hardy-Weinberg Equilibrium
based on the ternary plot. *Hum Hered* 65(2):77–84.

53. Graffelman J, Weir BS (2018) Multi-allelic exact tests for Hardy–Weinberg equilibrium that
account for gender. *Mol Ecol Resour* 18(3):461–473.

54. Graffelman J, Weir BS (2018) On the testing of Hardy-Weinberg proportions and equality of
allele frequencies in males and females at biallelic genetic markers. *Genet Epidemiol*
42(1):34–48.

55. Graffelman J, Weir B (2016) Testing for Hardy–Weinberg equilibrium at biallelic genetic
markers on the X chromosome. *Heredity* 116(6):558.

56. Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: a new series of programs to perform
population genetics analyses under Linux and Windows. *Mol Ecol Resour* 10(3):564–567.

57. Hanifin CT, Yotsu-Yamashita M, Yasumoto T, Brodie ED (1999) Toxicity of dangerous
prey: variation of tetrodotoxin levels within and among populations of the newt *Taricha
granulosa*. *J Chem Ecol* 25(9):2161–2175.

58. Williams BL, Brodie ED, Brodie ED (2002) Comparisons between toxic effects of
tetrodotoxin administered orally and by intraperitoneal injection to the garter snake
Thamnophis sirtalis. *J Herpetol* 36(1):112–115.

59. Brodie III ED, Brodie, Jr. ED (1991) Evolutionary response of predators to dangerous prey:
reduction of toxicity of newts and resistance of garter snakes in island populations.
Evolution 45(1):221–224.

60. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double digest RADseq:
An inexpensive method for de novo SNP discovery and genotyping in model and non-
model species. *PLoS ONE* 7(5):e37135–11.

61. Andrew S (2016) *FastQC* Available at:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

62. Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis
tool set for population genomics. *Mol Ecol* 22(11):3124–3140.

63. Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9(4):357–359.

64. Gruber B, Unmack PJ, Berry OF, Georges A (2017) dartr: An r package to facilitate analysis
of SNP data generated from reduced representation genome sequencing. *Mol Ecol Resour*.

65. R Core Team (2018) *R: A Language and Environment for Statistical Computing* (R
Foundation for Statistical Computing, Vienna, Austria) Available at: https://www.R-
project.org/.

66. Foll M, Gaggiotti O (2008) A genome-scan method to identify selected loci appropriate for
both dominant and codominant markers: a Bayesian perspective. *Genetics* 180(2):977–993.

67. Nei M (1987) *Molecular evolutionary genetics* (Columbia University Press).

68. Goudet J, Jombart T (2015) *hierfstat: Estimation and Tests of Hierarchical F-Statistics*
Available at: https://CRAN.R-project.org/package=hierfstat.

69. Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population
structure. *evolution* 38(6):1358–1370.

70. Pembleton LW, Cogan NOI, Forster JW (2013) StAMPP: an R package for calculation of
genetic differentiation and structure of mixed-ploidy level populations. *Mol Ecol Resour*
13:946–952.

71. Rousset F (1997) Genetic differentiation and estimation of gene flow from F-statistics under
isolation by distance. *Genetics* 145(4):1219–1228.

72. Legendre P, Fortin M-J (2010) Comparison of the Mantel test and alternative approaches for
detecting complex multivariate relationships in the spatial analysis of genetic data. *Mol
Ecol Resour* 10(5):831–844.

73. Meirmans PG (2015) Seven common mistakes in population genetics and how to avoid
them. *Mol Ecol* 24(13):3223–3231.

74. Oksanen J, et al. (2018) *vegan: Community Ecology Package* Available at: https://CRAN.R-
project.org/package=vegan.

75. Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using
multilocus genotype data. *Genetics* 155(2):945–959.

76. Falush D, Stephens M, Pritchard JK (2003) Inference of population structure using
multilocus genotype data: linked loci and correlated allele frequencies. *Genetics*
164(4):1567–1587.

77. Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet*
Resour 4(2):359–361.

78. Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using
the software structure: a simulation study. *Mol Ecol* 14(8):2611–2620.

79. Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program
for dealing with label switching and multimodality in analysis of population structure.
Bioinformatics 23(14):1801–1806.

80. Francis RM (2017) pophelper: An R package and web app to analyse and visualise
population structure. *Mol Ecol Resour* 17(1):27–32.

81. Manly BFJ (1986) Randomization and regression methods for testing for associations with
geographical, environmental and biological distances between populations. *Res Popul Ecol*
28(2):201–218.

82. Smouse PE, Long JC, Sokal RR (1986) Multiple regression and correlation extensions of the
mantel test of matrix correspondence. *Syst Zool* 35(4):627.

83. Legendre P, Lapointe F-J, Casgrain P (1994) Modeling brain evolution from behavior: a
permutational regression approach. *Evolution* 48(5):1487–1499.

84. Lichstein JW (2007) Multiple regression on distance matrices: a multivariate spatial analysis
tool. *Plant Ecol* 188(2):117–131.

85. Rosenblum EB (2005) Convergent evolution and divergent selection: lizards at the White
Sands ecotone. *Am Nat*.

86. Raufaste N, Rousset F (2001) Are partial mantel tests adequate? *Evolution* 55(8):1703–1705.

87. Rousset F, Waller D (2002) Partial Mantel tests: reply to Castellano and Balletto. *Evolution*
56(9):1874–1875.

88. Szymura JM, Barton NH (1986) Genetic analysis of a hybrid zone between the fire-bellied
toads, *Bombina bombina* and *B. variegata*, near Cracow in southern Poland. *Evolution*
40(6):1141–1159.

89. Szymura JM, Barton NH (1991) The genetic structure of the hybrid zone between the fire-
bellied toads *Bombina bombina* and *B. variegata*: comparisons between transects and
between loci. *Evolution* 45(2):237–261.

90. Derryberry EP, Derryberry GE, Maley JM, Brumfield RT (2014) HZAR: hybrid zone
analysis using an R software package. *Mol Ecol Resour* 14(3):652–663.

91. Baldassarre DT, White TA, Karubian J, Webster MS (2014) Genomic and morphological
analysis of a semipermeable avian hybrid zone suggests asymmetrical introgression of a
sexual signal. *Evolution* 68(9):2644–2657.

92. Scordato ES, et al. (2017) Genomic variation across two barn swallow hybrid zones reveals
traits associated with divergence in sympatry and allopatry. *Mol Ecol* 26(20):5676–5691.

93. Fox J, Weisberg S (2011) *An R companion to applied regression* (Sage, Thousand Oaks CA).
Second Available at: http://socserv.socsci.mcmaster.ca/jfox/Books/Companion.

94. Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R^2 from
generalized linear mixed-effects models. *Methods Ecol Evol* 4(2):133–142.

95. Bartoń K (2018) *MuMIn: Multi-Model Inference* Available at: https://CRAN.R-
project.org/package=MuMIn.

 FIGURE LEGENDS

Fig. 1. Matching phenotypes in prey and predator imply local co-adaptation in the arms race. (**A**) Population means of TTX levels ($\mu g/cm^2$) in newts and (**B**) phenotypic TTX resistance (50% MAMU dose) in snakes along the latitudinal transect. Error bars indicate 95% confidence intervals. The x-axis represents linear distance (km) from the northernmost sampling site (Clallam; 0 km). For snakes, the frequency of TTX-resistant alleles in the $Na_V1.4$ channel is shown with pie charts proportional to sample size. To the right, the schematic of $Na_V1.4$ shows the four domains of the channel (DI–DIV), with the extracellular pore loops (p-loops) highlighted with bold lines. Specific amino acid changes in the DIV p-loop are shown in their relative positions within the pore. The TTX-sensitive ancestral sequence (purple) is listed, followed by the two derived alleles known to confer increases in channel resistance in this lineage. (**C**) Map inset illustrates population estimates of prey toxin and predator resistance at each location in the geographic mosaic. Blue colors correspond to low estimates of TTX (squares) or resistance (circles), whereas red indicates escalated phenotypes in the arms race.

Fig. 2. Functional overlap in levels of prey toxin and predator resistance. The functional relationship of average adult newt total skin TTX (mg, log scale) is plotted against the oral dose of TTX (mg, log scale) required to reduce the speed of an average adult garter snake to 50% of baseline speed post-ingestion of TTX for each locality. Individual points are colorized by the average level of newt TTX as in Fig. 1C. Vertical bars represent the full range of newt TTX values observed for each population; horizontal bars represent the 95% confidence interval for the oral 50% dose of the corresponding snake population. The dashed 50% line represents a functional match between newt TTX and snake resistance: the dose of TTX in a newt that would reduce a snake of a given resistance to 50% of its performance. The solid 15% and 85% lines (calculated as best-fit regressions for each locality) delimit the range of functionally relevant TTX doses for snakes across the range of sampled localities. Below the 85% line, values of TTX resistance are high enough for snakes to consume co-occurring newts with no reduction in performance or fitness. Above the 15% line, doses of newt TTX are so high that any snake that ingests a newt would be completely incapacitated or killed. Localities with error bars that fall *entirely* outside the boundaries of these lines (none of which were observed) are considered mismatched and regions where reciprocal selection could not occur.

Fig. 3. Populations of prey and predator differ in geographic structure. Results from the principal coordinate (PCoA) and STRUCTURE analyses of neutral SNPs from newts and snakes. PCoA graphs are rotated 90º to emphasize the major axis of variation corresponding to latitude. The PCo1 values for each individual were used as a neutral expectation in the cline-fitting analyses. STRUCTURE plots are arranged latitudinally by population, in the same order as the map. Each horizontal bar represents the ancestry assignment of an individual, with populations separated by white dashed lines.

Fig. 4. Levels of prey toxin are best predicted by neutral population structure, whereas predator resistance is predicted by prey toxin. Cline-fitting results for phenotypic and genetic variation are shown, with error bars indicating confidence intervals surrounding the geographic cline centers. (**A**) Phenotypic clines of TTX levels ($\log[\text{TTX } \mu g/cm^2 + 0.1]$) and (**B**) TTX resistance ($\ln[\text{MAMU} + 1]$) are shown in red. For snakes, the frequency of TTX-resistant alleles

811   in the $Na_V1.4$ channel was also modeled (in black). Gray dashed lines represent the neutral

812   expectation for trait variation due to population structure, based on the PCoA. The cline center

813   points of TTX levels and neutral PCo1 in newts, and phenotypic resistance and TTX-resistant

814   alleles in snakes, are all located within in 84 km of each other along the 611 km transect.

TABLES

Table 1. Results from multiple regression of distance matrices (MRMs) comparing population divergence in phenotypic and genetic data.

Response Variable	Explanatory Variable(s)	Coefficient	p-value	R^2
	Neutral F_{ST} of newts	5.998	**0.002***	0.414
	TTX resistance of snakes	0.415	**0.019***	0.274
Newt TTX levels	Neutral F_{ST} + TTX resistance			
	Neutral F_{ST} of newts	4.827	**0.006***	0.501
	TTX resistance of snakes	0.253	0.065˙	
	Neutral F_{ST} of snakes	1.442	0.719	0.006
	TTX level of newts	0.662	**0.021***	0.274
Snake TTX resistance	Neutral F_{ST} + TTX toxicity			
	Neutral F_{ST} of snakes	-5.830	0.189	0.338
	TTX level of newts	0.873	**0.011***	
	Neutral F_{ST} of newts	4.632	**0.035***	0.155
Snake F_{ST} of DIV p-loop	Neutral F_{ST} of snakes	2.649	0.202	0.080
	Neutral F_{ST} of newts	3.196	**0.010***	0.311

 SUPPLEMENTAL INFORMATION

Supplemental Fig. S1. STRUCTURE results for K=2-4. The most likely number of genetic
cluster was K=2 for both species (Fig. 2). STRUCTURE plots are arranged latitudinally by
population, in the same order as the map. Each horizontal bar represents the ancestry assignment
of an individual, with populations separated by white dashed lines.

Supplemental Fig. S2. Comparison of neutral clines fit to (1) PCo1 values from the PCoA and
(2) average ancestry assignment values (K=2) from the STRUCTURE analysis. Error bars
indicate confidence intervals surrounding the geographic cline centers.

Supplemental Table S1. Datasets for each sampling location along the latitudinal transect. The total number of animals sampled in this study is shown first, followed by the number of individuals (n) included in each analysis. For estimates of phenotypic resistance in *Th. sirtalis*, we combined individuals from this study with previously published racetrack data from the same sampling locations (14, 45). In the DIV p-loop, the number (n) of alleles sampled accounts for females, the hemizygous sex with only one genetic copy of $Na_V1.4$. Results are shown for the joint test of Hardy-Weinberg Equilibrium (HWE) and Equality of Allele Frequencies (EAF).

Location	County	Total animals sampled in this study		*Ta. granulosa*				*Th. sirtalis*				
		Ta. granulosa	*Th. sirtalis*	TTX			ddRADseq	Phenotypic TTX resistance			DIV p-loop		ddRADseq
				n	mean (ug/cm²)	SE	n	n	50% MAMU	SE	n alleles	HWE & EAF test p-value	n
Clallam	Clallam, WA	12	11	12	0.731	0.307	12	53	4.163	0.076	14	NA	4
Cook Creek	Grays Harbor, WA	14	17	14	0.236	0.112	13	14	5.915	0.119	26	1.000	16
Potter's Slough	Pacific, WA	15	20	15	2.75	0.686	13	42	19.203	0.090	27	1.000	16
Warrenton	Clatsop, OR	15	24	14	1.63	0.241	14	323	17.357	0.038	36	0.257	20
Hebo	Tillamook, OR	15	15	15	4.65	0.822	12	2	43.413	0.085	21	0.708	14
Benton	Benton., OR	18	20	18	16.9	3.1	17	386	31.380	0.047	23	0.347	14
Ten Mile	Lane, OR	15	16	14	7.43	1.85	14	93	52.817	0.079	26	0.451	16
Lake Tahkenitch	Douglas, OR	14	29	13	10.7	2.56	12	17	14.552	0.227	42	0.182	15
Elk River	Curry, OR	20	17	15	6.52	0.612	16	43	12.935	0.063	20	0.324	17
Total		138	169	130			123	932			235		132

837 **Supplemental Table S2.** For each locality, functional estimates of adult *Ta. granulosa* total skin TTX (mg) and oral doses of TTX
838 (mg) required to reduce the speed of an average adult *Th. sirtalis* to 15, 50, and 85% of baseline speed post-ingestion of TTX.
839

Population	*Ta. granulosa*			*Th. sirtalis*		
	mean TTX (mg)	max TTX (mg)	min TTX (mg)	50% oral dose (mg)	15% oral dose (mg)	85% oral dose (mg)
Clallam	0.0176	0.0715	0.0001	0.1257	0.5710	0.0102
Cook Creek	0.0068	0.0422	0.0001	0.1787	0.4614	0.0585
Potter's Slough	0.0816	0.3813	0.0220	0.5800	1.9169	0.1610
Warrenton	0.0475	0.1263	0.0053	0.5243	2.2042	0.1074
Hebo	0.1933	0.5495	0.0698	1.3113	1.7279	0.9935
Benton	0.7107	1.9188	0.1097	0.9479	3.9647	0.2093
Ten Mile	0.3188	1.0166	0.0455	1.5954	11.1998	0.2051
Lake Tahkenitch	0.2590	0.8172	0.0468	0.4396	3.4492	0.0332
Elk River	0.2043	0.3495	0.1007	0.3907	1.4213	0.0918

841 **Supplemental Table S3.** Population genetic diversity statistics from neutral SNPs in each species. Sample size (n), average observed
842 heterozygosity (H_O), and expected heterozygosity (H_E) are shown, along with standard deviations (SD).
843

	Ta. granulosa					*Th. sirtalis*				
Population	*n*	H_O	H_O SD	H_E	H_S SD	*n*	H_O	H_O SD	H_E	H_S SD
Clallam	12	0.278	0.248	0.253	0.196	4	0.328	0.297	0.294	0.218
Cook Creek	13	0.324	0.237	0.294	0.178	16	0.311	0.231	0.288	0.177
Potters Slough	13	0.352	0.225	0.320	0.163	16	0.319	0.230	0.293	0.169
Warrenton	14	0.321	0.200	0.323	0.156	20	0.297	0.218	0.292	0.170
Hebo	12	0.330	0.205	0.332	0.159	14	0.292	0.218	0.302	0.168
Benton	17	0.335	0.198	0.330	0.154	14	0.337	0.270	0.288	0.184
Ten Mile	14	0.325	0.201	0.332	0.157	16	0.363	0.246	0.303	0.163
Tahkenitch	12	0.376	0.244	0.332	0.169	15	0.303	0.220	0.296	0.169
Elk River	16	0.300	0.204	0.318	0.174	17	0.306	0.232	0.285	0.180
Total	123	0.327	0.218	0.315	0.168	132	0.317	0.240	0.293	0.178

845 **Supplemental Table S4.** Pairwise F_{ST} statistics for the neutral SNP datasets of each species. For *Th. sirtalis*, F_{ST} divergence at the
846 DIV p-loop of the $Na_V1.4$ channel is also shown. F_{ST} values are shaded red to illustrate the extent of divergence among different
847 populations (white=0.00, red=1.00).
848

Ta. granulosa pairwise F_{ST} of neutral SNPs

	Clallam	Cook Creek	Potters Slough	Warrenton	Hebo	Benton	Ten Mile	Tahkenitch
Cook Creek	0.040							
Potters Slough	0.066	0.011						
Warrenton	0.077	0.024	0.004					
Hebo	0.111	0.058	0.031	0.019				
Benton	0.158	0.096	0.057	0.039	0.020			
Ten Mile	0.147	0.088	0.053	0.034	0.013	0.007		
Tahkenitch	0.144	0.088	0.055	0.040	0.022	0.016	0.009	
Elk River	0.218	0.154	0.116	0.101	0.081	0.066	0.065	0.067

Th. sirtalis pairwise F_{ST} of neutral SNPs

	Clallam	Cook Creek	Potters Slough	Warrenton	Hebo	Benton	Ten Mile	Tahkenitch
Cook Creek	0.044							
Potters Slough	0.058	0.025						
Warrenton	0.069	0.054	0.044					
Hebo	0.067	0.063	0.051	0.052				
Benton	0.120	0.104	0.084	0.082	0.045			
Ten Mile	0.066	0.070	0.058	0.062	0.024	0.056		
Tahkenitch	0.097	0.087	0.075	0.086	0.031	0.065	0.029	
Elk River	0.150	0.129	0.114	0.127	0.083	0.110	0.064	0.031

Th. sirtalis pairwise F_{ST} of DIV p-loop

	Clallam	Cook Creek	Potters Slough	Warrenton	Hebo	Benton	Ten Mile	Tahkenitch
Cook Creek	0.039							
Potters Slough	0.688	0.625						
Warrenton	0.722	0.665	0.014					
Hebo	0.575	0.497	0.034	0.154				
Benton	0.652	0.579	0.025	0.162	-0.034			
Ten Mile	0.593	0.514	0.105	0.251	-0.028	-0.019		
Tahkenitch	0.820	0.739	0.455	0.595	0.254	0.245	0.147	
Elk River	0.805	0.725	-0.010	-0.024	0.141	0.132	0.230	0.603

850 **Supplemental Table S5.** Results from cline-fitting analyses. The mean value and its geographic center point along the cline are
851 shown for each dataset, in addition to cline width. Confidence intervals (CI) are also listed.
852

		Mean	Center (km)	Lower CI	Upper CI	Width (km)	Lower CI	Upper CI
Ta. granulosa	**TTX**	0.118	193.184	168.778	200.283	284.949	207.732	363.750
	Neutral PCo1	-2.562	212.131	158.176	250.096	599.042	532.016	611.000
	STRUCTURE (K=2)	0.500	229.646	188.078	265.566	440.284	348.296	574.741
Th. sirtalis	**50% MAMU dose**	2.350	128.543	113.922	173.242	510.820	314.218	539.450
	Freq. of TTX resistant alleles	0.500	128.843	104.118	149.925	55.589	5.569	101.699
	Neutral PCo1	1.120	439.140	412.333	501.383	553.836	411.980	610.930
	STRUCTURE (K=2)	0.506	387.698	359.552	415.000	276.425	206.047	368.235

854 **Supplemental Table S6.** Results from the linear models (*Ta. granulosa*) and linear mixed models (*Th. sirtalis*) accompanying the
855 cline analyses. Separate models were fit to test whether variation in each coevolutionary trait is predicted by neutral population
856 structure from the PCoA (model 1) or phenotypic variation in the natural enemy (model 2). Model 3 includes all explanatory variables.
857 The TTX resistance of *Th. sirtalis* was measured as post-injection crawl speed, with baseline crawl speed and MAMU dose included
858 in the model as covariates.
859

TTX levels of *Ta. granulosa*	Model 1			Model 2			Model 3		
Explanatory variable	coefficient	t-value	p-value	coefficient	t-value	p-value	coefficient	t-value	p-value
PCo1	0.096	10.783	<0.001*	-	-	-	0.071	5.682	<0.001*
PCo2	-0.042	-2.575	0.011*	-	-	-	0.000	-0.003	0.998
TTX resistance of *Th. sirtalis*	-	-	-	0.559	9.028	<0.001*	0.276	2.803	0.006*
R^2	0.529			0.389			0.560		

TTX resistance of *Th. sirtalis*	Model 1			Model 2			Model 3		
Explanatory variable	coefficient	Wald χ^2	p-value	coefficient	Wald χ^2	p-value	coefficient	Wald χ^2	p-value
PCo1	-0.024	1.1743	0.279	-	-	-	0.039	0.8140	0.367
PCo2	-0.025	1.3458	0.246	-	-	-	0.034	0.6785	0.410
TTX toxicity of *Ta. granulosa*	-	-	-	-0.214	259.86	<0.001*	-0.365	2.9547	0.086 ˙
Baseline crawl speed	0.481	9.6090	0.002*	0.154	352.09	<0.001*	0.505	9.8270	0.002*
MAMU dose	0.257	46.5663	<0.001*	0.240	3212.54	<0.001*	0.285	57.5845	<0.001*
Conditional R^2	0.448			0.524			0.462		