Systematic Comparison of High-throughput Single-Cell and Single-Nucleus Transcriptomes during Cardiomyocyte Differentiation

Alan Selewa[1,4], Ryan Dohn[1], Heather Eckart[1], Stephanie Lozano[1], Bingqing Xie[1], Eric Gauchat[1,4], Reem Elorbany[2], Katherine Rhodes[2], Jonathan Burnett[2], Yoav Gilad[1,2], Sebastian Pott[2*], Anindita Basu[1,3*]

[1] Department of Medicine, University of Chicago
[2] Department of Human Genetics, University of Chicago
[3] Center for Nanoscale Materials, Argonne National Laboratory
[4] Biophysical Sciences Graduate Program, University of Chicago

* Correspondence: Sebastian Pott (spott@uchicago) and Anindita Basu (onibasu@uchicago.edu)

ABSTRACT

A comprehensive reference map of all cell types in the human body is necessary for improving our understanding of fundamental biological processes and in diagnosing and treating disease. High-throughput single-cell RNA sequencing techniques have emerged as powerful tools to identify and characterize cell types in complex and heterogeneous tissues. However, extracting intact cells from tissues and organs is often technically challenging or impossible, for example in heart or brain tissue. Single-nucleus RNA sequencing provides an alternative way to obtain transcriptome profiles of such tissues. To systematically assess the differences between high-throughput single-cell and single-nuclei RNA-seq approaches, we compared Drop-seq and DroNc-seq, two microfluidic-based 3' RNA capture technologies that profile total cellular and nuclear RNA, respectively, during a time course experiment of human induced pluripotent stem cells (iPSCs) differentiating into cardiomyocytes. Clustering of time-series transcriptomes from Drop-seq and DroNc-seq revealed six distinct cell types, five of which were found in both techniques. Furthermore, single-cell trajectories reconstructed from both techniques reproduced expected differentiation dynamics. We then applied DroNc-seq to *postmortem* heart tissue to test its performance on heterogeneous human tissue samples. We compared the detected cell types from primary tissue with iPSC-derived cardiomyocytes profiled with DroNc-seq. Our data confirm that DroNc-seq yields similar results to Drop-seq on matched samples and can be successfully used to generate reference maps for the human cell atlas.

Introduction

The identification and characterization of cell types from solid tissues and organs in the human body is the necessary basis for a comprehensive reference map of all human cells[1]. Such tissue atlases will provide a basis for understanding fundamental biological processes and to diagnose and treat disease. Single-cell RNA-sequencing (scRNA-seq) has emerged as a key tool to decompose complex tissues into cell types and states, and to investigate cellular heterogeneity[2–5]. Profiling cellular heterogeneity using thousands of cells and creating tissue level cellular maps require efficient and scalable scRNA-seq protocols. The development of microfluidic droplet-

based approaches, such as Drop-seq, has enabled transcriptional profiling of thousands of cells in
parallel[5,6]. Drop-seq has been used to characterize the cellular composition of a wide variety of
tissues and organisms, including the mouse retina[5], malaria parasites[7], and drosophila embryos[8].
However, Drop-seq requires suspensions of intact single cells for library preparation which cannot
be obtained for many tissues and cell types because of extra-cellular matrix that may be hard to
digest, fragile cell membranes, unusual cell morphology, or large cell-size. This challenge may be
addressed by adapting Drop-seq to single nuclei RNA-seq (DroNc-seq[9]). DroNc-seq obtains gene
expression profiles from isolated nuclei which are more amenable for direct dissociation from
tissues while maintaining membrane integrity. Both approaches can be used to characterize cellular
composition of complex tissues. Comparisons of low-throughput, high-coverage single cell and
single nucleus approaches suggest that both methods capture the cellular composition of
heterogeneous samples to a similar degree[10,11]. However, direct comparisons of Drop-seq and
DroNc-seq on matched samples have been limited to cell lines[9] and, more recently, samples from
mouse kidneys[12]. To establish a firm understanding of the differences and similarities of Drop-seq
and DroNc-seq, it is necessary to compare these technologies across a spectrum of different
biological conditions. A crucial aspect of single cell RNA-seq approaches is to capture cellular
heterogeneity associated with expression changes during dynamic processes, for example during
differentiation. We performed a systematic comparison of Drop-seq and DroNc-seq using time-
course data from human iPSCs differentiating into cardiomyocytes (CMs). This allowed us to
compare Drop-seq and DroNc-seq with respect to read depth, transcriptome composition, cell
types detected, and cellular differentiation trajectories. These assessments are important for
integrative analyses and interpretation of data produced using high-throughput single-cell and
single-nucleus RNA-seq in general, and with Drop-seq and DroNc-seq in particular. In addition,
we confirmed that inclusion of reads from intronic regions increases the sensitivity of DroNc-seq
and improves resolution in identifying cell types. Next, we applied DroNc-seq to frozen
postmortem human heart tissue to sample constituent cell types and compare them to CMs grown
in vitro from human iPSC. This work was conceived as part of benchmarking experiments to
establish the applicability of recent high-throughput single-nucleus RNA-seq for the Human Cell
Atlas (HCA)[1]. By identifying differences and similarities between Drop-seq and DroNc-seq, this
study will aid efforts such as the HCA that require the integration of single-cell and single-nucleus
RNA-seq data from various tissues and laboratories into a common platform.
78
79 **Results**
80
To quantitatively assess the similarities and differences in transcription profiles from single-cell
and single-nucleus RNA-seq, we performed Drop-seq and DroNc-seq, respectively, on cells
undergoing iPSC to CM differentiation, following an established protocol[13]. To compare Drop-
seq and DroNc-seq across samples with different cellular characteristics and degrees of
heterogeneity, we collected cells from multiple time-points throughout the differentiation process
(days 0, 1, 3, 7, and 15) (Figure 1A). For each technique, we obtained samples from two cell lines
per time-point, except for time-point day 15 which contains cells from a single cell line. DroNc-
seq also contains a single cell line for day 1. To approximate how many cell barcodes were
accidentally associated with 2 cells in our experiment (doublet rate), we mixed iPSCs from chimp
into the Drop-seq run from cell line 1 on day 7. These data confirmed a low doublet rate (<5%)
(Figure S1). The distributions of number of genes for each day of differentiation are shown in
Figure 1B. Overall, Drop-seq shows a higher number of genes and transcripts detected compared
with DroNc-seq, reflecting the greater abundance of transcripts in the intact cell, compared with
the nucleus alone. For our analyses, we selected cells and nuclei with at least 400 and 300 detected

genes (at least 1 UMI), respectively, and removed chimp cells from the day 7 sample. After
filtering, the mean number of genes detected per cell and per nucleus are 962 and 553, and the
mean number of UMI per cell, nucleus are 1474 and 721 for Drop-seq and DroNc-seq, respectively.
Based on the above cut-offs, we detected a total of 25,475 cells and 17,229 nuclei across all cell
lines and time-points for Drop-seq and DroNc-seq, respectively. Both cell lines were present at
each time-point in the filtered datasets (Figure 1C). Using raw RNA-seq reads, we found that top
expressed genes in Drop-seq comprised of mitochondrial and ribosomal genes, while the top gene
in DroNc-seq was the non-coding RNA, MALAT1 (Figure 1D).

103
104 Figure 1: Experimental design and preliminary data analyses. A) Two cell lines of iPSCs differentiating into CMs
105 over a 15-day time period underwent mRNA sequencing with Drop-seq and DroNc-seq. B) Boxplots showing the
106 distribution of number of genes in each day and cell line for Drop-seq (top) and DroNc-seq (bottom). C) Number of
107 cells present after applying quality control cut-offs. D) Percentage of counts for the top 15 genes in Drop-seq (left)
108 and DroNc-seq (right).
109
In addition to the differences in the number of genes detected in Drop-seq and DroNc-seq, DroNc-
seq captures a significantly higher fraction of intronic reads compared with Drop-seq (Figure 2A).
Up to 50% of the reads from DroNc-seq mapped to intronic regions, while for Drop-seq, only 7%
of reads were intronic. This discrepancy between the two techniques is expected and likely caused
by the sampling of unprocessed transcripts that are enriched in the nucleus. Intronic reads will be
detected if the transcript was not fully processed before capture by the polydT primer. In addition,
internal priming[14] on polyA stretches might lead to further sampling of introns. In order to
understand the sources of intronic reads in our dataset, we scanned the genome for polyA stretches
that are at least 5 bp long, and counted their frequency within and around each read with 20 bp
flanking regions. We found that approximately 40% of the intronic reads and their 20-bp flanking

regions contained at least one polyA stretches and that these polyA stretches were specifically
enriched towards the 3' end of reads (Figure S3). This suggests internal priming as a contributing
mechanism for intronic read sampling. RNA-seq reads aligning to introns have been used to
quantify gene expression levels previously[11,12]. Indeed, incorporating intronic reads to quantify
gene expression level improves the gene detection rate in DroNc-seq by ~2 times on average
(Figure 2B). This increase in detection rate leads to recovery of gene expression for cells which
would otherwise not be detected, as demonstrated by examples from mesoderm and cardiac genes
(Figure 2C). These data suggest that inclusion of introns can be used to compensate for the smaller
amount of nuclear RNA compared with whole cells. Accordingly, we incorporated intronic reads
into our analysis pipeline to improve gene detection rates in DroNc-seq. After intron inclusion, we
recovered 1.5 times more nuclei, bringing our total to 25,429 nuclei using a minimum of 300 genes
detected per nucleus. In addition, the mean number of UMI per cell increased from 721 to 918,
while the mean number of genes detected per cell increased from 553 to 672.

133
Figure 2: A) Distribution of reads across the genome in Drop-seq and DroNc-seq. B) Incorporating intronic reads in
quantifying gene expression increases each cell's gene detection rate by ~2X on average for DroNc-seq, enabling
detection of more genes per cell, compared with using exon reads only. C) Mesoderm and cardiac genes with
expression detected when incorporating intronic reads. D) Differential expression analysis between methods, days,
and cell lines. Genes with adjusted p-value < 0.05 and log-fold-change > 4 were kept. E) Proportion of differentially
expressed genes (DEGs) between Drop-seq and DroNc-seq associated with different gene categories.
140

To identify systematic differences in gene-specific detection rates between Drop-seq and DroNc-
seq, we obtained differentially expressed genes (DEGs) between the two techniques for matched
time-points and cell lines. As a comparison, we also performed differential gene expression
analyses between time-points and between cell lines within each technique. We detected
substantially more genes with differential expression between the two techniques than we observed
between different time-points or cell lines (Figure 2D). This phenomenon was most pronounced
for highly significant genes and became less pronounced at more lenient thresholds of log fold-
change (Figure S11). The differentially detected genes directly reflect the sampling differences in
cellular components for the two techniques. GO analysis on DEGs between Drop-seq and DroNc-
seq revealed functional annotations associated with the sampling of different cellular components
of the two techniques (Figure S5). In particular, 5% of genes detected at higher levels in DroNc-
seq were lncRNAs (compared to 1% in Drop-seq), while 20% and 6% of genes detected at higher
levels in Drop-seq were mitochondrial and ribosomal transcripts, respectively (Figure 2E).
154
Next, we tested if the differences between Drop-seq and DroNc-seq in the number of detected UMI
and enriched gene sets lead to inconsistent detection of cell types and variation in the inferred
differentiation trajectory. To infer cell types found with Drop-seq and DroNc-seq data, we
performed clustering of cells separately for each technique. We used the R package Seurat[15] to
perform normalization, dimensionality reduction, clustering, and visualization of individual cells,
grouped by cell types (see Methods). Cell types were assigned to clusters based on comparison of
genes that are significantly upregulated in the cluster to known marker genes. All genes were tested
for differential expression using a negative binomial likelihood ratio test within the Seurat package
and p-values were adjusted for multiple testing using Bonferroni correction. For each cluster, we
ordered genes by their average log-fold-change (logFC) in descending order to identify marker
genes, as genes associated with cell type have a large fold-change in expression. Note that p-values
(raw and adjusted) for all marker genes are small (adjusted $p < 10^{-5}$). We used the top marker genes
for each cluster to identify cell type specific genes (Figures S6 and S7). We found that the clusters
identified by Drop-seq and DroNc-seq captured the anticipated differentiation from iPSCs to CMs
over the course of 7 days (Figure 3A and B, Supplemental Figure 4). The cluster formed by cells
from early time-points day 0 and day 1 contained pluripotent stem cells (Figure 3A and B, 'iPSC',
orange cluster), in agreement with the expression of characteristic markers such as DPPA4. Cells
harvested on day 3 mostly formed a separate cluster ('Cardiac progenitors', green cluster)
composed of cells expressing markers concordant with cardiac progenitors (e.g. expression of
EOMES (logFC=1.08), a mesendoderm progenitor marker gene). For days 7 and 15 the clusters of
cells profiled by Drop-seq and DroNc-seq showed slight differences and we detected four clusters
in Drop-seq compared to three for DroNc-seq, indicating that Drop-seq might be more sensitive
towards detection. Drop-seq and DroNc-seq identified three clusters of ostensibly similar cell types.
Two of these clusters contained cells predominantly expressing markers of CMs, including *MYH6*,
TNNT2, MYL, and *MYBPC3* (Figure 3A, cyan cluster, 'Cardiomyocyte 1' and blue cluster,
'Cardiomyocyte 2'). We also detected a cell cluster that expressed cardiac markers alongside
markers of other lineages (e.g. *FOXA2* and *TTR*, pink cluster, 'Alternative lineage 1'). Drop-seq
revealed an additional smaller cluster (purple, 'Alternative lineage 2', expression of *FLT1*) for
which we did not find an equivalent cell population in DroNc-seq. These 'Alternative lineage'
clusters might represent cells at intermediate stages, failures of differentiation, or differentiation
towards alternative lineages. This heterogeneity and the detection of mesendodermal and
endodermal cell populations, including endothelial cells, is in agreement with previous scRNA-
seq data obtained during iPSC to cardiomyocyte differentiation[16].
188

Table S1 shows the marker genes used to identify each cell type and its corresponding cellular
prevalence. This comparison supported that both Drop-seq and DroNc-seq can identify the
predominant cell types expected in a population. Importantly, the identified clusters showed
expression of similar sets of genes in both techniques indicating that, despite differences in
detection rate between the techniques and preferential detection of specific subsets of genes the
identification of major cell types remained largely unaffected.
195
To test how concordant the cluster assignment of Drop-seq and DroNc-seq are with bulk RNA-
seq of similar cell types, we aggregated clusters representing iPSCs and iPSCs-CMs into pseudo-
bulk samples. We compared these pseudo-bulk data to bulk RNA-seq data obtained from a
previous study[17]. A total of 91 bulk RNA-seq samples composed of human iPSCs (n=18), iPSCs
differentiating into CMs (n=51), and adult primary heart tissue (n=22) were used for a correlation
analysis against pseudo-bulk iPSCs and CMs (Figure 3E). Drop-seq generally outperforms DroNc-
seq for all three sample types regardless of pseudo-bulk type by ~ 50%, which is expected as bulk
RNA-seq and Drop-seq both capture mRNA from whole cells. The iPSC pseudo-bulk samples of
both methods are best correlated with iPSCs, followed by iPSC-Cardiomyocytes and primary heart
tissue, as expected. For CM pseudo-bulk, both methods are best correlated with iPSC-
cardiomyocytes, followed by primary heart tissue, and iPSCs.
207
The time-series data allowed us to compare differentiation dynamics of iPSCs captured by Drop-
seq and DroNc-seq. We observed that several cell types were present in more than one time-point
(Figures 3 F, G). In particular, iPSCs were observed in days 0 and 1, while CMs are observed in
days 7 and 15 in both Drop-seq and DroNc-seq data. Detection of the same or similar cell types
across time-points should therefore enable us to reconstruct continuous single-cell differentiation
trajectories[14,18,19] in an unsupervised manner to characterize the temporal relationship between
different cell populations. Accordingly, we reconstructed differentiation trajectories of the cells
from DroNc-seq and Drop-seq data using Monocle[19]. In order to reduce computational time, we
selected the top 700 cells based on the number of genes detected at each time-point, for a total of
3,500 cells and used them to reconstruct the single-cell trajectory during iPSC to CM
differentiation.
219
Inferred trajectories from DroNc-seq and Drop-seq data show one and two branching points,
respectively. Coloring cells by cell type (Figures 3 H, I) and pseudo-time (Figure S9) confirms the
temporal order of cell types in Figures 3 F, G. Monocle places iPSCs at the beginning of the
trajectory, which has pseudo-time zero, followed by cardiac progenitors. Following cardiac
progenitors along the trajectory, we find one branching point in DroNc-seq which broadly
partitions CMs and the clusters associated with less well-defined cell types that might represent
alternative lineage decisions or incomplete differentiation (Figure 3). In Drop-seq, these immature
cells are on different branches and are both separated from the third branch containing CMs. These
differences might reflect the higher gene expression fold differences observed for the genes we
used to build the trajectories in Drop-seq compared to DroNc-seq. This might be a consequence of
the lower read depth observed for DroNc-seq. Both methods suggested the differentiation of iPSCs
into an intermediate cell type (cardiac progenitors), and finally a population of clearly identifiable
cardiomyocytes, based on the expression of TNNT2 and MYH6, and a divergent trajectory towards
alternative cell populations.

234
Figure 3: Cell type and single-cell trajectory analysis. A, B) Clustering results visualized with UMAP and colored by
inferred cell type for Drop-seq and DroNc-seq. C, D) Expression of marker genes overlaid on UMAP plots from A
and B for Drop-seq and DroNc-seq. E) Pearson correlation of DroNc-seq and Drop-seq pseudo-bulk against bulk
RNA-seq from iPSCs (n=18), iPSC-Cardiomyocytes (n=51), and primary heart tissue (n=22)[17]. F, G) Distribution of
cell types per time-point in Drop-seq and DroNc-seq, respectively. H, I) Inferred trajectories using Monocle with color
representing inferred cell types. A total of 3500 cells were used for the trajectory corresponding to 700 per time-point.
241

242 The comparison of Drop-seq and DroNc-seq data was motivated by the fact that Drop-seq cannot
243 be applied to generate single-cell RNA-seq data from adult primary heart tissue, but DroNc-seq
244 potentially can. Having established that DroNc-seq provides data ostensibly similar to Drop-seq
245 in our *in vitro* setup, we applied DroNc-seq to frozen human heart tissue to identify possible
246 cardiac cell sub-types and non-cardiac cells within the tissue.
247

We detected a total of 4,796 nuclei based on the presence of distinct cell barcodes using DroNc-
seq on tissue from an adult human male heart. We used both introns and exons to quantify number
of reads per nucleus, with mean number of genes and UMIs as 361 and 823, respectively. To focus
our analyses on good quality nuclei, our analyses used the top 30% (1,491) of cells based on the
number of genes detected. We performed cell type analysis on the heart cells using the same

procedure as described for the *in vitro* samples. As expected, the majority of cells (~82%) were
CMs and myofibroblasts (Fig. 4A). Figure 4B shows the distribution of marker genes for each cell
type obtained using negative binomial likelihood ratio test. A cluster was identified as CMs (Figure
4A, pink cluster) based on marker genes *TNNT2* (logFC=0.71), *MYH6* (logFC=0.87), and
MYBPC3 (logFC=1.38). A second cluster was identified as likely myofibroblasts (Figure 4A, dark-
green cluster) expressing the collagen genes *COL5A2* (logFC=1.95) and *COL6A3* (logFC=1.92)
and periostin (POSTN). Finally, a third cluster was identified as endothelial cells (Figure 4A, grey
cluster) based on vascular endothelial growth factor receptor *FLT1* (logFC=2.4) and blood clotting
protein *VWF* (logFC=1.77). A fourth clusr expressing *CPE* (logFC=2.4) and *ENPEP* (logFC=2.5)
was identified likely representing myofibroblasts (Figure 4A, black cluster). Additional marker
genes are listed in Figure S10, which shows the top 10 upregulated genes in terms of logFC in
each cluster.
265
To better understand the cell type composition of the primary heart tissue, we first aggregated data
from all nuclei into a pseudo-bulk heart sample and compared these to bulk data from iPSCs, iPSC-
CMs, and primary heart tissue as before. We found that the pseudo-bulk heart sample most closely
correlated with bulk RNA-seq data obtained from primary hearts, followed by iPSC-CMs. No
correlation was observed with bulk iPSCs (Figure 4C). Second, to compare the heart nuclei data
with the *in vitro* model we compared single nuclei of the heart to the DroNc-seq on iPSC-CMs
using correlation analysis. Figure 4D shows a bi-clustered heatmap of the Pearson correlation
coefficients with columns representing primary heart nuclei, and rows representing iPSC-CMs
nuclei. Interestingly, hierarchical clustering of each (heart) nuclei's Pearson values (columns)
confirms the clustering pattern found in Figure 4A, which demonstrates the presence of non-
cardiomyocytes within the single-nuclei primary heart sample. In particular, the cluster identified
as CMs (Figure 4A, pink cluster) has stronger correlation values with the iPSC-CMs than the
myofibroblasts and endothelial cells (Figure 4A, dark-green and grey clusters). Clustering the rows
also revealed the relative correlation strengths of the two iPSC-CMs clusters with the primary heart
nuclei. In particular, the 'Cardiomyocyte 2' cluster generally has stronger correlation with the
primary heart nuclei than the 'Cardiomyocyte 1' cluster. This is could potentially reflect the
observation that 'Cardiomyocyte 2' was associated with cells collected on day of our
differentiation protocol and therefore to closer towards the mature state of CMs. We used iPSCs
as an out-group for which we expect no correlation with primary heart nuclei, which is observed
to be the case (Figure 4D).

286
287 Figure 4: Application of DroNc-seq on human heart tissue. A) Cell type analysis visualized with UMAP. B)
288 Distribution of marker genes identified with differential expression analysis. All genes listed have p-values < 10^{-29}. C)
289 Pearson correlation of primary heart pseudo-bulk against bulk RNA-seq from iPSCs (n=18), iPSC-Cardiomyocytes
290 (n=51), and primary heart tissue (n=22)[17]. D) Bi-clustering on Pearson correlation values of primary heart nuclei with
291 nuclei from iPSCs and iPSC-derived cardiomyocytes.
292

Discussion

Building a cell atlas of the human body requires the expression profiling of all human tissues from
a range of different samples, including tissues that are hard to dissociate, composed of fragile cells,
and frozen specimens, all of which are incompatible with single-cell RNA sequencing. As an
alternative, DroNc-seq, a high-throughput single-nucleus RNA sequencing protocol, has the
potential to reveal tissue heterogeneity, at scale, based on *nuclear* RNA, and is being increasingly
used to profile primary tissue at high throughput. However, it is unclear how DroNc-seq compares
with earlier single-cell RNA-seq protocols like Drop-seq across a range of different cell types and
tissues. Previous studies have performed cell type comparisons using nuclear vs. whole-cell RNA
using full-length mRNA sequencing assays at low throughput[10,11]. Drop-seq and DroNc-seq have
been compared using adult mouse kidneys cells[12]. We performed a direct comparison of high-
throughput, single-cell (Drop-seq) and single-nucleus (DroNc-seq) RNA-seq using iPSCs
differentiating into CMs. Together with single-nucleus profiling of primary CMs from adult human
heart tissue, this study enabled us to compare cell type detection, transcriptome profiling and infer
cellular differentiation with two complementary high-throughput techniques, using an *in vitro*

model of CM differentiation, and compare them directly to human primary CMs obtained from a
frozen heart sample (see Methods) using DroNc-seq.
311
As expected, the number of UMIs per nucleus in DroNc-seq are lower than those for cells in Drop-
seq. Consequently, the gene detection rate in DroNc-seq was significantly lower than for Drop-
seq (Figure 1C). However, given the high number of reads in DroNc-seq that mapped to intronic
regions we reasoned that inclusion of such reads might increase the gene detection rate. Indeed,
intron inclusion significantly increased the sensitivity of DroNc-seq and improved cluster
separation and cell type identification, in agreement with previous studies[10–12]. We also found that
the inclusion of introns increased gene detection rate in single nuclei samples. Of note, a significant
proportion of the intronic reads seems to originate not from transcripts primed at the 3' end but
from direct priming to polyA stretches in introns[14] (Figure 2). While such reads still scale with the
expression level of a transcript, the assumption that transcript levels are uniquely quantified by a
single UMI may be violated in these cases.
323
Given the difference in input material, i.e., cellular vs. nuclear RNA, it is not surprising that we
found a significant proportion of genes that are differentially expressed between Drop-seq and
DroNc-seq samples. Some of the most highly enriched sets of genes reflected the technical
differences between the two technologies. Genes specifically enriched in Drop-seq are ribosomal
and mitochondrial. DroNc-seq presumably loses these transcripts that are predominantly localized
in the cytoplasm. Conversely, as a class, lncRNAs are enriched in DroNc-seq which agrees with
the nuclear localization of many of them.
331
Expression profiles in Drop-seq and DroNc-seq confirmed the differentiation of iPSCs into CMs
and revealed major cell types found within the *in vitro* differentiation model of iPSC-CMs. These
data also confirmed heterogeneity observed during differentiation. Drop-seq and DroNc-seq
detected a population of cardiac progenitors with cellular prevalence 23.3% and 18.2%,
respectively. They also both detected two clusters representing CMs: cardiomyocyte 1 (16.1% and
5.6% prevalence) and cardiomyocyte 2 (4.2% and 12.7% prevalence). Both methods also revealed
a population of cells, 'Alternative lineage 1', that might represent alternative fate or that failed to
reprogram fully, which accounted for 5.9% and 11.3% of all cells in Drop-seq and DroNc-seq,
respectively. The presence of non-CMs during late-stage is expected for the *in vitro* differentiation
model and has been observed previously[16]. Accordingly, the proportion of cells differentiating into
CMs expressing TNNT2, assessed by FACS, varies widely between 20-80%[13]. Based on our cell
type assignment in Drop-seq data, we obtained 28% and 29% cardiomyocytes on day 7 for the two
cell lines and 70% CMs on day 15 for cell line 2, which fall within the expected range.
345
Drop-seq revealed an additional smaller cluster (purple, 'Alternative lineage 2', expression of
FLT1 and comprising 1.4% of the total population) for which we did not find an equivalent cell
population in DroNc-seq. The reasons behind the failure of DroNc-seq to identify the small
fraction of cells identified as 'Alternative lineage 2' in Drop-seq may be due to the lower capture
rate of DroNc-seq (mean number of detected genes was 672) compared to Drop-Seq (mean number
of detected genes was 962) (Figure S8) which might result failure of the clustering approach to
resolve this sub-population in DroNc-seq, or due to the preferential loss of the particular cell type
arising from DroNc-seq's nuclei dissociation protocol. The mean number of genes detected in this
subpopulation in Drop-seq was 1032, representing the cluster with the highest gene detection rate.
It is possible that this facilitated the detection of this cluster in Drop-seq while the lower detection
rate in DroNc-seq combined with the small number of cells corresponding to this cluster in the

sample lead to the loss of this population during clustering. However, we cannot rule out specific
loss or selection biases for of the cell type introduced during DroNc-seq sample preparation.
359
We chose the iPSC-to-CM differentiation because in addition to cell type detection, the highly
heterogenous but temporally coordinated process allowed us to compare cellular lineages inferred
based on Drop-seq and DroNc-seq data, respectively. Indeed, we were able to infer similar
trajectories for both Drop-seq and DroNc-seq (Figure 3H and I). Both trajectories show continuous
differentiation of iPSCs into cardiac progenitors along a single path, which then branches into CM
and non-cardiac cells (progenitor cells and alternative lineages). This suggests that a substantial
proportion of cells identified as CM progenitors in our cluster analysis are diverging from the
differentiation trajectory relatively early on and ultimately are not becoming mature
cardiomyocytes[16]. In the case of Drop-seq 'Alternative lineage 1' and 'Cardiac progenitor' cells
are branching off on two separate points, while for DroNc-seq both populations are on one branch.
The additional branching point might reflect the higher resolution achieved by Drop-seq.
371
Compared with bulk samples, Drop-seq pseudo-bulk is closer to tissue-level expression than
DroNc-seq. This is expected as the tissue data represents RNA-seq data generated using whole
cells, rather than nuclei. However, this difference does not mask cell type specific differences in
the degree of correlation with bulk samples from iPSCs, iPSC-CM, and heart. Both Drop-seq and
DroNc-seq CM pseudo-bulk correlate the best with bulk iPSC-CMs samples followed by primary
heart tissue and iPSCs. While the iPSCs correlate best with the bulk iPSCs for both methods. The
comparison with bulk samples provides further evidence for the cell type labels that were assigned
based on marker genes.
380
Having demonstrated that Drop-seq and DroNc-seq performed similarly in detecting heart-like cell
types, we applied DroNc-seq to primary heart tissue from adult human male. As expected, cell
type analysis of the tissue revealed mostly CMs (43%) and (myo)fibroblasts (39%), as well as a
smaller population of endothelial cells (12%). Interestingly, TNNT2 was detected in all the cell
types but was significantly upregulated in the CM cluster. TNNT2 being a marker gene for CMs
suggested the possibility that all nuclei are of the same broad category of cell type. Correlating
transcription profiles from primary heart nuclei with the iPSC-derived CM nuclei further supports
the inferred cell types from the primary heart tissue. The transcriptome profiles of primary heart
nuclei that were assigned to the 'Cardiomyocyte' cluster are more strongly correlated with the
profile of iPSC-CMs compared with primary heart nuclei in other clusters.
391
Sequencing of additional cells and increased read depth will help to increase the resolution and
potentially lead to detection of additional cell types. However, it is important to keep in mind that
tissue samples are not uniform mixtures of cell types. Thus, the creation of comprehensive cell
maps likely requires sampling of a given tissue in multiple different locations, as seen from the
relatively low cell type complexity in DroNc-seq data on the human heart tissue when sampled
from only one anatomical region.
398
399 This comparison of Drop-seq and DroNc-seq demonstrates the capability of DroNc-seq in
400 dissecting the multicellular environment within complex tissue such as the heart, which would
401 otherwise not be possible with Drop-seq. We expect that DroNc-seq will be used to perform high-
402 throughput transcriptomic profiling of tissues for which it is difficult to obtain suspensions of intact
403 single cells and aid in initiatives such as the Human Cell Atlas and the Human Tumor Atlas.
404

Methods

Cell Culture and Differentiation

We used iPSCs from two individuals from a previously established panel of LCL-derived iPSCs[20]. iPSCs were seeded on 100 mm dishes 3-5 days prior to differentiation. At 70-100% confluency, growth media was replaced with heart media: RPMI (Thermo Fisher Scientific, 14-040-CM) supplemented with B-27 Supplement minus insulin (Thermo Fisher Scientific, A1895601), 2 mM GlutaMAX (Thermo Fisher Scientific, 35050-061), and 100 mg/mL Penicillin/Streptomycin (Corning, 30002Cl). A heart medium/Matrigel mix was made using this medium along with a 1:100 dilution of Matrigel (Corning, 35427) and 12 uM of the GSK-3 inhibitor CHIR99021 trihydrochloride (Tocris, 4953). This medium was changed to base heart media 24 hours later (Day 1). On Day 3, the previously described medium was replaced with heart medium containing 2 μM Wnt-C59 (Tocris, 5148). On days 5, 7, 10, 12 and 14 of the differentiation, media was refreshed with base heart media. Heart medium changes occurred daily. Beating CMs cells were observed around Day 7.

Cell Processing

At each time-point, cells were harvested from 100 mm plates by treating with Accutase (BD Biosciences, #561527) to generate a single cell suspension; from Day 7 onward, a cell scraper was also employed to release adherent cells from plates. Cells were centrifuged at 300 xg for 5 minutes and supernatant was aspirated off. Cells were washed 3 times with 1X PBS, 0.01% BSA (NEB, #B9000S), henceforth called PBS-BSA). 10 μL of cells was combined with trypan blue for counting in an NI hemocytometer (InCyto, DHC-N01-2). Viability of cells at each time point was recorded (see Table 1). Cells were also labelled with a combination of 4′,6-diamidino-2-phenylindole or DAPI (Sigma, Cat #D9542) and Wheat Germ Agglutinin (WGA; Thermo Fisher Scientific, W11262) to assess nucleus and cell membrane integrity under fluorescence imaging, as shown in Figure 5A. 400,000 cells were taken and suspended in 2 mL PBS-BSA (200,000 cells/mL) for Drop-seq, and the remaining cells were used for nuclei isolation for DroNc-seq.

Table 1: Viability of harvested cells from each iPSC-CMs differentiation time-point

Time Point	Date	Viability
Time Course 1 Day 0	11/16/2017	70%
Time Course 1 Day 1	11/15/2017	50%
Time Course 1 Day 3	11/17/2017	80%
Time Course 1 Day 7	11/21/2017	60%
Time Course 2 Day 0	1/22/2018	60%
Time Course 2 Day 1	1/23/2018	80%
Time Course 2 Day 3	1/25/2018	80%
Time Course 2 Day 7	1/29/2018	90%
Time Course 2 Day 15	2/6/2018	55%

Nuclei were isolated using the Nuclei EZ Prep isolation kit (Sigma, Cat #NUC-101). Briefly, cells were resuspended in 4 mL EZ Prep Lysis Buffer and incubated on ice for 10 minutes. After incubation, cells were agitated using a P1000 pipette and 10 μL of sample was imaged. DAPI (Sigma, Cat #D9542) and Wheat Germ Agglutinin (WGA; Thermo Fisher Scientific, W11262) were used to determine if the cellular membrane had properly lysed for each cell. If intact cells were still present, 2 mL of sample was moved to a glass dounce tissue grinder (Sigma, Cat #D8938)

and dounced 5 times. After douncing, another 10 μL sample was imaged under the microscope
with DAPI and WGA staining as before to determine if high-quality, intact nuclei were obtained
(see Figure 5B). We adjusted the number of dounces until only nuclei were found. As iPSCs
differentiated further into CMs, the number of required dounces needed to be increased. For
example, day 3 of differentiation required 5 dounces to obtain proper cell lysis and intact nuclei,
while Day 7 required 12 dounces. Nuclei were spun down at 500 xg for 5 minutes at 4 °C. After
centrifugation, the nuclei were washed with nuclei suspension buffer (NSB; 1X PBS, 0.01% BSA,
and 0.1% RNAse inhibitor (Lucigen, #F83923)), resuspended in 2 mL NSB and filtered using a
35 μm cell strainer (Corning, #352235). 10 μL of nuclei suspension was sampled using a NI
hemocytometer and the concentration adjusted to a final loading concentration of 300,000
nuclei/mL in NSB of which 2 mL was used for DroNc-seq.
453
<u>Microfluidic Co-encapsulation of Cells/Nuclei and Barcoded Beads</u>
For **Drop-seq**, 2 mL of cells at 200,000 cells/mL in PBS-BSA was loaded in a 3 mL syringe (BD,
#309657). A custom-built 90 μm Drop-seq microfluidic device (CAD file supplied separately) was
used for droplet generation, creating droplets smaller than the standard Drop-seq protocol.[5] We
chose to use the 90 μm droplets because the effective concentration of cellular RNA in the 90 um
drops is doubled, leading to better RNA capture, compared to 125 μm droplets used in Drop-seq.
Indeed, we see an increase in RNA capture for cells of smaller size, such as iPSC. We note that
the increase in capture efficiency often fails to translate to larger sized cells (~15 μm), likely due
to the higher concentrations of the lysed cell's endogenous RNase and lysosomes, etc. in the drop.
Cells at 200,000 cells/mL and ~2,600,000 droplets/mL (droplet volume is ~380 pL) amounts to a
Poisson loading distribution with $\lambda \approx 0.076$. DNA barcoded beads (ChemGenes, Macosko-2011-
10(V+)) were washed, filtered, and suspended in Drop-seq lysis buffer, also at 200,000 beads/mL
and kept in suspension under constant stirring using a magnetic tumble stirrer and flea magnet
(V&P Scientific, VP 710 Series, VP 782N-3-150). Beads and cells were co-flowed into the device,
each at 3 mL/hr, along with a surfactant-oil mix (BioRad, #1864006) at 12 mL/hr that was loaded
into a 10 mL syringe (BD, #302995) and used as the outer carrier oil phase. Reverse emulsions
droplets were generated at ~3000 drops/sec and collected in two batches of 20 minutes each in 50
mL tubes (Genesee Scientific, #28-106). After collection, the standard Drop-seq protocol for bead
recovery, washing, and reverse transcription was followed.[5] After washes and DNaseI treatment
as per Drop-seq protocol[5], cDNA amplification was performed on 75,000 RNA-DNA barcode
bead conjugates in a 96-well plate (Genesee Scientific, #24-302) loaded at 5000 beads per well,
for a total of 15 wells and amplified for 15 PCR cycles using template switching.[5] Post-PCR
cleanup was performed by removing the STAMPs (Single Transcriptome Attached to Micro-
Particles[5]) and pooling the supernatant from the wells together into a single 1.7 mL tube (Genesee
Scientific, #22-281LR) along with 0.6X Ampure XP beads (Beckman Coulter, #A63880). After
adding the Ampure beads to the PCR product, the tube was incubated at room temperature for 2
minutes on a thermomixer (Eppendorf Thermomixer C, #5382000023) set to 1250 rpm, and for
another 2 minutes on bench for stationary incubation. Next, the tube was placed on a magnet, and
4X 80% ethanol washes were performed with 1 mL ethanol added in each wash. cDNA was eluted
in 150 μL of water and the concentration and library size were measured using Qubit 3 fluorometer
(Thermo Fisher) and BioAnalyzer High Sensitivity Chip (Agilent, #5067-4626). A BioAnalyzer
trace is provided in Figure 5C as an example of the amplified transcriptome obtained from a Drop-
seq run. 450 pg of the cDNA library was used in Nextera Library prep, instead of 650 pg as
suggested in the Drop-seq protocol[5] to obtain Nextera libraries between 300 – 600 bp.
488

For **DroNc-seq**, a 75 μm microfluidic device[9] was used. 2 mL of nuclei at 300,000 nuclei/mL were
loaded into a 3 mL syringe and flowed at 1.5 mL/hr. Barcoded beads were filtered with a 40 μm
filter to select for smaller beads to prevent clogging events in the relatively smaller microfluidic
channels. 2 mL of beads were suspended at 350,000 beads/mL in Drop-seq lysis buffer, loaded in
a 3 mL syringe, kept suspended through a magnetic tumble stirrer, and flowed at 1.5 mL/hr, along
with carrier oil-surfactant mix loaded in 10 mL syringe and flowed at 12 mL/hr. Droplets were
generated at ~4,500 drops/sec and collected in 50 mL tubes in two batches for 22 minutes each.
After collection, the standard DroNc-seq protocol for bead recovery and reverse transcription was
followed.[9] cDNA amplification was performed on the STAMPs as above, for 15-20 wells at 5000
beads per well, for 15 PCR cycles. Cleanup was performed after removing the STAMPs and adding
0.6X Ampure XP beads (Beckman Coulter, #A63880) to the pooled supernatant followed by room
temperature incubation for 2-minutes on an Eppendorf thermomixer set to 1250 rpm and another
2-minute stationary incubation. Tubes were placed on a magnet and beads were allowed to migrate
prior to 4X washes in 80% ethanol. cDNA was eluted in 10 μL of water per well and DNA
concentration was measured using a Qubit 3 fluorometer (Thermo Fisher). 650 pg of DNA was
used in each Nextera reaction for fragmenting, tagging, and amplifying to create Nextera library.
Nextera library size and concentrations were determined using a BioAnalyzer DNA High
Sensitivity Chip (Agilent, #5067-4626).
507

508
509 Figure 5: Experimental quality control metrics. Images of Day 1 of differentiation of human iPSC derived
510 cardiomyocyte (iPSC-CM) cells- A) and nuclei- B) stained with DAPI and WGA; C) BioAnalyzer traces of WTA
511 product from Drop-seq on iPSC-CM Day 7, DroNc-seq on iPSC-CM Day 7, and DroNc-seq on archived adult male
512 heart tissue.
513

<u>Nuclei Isolation from Adult Human Heart Tissue</u>
Post-mortem human heart tissue was provided by the National Disease Research Interchange
(NDRI). The sample (m, 68 yrs) had been stored at -80ºC for 11 years before it was processed for
DroNc-seq. The frozen tissue sample was weighed and cut with a scalpel and 32.8 mg of sample
was processed, by mincing with the scalpel. The sample was placed into a glass dounce tissue
grinder (Sigma, Cat #D8938) with 2 mL of ice-cold EZ-Prep lysis buffer from the Nuclei EZ-prep
Isolation Kit (Sigma, Cat #NUC-101). The tissue was dounced 25 times with Pestle A, transferred
to a conical tube with an additional 2 mL lysis buffer, and incubated on ice for 5 minutes. Sample
was then centrifuged at 500 xg for 5 minutes at 4 ˚C. Supernatant was aspirated off and replaced
with 2 mL lysis buffer. Sample was transferred back to the tissue grinder and dounced 25 times
with Pestle B. Sample was then put back into a conical tube with an additional 2 mL lysis buffer,
centrifuged, and washed with 4 mL lysis buffer followed by 5-minute incubation on ice. 10 μL of
sample was taken and combined with DAPI and Wheat Germ Agglutinin (WGA) and put into an
NI hemocytometer (InCyto, DHC-N01-2) to check for nuclei quality. If whole cells were still

present, additional douncing with Pestle B was performed (additional 25-35 dounces expected)
before checking again using DAPI and WGA. The resulting nuclei were centrifuged, lysis buffer
was aspirated, and nuclei were washed and resuspended in Nuclei Suspension Buffer (NSB; 1x
PBS, 0.01% BSA, and 0.1% RNAse inhibitor (Lucigen, #F83923)). Nuclei were filtered once with
a 35 μm cell strainer (Corning, #352235), once with a 20 μm filter (pluriSelect, #43-50020-01),
and twice with a 10 μm filter (pluriSelect, #43-50010-01) and stored on ice for processing. Nuclei
were counted using an NI hemocytometer and brought to a final concentration of 300,000
nuclei/mL in 2 mL NSB for DroNc-seq. To assess the quality of RNA from the archived heart
tissue, we ran an independent experiment to extract total RNA using a Qiagen kit (Qiagen, #74004)
and measured using a BioAnalyzer RNA 6000 Pico kit (Agilent, #5067-1513). A RIN score of ~5
was obtained for this sample.
539
DroNc-seq on Nuclei Harvested from Heart Tissue
DroNc-seq was performed as previously described with a few exceptions: single 30-minute droplet
collection was performed using a 75 μm microfluidic device and flow rates mentioned previously.
During whole transcriptome amplification, 12 cycles of PCR were performed on 30 wells with
5000 barcoded beads per well. Clean-up was performed as described above. cDNA from each well
was eluted in 2 μL of water and pooled for quantification by BioAnalyzer (Figure 5C) and Qubit,
followed by Nextera library preparation.
547
Sequencing
Drop-seq and DroNc-seq samples for each differentiation time-point were sequenced in a single
run, with 150-200 million reads allocated per sample. Sample libraries were loaded at ~1.5 pM
concentration and sequenced on an Illumina NextSeq 500 using the NextSeq 75 cycle v3 kits for
paired-end sequencing. 20 bp were sequenced for Read 1, 60 bp for Read 2 using Custom Read 1
primer,   GCCTGTCCGCGGAAGCAGTGGTATCAACGCAGAGTAC[5],   according   to
manufacturer's instructions. Illumina PhiX Control v3 Library was added at 5% of the total loading
concentration for all sequencing runs.
556
RNA-Seq Data Processing and Analyses
The differentiating iPSCs were sampled at specific timepoints during a 15-day period (days 0, 1,
3, 7, 15) using both Drop-seq and DroNc-seq (Fig 1A). A total of 17 sequencing runs were
performed over the course of the differentiation. Each sequencing run produced paired-end reads,
with one pair representing the 12 bp cell barcode and 8 bp unique molecular identifier (UMI), and
the second pair representing a 60 bp mRNA fragment. We developed a Snakemake[21] protocol that
takes a FASTQ file with such paired-end reads as input and produces an expression matrix
corresponding to the UMI of each gene in each cell. The protocol initially performs *FastQC*[22] to
obtain a report of read quality. Next, it creates a whitelist of cell barcodes using *umi_tools*[23] *0.5.3*,
which is a list of cell barcodes with at least 30k reads. Next, each paired-end read is combined into
a single read where the read name contains the cell barcode and UMI extracted from paired end
read 1, and the sequence content corresponds to paired end read 2. This is done for every paired
end read and placed into a single "tagged" FASTQ file. The tagged FASTQ file contains only the
cell barcodes found in the whitelist. Finally, the protocol trims the ends of reads to remove polyA
sequences and adaptors using *cutadapt*[24] *1.15*. The tagged and trimmed FASTQ file is aligned to
the human reference genome (version GRCh38) using the *STAR*[25] aligner version 2.5.3, which
returns a BAM file sorted by coordinate. Next, we use *featureCounts*[26] version 1.6.0 to assign each
aligned read to a feature on the genome. Finally, we use the *count* function from *umi_tools* to
create a count matrix representing the frequency of each feature in the BAM file. The pipeline is

available at `github.com/aselewa/dropseq_pipeline`. A total of 17 count matrices
were produced by this pipeline, 9 of which correspond to Drop-seq and 8 correspond to DroNc-
seq. In order to incorporate introns into the counting process, the UMI count of a gene was
calculated as the sum of its exon and intron UMIs. This is particularly important for DroNc-seq as
approximately half the reads obtained come from intronic regions of pre-spliced mRNA.
GENCODE version 28 annotations contain exon features and gene features but do not contain
intron features. To derive an intron annotation file, we used exon and gene features. Exon regions
were subtracted from gene regions (on the same strand) and the remainder was counted as the
intron region for said gene. Then the expression level of a gene is given by the sum of the number
of intron and exons.
586
From each sequencing run, approximately 5000 cells were obtained with an average read depth of
30k – 40k per cell. Low quality cells were filtered based on the number of genes detected. A gene
was considered detected in a cell if there was at least 1 UMI present. Cells with less than 400 genes
and nuclei with less than 300 genes detected were removed. Low quality genes were also filtered
if they were not detected in at least 10 cells, in order to reduce noise and computation cost. The
total numbers of cells remaining were approximately 23,554 and 24,318 for Drop-seq and DroNc-
seq, respectively. After filtering, all expression matrices from Drop-seq experiments were merged
into a single expression matrix. The merging was done by taking the union of all genes. If a
particular dataset did not contain a gene that is expressed in another dataset, we set the expression
level to zero in the first dataset. Similarly, all expression matrices corresponding to DroNc-seq
were merged into a single expression matrix. Both merged matrices were processed and analyzed
separately downstream. Seurat[15] was used to perform normalization, clustering, and cell type
analysis. R scripts used for the analyses in this paper are documented at
`github.com/aselewa/czi`.
601
Internal Priming
We used the MEME[27] suite to find all 5 bp stretches of adenines using the human genome build
hg38. Next, we merged all 5 bp motifs in order to obtain all continuous polyA tracts. A total of ~2
$\times 10^7$ motifs at least 5 bp long were identified genome-wide. BAM files from each time-point
were merged and only intronic reads were kept. Intronic reads were extended by 20 bp on each
side and intersected with the adenine motifs in a strand-specific way. The motifs were centered by
the coordinates of the reads they intersect with and a histogram motif of 3' positions was obtained
(Figure S3).
610
Normalization and Scaling
Following the analysis procedure recommended by Seurat, we first normalize the count data. Each
cell's gene-specific UMIs were divided by the total number of UMI in the cell scaled to 10^4, which
yields TP10k (transcripts per 10k) values. Figure S2 shows the relationship between the mean
expression (mean TP10k) and the length of the gene. The relationship is relatively weak, therefore
normalizing by just the library size is sufficient. A pseudo-count of 1 was added to all scaled values
followed by a natural log transformation. After the log-transformation, the values were
standardized, i.e. mean-centered and scaled such that each gene has unit variance. These log-
normalized, and standardized data were used in downstream analyses to perform dimensionality
reduction and reconstruction of differentiation trajectories.
621
Dimensionality Reduction

The first step performed in dimensionality reduction is principal components analysis (PCA). Prior
to PCA, Seurat calculates the gene dispersion vs. mean expression in order to obtain a subset of
highly variable genes, which reduces the computational time of PCA compared with using the
entire subset of genes identified in the experiment. Highly variable genes were selected based on
a threshold of 1.5 for the dispersion level and a minimum expression level of 0.15 (on log scale)
yielding 400 genes and 350 genes with Drop-seq and DroNc-seq, respectively. These highly
variable genes were used to calculate principal components for Drop-seq and DroNc-seq data. The
top 7 principal components, which explained 60% and 70% of variation for Drop-seq and DroNc-
seq, respectively, were used to perform clustering and the results were visualized with the Uniform
Manifold Approximation and Projection (UMAP[28]), which produced a 2-dimensional visualization
of the data (Figures S4 A, B left). We also performed tSNE on the same data (Figures S4 A, B
right) using a perplexity of 50 and found that UMAP captures more of the global structure in the
data, as previously reported[29]. A minimum distance of 0.5 and 0.6 were used in UMAP for Drop-
seq and DroNc-seq, respectively.
637
Cell type Analysis
The principal components were used for graphical clustering using the *FindClusters* command of
Seurat. A resolution parameter of 0.13 is used to obtain 6 clusters in Drop-seq and 5 clusters in
DroNc-seq. In order to determine cell types from the clusters, we performed differential expression
analysis using the *FindAllMarkers* function and *negbinom* test in Seurat. This identifies
differentially expressed genes between every two groups of cells using a likelihood ratio test of
negative binomial generalized linear models. The Seurat's *negbinom* test yields relatively low false
positive rates for differential expression analyses, compared with other parametric methods[30]. The
p-values were adjusted for multiple testing using the Bonferroni correction. Furthermore, as we
were only interested in upregulated genes as these will define the cell type, we ordered genes in
each cluster, by their average log-fold-change (logFC) in descending order. Marker genes were
identified based on functional annotations as these genes associated with cell types have a large
fold-change in expression. Figures S6 and S7 show the top 10 differentially expressed genes in
each identified cluster for Drop-seq and DroNc-seq, respectively.
652
Pseudo-bulk Analysis
Raw RNA-seq counts were obtained from GEO accession GSE110471 and the human samples
were extracted from the population. The raw counts were converted into log-TP10k's. After
filtering low-quality cells, Drop-seq and DroNc-seq counts were aggregated (summed) for each
gene, and the resulting counts were converted to log (TP10k + 1). The Pearson correlation between
pseudo-bulk and each bulk RNA-seq sample was calculated using the *cor* function in *R 3.5.1* across
~6,000 genes.
660
Single-cell Trajectory Analysis
Monocle version 2.6.4 was used to construct single-cell differentiation trajectories. Computing the
trajectory of approximately 20,000 cells is computationally expensive and slow with Monocle. To
overcome this, we used the best 700 cells from each time-point. In particular, cells were ordered
by their detection rate (number of genes detected) and 700 cells with the highest detection rate
were chosen. The computation is also expensive and slow when the number of genes is high
(>10,000 genes). Selection of genes for trajectory analysis, or feature selection, is critical for
obtaining accurate trajectories. In our case, we used all of the differentially expressed genes in the
cell type analysis. The data given to Monocle are log-transformed TP10k values. The
reduceDimension function with the *DDRTree* method was used to obtain a 2-dimensional

representation of the developmental trajectories in each dataset. The cells were then ordered using
the *orderCell*s function, which infers the trajectory in reduced-dimension dataset using reserve
graph embedding[17]. A total of 3,500 cells (700 per time-point, 5 time-points in total) were used to
infer the trajectories in Drop-seq and DroNc-seq.
675
<u>Primary Heart Tissue Analysis</u>
A total of 4796 nuclei obtained from post-mortem adult human male heart tissue were profiled
using DroNc-seq. Genes were quantified using both introns and exons, with mean number of genes
and UMIs of 361 and 823, respectively. The top 30% of cells were chosen based on number of
genes detected, which corresponds to 1,491 cells. Transformation of data and cell type analysis
was performed in the manner described above. Next, we calculated the Pearson correlation
coefficient using the *cor* function in *R 3.5.1* between primary heart nuclei and the *in vitro* iPSC-
derived CMs profiled by DroNc-seq. We also used iPSCs profiled by DroNc-seq as an out-group.
A total of 200 iPSC-derived CMs and 50 iPSCs were used for the correlation analysis. For each
primary heart nuclei, a total of 250 correlation coefficients were calculated using ~2500 genes,
which we call the correlation profile of a cell. The resulting matrix of correlation values were
visualized and bi-clustered with the *heatmap.2* function in *R 3.5.1*.
688

689 **Data Availability**
690 All   raw   data   are   available   through   the   Human   Cell   Atlas   Portal
691 (https://prod.data.humancellatlas.org/explore/projects/c765e3f9-7cfc-4501-8832-79e5f7abd321).
692 All code used for analysis is available at github.com/aselewa/czi and github.com/aselewa/
693 dropseq_pipeline.
694
695 **Acknowledgement**
696
We thank Megan Rowton, Alex Guzzetta, and John Blischak for helpful comments on the
manuscript. This work was supported by the Chan-Zuckerberg Initiative pilot award #2017-
174052. RE was supported by the NIH MSTP Training Grant T32GM007281. KR was supported
by NIH GRTG 5T32GM007197 and AHA Predoctoral Fellowship 18PRE34030197. SP was
supported by the National Center for Advancing Translational Sciences of the NIH (K12
HL119995). This work was performed, in part, at the Center for Nanoscale Materials, a U.S.
Department of Energy Office of Science User Facility, and supported by the U.S. Department of
Energy, Office of Science, under Contract No. DE-AC02-06CH11357.

Supplementary Methods
705
<u>Species-mixing and single-cell specificity</u>
For the Drop-seq experiment on biological replicate #1, chimpanzee iPSCs[20] were mixed with
human iPSC-derived CMs from day 7 of the differentiation time-point, in order to assess the
frequency of doublets during cell encapsulation. We used chimpanzee cells for the species mixing
as these cells were grown using identical conditions as the human cells. The alignment protocol
was adjusted so that each read was aligned to both the human genome (GRCh38) and the chimp
genome (panTro5) separately. For each cell that passed quality control, we counted the number of
reads that aligned exclusively or with a better score to the genome of one of the species (Figure
S1). We then used the ratio of these counts as a 'species-specificity' score for each cell. We found
only a small number of cells with scores that could suggest mixing of cells from human and chimp
(< 5%), similar to previously reported estimates[5]. Cells with intermediate scores had typically

lower read counts and were thus removed by filtering based on read depth. We only kept cells with
a specificity score above 0.6 yielding ~739 cells. In agreement with our assignment, > 99% of
these cells were associated with clusters that we identified as CMs while none were associated
with iPSC clusters.

721
722   Figure S1: Scatterplot of number of reads assigned to hg38 vs panTro5 for each cell in Drop-seq day 7, cell line #1 as
723   part of a species-mixing experiment using human iPSC derived cardiomyocytes and chimpanzee iPSCs.
724
725
726

727
728   Figure S2: Mean expression (log) vs. gene length for Drop-seq (left) and DroNc-seq (right).
729
730
731

732
733 Figure S3: Density curves of the position of polyA at the 3' end. Green and red curves represent reads mapping to
734 the forward and reserve direction, respectively. The dashed line represents the average read length.
735
736

737
738 Figure S4: Dimensionality reduction for A) Drop-seq and B) DroNc-seq using tSNE (left) and UMAP (right). Color
739 represents the differentiation time point.

746
747 Figure S5: Gene enrichment analysis on differentially expressed genes between Drop-seq and DroNc-seq.
748
749
750
751

753 Figure S6: Heatmap of expression values of top 10 differentially expressed genes in each cell type cluster for Drop-
754 seq.
755
756

757
758 Figure S7: Heatmap of expression values of top 10 differentially expressed genes in each cell type cluster for DroNc-
759 seq.
760
761
762 Table S1: Breakdown of cell types and associated genes discovered in Drop-seq and DroNc-seq

Markers	Cell type	Prevalence (Drop-seq)	Prevalence (DroNc-seq)	Drop-seq Only Genes (top 5)	DroNc-seq Only Genes (top 5)
DPPA4	iPSC	48.9%	52%	SFRP2, AC025465.1, ESRG, CACNAD2D3, BDNF-AS	RIMS2, RPL8, GOLGA4, EIF4A2, SET
EOMES APLNR	Cardiac Progenitor	23.3%	18.2%	CER1, LHX1, CYP26A1, IRX3, MT-TS2	GRIB2B, AL3365295.1, IL1RAPL2, KCNQ5, NRX3
MYH6 TNNT2	Cardiomyocyte 1	16.1%	5.6%	MYL3, NPPA-AS1, NPPA,	AC012574.1, AC105233.5, MYO1D,

				ACTN2, TNNC1	ARHGAP42, CDK14
MYH6 TNNT2 AFP SERPINA1	Cardiomyocyte 2	4.2%	12.7%	AMBP, APOA2, SERPINF2, ITIH2, SERPINA5	KCNH7, ERBB4, ZBTB20, NRG3, KCNQ5
TTR FOXA2	Alternative Lineage 1	5.9%	11.3%	GATA3, S100A14, HHEX, FLIRT3, EPSTIL1	EWSR1, PTBP2, ZMYM2, LUC7L, LINC01876
CD34 SCARF1 FLT1	Alternative Lineage 2	1.4%	0%	CRHBP, GNG11, SOST, TFPI2, AC007319.1	None

768
769 Figure S8: Violin plots representing the of number of genes in each cluster for Drop-seq (top)
770 and DroNc-seq (bottom).

771
772 Figure S9: Cell differentiation trajectories constructed from Drop-seq (left), and DroNc-seq (right) using Monocle.
773 Each differentiation time-point sampled is labelled by the same color in both techniques. A, B) uses the time-point as
774 color, and C, D) shows the inferred pseudo-time as the color.
775

776
777 Figure S10: Top 10 upregulated genes identified in each cell type cluster using DroNc-seq on primary tissue from
778 archived adult human heart.
779

Figure S11: Differential expression analysis across time-points, cell-lines (biological replicates), and across Drop-seq and DroNc-seq using different thresholds for log-fold-change. All genes shown have adjusted p-value < 0.05.

References:

1. Rozenblatt-Rosen, O., Stubbington, M. J. T., Regev, A. & Teichmann, S. A. The Human Cell Atlas: From vision to reality. *Nature* (2017). doi:10.1038/550451a
2. Jaitin, D. A. *et al.* Massively parallel single-cell RNA-seq for marker-free decomposition of tissues into cell types. *Science (80-.).* (2014). doi:10.1126/science.1247651
3. Shalek, A. K. *et al.* Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells. *Nature* (2013). doi:10.1038/nature12172
4. Treutlein, B. *et al.* Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq. *Nature* (2014). doi:10.1038/nature13173
5. Macosko, E. Z. *et al.* Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161**, 1202–1214 (2015).
6. Klein, A. M. *et al.* Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic Stem Cells Accession Numbers GSE65525 Klein et al Resource Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic Stem Cells. *Cell* (2015). doi:10.1016/j.cell.2015.04.044
7. Poran, A. *et al.* Single-cell RNA sequencing reveals a signature of sexual commitment in malaria parasites. *Nature* (2017). doi:10.1038/nature24280
8. Karaiskos, N. *et al.* The Drosophila embryo at single-cell transcriptome resolution. *Science (80-.).* (2017). doi:10.1126/science.aan3235
9. Habib, N. *et al.* Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nat. Methods* **14**, 955–958 (2017).
10. Lake, B. B. *et al.* A comparative strategy for single-nucleus and single-cell transcriptomes confirms accuracy in predicted cell-type expression from nuclear RNA. *Sci. Rep.* (2017). doi:10.1038/s41598-017-04426-w
11. Bakken, T. E. *et al.* Single-nucleus and single-cell transcriptomes compared in matched cortical cell types. *PLoS One* (2018). doi:10.1371/journal.pone.0209648

811   12.   Wu, H., Kirita, Y., Donnelly, E. L. & Humphreys, B. D. Advantages of Single-Nucleus over Single-Cell
812   RNA Sequencing of Adult Kidney: Rare Cell Types and Novel Cell States Revealed in Fibrosis. *J. Am. Soc.*
813   *Nephrol.* (2018). doi:10.1681/asn.2018090912

814   13.   Banovich, N. E. *et al.* Impact of regulatory variation across human iPSCs and differentiated cells. *Genome*
815   *Res.* **28**, 1243–1252 (2017).

816   14.   La Manno, G. *et al.* RNA velocity of single cells. *Nature* (2018). doi:10.1038/s41586-018-0414-6

817   15.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data
818   across different conditions, technologies, and species. *Nat. Biotechnol.* (2018). doi:10.1038/nbt.4096

819   16.   Friedman, C. E. *et al.* Single-Cell Transcriptomic Analysis of Cardiac Differentiation from Human PSCs
820   Reveals HOPX-Dependent Cardiomyocyte Maturation. *Cell Stem Cell* (2018).
821   doi:10.1016/j.stem.2018.09.009

822   17.   Pavlovic, B. J., Blake, L. E., Roux, J., Chavarria, C. & Gilad, Y. A Comparative Assessment of Human and
823   Chimpanzee iPSC-derived Cardiomyocytes with Primary Heart Tissues. *Sci. Rep.* (2018).
824   doi:10.1038/s41598-018-33478-9

825   18.   Setty, M. *et al.* Wishbone identifies bifurcating developmental trajectories from single-cell data. *Nat.*
826   *Biotechnol.* (2016). doi:10.1038/nbt.3569

827   19.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by pseudotemporal
828   ordering of single cells. *Nat. Biotechnol.* (2014). doi:10.1038/nbt.2859

829   20.   Romero, I. G. *et al.* A panel of induced pluripotent stem cells from chimpanzees: A resource for
830   comparative functional genomics. *Elife* (2015). doi:10.7554/eLife.07103.001

831   21.   Köster, J. & Rahmann, S. Snakemake-a scalable bioinformatics workflow engine. *Bioinformatics* (2012).
832   doi:10.1093/bioinformatics/bts480

833   22.   Andrews, S. & Babraham Bioinformatics. FastQC: A quality control tool for high throughput sequence data.
834   *Manual* (2010). doi:citeulike-article-id:11583827

835   23.   Smith, T., Heger, A. & Sudbery, I. UMI-tools: Modeling sequencing errors in Unique Molecular Identifiers
836   to improve quantification accuracy. *Genome Res.* (2017). doi:10.1101/gr.209601.116

837   24.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*
838   (2011). doi:10.14806/ej.17.1.200

839   25.   Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* (2013).
840   doi:10.1093/bioinformatics/bts635

841   26.   Liao, Y., Smyth, G. K. & Shi, W. FeatureCounts: An efficient general purpose program for assigning
842   sequence reads to genomic features. *Bioinformatics* (2014). doi:10.1093/bioinformatics/btt656

843   27.   Bailey, T. L. *et al.* MEME Suite: Tools for motif discovery and searching. *Nucleic Acids Res.* (2009).
844   doi:10.1093/nar/gkp335

845   28.   McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold Approximation and
846   Projection. *J. Open Source Softw.* (2018). doi:10.21105/joss.00861

847   29.   Etienne, B. *et al.* Evaluation of UMAP as an alternative to t-SNE for single-cell data. *Development* (2018).
848   doi:10.1101/298430

849   30.   Soneson, C. & Robinson, M. D. Bias, robustness and scalability in single-cell differential expression
850   analysis. *Nat. Methods* (2018). doi:10.1038/nmeth.4612

851