1 Xrn1 activity broadly represses RNA polymerase II occupancy at mammalian but not viral
2 promoters during herpesvirus infection
3

4 Ella N Hartenian[1], Britt A Glaunsinger[1,2,3,*]
5
6 [1]Department of Molecular and Cell Biology, University of California Berkeley, CA, USA
7 [2]Department of Plant and Microbial Biology, University of California Berkeley, CA, USA
8 [3]Howard Hughes Medical Institute, University of California Berkeley, CA, USA
9
10 *Address correspondence to:
11 Britt Glaunsinger, glaunsinger@berkeley.edu

12
13
14
15

Abstract

In mammalian cells, widespread acceleration of cytoplasmic mRNA degradation is linked to

impaired RNA polymerase II (Pol II) transcription. This mRNA decay-induced transcriptional

repression occurs during infection with gammaherpesviruses including Kaposi's sarcoma-

associated herpesvirus (KSHV) and murine gammaherpesvirus 68 (MHV68), which encode an

mRNA endonuclease that initiates widespread RNA decay. Here, we show that MHV68-induced

mRNA decay leads to a genome-wide reduction of Pol II occupancy at mammalian promoters,

which requires the cellular 5'-3' exonuclease Xrn1. Viral genes, despite the fact that they require

Pol II for transcription, escape this transcriptional repression. Protection is not governed by viral

promoter sequences; instead, location on the viral genome is both necessary and sufficient to

escape the transcriptional repression effects of mRNA decay. We hypothesize that the ability to

escape from transcriptional repression is linked to the conformation of viral DNA in replication

compartments, providing a means for these viruses to counteract decay-induced viral transcript

loss.

Introduction

Regulating messenger RNA (mRNA) abundance is of central importance during both

cellular homeostasis and disease. While it is intuitive that transcriptional changes in the nucleus

impact RNA levels in the cytoplasm, evidence in both yeast and mammalian cells now indicates

that the converse is also true: altered cytoplasmic mRNA decay rates can broadly impact

transcription by RNA polymerase II (Pol II). In yeast, stabilizing the cytoplasmic mRNA pool by

removing mRNA decay factors such as Xrn1 causes a compensatory decrease in Pol II

transcription (Sun et al. 2012; Haimovich et al. 2013; Shalem et al. 2008; Shalem et al. 2011;

Dori-Bachash et al. 2012)Furthermore, reducing or slowing Pol II transcription leads to an

increase in overall mRNA stability (Sun et al. 2012) supporting a "buffering" model in which

yeast compensate for widespread perturbations to mRNA abundance by alternatively changing

rates of decay or transcription.

Broad changes in mRNA abundance are often triggered in mammalian cells during viral

infection. Numerous viruses, including alphaherpesviruses, gammaherpesviruses, SARS

coronavirus and influenza A virus drive accelerated mRNA decay during infection by expressing

mRNA specific endoribonucleases and/or activating host nucleases. This decay contributes to

viral immune evasion, increases the availability of translation machinery, and facilitates temporal

viral gene regulation (Richner et al. 2011; Liu et al. 2013; Parrish & Moss 2006; Shiflett & Read

2013; Dauber et al. 2011). Viral endonucleases target host and viral mRNAs for cleavage,

whereupon cellular exonucleases degrade the resulting mRNA fragments (Karr & Read 1999;

Glaunsinger et al. 2005; Huang et al. 2011; Kamitani et al. 2009; Jagger et al. 2012). This

strategy accelerates basal RNA decay by circumventing the typically rate limiting steps of

deadenylation and decapping (Covarrubias et al. 2011; Gaglia et al. 2012; Eckmann et al. 2010;

Gallouzi & Wilusz 2013; Schoenberg & Maquat 2012).

Although yeast and mammals share key proteins involved in cytoplasmic mRNA decay

and transcription, studies with RNA-decay inducing viruses in mammals have revealed that

unlike the yeast pathway, accelerated mRNA decay leads to an extensive shutoff of cellular gene

expression (Abernathy & Glaunsinger 2015; Gilbertson et al., 2018). Infection with the

gammaherpesviruses KSHV or MHV68 causes reduced Pol II recruitment at host promoters in a

manner dependent on mRNA cleavage by the viral endonucleases SOX or muSOX, respectively.

Thus, accelerated mRNA decay is not broadly counteracted by increased transcription-based

repopulation, but instead leads to a more extensive shutdown of cellular gene expression.

However, similar to yeast, the mammalian connection between decay and transcription requires

the activity of cellular exonucleases such as Xrn1, which degrade the viral endonuclease

cleavage products (Abernathy et al. 2015; Covarrubias et al. 2011). Pol II repression requires

exonucleolytic degradation of the mRNA cleavage fragments, as the initial endonucleolytic

mRNA cleavage without further fragment degradation is insufficient to activate this

signaling(Abernathy et al. 2015). Accelerated Xrn1-mediated degradation causes trafficking of

released RNA binding proteins (RBP) from the cytoplasm to the nucleus, which may provide a

means for the cell to detect and respond to increased RNA turnover (Gilbertson et al., 2018).

During MHV68 infection, in addition to host mRNA, cytoplasmic viral mRNA is

susceptible to cleavage by muSOX (Abernathy et al. 2014). However, measurements of nascent

transcript production from a subset of viral promoters suggested that viral genes are robustly

transcribed during the stage of infection when host transcription is reduced (Abernathy et al.

2015). A major outstanding question is how these viral promoters escape mRNA decay-induced

transcriptional repression despite being transcribed by RNA Pol II. Here, we measured Pol II

occupancy across the mouse and MHV68 genomes in infected cells to more comprehensively

define how accelerated mRNA decay impacts polymerase occupancy. Our data demonstrate a

genome-wide reduction in Pol II recruitment at host promoters under conditions of increased

mRNA decay, and underscore a key role for Xrn1 in this process. In contrast, Pol II occupancy

of the viral genome appears broadly resistant to the effects of mRNA degradation. This

protection is not conferred by viral promoter sequences; instead, location on the replicating viral

genome is both necessary and sufficient to escape the transcriptional effects of mRNA decay.

We propose a model in which DNA amplified in viral replication compartments, unlike the

cellular chromatin, is immune to transcriptional repression during accelerated cytoplasmic

mRNA decay. Thus, while both viral and cellular mRNA pools are susceptible to cytoplasmic

degradation, transcriptional repopulation selectively counteracts this loss for viral genes.

Results

Accelerated RNA decay broadly reduces Pol II occupancy

We previously reported that Pol II occupancy at several individual mammalian promoters

was significantly reduced during accelerated cytoplasmic mRNA decay(Abernathy et al. 2015;

Gilbertson et al., 2018.). To more comprehensively assess the extent of mRNA decay-induced

transcriptional repression, we evaluated global Pol II occupancy by chromatin

immunoprecipitation and deep sequencing (ChIP seq) in mock versus MHV68 infected MC57G

mouse fibroblasts at 24 hours post infection (hpi). As a control, we also infected cells with a

version of MHV68 containing the point mutation R443I in the viral muSOX endonuclease gene,

which reduces its mRNA cleavage activity (Richner et al. 2011). MHV68-infected cells

displayed less promoter-proximal Pol II at 86 percent of loci compared to mock-infected cells,

averaged across two biological replicates (Fig 1A, Suppl Fig 1A). Loss of Pol II occupancy was

primarily due to muSOX-induced mRNA decay, as promoter-proximal Pol ll recovered at 67

percent of these loci during infection with R443I MHV68 (Fig 1A, Supp Fig1A). Sixty-three

genes, or 1.3 percent of loci, showed higher Pol II occupancy in WT infection as compared to

mock infection, although these did not share any common GO terms (Supp Table 1). We

independently validated the ChIPseq results by ChIP-qPCR at the *Rplp0* and *Fus* promoters, both

of which showed a significant reduction of Pol II occupancy in cells infected with WT but not

R443I MHV68 (Fig. 1B). The genome-wide reduction in poised Pol II in WT but not R443I

infected cells was also evident when the sequencing reads from positions -2000 to +4000 around

the transcription start sites (TSS) were plotted as a histogram (Fig 1C, Supp Fig 1B). The

variability between replicates in R443I recovery is likely due to differences in the extent of host

shutoff reduction between replicates (Supp Figure 1C).

To assess the stage of transcription impacted by RNA decay, we compared the amount of

Pol II in promoter regions (-50 to +200) and the amount of Pol II in gene bodies (+200 to +5000)

between mock and infected samples. Consistent with the above analyses, MHV68 infected cells

contained markedly less promoter proximal Pol II than uninfected cells, and this signal was

partially recovered in the R443I infection (Fig 1D). Genes with impaired Pol II promoter binding

during MHV68 infection also had less Pol II within the gene body compared to mock infected

cells, which again was not observed during R443I infection (Fig 1D). Notably, the difference in

regression line slopes between MHV68 and mock infected cells was clearly more marked when

comparing Pol II promoter occupancy than gene body occupancy, suggesting that the primary

transcriptional defect is at the stage of initiation. This is consistent with data from previous

studies that found no defect in serine 2 phosphorylation of elongating Pol II and no consistent

change to Pol II elongation during MHV68 infection (Abernathy et al. 2015; Gilbertson et al.

2018.).

Finally, to assess if RNA decay affected transcription termination, we binned sequencing

reads between 1kb upstream and 1kb downstream of the transcription termination site (TTS) (Fig

1E, Supp Fig 1D). TTS-proximal Pol II shows no defect in termination in either infection

condition, suggesting Pol II removal from host transcripts is not impacted by MHV68 infection

or by RNA decay.

Pol II Loss is dependent on Xrn1

Our previous work implicated the catalytic activity of the host 5'-3' exoribonuclease

Xrn1 as necessary for the RNA-decay dependent loss of Pol II at several individual host genes

(Abernathy et al. 2015). To determine the extent to which mRNA decay-induced Pol II

repression is dependent on Xrn1, we measured Pol II occupancy by ChIPseq in mock and

MHV68 infected cells upon depletion of Xrn1 using siRNA (Fig 2A). Remarkably, knockdown

of Xrn1 resulted in a near complete recovery of Pol II promoter occupancy in MC57G cells

infected with WT MHV68 (Fig 2B, Supp Fig 2A). We validated these findings using

independent biological replicates at *Rplp0* and *Fus* by ChIP qPCR (Fig 2C). On a gene-by-gene

basis, we observed similar levels of Pol II at promoters in mock and infected cells depleted of

Xrn1 (Fig 2D, Supp Fig 2B). Fig 2E shows representative Pol II occupancy of *Srsf2* during

knockdown conditions. Similarly, comparing the amount of Pol II in gene bodies and at TTS of

Xrn1-depleted cells (as described in Fig 1) showed little difference between mock and MHV68

infected conditions (Figs 2F, 2G, Supp Fig 2C). Together, these data highlight the essentiality of

Xrn1 for the genome-wide repression of Pol II occupancy during virus-induced mRNA decay.

Pol II recruitment to viral genes is not affected by RNA decay

Herpesviral genes are transcribed in the nucleus by mammalian Pol II, and thus could be

similarly subjected to the mRNA decay-induced transcriptional repression observed across the

host genome. We therefore analyzed Pol II occupancy of the viral genome in WT MHV68 and

R443I infected cells to determine how viral genes respond to RNA decay. The average Pol II

signature across 82 viral ORFs between -500 and +500 basepairs (bp) around the TSS (O'Grady

et al., manuscript in preparation) was comparable in cells infected with WT or R443I MHV68,

with a slightly higher signal in the R443I infection (Fig 3A). Indeed, there was a robust and

widespread Pol II signal in both infection conditions across the viral genome (Fig 3B). Thus,

herpesviral genes appear to escape mRNA decay-induced transcriptional repression.

In contrast to host genes, we did not observe peaks of promoter-proximal Pol II on the

viral genome in our ChIPseq data (see Fig 3A, B). Pol II ChIP qPCR across various regions of

ORF54 and ORF37 also showed a similar Pol II signal throughout each gene body and at their

promoter. We did note a decrease in signal near the TTS of ORF54, however this was true for

both WT and R443I infections (Fig 3C). We hypothesize that pervasive transcription of the viral

genome at 24 hpi, compounded by the presence of TSSs on both Watson and Crick strands,

likely contributes to this phenotype.

PABPC is not excluded from viral replication compartments

We next considered how the viral genome maintains high levels of Pol II occupancy

during accelerated mRNA decay. During replication, viral DNA is transcribed within viral

replication compartments (RCs) in the nucleus, which exclude host DNA and thus stain poorly

with DAPI (Chang et al. n.d.; Taylor et al. 2003; Monier et al. 2000). Although they are not

membrane bound, these compartments selectively enrich for factors required for viral replication

(such as the viral processivity factor ORF59) and gene expression (such as Pol II) (Li et al. 2018;

Dembowski & DeLuca 2015; Rice et al. 1994). Virus-induced mRNA decay causes increased

trafficking of cytoplasmic poly(A) binding protein (PABPC) from the cytoplasm to the nucleus,

which we have recently linked to transcriptional repression (Kumar et al. 2011; Gilbertson et al.

2018). To determine whether replication compartments selectively exclude nuclear PABPC as a

means of avoiding transcriptional repression, we monitored its localization in cells lytically

infected with MHV68 or KSHV. As expected, the majority of infected cells displayed increased

nuclear PABPC compared to unreactivated or uninfected cells as measured by

immunofluorescence (IF). We further saw that PABPC was not excluded from replication

compartments (Fig. 4A, B, Supp Fig 3A). To determine PABPC localization relative to RCs, we

first identified cells with RCs by looking for regions of Pol II or ORF59 staining that excluded

DAPI. We then quantified nuclear PABPC pixel intensity, and counted cells as having an

overlapping RC/PABPC signal when the PABPC pixel intensity was at least 2x that of cells

without nuclear PABPC from the same image. Indeed, 85 percent of MHV68 infected cells

showed overlapping PABPC/Pol II signal and 90 percent of reactivated iSLKs showed an

overlapping PABPC/ORF59 signal (Fig 4C). Thus, PABPC exclusion from RCs is unlikely to

underlie viral promoter escape.

Viral promoters fail to escape transcriptional repression outside of the viral genome

We then tested whether viral promoter sequence elements were sufficient to enable

escape by querying whether they remained immune to transcriptional repression upon relocation

from the viral genome into the host DNA. We inserted the MHV68 late promoter M7 or the

KSHV LANA promoter upstream of a puromycin resistance cassette into 293T cell chromatin

using either lentiviral transduction or random plasmid DNA integration, respectively. We

induced mRNA decay and host transcriptional repression in these cells by expression of MHV68

muSOX or the herpes simplex virus type 1 (HSV-1) endonuclease vhs. Similar to the host

promoters *Gapdh* and *Rplp0*, Pol II recruitment to both integrated viral promoters was repressed

upon induction of mRNA decay (Fig 5A, 5B).

To confirm that viral promoters integrated into the host genome were also repressed in

the context of infection, we infected MC57G cells containing an integrated MHV68 M4

promoter-driven puromycin cassette with MHV68. Indeed, the M4 promoter was repressed to a

similar degree as host *Fus* and *Rplp0* promoters during infection with WT MHV68 but not the

mRNA decay-deficient MHV68 R443I (Fig 5C). In summary, these data argue against the

presence of protective elements within viral promoters, and instead suggest that the chromatin

context or location on the viral genome underlies escape from mRNA decay-induced

transcriptional repression.

Non-MHV68 promoters escape repression on the viral genome

A prediction from the above results is that non-MHV68 promoters would gain protection

from transcriptional repression when present on the replicating viral genome. To test this, we

measured the Pol II occupancy and transcriptional potential of the cytomegalovirus (CMV)

promoter, which drives green fluorescent protein (GFP) expression on the MHV68 genome.

Unlike alpha- and gammaherpesviruses, the betaherpesvirus CMV does not accelerate mRNA

decay. Furthermore, in either a plasmid context or when integrated into 293T cells, the CMV

promoter is strongly repressed during muSOX-induced mRNA decay (Fig 6A, B). However,

placed in the context of the replicating MHV68 genome, the CMV-GFP promoter had a Pol II

ChIPseq signal during both MHV68 WT and R443I infection that was not markedly different

from several other representative viral genes (ORFs 4, 25, 48, 52, 72, M11) (Fig 6C).

Finally, we evaluated whether lytically replicating viral DNA was necessary to confer

escape from transcriptional repression, or whether some other feature of the non-replicating viral

genome was important. To this end, we examined Pol II occupancy at promoters of several genes

expressed from the KSHV genome during latent infection in 293T cells, when the viral DNA is

maintained as an episome but not amplified in replication compartments. Latent infection does

not promote mRNA turnover, and thus we transfected muSOX into these cells to stimulate

mRNA turnover in the absence of lytic replication. Similar to the host genes *Gapdh* and *Rplp0*,

we observed an RNA decay-induced repression of Pol II occupancy at the latent viral promoter

LANA and the viral K1 promoter was repressed in 4 out of 5 experiments (Fig 6D). We also

observed repression of the human *Ef1α* promoter, which is present on the Bac16 KSHV viral

episome as a driver of the GFP reporter gene (Fig 6D). In sum, these observations indicate that

the viral genome per se does not confer protection from transcriptional repression; instead,

features linked to lytic genome amplification in replication compartments facilitate robust Pol II

recruitment under conditions of accelerated mRNA turnover.

Discussion

Here, we demonstrate that virus-induced mRNA decay broadly decreases Pol II recruitment to mammalian promoters, extending prior observations made with individual cellular genes to the genomic scale. Although the MHV68 endonuclease muSOX cleaves the majority of cytoplasmic mRNA, we found that the cellular 5'-3' exonuclease Xrn1, which degrades these cleaved fragments, is essential for inhibiting Pol II recruitment across the mammalian genome. In contrast, promoters on the replicating viral genome recruit Pol II with similar efficiency during normal or accelerated mRNA decay, indicating that they escape transcriptional inhibition. This protection is lost upon relocation of the viral promoters into the host genome or when the viral genome is in a latent state, emphasizing the importance of DNA context and replication state, rather than promoter sequence, in determining the transcriptional response to accelerated mRNA decay.

Across the host genome, Pol II occupancy appears to be most strongly affected at the stage of recruitment. The reduction in promoter-proximal Pol II is propagated into the gene body, although the differences there were less pronounced, suggesting that once transcription of a gene initiates there are not subsequent blocks to elongation. This agrees with prior studies of individual host loci in MHV68 infected or muSOX expressing cells (Abernathy et al. 2015; Gilbertson et al. 2018). We also did not observe a defect in transcription termination, or evidence of polymerases on the DNA after TSSs, suggesting that the gammaherpesvirus MHV68 does not induce the termination defect and run-on transcription of host genes recently described in HSV-1 (Rutkowski et al. 2015).

Two pieces of evidence indicate that reduction of Pol II at host genes is unlikely due to sequestration of or competition for the transcription machinery by the viral DNA. First, infection with the mRNA decay mutant virus R443I, which replicates its viral genome to WT levels (Richner et al. 2011), does not induce the widespread reduction in Pol II recruitment to host promoters. Second, transfection of muSOX alone (or other broad acting viral endonucleases) in the absence of viral infection is sufficient to induce transcriptional repression at host promoters. Therefore, we instead favor the hypothesis that one or more cellular proteins released from degrading mRNA fragments triggers a signaling event that restricts transcription machinery occupancy of mammalian chromatin (Gilbertson et al. 2018).

Reduced Pol II occupancy of the host genome during MHV68 infection parallels

infection data from the alphaherpesvirus HSV-1. HSV-1 infection results in a genome-wide

reduction in Pol II occupancy of host chromatin and reduced nascent RNA production of the

majority of host transcripts (Rutkowski et al. 2015; Abrisch et al. 2016; Spencer et al. 1997).

Furthermore, early data showed that while the endogenous mouse β-globin locus was

transcriptionally repressed during HSV-1 infection, its expression was rescued upon integration

of the β-globin gene into the HSV-1 genome (Smiley et al. 1987; Smibert & Smiley 1990).

Experiments with gene-specific null viruses have implicated immediate early (IE) proteins in the

HSV-1 transcriptional repression phenotype, including ICP27, which can accelerate mRNA

turnover (Spencer et al. 1997; Smiley 2004). Additionally, we have previously published, and

confirm in this report, that transfection of HSV-1 vhs, another broad acting endoribonuclease, is

sufficient to induce reduced Pol II occupancy of several host promoters (Abernathy et al.

2015).Thus, in addition to targeting host and viral mRNAs for cytoplasmic decay, both alpha and

gammaherpesviruses restrict Pol II occupancy of the host genome, although additional work is

needed to establish whether the underlying mechanism is similar.

During the lytic stage of herpesvirus infection, viral DNA is localized to replication

compartments. These non-membrane bound structures are sites of viral genome replication,

transcription, and packaging (Schmid et al. 2014).They are enriched for many host and viral

proteins involved in these processes, including chromatin modifying factors and Pol II

(Dembowski & DeLuca 2015; Rice et al. 1994; Li et al. 2018). Replication compartments grow

and coalesce over infection, excluding host chromatin, which becomes pushed to the nuclear

periphery (Chang et al. n.d.; Taylor et al. 2003; Monier et al. 2000). We recently showed that

muSOX and Xrn1-coordinated mRNA decay causes differential trafficking of many cellular

RNA binding proteins between the cytoplasm and nucleus (Gilbertson et al. 2018). Knockdown

and overexpression studies suggested that at least one of these proteins, PABPC, is associated

with transcriptional repression. Here, we detected PABPC in viral replication compartments of

infected cell nuclei, indicating that selective exclusion of PABPC from these compartments is

not the mechanism by which the viral genome escapes repression. Instead, we hypothesize that

either the state of viral DNA or other protective features of the replication compartment confers

escape from transcriptional repression.

Protection from mRNA decay-induced transcriptional repression was not conferred to

host or viral promoters present on the latent viral episome, indicating that lytic phase viral DNA

replication or replication compartment formation is key to this phenotype. The latent genomes of

gammaherpesviruses including KSHV and Epstein Bar Virus (EBV) are heavily methylated and

histone-rich, with the exception of promoters of genes expressed during latency, the latent origin

of replication, and CTCF binding sites (Günther & Grundhoff 2010; Szyf et al. 1985; Hilton et

al. 2013). Notably, during latency key lytic promoters like RTA contain both the activating

H3K4me3 and the repressive H3K27me3 histone marks, representing a 'poised' state (Toth et al.

2010). Rapid changes shortly after lytic cycle induction result in hypomethylation and removal

of repressive marks at early gene promoters(Günther & Grundhoff 2010; Toth et al. 2010) .At

later stages of infection, KSHV, EBV and HSV-1 genomes have been shown to be largely

nucleosome free, consistent with histones not being packaged in virions (Dembowski & DeLuca

2015; Monier et al. 2000; Hollingworth et al. 2017). For gammaherpesviruses, this is the stage at

which mRNA degradation is most pronounced. We therefore hypothesize that the 'open' state of

viral DNA during lytic replication plays an important role in facilitating Pol II recruitment under

conditions of accelerated cytoplasmic mRNA decay. Future work is geared towards exploring

whether widespread mRNA decay alters cellular chromatin states and DNA accessibility and, if

so, how this influences transcription factor recruitment.

In conclusion, decay-coupled transcriptional repression is a novel facet of viral

manipulation of the cellular gene expression landscape. We hypothesize that the ability to escape

from transcriptional repression is linked to the open conformation of viral DNA in replication

compartments, and that this helps these viruses counteract decay-induced viral transcript loss.

While widespread mRNA decay has been shown to facilitate immune evasion, gene expression

control, and latency establishment, the robust viral transcription we document here presents a

parsimonious explanation for a novel way in which the virus benefits from accelerated decay.

Materials and Methods

Plasmids and Primers

pCDEF3 muSOX and VHS and SOX have been previously published(Abernathy et al. 2015).

pCDEF3-GFP was used as a control.

Primers used for qPCR and for cloning are listed in Supp Table 2. M7 and M4-puro were

made by cloning the respective promoter sequence upstream of the puromycin cassette in a

pLKO-sgRNA vector. The promoter of M4 was defined as the 1131bp upstream of the M4

coding region not overlapping with other viral ORFs and the promoter sequence of M7 was

previously published (Wong-Ho et al. 2014). pGL3-kLANA-puromycin was made by

introducing the puromycin gene into the I-CeuI and XbaI sites of pGL3-kLANA via InFusion

(Clonetech) cloning. pGL3-kLANA was made by amplifying KSHV's LANA promoter

sequence (Jeong et al. 2001)and using InFusion cloning to introduce it into the XhoI and HindIII

sites of pGL3. pGL3-CMV was similarly made by introducing the amplified CMV promoter into

the same sites of pGL3.

Cells and transfections

MC57G mouse fibroblast cells (ATCC), 293T cells (ATCC) and NIH 3T3 cells (ATCC)

were maintained in DMEM (Invitrogen) supplemented with 10% fetal bovine serum (FBS).

iSLK BAC16 cells (Brulois et al. 2012) were maintained in DMEM supplemented with 10%

FBS and 1mg/mL hygromycin B. To reactivate iSLK cells, they were treated with 1ug/mL

doxycycline and 1mM sodium butyrate for 48h.

DNA transfections were carried out in 293T cells at 70-90% confluency in 10 cm plates

with 10 µg host-shutoff factor plasmid or vector control using PolyJet (SignaGen) 24h before

harvesting and again at 18h before harvesting.

Lentiviral transduction was carried out by spinfecting 1×10^6 293T or MC57G cells with

lentivirus made from 2^{nd} generation plasmids at a MOI <1 for 2h with 4ug/mL polybrene

(Fisher). Twenty-four hours later puromycin (1 µg/ml for 293T or 3 µg/ml for MC57G) was

added to select for transduced cells. 293T cells stably expressing the kLANA-puromycin

cassette were made by transfecting the cells with 1 µg pGL3-kLANA-puromycin using PolyJet,

then selecting with puromycin for random integration events.

For the CMV-luciferase promoter assay, populations of 293T cells expressing muSOX

were selected 24h post-transfection using the Miltenyi Biotec MACS cell separation system

according to the manufacturer's instructions as previously described (Gilbertson et al. 2018).

293T cells were infected with KSHV by reactivating iSLK-Bac16 cells for 48h and then

spinfecting cell-free virus onto a monolayer of 293T cells, as described for lentiviral production

but in the absence of polybrene.

3 million MC57G cells were nucleofected with a 100uL Neon tip (1400 Voltage/20

Width/2 Pulse) to knockdown Xrn1 using a pool of 4 siRNAs (Dharmacon M-046621-01-0005)

or a non-targeting control (Dharmacon D-001206-14-20) using 10uL of 20uM siRNA.

4SU labeling

Cells were incubated in suspension with 500 µM 4SU (Sigma T4509) in DMEM

supplemented with 10% FBS for 10 min then washed with PBS. RNA was extracted with TRIzol

followed by isopropanol purification. RNA (300 ug) was incubated in biotinylation buffer (50

mM HEPES [pH 7.5], 5 mM EDTA) and 5ug MTSEA-biotin (Biotium #90066), rotating at room

temperature in the dark for 30m. A no biotin control was made from an equal amount of total

RNA and incubated as above but without the addition of MTSEA-biotin. RNA was then

phenol:chloroform extracted and precipitated with isopropanol. The pellet was resuspended in

DEPC-treated water and mixed with 50 uL Dynabeads MyOne streptavidin C1 (Invitrogen) that

had been pre-washed twice with wash buffer (100 mM Tris [pH 7.5], 10 mM EDTA, 1 M NaCl,

0.1% Tween 20). Samples were rotated in the dark for 1hr at room temperature, then washed

twice with 65°C wash buffer and twice with room temperature wash buffer. Samples were eluted

twice with 100uL 5% Beta-mercaptoethanol (BME) in DEPC H_2O for 5m then RNA was

precipitated with isopropanol and quantified by RT-qPCR.

Viral mutagenesis, propagation and infections

The WT and R443I MHV68 bacterial artificial chromosome (BAC) were previously

described (Adler et al. 2000; Richner et al. 2011). MHV68 was produced by first making p0

virus by transfecting NIH 3T3 cells in 6-well plates with 2.5 µg BAC DNA using TransIT-X2

(Mirus Bio) for 24h. Five to seven days later, the cells were split into a 10 cm dish, then

harvested and frozen. After 5-7 days the majority of cells showed cytopathic effect (CPE). 30 µL

of p0 was then added per confluent 10 cm dish of NIH 3T12 cells, split 2 days later into four 10

cm dishes and harvested 4-6 days later when all cells were infected. Cells were collected by

centrifugation (5 min, 1500 rpm) and the pellet was dounced 10 times. The cell pellet and

supernatant were then ultracentrifuged for 2 h at 30,000 rpm. The resulting pellet was

resuspended and titered by plaque assay. Cells were infected with MHV68 at an MOI of 5 for

24h.

Western blotting

Cells were lysed in RIPA buffer (50 mM Tris-HCl pH7.6, 150 mM NaCl, 3 mM $MgCl_2$,

10% glycerol, 0.5% NP-40, cOmplete EDTA-free Protease Inhibitors [Roche]) and then clarified

by centrifugation at 15,000 rpm for 10 min at 4˚C. Whole cell lysate was quantified by Bradford

assay and resolved by SDS-PAGE. Antibodies for western blot are: Xrn1 (Bethyl A300-443A,

1:1000), and Gapdh (Abcam, ab8245, 1:2000).

RT-qPCR

RNA was reverse transcribed using AMV RT (Promega) with random 9-mer primers.

Total RNA was DNase treated with TURBO DNase (ThermoFisher) or DNase treated on column

with Zymo Direct-zol RNA MiniPrep Plus DNase. cDNA was quantified using iTaq Universal

SYBR Mastermix (BioRad) and transcript-specific primers. All qPCR results are normalized to

18S levels and WT or vector control set to 1.

Immunofluoresence

NIH 3T3 cells were plated on coverslips (7.5 x 10^4 cells/well of a 12-well), infected the

following day for 25-27 h, and then fixed in 4% formaldehyde for 10 min. iSLK-Bac16 cells

were plated on coverslips (5 x 10^4 cells/well of a 12-well), reactivated 24h later with doxycycline

and sodium butyrate for 48 hours and then fixed as above. Cells were permeabilized with ice-

cold methanol at -20°C for at least 20 min and incubated with anti-Pol II antibody (Biolegend,

8WG16 at 1:200), anti-PABPC antibody (Abcam ab21060 at 1:200 for mouse cells and Santa

Cruz sc32318 at 1:25 for human cells) or anti-ORF59 antibody (Advanced Biotechnologies 13-

211-100 at 1:200) in 5% BSA overnight at 4 °C. Secondary antibodies were added (1:1000) for 1

h at 37 °C. Coverslips were mounted in DAPI- containing Vectashield (VectorLabs). We

identified RCs by DAPI exclusion and Pol II recruitment or ORF59 staining (Rice et al. 1994;

Dai-Ju et al. 2006; Li et al. 2018). Images were collected on a Zeiss LSM 710 AxioObserver

with a 40x oil objective. Cells with RCs were first identified, then PABPC co-localization was

determined by counting cells with at least 2x nuclear pixel intensity relative to non-RC

containing cells in the same image (ImageJ).

Chromatin immunoprecipitation (ChIP)

ChIP was performed from 15 cm plates of cells. 3 x 10^6 MC57G or 293T cells were

plated 24 h before infection and then harvested 24 h after infection or transfection(s). Upon

harvesting, cells were washed with PBS, crosslinked in 1% formaldehyde at room temperature

for 10 min. Cells were quenched in 0.125 M glycine for 5 min and washed twice with cold PBS.

Crosslinked cell pellets were mixed with 1 mL fractionation buffer (5mM PIPES pH 8.0, 85mM

KCl, 0.5% NP-40 with cOmplete EDTA-free Protease Inhibitors [Roche]) and incubated on ice

for 10 min, during which time the lysate was dounce homogenized to release nuclei and spun at

4,000 rpm for 5 min at 4°C to pellet nuclei. Nuclei were then resuspended in 300 ul of nuclei

lysis buffer (50mM Tris-HCl pH 8.0, 0.3% SDS, 10mM EDTA, with cOmplete EDTA-free

Protease Inhibitors [Roche]) and rotated for 10 min at 4°C followed by sonication using a

QSonica Ultrasonicator. Sonicated chromatin was then spun at 13,000 rpm for 10 min at 4°C and

the pellet discarded. 40ug of chromatin was brought to 500uL with ChIP dilution buffer (16.7

mM Tris-HCl pH 8.0, 1.1% Triton X-100, 1.2 mM EDTA, 167 mM NaCl and cOmplete EDTA-

free Protease Inhibitors [Roche]) and incubated with 10 ug mouse monoclonal anti-RNAPII

(BioLegend, 8WG16) or rabbit IgG (Fisher Scientific) overnight, whereupon samples were

rotated with 20 ul protein G dynabeads (Thermofisher) for 2 h at 4°C. Beads were washed with

low salt immune complex (20 mM Tris pH 8.0, 1% Triton-x-100, 2 mM EDTA, 150 mM NaCl,

0.1% SDS), high salt immune complex (20 mM Tris pH 8.0, 1% Triton-x-100, 2 mM EDTA,

500 mM NaCl, 0.1% SDS), lithium chloride immune complex (10mM Tris pH 8.0, 0.25 M LiCl,

1% NP-40, 1% Deoxycholic acid, 1 mM EDTA), and Tris-EDTA for 5 min each at 4°C with

rotation. DNA was eluted from the beads using 100 ul of elution buffer (150 mM NaCl, 50 ug/ml

Proteinase K) and incubated at 55°C for 2h, then 65°C for 12h. DNA was purified using a Zymo

Oligo Clean & Concentrator kit, and quantified by qPCR using primers to the promoter regions

of indicated genes for 50 cycles. qPCR was simultaneously performed on input chromatin and

normalized to input.

ChIP-Sequencing and data analysis

446   Libraries were prepared for sequencing (Kapa Hyper Prep) and sequenced on a

447   HiSeq4000 with 100bp single end reads or 150bp reads (Xrn1kd second replicate only).

448   Sequencing quality was assessed with FastQC and trimmed with Sickle. A custom index was

449   made of the mm10 and MHV68 genomes with Bowtie2 build. Reads were mapped to that index

450   using Bowtie2 (2.3.0).

451   To visualize reads in the Integrative Genomics Viewer (IGV), tdf files were made from

bam files and indexes made from those bam files. To make tdf files for the viral genome, a

chrom.size file was created for MHV68 and then the count command was run. Analysis of

mapped reads was performed with HOMER (Hypergeometric Optimization of Motif

EnRichment) (Heinz et al. 2010). Tag directories were created from the bam files generated from

Bowtie with the makeTagDirectory command. This was done separately for the mouse and

MHV68 genomes. For mouse genes, the analyzeRepeats command was used to look at tag

density in the body of genes and the –c pausing option was used to compare pausing ratios.

annotatePeaks was run in both the –hist and –ghist mode to assess composite Pol II occupancy

across the mouse genome as well as individual gene occupancy.

461   Viral analyses were done by first creating a custom genome annotation in HOMER using

the loadGenome command. The TSS file was created from data generously shared by Scott

Tibbetts. annotatePeaks was run in the –hist mode to assess composite Pol II occupancy across

the viral genome as well as individual gene occupancy.

Acknowledgements

We thank all members of the Glaunsinger and Coscoy labs for their helpful suggestions and

discussions. We thank David McSwiggen for his microscopy assistance, Jennifer Kugel for her

guidance applying HOMER to the viral genome and Scott Tibbetts for sharing MHV68 TSS

data. This work used the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley,

supported by NIH S10 OD018174 Instrumentation Grant. Confocal imaging experiments

were conducted at the CRL Molecular Imaging Center on the Zeiss LSM 710, supported by the

Gordon and Betty Moore Foundation. We would like to thank Holly Aaron and Feather Ives for

their microscopy training and assistance. B.G. is an investigator of the Howard Hughes Medical

Institute, and is further supported by NIH R01 CA136367 for this work.

Figure Legends

Fig 1: Promoter-proximal Pol II recruitment to mammalian genes is RNA-decay dependent

(A) Pol II ChIP-seq signal profiles of host genes in mock-infected, MHV68-infected and R443I-infected cells. Each row of the heat map displays Pol II occupancy of one gene from -1000 to +1000 in 25bp bins. Genes are ranked by the Pol II-transcription start site (TSS) proximal signal in mock infected cells. (B) ChIP-qPCR validation of Pol II occupancy at the *Rplp0* and *Fus* promoters. Pol II ChIP was performed on mock, MHV68 WT or MHV68 R443I cells and Pol II levels were assayed near the TSS of two repressed host genes during MHV68 infection from the ChIP-seq data. (* $p < 0.05$, ** $p < 0.01$, students t-test) (C) Sequence tags were binned into 25-bp bins for -2000 to +4000 around the TSS. Mock (red), MHV68 (blue) and MHV68 R443I (green) traces are shown along with their input controls. (D) Scatterplot of Pol II occupancy of promoters and gene bodies, averaged across two replicate experiments with inputs subtracted out. Mock is compared to WT MHV68 infection on the left and to MHV68 R443I infection on the right. Linear regression lines are plotted in blue and equations are provided. A y=x line is plotted for reference in red. (E) Pol II transcription termination is not dependent on RNA decay. Sequence tags were binned into 25-bp bins for transcription termination sequence (TTS) proximal Pol II for -1000 to +1000 around the TTS with the same color scheme as (C).

Fig 2: Host promoter repression requires Xrn1

(A) Representative western blot showing the efficiency of Xrn1 knockdown (kd) in MC57G cells, with Gapdh serving as a loading control. (B) Sequence tags were binned into 25-bp bins for -2000 to +4000 around the transcription start site (TSS) in mock (purple) and MHV68 infected (green) Xrn1 kd samples for ChIPs and input controls. (C) ChIP-qPCR validation of Pol II occupancy at the *Rplp0* and *Fus* promoters during mock and MHV68 infection in Xrn1 kd conditions. (D) Pol II ChIP-seq profiles of host genes in mock-infected and MHV68-infected cells depleted of Xrn1. Each row of the heat map displays Pol II occupancy of one gene from -1000 to +1000 in 25 bp bins. (E) Pol II ChIP-seq coverage across the *Srsf2* gene for mock and MHV68-infected Xrn1 kd cells. Alignment files were converted to the tiled data file (tdf) format and visualized in the Integrative Genome Viewer (IGV). (F) Scatterplot of Pol II occupancy of the gene bodies comparing mock to WT MHV68-infected cells during Xrn1 kd with a linear

regression line plotted in blue and equation shown. A y=x line is plotted for reference in red. (G)

Sequence tags were binned into 25-bp bins for transcription termination sequence (TTS)

proximal Pol II for -1000 to +1000 around the TTS for mock (purple) and MHV68 infected

(green).

Fig 3: Viral genes are not susceptible to RNA decay dependent Pol II repression

(A) Viral gene Pol II recruitment is independent of RNA decay. Sequence tags were binned into

25-bp bins for -500 to +500 around the TSS. MHV68 WT (blue) and R443I (green) traces are

shown. (B) Pol II ChIP-seq coverage across 5.5kb of the MHV68 genome for MHV68 WT and

R443I in wild type (WT) cells and MHV68 infection in Xrn1 kd cells. Alignment files were

converted to tdf format and visualized in the IGV. (C) Pol II ChIP qPCR signal is similar across

gene bodies. Three regions of the ORF54 and ORF37 gene were assayed by Pol II ChIP-qPCR

during MHV68 WT or R443I infection to assess the relative levels of Pol II across the gene.

Fig 4: PABPC is not excluded from replication compartments

(A) Imunofluoresence (IF) was performed on MHV68 infected 3T3 cells at 27 hpi and stained

with PABPC, Pol II and DAPI. The MHV68 genome contains GFP, which served as a marker of

infection. Cells with RCs are denoted with red arrows. RCs were identified in cell nuclei with

PolII staining overlapping regions that are DAPI negative. The inset shows a merge of Pol II and

PABPC staining for several cells that co-stain for both proteins in RCs. (B) IF was performed on

KSHV-positive iSLK cells reactivated for 48 h and stained with PABPC, ORF59 and DAPI.

Cells with RCs are denoted with arrows and were identified in cell nuclei with ORF59 staining

that overlaps with DAPI negative regions. The inset shows a merge of ORF59 and PABPC

staining for several cells that co-stain for both proteins in RCs. (C) Percent of RC containing

cells that overlap with PABPC signal. Fractions of cells with PABPC signal in RCs over total

number of cells counted with RCs are displayed.

Fig 5 : Viral promoter sequences are insufficient to escape transcriptional regulation

(A) The MHV68 M7 viral promoter driving puromycin resistance was lentivirally integrated

into 293T cells. 24 h post transfection of muSOX or VHS, Pol II-ChIP qPCR was used to

measure Pol II levels at two host promoters (*Gapdh*, *Rplp0*) and the integrated M7 promoter. (* p

< 0.05, ** p < 0.01, *** p <0.001, **** p<0.0001 students t-test) (B) The KSHV LANA

promoter driving puromycin resistance was integrated into 293T cells by random incorporation.

Pol II-ChIP qPCR was used to measure Pol II levels 24 h post SOX or VHS transfection as

described above. (C) The MHV68 M4 viral promoter driving puromycin resistance was

lentivirally integrated into MC57G cells. The cells were then infected with WT MHV68 or

R443I at a MOI of 5 and Pol II levels at the indicated promoters were assayed by Pol II ChIP-

qPCR.

Fig 6 : Non-viral promoters escape repression on the viral genome

(A) 4SU-RNA levels of luciferase driven by CMV in muSOX and vector-control expressing

cells. Cells were separated using the co-expressed Thy1.1 marker to ensure pure populations of

transfected cells. (B) The CMV promoter driving puromycin resistance was lentivirally

integrated into MC57G cells. The cells were then infected with WT MHV68 or R443I at an MOI

of 5 and Pol II levels at the indicated promoters were assayed by Pol II ChIP-qPCR. (C) ChIP-

seq traces comparing WT MHV68 to R443I coverage from -300 to +200 around the TSS in 25

bp bins, averaged from two biological replicates. Six representative viral genes and CMV-GFP

are shown. (D) 293T cells were infected with KSHV and then transfected with muSOX twice

over a 24 h period to induce accelerated decay. Pol II levels at the indicated KSHV and human

promoters were measured by ChIP-qPCR. (** p < 0.01, students t-test)

Fig S1: ChIP-seq replicate with WT and R443I MHV68

A) Pol II ChIP-seq signal profiles of host genes are shown in mock-infected, MHV68-infected

and R443I-infected cells. Each row of the heat map displays Pol II occupancy of one gene from -

1000 to +1000 in 25bp bins. Genes are ranked by the Pol II-transcription start site (TSS)

proximal signal in mock infected cells. (B) Sequence tags were binned into 25-bp bins for -2000

to +4000 around the TSS. Mock, MHV68 (red) and MHV68 (blue) and R443I (green) traces are

shown along with their input controls. (C) Pol II ChIP-seq coverage across the *Srsf2* gene for

ChIP-seq replicate 1 and 2 with mock, WT MHV68 and R443I shown. Alignment files were

converted to the tiled data file (tdf) format and visualized in the Integrative Genome Viewer. (D)

Pol II transcription termination is not dependent on RNA decay. Sequence tags were binned into

25-bp bins for transcription termination sequence (TTS) proximal Pol II for -1000 to +1000 bp

around the TTS with the same color scheme as (B).

Fig S2: ChIP-seq replicate with WT MHV68 in Xrn1kd cells

(A) Sequence tags were binned into 25-bp bins for -4000 to +4000 around the transcription start

site (TSS) in mock (purple) and MHV68 infected (green) Xrn1 kd samples for Pol II ChIPs. (B)

Pol II ChIP-seq signal profiles of host genes in mock and MHV68-infected cells depleted of

Xrn1. Each row of the heat map displays Pol II occupancy of one gene from -1000 to +1000 in

25 bp bins. (C) Sequence tags were binned into 25-bp bins for transcription termination sequence

(TTS) proximal Pol II for -1000 to +1000 bp around the TTS.

Fig S3: Unreactivated iSLK cells show primarily cytoplasmic PABPC signal

(A)IF was performed on unreactivated iSLK cells using antibodies recognizing PABPC and

ORF59. Host DNA was stained with DAPI.

591   **References**
592

Abernathy, E. & Glaunsinger, B., 2015. Emerging roles for RNA degradation in viral replication
and antiviral defense. *Virology*, 479-480(C), pp.600–608.

Abernathy, E. et al., 2014. Gammaherpesviral Gene Expression and Virion Composition Are
Broadly Controlled by Accelerated mRNA Degradation L. F. van Dyk, ed. *PLoS Pathogens*,
10(1), pp.e1003882–14.

Abernathy, E. et al., 2015. Viral Nucleases Induce an mRNA Degradation- Transcription
Feedback Loop in Mammalian Cells. *Cell host & microbe*, 18(2), pp.1–12.

Abrisch, R.G. et al., 2016. Infection by Herpes Simplex Virus 1 Causes Near-Complete Loss of
RNA Polymerase II Occupancy on the Host Cell Genome. R. M. Sandri-Goldin, ed. *Journal
of Virology*, 90(5), pp.2503–2513.

Adler, H. et al., 2000. Cloning and mutagenesis of the murine gammaherpesvirus 68 genome as
an infectious bacterial artificial chromosome. *Journal of Virology*, 74(15), pp.6964–6974.

Brulois, K.F. et al., 2012. Construction and Manipulation of a New Kaposi's Sarcoma-Associated
Herpesvirus Bacterial Artificial Chromosome Clone. *Journal of Virology*, 86(18), pp.9708–
9720.

Chang, L. et al., Herpesviral replication compartments move and coalesce at nuclear speckles to
enhance export of viral late mRNA. *National Acad Sciences*

Covarrubias, S. et al., 2011. Coordinated Destruction of Cellular Messages in Translation
Complexes by the Gammaherpesvirus Host Shutoff Factor and the Mammalian Exonuclease
Xrn1 R. Renne, ed. *PLoS Pathogens*, 7(10), pp.e1002339–15.

Dai-Ju, J.Q. et al., 2006. ICP27 Interacts with the C-Terminal Domain of RNA Polymerase II
and Facilitates Its Recruitment to Herpes Simplex Virus 1 Transcription Sites, Where It
Undergoes Proteasomal Degradation during Infection. *Journal of Virology*, 80(7), pp.3567–
3581.

Dauber, B., Pelletier, J. & Smiley, J.R., 2011. The Herpes Simplex Virus 1 vhs Protein Enhances
Translation of Viral True Late mRNAs and Virus Production in a Cell Type-Dependent
Manner. *Journal of Virology*, 85(11), pp.5363–5373.

Dembowski, J.A. & DeLuca, N.A., 2015. Selective Recruitment of Nuclear Factors to
Productively Replicating Herpes Simplex Virus Genomes R. D. Everett, ed. *PLoS
Pathogens*, 11(5), pp.e1004939–35.

Dori-Bachash, M. et al., 2012. Widespread promoter-mediated coordination of transcription and
mRNA degradation. *Genome biology*, 13(12).

Eckmann, C.R., Rammelt, C. & Wahle, E., 2010. Control of poly(A) tail length. *Wiley
Interdisciplinary Reviews: RNA*, 2(3), pp.348–361.

Gaglia, M.M. et al., 2012. A Common Strategy for Host RNA Degradation by Divergent
Viruses. *Journal of Virology*, 86(17), pp.9527–9530.

Gallouzi, I.E. & Wilusz, J., 2013. A DIStinctively novel exoribonuclease that really likes U. *The
EMBO journal*, 32(13), pp.1799–1801.

Gilbertson, S. et al., 2018. Changes in mRNA abundance drive shuttling of RNA binding
proteins, linking cytoplasmic RNA degradation to transcription. *eLife*.

Glaunsinger, B., Chavez, L. & Ganem, D., 2005. The Exonuclease and Host Shutoff Functions
of the SOX Protein of Kaposi's Sarcoma-Associated Herpesvirus Are Genetically Separable.
Journal of Virology, 79(12), pp.7396–7401.

Günther, T. & Grundhoff, A., 2010. The Epigenetic Landscape of Latent Kaposi Sarcoma-
Associated Herpesvirus Genomes P. Kellam, ed. *PLoS Pathogens*, 6(6), pp.e1000935–19.

Haimovich, G. et al., 2013. Gene Expression Is Circular: Factors for mRNA Degradation Also
Foster mRNA Synthesis. *Cell*, 153(5), pp.1000–1011.

Heinz, S. et al., 2010. Simple Combinations of Lineage-Determining Transcription Factors Prime
cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell*,
38(4), pp.576–589.

Hilton, I.B. et al., 2013. The Open Chromatin Landscape of Kaposi's Sarcoma-Associated
Herpesvirus. *Journal of Virology*, 87(21), pp.11831–11842.

Hollingworth, R. et al., 2017. Localization of Double-Strand Break Repair Proteins to Viral
Replication Compartments following Lytic Reactivation of Kaposi's Sarcoma-Associated
Herpesvirus J. U. Jung, ed. *Journal of Virology*, 91(22), pp.1186–21.

Huang, C. et al., 2011. SARS Coronavirus nsp1 Protein Induces Template-Dependent
Endonucleolytic Cleavage of mRNAs: Viral mRNAs Are Resistant to nsp1-Induced RNA
Cleavage R. S. Baric, ed. *PLoS Pathogens*, 7(12), pp.e1002433–18.

Jagger, B.W. et al., 2012. An Overlapping Protein-Coding Region in Influenza A Virus Segment
3 Modulates the Host Response. *Science*, 337(6091), pp.199–204.

Jeong, J., Papin, J. & Dittmer, D., 2001. Differential Regulation of the Overlapping Kaposi's
Sarcoma-Associated Herpesvirus vGCR (orf74) and LANA (orf73) Promoters. *Journal of
Virology*, 75(4), pp.1798–1807.

Kamitani, W. et al., 2009. A two-pronged strategy to suppress host protein synthesis by SARS
coronavirus Nsp1 protein. *Nature Structural & Molecular Biology*, 16(11), pp.1134–1140.

Karr, B.M. & Read, G.S., 1999. The virion host shutoff function of herpes simplex virus
degrades the 5" end of a target mRNA before the 3" end. *Virology*, 264(1), pp.195–204.

Kumar, G.R., Shum, L. & Glaunsinger, B.A., 2011. Importin alpha-mediated nuclear import of
cytoplasmic poly(A) binding protein occurs as a direct consequence of cytoplasmic mRNA
depletion. *Molecular and Cellular Biology*, 31(15), pp.3113–3125.

Li, D., Fu, W. & Swaminathan, S., 2018. Continuous DNA replication is required for late gene
transcription and maintenance of replication compartments in gammaherpesviruses E. S.
Robertson, ed. *PLoS Pathogens*, 14(5), pp.e1007070–25.

Liu, S.W. et al., 2013. The D10 Decapping Enzyme of Vaccinia Virus Contributes to Decay of
Cellular and Viral mRNAs and to Virulence in Mice. *Journal of Virology*, 88(1), pp.202–
211.

Monier, K. et al., 2000. Annexation of the interchromosomal space during viral infection. *Nature
Cell Biology*, 2(9), pp.661–665.

O'Grady et al., 2019. *Manuscript in revision*.

Parrish, S. & Moss, B., 2006. Characterization of a vaccinia virus mutant with a deletion of the
D10R gene encoding a putative negative regulator of gene expression. *Journal of Virology*,
80(2), pp.553–561.

Rice, S.A. et al., 1994. Rna-Polymerase-Ii Is Aberrantly Phosphorylated and Localized to Viral
Replication Compartments Following Herpes-Simplex Virus-Infection. *Journal of Virology*,
68(2), pp.988–1001.

Richner, J.M. et al., 2011. Global mRNA Degradation during Lytic Gammaherpesvirus Infection
Contributes to Establishment of Viral Latency S. H. Speck, ed. *PLoS Pathogens*, 7(7),
pp.e1002150–13.

Rutkowski, A.J. et al., 2015. Widespread disruption of host transcription termination in HSV-1
infection. *Nature Communications*, 6, p.7126.

Schmid, M. et al., 2014. DNA Virus Replication Compartments. *Journal of Virology*, 88(3),
pp.1404–1420.

Schoenberg, D.R. & Maquat, L.E., 2012. Regulation of cytoplasmic mRNA decay. *Nature
Publishing Group*, 13(4), pp.246–259.

Shalem, O. et al., 2011. Transcriptome Kinetics Is Governed by a Genome-Wide Coupling of
mRNA Production and Degradation: A Role for RNA Pol II G. S. Barsh, ed. *PLOS Genetics*,
7(9), pp.e1002273–10.

Shalem, O. et al., 2008. Transient transcriptional responses to stress are generated by opposing
effects of mRNA production and degradation. *Molecular Systems Biology*, 4(1), pp.223–10.

Shiflett, L.A. & Read, G.S., 2013. mRNA decay during herpes simplex virus (HSV) infections:
mutations that affect translation of an mRNA influence the sites at which it is cleaved by the
HSV virion host shutoff (Vhs) protein. *Journal of Virology*, 87(1), pp.94–109.

Smibert, C.A. & Smiley, J.R., 1990. Differential Regulation of Endogenous and Transduced
Beta-Globin Genes During Infection of Erythroid-Cells with a Herpes-Simplex Virus Type-1
Recombinant. *Journal of Virology*, 64(8), pp.3882–3894.

Smiley, J.R., 2004. Herpes simplex virus virion host shutoff protein: immune evasion mediated
by a viral RNase? *Journal of Virology*, 78(3), pp.1063–1068.

Smiley, J.R., SMIBERT, C. & EVERETT, R.D., 1987. Expression of a Cellular Gene Cloned in
Herpes-Simplex Virus - Rabbit Beta-Globin Is Regulated as an Early Viral Gene in Infected
Fibroblasts. *Journal of Virology*, 61(8), pp.2368–2377.

Spencer, C.A., Dahmus, M.E. & Rice, S.A., 1997. Repression of host RNA polymerase II
transcription by herpes simplex virus type 1. *Journal of Virology*, 71(3), pp.2031–2040.

Sun, M. et al., 2012. Comparative dynamic transcriptome analysis (cDTA) reveals mutual
feedback between mRNA synthesis and degradation. *Genome Research*, 22(7), pp.1350–
1359.

Szyf, M. et al., 1985. Cellular and Viral-Dna Hypomethylation Associated with Induction of
Epstein-Barr Virus Lytic Cycle. *Proceedings of the National Academy of Sciences*, 82(23),
pp.8090–8094.

Taylor, T.J. et al., 2003. Herpes simplex virus replication compartments can form by coalescence
of smaller compartments. *Virology*, 309(2), pp.232–247.

Toth, Z. et al., 2010. Epigenetic Analysis of KSHV Latent and Lytic Genomes P. Kellam, ed.
PLoS Pathogens, 6(7), pp.e1001013–17.

Wong-Ho, E. et al., 2014. Unconventional Sequence Requirement for Viral Late Gene Core
Promoters of Murine Gammaherpesvirus 68. *Journal of Virology*, 88(6), pp.3411–3422.

Figure 1

a
mock MHV68 R443I

host genes

distance from TSS
(-1000,1000)

b
Pol II ChIP of MHV68 and R443I

relative % input

Mock
MHV68
R443I
IgG

Rplp0   Fus

c
Pol II signal around the TSS

ChIP fragment coverage

samples
Mock
MHV68
R443I

Distance from TSS

d
Promoter (-50,+200) Pol II Occupancy

MHV68

Mock

y = 0.37x + 0.48

Promoter (-50,+200) Pol II Occupancy

R443I

Mock

y = 0.70x + 0.39

Gene body (+200,+5000) Pol II Occupancy

MHV68

Mock

y = 0.83x + 1.69

Gene body (+200,+5000) Pol II Occupancy

R443I

Mock

y = 0.96x + 0.48

e
Pol II signal around the TTS

ChIP fragment coverage

samples
Mock
MHV68
R443I

Distance from TTS

Figure 2

Figure 3

a

Pol II signal at viral TSS

b

c

Pol II ChIP of MHV68 ORF54

Pol II ChIP of MHV68 ORF37

Figure 4

a MHV68 infection of 3T3s

b KSHV reactivation of iSLKs

c

Percent of cells co-staining for PABPC
and replication compartments

Figure 5

Figure 6

Supp Figure 1

a mock MHV68 R443I

host genes

distance from TSS
(-1000 to 1000)

b Pol II signal around TSS

ChIP fragment coverage

Distance from TSS

samples
Mock
MHV68
R443I

d Pol II signal around TTS

ChIP fragment coverage

Distance from TTS

samples
Mock
MHV68
R443I

c

mock 1 [0 - 75]
MHV68 1 [0 - 75]
R443I 1 [0 - 75]
mock 2 [0 - 75]
MHV68 2 [0 - 75]
R443I 2 [0 - 75]

Srsf2

Supp Figure 2

a Pol II signal around TSS

b Xrn1 kd MC57G
mock   MHV68

host genes

distance from TSS (-1000 to 1000)

c Pol II signal around TTS

Supp Figure 3

a

Unreactivated iSLKs harboring latent KSHV