1   Herpesvirus infection reduces Pol II occupancy of host promoters but spares viral promoters

13   Ella N Hartenian[1], Britt A Glaunsinger[1,2,3,*]

14
15   [1]Department of Molecular and Cell Biology, University of California Berkeley, CA, USA

16
17   [2]Department of Plant and Microbial Biology, University of California Berkeley, CA, USA

18
19   [3]Howard Hughes Medical Institute, University of California Berkeley, CA, USA

20
21   *Corresponding author

22
23   Email: glaunsinger@berkeley.edu

24
25
26
27

Abstract

In mammalian cells, widespread acceleration of cytoplasmic mRNA degradation is linked to

impaired RNA polymerase II (Pol II) transcription. This mRNA decay-induced transcriptional

repression occurs during infection with gammaherpesviruses including Kaposi's sarcoma-

associated herpesvirus (KSHV) and murine gammaherpesvirus 68 (MHV68), which encode an

mRNA endonuclease that initiates widespread RNA decay. Here, we show that MHV68-induced

mRNA decay leads to a genome-wide reduction of Pol II occupancy at mammalian promoters.

Viral genes, despite the fact that they require Pol II for transcription, escape this transcriptional

repression. Protection is not governed by viral promoter sequences; instead, location on the viral

genome is both necessary and sufficient to escape the transcriptional repression effects of mRNA

decay. We hypothesize that the ability to escape from transcriptional repression is linked to the

localization of viral DNA in replication compartments, providing a means for these viruses to

counteract decay-induced viral transcript loss.

Introduction

Regulating messenger RNA (mRNA) abundance is of central importance during both

cellular homeostasis and disease. While it is intuitive that transcriptional changes in the nucleus

impact RNA levels in the cytoplasm, evidence in both yeast and mammalian cells now indicates

that the converse is also true: altered cytoplasmic mRNA decay rates can broadly impact

transcription by RNA polymerase II (Pol II). In yeast, stabilizing the cytoplasmic mRNA pool by

removing mRNA decay factors such as Xrn1 causes a compensatory decrease in Pol II

transcription (1-5). Furthermore, reducing or slowing Pol II transcription leads to an increase in

50 overall mRNA stability (1), supporting a "buffering" model in which yeast can compensate for

51 widespread perturbations to mRNA abundance by alternatively changing rates of transcription.

52   Broad changes in mRNA abundance are often triggered in mammalian cells during viral

53 infection. Numerous viruses, including alphaherpesviruses, gammaherpesviruses, SARS

54 coronavirus and influenza A virus drive accelerated mRNA decay during infection by expressing

55 mRNA specific endoribonucleases and/or activating host nucleases. This decay contributes to

56 viral immune evasion, increases the availability of host translation machinery, and facilitates

57 temporal viral gene regulation (6-10). Viral endonucleases target host and viral mRNAs for

58 cleavage, whereupon cellular exonucleases degrade the resulting mRNA fragments (11-15). This

59 strategy accelerates basal RNA decay by circumventing the typically rate limiting steps of

60 deadenylation and decapping (16-20).

61   Although yeast and mammals share key proteins involved in cytoplasmic mRNA decay

62 and transcription, studies with RNA-decay inducing viruses in mammals have revealed that

63 unlike the yeast pathway, accelerated mRNA decay leads to an extensive shutoff of cellular gene

64 expression (21,22). Infection with the gammaherpesviruses KSHV or MHV68 causes reduced

65 Pol II recruitment at host promoters in a manner dependent on mRNA cleavage by the viral

66 endonucleases SOX or muSOX. Thus, accelerated mRNA decay is not broadly counteracted by

67 increased transcription-based repopulation, but instead leads to a more extensive shutdown of

68 cellular gene expression. However, similar to yeast, the mammalian connection between decay

69 and transcription requires the activity of cellular exonucleases such as Xrn1, which degrade the

70 viral endonuclease cleavage products (16,21). Accelerated Xrn1-mediated degradation causes

71 trafficking of released RNA binding proteins (RBP) from the cytoplasm to the nucleus, which

72 may provide a means for the cell to detect and respond to increased RNA turnover (22).

During MHV68 infection, in addition to host mRNA, cytoplasmic viral mRNA is

susceptible to cleavage by muSOX (23). However, measurements of nascent transcript

production from a subset of viral promoters suggested that viral genes are robustly transcribed

during the stage of infection when host transcription is reduced (24). An outstanding question is

how these viral promoters escape mRNA decay-induced transcriptional repression despite being

transcribed by RNA Pol II. Here, we measured Pol II occupancy across the mouse and MHV68

genomes in infected cells to more comprehensively define how accelerated mRNA decay

impacts polymerase occupancy. Our data demonstrate a genome-wide reduction in Pol II

recruitment at host promoters under conditions of increased mRNA decay. In contrast, Pol II

occupancy of the viral genome appears broadly resistant to the effects of mRNA degradation.

This protection is not conferred by viral promoter sequences; instead, location on the replicating

viral genome is both necessary and sufficient to escape the transcriptional effects of mRNA

decay. We propose a model in which DNA amplified in viral replication compartments, unlike

the cellular chromatin, is immune to transcriptional repression during accelerated cytoplasmic

mRNA decay. Thus, while both viral and cellular mRNA pools are susceptible to cytoplasmic

degradation, transcriptional repopulation selectively counteracts this loss for viral genes.

Results

Accelerated RNA decay broadly reduces Pol II occupancy

We previously reported that Pol II occupancy at several individual mammalian promoters

was significantly reduced during accelerated cytoplasmic mRNA decay (22,24). To more

comprehensively assess the extent of mRNA decay-induced transcriptional repression, we

evaluated global Pol II occupancy by chromatin immunoprecipitation and deep sequencing

(ChIP-seq) in mock versus MHV68 infected MC57G mouse fibroblasts at 24 hours post

infection (hpi). As a control, we also infected cells with a version of MHV68 containing the

point mutation R443I in the viral muSOX endonuclease gene, which reduces its mRNA cleavage

activity. R443I retains a similar lytic cycle and infectious capability to wildtype (WT) MHV68 *in*

vivo and in vitro its infectious capability is similar to MHV68 in mouse fibroblasts (6,23).

MHV68-infected cells displayed promoter-proximal loss of Pol II at 86% of loci compared to

mock-infected cells, averaged across two biological replicates and subtracting input signal (Fig

1A and S1 Fig A). Loss of Pol II occupancy was primarily due to muSOX-induced mRNA

decay, as this signal loss either did not occur or was reduced at 67% of these loci during

infection with R443I MHV68 (Fig 1A and S1 Fig A). Sixty-three genes, or 1.3% of loci, showed

higher Pol II occupancy in WT infection as compared to mock infection, although these did not

share any common GO terms (S1 Table). A representative genome browser view shows reduced

Pol II signal throughout *Srsf2* in the MHV68 condition of two replicate ChIP experiments. We

independently validated the ChIP-seq results by ChIP-qPCR at the *Rplp0* and *Fus* promoters,

both of which showed a significant reduction of Pol II occupancy in cells infected with WT but

not R443I MHV68 (Fig 1C), as did a validation using a second antibody that recognizes the N-

terminus of the major subunit of Pol II, Rpb1 (Fig1D). The genome-wide reduction in poised Pol

II in WT but not R443I infected cells was also evident when the sequencing reads from positions

-2000 to +4000 around the transcription start sites (TSS) were plotted as a histogram (Fig 1G and

S1 Fig B). The variability between replicates in R443I recovery (Fig 1B) is likely due to

differences in the extent of host shutoff reduction between replicates due to R443I reducing

muSOX activity while not being a catalytic mutant.

To assess the stage of transcription impacted by RNA decay, we compared the amount of

Pol II in promoter regions (-50 to +200) and the amount of Pol II in gene bodies (+200 to +5000)

between mock and infected samples. Consistent with the above analyses, MHV68 infected cells

contained markedly less promoter proximal Pol II than uninfected cells, and this signal was

partially recovered in the R443I infection (Fig 1E). Genes with impaired Pol II promoter binding

during MHV68 infection also had less Pol II within the gene body compared to mock infected

cells, which again was not observed during R443I infection (Fig 1F). Notably, the difference in

regression line slopes between MHV68 and mock infected cells was clearly more marked when

comparing Pol II promoter occupancy, suggesting that the primary transcriptional defect is at the

stage of initiation. This is consistent with data from previous studies that found no defect in

serine 2 phosphorylation of elongating Pol II and no consistent change to Pol II elongation

during MHV68 infection (22,24).

Finally, to assess if RNA decay affected transcription termination, we plotted sequencing

reads between 1kb upstream and 1kb downstream of the transcription termination site (TTS) as a

histogram (Fig 1H and S1 Fig D). TTS-proximal Pol II shows no defect in termination in either

infection condition, suggesting Pol II removal from host transcripts is not impacted by MHV68

infection or by RNA decay.

Pol II recruitment to viral genes is not affected by RNA decay

Herpesviral genes are transcribed in the nucleus by mammalian Pol II, and thus could be

similarly subjected to the mRNA decay-induced transcriptional repression observed across the

host genome. We therefore analyzed Pol II occupancy of the viral genome in WT MHV68 and

R443I infected cells to determine how viral genes respond to RNA decay. The average Pol II

signature across 82 viral ORFs between -500 and +500 around the TSS was comparable in cells

infected with WT or R443I MHV68, with a slightly higher signal in the R443I infection (Fig

2A). Indeed, there was a robust and widespread Pol II signal in both infection conditions across

the viral genome (Fig 2B). Thus, herpesviral genes appear to escape mRNA decay-induced

transcriptional repression.

In contrast to host genes, we did not observe peaks of promoter-proximal Pol II on the

viral genome in our ChIP-seq data (Fig 2A and B). Pol II ChIP qPCR across various regions of

ORF54 and ORF37 also showed a similar Pol II signal throughout each gene body and at their

promoter. We did note a decrease in signal near the TTS of ORF54, however this was true for

both WT and R443I infections (Fig 2C). We hypothesize that pervasive transcription of the viral

genome at 24 hpi, compounded by the presence of TSSs on both Watson and Crick strands,

likely contributes to this phenotype.

PABPC is not excluded from viral replication compartments

We next considered how the viral genome maintains high levels of Pol II occupancy

during accelerated mRNA decay. During replication, viral DNA is transcribed within viral

replication compartments (RCs) in the nucleus, which exclude host DNA and thus stain poorly

with DAPI (25-27). Although they are not membrane bound, these compartments selectively

enrich for factors required for viral replication (such as the viral processivity factor ORF59) and

gene expression (such as Pol II) (28-30). Virus-induced mRNA decay causes increased

trafficking of cytoplasmic poly(A) binding protein (PABPC) from the cytoplasm to the nucleus,

which we have recently linked to transcriptional repression (22,31). To determine whether

replication compartments selectively exclude nuclear PABPC as a means of avoiding

transcriptional repression, we monitored its localization in cells lytically infected with MHV68

or KSHV. As expected, the majority of infected cells displayed increased nuclear PABPC

compared to unreactivated or uninfected cells as measured by immunofluorescence (IF). We

further saw that PABPC was not excluded from replication compartments (Fig 3A, B and S2 Fig

A). To determine PABPC localization relative to RCs, we first identified cells with RCs by

looking for regions of Pol II or ORF59 staining that excluded DAPI. We then quantified nuclear

PABPC pixel intensity, and counted cells as having an overlapping RC/PABPC signal when the

PABPC pixel intensity was at least 2x that of cells without nuclear PABPC from the same image.

Indeed, 85 percent of MHV68 infected cells showed overlapping PABPC/Pol II signal and 90

percent of reactivated iSLKs showed an overlapping PABPC/ORF59 signal (Fig 3C). Thus,

PABPC exclusion from RCs is unlikely to underlie viral promoter escape.

Viral promoters fail to escape transcriptional repression outside of the viral genome

We then tested whether viral promoter sequence elements were sufficient to enable

escape by querying whether they conferred resistance to transcriptional repression upon

relocation from the viral genome into the host DNA. We inserted the MHV68 late promoter M7

or the KSHV LANA promoter upstream of a puromycin resistance cassette into 293T cell

chromatin using either lentiviral transduction or random plasmid DNA integration, respectively.

We induced mRNA decay and host transcriptional repression in these cells by expression of

MHV68 muSOX or the herpes simplex virus type 1 (HSV-1) endonuclease vhs. Similar to the

host promoters *Gapdh* and *Rplp0*, Pol II recruitment to both integrated viral promoters was

repressed upon infection with MHV68 but not with R443I (Fig 4A and 5B).

To confirm that viral promoters integrated into the host genome were also repressed in

the context of infection, we infected MC57G cells containing an integrated MHV68 M4

promoter-driven puromycin cassette with MHV68. Indeed, the M4 promoter was repressed to a

similar degree as host *Fus* and *Rplp0* promoters during infection with WT MHV68 but not the

mRNA decay-deficient MHV68 R443I (Fig 4C). In summary, these data argue against the

presence of protective elements within viral promoters, and instead suggest that the chromatin

context or location on the viral genome underlies escape from mRNA decay-induced

transcriptional repression.

Non-MHV68 promoters escape repression on the viral genome

A prediction from the above results is that non-MHV68 promoters would gain protection

from transcriptional repression when present on the replicating viral genome. To test this, we

measured the Pol II occupancy and transcriptional potential of the cytomegalovirus (CMV)

promoter, which drives green fluorescent protein (GFP) expression on the MHV68 genome.

Unlike alpha- and gammaherpesviruses, the betaherpesvirus CMV does not accelerate mRNA

decay. Furthermore, in either a plasmid context or when integrated into 293T cells, the CMV

promoter is strongly repressed during muSOX-induced mRNA decay. This is seen by 4SU

incorporation measuring nascent RNA production - similar to the control *ActB* gene - and by Pol

II recruitment, similar to the control *Fus* and *Rplp0* promoters (Fig 5A and B). However, in the

context of the replicating MHV68 genome, the CMV-GFP promoter had a Pol II ChIP-seq signal

during both MHV68 WT and R443I infection that was not markedly different from several other

viral genes (ORFs 4, 25, 48, 52, 72, M11) (Fig 5C).

Finally, we evaluated whether the lytic stage of infection was necessary to confer escape

from transcriptional repression, or whether some other feature of the non-replicating viral

genome was important. To this end, we examined Pol II occupancy at promoters of several genes

expressed from the KSHV genome during latent infection in 293T cells, when the viral DNA is

maintained as an episome but not amplified in replication compartments. Latent infection does

not promote mRNA turnover, and thus we transfected muSOX into these cells to stimulate

mRNA turnover in the absence of lytic replication. Similar to the host genes *Gapdh* and *Rplp0*,

we observed an RNA decay-induced repression of Pol II occupancy at the latent viral promoter

LANA while the viral K1 promoter was repressed in 4 out of 5 experiments (Fig 5D). We also

observed repression of the human *Ef1α* promoter, which is present on the KSHV viral episome

as a driver of the GFP reporter gene (Fig 5D). In sum, these observations indicate that the viral

genome per se does not confer protection from transcriptional repression; instead, features linked

to lytic genome amplification in replication compartments facilitate robust Pol II recruitment

under conditions of accelerated mRNA turnover.

Discussion

Here, we demonstrate that virus-induced mRNA decay broadly decreases Pol II

recruitment to mammalian promoters, extending prior observations made with individual cellular

genes to the genomic scale. In contrast, promoters on the replicating viral genome recruit Pol II

with similar efficiency during normal or accelerated mRNA decay, indicating that they escape

transcriptional inhibition. This protection is lost upon relocation of the viral promoters into the

host genome or when the viral genome is in a latent state, emphasizing the importance of DNA

context and replication state, rather than promoter sequence, in determining the transcriptional

response to accelerated mRNA decay.

Across the host genome, Pol II appears to be most strongly affected at the stage of

recruitment. The reduction in promoter-proximal Pol II is propagated into the gene body,

although the differences there were less pronounced, suggesting that once transcription of a gene

initiates there are not subsequent blocks to elongation. This agrees with prior studies of

individual host loci in MHV68 infected or muSOX expressing cells (22,24). The population of

Pol II that is promoter-proximal and measured during a ChIP experiment has been shown to

comprise a combination of actively engaged Pol II as well as Pol II transiently binding to the

DNA. Thus, a large reduction in promoter-proximal Pol II may not be expected to be borne out

in the gene body (32). Additionally, the antibody we use in this study binds with higher affinity

to hypophosphorylated Pol II(33), likely affecting our ability to pull down elongating and

phosphorylated forms of Pol II. We also did not observe a defect in transcription termination, or

evidence of polymerases on the DNA after TTSs, suggesting that the gammaherpesvirus MHV68

does not induce the termination defect and run-on transcription of host genes recently described

in HSV-1 (34).

Two pieces of evidence indicate that reduction of Pol II at host genes is unlikely due to

sequestration of or competition for the transcription machinery by the viral DNA. First, infection

with the mRNA decay mutant virus R443I, which replicates its viral genome to WT levels (6),

does not induce the widespread reduction in Pol II recruitment to host promoters. Second,

transfection of muSOX alone (or other broad acting viral endonucleases) in the absence of viral

infection is sufficient to induce transcriptional repression at host promoters. Therefore, we

instead favor the hypothesis that one or more cellular proteins released from degrading mRNA

fragments triggers a signaling event that restricts transcription machinery occupancy of

mammalian chromatin (22).

Reduced Pol II occupancy of the host genome during MHV68 infection parallels

infection data from the alphaherpesvirus HSV-1. HSV-1 infection results in a genome-wide

reduction in Pol II occupancy of host chromatin and reduced nascent RNA production of the

majority of host transcripts (34-36). Furthermore, early data showed that while the endogenous

mouse β-globin locus was transcriptionally repressed during HSV-1 infection, its expression was

rescued upon integration of the β-globin gene into the HSV-1 genome (37,38). Experiments with

gene-specific null viruses have implicated immediate early (IE) proteins in the HSV-1

transcriptional repression phenotype, including ICP27, which can accelerate mRNA turnover

(36,39). Additionally, we have previously published, and confirm in this report, that transfection

of HSV-1 vhs, another broad acting endoribonuclease, is sufficient to induce reduced Pol II

occupancy of several host promoters (21). Thus, in addition to targeting host and viral mRNAs

for cytoplasmic decay, both alpha and gammaherpesviruses restrict Pol II occupancy of the host

genome, although additional work is needed to establish whether the underlying mechanism is

similar.

During the lytic stage of herpesvirus infection, viral DNA is localized to replication

compartments. These non-membrane bound structures are sites of viral genome replication,

transcription, and packaging (40). They are enriched for many host and viral proteins involved in

these processes, including chromatin modifying factors and Pol II (28-30). Replication

compartments grow and coalesce over infection, excluding host chromatin, which becomes

pushed to the nuclear periphery (25-27). We recently showed that muSOX and Xrn1-coordinated

mRNA decay causes differential trafficking of many cellular RNA binding proteins between the

cytoplasm and nucleus. Knockdown and overexpression studies suggested that at least one of

these proteins, PABPC, is associated with transcriptional repression (22). Here, we detected

PABPC in viral replication compartments of infected cell nuclei, indicating that selective

exclusion of PABPC from these compartments is not the mechanism by which the viral genome

escapes repression. Instead, we hypothesize that either the state of viral DNA or other protective

features of the replication compartment confers escape from transcriptional repression.

Protection from mRNA decay-induced transcriptional repression was not conferred to

non-MHV68 or viral promoters present on the latent viral episome, indicating that lytic phase

viral DNA replication or replication compartment formation is key to this phenotype. The latent

genomes of gammaherpesviruses including KSHV and Epstein Bar Virus (EBV) are heavily

methylated and histone-rich, with the exception of promoters of genes expressed during latency,

the latent origin of replication, and CTCF binding sites (41-43). Notably, during latency key lytic

promoters like RTA contain both the activating H3K4me3 and the repressive H3K27me3 histone

marks, representing a 'poised' state (44). Rapid changes shortly after lytic cycle induction result

in hypomethylation and removal of repressive marks at early gene promoters (41,44). At later

stages of infection, KSHV, EBV and HSV-1 genomes have been shown to be largely

nucleosome free, consistent with histones not being packaged in virions (27,29,45). For

gammaherpesviruses, this is the stage at which mRNA degradation is most pronounced. We

therefore hypothesize that the 'open' state of viral DNA during lytic replication plays an

important role in facilitating Pol II recruitment under conditions of accelerated cytoplasmic

mRNA decay. Future work is geared towards exploring whether widespread mRNA decay alters

cellular chromatin states and DNA accessibility and, if so, how this influences transcription

factor recruitment.

In conclusion, decay-coupled transcriptional repression is a novel facet of viral

manipulation of the cellular gene expression landscape. We hypothesize that the ability to escape

from transcriptional repression is linked to the open conformation of viral DNA in replication

compartments, and that this helps these viruses counteract decay-induced viral transcript loss.

While widespread mRNA decay has been shown to facilitate immune evasion, gene expression

control, and latency establishment, the robust viral transcription we document here presents a

parsimonious explanation for a novel way in which the virus benefits from accelerated decay.

Materials and Methods

Plasmids and Primers

pCDEF3 muSOX and VHS and SOX have been previously published (24). pCDEF3-GFP

was used as a control.

Primers used for qPCR and for cloning are listed in S2 Table. M7, M4 and CMV-puro were

made by cloning the respective promoter sequence upstream of the puromycin cassette in a

pLKO-sgRNA vector. The promoter of M4 was defined as the 1131bp upstream of the M4

coding region not overlapping with other viral ORFs and the promoter sequence of M7 was

previously published (46). pGL3-kLANA-puromycin was made by introducing the puromycin

gene into the I-CeuI and XbaI sites of pGL3-kLANA via InFusion (Clonetech) cloning. pGL3-

kLANA was made by amplifying KSHV's LANA promoter sequence (47) and using InFusion

cloning to introduce it into the XhoI and HindIII sites of pGL3.

Cells and transfections

MC57G mouse fibroblast cells (ATCC), 293T cells (ATCC) and NIH 3T3 cells (ATCC)

were maintained in DMEM (Invitrogen) supplemented with 10% fetal bovine serum (FBS).

iSLK BAC16 cells (48) were maintained in DMEM supplemented with 10% FBS and 1 mg/mL

hygromycin B. To reactivate iSLK cells, they were treated with 1 μg/mL doxycycline and 1 mM

sodium butyrate for 48 hours.

DNA transfections were carried out in 293T cells at 70-90% confluency in 10 cm plates

with 10 μg host-shutoff factor plasmid or vector control using PolyJet (SignaGen) 24 hours

before harvesting and again at 18 hours before harvesting.

Lentiviral transduction was carried out by spinfecting $1x10^6$ 293T or MC57G cells with

lentivirus made from 2nd generation plasmids at a MOI <1 for 2 hours with 4 μg/mL polybrene

(Fisher). Twenty-four hours later puromycin (1 μg/ml for 293T or 3 μg/ml for MC57G) was

added to select for transduced cells. 293T cells stably expressing the kLANA-puromycin cassette

were made by transfecting the cells with 1 μg pGL3-kLANA-puromycin using PolyJet, then

selecting with puromycin for random integration events.

For the CMV-luciferase promoter assay, populations of 293T cells expressing muSOX

were selected 24 hours post-transfection using the Miltenyi Biotec MACS cell separation system

according to the manufacturer's instructions as previously described (22).

293T cells were infected with KSHV by reactivating iSLK-Bac16 cells for 48 hours and

then spinfecting cell-free virus onto a monolayer of 293T cells, as described for lentiviral

production but in the absence of polybrene.

3 million MC57G cells were nucleofected with a 100uL Neon tip (1400 Voltage/20

Width/2 Pulse) to knockdown Xrn1 using a pool of 4 siRNAs (Dharmacon M-046621-01) or a

non-targeting control (Dharmacon D-001206-14) using 10 μL of 20 μM siRNA.

4SU labeling

Adapted from (49).Cells were incubated in suspension with 500 μM 4SU (Sigma T4509) in DMEM supplemented with 10% FBS for 10 min then washed with PBS. RNA was extracted with TRIzol followed by isopropanol purification. RNA (300 μg) was incubated in biotinylation buffer (50 mM HEPES [pH 7.5], 5 mM EDTA) and 5 μg MTSEA-biotin (Biotium #90066), rotating at room temperature in the dark for 30 min. A no biotin control was made from an equal amount of total RNA and incubated as above but without the addition of MTSEA-biotin. RNA was then phenol:chloroform extracted and precipitated with isopropanol. The pellet was resuspended in DEPC-treated water and mixed with 50 μL Dynabeads MyOne streptavidin C1 (Invitrogen) that had been pre-washed twice with wash buffer (100 mM Tris [pH 7.5], 10 mM EDTA, 1 M NaCl, 0.1% Tween 20). Samples were rotated in the dark for 1 hour at room temperature, then washed twice with 65°C wash buffer and twice with room temperature wash buffer. Samples were eluted twice with 100 μL 5% Beta-mercaptoethanol (BME) in DEPC H_2O for 5 min then RNA was precipitated with isopropanol and quantified by RT-qPCR.

Viral mutagenesis, propagation and infections

The WT and R443I MHV68 bacterial artificial chromosome (BAC) were previously described (6,50). MHV68 was produced by first making p0 virus by transfecting NIH 3T3 cells in 6-well plates with 2.5 μg BAC DNA using TransIT-X2 (Mirus Bio) for 24h. Five to seven days later, the cells were split into a 10 cm dish, then harvested and frozen. After 5-7 days the majority of cells showed cytopathic effect (CPE). 30 μL of p0 was then added per confluent 10 cm dish of NIH 3T12 cells, split 2 days later into four 10 cm dishes and harvested 4-6 days later when all cells were infected. Cells were collected by centrifugation (5 min, 1500 rpm) and the

pellet was dounced 10 times. The cell pellet and supernatant were then ultracentrifuged for 2 h at

30,000 rpm. The resulting pellet was resuspended and titered by plaque assay. Cells were

infected with MHV68 at an MOI of 5 for 24 hours.

Western blotting

Cells were lysed in RIPA buffer (50 mM Tris-HCl pH7.6, 150 mM NaCl, 3 mM $MgCl_2$,

10% glycerol, 0.5% NP-40, cOmplete EDTA-free Protease Inhibitors [Roche]) and then clarified

by centrifugation at 21,000 x g for 10 min at 4˚C. Whole cell lysate was quantified by Bradford

assay and resolved by SDS-PAGE. Antibodies for western blot are: Xrn1 (Bethyl A300-443A,

1:1000), and Gapdh (Abcam, ab8245, 1:2000).

RT-qPCR

RNA was reverse transcribed using AMV RT (Promega) with random 9-mer primers.

Total RNA was DNase treated with TURBO DNase (ThermoFisher) or DNase treated on column

with Zymo Direct-zol RNA MiniPrep Plus DNase. cDNA was quantified using iTaq Universal

SYBR Mastermix (BioRad) and transcript-specific primers. All qPCR results are normalized to

18S levels and WT or vector control set to 1.

Immunofluoresence

NIH 3T3 cells were plated on coverslips (7.5 x 10^4 cells/well of a 12-well), infected the

following day for 25-27 h, and then fixed in 4% formaldehyde for 10 min. iSLK-Bac16 cells

were plated on coverslips (5 x 10^4 cells/well of a 12-well), reactivated 24h later with doxycycline

and sodium butyrate for 48 hours and then fixed as above. Cells were permeabilized with ice-

cold methanol at -20°C for at least 20 min and incubated with anti-Pol II antibody (Biolegend,

8WG16 at 1:200), anti-PABPC antibody (Abcam ab21060 at 1:200 for mouse cells and Santa

Cruz sc32318 at 1:25 for human cells) or anti-ORF59 antibody (Advanced Biotechnologies 13-

211-100 at 1:200) in 5% BSA overnight at 4 °C. Secondary antibodies were added (1:1000) for 1

h at 37 °C. Coverslips were mounted in DAPI- containing Vectashield (VectorLabs). We

identified RCs by DAPI exclusion and Pol II recruitment or ORF59 staining (28,30,51). Images

were collected on a Zeiss LSM 710 AxioObserver with a 40x oil objective. Cells with RCs were

first identified, then PABPC co-localization was determined by counting cells with at least 2x

nuclear pixel intensity relative to non-RC containing cells in the same image (ImageJ).

Chromatin immunoprecipitation (ChIP)

ChIP was performed from 15 cm plates of cells. 3×10^6 MC57G or 293T cells were

plated 24 h before infection and then harvested 24 h after infection or transfection(s). Upon

harvesting, cells were washed with PBS, crosslinked in 1% formaldehyde at room temperature

for 10 min. Cells were quenched in 0.125 M glycine for 5 min and washed twice with cold PBS.

Crosslinked cell pellets were mixed with 1 mL fractionation buffer (5mM PIPES pH 8.0, 85 mM

KCl, 0.5% NP-40 with cOmplete EDTA-free Protease Inhibitors [Roche]) and incubated on ice

for 10 min, during which time the lysate was dounce homogenized to release nuclei and spun at

4,000 rpm for 5 min at 4°C to pellet nuclei. Nuclei were then resuspended in 300 µl of nuclei

lysis buffer (50mM Tris-HCl pH 8.0, 0.3% SDS, 10mM EDTA, with cOmplete EDTA-free

Protease Inhibitors [Roche]) and rotated for 10 min at 4°C followed by sonication using a

QSonica Ultrasonicator. Sonicated chromatin was then spun at 13,000 rpm for 10 min at 4°C and

the pellet discarded. 40 µg of chromatin measured with the Qubit broad range DNA reagent

(ThermoFisher) was brought to 500 μL with ChIP dilution buffer (16.7 mM Tris-HCl pH 8.0,

1.1% Triton X-100, 1.2 mM EDTA, 167 mM NaCl and cOmplete EDTA-free Protease Inhibitors

[Roche]) and incubated with 10 μg mouse monoclonal anti-RNAPII (BioLegend, 8WG16) or

mouse IgG (Fisher Scientific) overnight, whereupon samples were rotated with 20 μl protein G

dynabeads (Thermofisher) for 2 hours at 4°C. Beads were washed with low salt immune complex

(20 mM Tris pH 8.0, 1% Triton-x-100, 2 mM EDTA, 150 mM NaCl, 0.1% SDS), high salt

immune complex (20 mM Tris pH 8.0, 1% Triton-x-100, 2 mM EDTA, 500 mM NaCl, 0.1%

SDS), lithium chloride immune complex (10 mM Tris pH 8.0, 0.25 M LiCl, 1% NP-40, 1%

Deoxycholic acid, 1 mM EDTA), and Tris-EDTA for 5 min each at 4°C with rotation. DNA was

eluted from the beads using 100 μl of elution buffer (150 mM NaCl, 50 μg/ml Proteinase K) and

incubated at 55°C for 2 hours, then 65°C for 12 hours. DNA was purified using a Zymo Oligo

Clean & Concentrator kit, and quantified by qPCR using primers to the promoter regions of

indicated genes for 50 cycles. qPCR was simultaneously performed on input chromatin and

normalized to input. Raw percent input values are normalized by setting mock infected or a

transfection control to 1 and are plotted as relative % input. IgG pull downs were performed with

the MHV68 infected sample under scramble conditions, or the muSOX or SOX transfected

sample for transfection experiments.

ChIP Sequencing and data analysis

Libraries were prepared for sequencing (Kapa Hyper Prep) using equal amounts of

chromatin within each experiment, amplified for 10-12 cycles based on Kapa's recommendations

and sequenced on a HiSeq4000 with 100bp single end reads or 150bp reads (Xrn1kd second

replicate only). Sequencing quality was assessed with FastQC and trimmed with Sickle. A

custom index was made of the mm10 and MHV68 genomes with Bowtie2 build. Reads were

mapped to that index using Bowtie2 (2.3.0).

To visualize reads in the Integrative Genomics Viewer (IGV), tdf files were made from

bam files and indexes made from those bam files. To make tdf files for the viral genome, a

chrom.size file was created for MHV68 and then the count command was run. Analysis of

mapped reads was performed with HOMER (Hypergeometric Optimization of Motif

EnRichment) (52). Tag directories were created from the bam files generated from Bowtie with

the makeTagDirectory command. This was done separately for the mouse and MHV68 genomes.

For mouse genes, the analyzeRepeats command was used to look at tag density in the body of

genes and the –c pausing option was used to compare pausing ratios. annotatePeaks was run in

both the –hist and –ghist mode to assess composite Pol II occupancy across the mouse genome as

well as individual gene occupancy.

Viral analyses were done by first creating a custom genome annotation in HOMER using

the loadGenome command. The TSS file was created from data generously shared by Scott

Tibbetts (University of Florida) including portions of the viral BAC provided by Laurie Krug

(Stony Brook University). annotatePeaks was run in the –hist mode to assess composite Pol II

occupancy across the viral genome as well as individual gene occupancy.

Acknowledgements

We thank all members of the Glaunsinger and Coscoy labs for their helpful suggestions and

discussions. We thank David McSwiggen for his microscopy assistance, Jennifer Kugel for her

guidance applying HOMER to the viral genome, Scott Tibbetts for sharing MHV68 TSS data and

Laurie Krug for providing a KSHV BAC sequence. This work used the Vincent J. Coates

Genomics Sequencing Laboratory at UC Berkeley.

Figure Legends

Fig 1: Promoter-proximal Pol II recruitment to mammalian genes is RNA-decay dependent

(A) Pol II ChIP-seq signal profiles of host genes in mock-infected, MHV68-infected and R443I-

infected MC57G cells. Each row of the heat map displays Pol II occupancy of one gene from -

1000 to +1000 in 25 bp bins. Genes are ranked by the Pol II-transcription start site (TSS)

proximal signal in mock infected cells. (B) ChIP-qPCR validation of Pol II occupancy at the

Rplp0 and *Fus* promoters. Pol II ChIP was performed on mock, MHV68 WT or MHV68 R443I

infected MC57G cells and Pol II levels were assayed near the TSS of two repressed host genes

during MHV68 infection from the ChIP-seq data. (* $p < 0.05$, ** $p < 0.01$, students t-test) (C)

Sequence tags were plotted as a histogram with 25 bp bins for -2000 to +4000 around the TSS.

Mock (red), MHV68 (blue) and MHV68 R443I (green) traces are shown along with their input

controls. (D) Scatterplot of Pol II occupancy of promoters, averaging the sum of ChIP-seq tags

from -50 to +200 across two replicate experiments with inputs subtracted out. Mock is compared

to WT MHV68 infection on the left and to MHV68 R443I infection on the right. Linear

regression lines are plotted in blue and equations are provided. A y=x line is plotted for reference

in red. (E) Scatterplot of Pol II occupancy of gene bodies, averaging the sum of ChIP-seq tags

from -50 to +200 across two replicate experiments with inputs subtracted out. Data as described

for (D) (E) Pol II transcription termination is not dependent on RNA decay. Sequence tags were

plotted as a histogram in 25 bp bins for transcription termination sequence (TTS) proximal Pol II

for -1000 to +1000 around the TTS with the same color scheme as (C).

Fig 2: Viral genes are not susceptible to RNA decay dependent Pol II repression

(A) Viral gene Pol II recruitment is independent of RNA decay. Sequence tags were plotted as a histogram in 25 bp bins for -500 to +500 around the TSS. MHV68 WT (blue) and R443I (green) traces are shown from infected MC57G cells. (B) Pol II ChIP-seq coverage across 5.5 kb of the MHV68 genome for MHV68 WT and R443I in wild type (WT) cells and MHV68 infection in Xrn1 kd cells. Alignment files were converted to tdf format and visualized in the IGV. (C) Pol II ChIP qPCR signal is similar across gene bodies. Three regions of the ORF54 and ORF37 gene were assayed by Pol II ChIP-qPCR during MHV68 WT or R443I infection to assess the relative levels of Pol II across the gene.

Fig 2: PABPC is not excluded from replication compartments

(A) Imunofluoresence (IF) was performed on MHV68 infected 3T3 cells at 27 hpi and stained with PABPC, Pol II and DAPI. The MHV68 genome contains GFP, which served as a marker of infection. Cells with RCs are denoted with red arrows. RCs were identified in cell nuclei with PolII staining overlapping regions that are DAPI negative. The inset shows a merge of Pol II and PABPC staining for several cells that co-stain for both proteins in RCs. (B) IF was performed on KSHV-positive iSLK cells reactivated for 48 h and stained with antibodies against PABPC, ORF59 and DAPI. Cells with RCs are denoted with arrows and were identified in cell nuclei with ORF59 staining that overlaps with DAPI negative regions. The inset shows a merge of ORF59 and PABPC staining for several cells that co-stain for both proteins in RCs. (C) Percent of RC containing cells that overlap with PABPC signal. Fractions of cells with PABPC signal in RCs over total number of cells counted with RCs are displayed.

Fig 3: Viral promoter sequences are insufficient to escape transcriptional regulation

(A) The MHV68 M7 viral promoter driving puromycin resistance was lentivirally integrated into 293T cells. 24 h post transfection of muSOX or VHS, Pol II-ChIP qPCR was used to measure Pol II levels at two host promoters (*Gapdh*, *Rplp0*) and the integrated M7 promoter. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ students t-test) (B) The KSHV LANA promoter driving puromycin resistance was integrated into 293T cells by random incorporation. Pol II-ChIP qPCR was used to measure Pol II levels 24 h post SOX or VHS transfection as described above. (C) The MHV68 M4 viral promoter driving puromycin resistance was lentivirally integrated into MC57G cells. The cells were then infected with WT MHV68 or R443I at a MOI of 5 and Pol II levels at the indicated promoters were assayed by Pol II ChIP-qPCR.

Fig 4: Non-MHV68 promoters escape repression on the viral genome

(A) 4SU incorporated RNA levels of luciferase driven by CMV in muSOX and vector-control expressing 293T cells. Cells were separated using the co-expressed Thy1.1 marker to ensure pure populations of transfected cells(22). (B) The CMV promoter driving puromycin resistance was lentivirally integrated into MC57G cells. The cells were then infected with WT MHV68 or R443I at an MOI of 5 and Pol II levels at the indicated promoters were assayed by Pol II ChIP-qPCR. (C) ChIP-seq traces comparing WT MHV68 to R443I coverage from -300 to +200 around the TSS in 25 bp bins, averaged from two biological replicates of infected MC57G cells. Six representative viral genes and CMV-GFP are shown. (D) 293T cells were infected with KSHV and then transfected with muSOX twice over a 24 h period to induce accelerated decay. Pol II

levels at the indicated KSHV and human promoters were measured by ChIP-qPCR. (** p < 0.01,

students t-test)

Supporting information

S1 Fig ChIP-seq replicate with WT and R443I MHV68

A) Pol II ChIP-seq signal profiles of host genes are shown in mock-infected, MHV68-infected

and R443I-infected cells. Each row of the heat map displays Pol II occupancy of one gene from -

1000 to +1000 in 25 bp bins. Genes are ranked by the Pol II-transcription start site (TSS)

proximal signal in mock infected cells. (B) Sequence tags were plotted as a histogram in 25 bp

bins for -2000 to +4000 around the TSS. Mock, MHV68 (red) and MHV68 (blue) and R443I

(green) traces are shown along with their input controls. (C) Pol II ChIP-seq coverage across the

Srsf2 gene for ChIP-seq replicate 1 and 2 with mock, WT MHV68 and R443I shown. Alignment

files were converted to the tiled data file (tdf) format and visualized in the Integrative Genome

Viewer. (D) Pol II transcription termination is not dependent on RNA decay. Sequence tags were

plotted as a histogram in 25 bp bins for transcription termination sequence (TTS) proximal Pol II

for -1000 to +1000 around the TTS with the same color scheme as (B).

S2 Fig Unreactivated iSLK cells show primarily cytoplasmic PABPC signal

(A)IF was performed on unreactivated iSLK cells using antibodies recognizing PABPC and

ORF59. Host DNA was stained with DAPI.

S3? Fig 4SU-enriched RNA from MHV68 and R443I infected MC57G cells

(A) 4SU incorporated RNA levels of 7 viral genes in cells infected with MHV68 and R443I.

551
552
553   **References**
554

1.   Sun M, Schwalb B, Schulz D, Pirkl N, Etzold S, Lariviere L, et al. Comparative dynamic
transcriptome analysis (cDTA) reveals mutual feedback between mRNA synthesis and
degradation. Genome Research. 2012 Jul 2;22(7):1350–9.

2.   Haimovich G, Medina DA, Causse SZ, Garber M, Millán-Zambrano G, Barkai O, et al.
Gene Expression Is Circular: Factors for mRNA Degradation Also Foster mRNA
Synthesis. Cell. Elsevier Inc; 2013 May 23;153(5):1000–11.

3.   Shalem O, Dahan O, Levo M, Martinez MR, Furman I, Segal E, et al. Transient
transcriptional responses to stress are generated by opposing effects of mRNA production
and degradation. Mol Syst Biol. EMBO Press; 2008;4(1):223–10.

4.   Shalem O, Groisman B, Choder M, Dahan O, Pilpel Y. Transcriptome Kinetics Is
Governed by a Genome-Wide Coupling of mRNA Production and Degradation: A Role
for RNA Pol II. Barsh GS, editor. PLoS Genet. Public Library of Science; 2011 Sep
8;7(9):e1002273–10.

5.   Dori-Bachash M, Shalem O, Manor YS, Pilpel Y, Tirosh I. Widespread promoter-
mediated coordination of transcription and mRNA degradation. Genome Biol. BioMed
Central; 2012;13(12).

6.   Richner JM, Clyde K, Pezda AC, Cheng BYH, Wang T, Kumar GR, et al. Global mRNA
Degradation during Lytic Gammaherpesvirus Infection Contributes to Establishment of
Viral Latency. Speck SH, editor. PLoS Pathog. 2011 Jul 21;7(7):e1002150–13.

7.   Liu SW, Wyatt LS, Orandle MS, Minai M, Moss B. The D10 Decapping Enzyme of
Vaccinia Virus Contributes to Decay of Cellular and Viral mRNAs and to Virulence in
Mice. Journal of Virology. 6 ed. 2013 Dec 18;88(1):202–11.

8.   Parrish S, Moss B. Characterization of a vaccinia virus mutant with a deletion of the
D10R gene encoding a putative negative regulator of gene expression. Journal of
Virology. American Society for Microbiology; 2006 Jan;80(2):553–61.

9.   Read GS. Virus-encoded endonucleases: expected and novel functions. WIREs RNA. 1st
ed. 2013 Jul 30;4(6):693–708.

10.   Dauber B, Poon D, Santos dos T, Duguay BA, Mehta N, Saffran HA, et al. The Herpes
Simplex Virus Virion Host Shutoff Protein Enhances Translation of Viral True Late
mRNAs Independently of Suppressing Protein Kinase R and Stress Granule Formation.
Sandri-Goldin RM, editor. Journal of Virology. 2016 Jun 10;90(13):6049–57.

11.   Karr BM, Read GS. The virion host shutoff function of herpes simplex virus degrades the
5" end of a target mRNA before the 3" end. Virology. 1999 Nov 10;264(1):195–204.

12.   Glaunsinger B, Chavez L, Ganem D. The Exonuclease and Host Shutoff Functions of the
SOX Protein of Kaposi's Sarcoma-Associated Herpesvirus Are Genetically Separable.
Journal of Virology. 2005 May 26;79(12):7396–401.

13.   Huang C, Lokugamage KG, Rozovics JM, Narayanan K, Semler BL, Makino S. SARS
Coronavirus nsp1 Protein Induces Template-Dependent Endonucleolytic Cleavage of
mRNAs: Viral mRNAs Are Resistant to nsp1-Induced RNA Cleavage. Baric RS, editor.
PLoS Pathog. 2011 Dec 8;7(12):e1002433–18.

14.   Kamitani W, Huang C, Narayanan K, Lokugamage KG, Makino S. A two-pronged
strategy to suppress host protein synthesis by SARS coronavirus Nsp1 protein. Nature
Structural & Molecular Biology. Nature Publishing Group; 2009 Oct 18;16(11):1134–40.

15.   Jagger BW, Wise HM, Kash JC, Walters K-A, Wills NM, Xiao Y-L, et al. An
Overlapping Protein-Coding Region in Influenza A Virus Segment 3 Modulates the Host
Response. Science. 2012;337(6091):199–204.

16.   Covarrubias S, Gaglia MM, Kumar GR, Wong W, Jackson AO, Glaunsinger BA.
Coordinated Destruction of Cellular Messages in Translation Complexes by the
Gammaherpesvirus Host Shutoff Factor and the Mammalian Exonuclease Xrn1. Renne R,
editor. PLoS Pathog. 2011 Oct 27;7(10):e1002339–15.

17.   Gaglia MM, Covarrubias S, Wong W, Glaunsinger BA. A Common Strategy for Host
RNA Degradation by Divergent Viruses. Journal of Virology. 2012 Aug 9;86(17):9527–
30.

18.   Eckmann CR, Rammelt C, Wahle E. Control of poly(A) tail length. WIREs RNA. 2010
Nov 17;2(3):348–61.

19.   Gallouzi IE, Wilusz J. A DIStinctively novel exoribonuclease that really likes U. EMBO
J. Nature Publishing Group; 2013 Jun 11;32(13):1799–801.

20.   Schoenberg DR, Maquat LE. Re-capping the message. Trends in Biochemical Sciences.
2009 Sep;34(9):435–42.

21.   Abernathy E, Glaunsinger B. Emerging roles for RNA degradation in viral replication and
antiviral defense. Virology. Elsevier; 2015 May 1;479-480(C):600–8.

22.   Gilbertson S, Federspiel JD, Hartenian E, eLife IC, 2018. Changes in mRNA abundance
drive shuttling of RNA binding proteins, linking cytoplasmic RNA degradation to
transcription. eLife.

23.   Abernathy E, Clyde K, Yeasmin R, Krug LT, Burlingame A, Coscoy L, et al.
Gammaherpesviral Gene Expression and Virion Composition Are Broadly Controlled by

621   Accelerated mRNA Degradation. van Dyk LF, editor. PLoS Pathog. 2014 Jan
622   16;10(1):e1003882–14.

623   24.   Abernathy E, Gilbertson S, Alla R, Glaunsinger B. Viral Nucleases Induce an mRNA
624   Degradation- Transcription Feedback Loop in Mammalian Cells. Cell Host Microbe.
625   Elsevier Inc; 2015 Jul 21;18(2):1–12.

626   25.   Chang L, Godinez WJ, Kim, I, 2011. Herpesviral replication compartments move and
627   coalesce at nuclear speckles to enhance export of viral late mRNA. Proceedings of the
628   National Academy of Sciences. 2011 May;108(21):E136-E144.

629   26.   Taylor TJ, McNamee EE, Day C, Knipe DM. Herpes simplex virus replication
630   compartments can form by coalescence of smaller compartments. Virology. 2003
631   May;309(2):232–47.

632   27.   Monier K, Armas JCG, Etteldorf S, Ghazal P, Sullivan KF. Annexation of the
633   interchromosomal space during viral infection. Nat Cell Biol. Nature Publishing Group;
634   2000 Sep 1;2(9):661–5.

635   28.   Li D, Fu W, Swaminathan S. Continuous DNA replication is required for late gene
636   transcription and maintenance of replication compartments in gammaherpesviruses.
637   Robertson ES, editor. PLoS Pathog. 2018 May 29;14(5):e1007070–25.

638   29.   Dembowski JA, DeLuca NA. Selective Recruitment of Nuclear Factors to Productively
639   Replicating Herpes Simplex Virus Genomes. Everett RD, editor. PLoS Pathog. 2015 May
640   27;11(5):e1004939–35.

641   30.   Rice SA, Long MC, LAM V, Spencer CA. Rna-Polymerase-Ii Is Aberrantly
642   Phosphorylated and Localized to Viral Replication Compartments Following Herpes-
643   Simplex Virus-Infection. Journal of Virology. American Society for Microbiology
644   (ASM); 1994 Feb;68(2):988–1001.

645   31.   Kumar GR, Shum L, Glaunsinger BA. Importin alpha-mediated nuclear import of
646   cytoplasmic poly(A) binding protein occurs as a direct consequence of cytoplasmic
647   mRNA depletion. Molecular and Cellular Biology. American Society for Microbiology;
648   2011 Aug;31(15):3113–25.

649   32.   Darzacq X, Shav-Tal Y, de Turris V, Brody Y, Shenoy SM, Phair RD, et al. In vivo
650   dynamics of RNA polymerase II transcription. Nat Struct Mol Biol. 2007 Aug
651   5;14(9):796–806.

652   33.   Nojima T, Gomes T, Grosso ARF, Kimura H, Dye MJ, Dhir S, et al. Mammalian NET-
653   Seq Reveals Genome-wide Nascent Transcription Coupled to RNA Processing. Cell. The
654   Authors; 2015 Apr 23;161(3):526–40.

655   34.   Rutkowski AJ, Erhard F, L'Hernault A, Bonfert T, Schilhabel M, Crump C, et al.
656   Widespread disruption of host transcription termination in HSV-1 infection. Nature
657   Communications. 2015;6:7126.

35.　Abrisch RG, Eidem TM, Yakovchuk P, Kugel JF, Goodrich JA. Infection by Herpes
Simplex Virus 1 Causes Near-Complete Loss of RNA Polymerase II Occupancy on the
Host Cell Genome. Sandri-Goldin RM, editor. Journal of Virology. American Society for
Microbiology; 2016 Mar;90(5):2503–13.

36.　Spencer CA, Dahmus ME, Rice SA. Repression of host RNA polymerase II transcription
by herpes simplex virus type 1. Journal of Virology. 1997 Mar;71(3):2031–40.

37.　Smiley JR, Smibert C, Everett RD. Expression of a Cellular Gene Cloned in Herpes-
Simplex Virus - Rabbit Beta-Globin Is Regulated as an Early Viral Gene in Infected
Fibroblasts. Journal of Virology. American Society for Microbiology (ASM); 1987
Aug;61(8):2368–77.

38.　Smibert CA, Smiley JR. Differential Regulation of Endogenous and Transduced Beta-
Globin Genes During Infection of Erythroid-Cells with a Herpes-Simplex Virus Type-1
Recombinant. Journal of Virology. American Society for Microbiology Journals; 1990
Aug;64(8):3882–94.

39.　Smiley JR. Herpes simplex virus virion host shutoff protein: immune evasion mediated by
a viral RNase? Journal of Virology. American Society for Microbiology Journals; 2004
Feb;78(3):1063–8.

40.　Schmid M, Speiseder T, Dobner T, Gonzalez RA. DNA Virus Replication Compartments.
Journal of Virology. 4 ed. 2014 Jan 14;88(3):1404–20.

41.　Günther T, Grundhoff A. The Epigenetic Landscape of Latent Kaposi Sarcoma-
Associated Herpesvirus Genomes. Kellam P, editor. PLoS Pathog. 2010 Jun
3;6(6):e1000935–19.

42.　Szyf M, Eliasson L, Mann V, Klein G, Razin A. Cellular and Viral-Dna Hypomethylation
Associated with Induction of Epstein-Barr Virus Lytic Cycle. Proc Natl Acad Sci USA.
National Academy of Sciences; 1985;82(23):8090–4.

43.　Hilton IB, Simon JM, Lieb JD, Davis IJ, Damania B, Dittmer DP. The Open Chromatin
Landscape of Kaposi's Sarcoma-Associated Herpesvirus. Journal of Virology. 2013 Oct
3;87(21):11831–42.

44.　Toth Z, Maglinte DT, Lee SH, Lee H-R, Wong L-Y, Brulois KF, et al. Epigenetic
Analysis of KSHV Latent and Lytic Genomes. Kellam P, editor. PLoS Pathog. Public
Library of Science; 2010 Jul 22;6(7):e1001013–7.

45.　Hollingworth R, Horniblow RD, Forrest C, Stewart GS, Grand RJ. Localization of
Double-Strand Break Repair Proteins to Viral Replication Compartments following Lytic
Reactivation of Kaposi's Sarcoma-Associated Herpesvirus. Jung JU, editor. Journal of
Virology. 2017 Oct 27;91(22):1186–21.

46.   Wong-Ho E, Wu TT, Davis ZH, Zhang B, Huang J, Gong H, et al. Unconventional
Sequence Requirement for Viral Late Gene Core Promoters of Murine Gammaherpesvirus
68. Journal of Virology. 2014 Feb 24;88(6):3411–22.

47.   Jeong J, Papin J, Dittmer D. Differential Regulation of the Overlapping Kaposi's Sarcoma-
Associated Herpesvirus vGCR (orf74) and LANA (orf73) Promoters. Journal of Virology.
2001 Feb 15;75(4):1798–807.

48.   Brulois KF, Chang H, Lee ASY, Ensser A, Wong LY, Toth Z, et al. Construction and
Manipulation of a New Kaposi's Sarcoma-Associated Herpesvirus Bacterial Artificial
Chromosome Clone. Journal of Virology. 2012 Aug 23;86(18):9708–20.

49.   Duffy EE, Rutenberg-Schoenberg M, Stark CD, Kitchen RR, Gerstein MB, Simon MD.
Tracking Distinct RNA Populations Using Efficient and Reversible Covalent Chemistry.
Molecular Cell. 2015 Sep;59(5):858–66.

50.   Adler H, Messerle M, Wagner M, Koszinowski UH. Cloning and mutagenesis of the
murine gammaherpesvirus 68 genome as an infectious bacterial artificial chromosome.
Journal of Virology. 2000 Aug;74(15):6964–74.

51.   Dai-Ju JQ, Li L, Johnson LA, Sandri-Goldin RM. ICP27 Interacts with the C-Terminal
Domain of RNA Polymerase II and Facilitates Its Recruitment to Herpes Simplex Virus 1
Transcription Sites, Where It Undergoes Proteasomal Degradation during Infection.
Journal of Virology. 2006 Mar 14;80(7):3567–81.

52.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple Combinations of
Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for
Macrophage and B Cell Identities. Molecular Cell. Elsevier Inc; 2010 May 28;38(4):576–
89.

Figure 1

a Pol II ChIP of MC57G cells
mock   MHV68   R443I

host genes

distance from TSS
(-1000,1000)

b Genome Browser
1kb

mock 1
MHV68 1
R443I 1
mock 2
MHV68 2
R443I 2

Srsf2   Msfd11

c Pol II ChIP of MHV68 and R443I
antibody : 8WG16

relative % input

Mock
MHV68
R443I
IgG

Rplp0   Fus

d Pol II ChIP of MHV68 and R443I
antibody : Rpb1 N-term

relative % input

Mock
MHV68
R443I
IgG

Rplp0 prom   Rplp0 internal   Fus prom

e Promoter (-50, +200) Pol II Occupancy

MHV68 ChIP Tag Sum

Mock ChIP Tag Sum

y = 0.37x + 0.48

R443I ChIP Tag Sum

Mock ChIP Tag Sum

y = 0.70x + 0.39

f Gene body (+200, +5000) Pol II Occupancy

MHV68 ChIP Tag Sum

Mock ChIP Tag Sum

y = 0.83x + 1.69

R443I ChIP Tag Sum

Mock ChIP Tag Sum

y = 0.96x + 0.48

g Pol II signal around the TSS

ChIP tag coverage

samples
Mock
MHV68
R443I

Distance from TSS (bp)

h Pol II signal around the TTS

ChIP tag coverage

samples
Mock
MHV68
R443I

Distance from TSS (bp)

Figure 2

a

Pol II ChIP signal at viral TSS

ChIP tag coverage

— MHV68
— R443I

Distance from TSS (bp)

b

Genome browser of a representative genomic region

MHV68

R443I

Xrn1 kd MHV68

ORF50   M7   ORF52   ORF53   ORF54   ORF55   ORF56

c

Pol II ChIP of MHV68

relative % input

■ MHV68
■ R443I
■ IgG

TSS   middle   TTS

Pol II ChIP of MHV68

relative % input

■ MHV68
■ R443I
■ IgG

TSS   middle   TTS

Figure 3

a   MHV68 infection of 3T3s

b   KSHV reactivation of iSLKs

c

Percent of cells co-staining for PABPC
and replication compartments

Figure 4

a 293T cells with integrated MHV68 M7 promoter

Legend: vector, muSOX, vhs, IgG

b 293T cells with integrated KSHV LANA promoter

Legend: vector, SOX, vhs, IgG

c MC57G cells with integrated MHV68 M4 promoter

Legend: control, MHV68, R443I, IgG

Figure 5

Supp Figure 1

a mock MHV68 R443I

host genes

distance from TSS
(-1000 to 1000)

b Pol II signal around TSS

ChIP fragment coverage

Distance from TSS

samples
Mock
MHV68
R443I

d Pol II signal around TTS

ChIP fragment coverage

Distance from TTS

samples
Mock
MHV68
R443I

Supp Figure 2

a Unreactivated iSLKs harboring latent KSHV

Supp Figure 3

a

4SU-enriched RNA from
MC57Gs infected with MHV68 or R443I

Relative nascent RNA/18s

- scr MHV68
- scr R443I
- no biotin

M1 ORF8 ORF37 ORF54 ORF75a ORF24 GP70