Global biogeographical regions reveal a signal of past human impacts

Marta Rueda[1,2]*, Manuela González-Suárez[3], Eloy Revilla[2]

1. Department of Plant Biology and Ecology, University of Seville, Seville, Spain.

2. Department of Conservation Biology, Estación Biológica de Doñana (EBD-CSIC), Seville,

Spain.

3. Ecology and Evolutionary Biology, School of Biological Sciences, University of Reading,

Reading, UK.

Corresponding author:

Marta Rueda

Tel: +34656532402

ORCID: https://orcid.org/0000-0002-9754-6258

Email: mruedag@us.es

Abstract

Abstract

Ecologists have long documented that the world´s biota is spatially organized in regions with

boundaries shaped by processes acting on geological and evolutionary timescales. Although

growing evidence suggests that human impact has been key in how biodiversity is currently

assembled, its role as a driver of the geographical organization of biodiversity remains unclear.

Here, we quantify the relative importance of human land use during the Late Holocene to

predict the current assemblage of terrestrial mammals in biogeographical regions across the

Earth. Results show that human land use 2,000 years ago has left an imprint on the taxonomic

differentiation of some of the largest biogeographical realms, whereas land use at present

stands out as a driver of the taxonomic differences between medium-sized subregions, i.e.,

within and among continents. Our findings highlight the far-reaching effect that past

anthropogenic actions have had on the organization of biodiversity globally and warn us about

future impacts.

Introduction

How is the world's biodiversity organized, and why do large-scale patterns of taxonomic diversity change through natural geographic regions? These questions have attracted the attention of naturalists since the early 19th century[1-5] and are now critical to forecast the future of biodiversity in the face of global change[6]. A key step in understanding the organization of biodiversity is to identify the assemblage of regions based on their shared elements[4]. Alfred R. Wallace was among the first to propose that the world's fauna is organized hierarchically in broad regions shaped by geographic and climatic factors[1]. About 150 years later, the development of multivariate analytical techniques has led to revaluations of Wallace's proposal[7-10] and the refinement of our understanding regarding the extrinsic determinants driving the main dissimilarities among bioregions[11]. Broadly, what we know is that the interplay of multiple factors has jointly contributed to the formation of bioregions, but their prominence varies across the globe[11-13]. Processes acting deeper in the past, like plate tectonics, are the most important in explaining the separation between strongly divergent biogeographical realms. Factors representing present-day ecological barriers, like climatic heterogeneity, determine the separation between less dissimilar regions, i.e., within and among continents; while mountains - the third factor in importance - operate at all levels of biogeographical differentiation[11]. However, although there is clear evidence that anthropogenic actions are the main driver of the extinction processes acting at multiple scales[14,15], and, thereby, of the biotic homogenization of the ecological communities worldwide[16,17], we do not yet fully understand whether and how anthropogenic impact has caused changes at a biogeographical level.

Anthropogenic impact has usually not been contemplated in the biogeographical scenario likely

because extant biogeographical regions have traditionally been assumed to reflect the natural

organization of biodiversity that resulted from ecological, historical, and evolutionary

processes acting over millions of years[12]. However, rising evidence showing how human-

mediated species introductions are already affecting extant biogeographic regions[18,19]

challenges this view. Furthermore, accumulating facts indicate that Quaternary human

activities induced shifts in the plant and animal communities we see today[20-22], and there is

little doubt that modern humans have been a major driver in the extinction of large mammals

during the late Pleistocene and early Holocene[21,22]. Moreover, archaeological, palaeoecological

and genetic data suggest that cumulative human transformation of ecosystems over millennia

has resulted in dramatic changes in composition, community structure, richness and genetic

diversity of a diverse array of organisms across taxonomic groups[23,24]. This body of evidence

raises the question of whether anthropogenic activities may have been large enough to leave an

imprint detectable today on biogeographical assemblages globally.

To answer this question, we set up a bioregionalization and analytical procedure focused on

two data sets; one describing the current distributions of global terrestrial mammals and

another inferring their natural distributions (distribution of terrestrial mammals from the

Pleistocene to the present in the absence of human impact[25]). We then build hierarchical

bioregionalizations, upscaling from the smallest detectable bioregions to the largest realms,

applying an affinity propagation clustering algorithm[26] to the corresponding co-occurring

species matrix of global terrestrial mammals. Finally, we compute random forest classification

models to identify those variables that best predict bioregion assignation for all cells across the

globe. In the models, we include predictors previously identified as important determinants of

biogeographical boundaries[11], and predictors associated with anthropogenic activities over the

Holocene, specifically, the human land use at present, as well as 2,000 years ago.

Bioregionalizations and random forest models are also carried out using only those extant

species whose ranges of distribution are known to have undergone shifts due to anthropogenic

actions[25], as we expect a biogeographical imprint to be more apparent in this group. We

hypothesized that if anthropogenic actions have had a noteworthy effect on the biogeographical

assemblages and boundaries as we know them nowadays, then (1) the biogeographic

configurations built using current mammal's distributions will differ from those based on

natural distributions; and (2) human land use, present and past, should predict the

biogeographic patterns obtained using current distributions of mammals, but not patterns

obtained using natural distributions. Our analyses reveal that human actions have left an

imprint on how diversity is structured on Earth at a biogeographical scale. Our findings are

consistent with recent evidence showing that the impact of past human actions on ecosystems

and organisms has been greater than previously thought, and warn us about the effect of future

impacts.

Results

Bioregionalizations

Both for the current and natural bioregionalizations we identified a hierarchical system of

biogeographic regions with four levels showing that global biodiversity patterns can be

cohesively shaped from local to regional and realm scales (Fig. S1). For current mammalian

distributions, the broadest delineation of nine large bioregions was mostly consistent with the

original maps of Wallace´s realms and subregions[1,10] with some particular differences including

the limits of the Palearctic[7,9,10], and an extension towards the arid steppes of Mongolia of what

we have named the 'Saharo-Arabian' region[7,10] (Fig. 1a). For natural mammalian distributions,

the eight bioregions obtained showed a different biogeographic arrangement with respect to

the nine large current ones, especially in the Northern Hemisphere (Fig. 1b) (Table 1).

Differences between current and natural bioregions extended to the subregions or medium-

sized bioregions and smaller bioregions (Fig. S1), and were confirmed for the

bioregionalizations carried out with the 536 species whose range of distribution have been

modified by anthropogenic actions[25] (Fig. S2) (Table S1).

Global importance of predictors of the biogeographical patterns

For the broader bioregions, human land use 2,000 years ago was the third most important

driver of the biogeographic structure of current realms behind plate tectonics and seasonality

in temperature and ahead of mountains (Fig. 2c), with a moderate but non-negligible predictive

value (~22% of cells classified correctly when the model was run just with this predictor, see

Fig. 3a). Human land use at present had less relevance in the classification models (Fig. 2a). In

accordance with our prediction, past and present human land use were poorer predictors of

natural bioregions (Fig. 2b, d). Results were consistent also if we excluded Australia and

Madagascar, two biogeographic regions with minimal human land use before 2,000 years ago

(Fig. S3). Models run with the unique 536 extant terrestrial mammals whose range of

distribution has been affected by human impact revealed present and past human land use as

the second predictor after plate tectonics but with little relevance (Fig. S4). The

misclassification rate (OBB) for these set of models was quite low, never exceeding 3%.

For medium-sized bioregions or subregions, plate tectonics continued to be the most important

determinant of the biogeographic organization of both current and natural bioregions (Fig. 4;

OBB values in general < 7.50%), with human land use at present and 2,000 years ago as the

second predictors in importance for the current bioregions (Fig. 4a, c). As hypothesized, the

importance of land use was smaller for the natural bioregions, although its past land use

remained a relatively strong predictor (Fig. 4 b, d). Results were consistent for the models run

with the unique 536 extant terrestrial mammals influenced by human actions (Fig. S5; OBB <

8.5%).

As bioregions became smaller, many more variables were necessary to predict the classification

of cells into bioregions, and results for the current and natural bioregions became similar (Fig.

S6 and S7; OBB < 15.1% and < 35.6% for small and very small bioregions, respectively). Overall,

human land use at present has minor relevance in the classification models, whereas human

land use 2,000 years ago remained relatively high, being third or fourth in importance (Fig. S6c,

d and S7c, d).

Local importance of predictors of the biogeographical patterns

Maps showing local importance scores revealed large differences in the location of grid cells

where human land use at present and 2,000 years ago have been analytically important for the

classification of the broader current bioregions (Fig. 5). The cells where human land use at

present was most relevant are located in southern Africa, the easternmost boundary between

the Palearctic and the Oriental regions, part of the northern boundary between the Palearctic

and 'Saharo-Arabian' regions, and much of the arctic zone of the Palearctic (Fig. 5a), although

note that grid cells of the two latter areas are important because of their lack of human land use

(Fig. 5c). The cells where human land use 2,000 years ago was more important are located

mainly throughout Central America and the Andes mountain range, Europe except for the

northernmost part, most of the Oriental region, the boundary between the Palearctic and the

Oriental regions, as well as Madagascar and much of the arctic zone of the Palearctic due to lack

of human land use (Fig. 5b). These areas mostly coincided for the broader natural bioregions

(Fig. 5e, f).

For the medium current bioregions, the cells where the human land use at present was most

relevant for the classification of bioregions included those highlighted for the broader

bioregions plus most part of Europe, southern India, eastern Australia, a large area of the

Nearctic, and the Sahara Desert (Fig. 6a, c). When focusing on the human land use 2,000 years

ago, the areas affected by human land use were similar, with some exceptions (e.g., Sahara

Desert), although their importance scores were much higher (Fig. 6, b, d). The location of cells

affected by human land use for the medium current bioregions quite coincided with that for

natural bioregions (Fig. 6e, f).

Discussion

Our study shows that human activity during the last few millennia has left its mark on the

spatial organization of the Earth´s biodiversity. First, we show that biogeographical

configurations based on current and natural mammal distributions differ notably, particularly

in the Northern Hemisphere. These differences were not solely the result of including extinct

megafauna - with huge distributions ranges – in the natural biogeographic regions, as

biogeographical patterns for the 536 extant terrestrial mammals whose ranges of distribution

have been affected by anthropogenic impacts confirmed our findings. Second, we show that

human land use is an important predictor of current bioregions but does not largely contribute

to predict natural bioregions. For example, for the broader regions or realms, human land use

2,000 years ago was the third most important predictor behind plate tectonics and temperature

seasonality and ahead of orography, but has minor importance on natural regions. Past

anthropogenic actions have seemingly left a perceptible biogeographical imprint on the

assembly of the global realms we recognize today. Moreover, for the medium-sized subregions,

human land use at present was the second most important predictor suggesting that recent

human activity is already causing changes in the biogeographical assemblages at the

subregional scale, i.e., within and among continents.

The mammalian distributions that we know today are not only a reflection of the most

recent human actions but also those exerted during the last few millennia[27,28]. Recent studies

had already shown that human actions in the present can be evident at biogeographical

scales[18,19], but here we also show evidence of an effect of past human land use. What did

happen 2,000 years ago that left such as noticeable footprint in the Earth's biogeographical

regions? This period coincides with the settlement of densely populated areas after the

Neolithic agricultural revolution[29]. Long-term impacts to enhance agricultural productivity,

including forest clearing, increased fire frequencies, megafauna extinctions, species invasions,

and soil erosion, were already apparent in some regions at this time[20,23,24]. In fact,

reconstructions suggest that >20% of Earth's temperate woodlands had been already impacted

209 by humans by the late-Holocene[23]. Indeed, our results on local importance show the match

210 between some of the biogeographical boundaries and regions of the New World and the

211 distribution – and presumed impact – of ancient Mesoamerican and South American pre-

212 European civilizations[30,31]. In the Old World, we note the prominence of past human land use

213 covering most of the Oriental realm territory, likely associated to the rise of rice cultivation in

214 eastern, southern and southeast Asia over the Holocene[32]. Indeed, cumulative archaeological

215 data show that the scale of agriculture and land use in these regions was significant; expansion

216 of land area used for livestock and rice (*Oryza sativa*) paddy agriculture in southern Asia was

217 sufficient to increase atmospheric methane emissions between 4000 and 1000 years ago[33].

218 Moreover, the atmospheric CO_2 decline registered between A.D. 1570 and 1620 has been

219 attributed to the uptake by vegetation following the agriculture abandonment caused by the

220 population crash after the arrival of Europeans to America[34]. Human land use 2,000 years ago

221 can also partly explain differences between the southern Palearctic boundary and the

222 Afrotropic below the Sahara. Both can be related to the transformation of European and Near

223 Eastern landscapes during the Roman period[35], and the expansion of sorghum (*Sorghum*

224 *bicolor*) cultivation 3000 to 2000 years ago in the sub-Saharan Africa[34].

226 While we found differences in importance of human land use 2,000 years ago between current

227 and natural medium-sized bioregions, we were surprised that past human land use did partly

228 predict natural bioregions, which were based on estimated distributions of mammals in the

229 absence of human impact. It is possible that the natural distribution maps did not fully exclude

230 human impacts as for 3,424 of the extant mammals analyzed, natural and current distributions

231 were identical. In addition, these results could also reflect an overlap between human land use

and biogeography. If the distribution of humans 2,000 years ago responded to the same drivers

as the distributions of terrestrial mammals, human land use could appear as a predictor even if

it had no causal effect. In fact, the importance of this variable appears to be a consequence of

how human land use was distributed globally 2,000 years ago, or more accurately where

human activity was largely absent. While current human land use is distributed over wide areas

of the planet, that of 2,000 years ago was centralized in specific areas coinciding with the

distribution of ancient civilizations. This means that on a subregional scale, many bioregions

can be predicted by the absence of human land use. Indeed, an additional analysis that excluded

subregions with no reported human land use 2,000 years ago showed very little importance of

past human land use (Fig. S8).

In summary, our results show the value of considering impacts of past human actions to

understand the current organization of biodiversity globally. Previous studies have

documented lasting effects of human land use changes during the last millennia on current

biodiversity patterns[36-38], but this is the first time the signal has been recognized on the

taxonomic differentiation of the largest realms. This would mean that some of the

biogeographical boundaries proposed by Wallace 140 years ago[1] were already showing the

influenced of anthropogenic actions. The human transformation of ecosystems that occurred

in large terrestrial extensions coinciding with the distribution of ancient civilizations, makes

past human land use important along with orographic processes and temperature

seasonality to discern taxonomic differences among the large biogeographical zones of the

world. It is certainly difficult to disentangle direct causal effects from correlative

relationships in non-experimental large-scale studies such as this. We know the same

geographic and environmental characteristics that influenced the distribution across space

and time of mammals likely also influenced the distribution of our ancestors[39]. However, we

considered several of those predictors and still found a global imprint of past human actions

on the patterns of distribution of global diversity. Our results force us to reflect on how the

much more widespread and severe changes that have occurred since the beginning of the

industrial revolution will affect the organization of biodiversity in the future. We show here

the effects of recent human actions are already detectable at the subregional scale, and

possible at the broader biogeographical scale. Although it may seem like an unlikely or

distant future, the possibility that human activities become the main driver, above historical

geomorphological or climatic processes, that shapes Earth´s biodiversity cannot be ignored.

Methods

Generating presence-absence matrices using range distributions. Current and natural

distribution ranges of non-volant terrestrial mammals were obtained from PHYLACINE 1.2

database[25]. This database contains range maps for all 5,831 known mammal species that lived

since the last interglacial (~130,000 years ago until present). Briefly, ranges for current extant

species contained in PHYLACINE are IUCN distribution maps[40] projected to Behrmann

cylindrical equal area rasters with a cell size of 96.5 km by 96.5 km at 30° North and 30° South.

Only current and natural ranges are considered (i.e. introduced, extinct or probably extinct

ranges have been excluded). Moreover, PHYLACINE also contains present natural ranges, which

represent estimates of where species would live if they had never experience anthropogenic

pressures, and – in the case of extinct species – had not gone extinct. We generated two

presence-absences matrices, one for the current and one for the natural ranges, in which every

row represents a grid cell and every column a species. From theses matrices we then excluded

(1) *Homo* species; (2) bats and marine mammals; (3) cells containing less than 50% of land area

to approximate equal-size samples, as a result small islands and coastal cells were not included

in the analyses, and, (4) cells containing fewer than five species to reduce potential distortions

caused by having few taxa[9,10]. These exclusion criteria rendered a total of 14,087 cells

representing 3,960 terrestrial mammal species for the current matrix, and 14,151 cells

representing 4,306 for the natural matrix. With the aim of conducting complementary analyses

to support our results we also generated two additional matrices using distributions of those

extant species that have undergone some modification in their range of distribution due to

anthropogenic pressure, i.e., they have an estimated distribution range in PHYLACINE different

from that of the IUCN, representing 536 species of non-volant terrestrial mammals.

Building biogeographical regionalizations. To each of the presence-absence matrices we

applied a machine-learning algorithm referred to as affinity propagation[26] to build

biogeographical regionalizations at different biogeographical resolution. Affinity propagation

(AP hereafter) is a powerful clustering algorithm extensively used in bioinformatics and

astrophysics and is making inroads in ecology[41] and biogeography[10]. One of its main advantages

is that it can compress massive data sets very efficiently (i.e. with lower error). The algorithm

detects special data points called exemplars, and by a message-passing procedure it iteratively

connects every data point to the exemplar that best represents it until an optimal set of

exemplars and clusters emerges. Contrary to algorithms in which exemplars are found by

randomly choosing an initial subset of data points, AP takes as input measures of 'similarities'

between pairs of data points (grid cells here) and simultaneously considers all the points as

potential exemplars. The optimal set of exemplars is the one for which the sum of similarities of

each point to its exemplar is maximized. Hence, detecting exemplars goes beyond simple

clustering because the exemplars themselves store compressed information[42].

We used a protocol based on a successive application of AP[10] to obtain a biogeographical

upscaling from the smallest possible bioregions (i.e. the highest biogeographical resolution) to

the largest ones. We first used the presence-absence matrix to calculate pairwise similarities

between pairs of cells. As with clustering algorithms, there are many similarities indices or

taxonomic/phylogenetic turnover metrics to choose from, and none can be said to be perfect

from a biogeographical perspective. Here we selected Hellinger distance[43], which is calculated

by first modifying the species-presence data and then computing the Euclidean distance among

pairs of cells based on the modified data[44]. The Hellinger distance is used to avoid the 'double-

zero' problem, i.e. when two sites or grid cells that have no species in common are assigned the

same distance as two sites that share the same species; and the 'species-abundance paradox',

which frequently occurs when two sites share only a small fraction of all the species in the same

regional pool. This is expected to be a particular problem at the margins of biogeographical

regions where sites may be quite different from one another rather than in the centre of a

region where sites are likely to be very similar in their species assemblages[45,46]. We performed

an initial AP analysis involving all grid cells of the similarity matrix. This first AP run generates

the optimal solution of the highest resolution bioregions, while also identifying its exemplars.

We obtained 1,128 clusters and exemplars for the current distributions and 1,053 for the

natural distributions. Then, using the exemplars as the new units of analysis we conducted

again an AP, i.e., we calculated a new similarity matrix and re-run a new AP. This process was

repeated until a small and coherent number of large clusters emerged. Finally, to obtain maps

of each clustering result, we classified each grid cell (row) of every presence-absence matrix

according to the cluster to which they were assigned in its corresponding AP analysis. AP

analyses were performed using the '*APCluster*' package[47] in R Version 3.6.1[48].

Assessing differences in biogeographical configurations. The degree of similarity between

clusters (i.e. regions) of current and natural bioregions was estimated using the Jaccard index.

For that, we used the *cluster_similarity* function of the '*clusteval*' package[49] in R Version 3.6.1[48].

This function computes the similarity between two clusterings of the same data. Jaccard index

ranges between 0 (no similarity) to 1 (perfect match).

Predictors. To each of the cells of the presence-absence matrices we assigned a mean value of

several predictors. We considered predictors previously tested as determinants of

biogeographical assemblages and boundaries[11] related to climatic heterogeneity, orographic

barriers, tectonic movements, and instability of past climate, plus predictors related to

anthropogenic land use over the Holocene. In particular, we tested four climatic variables:

annual total precipitation, mean annual temperature, seasonality in temperature and

seasonality in precipitation. All climatic variables were extracted from the WorldClim dataset[50]

up-scaled at a 96.5 km resolution. These variables represent both average conditions and

variability within years and have been shown to be determinants of vertebrate distributions[51].

Climatic conditions have strongly shifted over the Quaternary period, and have been shown to

play a role in the present-day species distributions, endemism, and assemblages[52-55]. So, to test

for the potential effect of past climate change or stability, we calculated for each cell the

average velocity of climate change since the Last Glacial Maximum (LGM; ~22,000 years ago)

and since the Mid Holocene (MH; ~ 6000 years ago)[54]. To carry out these calculations, we used

the mean annual temperature and annual total precipitation for the MH and the LGM calculated

by means of the Model for Interdisciplinary Research on Climate (MIROC-ESM)[56] and available

in the WorldClim dataset[50]. Mountain ranges represent major barriers to dispersal for most

mammals, whereas plate tectonics are responsible of the long-term isolation of the biotas[11,12].

We use the GTOPO30 to calculate the mean elevation per grid cell. Plate tectonics were

obtained from Bird's dataset[57]. Each grid cell was assigned the tectonic plate to which it

belongs. When a cell was represented by more than one tectonic plate, it was assigned the one

that occupied a greater percentage of the cell. Finally, past anthropogenic land use was

obtained from Ellis et al.' dataset[23]. We chose the more realistic KK_{10} model, which assumes that

humans use land more intensively when population density is high and land scarce[58]. In

counterpoint, the most popular HYDE model omits land-use intensification and predicts that

except for the developed regions of Europe, human use of land was insignificant in every biome

and region before A.D. 1750. We calculated human pressures at four different time spans:

8,000, 5,000, and 2,000 years ago, as well as at the present time, although in the final analyses

only human land use at present and 2,000 years ago were used (see below).

Evaluating potential drivers of the biogeographical configurations. We used random

forest[59,60] models of 5,000 classification trees to assess the factors that may predict the

classification of cells in bioregions at different biogeographical resolutions and to estimate the

relative importance of the predictors. Random forest is a machine learning method based on a

combination of a large set of decision trees. Each tree is trained by selecting a random set of

variables and a random sample from the training dataset (i.e., the calibration data set). The

number of variables randomly sampled as candidates at each split, *mtry*, is a tuning parameter

of random forest models. The default value for random forest classification is the squared root

of the number of variables (i.e., the squared root of 9 resulting in a *mtry* of 3). We also used the

tuneRF function[61] to look for the optimal *mtry* parameter. We ran the code 5 times for each

biogeographic scale and the results varied between 3 and 8, although the trend was always

higher towards 3, which consequently was the *mtry* we finally chose. To calculate the accuracy

of the models we used the out-of-bag (OOB), an estimate of the misclassification rate that

represents and unbiased error rate of the model that is calculated by counting how many cases

in the training set are misclassified and dividing the number by the total number of

observations.

Random forests are able to disentangle interacting effects and identify nonlinear and scale-

dependent relationships that often occur at the scale of the analysis performed here among

multiple correlated predictors[60]. Although random forests are generally assumed to not be

affected by highly correlated predictor variables, we eliminated some predictors showing a

moderate to high correlation ($r > 0.50$, Table S1) as some evidence from genomic studies

suggests that variable importance measures may show a bias towards correlated predictor

variables[62]. So, our final models included seasonality in temperature and precipitation as

representatives of climate heterogeneity; velocity of climate change since the LGM and the MH

to the present as representatives of past climate change; plate tectonics; mean elevation

representing mountains; and human land use at the present and 2,000 years ago. We decided to

not include human land use 8,000 years ago in the models due to its low representation at

global level (Fig. S9). We ran separate models for human land use at present and human land

use 2,000 years ago. Although globally the correlation between both variables is relatively low

(r = 0.29), we made the decision to perform independent models because for certain bioregions

both human land use at present and 2,000 years ago can be highly correlated. We also ran

additional models with the 536 extant species that have undergone some modification in their

distribution range due to anthropogenic pressures. Finally, the algorithm used in the process of

regionalization only uses species presence-absences, which means that humans can only have a

role if present and if causing local extinctions, thereby changing the amount of shared species

between grid cells. At the largest biogeographical scale two bioregions, Madagascar and

Australia, are characterised by an extremely low historical human land use that could influence

results. Therefore, we also ran the models excluding Australia and Madagascar. Random forest

models were computed using the package '*RandomForest*'[61] in R Version 3.6.1[48].

Calculating global and local predictor importance. Random forest also calculates estimates

of global variable importance for classification, which are very useful to interpret the relevance

of variables for the dataset under study. We measured variable importance using the mean

decrease in accuracy: variables with a large mean decrease in accuracy are more important for

classification of the data. This measure is obtained by permuting randomly each variable and

assessing the decrease in classification accuracy of the model[61]. Thus, the more the accuracy of

the random forest decreases due to the permutation (or exclusion) of a single variable, the

more important that variable is considered.

Moreover, the local variable importance defines the importance of a variable in the

classification of a single sample (grid cell here) and therefore shows a direct link between

variables and samples[63]. The local importance score is derived from all trees for which the

sample was not used to train the tree (i.e. its value is OBB). The percentage of correct votes for

the correct class in the permuted OBB data is subtracted from the percentage of votes for the

correct class in the original OOB data to assign a local importance score for the variable for

which the values were permuted. The score reflects the impact on correct classification of a

given sample: negative, 0 (the variable is neutral) and positive. Given that local importance

values are noisier than global importance ones we run the same classification 5 times (5 per

biogeographical scale) and averaged the local importance scores to obtain a robust estimation

of local importance values[63]. Since one value per cell was obtained, these were used to create

maps of the local importance of present and past human land use. However, the values of local

importance do not distinguish between areas where the human land use was high and areas

where it was negligible, i.e., both are equally valuable for the classification analysis. So, to help

in the interpretation we also represent in maps the local importance value for those grid cells

where the percentage of land used was greater than 10%.

References

1. Wallace, A. R. *The geographical distribution of animals* (Harper & Brothers, New York, 1876).

2. von Humboldt, A. *Essai sur la geographie des plantes; accompagné d'un tableau physique des*

régions équinoxales, accompagné d'un tableau physique des régions équinoctiales (Schoel &

Co., Paris, 1806).

3. Ricklefs, R. E. A comprehensive framework for global patterns in biodiversity. *Ecol. Lett.* **7,** 1-

15 (2004).

4. Wiens, J. J. The niche, biogeography and species interactions. *Philos. Trans. R. Soc. Lond. B Biol.*

Sci. **366,** 2336-2350 (2011).

5. Daru, B. H., Elliot, T. L., Park, D. S. & Davies, T. J. Understanding the processes underpinning

patterns of phylogenetic regionalization. *Trends Ecol. Evolut.* **32,** 845-860 (2017).

6. Botkin, D. B. et al*.* Forecasting the effects of global warming on biodiversity. *Bioscience* **57,**

227-236 (2007).

7. Holt, B. et al*.* An update of Wallace´s zoogeographic regions of the world. *Science* **339,** 74–78

(2013).

8. Proches, S. & Ramdhani, S. The world´s zoogeographical regions confirmed by cross-taxon

analyses. *Bioscience* **62,** 260–270 (2012).

9. Kreft, H. & Jetz. W. A framework for delineating biogeographical regions based on species

distributions. *J. Biogeogr.* **37,** 2029-2053 (2010).

10. Rueda, M., Rodriguez, M. A. & Hawkins, B. A. Identifying global zoogeographical regions:

lessons from Wallace. *J. Biogeogr.* **40,** 2215–2225 (2013).

11. Ficetola, G. F., Mazel, F. & Thuiller, W. Global determinants of zoogeographical boundaries.

Nature Ecology & Evolution **1,** 0089 (2017).

12. Lomolino, M. V., Riddle, B. R., Whittaker, R. J. & Brown, J. H. *Biogeography* (Sinauer

Associates, 2010).

13. Riddle, B. R. & Hafner, D. J. Integrating pattern with process at biogeographic boundaries:

the legacy of Wallace. *Ecography* **33,** 321–325 (2010).

14. Barnosky et al. Has the Earth´s sixth extinction already arrived? *Nature* **471**, 51-57 (2011).

15. Ceballos, G., Ehrlich, P. R. & Dirzo, R. Biological annihilation via the ongoing sixth mass

extinction signaled by vertebrate population losses and declines. *Proc. Natl. Acad. Sci. USA*

114, E6089-E6096 (2017).

16. Baiser, B., Olden, J. D., Record, S., Lockwood, J. L. & McKinney, M. L. Pattern and process of

biotic homogenization in the new Pangea. *Proc. Royal Soc. B* **279,** 4772-4777 (2012).

17. Olden, J. D. Biotic homogenization: a new research agenda for conservation biogeography. *J.*

Biogeogr. **33**, 20-27-2039 (2006).

18. Capinha, C. et al. The dispersal of alien species redefines biogeography in the Anthropocene.

Science **348,** 1248-1251 (2015).

19. Bernardo-Madrid, R. et al. Human activity is altering the world´s zoogeographical regions.

Eco Lett, https:/doi.org/10.1111/ele.13321 (2019).

20. Lyons, S. K. et al. Holocene shifts in the assembly of plant and animal communities implicate

human impacts. *Nature* **529,** 80-85 (2016).

21. Sandom, C., Faurby, S., Sandel, B. & Svenning, J-C. Global late Quaternary megafauna

extinctions linked to humans, not climate change. *Proc. Royal Soc. B* **281,** 20133254 (2014).

22. Smith, F., Elliott Smith, R. E., Lyons, S. K. & Payne, J. Body size downgrading of mammals

over the late Quaternary. *Science* **360**, 310-313 (2018).

23. Ellis, E. C. et al. Used planet: a global history. *Proc. Natl. Acad. Sci. USA* **110,** 7978-7985

(2013).

24. Boivin, N. L. et al. Ecological consequences of human niche construction: Examining long-

tern anthropogenic shaping of global species distributions. *Proc. Natl. Acad. Sci. USA* **113**,

6388-6296 (2016).

25. Faurby, S., Davis, M., Pedersen, R. O., Schowannek, S. D., Antonelli, A. & Svenning, J-C.

PHYLACINE 1.2: The Phylogenetic atlas of mammal macroecology. *Ecology* **99**, 2626 (2018).

26. Frey, B. J. & Dueck, D. Clustering by passing messages between data points. *Science* **315,**

972-976 (2007).

27. Faurby, S. & Svenning, J-C. Historic and prehistoric human-driven extinctions have reshaped

global mammal diversity patterns. *Diversity Distrib.* **21**, 1155-1166 (2015).

28. Polaina, E., González-Suárez, M. & Revilla, E. The legacy of past human land use in current

patterns of mammal distribution. *Ecography* **42**, 1623-1635 (2020).

29. Bocquet-Appel, J-P. When the world´s population took off: the springboard of the Neolithic

demographic transition. *Science* **333**, 560-561 (2011).

30. Richard, E. W. A. *Ancient civilizations of the new world: Essays in world history* (Westview

Press, Boulder, Co, 1997).

31. Dunning, N. P., Beach, T. P. & Luzzadder-Beach, S. Kax and kol: Collapse and resilience in

lowland Maya civilization. *Proc Natl Acad Sci USA* **109**, 3652-3657 (2013).

32. Silva, F. et al. Modelling the geographical origin of rice cultivation in Asia using the rice

archaeological database. *PLoS ONE* **10**, e0137024 (2015).

33. Fuller, D. Q. et al. The contribution of rice agriculture and livestock pastoralism to

prehistoric methane levels. An archaeological assessment. *Holocene* **21**, 743-759 (2011).

34. Lewis, S. L. & Maslin, M. A. Defining the Anthropocene. *Nature* **519**, 171-180 (2015).

35. Butzer, K. W. Environmental history in the Mediterranean world: cross-disciplinary

investigation of cause-and-effect for degradation and soil erosion. *J. Archaeol Sci* **32**, 1773-

1800 (2005).

36. Dambrine, E. et al. Present forest biodiversity patterns in France related to former Roman

agriculture. *Ecology* **88,** 1430-1439 (2007).

37. van der Sande, M. et al. A 7000-year history of changing plant trait composition in an

Amazonian landscape; the role of humans and climate. *Ecol. Lett.* **22**, 925-935 (2019).

38. Haberle, S. G. Prehistoric human impact on rainforest biodiversity in highland New Guinea.

Phil. Trans. R. Soc. B **362**, 219-228 (2007).

39. Lomolino, M. V. On teaching "…that grand subject, …". *Front. Biogeogr.* **10**, e37812 (2018).

40. IUCN 2016. IUCN red list of threatened species. Version 2016-3. http://www.iucnredlist.org

41. Cardille, J. A. & Lambois, M. From the redwood forest to the Gulf Stream waters: human

signature nearly ubiquitous in representative US landscapes. *Front. Ecol. Environ.* **8,** 130–134

(2009).

42. Mézard, M. Where are the exemplars? *Science* **315,** 949-951 (2007).

43. Rao, C. R. A review of canonical coordinates and an alternative to correspondences analysis

using Hellinger distance. *Qüestiió* **19,** 23–63 (1995).

44. Legendre, P. & Gallagher, E. D. Ecological meaningful transformations for ordination of

species data. *Oecologia* **129,** 271-280 (2001).

45. Legendre, P. & Legendre, L. *Numerical ecology*, (Elsevier, Amsterdam, 1998).

46. Gagné, S. & Proulx, R. Accurate delineation of biogeographical regions depends on the use of

an appropriate distance measure. *J. Biogeogr.* **36,** 561–567 (2009).

47. Bodenhofer, U., Kothmeier, A. & Hochreiter, S. APCluster: a R package for affinity

propagation clustering. *Bioinformatics* **27,** 2463–2464 (2011).

48. R Core Team. R: *A Language and Environment for Statistical Computing, 3.6.1* (R Foundation

for Statistical Computing, 2019).

49. Ramey, J. A. Clusteval: Evaluation of Clustering Algorithms. R package version 0.1. https:

//CRAN-R-project.org/package=clusteval (2012).

50. Fick, S. E. & Hijmans, R. J. Worldclim 2: New 1-km spatial resolution climate surfaces for

global land areas. *Int. J. Climat.* **37,** 4302-4315 (2017).

51. Sexton, J. P., McIntyre, P. J., Angert, A. L. & Rice, K. J. Evolution and ecology of species range

limits. *Ann. Rev. Ecol. Evol. Syst.* **40,** 415-436 (2009).

52. Graham, R. W. et al. Spatial response of mammals to Late Quaternary environmental

fluctuations. *Science* **272,** 1601-1606 (1996).

53. Davis, M. B. & Shaw, R. G. Range shifts and adaptive responses to Quaternary climate

change. *Science* **292,** 673–679 (2001).

54. Sandel, B. et al. The influence of late Quaternary climate-change velocity on species

endemism. *Science* **334,** 660-664 (2011).

55. Araujo, M. B. et al. Quaternary climate changes explain diversity among reptiles and

amphibians. *Ecography* **31,** 8-15 (2008).

56. Watanabe, S. et al. MIROC-ESM 2010: model description and basic results of CMIP5-20c3m

experiments. *Geosci. Model Dev.* **4,** 845–872 (2011).

57. Bird, P. An updated digital model of plate boundaries. *Geochemistry, Geophysics, Geosystems*

4, 1027, doi:10.1029/2001GC000252 (2003)

58. Kaplan, J. O. et al. Holocene carbon emissions as a result of anthropogenic land cover

change. *Holocene* **21,** 775–791 (2011).

59. Breiman, L. Random forests. *Machine Learning* **45,** 5–32 (2001).

60. Cutler, D. R. et al. Random forests for classification in ecology. *Ecology* **88,** 2783–2792

(2007).

61. Liaw, A. & Wiener, M. Classification and regression by RandomForest. *R News* **2,** 18–22

(2002).

62. Nicodemus, K. K. et al. The behaviour of random forest permutation-based variable

importance measures under predictor correlation. *BMC Bioinformatics* **11**, 110 (2010).

63. Touw, W. G. et al. Data mining in the Life Sciences with Random Forest: a walk in the park or

lost in the jungle. *Briefings in Bioinformatics* **14**, 315-326 (2012).

Acknowledgements

This work was supported by the project 2020/125 (Programa Operativo FEDER Andalucía 2014-

2020) to MR, and by CGL2017-83045-R AEI/FEDER EU (Agencia Estatal de Investigación from

the Ministry of Economy, Industry and Competitiveness, Spain co-financed with FEDER) to ER.

Author contributions

MR, MG-S and ER designed the study; MR analyzed the data and led the writing, with input and

edits from all authors.

Competing interests

The authors declare no competing interests.

Material and Correspondence

Marta Rueda (mruedag@us.es)

Fig. 1. Current and natural broader regions or realms. Only results for the broader regions are shown here: 9 current bioregions (a), and 8 natural bioregions (b). The greatest differences between both biogeographical configurations are more relevant in the Northern Hemisphere, where for the natural bioregions we found that the Palearctic and the Nearctic are united forming a great extension, the typical biographical boundary between the Nearctic and the Neotropical realms disappears and extends towards the north, and India is no longer part of the Oriental region and is included instead within the 'Saharo-Arabian' region.

CURRENT

NATURAL

a) HUMAN LAND USE AT PRESENT

Plate tectonics	619.7
Temp seasonality	341.3
Mean elevation	187.6
Land Use present	186.7
VCC-MH mean T	166.8
VCC-LGM annual P	162.6
Precip seasonality	147.6
VCC-MH annual P	140.6
VCC-LGM mean T	96.8

OBB = 2.85%

b) HUMAN LAND USE AT PRESENT

Plate tectonics	498.3
Temp seasonality	345.2
VCC-LGM annual P	168.3
Precip seasonality	167.3
Mean elevation	156.4
Land Use present	152.1
VCC-MH annual P	139.6
VCC-LGM mean T	137.5
VCC-MH mean T	99.3

OBB = 2.56%

c) HUMAN LAND USE 2000 YA

Plate tectonics	656.9
Temp seasonality	328.7
Land Use 2000 ya	244.7
Mean elevation	174.3
VCC-LGM annual P	166.9
Precip seasonality	133.2
VCC-MH mean T	117.2
VCC-MH annual P	103.1
VCC-LGM mean T	86.6

OBB = 2.43%

d) HUMAN LAND USE 2000 YA

Plate tectonics	494.6
Temp seasonality	350.3
VCC-LGM annual P	197.8
Precip seasonality	176.1
Land Use 2000 ya	165.4
Mean elevation	154.0
VCC-LGM mean T	136.3
VCC-MH annual P	115.4
VCC-MH mean T	101.8

OBB = 2.63%

Fig. 2. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural broader bioregions. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. Importance was measured by the drop-in classification accuracy after predictor randomization in random forests of 5,000 trees. Higher values of mean decreased in accuracy indicate variables that are more important to the classification. OBB (Out-of-bag) represents the percentage of cells misclassified in each model. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

9 CURRENT BIOREGIONS   **8 NATURAL BIOREGIONS**

Fig. 3. Venn diagrams describing the accuracy in classification for models including the three most important drivers of the current and natural broader bioregions. Values indicate the percentage of cells correctly classified (1-OBB error) in their bioregion, and are obtained from running random forest classification models for each individual variable, by pairs and for the three most important factors together.

Fig. 4. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural medium-sized bioregions or subregions. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. Specifications as in Fig. 2.

Fig. 5. Maps of the local importance of present (left panels) and past (right panels)

human land use for current (the four maps above) and natural (the four maps below)

broader bioregions. The local importance shows the importance of the variable for the

classification of a single sample (grid cells here). The score of the legends shows the impact on

correct classification of grid cells, with blue colors indicating from negative to 0 values

(incorrect to neutral classification), and yellow, orange and red indicating positive values

613   (correct classification). Since the values of local importance do not distinguish cells where the

614   human land use was significant and areas where it was negligible, i.e., both can be important for

615   the classification process, we have also mapped the importance values for those grid cells

616   where the percentage of land use at present and 2,000 years ago was >10% (c, d, g, h). In these

617   maps, white cells imply human land use values <10%.

Fig. 6. Maps of the local importance of present (left panels) and past (right panels) human land use for current and natural medium-sized subregions. Specifications as in Fig. 5.

625 **Table 1**. **Values of Jaccard index between current and natural hierarchical bioregions**.

626 Jaccard index assesses the degree of similarity between clusters (bioregions) and ranges

627 between 0 (no similarity) and 1 (perfect match). The number of bioregions obtained during

628 each bioregionalization processes is also provided.

CURRENT	Jaccard coefficient	NATURAL
9 broader realms	0.57	8 broader realms
27 subregions	0.58	25 subregions
141 small bioregions	0.42	121 small bioregions
1,128 smaller bioregions	0.28	1,051 smaller bioregions

Supplementary Materials for

Global biogeographical regions reveal a signal of past human impacts

Marta Rueda*, Manuela González-Suárez, Eloy Revilla

Corresponding author. Email: mruedag@us.es

This PDF files includes:

a) CURRENT HIERARCHICAL BIOREGIONS

b) NATURAL HIERARCHICAL BIOREGIONS

Fig. S1. Nested global bioregionalizations for non-volant terrestrial mammals. The hierarchical application of the affinity propagation algorithm resulted in a nested global bioregionalization containing for the current hierarchical bioregions (a): 9 (filled colours), 27 (red boundaries), 141 (black boundaries) and 1128 (grey boundaries) bioregions from the largest extant realms to the smallest obtained bioregions; and for the natural hierarchical bioregions (b): 8 (filled colours), 25 (red boundaries), 121 (black boundaries), and 1,051 (grey boundaries) bioregions.

a) CURRENT BIOREGIONS

b) NATURAL BIOREGIONS

Fig. S2. Broader regions and medium-sized bioregions for the 536 extant non-volant terrestrial mammals whose distribution ranges have shifted due to human actions. The affinity propagation clustering algorithm resulted in 6 broader bioregions (filled colours), and 18 subregions (black boundaries) for (a) Current bioregions; and 6 broader bioregions (filled colours) and 17 subregions (black boundaries) for (b) Natural bioregions. Although not represented in the map, the clustering algorithm resulted in 93 and 1,333 smaller regions for the current, and 98 and 997 smaller regions for the natural bioregions.

Fig. S3. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural broader bioregions without Australia and Madagascar. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. Importance was measured by the drop-in classification accuracy after predictor randomization in random forests of 5,000 trees. Higher values of mean decreased in accuracy indicate variables that are more important to the classification. OBB (Out-of-bag) represents the percentage of cells misclassified in each model. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

Fig. S4. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural broader bioregions for the 536 extant non-volant terrestrial mammals whose distribution ranges have shifted due to human actions. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

CURRENT   **NATURAL**

Fig. S5. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural medium-sized subregions for the 536 extant non-volant terrestrial mammals whose distribution ranges have shifted due to human actions. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

Fig. S6. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural small subregions. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

Fig. S7. Ranking of importance values for the drivers of taxonomic differentiation for the current and natural smaller subregions. Above panels show importance values from models with human land use at present for (a) current and (b) natural bioregions, while the panels below show importance values from models with human land use 2,000 years ago for (c) current and (d) natural bioregions. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

Fig. S8. Ranking of importance values for the drivers of taxonomic differentiation for the current (a) and natural (b) medium-sized subregions when subregions without human land use 2,000 years ago are removed from the analysis. Subregions removed were Madagascar, the two Australian subregions, the Sahara Desert (only for the current subregions), the Tibetan plateau, and the American Arctic. VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; annual P = annual precipitation; mean T = mean annual temperature; ya = years ago.

Fig. S9. Percentage of human land used at present, 2,000 and 8,000 years ago for broader regions and medium-sized bioregions.

Table S1. Jaccard index for current and natural hierarchical bioregions built with the 536 extant mammals whose distributions ranges have shifted due to human impact. Jaccard index assesses the degree of similarity between clusters (bioregions) and ranges between 0 (no similarity) and 1 (perfect match). The number of bioregions obtained during each of the bioregionalization processes is also provided.

CURRENT	Jaccard coefficient	NATURAL
6 broader realms	0.48	6 broader realms
18 subregions	0.33	17 subregions
93 small bioregions	0.20	98 small bioregions
1,333 smaller bioregions	0.19	997 smaller bioregions

Table S2. Pearson's correlations between continuous determinants. MAT = Mean annual temperature; VCC = velocity of climate change; MH = Mid-Holocene; LGM = Last Glacial Maximum; ya = years ago.

	Elevation	MAT	Annual precip	Temp season	Precipit season	VCC-LGM Temp	VCC-MH Temp	VCC-LGM Precip	VCC-MH Precip	Land use present	Land use 2,000 ya	Land use 5,000 ya	Land use 8,000 ya
Elevation	1												
MAT	-0.21	1											
Annual precipitation	-0.15	0.35	1										
Temp seasonality	-0.007	-0.84	-0.55	1									
Precipit seasonality	0.21	0.30	-0.18	-0.18	1								
VCC-LGM Temp	-0.27	-0.41	-0.12	0.44	-0.29	1							
VCC-MH Temp	-0.21	0.13	0.03	-0.06	0.09	0.32	1						
VCC-LGM Precip	-0.17	-0.16	0.33	0.04	-0.19	0.37	0.04	1					
VCC-MH Precip	-0.09	0.36	0.24	-0.36	0.37	-0.17	0.19	0.11	1				
Land use present	0.07	0.20	-0.06	-0.10	0.13	-0.08	-0.05	-0.12	0.13	1			
Land use 2,000 ya	-0.0007	0.12	0.07	-0.15	0.13	-0.01	-0.01	-0.08	0.04	0.29	1		
Land use 5,000 ya	0.016	0.07	0.05	-0.06	0.11	-0.04	-0.03	-0.04	-0.01	0.22	0.87	1	
Land use 8,000 ya	0.006	0.06	0.008	-0.04	0.08	-0.01	-0.03	-0.04	-0.03	0.13	0.63	0.86	1