Supervised dimension reduction for large-scale "omics" data with censored survival outcomes under possible non-proportional hazards

Lauren Spirko-Burns[1], Karthik Devarajan[2]

[1]*Department of Statistical Science, Temple University, Philadelphia, PA*
[2]*Department of Biostatistics & Bioinformatics, Fox Chase Cancer Center,*
Temple University Health System, Philadelphia, PA

Abstract

The past two decades have witnessed significant advances in high-throughput "omics" technologies such as genomics, proteomics, metabolomics, transcriptomics and radiomics. These technologies have enabled the simultaneous measurement of the expression levels of tens of thousands of "omic" features from individual patient samples and have generated enormous amounts of data that require analysis and interpretation. One specific area of interest has been in studying the relationship between these features and patient outcomes such as overall and recurrence-free survival with the goal of developing a predictive "omics" profile. In this paper, we propose a supervised dimension reduction method for feature selection and survival prediction. Our approach utilizes continuum power regression - a framework that includes ordinary least squares, principal components regression and partial least squares - in conjunction with the parametric or semi-parametric accelerated failure time model and enables feature selection under possible non-proportional hazards. The proposed approach can handle censored observations using robust Buckley-James estimation in this high-dimensional setting and the parametric version employs the flexible generalized F model that encompasses a wide spectrum of well known survival models. We evaluate the predictive performance of our methods via extensive simulation studies and compare it to existing methods using publicly available data sets in cancer genomics.

Keywords: high-dimensional data, cancer genomics, high-throughput "omics", accelerated failure time model, continuum power regression, generalized F model, non-proportional hazards, censored survival data

1 Introduction

Advances in high-throughput technologies in the past two decades have enabled large-scale "omics" studies that generate enormous amounts of data that are measured on a variety of scales. Examples include, but are not limited to, genomic studies such as next-generation sequencing, methylation, allele-specific expression, microarrays, and DNA copy number as well as studies involving genome-wide association, proteomics, metabolomics, transcriptomics and radiomics. Genomic studies, for instance, enable the simultaneous measurement of the expression profiles of tens of thousands of genomic features, often from a relatively small number of individual patient samples. Such studies result in massive quantities of data requiring analysis and interpretation while offering tremendous potential for growth in our understanding of the pathophysiology of many diseases. When information on an outcome

variable such as time to an event (or survival time) is available, one of the goals of an investigator is to understand how the expression levels of genomic features, and clinical and demographic variables (covariates) relate to an individual's survival over the course of a disease. The number of covariates (n) far exceeds the number of observations (p), typically, in these large-scale genomic studies. With the tremendous volume of information available, the investigator can now estimate and attempt to understand the effects of specific genomic features on various diseases with the ultimate goal of developing a prognostic profile of patient survival. In this context, biomarker discovery poses many challenges and plays a pivotal role in the search for more precise treatments. The role and significance of the analysis of time-to-event data in cancer research cannot be overstated where current efforts focus on predicting therapeutic responses of patients with a view to personalizing cancer treatment.

The ill-conditioned problem of predicting the survival probability when $p >> n$ is further compounded by the presence of censored survival times. In this high-dimensional setting, one is often interested in building a genomic profile that is predictive of the survival probability for a new patient. The Cox proportional hazards (PH) model is the most celebrated and widely used statistical model linking survival time to covariates (Cox, 1972). It is a multiplicative hazards model that implies constant hazard ratio (HR); that is, it postulates that the risk (or hazard) of death of an individual given their covariates is simply proportional to their baseline risk in the absence of any covariate. While this model has proved to be very useful in practice due to its simplicity and interpretability, the assumption of constant HR has been shown to be invalid in a variety of situations in medical studies (Devarajan & Ebrahimi, 2011; Peri et al., 2013). When applied to our problem, the PH model would implicitly assume a constant effect of genomic feature expression on survival over the entire period of follow-up in a study, a supposition that is neither verifiable nor likely for each feature. For example, non-proportional hazards (NPH) can occur when the effect of a genomic feature increases or decreases over time leading to converging or diverging hazards (CH or DH), a situation that cannot be handled by the PH model (Bhattacharjee et al., 2001; Xu et al., 2005; Dunkler et al., 2010; Rouam et al., 2011). In addition, NPH can result from model misspecification such as from omitting a strong clinical covariate (for instance, age at diagnosis or stage of disease) or another genomic feature. Another scenario encountered in practice is the case of dependence between covariates and the censoring time distribution (Chen et al., 2002). These scenarios require more general survival models that consider time-varying covariate effects. Examples of such models include the Accelerated Failure Time (AFT) and Proportional Odds (PO) models, among others (Buckley & James, 1979; Jin et al., 2006; Martinussen & Scheike, 2006; Yang & Prentice, 2005; Devarajan & Ebrahimi, 2011). The AFT model is a censored linear regression model in which the covariates cause an acceleration (linear transformation) of the time scale while the PO model postulates that the odds of death for an individual, given their covariates, is simply proportional to their baseline odds - a situation typically encountered when the effect of a genomic feature decreases with time leading to diverging hazards. Unlike PH, these models do not imply a constant HR and, interestingly, both PH and PO models intersect with the AFT model. Moreover, the AFT model can accommodate a variety of well known survival time distributions - such as the lognormal, log-logistic, Weibull and exponential, to name a few - useful for modeling censored survival data in practice or accommodate a completely distribution-free approach with no prior assumption on the data generating mechanism (Kalbfleisch & Prentice, 2002; Jin et

al., 2006). These attractive properties provide modeling flexibility and make the AFT model a versatile alternative useful for handling a variety of data structures.

As evidenced by the following literature survey, very little research has been done to account for the time-varying effect of genomic features or to study the consequences of NPH on feature selection and survival prediction, despite its clear importance in translational medicine. Within a broader context, these shortcomings extend to the many types of high-throughput "omics" studies outlined earlier. In this paper, we generically use the term "omics" to represent this variety and the term feature to denote the appropriate "omic" feature of interest. The rest of the paper is organized as follows. In §2, we survey existing methods for supervised dimension reduction within the context of high-throughput "omics" data and censored survival outcomes, and discuss their weaknesses. Section 3 begins with a motivation of the need for a flexible method using real-life cancer genomic data. We then propose an approach that combines continuum power regression with the parametric or semi-parametric AFT model. In §4, we develop a prognostic index and survival prediction algorithm using this approach. Section 5 is devoted to extensive simulation studies for evaluating the proposed methods while §6 focuses on application of these methods to several publicly available data sets in cancer genomics. Last but not least, §7 provides a summary and discussion including future work.

2 A brief survey of existing methods and their limitations

A variety of methods are currently available in the literature for handling a large number of features in conjunction with censored survival outcomes. These include methods based on principal components regression (PCR) (Li & Li, 2004; Bair et al., 2006; Ma et al., 2006), partial least squares (PLS) (Park et al., 2002; Li & Gui, 2004; Nguyen & Rocke, 2002; Nguyen, 2005; Nygard et al., 2008; Boulesteix & Strimmer, 2007; Bastien et al., 2015) and regularization such as the ridge regression, least absolute shrinkage and selection operator (LASSO), least angle regression, elastic net or related variants (Verweij & van Houwelingen, 1994; Tibshirani, 1997; Fan & Li, 2002; Gui & Li, 2005; Park and Hastie, 2006; Zhang & Lu, 2007; Segal, 2008; Zou, 2008; Engler & Yi, 2009; Wang et al., 2009; Kaneko et al., 2012). Other available methods in this context include those based on boosting (Li & Luan, 2005; Wei & Li, 2007), sure screening procedures (Fan et al., 2010), Cox assisted clustering (Eng & Hanlon, 2012), networks (Zhang et al., 2013), kernel methods (Li & Luan, 2003) and nested cross-validation (Laimighofer et al., 2016).

A comparison of various existing methods has revealed that those based on a linear combination of features (such as PCR and PLS) or regularization (such as LASSO etc.) showed overall superior performance (Bovelstad et al., 2007; van Wieringen et al., 2009; Witten & Tibshirani, 2008). Methods based on PLS and PCR typically utilize all features for prediction and cannot directly specify relevant features that are associated with survival. Regularization methods generally perform well in this setting by identifying unimportant features from the large number of features present by shrinking their coefficients to exactly zero. However, a method such as LASSO suffers from some fundamental limitations due to

the L_1 penalty. For instance, the number of non-zero coefficients can be at most n, i.e., the number of features that can be selected by LASSO is bounded by the sample size of the data set (Rosset et al., 2004). In large-scale genomic studies, this can lead to the unrealistic conclusion that no more than n genomic features are relevant to survival in a complex biological process where $p \gg n$ are actually present. This is further compounded by the relatively small number of observations often seen in these biomedical studies. Moreover, the expression levels of features sharing a particular biological pathway can be highly correlated. It is therefore desirable to have a method that automatically selects the entire set of correlated features; however, LASSO can usually select only one feature in this situation. On the other hand, a method such as ridge regression necessarily selects all features in a data set. These issues can pose serious problems particularly when dealing with the ultra high-dimensional data sets obtained in modern "omics" studies (Wang et al., 2008).

An inherent weakness of these methods is the assumption of PH in their formulation which does not permit the incorporation and, therefore, the detection of time-dependent covariate effects. However, there exist methods based on alternate survival models such as the AFT (Huang & Harrington, 2004; Huang et al., 2006; Wang & Leng, 2007; Datta et al., 2007; Leng & Ma, 2007; Wang et al., 2008; Luan & Li, 2008; Johnson, 2008a,b, 2009; Johnson et al., 2008; Cai et al., 2009; Wang & Wang, 2010; Engler & Yi, 2009; Liu et al., 2010; Devarajan et al., 2010), PO (Lu & Li, 2008), non-linear transformation (Lu & Li, 2008), additive hazards (Ma et al., 2006) or that are model-free (Van Belle et al., 2011; Geng et al., 2014; Pang et al., 2012). Huang et al. (2006) combine the AFT model with LASSO or threshold-gradient-directed regularization (TGDR) using Stute's estimator (Stute, 1993), thereby providing flexible methods for handling NPH and high-dimensionality. In addition to known limitations, their LASSO approach has been known to result in inferior prediction accuracy in empirical studies. Furthermore, TGDR is sensitive to the choice of a parameter value that could significantly alter the number of features selected and thus lead to overestimation of the number of non-zero coefficients, potentially further reducing the number of features selected (Wang & Wang, 2010). Datta et al. (2007) developed an approach that combines standard PLS or LASSO with the AFT model after mean imputation of censored observations. This approach does not improve upon these existing methods and suffers from the limitations of LASSO. Wang et al. (2008) proposed a doubly penalized method based on the AFT model for estimation, feature selection and survival prediction by extending elastic net regression for linear models to censored survival data. Unlike LASSO, this approach can select an arbitrary number of highly correlated features with non-zero coefficients; however, it involves the selection of tuning parameters and can be computationally slow. Engler & Yi (2009) proposed an elastic net approach with mean imputation in conjunction with the Cox PH or AFT model and showed that the AFT version showed better performance. Existing model-free methods provide a flexible alternative that can account for linear and non-linear covariate effects; however, they tend to be computationally infeasible and typically require the choice of various tuning parameters.

3 Motivation for the proposed methods

In the high-dimensional setting, incorporating time-dependent covariates in the Cox PH model, use of stratification or separate modeling for different time periods in order to account for NPH are prohibitive and infeasible. As outlined in the literature survey above, computational speed and/or infeasibility, number and choice of tuning parameters, restrictions on the number of "omic" features that can be selected, over-estimation of the number of relevant features and poor predictive performance are some of the noteworthy limitations when regularization is used on more general survival models that account for NPH or when a completely model-free method (such as SVM, random forests etc.) is used. Unlike the PH model, the AFT is built on the linear regression model for censored survival data and is a viable alternative to it since it directly models survival time and, thus, has a simpler and more intuitive interpretation. More importantly, it allows crossing hazard and survival curves, a useful property for modeling large-scale "omics" data with tens of thousands of features. In this paper, we adopt a more pragmatic approach for initially identifying the number of relevant features in a data set by simultaneously utilizing model significance and model fit based on different criteria. In addition, we adjust for potential confounders such as age of diagnosis and stage of disease with the goal of further eliminating spurious features. Such supervised marginal screening ensures that each feature selected for inclusion in the development of a prediction model actually fits the model of interest *and* has a statistically significant effect on survival.

Let Y denote the survival time of a typical subject in the study, the length of time entry into the study until a prescribed endpoint is attained. This endpoint may be the onset of a disease or event associated with it, or death itself. In addition, we let C be the duration of observation of the subject, i.e. the time from entry into the study until removal. The random variable C is referred to as a censoring variable. In general, we assume that both Y and C are non-negative random variables of which only the first one to occur is observed. Thus, an observation consists of the pair (T, δ), where $T = \min(Y, C)$ and $\delta = I(T = Y)$. We also have data on p covariates from each subject. It is assumed that censoring is non-informative, i.e. the survival time Y and the censoring mechanism C are independent, and that the covariates do not provide information about the censoring time C. Survival data usually consists of N samples, each containing the triple $(T_i, \delta_i, \mathbf{z}_i)$ for $i = 1, \cdots, n$, where $\mathbf{z}_i = (z_{i1}, \cdots, z_{ip})$ is the covariate vector or profile of the i-th subject, T_i is the survival time if $\delta_i = 1$ and it is the right censored time if $\delta_i = 0$. The AFT model postulates a log-linear relationship between time and covariates given by,

$$\log Y = \boldsymbol{\beta}'\mathbf{z} + \sigma\epsilon, \tag{3.1}$$

where $\boldsymbol{\beta}'$ is a vector of regression coefficients, $\mathbf{z}$ is the vector of covariates, σ is a scale parameter and ϵ is the error term whose distribution is either pre-specified or is left completely unspecified, thus resulting in parametric or semi-parametric versions of the model (Kalbfleisch & Prentice, 2002; Jin et al., 2006). The intercept can be absorbed into $\boldsymbol{\beta}'$ and in the semi-parametric version, $\sigma \equiv 1$ without loss of generality. Its log-linear form enables the measurements of the direct effect of features on survival time instead of the hazard; moreover, the regression coefficients can be interpreted in a similar fashion to that of multiple linear regression. In this model, effect size is measured as the ratio of expected survival times

between two groups, say, patients exhibiting low and high expression of a particular feature or a set of features. In the clinical setting, it quantifies the effect of a feature on the expected duration of illness for a patient. This has lead many prominent statisticians, most notably Sir D. R. Cox, to observe that the AFT model and its estimated regression coefficients to have a rather direct physical interpretation' (Reid, 1994). Moreover, it is well-known that the PH and AFT models cannot simultaneously hold except in the case of extreme value error distributions. Therefore, the AFT model assumptions can hold when the PH model assumptions fail.

The semi-parametric AFT ($sAFT$) model is particularly attractive due to its distribution-free nature. Rank-based inference for this model is described in Jin et al. (2003), and regularized estimation is described in Cai et al. (2009) for high-dimensional data. An iterative solution has been developed to estimate the regression parameters (Jin et al., 2006). This procedure is based on the least-squares principle while accounting for censoring; however, it is computationally slow which can be problematic in the high-dimensional setting. The scope and applicability of AFT models can be significantly broadened by use of the generalized F distribution ($GenF$) (Ciampi et al. (1986) and more recently by Cox (2008)). $GenF$ has the form seen in Equation 4.4. Estimation for this model is based on maximum likelihood and, thus, offers a flexible and computationally efficient alternative to the $sAFT$ model. Although $GenF$ spans a variety of well known and lesser known models that are appropriate for modeling survival data, it has received little recognition in the literature. Its benefit lies in its umbrella structure and it includes the Weibull (exponential), generalized gamma (Weibull, gamma, Maxwell-Boltzmann, generalized normal, half-normal, Chi and Raleigh), log-normal and log-logistic models, among many others and provides a flexible approach to modeling patient survival in conjunction with large-scale "omics" data. There are several advantages to using $GenF$. As alluded to in §1, the AFT model intersects with the PH and PO models when the underlying data distribution is Weibull and log-logistic, respectively. The Weibull model with its monotonic hazard function and the log-normal model due to its mathematical intractability in dealing with censored observations offer limited potential for modeling survival data. Although the log-logistic model is similar in shape to the log-normal, its non-monotonic hazard function allows hazard curves to converge with time thereby incorporating a particular type of NPH and making it suitable for modeling cancer survival. It can be used if the course of the disease is such that mortality peaks after a finite time period and then slowly declines.

We motivate the utility of the AFT model for our problem using three data sets from large-scale cancer genomic studies that are detailed in §7.1. In this analysis, we fit semi-parametric PH, PO and AFT models to each feature, after adjusting for clinical covariates such as age at diagnosis and stage of cancer, and evaluate their goodness-of-fit (GOF) using appropriate methods (Therneau & Grambsch, 1994; Martinussen & Scheike, 2006; and Novak, 2010). For each model, the q-value method was used to account for multiple testing (Storey, 2002). The goal is to identify features that exhibit some form of NPH, thus demonstrating the need for alternatives to the PH model and, in particular, providing the rationale for a flexible model like AFT.

The results are summarized in Table 1 where A and B refer to sets of features for which the PH and PO model do not fit, respectively, and C refers to the set of features for which the AFT model fits, at the 5% significance level. Typically, there is a large number of features

for which the PH or PO model does not fit. More importantly, there is a significantly large fraction of features for which the AFT model fits (median of 95%). After correction for multiple testing, these observations are further corroborated by the corresponding q-values where a smaller value indicates evidence of lack of fit for a given model. The intersections of these sets is particularly revealing where we observe that the AFT model fits a large fraction of features for which the PH or PO model do not provide a good fit (median of 94%). Thus, it would be beneficial to develop methods based on the more general AFT model, which overlaps with the PH and PO models, due to its inherent ability to account for crossing hazards.

Table 1: Summary of model fits

Data Set	HNSCC	GBM	Ovarian cancer	Oral cancer
Number of features	19,341	9,452	32,575	12,776
Number of observations	221	280	276	86
% Censored	62%	26%	59%	59%
A: PH (lack of fit)	752 (4%)^a	217 (2%)^c	4,617 (14%)^b	1,810 (14%)^b
B: PO (lack of fit)	1,992 (10%)^b	744 (8%)^b	3,004 (9%)^b	1,232 (10%)^b
C: AFT (good fit)	19,090 (99%)^a	9,380 (99%)^a	29,902 (92%)^c	9,753 (76%)^b
A ∩ C	733	212	4,224	1,379
B ∩ C	1,966	725	2,655	1,023
A ∩ B (∩ C)	464 (451)	47 (46)	1,892 (1,685)	863 (694)

q-value: $0.99^a, \leq 0.25^b, \approx 0.5^c$

4 Continuum power regression for large-scale "omics" data

We develop analytical methods for large-scale "omics" data using continuum power regression (CPR) - a unified framework for supervised dimension reduction - in conjunction with the AFT model. CPR embeds a spectrum of regression methods into a single framework that includes well known methods such as ordinary least squares (OLS), partial least squares (PLS) and principal components regression (PCR) as special cases. Stone & Brooks (1990) first proposed continuum regression (CR) and showed that OLS, PLS and PCR differed only in the target quantity being maximized in the process of extracting latent components that are linear combinations of these high-dimensional covariates. CR aims to maximize a quantity that includes the variation in covariates as well as the correlation of response with covariates, the relative proportions of which are controlled via a single parameter γ. At the extremes of this continuum, OLS maximizes correlation and PCR extracts orthogonal components by maximizing variance, while PLS lies in-between and maximizes the covariance

between response and covariates. The numerical instability suffered by OLS due to multi-collinearity and high-dimensionality are circumvented by the unsupervised and supervised approaches provided by PCR and PLS, respectively, while the choice of γ provides further modeling flexibility.

Given an $n \times p$ matrix $\mathbf{Z}$ of predictors and an n-vector $\mathbf{t}$ of quantitative responses of an outcome, the objective function for constructing reduced components in CR can be expressed in terms of the objective functions for OLS (correlation, R^2), PLS (covariance, Cov) and PCR (variance, Var) as

$$\ell_\gamma(\boldsymbol{\nu}) = g_\gamma(R^2(\mathbf{Z}\boldsymbol{\nu}, \mathbf{t}), Var(\mathbf{Z}\boldsymbol{\nu})) = R^2(\mathbf{Z}\boldsymbol{\nu}, \mathbf{t})[Var(\mathbf{Z}\boldsymbol{\nu})]^\gamma \propto Cov(\mathbf{Z}\boldsymbol{\nu}, \mathbf{t})^2[Var(\mathbf{Z}\boldsymbol{\nu})]^{\gamma-1}.$$

(4.1)

The optimization criterion is $\max_{||\boldsymbol{\nu}||=1} \ell_\gamma(\boldsymbol{\nu})$ subject to $\boldsymbol{\nu}'\mathbf{S}\boldsymbol{\nu}_j = 0, j = 1, ..., K$, where $\mathbf{S} = \mathbf{Z}'\mathbf{Z}$ is the covariance of $\mathbf{Z}$, the columns of $\boldsymbol{\nu}$ are weight vectors, $\gamma \geq 0$ and K is the number of components. CR reduces to OLS ($\gamma = 0$), PLS ($\gamma = 1$) and PCR ($\gamma \to \infty$) and can be shown to be closely related to ridge regression (Sundberg, 2002).

CPR is a variant of CR that is defined by the algorithm and not as the solution to the optimization problem in eqn. (3.2). In CPR, the PLS estimate $\boldsymbol{\tau} \propto \mathbf{Z}\mathbf{Z}'\mathbf{t}$ is generalized to $\boldsymbol{\tau} \propto (\mathbf{Z}\mathbf{Z}'\mathbf{t})^\gamma$ for $\gamma \geq 0$ where $\mathbf{Z}$ is modified into its powered version $\mathbf{Z}^{(\gamma)}$ via the SVD of $\mathbf{Z}$, i.e., $\mathbf{Z}^{(\gamma)} = UL^{\gamma/2}V'$. CPR simplifies similar methods by requiring only one SVD after which standard PLS can be applied to $\mathbf{Z}^{(\gamma)}$, thus significantly improving computational speed and ease of interpretation (de Jong et al., 2001; Lorber et al., 1987). CPR coincides with CR for the special cases ($\gamma = 0, 1$ and ∞). It has been suggested that the continuity parameter γ and dimensionality K can play similar roles (Stone & Brooks, 1990; Frank & Friedman, 1993). There is also evidence to suggest that it is sufficient to consider only the three important special cases (OLS, PLS and PCR) and that the continuum may be unnecessary in CR under certain conditions (Chen & Cook, 2010). Given the fact that at three points CR is identical to CPR and that K and γ have similar effects, the simplicity, modeling flexibility and speed of CPR confer significant advantages over CR. In general, PLS requires fewer components than PCR; this is because the components from the latter need not necessarily be correlated with time-to-event, whereas all PLS components must be. PLS may be regarded as a compromise between OLS and PCR.

4.1 The CPR-AFT model

For a given application, CPR has the potential to offer insight into the underlying model. The ability of AFT to incorporate crossing hazard curves offers unparalleled flexibility for modeling large-scale "omics" data. As both CPR and AFT fall within the linear models framework, it seems natural to consider a hybrid model that combines their strengths. By combining AFT with CPR in a two-step procedure we develop supervised dimension reduction methods, which we call Adjusted CPR-AFT or (A)CPR-AFT, that adjust for censored observations. (A)CPR-AFT provide a powerful array of solutions for this problem and enables identification of an "omic" profile that is predictive of a patient's response to a specific treatment under a variety of scenarios encountered in practice.

(A)CPR-AFT has a distinct advantage over other methods in the literature because it directly addresses the three main issues with the application of survival analysis to "omics"

data. First, it addresses the issue of high-dimensionality using CPR, reducing the number of "omic" features into a smaller number of CPR components that are linear combinations of these features. Second, it addresses the issue of NPH by using the AFT model, a model that does not assume PH but partly overlaps with the PH model. Lastly, it addresses the issue of censoring by imputing the censored observations using the extracted CPR components and the fitted AFT model. In the literature survey in §2 we noted other dimension reduction methods and, while some utilized either PLS or AFT, none of the methods addressed the issue of censoring directly. A large number of published large-scale genomic studies with censored survival outcomes seem to indicate that the proportion of censored observations is in the 60-80% range (Rouam et al., 2011, TCGA Research Network). Thus, in this application, having a method like ACPR-AFT that adjusts for censored data is not only beneficial but also desirable. Furthermore, we explore the utility of CPR coefficients, ω, which are computed as $\omega = \boldsymbol{\nu}\mathbf{c}'$ where the columns of $\boldsymbol{\nu}$ are weights vectors and $\mathbf{c}'$ are the loadings, in developing a survival prediction model and for feature ranking and selection.

4.1.1 Supervised extraction of CPR components

The first step in the (A)CPR-AFT algorithm is to apply CPR which finds weight vectors, columns of $\boldsymbol{\nu}$, such that the linear combinations $\mathbf{Z}^{(\gamma)}\boldsymbol{\nu}$ maximize the objective function $R^2(\mathbf{Z}^{(\gamma)}\boldsymbol{\nu}, \tilde{\mathbf{t}})[\mathrm{Var}(\mathbf{Z}^{(\gamma)}\boldsymbol{\nu})]^{\gamma}$ where $\tilde{\mathbf{t}} = (\log t_1, \ldots, \log t_n)$ are the log transformed observed event times subject to the constraints outlined earlier. Here, $\mathbf{Z}^{(\gamma)}$ is found via the spectral decomposition of $\mathbf{Z}$; after this step, standard PLS can be applied to $\mathbf{Z}^{(\gamma)}$. Let $u_{ik} = \mathbf{z}_i^*\boldsymbol{\nu}_k, i = 1, \ldots, N, k = 1, \ldots, K$, denote the linear combinations selected by CPR where $\mathbf{z}_i^*$ denotes the i^{th} row of $\mathbf{Z}^{(\gamma)}$ and $\boldsymbol{\nu}_k$ denotes the k^{th} column of $\boldsymbol{\nu}$. These represent the CPR components, and the number of components $K < p$ is chosen based on leave-one-out cross validation (LOOCV) to minimize the predicted residual sum of squares (PRESS) statistic. We use the reparametrization $\gamma \equiv \alpha/(1 - \alpha)$ where α values of 0, 1/2 and 1 correspond to OLS, PLS and PCR, respectively. In subsequent sections, we outline how (A)CPR-AFT can be used to select K, the optimal number of components, and α, the CPR parameter. PLS is an important special case of CPR which maximizes the covariance between $\mathbf{Z}\boldsymbol{\nu}$ and $\log(t)$ (Devarajan et al., 2010).

4.1.2 Fitting the AFT model to CPR components

The next step in the (A)CPR-AFT methods involve fitting the AFT model in equation 3.1. We propose flexible parametric and distribution-free versions of (A)CPR-AFT using the *GenF* and *sAFT* models, respectively. The flexible parametric AFT model, *GenF*, is given by

$$\log Y_i = \boldsymbol{\beta}'\mathbf{u}_i + \sigma^*W_i, i = 1, \ldots, N, \tag{4.2}$$

where Y_i is the survival time for the i-th subject, $\mathbf{u}_i' = (u_{i0}, \ldots, u_{iK})$ is a $(K+1)$ vector for the i-th subject, $\boldsymbol{\beta} = (\beta_0, \beta_1, \ldots, \beta_K)$ is the $(K+1)$ vector of unknown regression parameters, W_i are independent error terms with a common distribution $F_W \sim GenF$ and σ^* is the scale parameter. In this setting, $u_{ik}, k = 1, \ldots, K$ represent the K CPR components for the i-th subject, $i = 1, \ldots, N$. The semi-parametric AFT model, *sAFT*, has the form

$$\log Y_i = \boldsymbol{\beta}'\mathbf{u}_i + W_i, i = 1, \ldots, N, \tag{4.3}$$

where the terms are as defined in equation 4.2. Here, W_i are independent error terms with unknown distribution F. Given a feature expression vector $\mathbf{z}$ and PLS component $\mathbf{u}$, using equation 4.2 the survival function of Y is given by

$$
\begin{aligned}
S(y|\mathbf{z}) = P(Y > y|\mathbf{z}) &\approx P(\log Y > \log y|\mathbf{u}) \\
&= P(\boldsymbol{\beta}'\mathbf{u} + \sigma^* W > \log y|\mathbf{u}) \\
&= P(W > (\log y - \boldsymbol{\beta}'\mathbf{u})/\sigma^*) \\
&= 1 - F_W\left(\frac{\log y - \boldsymbol{\beta}'\mathbf{u}}{\sigma^*}\right),
\end{aligned}
$$

and is estimated by replacing the unknown parameters with their maximum likelihood estimates.

4.2 The (A)CPR-AFT algorithms

Below, we outline the CPR-AFT and ACPR-AFT algorithms for a pre-specified CPR parameter α. **CPR-AFT** treats ignores censoring and treats those observations as complete while **ACPR-AFT** imputes censored observations using mean residual life based on available data.

Algorithm 1 CPR-AFT

1: Choose the parameter α.
2: Perform CPR using the chosen α to obtain weight vectors $\boldsymbol{\nu}_k, k = 1, \ldots, K$, for each value of the number of CPR components, K, in pre-specified range. Choose K using the PRESS statistic based on LOOCV. These weight vectors are used to compute the CPR coefficients ω as described earlier.
3: Use the observed time to event data (t_i, δ_i) and covariates $\mathbf{u_i} = (u_{i1}, \ldots, u_{ik})$ (components), to model the effect of these covariates on survival time by fitting the $GenF$ or $sAFT$ model (Equation 4.2 or 4.3) via maximization of the appropriate likelihood function.

Algorithm 2 ACPR-AFT

1: Repeat Steps 1 and 2 of CPR-AFT (Algorithm 1).
2: Use uncensored data to obtain CPR components where the number of components, K, is chosen as specified in Step 2 of the CPR-AFT algorithm. For $GenF$, use these components as covariates for the model in equation 4.2 and obtain estimates for $\boldsymbol{\beta}$ and σ^*; and for $sAFT$, use these components as covariates for the model in equation 4.3 and obtain the estimate for $\boldsymbol{\beta}$. It is important to note that the components obtained in this step are used only to estimate $\boldsymbol{\beta}$ and/or σ^*.
3: Let $v_i = \log t_i$. Replace v_i by $v_i^* = \delta_i v_i + (1-\delta_i)\hat{E}(\log Y_i | \log Y_i > v_i), i = 1,\ldots,N$. Under the $GenF$ model in equation (4.2), $\hat{E}(\log Y_i | \log Y_i > v_i) = \hat{\boldsymbol{\beta}}\mathbf{u_i} + \hat{\sigma}^* E\left(W | W > \frac{v_i - \hat{\boldsymbol{\beta}}\mathbf{u_i}}{\hat{\sigma}^*}\right)$, where $\hat{\boldsymbol{\beta}}$ and $\hat{\sigma}^*$ are obtained in Step 2 and W is the error term in equation 4.2. Under the $sAFT$ model (4.3), $\hat{E}(\log Y_i | \log Y_i > v_i) = \hat{\boldsymbol{\beta}}\mathbf{u_i} + \hat{\sigma}^* E\left(W | W > v_i - \hat{\boldsymbol{\beta}}\mathbf{u_i}\right)$, where $\hat{\boldsymbol{\beta}}$ is obtained in Step 2 and W is the error term in equation 4.3. The calculation of this conditional expectation for $GenF$ and $sAFT$ models are outlined in §4.3 and §4.4, respectively. Here, $\boldsymbol{u}_i$ are the CPR components obtained in Step 2 of CPR-AFT.
4: Use $v_1^*, \ldots, v_N^*$ from Step 2 to construct new CPR components. The number of components, K, for the adjusted survival data is determined using PRESS based on LOOCV as outlined in Step 2 of CPR-AFT. The weight vectors $\boldsymbol{\nu}_k$ corresponding to these new CPR components are used to compute the CPR coefficients ω.
5: Use the new CPR components to build the final model. This is discussed in detail in §5.

For a pre-specified CPR parameter α, the (A)CPR-AFT algorithms outlined above facilitate the efficient extraction of the optimal number, K, of CPR components as determined by PRESS and LOOCV. ACPR-AFT has a significant advantage over CPR-AFT because it adjusts for the censored observations. In §5, we propose a survival prediction algorithm based on (A)CPR-AFT that simultaneously allows the optimal choice of α to be chosen in addition to K and utilizes the CPR coefficients, ω, from the final model. Furthermore, in §6, we show that (A)CPR-AFT can be utilized for feature ranking and selection.

4.3 A flexible parametric approach to (A)CPR-AFT

In equation 4.2, if $Y \sim GenF$ then the density of can be written as

$$f_Y(y|\mu,\sigma,n,m) = \frac{1}{\sigma\beta(n,m)}e^{\frac{-\mu n}{\sigma}}y^{\frac{n}{\sigma}-1}\left(\frac{n}{m}\right)^n\left[1 + \left(\frac{n}{m}\right)(e^{-\mu}y)^{\frac{1}{\sigma}}\right]^{-(n+m)}, y > 0. \quad (4.4)$$

As shown in Ciampi et al. (1986) and Cox (2008), this model has an umbrella structure with many special cases, where choosing specific parameter values will result in a particular model of interest. A few special cases are described in Table 2. Many other models fall under this umbrella including gamma, Weibull (exponential), log-normal, log-logistic, and Burr-type distributions. Here, we propose a generalization of (A)CPR-AFT using the $GenF$ model, which encompasses many important special cases and is, therefore, a flexible alternative for modeling censored survival data in conjunction with large-scale "omics" data. To this end, we first obtain the density of $W = \frac{\log Y - \mu}{\sigma}$ in equation 4.2 and use it to derive the expression for

Table 2: *GenF* Model: Some special Cases

Model	Parameters
Log-Normal	$m \to \infty$, $n \to \infty$
Weibull	$m \to \infty$, $n = 1$
Log Logistic	$m = 1$, $n = 1$

the conditional expectation $E\left(W|W > \frac{v_i - \hat{\beta}u_i}{\hat{\sigma}^*}\right)$ for this model in the ACPR-AFT algorithm. Using equation 4.4, the density of W can be derived to be

$$f_W(w) = \frac{1}{\beta(n,m)} \left(\frac{n}{m}\right)^n e^{nw} \left[1 + \left(\frac{n}{m}\right)e^w\right]^{-(n+m)}, \quad -\infty < w < \infty.$$

Using the density of W,

$$E(W|W > z) = \frac{\int_z^\infty w f_W(w)\, dw}{1 - F(z)} = \frac{\int_{\frac{(\frac{n}{m})e^z}{1+(\frac{n}{m})e^z}}^1 \ln\left[\frac{m}{n}\left(\frac{v}{1-v}\right)\right] f_V(v)\, dv}{1 - \int_0^{\frac{(\frac{n}{m})e^z}{1+\frac{n}{m}e^z}} f_V(v)\, dv}, \tag{4.5}$$

where $f_V(v) \sim Beta(n,m) = \frac{1}{\beta(n,m)} v^{n-1}(1-v)^{m-1}$ and $z = \frac{v_i - \hat{\beta}u_i}{\hat{\sigma}^*}$.

In CPR-AFT, *GenF* is used for model fitting in Step 3 while in ACPR-AFT, it is used to use to compute the expression in equation 4.5 in Step 3 and to fit appropriate models to estimate the parameters in Steps 2 and 5. The *GenF* model can be fitted using the R package flexsurvreg described in Cox (2008). In addition to *GenF*, we are also interested in comparing its performance to important special cases such as the log-normal, log logistic, and Weibull. To obtain the conditional expectation, $E(W|W > z)$, for these special cases, one just needs to replace the parameters in equation 4.5 with those listed in Table 2 for the respective model.

4.4 A semi-parametric approach to (A)CPR-AFT

Although the generalization based on *GenF* is parametric in nature, it offers tremendous modeling flexibility. Here, we further extend our (A)CPR-AFT approach using the *sAFT* model which has no distributional assumption for the error term. This is an attractive property as it does not force the choice of a specific model, thus providing more flexibility in the application of the method. The *sAFT* model has the form given in equation 4.2. Following Jin et al. (2006), the conditional expectation in Step 3 of ACPR-AFT is obtained as

$$E(W|W > z) = \frac{\int_{e_i(\beta)}^\infty u\, d\hat{F}_\beta(u)}{1 - \hat{F}_\beta\{e_i(\beta)\}}, \tag{4.6}$$

where $z = e_i(\beta) = v_i - \hat{\beta}u_i$ and $\hat{F}_\beta(t) = 1 - \prod_{i:e_i(\beta)<t}\left(1 - \frac{\delta_i}{\sum_{j=1}^n I\{e_j(\beta)\geq e_i(\beta)\}}\right)$ is the Kaplan Meier estimator of F based on $\{e_i(\beta), \delta_i\}$. In CPR-AFT, *sAFT* is used for model fitting in

Step 3 while in ACPR-AFT, it is used to compute the expression in equation 4.6 in Step 3 and to fit appropriate models to estimate the parameters in Steps 2 and 5. Fitting the semi-parametric AFT model is based on the Buckley-James (BJ) type estimator developed by Jin et al. (2006) and is implemented in the R package lss.

Merits of BJ Estimation The BJ estimation method (Buckley & James, 1979) is an iterative least squares approach that is closely related to OLS without censoring and, thus, provides a more accessible interpretation to practitioners. It has been utilized in a variety of applications involving many areas such as medicine (Hammer et al., 2002), genetics (Bautista et al., 2008), astronomy (Steffen et al., 2006) and economics (Deaton and Irish, 1984; Calli and Wever bergh, 2009), and has been shown to be the preferred estimation approach in a comparison study (Heller & Simonoff, 1990). In contrast to methods that assume independence between the censoring mechanism and covariates, the BJ approach requires weaker assumptions and, in conjunction with boosting, has been shown to be superior to LASSO-type methods and to generate sparser models (Wang & Wang, 2010). It utilizes Kaplan-Meier estimates and is readily available in statistical software such as R (R Core Team, 2018); moreover, BJ estimation for the AFT model can be conveniently extended to describe more complex data structures with existing software, such as MART and MARS (Friedman, 1991; 2001). Hence, we use BJ estimation in the $sAFT$ model.

5   Supervised dimension reduction

A natural approach to build a final model based on (A)CPR-AFT is using the K CPR components from ACPR-AFT as covariates in equation 4.2 and following Step 3 of CPR-AFT. It would result in an AFT model based on reduced components from ACPR-AFT. A prognostic index can be defined as $\eta = \mathbf{u}\beta$, where $\mathbf{u}$ is the $n \times K$ matrix whose columns contain the K CPR components and β is the K-vector of coefficients from this final AFT model fit. A major disadvantage of this approach is that component information is not available for new subjects and, therefore, it is not possible to develop a prediction model that can be used on future subjects with feature expression profiles. Recall that $\omega_{p \times 1} = \boldsymbol{\nu}_{p \times K} \mathbf{c}'_{K \times 1}$ where the K columns of $\boldsymbol{\nu}$ contain weight vectors and $\mathbf{c}'$ contains the loadings associated with the K CPR components which are contained in the columns of $\mathbf{u}_{n \times K}$. Hence, we propose an approach based on the CPR coefficients, ω, rather than directly using the reduced CPR components, $\mathbf{u}$, for predicting the survival probability of a future subject whose feature expression profile is readily available. As shown in the next section, some important differences exist between this approach and the final AFT model discussed above with vastly different implications for prediction.

5.1   Developing a prognostic index

We use the CPR coefficients to devise an approach based on the weighted average of feature expressions and illustrate its utility in developing a survival prediction model. Using the vector of CPR coefficients, ω, for the p features from a particular model of interest and the

$n \times p$ feature expression matrix, $\mathbf{Z}$, the weighted average, η, is calculated as

$$\eta = \mathbf{Z}\omega. \tag{5.1}$$

This results in an n-dimensional vector, which we call the *prognostic index (PI)*, where each element represents a subject's weighted average feature expression. When utilizing all features in the calculation of η, a heavier weight is placed on features deemed significant; however, η can be calculated using only a subset of features of interest. It is worth noting that PI represents the predicted (log) survival times and, thus, enables the development of a prediction model using (A)CPR-AFT as outlined in the next section.

5.2   A survival prediction algorithm using (A)CPR-AFT

We develop a survival prediction algorithm using the CPR coefficients ω from (A)CPR-AFT, separately for $GenF$ and $sAFT$, by adopting a flexible approach that simultaneously chooses the optimal α in addition to K, the number of CPR components. The proposed approach utilizes (A)CPR-AFT for several choices of α that represent a variety of scenarios: the midpoint of the trajectory from OLS and PLS ($\alpha = 0.25$), PLS (0.5), the midpoint of the trajectory from PLS to PCR (0.75) and PCR (0.95). Since $p \gg n$, OLS ($\alpha = 0$) is not a useful option in our application and PCR ($\alpha = 1$) requires a value of α close to 1 in order to avoid numerical instability. In this approach, the final model is chosen based on the optimal (K, α) combination that results in the smallest PRESS using LOOCV after applying (A)CPR-AFT, and the corresponding CPR coefficients ω are used to develop the PI for evaluating the performance of this prediction model. The CPR parameter α for the unadjusted CPR-AFT cannot be optimally chosen. Hence, we consider two different unadjusted methods in our comparisons, one each based on the chosen value of α from $GenF$ and $sAFT$ in ACPR-AFT.

Prior to the application of (A)CPR-AFT, supervised marginal screening procedures were used to narrow down the number of features. These methods ensure that features used for prediction demonstrate an association with survival, at the univariate level, after adjusting for potential confounders such as age of diagnosis and stage of disease. An added benefit of such pre-filtering is that it significantly reduces computation time. Supervised marginal screening was performed to select (i) features that fit the $sAFT$ model and had a statistically significant effect on survival (sAFT), (ii) features that had a significant effect on survival using concordance regression (CON) (Dunkler et al., 2010), or (iii) features that fit the $GenF$ model ($R^2 > 0.80$) and had a statistically significant effect on survival (GenF), a t the 0.05 significance level. Once a subset is selected, (A)CPR-AFT is applied and the optimal (α, K) is chosen. The CPR coefficients, ω, are retained for the adjusted (ACPR-AFT) and unadjusted (CPR-AFT) methods (based on $GenF$ and $sAFT$) and used to predict the logarithm of survival time for each subject given their feature expression profile $\mathbf{Z}$ using the prognostic index, $PI = \eta = \mathbf{Z}\omega$. We use italicized notation $sAFT$ and $GenF$ to denote the ACPR-AFT methods while sAFT and GenF are used to denote the respective marginal screening methods.

The following cross validation approach is used to build and evaluate the prediction models. The data is first split into training and test sets roughly in a 2:1 ratio, where ω_{tr}

represents the vector of CPR coefficients corresponding to the training set and is used to predict the logarithm of survival time in the test set. Thus, $PI = \mathbf{Z}_{te}\omega_{tr}$, where $\mathbf{Z}_{te}$ is from the test set, and the model is evaluated for prediction accuracy. We utilize the following measures of prediction accuracy to evaluate and compare the predictive performance of ACPR-AFT, using $GenF$ or $sAFT$ models, to the unadjusted CPR-AFT approach: (i) R^2, the fraction of variation that is explained by the K CPR components in the final (A)CPR-AFT model, (ii) Mean Squared Error, $MSE = \frac{1}{n^*}\sum_{i=1}^{n}\delta_i(PI_i - \log T_i)^2$ where PI_i is the prognostic index for the i^th subject, $n^* = \sum_{i=1}^{n}\delta_i$ and $\delta_i = 1$ implies the event was observed; MSE is calculated for both the training set, MSE_{TR}, and the test set, MSE_{TE}, and (iii) area under the time-dependent receiver operating characteristic curve (AUC) which quantifies a method's ability to predict survival at varying time points such as 2, 3 or 5 years and is implemented in the R package survivalROC (Haegerty et al., 2000) . An AUC close to 1 indicates better prediction accuracy (Blanche et al., 2013). In summary, the survival prediction algorithm involves the following steps:

Algorithm 3 Survival Prediction Algorithm

1: Use supervised marginal screening to filter features using sAFT, CON or GenF, as outlined above.
2: Randomly split the filtered data set into training (67% of subjects) and test sets (33% of subjects).
3: Apply (A)CPR-AFT ($GenF$ and $sAFT$) to the training set using $\alpha = (.25, .5, .75, .95)$.

- Choose optimal (α, K) combination.
- Retain the CPR regression coefficients, ω_{tr}.

4: Use ω_{tr} from Step 3 to predict (log) survival times in the test set, i.e., calculate $PI = \mathbf{Z}_{te}\omega_{tr}$, where $\mathbf{Z}_{te}$ is from the test set.
5: Evaluate the prediction models using the measures of prediction accuracy outlined above.
6: Repeat Steps 2-5 25 times. Median values of the prediction accuracy measures from step 5 are reported.

---

6 Application to simulated data

6.1 Simulation schemes

We considered two different simulation schemes to generate artificial survival and feature expression data sets based on the approach outlined in Dunkler et al. (2010). In order to account for various types of hazards, survival times $Y_i, i = 1, \ldots, n$, were generated from 5 different models - standard log-normal (LN), log-logistic (LL1, LL2), and Weibull (W1, W2). These models are specified as follows:

- LN ($\mu = 0$, $\sigma = 1$)

- LL1 ($\alpha_1 = 2, \lambda_1 = 2, \lambda_2 = 4$)

- LL2 ($\alpha_1 = 3, \alpha_2 = 4, \lambda_1 = 1, \lambda_2 = 2$)

- W1 ($\alpha_1 = 1, \lambda_1 = \frac{1}{2}$)

- W2 ($\alpha_1 = 3, \alpha_2 = 2, \lambda_1 = 1, \lambda_2 = \frac{1}{2}$)

where (α_1, α_2) and (λ_1, λ_2) are the shape and scale parameters, respectively. In the LN model, μ and σ are the location and scale parameters, respectively. Here, LL1 and W1 refer to the respective models where the shape parameters are the same but the scale parameters differ, and LL2 and W2 refer to the respective models where both the shape and scale parameters differ. We use a more informed approach that is broader in scope compared to that of Dunkler et al. (2010), who only considered W1 in their simulations. Here, LN, LL2 and W2 cases are of particular interest because of their ability to simulate crossing hazards. To simulate censoring, we drew random samples with uniform follow-up times C from $U(0, \tau)$ and defined the observed survival time as $T = \min(Y, C)$ with censoring indicator $\delta = I(T = Y)$. We chose τ to get censoring proportions of 33, 67% and 80%.

For each model, we simulated censored survival times and feature expression data for $N = 200$ subjects and $p = 5000$ mock features where feature expression is linked to survival time based on the logarithm of the hazard ratio (HR), $\beta_g(t) = \beta_0 \log(HR)$. Feature expression data was generated from the standard normal model. Following Klein and Moeschberger (2003), $\log(HR)$ was calculated based on the respective model of interest. For LN, we used $\beta_g(t) = \beta_0(t^2 - 1)$ to simulate crossing hazards similar to what was done in Dunkler et al. (2010). Then, β_0 was chosen so that only the first 400 features were assumed to have an effect on survival time, with 200 having a large effect and 200 having a small effect. In Scheme 1, we adopt a univariate approach where feature expression is linked to survival one feature at a time, and in Scheme 2 we adopt a multivariate approach that incorporates correlations between features. More details on these steps can be found in Dunkler et al. (2010).

6.2 Feature ranking and selection using (A)CPR-AFT

In this section, we discuss the utility of (A)CPR-AFT for feature ranking and selection by comparing the performance of (A)CPR-AFT algorithms based on log-normal (LN), log-logistic (LL), Weibull (W), $GenF$ and $sAFT$ models. Wold et al. (2002), Devarajan et al. (2010) and Mehmood et al. (2012) discuss the use of PLS coefficients, ω, and variable importance projection (VIP), denoted by ξ, as measures for feature selection in PLS. Note that the PLS coefficients arise as a special case of CPR when $\alpha = 0.5$. For example, features could be ranked based on the absolute value of CPR coefficients, ω, which can take on values on the entire real line, or directly using the VIP, ξ, which is a non-negative quantity.

VIP accumulates the importance of each feature as reflected by the weight $\boldsymbol{\nu}_k$ from each component. Essentially, it is a measure of the contribution of each feature according to the variance explained by each component. The VIP value, ξ, for feature j is calculated as

$$\xi_j = \sqrt{p \sum_{k=1}^{K} \left[SS_k \left(\nu_{jk} / ||\boldsymbol{\nu}_k|| \right)^2 \right] / \sum_{k=1}^{K} SS_k}, \, j = 1, \ldots, p, \tag{6.1}$$

where p is the number of features, K is the number of components, $\boldsymbol{\nu}_k$ is the weight vector for the k-th component, and SS_k is the sum of squares explained by the k-th component. In other words, CPR produces K vectors of weights, each of which has p elements corresponding to the p features. In the VIP calculation, $(\nu_{jk}/||\boldsymbol{\nu}_k||)^2$ represents the importance of the k-th component. Thus, ξ_j is calculated for feature j, $j = 1 \ldots p$, and then features are selected based on a threshold pre-specified by the user. A popular threshold is 1 (i.e., $\xi > 1$) and is discussed in Mehmood et al. (2012). Since the number of relevant features is pre-determined in simulations, a threshold is not relevant in our application where features are ranked separately based on decreasing VIP and absolute value of ω. Although choice of a threshold is unclear in the analysis of real data, these measures can be used for feature ranking and selection. We plan to investigate choice of the ξ threshold in future work. Our analyses using a variety of large-scale "omics" data sets showed that $\alpha = 0.5$ resulted in overall better performance in terms of model fit and prediction accuracy; hence, we focus on this special case for the simulation studies. As mentioned earlier, high censoring appears to be a common theme in many large-scale "omics" studies involving censored survival outcomes (Bhattacharjee et al., 2001; Beer et al., 2002; Tothill et al., 2008; Saintigny et al., 2011; Rouam et al., 2011; TCGA Research Network); hence, this scenario is also of particular interest in the simulations. As evidenced in the following sections, our studies indicate that VIP is a more useful quantity for feature ranking but less useful for the purposes of developing a prediction model where the CPR coefficients play a significant role.

For each simulation scheme and censoring combination, 200 randomly generated data sets were created and assessed. The (A)CPR-AFT algorithm was applied to each simulated data set and mock features were ranked separately using the absolute value of PLS coefficients, ω, and VIP, ξ. In the remainder of this section, we will only use ω or ξ to denote these methods. For CPR-AFT, ω and ξ are calculated in Step 1 of the algorithm, and therefore, are not model specific. On the other hand, in ACPR-AFT these quantities are computed after adjusting censored observations in Step 4, thus taking the pre-specified model into account in the parametric version or in a completely distribution-free manner in the case of $sAFT$. In the parametric version, we considered $GenF$ and three of its well-known special cases - LN, LL and W. For each method, the ranked lists were used to compute mean values of sensitivity, specificity, Youden Index (=sensitivity+specificity-1) (Youden, 1950) and area under the receiver operating characteristic (ROC) curve (AUC) across the 200 data sets. The purpose of this analysis is to compare the performance of (i) the two ranking methods, ω and ξ, (ii) $GenF$ vs. its special cases in ACPR-AFT, and (iii) CPR-AFT vs. ACPR-AFT for $GenF$ and $sAFT$ models under both simulation schemes.

6.2.1 Comparison of $GenF$ and its special cases in ACPR-AFT

We examined the performance of $GenF$ and its special cases in ACPR-AFT using AUC and the Youden Index for 33% and 80% censoring. Figure 1 depicts the ROC curves for 33% and 80% censoring for ω and ξ using simulation scheme 1. In all four cases - shown in panels (a)-(d) - we observe that $GenF$ outperforms LN, W, and LL. The corresponding AUCs and Youden Indices are reported in Table 3. The AUCs for $GenF$ are higher than its special cases in each situation, and the differences increase as censoring increases. We note that the unadjusted CPR-AFT results in the lowest Youden index and AUC in every scenario and

performs significantly worse than ACPR-AFT using $GenF$. In addition, we observe that ξ is superior to the use of ω in terms of AUC and the Youden Index, thus indicating that VIP is a better tool for feature ranking and selection.

Next, we evaluated the performance using simulation scheme 2 for 33% and 80% censoring. Panels (a) and (b) of Figure 2 show the ROC curves for these censoring proportions, respectively, comparing the performance of ω and ξ for $GenF$. Similar to scheme 1, we observe that ξ significantly outperforms ω. Panels (c) and (d) of Figure 3 show the ROC curves for 33% and 80%, respectively, comparing the performance of $GenF$ and CPR-AFT for ξ. Again, it is evident that $GenF$ outperforms the unadjusted method. Although not shown, scheme 2 showed similar results for $GenF$ compared to its special cases LN, LL and W. At each censoring level, $GenF$ had the largest AUC and Youden index, similar to what was observed for scheme 1 in Figure 1 and Table 3. Since $GenF$ outperformed its special cases, we will focus only on $GenF$, $sAFT$ and the unadjusted method for the remainder of the paper. A very similar performance was observed across multiple simulated data sets and thus, results are reported only for a single, representative data set.

6.2.2 Simulation Scheme 1

In this section, we evaluate the performance of $GenF$- and $sAFT$- based ACPR-AFT against unadjusted CPR-AFT using data simulated from scheme 1 as described in §5.1. Table 4 shows the AUCs using ξ as the feature selection tool. Across each scheme and censoring level, we observe that the proposed ACPR-AFT method (both $GenF$- and $sAFT$-based) have a clear advantage over CPR-AFT, and as the censoring increases, the differences in performance become stronger. In fact, CPR-AFT has the lowest AUC in every single scheme for the 67% and 80% censoring cases. Thus, we observe a clear benefit when imputing censored observations using ACPR-AFT. Next, we examine differences between $GenF$- and $sAFT$-based ACPR-AFT. Both $GenF$ and $sAFT$ perform similarly in many instances, but there are particular schemes where one outperforms the other. For example, AUCs of $GenF$ are higher for the LN, LL1 and LL2 cases, and AUCs of $sAFT$ are higher in the W1 and W2 cases. In Table 5, AUCs using ω are shown and observed to be lower than those of ξ in each case. Thus, ξ appears to be the optimal approach for feature selection; however, even when ω is used, $GenF$ and $sAFT$ outperform CPR-AFT.

Next, we focus our attention on the Youden index displayed in Table 6 for each method and censoring level using both ω and ξ as ranking methods. First, we note that Youden indices based on ω are lower than those based on ξ in every single case. Thus, as expected, ξ significantly outperforms ω in ranking features in every scenario. Furthermore, we observe that $GenF$- and $sAFT$-based ACPR-AFT outperform CPR-AFT in almost every scenario, and just as we had observed with the AUCs, the differences in performance become larger as censoring increases. Thus, we notice a clear improvement due to ACPR-AFT. $GenF$ and $sAFT$ perform similarly in many cases, but as indicated by the AUC results, one occasionally outperforms the other.

6.2.3 Simulation Scheme 2

We now examine simulation scheme 2, which introduces correlations between features. Recall that in simulation scheme 1, $GenF$ and $sAFT$ performed similarly and better than the unadjusted approach, and ξ outperformed ω. In this section, we demonstrate that the same conclusions can be made for scheme 2. However, although the results have a similar trend, the AUC and Youden values for scheme 2 are lower than those in scheme 1 likely due to the complexity in the data introduced by the correlation structure between features. The AUCs for scheme 2 using ξ are shown in Table 7. Similar to scheme 1, we observe that our adjusted methods have a clear advantage over the unadjusted method and this difference becomes greater as the censoring fraction increases. We observe that $GenF$ results in higher AUCs for LN and LL2, which differs slightly from scheme 1 results, but $GenF$ and $sAFT$ perform very similarly in the remaining schemes. The AUCs using ω were observed to be much lower than those obtained using ξ in each case (data not shown); in fact, the AUCs for ω ranged from 0.50 to 0.57 which suggests that its selection capability is only slightly better than a coin flip. Thus, ξ is seen to be optimal approach for feature selection. Next, we examine the Youden index displayed in Table 8. We note that the Youden indices based on ω are lower than those based on ξ in every single case, an observation similar to that in scheme 1. In fact, in many cases, the Youden index for ω is 0. We also observe from these results that $GenF$ and $sAFT$ outperform the unadjusted method in practically every scenario and particularly for higher censoring. $GenF$ and $sAFT$ perform similarly in many cases, but as indicated by the AUC results, one occasionally outperforms the other. The simulation results from both schemes showed $GenF$ either outperformed or matched the performance of LN, LL and W. Hence, we focus only on $GenF$ in the examples in the remainder of this paper.

6.3 Evaluating the prediction algorithm

Simulated data was generated using the same approach outlined in §5.1. Simulation schemes 1 and 2 were utilized for LN, LL1, LL2, W1 and W2 models for 33%, 67% and 80% censoring. Each data set contained 200 observations and 5,000 mock features and model parameters were chosen appropriately to result in survival times that mimicked actual survival times (say in months or years). Training and test sets were obtained using a 2:1 split. For each combination of simulation scheme, model and censoring proportion, a total of 25 different random splits were generated and the predictive performance of ACPR-AFT (using $\alpha = 0.5$) was evaluated on each data set using $GenF$ or $sAFT$ and compared with the unadjusted method, CPR-AFT. Median summaries for different measures of prediction accuracy are presented in Tables 9-14. It is not surprising to note that the overall performance of training sets is better than that of test sets across all parameters considered for both CPR-AFT and ACPR-AFT. However, as the censoring fraction increases, a significant improvement is noted in the predictive performance (AUC and MSE) of ACPR-AFT (both $GenF$ and $sAFT$) compared to CPR-AFT for the test sets. This improved performance is observed under both simulation schemes for each of the five different models under consideration. These results, thus, highlight the superiority of ACPR-AFT under a variety of data generating mechanisms encountered in practice.

7 Application to high-throughput "omics" data

We demonstrate the utility of (A)CPR-AFT in supervised dimension reduction and developing a survival prediction model using the following publicly available data sets in cancer genomics.

7.1 Data sets

- Head & Neck squamous cell carcinoma (HNSCC): This data set was published by the TCGA Research Network (http://cancergenome.nih.gov/) and contains the expression profiles of 19,341 genes obtained using RNA sequencing for 221 subjects with cancers of the oral cavity, a subgroup of HNSCC. Pre-processed gene expression data (RSEM values, Li & Dewey, 2011) was downloaded from the Broad Institute (http://gdac.broadinstitute.org/) and further normalized using the $\log_2(x+1)$ transformation which accounts for exact zeros. A gene was included in the analyses only if (i) if 50% of patients have expression values for that gene, and (ii) protein expression of that gene was observed in at least one head and neck cancer sample in the Human Protein Atlas database (Uhlen et al., 2015).

- Glioblastoma (GBM): This data set was published by the TCGA Research Network (http://cancergenome.nih.gov/) and contains the methylation profiles (beta values) for 280 tumor samples obtained using the Infinium HumanMethylation27 platform. The beta values were normalized using the logit transformation. For genes with multiple methylation probes, the probe most negatively correlated with expression is used.

- Ovarian cancer: Tothill et al. (2008) studied the relationship between survival and gene expression in ovarian cancer using tumor samples from 276 subjects and Affymetrix U133 Plus 2 microarrays. This RMA normalized and $\log_2$ transformed (Irizarry et al., 2003) data set was filtered using a coefficient of variation threshold of 5% to remove probe sets with low expression, thus resulting in a data set containing 32,575 probe sets.

- Oral cancer: Saintigny et al. (2011) studied 86 subjects enrolled in a clinical chemoprevention trial where the primary endpoint of interest was the development of oral cancer. This RMA normalized and $\log_2$ transformed data set (Irizarry et al., 2003) contains the expression profiles of 12,776 probe sets obtained using the Human Gene 1.ST platform.

7.2 Extracting genomic components using (A)CPR-AFT

For the purpose of illustrating the application of (A)CPR-AFT in supervised dimension reduction, we resort to the PLS special case corresponding to $\alpha = 0.5$. CPR-AFT and ACPR-AFT were applied to the four genomic data sets outlined above where the latter method was based on $GenF$ or $sAFT$. In each case, the optimal number of CPR components, K, was determined using the PRESS statistic based on LOOCV by considering ranks $k = 2, \ldots, 15$. The results, summarized in Table 15, indicate that for all data sets $GenF$- and

sAFT-based APCRAFT generally outperform the unadjusted method by explaining a higher proportion of the variation in the data at the chosen optimal K or by choosing a smaller optimal K. In some cases, ACPR-AFT is observed to choose more than or at least as many components as the unadjusted method; however, even at the optimal K chosen by the unadjusted method (ovarian and oral data) or at a lower rank (HNSCC data), ACPR-AFT explains a significantly higher fraction of the variation compared to the unadjusted method (as shown in the last three rows of Table 15). For the GBM data set, ACPR-AFT performs as well as the unadjusted method; this is likely due to the relatively small fraction (26%) of censored observations in this set compared to the others where censoring ranges from 59-62%. These examples suggest that it is possible to choose a more parsimonious model than that provided by ACPR-AFT while still explaining most of the variation in the data. Moreover, they highlight the utility of ACPR-AFT for handling censored data. The CPR coefficients and VIP obtained from ACPR-AFT can also be used for feature ranking purposes as demonstrated by our simulations.

7.3   Interpreting the prognostic index

The prognostic index, $PI = \eta = \mathbf{Z}\omega$ can be interpreted as a linear predictor and is particularly relevant within the context of (A)CPR-AFT which combines two linear models. The predicted survival curves obtained from *GenF*- and *sAFT*-based ACPR-AFT using η, plotted in Figure 4 for each data set, thus illustrate the use of CPR coefficients, ω, in building a prediction model. In each case, the corresponding model under consideration (*GenF* or *sAFT*) was statistically significant at the 5% significance level. Furthermore, GOF tests performed using the methods outlined in §3 revealed a good fit for the PH, PO and AFT models across different methods and data sets. The only exception to this is a lack of fit for the PH model for *sAFT*-based ACPR-AFT for the ovarian data set which indicates an overall time-varying effect due to features selected by the model. While a weighted average using VIP, ξ, could serve as a prognostic index, it does not directly predict survival time and, hence, its interpretation is unclear. These examples thus serve to illustrate the utility of η in elucidating the relationship between feature expression and patient survival and to account for time-varying effects of features.

7.4   Evaluating the prediction algorithm

The prediction algorithm utilizing (A)CPR-AFT was evaluated using these four genomic data sets. The filtering mechanisms help reduce the data by a significant fraction by eliminating features that are not associated with patient survival according to different criteria. features could be removed using the sAFT or CON filter as outlined in §4.2; in addition, a VIP threshold such as $\xi > 1$ could be imposed as discussed in §5.2. We observed that the range of ξ was quite similar for both filters for each method (*sAFT* or *GenF*) as well as between methods, and the median value was around 1 across these data sets. Thus, the filtered sets used for model building contained statistically significant features from sAFT or CON where approximately half of the features had $\xi > 1$.

After filtering, the prediction algorithm was applied separately to each training and test set. Median values of different measures of prediction accuracy, based on 25 random training

and test sets split in a 2:1 ratio, are summarized in Tables 16-19. Overall, less variation in these measures was observed for ACPR-AFT across the training and test sets and for both filters used compared to CPR-AFT; in particular, $GenF$ showed much less variation in R^2 and MSE compared to $sAFT$ while the opposite effect was observed for AUC. In each run, the optimal (K, α) combination was chosen as outlined earlier for ACPR-AFT. Since CPR-AFT does not involve choosing α based on adjustment (as in Step 4 of ACPR-AFT), the optimal α chosen by $GenF$- and $sAFT$-based ACPR-AFT is used for comparison purposes.

The fraction of variation in the data explained by the CPR components extracted by a particular model, quantified by R^2, are significantly higher for $GenF$- and $sAFT$-based ACPR-AFT compared to the unadjusted CPR-AFT. Since computation of R^2 requires model fitting, it is relevant only to the training set. Not surprisingly, MSE is generally larger for the test set compared to the training set across all filtering mechanisms and data sets. However, for both the training and test sets, we consistently observe smaller MSE for ACPR-AFT methods compared to CPR-AFT across all four data sets which indicates that both $GenF$ and $sAFT$ result in more accurate predictions. In particular, a significant reduction in MSE between ACPR-AFT and CPR-AFT is observed for the oral test sets, ranging from 58-91% across both filters. For the ovarian and HNSCC test sets, the CON (57%) and sAFT (26%) filters result in maximum reduction.

In addition, we examine the AUCs calculated for 2, 3 and 5 year survival. Once again, we note that the training set AUCs are larger than those of the test set in each case. More importantly, in both the training and test sets and across both filtering mechanisms, we observe larger AUCs for $GenF$ and $sAFT$ compared to the unadjusted methods. The improvements observed in the test sets are particularly relevant and are noteworthy for all data sets. For example, using the sAFT and CON filters, we observe an AUC range of 0.75-0.86 for $GenF$ and $sAFT$ for the ovarian test sets while the unadjusted methods range only from 0.64-0.74. The performance of the two filters was similar for the ovarian and HNSCC data sets and resulted in improvements of 0.12 and 0.08 in AUC, respectively; for the GBM data set, however, CON provided substantial improvement in AUC of 0.09 compared to 0.01 for sAFT and 0.05 overall (median values). In addition, a statistically significant difference was generally observed between the AUCs from time-dependent ROC curves for ACPR-AFT and CPR-AFT (data not shown).

Furthermore, the prediction algorithm was evaluated using the HNSCC data set (all observations) and the results are summarized in Table 20. Here, the entire data set is treated as a single, large training set. Three different filters were considered in this analysis - sAFT, CON and GenF - as outlined in §4.2. The optimal value of α chosen by ACPR-AFT for various filters is indicated in the footnote of Table 20 and this value was used for the corresponding unadjusted method. A significant improvement in the performance of the algorithm is noted across all three filters and both adjusted methods compared to the unadjusted method; in particular, the parametric GenF filter results in the best overall performance when used in conjunction with either adjusted method. These results are similar to that seen in Table 16 and further demonstrate the predictive performance of ACPR-AFT compared to CPR-AFT.

8   Conclusions and discussion

In this paper, we proposed supervised dimension reduction methods for analyzing large-scale "omics" data in conjunction with censored survival outcomes. Our methods combine CPR - a unified framework that includes OLS, PCR and PLS as special cases - for dimension reduction with the AFT model - a censored linear regression model - for handling survival data, and offer distinct advantages relative to currently available approaches. The versatility afforded by the parametric ($GenF$) and semi-parametric ($sAFT$) versions of the AFT model and its partial overlap with the widely used PH and PO models allow a variety of time-varying feature effects to be incorporated. Moreover, both CPR and AFT fall within the linear models framework and the proposed hybrid model, (A)CPR-AFT, combines their strengths in a unique fashion that does not match any other available method. The umbrella structure of $GenF$ provides a fully parametric, yet tremendously flexible, approach for modeling survival data while $sAFT$ utilizes BJ estimation which has been shown to be a robust method. A particularly attractive characteristic of ACPR-AFT is its ability to account for censored observations common to studies with survival endpoints. Many large-scale "omics" studies involving survival outcomes of interest tend to contain a significant fraction of censored observations and an appropriate method for handling these incomplete observations has been lacking. The simulation results demonstrated the superior predictive performance of ACPR-AFT over CPR-AFT under a plethora of data generating mechanisms particularly as the fraction of censored observations increased, thus making it a practically useful tool for data analysis. These results were corroborated using publicly available data sets in cancer genomics where the performance of the proposed survival prediction algorithm was shown to improve significantly when censoring was accounted for in this manner.

The ability of the proposed methods to handle NPH within this context is unparalleled and it offers a robust and flexible approach for predictive modeling of a wide variety of large-scale "omics" data. The CPR coefficients and VIP play complementary roles and serve different purposes. The former is useful for computing the PI which was used to develop and evaluate the prediction algorithm while the latter was shown to be a superior measure for feature ranking. However, choosing an appropriate threshold for these measures is an important consideration and could form part of future work on this topic. When combined with an appropriate marginal screening method, this approach could serve as a useful feature selection tool by significantly reducing the number of relevant features in the prediction algorithm which could, in turn, not only improve its performance further but also help develop a feature signature that is predictive of survival.

Furthermore, the proposed methods are broadly applicable to a variety of high-throughput "omics" data such as feature expression data arising from next-generation sequencing, allele-specific expression, methylation, microarrays and SNP arrays as well as large-scale data from proteomics, metabolomics and DNA copy number studies, many of which have been utilized in this study. There has been a recent surge in integrative "omic" analyses that simultaneously involve different data types as well as other quantitative outcome variables using publicly available data from repositories such as TCGA and GEO (Ramakodi et al., 2016; Li et al., 2013; Lawrenson et al., 2015). Within this context, the proposed unifying framework offers a robust platform for analysis and interpretation.

Acknowledgement The work of KD was funded in part through the NIH/NCI Cancer Center Support Grant P30 CA006927.

References

[1] Bair, E., Hastie, T., Paul, D., & Tibshirani, R. (2006). Prediction by Supervised Principal Components. *Journal of the American Statistical Association*, **101(473)**, 119-137.

[2] Bastien, P., Bertrand, F., Meyer, N. & Maumy-Bertrand, M. (2015). Deviance residuals-based sparse PLS and sparse kernel PLS regression for censored data. *Bioinformatics*, **31(3)**, 397-404.

[3] Bautista, L. E.,Vargas, C. I., Orostegui, M., & Gamarra, G. (2008). Population-based case-control study of renin-angiotensin system features polymorphisms and hypertension among Hispanics. *Hypertension Research*, **31(3)**, 401-408.

[4] Beer, D.G., Kardia, S.L., Huang, C.C., Giordano, T.J., Levin, A.M., Misek, D.E., . . ., Hanash, S. (2002) Gene-expression profiles predict survival of patients with lung adenocarcinoma. *Nature Medicine*, **8**.

[5] Bhattacharjee, A., Richards, W.G., Staunton, J., Li, C., Monti, S., Vasa, P., . . ., Meyerson, M. (2001). Classification of human lung carcinomas by mRNA expression profiling reveals distinct adenocarcinoma subclasses. *PNAS*, **98(24)**, 13790-13795.

[6] Blanche, P., Dartigues, J., & Jacqmin-Gadda, H. (2013). Estimating and comparing time-dependent areas under receiver operating characteristic curves for censored event times with competing risks. *Statistics in Medicine.*

[7] Boulesteix, A.L. & Strimmer, K. (2007). Partial least squares: a versatile tool for the analysis of high dimensional genomic data. *Bioinformatics*, **8**, 32-44.

[8] Bøvelstad, H.M., Nygård, S., Størvold, H.L., Alrdrin, M., Borgan, Ø., Frigessi, A., & Lingjaerde, O.C. (2007). Predicting survival from microarray data - a comparative study. *Bioinformatics*, **23**, 2080-2087.

[9] Buckley, J. & James, I. (1979). Linear Regression with Censored Data. *Biometrika*, **66(3)**, 429-436.

[10] Cai, T., Huang, J., & Tian, L. (2009). Regularized estimation for the accelerated failure time model. *Biometrics*, **65(2)**, 394-404.

[11] Calli, M. K. & Wever bergh, M. (2009). Forecasting newspaper demand with censored regression. *Journal of the Operational and Research Society*, **60(7)**, 944951.

[12] Chen, X. & Cook, D.R. (2010). Some insights into continuum regression and its asymptotic properties. *Biometrika*, **97(4)**, 985-989.

[13] Ciampi, A., Hogg, S.A., & Kates, L. (1986). Regression Analysis of Censored Survival Data with the Generalized F Family–An Alternative to the Proportional Hazards Model. *Statistics in Medicine*, **5**, 85-96.

[14] Chen, K., Zhezhen, J., & Ying, Z. (2002). Semiparametric analysis of transformation models with censored data. *Biometrika*, **89(3)**, 659-668.

[15] Cox, C. (2008). The generalized F distribution: An umbrella for parametric survival analysis. *Statistics in Medicine*, **27**, 4301-4312.

[16] Cox, D.R. (1972). Regression models and life-tables. *Journal of the Royal Statistical Society*, **34**, 187-220.

[17] Datta, S., Le-Rademacher, J., & Datta, S. (2007). Predicting patient survival from microarray data by accelerated failure time modeling using partial least squares and LASSO. *Bioinformatics*, **63**, 259-271.

[18] Deaton, A. & Irish, M. (1984). Statistical models for zero expenditures in household budgets. *Journal of Public Economics*, **23(1-2)**, 5980.

[19] de Jong, S., Wise, B.M., & Ricker, N.L. (2001). Canonical partial least squares and continuum power regression. *Journal of Chemometrics*, **15**, 85-100.

[20] Devarajan, K., Zhou, Y., Chachra, N., & Ebrahimi, N. (2010). A supervised approach for predicting patient survival with feature expression data. *Proc IEEE Int Symp Bioinformatics Bioeng*, **5521718**, 26-31.

[21] Devarajan, K. & Ebrahimi, N. (2011). A semi-parametric generalization of the Cox proportional hazards regression model: Inference and applications. *Computational Statistics and Data Analysis*, **55**, 667-676.

[22] Dunkler, D., Schemper, M., & Heinze, G. (2010). Gene selection in microarray survival studies under possibly non-proportional hazards. *Bioinformatics*, **26**, 784-790.

[23] Eng, K. H. & Hanlon, B. M. (2012). Discrete mixture regression models for heterogeneous time-to-event data: Cox Assisted Clustering. *Bioinformatics*, **30(12)**, 1690-1697.

[24] Engler, D. & Li, Yi. (2009). Survival Analysis with High-Dimensional Covariates: An Application in Microarray Studies. *Stat Appl Genet Mol Biol*, **8(1)**, 1544-6115.

[25] Fan, J. & Li, R. (2002). Variable selection for Cox's proportional hazards model and frailty model. *Ann. Statist.*, **30**, 7499.

[26] Fan, J., Feng, Y., & Wu, Y. (2010). High-dimensional variable selection for Cox's proportional hazards model. *Institute of Mathematical Statistics*, **6**, 70-86.

[27] Frank, I. E., & Friedman, J. H. (1993). A statistical view of some chemometrics regression tools (with discussion). *Technometrics*, **35(2)**:109148.

[28] Friedman, J. (1991). Multivariate adaptive regression splines (with discussion). *The Annals of Statistics*, **19**, 1-141.

[29] Friedman, J. (2001). Greedy function approximation: A gradient boosting machine. *The Annals of Statistics*, **29(5)**, 1189-1232.

[30] Geng, Y., Lu, W., & Zhang, H.H. (2014). A Model-Free Machine Learning Method for Risk Classification and Survival Probability Prediction. *Stat*, **3(1)**, 337-350.

[31] Gui, J. & Li, H. (2005). Penalized Cox regression analysis in the high-dimensional and low-sample size settings, with applications to microarray feature expression data. *Bioinformatics*, **21(13)**, 3001-3008.

[32] Haegerty, P.J., Lumley, T., & Pepe, M.S. (2000). Time-dependent ROC curves for censored survival data and a diagnostic marker. *Biometrics*, **56(2)**, 337-44.

[33] Hammer, S.,Vaida, F., Bennett, K., Holohan, M., Sheiner, L., Eron, J.,..., Mellors, J. (2002). Dual vs single protease inhibitor therapy following antiretroviral treatment failure: a randomized trial. *Journal of the American Medical Association*, **288**, 169180.

[34] Heller, G. & Simonoff, J. S. (1990). A comparison of estimators for regression with a censored response variable. *Biometrika*, **77**, 515520.

[35] Huang, J. & Harrington, D. (2004). Dimension reduction in the linear model for right-censored data: Predicting the change of HIV-I RNA levels using clinical and protease feature mutation data. *Lifetime Data Analysis*, **10**, 425-443.

[36] Huang, J., Ma. S., & Xie, H. (2006). Regularized Estimation in the Accelerated Failure Time Model with High-Dimensional Covariates. *Biometrics*, **62(3)**, 813-820.

[37] Irizarry, R.A., Hobbs, B., Collin, F., Beazer-Barclay, Y.D., Antonellis, K.J., Scherf, U., & Speed, T.P. (2003). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics*, **4(2)**, 249-264.

[38] Jin, Z., Lin, D.Y., Wei, L.J., & Ying, Z. (2003). Rank-based inference for accelerated failure time model. *Biometrika*, **90**, 341-353.

[39] Jin, Z., Lin, D.Y., & Ying, Z. (2006). On least-squares regression with censored data. *Biometrika*, **93**, 147-161.

[40] Johnson, B. A. (2009). On lasso for censored data. *Electronic Journal of Statistics*, **3**, 485-506.

[41] Johnson, B. A. (2008a). Variable selection in semiparametric linear regression with censored data . *Journal of the Royal Statistical Society*, **70**, 351-370.

[42] Johnson, B. A. (2008b). Estimation in the L_1-Regularized Accelerated Failure Time Model. *Technical Report, Emory University*.

[43] Johnson, B. A., Lin, D.Y., & Zeng, D. (2008). Penalized Estimating Functions and Variable Selection in Semiparametric Regression Models. *Journal of the American Statistical Association*, **103**, 672-680.

[44] Kalbfleisch, J.D. & Prentice, R.L. (2002) The statistical analysis of failure time data. *Wiley.*

[45] Kaneko, S., Hirakawa, A., & Hamada, C. (2012). Gene Selection using a High-Dimensional Regression Model with Microarrays in Cancer Prognostic Studies. *Cancer Informatics*, **11**, 29-39.

[46] Klein, P. J. & Moeschberger, L. M. (2003). Survival Analysis: Techniques for censored and truncated data. *New York: Springer.*

[47] Laimighofer, M., Krumsiek, J., Buettner, F. & Theis, F.J. (2016). Unbiased Prediction and Feature Selection in High-Dimensional Survival Regression. *Journal of Computational Biology*, **23(4)**, 1-12.

[48] Lawrenson, K., Li, Q., Kar, S., Seo, J.H., Tyrer, J., Spindler, T.J., . . . , Freedman, M.L. (2015). Cis-eQTL analysis and functional validation of candidate susceptibility features for high-grade serous ovarian cancer. *Nat Commun*, **6**, 8234.

[49] Leng, C. & Ma, S. (2007). Accelerated failure time models with nonlinear covariates effects. *Aust. N. Z. J. Stat*, **49(2)**, 155-172.

[50] Li, B. & Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference group. *BMC Bioinformatics*, **4(12)**, 323.

[51] Li. H. & Gui, J. (2004). Partial Cox regression analysis for high-dimensional microarray feature expression data. *Bioinformatics*, **20(1)**, 208-215.

[52] Li, L. & Li, H. (2004). Dimension reduction methods for microarrays with application to censored survival data. *Bioinformatics*, **20(18)**, 3406-3412.

[53] Li, H. & Luan, Y. (2003). Kernel Cox regression models for linking feature expression profiles to censored survival data. *Pacific Symposium of Biocomputing*, **8**, 65-76.

[54] Li, H. & Luan, Y. (2005). Boosting proportional hazards models using smoothing splines, with applications to high-dimensional microarray data. *Bioinformatics*, **21(10)**, 2403-2409.

[55] Li, Q., Seo, J.H., Stranger, B., McKenna, A., Pe'er, I., Laframboise, T., . . . , Freedman, M.L. (2013). Integrative eQTL-based analyses reveal the biology of breast cancer risk loci. *Cell*, **152(3)**, 633-641.

[56] Liu, Z., Chen, D., Tan, M., Jiang, F. & Gartenhaus, R. (2010). Kernel based methods for accelerated failure time model with ultra-high dimensional data. *BMC Bioinformatics*, **11**, 606.

[57] Lu, W. & Li, L. (2008). Boosting method for nonlinear transformation models with censored survival data. *Biostatistics*, **9**, 658-667.

[58] Lorber, A., Wangen, L. E. & Kowalski, B. R. (1987). A theoretical foundation for the PLS algorithm. *J. Chemometrics*, **1**, 19-31.

[59] Luan,Y. & Li, H. (2008). Group additive regression models for genomic data analysis. *Biostatistics*, **9(1)**, 100113.

[60] Ma, S., Kosorok, M. R. & Fine, J. P. (2006). Additive risk models for survival data with high-dimensional covariates. *Biometrics*, **62**, 202-210.

[61] Martinussen & Scheike (2006). Dynamic Regression Models for Survival Data. *Statistics for Biology and Health.*

[62] Mehmood, T., Liland, K.H., Snipen, L. & Saebo, S. (2012). A review of variable selection methods in Partial Least Squares Regression. *Chemometrics and Intelligent Laboratory Systems*, **118**, 62-69.

[63] Nguyen, D. V. & Rocke, D. M. (2002). Partial least squares proportional hazard regression for application to DNA microarray survival data. *Bioinformatics*, **18**, 1625-1632.

[64] Nguyen, D. V. (2005). Partial least squares dimension reduction for microarray feature expression data with censored response. *Mathematical Biosciences*, **193**, 119-137.

[65] Novak, P. (2010). Checking goodness-of-fit of the accelerated failure time model for survival data. *WDS'10 Proceedings of Contributed Papers*, **I**, 189-194.

[66] Nygard, S., Borgan, O., Kingjaerde, O.C., & Storvold, H.L. (2008). Partial least squares Cox regression for genome-wide data. *Lifetime Data Anal*, **14**, 179-195.

[67] Pang, H., George, S.L., Hui, K., & Tong, T. (2012). Gene selection using iterative feature elimination random forests for survival outcomes. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, **9(5)**, 1422-1431.

[68] Park, P., Tian, L., & Kohane, I. (2002). Linking feature expression data with patient survival times using partial least squares. *Bioinformatics*, **18**, 120-127.

[69] Park, M. Y. & Hastie, T. (2006). An L_1 regularization path algorithm for generalized linear models. *Technical report, Stanford University.*

[70] Peri, S., Devarajan, K., Yang, D.H., Knudson, A.G., & Balachandran, S. (2013). Meta-Analysis Identifies NF-B as a Therapeutic Target in Renal Cancer. *PLoS ONE*, **8(10)**, e76746.

[71] R Core Team (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

[72] Ramakodi, M.P., Devarajan, K., Blackman, E., Gibbs, D., Luce, D., Deloumeaux, J., ..., Ragin, C.C. (2016). Integrative genomic analysis identifies ancestry-related expression quantitative trait loci on DNA polymerase beta and supports the association of genetic ancestry with survival disparities in head and neck squamous cell carcinoma. *Cancer*, **123(5)**, 849-860.

[73] Reid, N. (1994). A Conversation with Sir David Cox. Statistical Science, **9(3)**, 439-455.

[74] Rosset, S., Zhu, J., & Hastie, T. (2004). Boosting as a regularized path to a maximum margin classifier. *Journal of Machine Learning Research*, **5**, 941-973.

[75] Rouam, S., Moreau, T., & Broet, P. (2011). A pseudo-R^2 measure for selecting genomic markers with crossing hazards functions. *BMC Medical Research Methodology*, **11**, 28.

[76] Saintigny, P., Zhang, L., Fan, Y.H., El-Naggar, A.K., Papadimitrakopoulou, V.A., Feng, L., ..., Mao, L. (2011). feature expression profiling predicts the development of oral cancer. *Cancer Prev Res (Phila)*, **4**, 218-229.

[77] Segal, M.R. (2006). Microarray feature expression data with linked survival phenotypes: diffuse large-B-cell lymphoma revisited. *Biostatistics* **7(2)**: 268285. doi:10.1093/biostatistics/kxj006

[78] Stone, M. & Brooks, R.J. (1990). Continuum Regression: Cross-Validated Sequentially Constructed Prediction Embracing Ordinary Least Squares, Partial Least Squares and Principal Components Regression. *Journal of the Royal Statistical Society*, **52(2)**, 237-269.

[79] Stute, W. (1993). Consistent estimation under random censorship when covariables are present. *Journal of Multivariate Analysis*, **45(1)**, 89-103.

[80] Sundberg, R. (2002). Continuum Regression. *Encyclopedia of Statistical Science*, **2nd Ed**.

[81] Steffen, A.T., Strateva, I., Brandt, W. N., Alexander, D. M., Koekemoer, A. M., Lemmie, B. D., Schneider, D. P., and Vignali, C. (2006). The X-ray-to-optical properties of optically selected active galaxies over wide luminosity and redshift ranges. *The Astronomical Journal*, **131**, 28262842.

[82] Storey, J. D. (2002). A direct approach to false discovery rates. *Royal Statistical Society*, **64(3)**, 479-498.

[83] The Cancer Genome Atlas (TCGA) Network: http://cancergenome.nih.gov/

[84] Therneau, T. & Grambsch, P. (1994). Proportional hazards tests and diagnostics based on weighted residuals. *Biometrika*, **81**, 515-526.

[85] Tibshirani, R. (1997). The LASSO method for variable selection in the Cox model. *Statistics in Medicine*, **16**, 385-395.

[86] Tothill, R.W., Tinker, A.V., George, J., Brown, R., Fox, S.B., Lade, S., ..., Bowtell, D.D. (2008). Novel molecular subtypes of serous and endometroid ovarian cancer linked to clinical outcome. *Clinical Cancer Research*, **14**, 5198-5208.

[87] Uhlen, M., Fagerberg, L., Hallstrom, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., ..., Nielsen, J. (2015). Proteomics. Tissue-based map of the human proteome. *Science*, **347(6220)**, 1260419.

[88] Van Belle, V., Pelckmans, K., Van Huffel, S., & Suykens, J.A. (2011). Improved performance on high-dimensional survival data by application of Survival-SVM. *Bioinformatics*, **21(1)**, 87-94.

[89] Verweij, P. & Houwelingen, H. (1994). Penalized likelihood in Cox regression. *Statistics in Medicine*, **13**, 2427-2436.

[90] Wang, H. & Leng, C. (2007). Unified LASSO Estimation by Least Squares Approximation. *Journal of the American Statistical Association*, **102**, 1039-1048.

[91] Wang, S., Nan, B., Zu, J. & Beer, D.G. (2008). Doubly penalized Buckley- James method for survival data with high-dimensional covariates. *Biometrics*, **64**, 132-140.

[92] Wang, S., Nan, B., Zhou, N., & Zhu, J. (2009). Hierarchically penalized Cox regression with grouped variables. *Biometrika*, **96**, 241-247.

[93] Wang, Z. & Wang, C. Y. (2010). Buckley-James boosting for survival analysis with high-dimensional biomarker data. *Statistical Applications in Genetic Molecular Biology*, **9(24)**.

[94] Wei, Z. & Li, H. (2007). Nonparametric pathway-based regression models for analysis of genomic data. *Biostatistics*, **8**, 265-284.

[95] van Wieringen, W.N., Kun, D., Hampel, R., & Boulesteix, A. (2009). Survival prediction using feature expression data: A review and comparison. *Computational Statistics and Data Analysis*, **53**, 1590-1603.

[96] Witten, D. & Tibshirani, R. (2008). Testing significance of features by lassoed principal components. *The Annals of Applied Statistics*, **2(3)**, 9861012.

[97] Wold, S., Sjöström, M., & Eriksson, L. (2002). Partial Least Squares Projections to Latent Structures (PLS) in Chemistry. *Encyclopedia of Computational Chemistry*.

[98] Xu, J., Yang, Y., & Ott, J. (2005). Survival analysis of microarray expression data by transformation models. *Computational Biology and Chemistry*, **29**, 91-94.

[99] Yang, S. & Prentice, R. L. (2005). Semiparametric analysis of short-term and long-term hazard ratios with two-sample survival data. *Biometrika*, **92**, 1-17.

[100] Youden, W. J. (1950). Index for rating diagnostic tests. *Cancer*, **3(1)**, 32-35.

[101] Zhang, H. & Lu, W. (2007). Adaptive-Lasso for Cox's Proportional Hazards Model. *Biometrika*, **94**, 691-703.

[102] Zhang, W., Ota, T., Shridhar, V., Chien, J., & Kuang, R. (2013). Network-based Survival Analysis Reveals Subnetwork Signatures for Predicting Outcomes of Ovarian Cancer Treatment. *PLoS Comput Biol*, **9(3)**, e1002975.

[103] Zou, H. (2008). A Note on Path-Based Variable Selection in the Penalized Proportional Hazards Model. *Biometrika*, **95**, 241-247.

Table 3: Simulation Scheme 1

Censoring	Method	ω		ξ	
		Youden	AUC	Youden	AUC
33 %	ACPR-AFT LN	.19	.67	.65	.94
	ACPR-AFT W	.21	.68	.65	.94
	ACPR-AFT LL	.20	.67	.65	.94
	ACPR-AFT *GenF*	.33	.75	.74	.96
	CPR-AFT	.12	.64	.48	.88
80%	ACPR-AFT LN	.16	.63	.30	.74
	ACPR-AFT W	.16	.63	.30	.74
	ACPR-AFT LL	.16	.63	.30	.74
	ACPR-AFT *GenF*	.44	.80	.49	.84
	CPR-AFT	.03	.51	.09	.62

Table 4: Simulation Scheme 1: AUC (ξ)

Censoring	Method	LN	LL1	LL2	W1	W2
33 %	ACPR-AFT *GenF*	.91	.92	.69	.95	.89
	ACPR-AFT *sAFT*	.91	.92	.76	.97	.92
	CPR-AFT	.88	.89	.74	.93	.89
67 %	ACPR-AFT *GenF*	.87	.89	.72	.90	.76
	ACPR-AFT *sAFT*	.86	.84	.71	.95	.84
	CPR-AFT	.80	.78	.65	.85	.72
80 %	ACPR-AFT *GenF*	.85	.84	.75	.87	.64
	ACPR-AFT *sAFT*	.81	.77	.68	.92	.70
	CPR-AFT	.72	.66	.59	.77	.56

Table 5: Simulation Scheme 1: AUC (ω)

Censoring	Method	LN	LL1	LL2	W1	W2
	ACPR-AFT *GenF*	.67	.64	.52	.66	.56
33 %	ACPR-AFT *sAFT*	.68	.64	.55	.71	.54
	CPR-AFT	.60	.57	.53	.62	.54
	ACPR-AFT *GenF*	.71	.63	.57	.62	.52
67 %	ACPR-AFT *sAFT*	.66	.55	.56	.73	.55
	CPR-AFT	.54	.51	.53	.54	.50
	ACPR-AFT *GenF*	.68	.63	.60	.62	.51
80 %	ACPR-AFT *sAFT*	.59	.54	.54	.75	.53
	CPR-AFT	.51	.51	.51	.53	.50

Table 6: Simulation Scheme 1: Youden Index

Model	Method	33% cens.		67% cens.		80% cens.	
		ω	ξ	ω	ξ	ω	ξ
	ACPR-AFT *GenF*	.21	.52	.29	.55	.22	.52
LN	ACPR-AFT *sAFT*	.21	.52	.19	.52	.10	.48
	CPR-AFT	.10	.50	.03	.41	.01	.17
	ACPR-AFT *GenF*	.15	.56	.15	.54	.14	.45
LL1	ACPR-AFT *sAFT*	.15	.56	.05	.43	.04	.24
	CPR-AFT	.06	.50	.01	.21	.01	.07
	ACPR-AFT *GenF*	.02	.20	.07	.23	.11	.31
LL2	ACPR-AFT *sAFT*	.05	.29	.06	.20	.04	.14
	CPR-AFT	.03	.23	.03	.10	.01	.05
	ACPR-AFT *GenF*	.18	.62	.12	.52	.14	.50
W1	ACPR-AFT *sAFT*	.24	.70	.31	.65	.34	.61
	CPR-AFT	.11	.56	.04	.40	.03	.21
	ACPR-AFT *GenF*	.04	.45	.02	.22	.01	.10
W2	ACPR-AFT *sAFT*	.07	.52	.05	.39	.04	.18
	CPR-AFT	.03	.43	.01	.12	.01	.02

Table 7: Simulation Scheme 2: AUC (ξ)

Censoring	Method	LN	LL1	LL2	W1	W2
	ACPR-AFT *GenF*	.58	.82	.66	.68	.61
33 %	ACPR-AFT *sAFT*	.57	.89	.66	.67	.60
	CPR-AFT	.54	.84	.67	.67	.62
	ACPR-AFT *GenF*	.82	.88	.70	.79	.67
67 %	ACPR-AFT *sAFT*	.75	.88	.69	.69	.68
	CPR-AFT	.63	.80	.70	.69	.66
	ACPR-AFT *GenF*	.84	.86	.84	.83	.70
80 %	ACPR-AFT *sAFT*	.76	.88	.76	.74	.71
	CPR-AFT	.68	.78	.72	.69	.67

Table 8: Simulation Scheme 2: Youden Index

Model	Method	33% cens.		67% cens.		80% cens.	
		ω	ξ	ω	ξ	ω	ξ
LN	ACPR-AFT $GenF$	0	.01	0	.11	0	.13
	ACPR-AFT $sAFT$	.01	.01	.01	.09	0	.09
	CPR-AFT	0	.01	0	.01	0	.03
LL1	ACPR-AFT $GenF$	.05	.19	.06	.24	.06	.20
	ACPR-AFT $sAFT$	.06	.24	.04	.23	.04	.22
	CPR-AFT	.02	.16	.01	.10	0	.08
LL2	ACPR-AFT $GenF$	0	.03	0	.04	0	.13
	ACPR-AFT $sAFT$	0	.03	0	.03	0	.06
	CPR-AFT	0	.03	0	.03	0	.05
W1	ACPR-AFT $GenF$	.01	.04	.01	.09	.02	.15
	ACPR-AFT $sAFT$	.01	.03	.02	.05	.01	.07
	CPR-AFT	0	.03	.01	.04	0	.04
W2	ACPR-AFT $GenF$	.01	.02	0	.03	0	.04
	ACPR-AFT $sAFT$	.01	.02	0	.04	0	.03
	CPR-AFT	0	.02	0	.02	0	.02

Table 9: Simulation Scheme 1: 33% Censoring

Model	Measure	ACPR-AFT *GenF*	ACPR-AFT *sAFT*	CPR-AFT
LN, Scheme 1	R^2_{COMP}	100	99.9	99.9
	MSE_{TR}	0.007	0.009	0.008
	AUC, t = 2	0.94	0.96	0.92
	AUC, t = 3	0.91	0.93	0.89
	MSE_{TE}	12.7	12.8	13.4
	AUC, t = 2	0.94	0.92	0.91
	AUC, t = 3	0.93	0.95	0.94
LL1, Scheme 1	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0005	0.0006	0.0006
	AUC, t = 2	0.99	0.99	0.99
	AUC, t = 3	0.99	0.99	0.99
	MSE_{TE}	2.05	2.05	2.06
	AUC, t = 2	0.97	0.97	0.97
	AUC, t = 3	0.97	0.96	0.96
LL2, Scheme 1	R^2_{COMP}	99.8	100	99.8
	MSE_{TR}	0.03	0.001	0.03
	AUC, t = 2	0.98	0.99	0.97
	AUC, t = 3	0.98	0.98	0.97
	MSE_{TE}	2.93	2.95	3.28
	AUC, t = 2	0.88	0.88	0.88
	AUC, t = 3	0.93	0.95	0.94
W1, Scheme 1	R^2_{COMP}	100	100	99.5
	MSE_{TR}	0.001	0.001	0.001
	AUC, t = 2	0.98	0.98	0.99
	AUC, t = 3	0.98	0.99	0.98
	MSE_{TE}	4.21	4.14	4.94
	AUC, t = 2	0.99	0.99	0.98
	AUC, t = 3	0.98	0.98	0.98
W2, Scheme 1	R^2_{COMP}	100	100	98.9
	MSE_{TR}	0.001	0.001	0.05
	AUC, t = 2	0.98	0.98	0.98
	AUC, t = 3	0.97	0.98	0.96
	MSE_{TE}	7.52	7.66	8.68
	AUC, t = 2	0.86	0.87	0.86
	AUC, t = 3	0.92	0.94	0.92

All methods are based on $\alpha = 0.5$.

Table 10: Simulation Scheme 2: 33% Censoring

Model	Measure	ACPR-AFT GenF	ACPR-AFT sAFT	CPR-AFT
LN, Scheme 2	R^2_{COMP}	100	98.2	98.1
	MSE_{TR}	0.006	0.179	0.228
	AUC, t = 2	0.75	0.76	0.76
	AUC, t = 3	0.89	0.89	0.9
	MSE_{TE}	9.3	9.3	9.2
	AUC, t = 2	0.89	0.91	0.91
	AUC, t = 3	0.88	0.9	0.84
LL1, Scheme 2	R^2_{COMP}	99.9	99.9	100
	MSE_{TR}	0.002	0.002	0.001
	AUC, t = 2	0.71	0.7	0.66
	AUC, t = 3	0.71	0.71	0.67
	MSE_{TE}	3.12	3.11	3.14
	AUC, t = 2	0.99	0.98	0.97
	AUC, t = 3	0.97	0.97	0.96
LL2, Scheme 2	R^2_{COMP}	97.7	98	97.6
	MSE_{TR}	0.056	0.097	0.092
	AUC, t = 2	0.56	0.56	0.56
	AUC, t = 3	0.59	0.58	0.57
	MSE_{TE}	2.3	2.3	2.3
	AUC, t = 2	0.98	0.97	0.95
	AUC, t = 3	0.86	0.96	0.89
W1, Scheme 2	R^2_{COMP}	98.2	98	98.1
	MSE_{TR}	0.08	0.1	0.09
	AUC, t = 2	0.84	0.84	0.85
	AUC, t = 3	0.79	0.8	0.8
	MSE_{TE}	4.36	4.04	4.02
	AUC, t = 2	0.98	0.99	0.99
	AUC, t = 3	0.95	0.98	0.98
W2, Scheme 2	R^2_{COMP}	99.9	98	98
	MSE_{TR}	0.005	0.127	0.114
	AUC, t = 2	0.91	0.91	0.92
	AUC, t = 3	0.87	0.88	0.88
	MSE_{TE}	6.12	5.29	5.3
	AUC, t = 2	0.98	0.98	0.98
	AUC, t = 3	0.98	0.98	0.97

All methods are based on $\alpha = 0.5$.

Table 11: Simulation Scheme 1: 67% Censoring

Model	Measure	ACPR-AFT GenF	ACPR-AFT sAFT	CPR-AFT
LN, Scheme 1	R^2_{COMP}	100	100	99.9
	MSE_{TR}	0.003	0.006	0.007
	AUC, t = 2	0.82	0.88	0.81
	AUC, t = 3	0.82	0.86	0.79
	MSE_{TE}	10.9	11.6	12.7
	AUC, t = 2	0.86	0.84	0.84
	AUC, t = 3	0.85	0.85	0.84
LL1, Scheme 1	R^2_{COMP}	100	100	100
	MSE_{TR}	0.001	0.00001	0.003
	AUC, t = 2	0.98	0.97	0.97
	AUC, t = 3	0.96	0.96	0.94
	MSE_{TE}	1.35	1.25	1.36
	AUC, t = 2	0.94	0.86	0.84
	AUC, t = 3	0.91	0.84	0.79
LL2, Scheme 1	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0002	0.0003	0.0004
	AUC, t = 2	0.87	0.89	0.85
	AUC, t = 3	0.93	0.94	0.91
	MSE_{TE}	1.93	1.89	1.99
	AUC, t = 2	0.88	0.86	0.84
	AUC, t = 3	0.8	0.84	0.79
W1, Scheme 1	R^2_{COMP}	100	100	99.5
	MSE_{TR}	0.003	0.0003	0.01
	AUC, t = 2	0.97	0.97	0.96
	AUC, t = 3	0.93	0.93	0.92
	MSE_{TE}	3.7	3.79	4.09
	AUC, t = 2	0.92	0.9	0.86
	AUC, t = 3	0.92	0.88	0.74
W2, Scheme 1	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0004	0.00001	0.0005
	AUC, t = 2	0.96	0.97	0.96
	AUC, t = 3	0.98	0.97	0.97
	MSE_{TE}	5.52	5.51	5.57
	AUC, t = 2	0.89	0.89	0.86
	AUC, t = 3	0.9	0.85	0.79

All methods are based on $\alpha = 0.5$.

Table 12: Simulation Scheme 2: 67% Censoring

Model	Measure	ACPR-AFT GenF	ACPR-AFT sAFT	CPR-AFT
LN, Scheme 2	R^2_{COMP}	100	100	100
	MSE_{TR}	0.000002	0.000002	0.000002
	AUC, t = 2	0.8	0.79	0.79
	AUC, t = 3	0.92	0.92	0.92
	MSE_{TE}	4.9	4.9	5
	AUC, t = 2	0.85	0.87	0.76
	AUC, t = 3	0.87	0.87	0.83
LL1, Scheme 2	R^2_{COMP}	100	100	100
	MSE_{TR}	0.00003	0.00004	0.0009
	AUC, t = 2	0.94	0.9	0.79
	AUC, t = 3	0.79	0.8	0.74
	MSE_{TE}	1.12	1.11	1.14
	AUC, t = 2	0.95	0.9	0.89
	AUC, t = 3	0.87	0.84	0.8
LL2, Scheme 2	R^2_{COMP}	97.5	97.3	97.5
	MSE_{TR}	0.016	0.046	0.061
	AUC, t = 2	0.7	0.7	0.56
	AUC, t = 3	0.72	0.7	0.58
	MSE_{TE}	1.45	1.49	1.52
	AUC, t = 2	0.95	0.86	0.76
	AUC, t = 3	0.73	0.87	0.78
W1, Scheme 2	R^2_{COMP}	100	100	100
	MSE_{TR}	0.001	0.001	0.001
	AUC, t = 2	0.98	0.98	0.97
	AUC, t = 3	0.96	0.96	0.95
	MSE_{TE}	3.88	3.96	3.97
	AUC, t = 2	0.94	0.92	0.9
	AUC, t = 3	0.9	0.89	0.83
W2, Scheme 2	R^2_{COMP}	97.7	97.5	97.5
	MSE_{TR}	0.05	0.06	0.08
	AUC, t = 2	0.92	0.91	0.86
	AUC, t = 3	0.95	0.93	0.89
	MSE_{TE}	4.98	5	5
	AUC, t = 2	0.98	0.93	0.93
	AUC, t = 3	0.92	0.91	0.9

All methods are based on $\alpha = 0.5$.

Table 13: Simulation Scheme 1: 80% Censoring

Model	Measure	ACPR-AFT GenF	ACPR-AFT sAFT	CPR-AFT
LN, Scheme 1	R^2_{COMP}	100	100	99.9
	MSE_{TR}	0.000004	0.0001	0.0001
	AUC, t = 2	0.93	0.88	0.76
	AUC, t = 3	0.93	0.92	78
	MSE_{TE}	10	10	11.4
	AUC, t = 2	0.74	0.77	0.65
	AUC, t = 3	0.75	0.75	0.64
LL1, Scheme 1	R^2_{COMP}	99.2	99.3	99
	MSE_{TR}	0.002	0.008	0.01
	AUC, t = 2	0.93	0.97	0.93
	AUC, t = 3	0.87	0.92	0.84
	MSE_{TE}	0.61	0.66	0.7
	AUC, t = 2	0.9	0.76	0.65
	AUC, t = 3	0.71	0.82	0.6
LL2, Scheme 1	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0001	0.0002	0.0002
	AUC, t = 2	0.95	0.97	0.98
	AUC, t = 3	0.92	0.95	0.98
	MSE_{TE}	0.91	0.88	1.07
	AUC, t = 2	0.8	0.86	0.75
	AUC, t = 3	0.85	0.84	0.83
W1, Scheme 1	R^2_{COMP}	100	100	99.5
	MSE_{TR}	0.0002	0.0001	0.006
	AUC, t = 2	0.92	0.9	0.93
	AUC, t = 3	0.92	0.9	0.93
	MSE_{TE}	2.39	2.16	2.6
	AUC, t = 2	0.75	0.78	0.65
	AUC, t = 3	0.75	0.78	0.65
W2, Scheme 1	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0003	0.0003	0.0004
	AUC, t = 2	0.91	0.91	0.87
	AUC, t = 3	0.93	0.91	0.89
	MSE_{TE}	3.99	3.98	4.01
	AUC, t = 2	0.89	0.81	0.81
	AUC, t = 3	0.86	0.81	0.74

All methods are based on $\alpha = 0.5$.

Table 14: Simulation Scheme 2: 80% Censoring

Model	Measure	ACPR-AFT $GenF$	ACPR-AFT $sAFT$	CPR-AFT
LN, Scheme 2	R^2_{COMP}	98.9	98.4	98.1
	MSE_{TR}	0.035	0.12	0.14
	AUC, t = 2	0.73	0.74	0.73
	AUC, t = 3	0.83	0.82	0.79
	MSE_{TE}	3.76	3.59	3.76
	AUC, t = 2	0.89	0.95	0.84
	AUC, t = 3	0.95	0.89	0.86
LL1, Scheme 2	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0002	0.0002	0.0005
	AUC, t = 2	0.95	0.97	0.93
	AUC, t = 3	0.94	0.96	0.92
	MSE_{TE}	1.72	1.72	1.75
	AUC, t = 2	0.68	0.79	0.66
	AUC, t = 3	0.7	0.83	0.55
LL2, Scheme 2	R^2_{COMP}	100	100	97.7
	MSE_{TR}	0.0003	0.00001	0.026
	AUC, t = 2	0.86	0.89	0.68
	AUC, t = 3	0.81	0.9	0.63
	MSE_{TE}	1.23	1.13	1.33
	AUC, t = 2	0.71	0.84	0.61
	AUC, t = 3	0.62	0.91	0.61
W1, Scheme 2	R^2_{COMP}	100	100	100
	MSE_{TR}	0.00002	0.00003	0.00002
	AUC, t = 2	0.96	0.95	0.93
	AUC, t = 3	0.96	0.95	0.93
	MSE_{TE}	6.17	4.8	4.55
	AUC, t = 2	0.75	0.88	0.68
	AUC, t = 3	0.75	0.88	0.68
W2, Scheme 2	R^2_{COMP}	100	100	100
	MSE_{TR}	0.0006	0.0007	0.0009
	AUC, t = 2	0.98	0.99	0.98
	AUC, t = 3	0.98	0.98	0.97
	MSE_{TE}	4.73	4.29	5.25
	AUC, t = 2	0.9	0.9	0.86
	AUC, t = 3	0.79	0.82	0.66

All methods are based on $\alpha = 0.5$.

Table 15: (A)CPR-AFT: Summary of Results

Method	HNSCC		GBM		Ovarian		Oral	
	K	R^2	K	R^2	K	R^2	K	R^2
ACPR-AFT $GenF$	8	100%	5	100%	15	100%	9	100%
ACPR-AFT $sAFT$	8	100%	5	100%	15	100%	5	97%
CPR-AFT	12	100%	5	100%	2	70%	5	84%
	2	92%	2	93%				
ACPR-AFT $GenF$	2	96%	2	93%	2	95%	5	97%
ACPR-AFT $sAFT$	2	98%	2	94%	2	93%	5	97%

Table 16: HNSCC Data: Summary of results on training and test sets

Marginal screening method (Number of features)	Set	Measure	ACPR-AFT $GenF$	ACPR-AFT $sAFT$	CPR-AFT $GenF^a$	CPR-AFT $sAFT^a$
sAFT (2,336)	Training	R^2	99.0	99.7	96.9	91.7
		MSE_{TR}	0.66	0.19	0.86	1.08
		AUC, t = 2	0.88	0.91	0.84	0.77
		AUC, t = 3	0.87	0.96	0.84	0.75
		AUC, t = 5	0.79	0.93	0.74	0.65
	Test	MSE_{TE}	1.35	1.47	1.82	1.5
		AUC, t = 2	0.72	0.76	0.66	0.64
		AUC, t = 3	0.67	0.72	0.61	0.62
		AUC, t = 5	0.61	0.67	0.55	0.56
CON (2,254)	Training	R^2	99.0	99.2	91.8	91.8
		MSE_{TR}	0.96	0.57	1.06	1.19
		AUC, t = 2	0.76	0.97	0.72	0.73
		AUC, t = 3	0.73	0.96	0.71	0.71
		AUC, t = 5	0.73	0.94	0.66	0.69
	Test	MSE_{TE}	1.30	1.27	1.40	1.44
		AUC, t = 2	0.67	0.7	0.62	0.62
		AUC, t = 3	0.68	0.72	0.65	0.62
		AUC, t = 5	0.65	0.71	0.57	0.58

a: based on α chosen by the corresponding ACPR-AFT method.

Table 17: GBM Data: Summary of results on training and test sets

Marginal screening method (Number of features)	Set	Measure	ACPR-AFT $GenF$	ACPR-AFT $sAFT$	CPR-AFT $GenF^a$	CPR-AFT $sAFT^a$
sAFT (3,605)	Training	R^2	87.9	90.3	85.0	85.6
		MSE_{TR}	1.10	0.89	0.90	0.90
		AUC, t = 2	0.63	0.74	0.63	0.71
		AUC, t = 3	0.70	0.83	0.70	0.69
		AUC, t = 5	0.95	0.94	0.80	0.86
	Test	MSE_{TE}	1.42	1.33	1.45	1.35
		AUC, t = 2	0.61	0.62	0.61	0.61
		AUC, t = 3	0.69	0.66	0.68	0.67
		AUC, t = 5	0.82	0.82	0.82	0.77
CON (1,125)	Training	R^2	88.7	90.5	83.5	80.1
		MSE_{TR}	0.85	1.08	0.90	1.12
		AUC, t = 2	0.77	0.74	0.75	0.69
		AUC, t = 3	0.78	0.85	0.76	0.79
		AUC, t = 5	0.93	0.94	0.91	0.90
	Test	MSE_{TE}	1.31	1.22	1.33	1.35
		AUC, t = 2	0.63	0.65	0.57	0.58
		AUC, t = 3	0.79	0.74	0.69	0.69
		AUC, t = 5	0.90	0.93	0.79	0.78

a: based on α chosen by the corresponding ACPR-AFT method.

Table 18: Ovarian Cancer Data: Summary of results on training and test sets

Marginal screening method (Number of features)	Set	Measure	ACPR-AFT $GenF$	ACPR-AFT $sAFT$	CPR-AFT $GenF^a$	CPR-AFT $sAFT^a$
sAFT (3,550)	Training	R^2	95.5	99.8	72.3	71.4
		MSE_{TR}	0.07	0.02	0.44	0.47
		AUC, t = 2	0.95	1.00	0.80	0.80
		AUC, $t = 3$	0.85	1.00	0.80	0.84
		AUC, t = 5	0.88	1.00	0.76	0.83
	Test	MSE_{TE}	0.53	0.5	0.58	0.51
		AUC, t = 2	0.77	0.83	0.71	0.74
		AUC, $t = 3$	0.77	0.86	0.71	0.73
		AUC, $t = 5$	0.75	0.83	0.66	0.66
CON (1,858)	Training	R^2	94.3	96.0	69.2	69.6
		MSE_{TR}	0.24	0.48	0.54	0.58
		AUC, t = 2	0.82	0.90	0.72	0.76
		AUC, $t = 3$	0.89	0.93	0.76	0.76
		AUC, t = 5	0.87	0.93	0.75	0.76
	Test	MSE_{TE}	0.29	0.66	0.67	0.73
		AUC, t = 2	0.78	0.78	0.64	0.67
		AUC, $t = 3$	0.84	0.82	0.69	0.69
		AUC, $t = 5$	0.82	0.83	0.70	0.72

a: based on α chosen by the corresponding ACPR-AFT method.

Table 19: Oral Cancer Data: Summary of results on training and test sets

Marginal screening method (Number of features)	Set	Measure	ACPR-AFT $GenF$	ACPR-AFT $sAFT$	CPR-AFT $GenF^a$	CPR-AFT $sAFT^a$
sAFT (3,065)	Training	R^2	94.6	99	93.7	95.1
		MSE_{TR}	0.03	0.02	0.21	0.14
		AUC, t = 2	0.98	1.00	0.97	0.97
		AUC, t = 3	0.97	1.00	0.96	0.98
		AUC, t = 5	0.96	1.00	0.97	0.98
	Test	MSE_{TE}	0.10	0.34	1.08	0.99
		AUC, t = 2	0.89	0.87	0.86	0.85
		AUC, t = 3	0.89	0.92	0.89	0.89
		AUC, t = 5	0.94	0.95	0.93	0.94
CON (4,652)	Training	R^2	95.6	95.6	89.3	88.3
		MSE_{TR}	0.01	0.12	0.40	0.4
		AUC, t = 2	0.96	0.97	0.95	0.94
		AUC, t = 3	0.98	0.99	0.97	0.97
		AUC, t = 5	0.97	0.99	0.96	0.98
	Test	MSE_{TE}	0.36	0.44	1.20	1.17
		AUC, t = 2	0.81	0.83	0.82	0.82
		AUC, t = 3	0.86	0.88	0.86	0.86
		AUC, t = 5	0.90	0.92	0.90	0.91

a: based on α chosen by the corresponding ACPR-AFT method.

Table 20: HNSCC Data: Summary of Prediction Measures

Marginal screening method (Number of features)	Measure	ACPR-AFT $GenF^a$	ACPR-AFT $sAFT^b$	CPR-AFT^c
sAFT (2,336)	R^2	100	99.8	98.3
	MSE	0.004	0.07	0.53
	AUC, t = 2	0.81	0.96	0.73
	AUC, t = 3	0.85	0.95	0.71
	AUC, t = 5	0.67	0.89	0.67
CON (2,254)	R^2	99.0	99.1	91.3
	MSE	0.95	0.62	1.18
	AUC, t = 2	0.69	0.92	0.69
	AUC, t = 3	0.70	0.93	0.69
	AUC, t = 5	0.67	0.9	0.66
GenF (2,530)	R^2	99.7	99.1	91.3, 90.4
	MSE	0.14	0.62	1.15, 1.10
	AUC, t = 2	0.94	0.94	0.71, 0.67
	AUC, t = 3	0.89	0.92	0.70, 0.69
	AUC, t = 5	0.82	0.82	0.58, 0.61

$a : \alpha = 0.5$ for all filters; $b : \alpha = 0.5$ for AFT/CON and $\alpha = 0.75$ for $GenF$; $c :$ optimal α chosen by the corresponding ACPR-AFT method used.

(a) 33% Censoring, ω (b) 80% Censoring, ω

(c) 33% Censoring, ξ (d) 80% Censoring, ξ

Figure 1: ROC Curves for Simulation Scheme 1

(a) 33% censoring

(b) 80% censoring

(c) 33% censoring

(d) 80% censoring

Figure 2: ROC Curves for Simulation Scheme 2, ξ vs. ω for $GenF$ (a, b) and $GenF$ vs. $\text{PLS}_{\text{unadj}}$ using ξ (c, d)

(a) HNSCC data

(b) GBM data

(c) Ovarian data

(d) Oral data

Figure 3: Predicted survival curves based on PI